

Preprint no. NJU-INP 011/19

Two photon transition form factors of neutral pseudoscalar mesons

Minghui Ding* and Daniele Binosi

European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT) and Fondazione Bruno Kessler*

Villa Tambosi, Strada delle Tabarelle 286, I-38123 Villazzano (TN) Italy

**E-mail: mding@ectstar.eu*

Khépani Raya and Lei Chang

School of Physics, Nankai University, Tianjin 300071, China

Craig D. Roberts

School of Physics, Nanjing University, Nanjing, Jiangsu 210093, China

Institute for Nonperturbative Physics, Nanjing University, Nanjing, Jiangsu 210093, China

Starting from the axial vector Ward-Green-Takahashi identity, and the necessity of preserving this identity when attempting to calculate the properties describing pseudoscalar mesons, we describe a Bethe-Salpeter kernel which takes into account the non-Abelian anomaly and therefore is suitable to describe the flavor mixing states η, η' . We then report results for the two photon transition form factors of η, η' , and compare them with existing experimental data.

Keywords: Pseudoscalar meson; Transition form factor; Non-Abelian anomaly.

1. Ward-Green-Takahashi identity

Ward's identity¹ and its generalization by Green² and Takahashi³, ought to be preserved by any approach that is willing to capture QCD's fundamental features. In its axial vector formulation this identity reads

$$P^\mu \Gamma_{5\mu}^a(k; P) = S^{-1}(k_+) i\gamma_5 \mathcal{F}^a + i\gamma_5 \mathcal{F}^a S^{-1}(k_-) - 2i\mathcal{M}^{ab} \Gamma_5^b(k; P) - \mathcal{A}^a(k; P), \quad (1)$$

where: $P(k)$ is the total (relative) momentum between the amputated quark legs; $\{\mathcal{F}^a | a = 0, \dots, N_f^2 - 1\}$ are the generators of $U(N_f)$ in the fundamental representation, orthonormalized according to $\text{tr} \mathcal{F}^a \mathcal{F}^b = \frac{1}{2} \delta^{ab}$; $\mathcal{M}^{ab} = \text{tr}_F [\{\mathcal{F}^a, \mathcal{M}^{\text{bm}}\} \mathcal{F}^b]$, with $\mathcal{M}^{\text{bm}} = \text{diag}[m_u^{\text{bm}}, m_d^{\text{bm}}, m_s^{\text{bm}}, \dots]$ the current-quark bare masses matrix; $\mathcal{S} = \text{diag}[S_u, S_d, S_s, \dots]$ the dressed-quark propagator matrix, with nonzero entries determined by the corresponding Dyson-Schwinger equation; $\Gamma_{5\mu}^a$ and Γ_5^a the axial-vector and

pseudoscalar vertex respectively, both satisfying the corresponding inhomogeneous Bethe-Salpeter equations; and, finally, \mathcal{A}^a in an expression of the non-Abelian axial anomaly and can be defined through the topological charge density operator. Notice that only $\mathcal{A}^{a=0}$ is nonzero: therefore, flavor-nonsinglet neutral pseudoscalar mesons satisfy the standard identity, while flavor-singlet state satisfies the anomalous one⁴.

One of the far-reaching consequences of the Ward-Green-Takahashi (WGT) identity (1) is a mass formula for pseudoscalar mesons which is valid irrespectively from the magnitude of the current-quark masses. It reads

$$m_M^2 f_M^a = 2 \mathcal{M}^{ab} \rho_M^b + \delta^{a0} n_M, \quad (2)$$

with: m_M the meson mass; f_M^a the decay constant; ρ_M^b the pseudoscalar projection of the meson's Bethe-Salpeter wave function onto the origin in configuration space; and n_M a quantity derived from topological charge density operator. This one single equation has a wide range of effects on Goldstone modes, excited states as well as system(s) described by the non-Abelian axial anomaly⁴⁻⁶; in particular, it shows that:

- The Nambu-Goldstone pion is a consequence of Dynamical Chiral Symmetry Breaking (DCSB).
- In the chiral limit, the decay constants of the pion's radial excitations are zero.
- η' is split from the octet pseudoscalars by an amount that depends on QCD's topological susceptibility.

2. Kernels for the bound-state equations

Yet another consequence of Eq. (1) is that it leads to a symmetry-preserving truncation scheme, the simplest of which is the Rainbow-Ladder (RL) approximation. RL truncation is known to be accurate for ground-state light-quark vector- and isospin-nonzero-pseudoscalar-mesons⁷⁻¹⁰, and has been used in a recent study on pion parton distribution functions¹¹. Implemented in the Bethe-Salpeter and gap equations, RL truncation is expressed via the following kernels, respectively:

$$[\mathcal{K}_N(k, q; P)]_{\ell_1 \ell_2}^{\ell_1' \ell_2'} = -\frac{4}{3} \mathcal{G}_{\mu\nu}(t) [i\gamma_\mu]_{\ell_1 \ell_1'} [i\gamma_\nu]_{\ell_2' \ell_2}, \quad (3a)$$

$$\mathcal{G}_{\mu\nu}(t = k - q) = \tilde{\mathcal{G}}(t^2) T_{\mu\nu}(t), \quad (3b)$$

where: $t^2 T_{\mu\nu} = t^2 \delta_{\mu\nu} - t_\mu t_\nu$; and $\tilde{\mathcal{G}}(t^2)$ is the interaction consistent with that determined in studies of QCD's gauge sector¹².

However, RL truncation fails for the η - and η' -mesons as they do not produce vertices that satisfy the anomalous axial-vector WGT identity. Indeed, whilst the RL kernel is constrained by a large body of successful phenomenology, the form of the anomalous kernel is unknown. Considering the structure of the non-Abelian anomaly, it readily becomes apparent that no related contribution to the Bethe-Salpeter kernel can contain external quark or antiquark lines which are simply connected to the internal lines: purely gluonic configurations must mediate. Moreover, no finite sum of diagrams can be sufficient⁴. On general grounds, its contribution to the Bethe-Salpeter equation for pseudoscalar mesons must take the following form¹³:

$$[\mathcal{K}_A(k, q; P)]_{\ell_1 \ell_2}^{\ell'_1 \ell'_2} = - \sum_{i=1}^4 \sum_{f=l,s} a_i^f(k, q; P) \times [\mathcal{F}^f \mathcal{D}_i(q; P)]_{\ell'_2 \ell'_1} [\mathcal{F}^f \mathcal{D}_i(k; P)]_{\ell_1 \ell_2}, \quad (4)$$

where: $\{\mathcal{D}_i(k; P) | i = 1, \dots, 4\}$ constitute a Dirac basis of the pseudoscalar meson Bethe-Salpeter wave function; and $\{a_i^f(k, q; P) | i = 1, \dots, 4; f = l, s\}$ are scalar functions. An educated Ansatz for \mathcal{K}_A had been put forward in Ref. 13, which being essentially nonperturbative in nature, provided material support at infrared momenta. In this way the corresponding Bethe-Salpeter kernel becomes the sum of two terms: $\mathcal{K} = \mathcal{K}_V + \mathcal{K}_A$, and with the bound-state kernel thus defined, results for the $\gamma^* \gamma \rightarrow \eta, \eta'$ transition form factors have been obtained.

3. Two-photon transition form factors

The two-photon transition form factor provides a window for exploring the internal structure of neutral pseudoscalar meson^{14–16}. The parameters in the gap and Bethe-Salpeter equations can be fixed by requiring that the solutions of the coupled-channels bound-state problems reproduce the empirical η, η' masses and the four phenomenologically-determined values of the light- and strange-quark η, η' decay constants¹³. We find $m_\eta = 0.56$ GeV and $m_{\eta'} = 0.96$ GeV, to be compared with the experimental values¹⁷ 0.55 GeV and 0.96 GeV, respectively; other observables like decay constants, mixing angle and two-photon decay widths are reported in Ref. 13.

The transition form factors $F_{\eta, \eta'}(Q^2)$ obtained within the framework sketched above, are shown in Fig. 1 and compared with the asymptotic value of the π^0 form factor, $2f_\pi = 0.186$ GeV, drawn as the dotted (red) curve in both panels. There are marked similarities between $F_\eta(Q^2)$ and the π^0 tran-




Fig. 1. $\gamma^*\gamma \rightarrow \eta, \eta'$ transition form factors: left panel, η ; right panel, η' . Curves: solid (green), our prediction with complete evolution, with the shaded (green) band indicating an uncertainty in our prediction owing to variations in the value of f_η^s ; dashed (blue), asymptotic limit, with the banded-shading (blue) region indicating the impact of uncertainty in f_η^s on the asymptotic behaviour. The broader, shaded (grey) band combines this with the uncertainty owing to omission of φ_M^0 - φ_M^g mixing (φ_M^0 is the flavor singlet distribution amplitude and φ_M^g is the two-gluon distribution amplitude). In both panels, the dotted (red) curve is the π^0 asymptotic limit, $2f_\pi$. Experimental data are¹⁸: diamonds (blue) “CLEO”; circles (red) “BaBar”.

sition form factor (see, for example, Ref. 14, Fig. 2); furthermore, our full $F_{\eta, \eta'}(Q^2)$ result (solid, green curve) agree well with existing data. Finally, at asymptotically large momentum transfers, *i.e.* on $\tau^2 := \Lambda_{\text{QCD}}^2/Q^2 \simeq 0$, our full results meet the asymptotic trajectory (dashed blue curve).

The scope of the analysis sketched herein could also be extended to include the doubly off-shell transition form factors, for which the first data $\gamma^*(k_1)\gamma^*(k_2) \rightarrow \eta'$ now exist¹⁹.

Acknowledgments: Work partially supported by Jiangsu Province *Hundred Talents Plan for Professionals*.

References

1. J. C. Ward, An Identity in Quantum Electrodynamics, *Phys. Rev.* **78**, 182 (1950).
2. H. S. Green, A Pre-renormalized quantum electrodynamics, *Proc. Phys. Soc.* **A66**, 873-880 (1953).
3. Y. Takahashi, On the generalized Ward identity, *Nuovo Cim.* **6**, 371 (1957).
4. M. S. Bhagwat, L. Chang, Y. X. Liu, C. D. Roberts, and P. C. Tandy, Flavour symmetry breaking and meson masses, *Phys. Rev.*

- C76**, 045203 (2007).
5. P. Maris, C. D. Roberts, and P. C. Tandy, Pion mass and decay constant, *Phys. Lett.* **B420**, 267-273 (1998).
 6. A. Holl, A. Krassnigg, C. D. Roberts, and S. V. Wright, On the complexification of pseudoscalar mesons, *Int. J. Mod. Phys.* **A20**, 1778-1784 (2005).
 7. T. Horn and C. D. Roberts, The pion: an enigma within the Standard Model, *J. Phys.* **G43**, no.7, 073001 (2016).
 8. G. Eichmann, H. Sanchis-Alepuz, R. Williams, R. Alkofer, and C. S. Fischer, Baryons as relativistic three-quark bound states, *Prog. Part. Nucl. Phys.* **91**, 1-100 (2016).
 9. C. S. Fischer, QCD at finite temperature and chemical potential from DysonSchwinger equations, *Prog. Part. Nucl. Phys.* **105**, 1-60 (2019).
 10. V. D. Burkert, and C. D. Roberts, Colloquium : Roper resonance: Toward a solution to the fifty year puzzle, *Rev. Mod. Phys.* **91**, 011003 (2019).
 11. M. Ding, K. Raya, D. Binosi, L. Chang, C. D. Roberts and S. M. Schmidt, Symmetry, symmetry breaking, and pion parton distributions, (arXiv:1905.05208 [nucl-th]).
 12. D. Binosi, L. Chang, J. Papavassiliou and C. D. Roberts, Bridging a gap between continuum-QCD and ab initio predictions of hadron observables, *Phys. Lett.* **B742**, 183-188 (2015).
 13. M. Ding, K. Raya, A. Bashir, D. Binosi, L. Chang, M. Chen and C. D. Roberts, $\gamma^*\gamma \rightarrow \eta, \eta'$ transition form factors, *Phys. Rev.* **D99**, 014014 (2019).
 14. K. Raya *et al.*, Structure of the neutral pion and its electromagnetic transition form factor, *Phys. Rev.* **D93**, 074017 (2016).
 15. K. Raya, M. Ding, A. Bashir, L. Chang, and C. D. Roberts, Partonic structure of neutral pseudoscalars via two photon transition form factors, *Phys. Rev.* **D95**, 074014 (2017).
 16. J. Chen, M. Ding, L. Chang and Y. X. Liu, Two Photon Transition Form Factor of $\bar{c}c$ Quarkonia, *Phys. Rev.* **D95**, 016010 (2017).
 17. M. Tanabashi *et al.*, Review of Particle Physics, *Phys. Rev.* **D98**, 030001 (2018).
 18. P. del Amo Sanchez *et al.*, Measurement of the $\gamma\gamma^* \rightarrow \eta$ and $\gamma\gamma^* \rightarrow \eta'$ transition form factors, *Phys. Rev.* **D84**, 052001 (2011).
 19. J. P. Lees *et al.*, Measurement of the $\gamma^*\gamma^* \rightarrow \eta'$ transition form factor, *Phys. Rev.* **D98**, 112002 (2018).