



XXVIIIth International Conference on Ultrarelativistic Nucleus-Nucleus Collisions
(Quark Matter 2019)

Search for jet quenching effects in high multiplicity pp collisions at $\sqrt{s} = 13$ TeV

P. M. Jacobs for the ALICE Collaboration

Lawrence Berkeley National Laboratory

Abstract

The ALICE Collaboration reports a search for jet quenching effects in pp collisions at $\sqrt{s} = 13$ TeV, in events selected on high multiplicity compared to the minimum bias population. The measurement is based on the semi-inclusive acoplanarity distribution of jets recoiling from a high- p_T trigger hadron. Significant broadening of the recoil jet acoplanarity distribution is observed in high multiplicity pp collisions, in both data and in simulations based on the PYTHIA model. Analysis is ongoing to elucidate the origin of this effect.

1. Introduction

High-energy collisions of small systems, in which one or both projectiles is a proton or light nucleus, exhibit evidence of collective flow [1, 2, 3], raising the question whether a Quark-Gluon Plasma (QGP) is produced in such collisions [4, 5]. Jet quenching, the modification of a jet shower due to its interaction with dense QCD matter, is a necessary consequence of the formation of a QGP. Observation of jet quenching effects in small collision systems would provide compelling evidence of the formation of a QGP.

Jet quenching generates multiple related phenomena: yield suppression of high- p_T hadrons and of jets with finite radius R ; modification of jet substructure; and jet centroid deflection (medium-induced acoplanarity). Experimentally, observables based on these various phenomena have different sensitivities to jet quenching effects, depending upon collision system and kinematics.

The search for jet quenching effects in small systems is carried out by comparing distributions from collisions with small and large event activity (EA), measured by forward multiplicity or E_T . Inclusive hadron and jet production measurements as a function of EA in p–A collisions yield no significant evidence of yield suppression due to quenching (e.g. [6]). However, such inclusive yield measurements have limited precision due to systematic uncertainties in calculating Glauber scaling factors in small systems [7], and coincidence observables are required for more precise quenching measurements in small systems. The first such search set a limit of 400 MeV (90% CL) for medium-induced out-of-cone energy transport in high vs low EA p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV [8]. Further progress requires yet better precision for measurements such as these, together with measurements based on new observables and additional small collision systems.

In these proceedings, we present the first search for jet quenching in pp collisions. Since Glauber scaling between pp event populations with different EA is not defined, jet quenching measurements based on inclusive yield suppression are not possible. In this analysis we focus on medium-induced acoplanarity [9, 10], which is measured via the semi-inclusive distribution of jets recoiling from a high- p_T charged hadron [11, 12]. There is significant acoplanarity even in the absence of in-medium broadening [13], and medium-induced acoplanarity is identified by additional broadening relative to this vacuum effect. The search for jet quenching using acoplanarity is carried out by comparing populations with different EA, high multiplicity (HM) vs. minimum bias (MB).

2. Analysis

Jets are reconstructed from charged tracks in the ALICE acceptance, using the anti- k_T algorithm with $R = 0.4$. We utilize the observable Δ_{recoil} [11], which is the difference of two semi-inclusive distributions with two widely different ranges of trigger hadron kinematics $p_{T,\text{trig}}$ (TT),

$$\Delta_{\text{recoil}}(p_{T,\text{jet}}^{\text{ch, reco}}, \Delta\varphi) = \frac{1}{N_{\text{trig}}} \frac{d^2 N_{\text{jets}}}{dp_{T,\text{jet}}^{\text{ch, reco}} d\Delta\varphi} \Big|_{p_{T,\text{trig}} \in \text{TT}_{\text{Sig}}} - c_{\text{Ref}} \cdot \frac{1}{N_{\text{trig}}} \frac{d^2 N_{\text{jets}}}{dp_{T,\text{jet}}^{\text{ch, reco}} d\Delta\varphi} \Big|_{p_{T,\text{trig}} \in \text{TT}_{\text{Ref}}}, \quad (1)$$

where $\Delta\varphi$ is the azimuthal angle between trigger hadron and recoil jet. The trigger hadron selections are TT_{Sig} : $20 < p_{T,\text{trig}} < 30$ GeV/ c (denoted TT{20,30}); and TT_{Ref} : $6 < p_{T,\text{trig}} < 7$ GeV/ c (denoted TT{6,7}). The standard area-based approach is applied to adjust $p_{T,\text{jet}}^{\text{ch}}$ for the median underlying-event density ($p_{T,\text{jet}}^{\text{ch, reco}}$) [14]. The constant c_{Ref} is extracted from data [11], with value $\sim 0.95 - 1.0$. The Δ_{recoil} observable provides fully data-driven suppression of background jet yield that is uncorrelated with the hadron trigger. This is of particular importance for acoplanarity measurements, in order to suppress the effects of high- Q^2 Multiple Parton Interactions (MPIs) (see [11] for discussion).

The data were recorded by ALICE during LHC Run 2 with pp collisions at $\sqrt{s} = 13$ TeV. Two on-line triggers were applied: an MB trigger collected 3.2B events (equivalent to 0.098 pb^{-1}), and a high-multiplicity trigger (HM–Online), based on the forward scintillators V0A ($2.8 < \eta < 5.1$) and V0C ($-3.7 < \eta < -1.7$), sampled 13 pb^{-1} . Events are classified according to their summed V0 signal, $\text{VOM} = \text{V0A} + \text{V0C}$. In order to combine runs with different V0 gain and to facilitate comparison to theoretical calculations, we utilize the scaled multiplicity $\text{VOM}/\langle \text{VOM} \rangle$, where the denominator is the mean of VOM for the MB population.




Fig. 1. Scaled multiplicity distribution recorded by the MB trigger. Vertical dashed lines indicate the High Multiplicity (HM) offline selection.

Figure 1 shows the uncorrected $\text{VOM}/\langle \text{VOM} \rangle$ distribution measured by the MB trigger (HM–Online distribution not shown). In this analysis we compare the MB population to an offline-selected HM population defined by $5 < (\text{VOM}/\langle \text{VOM} \rangle) < 9$, indicated by the vertical dashed lines. The HM selection

corresponds to 0.1% of the MB cross section. The figure also shows the $V0M/\langle V0M \rangle$ distribution calculated by PYTHIA Monash at the particle level, showing qualitative agreement with data (albeit at the particle level) for $(V0M/\langle V0M \rangle) < 3$, but lower yield than data at higher $V0M/\langle V0M \rangle$.





Fig. 2. Semi-inclusive distributions of recoil jets in HM-selected events for TT{6,7}, TT{20,30}, and the distribution of Δ_{recoil} (Eq. 1). Left: vs. $p_{T,\text{jet}}^{\text{ch, reco}}$ for $\Delta\varphi > \pi - 0.6$; right: vs. $\Delta\varphi$ for $20 < p_{T,\text{jet}}^{\text{ch, reco}} < 30$ GeV/c.

Figure 2 shows the distributions of Δ_{recoil} and its two components as a function of $p_{T,\text{jet}}^{\text{ch, reco}}$ and $\Delta\varphi$. There is significant correlated yield at large recoil angles relative to $\Delta\varphi = \pi$, following the subtraction in Eq. 1.

3. Results



ALI-PREL-339740

Fig. 3. Uncorrected acoplanarity distributions for intervals in $p_{T,\text{jet}}^{\text{ch, reco}}$, for MB and HM selections in EA. Left: ALICE data; right: PYTHIA Monash particle-level simulation.

Figure 3, left panels, show $\Delta\varphi$ distributions in intervals of $p_{T,\text{jet}}^{\text{ch, reco}}$ for both MB and HM-selected ALICE data, not corrected for instrumental and background smearing effects. They exhibit a striking effect: for

$p_{T,\text{jet}}^{\text{ch, reco}} < 40 \text{ GeV}/c$, the HM acoplanarity distributions are significantly suppressed at $\Delta\varphi \sim \pi$ and broadened away from the peak. This is qualitatively what is expected from jet quenching in high-EA collisions.

Before concluding that jet quenching has been observed in HM pp collisions, all other potential sources must be excluded. Azimuthal broadening due to uncorrelated background was measured by embedding PYTHIA Monash events (detector-level) in HM-selected real events and repeating the analysis; negligible broadening is observed, so that the effect is not due to high track density. Likewise, the effect is observed qualitatively when comparing only the Signal distributions in Eq. 1, so that the effect is not due to the subtraction in Δ_{recoil} . The broadening is a significant physical effect.

Figure 3, right panels, show similar distributions calculated using PYTHIA Monash at the particle level, in this case with HM corresponding to $4 < (V0M/\langle V0M \rangle) < 9$; qualitative comparison only may be made to data. Nevertheless, the PYTHIA-generated distributions are qualitatively similar to those of the data, with broadened acoplanarity for HM-selected events.

Finally, we have explored the evolution of the properties of EA-selected pp collisions by measuring the distribution of forward-backward multiplicity asymmetry as a function of $V0M/\langle V0M \rangle$ (figures not shown). A narrowing of the asymmetry distribution is observed as EA increases; scaling of its width with EA is consistent with counting statistics being the mechanism driving the change in width, with clustering (e.g. due to enhanced multi-jet production at high EA) playing a lesser or negligible role.

4. Outlook

A significant broadening of the recoil jet acoplanarity distribution in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ has been observed in High EA-selected events compared to the MB population, for $p_{T,\text{jet}}^{\text{ch, reco}} < 40 \text{ GeV}/c$. While this effect is characteristic of jet quenching, qualitatively similar features are observed in PYTHIA simulations. Additionally, fluctuation properties of the multiplicity distribution do not change significantly as a function of EA, suggesting similar production mechanisms at low and high-EA. Determination of the origin of the broadening for high-EA requires new calculations based on PYTHIA and other Monte Carlo generators, in which candidate mechanisms are selectively modified, with additional constraints imposed by measurements of forward/backward multiplicity fluctuations. Such calculations will help elucidate the role of jet quenching in the observed acoplanarity broadening.

References

- [1] V. Khachatryan, et al., Observation of Long-Range Near-Side Angular Correlations in Proton-Proton Collisions at the LHC, *JHEP* 1009 (2010) 091. [arXiv:1009.4122](#).
- [2] B. Abelev, et al., Long-range angular correlations on the near and away side in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$, *Phys.Lett. B* 719 (2013) 29–41. [arXiv:1212.2001](#).
- [3] C. Aidala, et al., Creation of quark-gluon plasma droplets with three distinct geometries, *Nature Phys.* 15 (3) (2019) 214–220. [arXiv:1805.02973](#).
- [4] R. D. Weller, P. Romatschke, One fluid to rule them all: viscous hydrodynamic description of event-by-event central p+p, p+Pb and Pb+Pb collisions at $\sqrt{s} = 5.02 \text{ TeV}$, *Phys. Lett. B* 774 (2017) 351–356. [arXiv:1701.07145](#).
- [5] M. Mace, V. V. Skokov, P. Tribedy, R. Venugopalan, Systematics of azimuthal anisotropy harmonics in proton-nucleus collisions at the LHC from the Color Glass Condensate, *Phys. Lett. B* 788 (2019) 161–165. [arXiv:1807.00825](#).
- [6] J. Adam, et al., Centrality dependence of charged jet production in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$, *Eur.Phys.J. C* 76 (2016) 271. [arXiv:1603.03402](#).
- [7] J. Adam, et al., Centrality dependence of particle production in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$, *Phys. Rev. C* 91 (6) (2015) 064905. [arXiv:1412.6828](#).
- [8] S. Acharya, et al., Constraints on jet quenching in p–Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ measured by the event-activity dependence of semi-inclusive hadron-jet distributions, *Phys. Lett. B* 783 (2018) 95–113. [arXiv:1712.05603](#).
- [9] D. A. Appel, Jets as a Probe of Quark - Gluon Plasmas, *Phys. Rev. D* 33 (1986) 717.
- [10] J. P. Blaizot, L. D. McLerran, Jets in Expanding Quark–Gluon Plasmas, *Phys. Rev. D* 34 (1986) 2739.
- [11] J. Adam, et al., Measurement of jet quenching with semi-inclusive hadron-jet distributions in central Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$, *JHEP* 09 (2015) 170. [arXiv:1506.03984](#).
- [12] L. Adamczyk, et al., Measurements of jet quenching with semi-inclusive hadron+jet distributions in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$, *Phys. Rev. C* 96 (2) (2017) 024905. [arXiv:1702.01108](#).
- [13] L. Chen, G.-Y. Qin, S.-Y. Wei, B.-W. Xiao, H.-Z. Zhang, Probing Transverse Momentum Broadening via Dihadron and Hadron-jet Angular Correlations in Relativistic Heavy-ion Collisions, *Phys. Lett. B* 773 (2017) 672–676. [arXiv:1607.01932](#).
- [14] M. Cacciari, G. P. Salam, Pileup subtraction using jet areas, *Phys. Lett. B* 659 (2008) 119–126. [arXiv:0707.1378](#).