## A Sufficient Condition For An Operator To Map $uL^{\infty}$ to $BMO_u$

Sakin Demir

February 1, 2022

## Abstract

Let T be an operator and suppose that there exists a positive constant C such that

$$\left(\int_{I} |Tf(x)|^{q} dx\right)^{1/q} \le C \left(\int_{I} |f(x)|^{q} dx\right)^{1/q}$$

for every q which is near enough to 1 and for every interval I in  $\mathbb{R}$  and  $f \in L^{\infty}(\mathbb{R})$ . Then we show that T maps  $uL^{\infty}$  to BMO<sub>u</sub>.

Mathematics Subject Classification: 47B38. Keywords: Weighted BMO Space,  $L^{\infty}$  Space.

We say that a non-negative function  $w \in L^1_{loc}(\mathbb{R})$  is an  $A_p$  weight for some 1 if

$$\sup_{I} \left[ \frac{1}{|I|} \int_{I} w(x) \, dx \right] \left[ \frac{1}{|I|} \int_{I} [w(x)]^{-1/(p-1)} \, dx \right]^{p-1} \le C < \infty$$

The supremum is taken over all intervals  $I \subset \mathbb{R}^n$ ; |I| denotes the measure of I.

w is an  $A_{\infty}$  weight if given an interval I there exists  $\delta > 0$  and  $\epsilon > 0$  such that for any measurable set  $E \subset I$ ,

$$|E| < \delta \cdot |I| \implies w(E) < (1 - \epsilon) \cdot w(I).$$

Here

$$w(E) = \int_E w(x) \, dx.$$

 $w \in A_1$  if

$$I(w) = \frac{1}{|I|} \int_{I} w(x) \, dx \le C \operatorname{ess\,inf}_{I} w$$

for all intervals I.

It is well known and can easily be seen that  $w \in A_{\infty}$  implies  $w \in A_p$  if 1 .

Before presenting our main result let us first mention some of the well known properties of the functions in  $A_p$ .

**Lemma 1** (Reverse Hölder Inequality). Let w satisfy  $A_p$ , where  $1 \le p < \infty$ . Then the inequality

$$\left(\frac{1}{|I|}\int_{I} (w(x))^{1+\delta} dx\right)^{\frac{1}{1+\delta}} \le C\left(\frac{1}{|I|}\int_{I} w(x) dx\right)$$

holds for all intervals  $I \subset \mathbb{R}$ , with constants  $C, \delta > 0$  independent of I.

*Proof.* This lemma is a part of the Corollary 2.13 in J. Garcia-Cuerva and J. L. Rubio de Francia [1] on page 403.  $\Box$ 

Note also that it is well known and easy to see that  $A_p \subset A_r$  for p < r. Let |I| denote the Lebesgue measure of the interval I and let

$$I(f) = \frac{1}{|I|} \int_{I} f.$$

We say that  $f \in BMO_u$  if

$$\frac{1}{|I|} \int_{I} |f - I(f)| \le CI(u)$$

for all intervals I in  $\mathbb{R}$ .

The sharp maximal function is defined as

$$f^{\sharp}(x) = \sup\left\{\frac{1}{|I|} \int_{I} |f - I(f)| : x \in I, \ I \text{ an interval}\right\}.$$

If 1 , then

 $||f^{\sharp}||_p \approx ||f||_p,$ 

provided  $||f||_p$  is finite.

Let T be an operator. We say that  $T: uL^{\infty} \to BMO_u$  provided

$$\frac{1}{|I|} \int_{I} |T(fu) - I(Tfu)| \le CI(u) ||f||_{\infty}$$

with C independent of I, and we say that  $T: L^p(u) \to L^p(u)$  if

$$||Tf||_{L^p(u)} \le C ||f||_{L^p(u)}.$$

We can now state and prove our main result:

**Theorem 1.** Let T be an operator and suppose that there exists a positive constant C such that

$$\left(\int_{I} |Tf(x)|^{q} dx\right)^{1/q} \leq C \left(\int_{I} |f(x)|^{q} dx\right)^{1/q}$$

for every q which is near enough to 1 and for every interval I in  $\mathbb{R}$  and  $f \in L^{\infty}(\mathbb{R})$ . Then T maps  $uL^{\infty}$  to  $BMO_u$  for all  $u \in A_1$ .

*Proof.* For this let  $u \in A_1$  and  $f \in L^{\infty}$ . We need to show that given an interval I in  $\mathbb{R}$ 

$$\frac{1}{|I|} \int_{I} |T(fu) - I(Tfu)| \le CI(u) ||f||_{\infty},$$

with a positive constant C independent of I.

Let q > 1 be near enough to 1 so that reverse Hölder inequality holds for u

with exponent q. Then we have

$$\begin{split} \frac{1}{|I|} \int_{I} |T(fu) - I(Tfu)| &\leq \frac{1}{|I|} \int_{I} |T(fu)| + \frac{1}{|I|} \int_{I} |I(Tfu)| \\ &\leq \frac{1}{|I|} \int_{I} |T(fu)| + \frac{1}{|I|} \int_{I} |(Tfu)| \\ &= \frac{2}{|I|} \int_{I} |T(fu)| \\ &\leq 2 \left( \frac{1}{|I|} \int_{I} |T(fu)|^{q} \right)^{1/q} \\ &\leq 2C \left( \frac{1}{|I|} \int_{I} |f|^{q} u^{q} \right)^{1/q} \quad \text{(by the hypothesis)} \\ &\leq 2C \|f\|_{\infty} \left( \frac{1}{|I|} \int_{I} u^{q} \right)^{1/q} \\ &\leq C_{1} \|f\|_{\infty} \frac{1}{|I|} \int_{I} u \quad \text{(by reverse Hölder inequality)} \\ &= C_{1} \|f\|_{\infty} I(u), \end{split}$$

where  $C_1$  is a positive constant independent of I. This shows that T maps  $uL^{\infty}$  to BMO<sub>u</sub>.

## References

[1] J. Garcia-Cuerva and J. L. Rubio de Francia, Weighted norm inequalities and related topics, North-Holland, Mathematics Studies 116, 1985.

Sakin Demir E-mail: sakin.demir@gmail.com Address: İbrahim Çeçen University Faculty of Education 04100 Ağrı, TURKEY.