On the global attractivity of non-autonomous neural networks with a distributed delay

Leonid Berezansky¹ and Elena Braverman²

¹ Dept. of Math, Ben-Gurion University of Negev, Beer-Sheva 84105, Israel

E-mail: brznsky@math.bgu.ac.il ² Dept. of Math & Stats, University of Calgary, 2500 University Dr. NW, Calgary, AB, Canada T2N 1N4

E-mail: maelena@math.ucalgary.ca

Abstract. We consider a system of several nonlinear equations with a distributed delay and obtain absolute asymptotic stability conditions, independent of the delay. The ideas of the proofs are based on the notion of a strong attractor. The results are applied to Hopfield neural networks, Nicholson's blowflies type system, and compartment models of population dynamics.

AMS Subject Classification: 34K20, 34K25, 92B20, 37C70

Keywords: distributed delay, non-autonomous systems, neural networks, global attractivity, Nicholson's blowflies model

1. Introduction

In many applications, systems experiencing delay in fact involve distributed delays. If the rate of change depends on the past, for example, reporting delays in economics or maturation delays in ecological systems, assuming constant concentrated delays is a significant simplification. Generally, the dependency will be on a certain segment of prehistory of the process, and, in certain cases, on the whole of it, leading to infinite delays. This is the reason why stability of systems with a distributed delay has been intensively investigated, let us mention some recent publications [1, 4, 17, 18, 20, 24, 25, 28, 30, 31], see also references therein. In biological models [21, 22, 32], a distributed delay sometimes is accepted by default. A neural network [23] is one of the most important applications where delays, in particular distributed, occur.

For scalar equations with a distributed delay, stability, either dependent or independent of the delay distribution, has been studied by several authors [5, 6, 8, 15, 33]. A particular case of a system of two equations was explored in [7, 16].

The purpose of the present paper is to obtain delay-independent stability conditions for a system of differential equations with a distributed delay

$$\frac{dX}{dt} = G(t) \left[\int_{H(t)} d_{\tau} R(t,\tau) F(X(\tau)) - X(t) \right], \qquad (1.1)$$

where $X : \mathbb{R} \to \mathbb{R}^s$, $s \in \mathbb{N}$, X is a column vector function $X = (x_1, x_2, \dots, x_s)$, the column vector function of s variables $F : \mathbb{R}^s \to \mathbb{R}^s$ is

$$F = (f_1, f_2, \dots f_s), \tag{1.2}$$

 $G: \mathbb{R} \to \mathbb{R}^s_+ \times \mathbb{R}^s_+, \mathbb{R}^+ = [0, +\infty)$, and $R: \mathbb{R}^{s+1} \to \mathbb{R}^s \times \mathbb{R}^s$ are matrix functions: G is diagonal with g_i on the diagonal, and $r_{ij}(t, \tau)$ are entries of R. We consider the case when the s-dimensional domain H(t) is $H(t) = [h_1(t), t] \times [h_2(t), t] \times \cdots \times [h_s(t), t], h_j(t) \leq t$, and the volume integral can be iterated, where system (1.1) can be rewritten as a collection of s equations for $i = 1, 2, \ldots, s$,

$$\frac{dx_i}{dt} = g_i(t) \left[\int_{h_{i1}(t)}^t d_{\tau_1} r_{i1}(t,\tau_1) \cdots \int_{h_{is}(t)}^t d_{\tau_s} r_{is}(t,\tau_s) f_i(x_1(\tau_1), x_2(\tau_2), \dots, x_s(\tau_s)) - x_i(t) \right].$$
(1.3)

In particular, let $h_{ij}(t) \leq t, i, j = 1, ..., s$ be measurable functions, and r_{ij} be step functions taking the value of one on half-open intervals $(h_{ij}(t), +\infty)$:

$$r_{ij}(t,\zeta) = \chi_{(h_{ij}(t),\infty)}(\zeta), \quad \chi_J(\zeta) := \begin{cases} 1, & \zeta \in J, \\ 0, & \zeta \notin J. \end{cases}$$

Then, (1.3) has the form

$$\frac{dx_i}{dt} = g_i(t) \left[f_i(x_1(h_{i1}(t)), x_2(h_{i2}(t)), \dots, x_s(h_{is}(t))) - x_i(t) \right], \quad i = 1, 2, \dots, s.$$
(1.4)

The Hopfield neural network [23]

$$x'_{i}(t) = -b_{i}x_{i}(t) + \sum_{j=1}^{s} c_{ij}\tilde{f}_{j}(x_{j}(t-\tau_{ij})), \quad t \ge 0, \ i = 1, \dots, s$$
(1.5)

is a particular case of (1.4) for

$$g_i(t) \equiv b_i, \quad f_i(x_1, x_2, \dots, x_n) = \frac{1}{b_i} \sum_{j=1}^s c_{ij} \tilde{f}_j(x_j), \quad h_{ij}(t) \equiv t - \tau_{ij}, \quad i, j = 1, \dots, s.$$

In the case of absolutely continuous in ζ functions $r_{ij}(t,\zeta)$,

$$\frac{\partial}{\partial \zeta} r_{ij}(t,\zeta) = k_{ij}(t,\zeta), \quad i,j = 1,\dots,s, \quad K_i(t,\tau_1,\tau_2,\dots,\tau_s) = \prod_{j=1}^s k_{ij}(t,\tau_j),$$

(1.3) becomes

$$\frac{dx_i}{dt} = g_i(t) \left[\int_{H(t)} K_i(t, \tau_1, \dots, \tau_s) f_i(x_1(\tau_1), \dots, x_s(\tau_s)) \ d\tau_1 \dots d\tau_s - x_i(t) \right], \ i = 1, \dots, s.$$
(1.6)

In future, we consider each of the equations separately, as in (1.3). However, due to the length of (1.3), a shorter notation will be used, aligned with (1.1)

$$\frac{dx_i}{dt} = g_i(t) \left[\int_{H(t)} d_\tau R_i(t,\tau) f_i(X(\tau)) - x_i(t) \right], \qquad (1.7)$$

where $\tau = (\tau_1, \ldots, \tau_s)$.

The purpose of the present paper is to explore global asymptotic stability of cooperative systems with distributed delays, which include (1.5) and (1.6) as special cases. In addition to being distributed, the delays can change with time. Distributed delays describe a feasible fact that any interval for delay values has some probability, such models include equations with concentrated (either constant or variable) delays.

Compared to most previous works, main differences are outlined below.

- Distributed delays can, as particular cases, include systems with variable concentrated delays, integral terms (used in most papers on distributed delays), their combinations, and some other models (for example, the Cantor function as a distribution). Moreover, argument deviations can be Lebesgue measurable, not necessarily continuous, functions. This is the reason why methods for continuous delays do not work in this setting.
- Delay distributions can be non-autonomous. If we describe these distributions as a probability that a delay takes a greater than a given value, this corresponds to time-dependent delay. In applications, this allows to consider, for example, seasonal changes

in delay distributions. To some extent, we explore the most general system with a unique positive equilibrium, and justify global stability of this equilibrium, once delays are involved only in those terms which describe cross-influences. The present paper answers the question when delays do not have any destabilizing effect on a non-autonomous system.

• On the other hand, many of the previous papers on distributed delay describe much more complicated dynamics than absolute global stability established in the present paper. For example, delay dependence of stability properties was studied in [13], while possible multistability considered in [6]. However, the study of systems which can be destabilized by large enough delay are not in the framework of the present paper. Here we restrict ourselves to "absolutely stable" systems, where no type or size of a finite delay can destabilize it, as long as the initial conditions belong to the "attraction set".

The plan of the paper is as follows. After some preliminaries and an auxiliary statement in Section 2, we get stability results for systems with a distributed delay in Section 3. These theorems are later applied to particular cases of neural networks and models of population dynamics in Section 4. Finally, the results are discussed, and some open problems and directions of research are outlined in Section 5.

2. Preliminaries

Consider a system with distributed delays (1.1), under the initial condition

$$X(t) = \Phi(t), \ t \le t_0,$$
 (2.1)

where $\Phi(t)$ is a bounded vector function.

Definition 2.1 A vector function X(t) is a solution of system (1.1),(2.1) if it satisfies (1.1) for almost all $t \ge t_0$ and (2.1) for $t \le t_0$.

In particular, (1.3) can be written in matrix form (1.1), where

$$H(t) = (h_1(t), t] \times (h_2(t), t] \times \dots \times (h_s(t), t].$$
(2.2)

Problems (1.3),(2.1) and (1.1),(2.1) will be investigated under some of the following assumptions.

- (a1) There is a domain $D \subset \mathbb{R}^s$ such that all $f_i : D \to \mathbb{R}$ are continuous functions, $i = 1, \ldots, s$.
- (a2) Any scalar delay function $h : \mathbb{R}^+ \to \mathbb{R}$ considered in the paper $(h_{ij} \text{ in particular})$ is Lebesgue measurable, $h(t) \leq t$ and $\lim_{t \to +\infty} h(t) = +\infty$.

(a3) The entries of the matrix $R(t,\tau) = (r_{ij}(t,\tau))_{i,j=1}^{s}$, $r_{ij}(t,\cdot)$, $i,j = 1,\ldots,s$ are left continuous non-decreasing functions for any $t, r_{ij}(\cdot,\tau)$ are locally integrable for any $s, r_{ij}(t,\tau) = 0, \tau \leq h_i(t), r_{ij}(t,t^+) = 1, i, j = 1,\ldots,s$, here all the integrals are understood in the sense

$$\int_{h(t)}^{t} f(\zeta) \, dr(\zeta) = \int_{h(t)}^{t+} f(\zeta) \, dr(\zeta), \quad \int_{h(t)}^{t} f(\zeta) \, d\chi_{[t,+\infty)}(\zeta) = f(t),$$

where $u(t^+)$ is the right-side limit of the function u at point t.

(a4) $G(t) = \text{diag}\{g_1(t), \dots, g_s(t)\}, g_i(t)$ are Lebesgue measurable essentially bounded on \mathbb{R}^+ functions, $g_i(t) \ge 0, i = 1, \dots, s, \int_0^{+\infty} g_i(s) \, ds = +\infty, i = 1, \dots, s.$

(a5) $\Phi: (-\infty, 0] \to \mathbb{R}^s$ is a continuous bounded vector function.

Examples of (1.3) include a system with several concentrated delays

$$\frac{dx_i}{dt} = g_i(t) \left[\sum_{j=1}^{n_i} \alpha_{ij} f_i \left(x_1(h_{i1j}(t)), \dots, x_s(h_{isj}(t)) \right) - x_i(t) \right],$$
(2.3)

where h_{ikj} satisfy (a2), $\sum_{j=1}^{n_i} \alpha_{ij} = 1, i = 1, ..., s$, as well as a system of integro-differential equations

$$\frac{dx_i}{dt} = g_i(t) \left[\sum_{j=1}^s \int_{h_{ij}(t)}^t K_{ij}(t,\tau) f_i(x_1(\tau),\dots,x_s(\tau)) \ d\tau - x_i(t) \right],$$
(2.4)

with h_{ij} satisfying (a2),

$$\sum_{j=1}^{s} \int_{h_{ij}(t)}^{t} K_{ij}(t,\tau) \ d\tau \equiv 1, \quad K_{ij}(t,\tau) \ge 0, \quad i,j = 1,\dots,s.$$
(2.5)

Definition 2.2 (see [26]) Let $D = (a_1, b_1) \times (a_2, b_2) \times \cdots \times (a_s, b_s)$, $I_n = [a_{1n}, b_{1n}] \times [a_{2n}, b_{2n}] \times \cdots \times [a_{sn}, b_{sn}]$, $F : \mathbb{R}^s \to \mathbb{R}^s$. An equilibrium $z_* \in D$ is a strong attractor in D of the difference system

$$X(n+1) = F(X(n)), \quad n = 0, 1, \dots$$
(2.6)

if there exists a sequence of sets $\{I_n\}, n = 0, 1, \dots$, such that

$$Int(I_0) = D, \ F(I_n) \subset I_{n+1} \subset Int(I_n), \ n = 0, 1, \dots, \ \bigcap_{n=1}^{+\infty} I_n = z_*.$$
(2.7)

Note that, once z_* is a strong attractor of F in D, it is unique, moreover, there are no other equilibrium points of F in D.

Lemma 2.3 Let F be a continuous function, where F is defined in (1.2), and $F(I_k) \subset I_{k+1} \subset \text{Int}(I_k), F(I_{k+1}) \subset I_{k+2} \subset \text{Int}(I_{k+1}), \text{ for some } k \in \mathbb{N}, \text{ where } I_k = [a_{1k}, b_{1k}] \times [a_{2k}, b_{2k}] \times \cdots \times [a_{sk}, b_{sk}].$

Then there exist $\overline{J} = [\overline{c}_1, \overline{d}_1] \times [\overline{c}_2, \overline{d}_2] \times \cdots \times [\overline{c}_s, \overline{d}_s]$ and $\underline{J} = [c_1, d_1] \times [c_2, d_2] \times \cdots \times [c_s, d_s]$ such that $\overline{J} \subset \operatorname{Int}(I_k), I_{k+1} \subset \operatorname{Int}\overline{J}, I_{k+2} \subset \operatorname{Int}\underline{J}, \underline{J} \subset \operatorname{Int}(I_{k+1})$ and $F(\overline{J}) \subset \underline{J}$.

Proof. For simplicity, we choose $c_i = 0.5(a_{ik+1} + a_{ik+2})$, $d_i = 0.5(b_{ik+1} + b_{ik+2})$, then $I_{k+2} \subset \operatorname{Int} \underline{J}$ and $\underline{J} \subset \operatorname{Int}(I_{k+1})$ are obviously satisfied.

Next, introduce a family $\{J_{\alpha}\}$ of compact subsets of the interior of I_k as

$$J_{\alpha} = [\alpha a_{1k} + (1 - \alpha)a_{1k+1}] \times [\alpha a_{2k} + (1 - \alpha)a_{2k+1}] \times \dots \times [\alpha a_{sk} + (1 - \alpha)a_{sk+1}], \quad \alpha \in [0, 1]$$

and notice that for $\alpha = 0$, $J_0 = I_{k+1}$, $F(J_0) \subset I_{k+2}$, and $I_{k+2} \subset \text{Int} \underline{J}$. Thus there exists $\alpha_0 \in [0, 1]$ such that

$$\alpha_0 = \inf \left\{ \alpha \in [0, 1] : F(J_\alpha) \subset \underline{J} \right\},\$$

as the set in the right-hand side is non-empty. If $\alpha_0 > 0$, we choose $\alpha = \min\{\alpha_0, \frac{1}{2}\}$ (to avoid $\alpha = 1$) and denote $\overline{J} = J_{\alpha}$. Then $\overline{J} \subset \operatorname{Int}(I_k)$, $I_{k+1} \subset \operatorname{Int}\overline{J}$, $I_{k+2} \subset \operatorname{Int}\underline{J}$, $\underline{J} \subset \operatorname{Int}(I_{k+1})$ and $F(\overline{J}) \subset \underline{J}$, and the proof is complete.

It remains to exclude the case $\alpha_0 = 0$. If for any positive α , $F(J_{\alpha}) \not\subset \underline{J}$, we choose a sequence $\alpha_n = \frac{1}{n}$; by our assumption, there is a sequence of points $z_n \in J_{1/n}$ such that $F(z_n) \not\in \underline{J}$. By definition, all $z_n \in I_k$ which is a compact set, thus there is a subsequence convergent to some \overline{z} , and $F(\overline{z})$ does not belong to the interior of \underline{J} . However, as $\alpha_n \to 0$, this limit point \overline{z} belongs to $J_0 = I_{k+1}$. However, $F(I_{k+1}) \subset I_{k+2}$ and $\operatorname{Int}(\underline{J}) \subset I_{k+2}$, thus $F(\overline{z}) \in \operatorname{Int}(\underline{J})$, which is a contradiction. Thus there exists $\alpha_0 > 0$, and the proof is complete.

3. Main Results

Now, we are in a position to prove the main statement of the paper.

Theorem 3.1 Let $D = (a_1, b_1) \times (a_2, b_2) \times \cdots \times (a_s, b_s)$, $F : \mathbb{R}^s \to \mathbb{R}^s$. Suppose (a1)-(a5) hold, and z_* is a strong attractor of F in D, with $I_n = [a_{1n}, b_{1n}] \times [a_{2n}, b_{2n}] \times \cdots \times [a_{sn}, b_{sn}]$, $n \in \mathbb{N}_0 = \{0\} \cup \mathbb{N}$.

Then for any initial function such that

$$\Phi \in C((-\infty, t_0), D), \tag{3.1}$$

the solution of (1.1),(2.1) satisfies $\lim_{n \to +\infty} X(t) = z_*$.

Proof. We prove that for any initial function satisfying (3.1), first, $X(t) \in Int(I_0)$ for any $t \ge t_0$ and, in addition, there is $t_1 \ge t_0$ such that $X(t_1) \in Int(I_1)$. Moreover, $X(t) \in Int(I_1)$ for any $t \ge t_1$.

Assume that with (3.1) satisfied, there is the first point t^* such that $X(t^*) \notin \text{Int}(I_0)$, i.e. $X(t^*)$ is on the boundary ∂I_0 . Then there exists $i \in \{1, \ldots, s\}$ such that either $x_i(t^*) = a_{i0}$ or $x_i(t^*) = b_{i0}$, and $F(X(t)) \in I_1$, $f_i(X(t)) > a_{i1}$, $t \in [t_0, t^*)$. In the former case, due to continuity of x_i , there is a $t_0^* \in [t_0, t^*)$ such that $x_i(t) < 0.5(a_{i0} + a_{i1})$ for $[t_0^*, t^*)$. Hence, using the notation of (1.7), we get

$$\begin{aligned} x_i(t^*) &= x_i(t_0^*) + \int_{t_0^*}^{t^*} g_i(t) \left[\int_{H(t)} d_\tau R_i(t,\tau) f_i(X(\tau)) - x_i(t) \right] dt \\ &> a_{i0} + \int_{t_0^*}^{t^*} g_i(t) \left[a_{i1} \int_{H(t)} d_\tau R_i(t,\tau) - \frac{a_{i0} + a_{i1}}{2} \right] \\ &> a_{i0} + \int_{t_0^*}^{t^*} g_i(t) \frac{a_{i1} - a_{i0}}{2} dt > a_{i0}, \end{aligned}$$

which contradicts to the assumption $x_i(t^*) = a_{i0}$.

Similarly, in the latter case, assuming that $x_i(t^*) = b_{i0}$ and $x_i(t) > 0.5(b_{i1} + b_{i0})$ for $t \in [t_1^*, t^*]$, we obtain

$$\begin{aligned} x_i(t^*) &= x_i(t_1^*) + \int_{t_1^*}^{t^*} g_i(t) \left[\int_{H(t)} d_\tau R_i(t,\tau) f_i(X(\tau)) - x_i(t) \right] dt \\ &< b_{i0} + \int_{t_1^*}^{t^*} g_i(t) \left[b_{i1} \int_{H(t)} d_\tau R_i(t,\tau) - \frac{b_{i1} + b_{i0}}{2} \right] \\ &< b_{i0} + \int_{t_1^*}^{t^*} g_i(t) \frac{b_{i0} - b_{i1}}{2} dt < b_{i0}, \end{aligned}$$

again leading to a contradiction. Thus, $X(t) \in Int(I_0)$ for any $t \ge t_0$.

Note that by Lemma 2.3, there exist $\overline{J} = [\overline{c}_1, \overline{d}_1] \times [\overline{c}_2, \overline{d}_2] \times \cdots \times [\overline{c}_s, \overline{d}_s]$ and $\underline{J} = [c_1, d_1] \times [c_2, d_2] \times \cdots \times [c_s, d_s]$ such that $\overline{J} \subset \operatorname{Int}(I_0), I_1 \subset \operatorname{Int}\overline{J}, I_2 \subset \operatorname{Int}\underline{J}, \underline{J} \subset \operatorname{Int}(I_1)$ and $F(\overline{J}) \subset \underline{J}$.

The proof that there exists $t_1 > t_0$, such that $X(t) \in \text{Int}(I_1)$ for $t \ge t_1$ will consist of two parts. First, we prove that there is a \overline{t} such that $X(\overline{t}) \in \text{Int}(\overline{J})$; moreover, $X(t) \in \text{Int}(\overline{J})$ for any $t \ge \overline{t}$. Second, we find $t_1 \ge \overline{t}$ for which $X(t_1) \in \text{Int}(I_1)$, and also justify that $X(t) \in \text{Int}(I_1), t \ge t_1$.

Since $I_1 \subset \text{Int}\overline{J}$, we have $\overline{c}_i < a_{i1} < b_{i1} < \overline{d}_i$. Denote

$$\delta := \min\left\{\min_{1 \le i \le s} \left(a_{i1} - \bar{c}_i\right), \min_{1 \le i \le s} \left(\bar{d}_i - b_{i1}\right)\right\} > 0,$$
(3.2)

which is positive as a minimum of 2s positive values.

For any $X(t) \in \text{Int}(I_0)$ we have $F(X(t)) \in I_1$. Assume that this does not hold for some i. Let t be such that the ith component of X(t) satisfies $x_i \leq \bar{c}_i$. We prove that there is a moment of time t_{1i} such that $x_i(t_{1i}) > \bar{c}_i$.

We recall that

$$a_{i0} < \bar{c}_i < a_{i1} < c_i < a_{i2} < b_{i2} < d_i < b_{i1} < \bar{d}_i < b_{i0}, \quad i = 1, \dots, s.$$
 (3.3)

As long as $x_i(t) \leq \bar{c}_i$, we have, by (3.2),

$$\begin{aligned} x_i'(t) &= g_i(t) \left[\int_{H(t)} d_\tau R_i(t,\tau) f_i(X(\tau)) - x_i(t) \right] \\ &\geq g_i(t) \left[\int_{H(t)} a_{i1} d_\tau R_i(t,\tau) - \bar{c}_i \right] = g_i(t) (a_{i1} - \bar{c}_i) \geq \delta g_i(t). \end{aligned}$$

By (a4), the integral of the positive right-hand side diverges, thus there is t_{i1} such that $x_i(t_{i1}) > \overline{c}_i$. Since there is $t < t_{i1}$ such that $x_i(t_{i1}) \leq \overline{c}_i < a_{i1}$, without loss of generality we can assume $x_i(t) < a_{i1}$, $t < t_{i1}$. Next, note that $x_i(t) > \overline{c}_i$ for any $t \geq t_{i1}$. In fact, assuming the contrary that $t^* > t_{i1}$ is the smallest value exceeding t_{i1} at which $x(t^*) = \overline{c}_i$, we get $x_i(t) \in [\overline{c}_i, a_{i1}]$ for $t \in [t_{i1}, t^*]$. Recall that $f_i(X(t)) \geq a_{i1}$, $t \in [t_{i1}, t^*]$, therefore

$$x_{i}(t^{*}) = x(t_{i1}) + \int_{t_{i1}}^{t^{*}} g_{i}(\zeta) \left[\int_{H(\zeta)} d_{\tau} R_{i}(\zeta, \tau) f_{i}(X(\tau)) - x_{i}(\zeta) \right] d\zeta$$

$$\geq x(t_{i1}) + \int_{t_{i1}}^{t^{*}} g_{i}(\zeta) \left[a_{i1} - a_{i1} \right] d\zeta = x(t_{i1}).$$

This contradicts to our assumption that $x(t^*) = \bar{c}_i < x(t_{i1})$.

Similarly, for $x_i(t) \ge \bar{d}_i$,

$$\begin{aligned} x'_{i}(t) &= g_{i}(t) \left[\int_{H(t)} d_{\tau} R_{i}(t,\tau) f_{i}(X(\tau)) - x_{i}(t) \right] \\ &\leq g_{i}(t) \left[\int_{H(t)} b_{i1} d_{\tau} R_{i}(t,\tau) - \bar{d}_{i} \right] = g_{i}(t) (b_{i1} - \bar{d}_{i}) \leq -\delta g_{i}(t). \end{aligned}$$

Hence there is t_{i2} such that $x_i(t_{i2}) < \bar{d}_i$. Again, we justify that $x_i(t) < \bar{d}_i$ for any $t \ge t_{i2}$. Choosing

$$\bar{t} = \max\left\{\max_{1 \le i \le s} t_{i1}, \max_{1 \le i \le s} t_{i2}\right\},\,$$

we conclude that $X(t) \in Int(\overline{J})$ for any $t \ge \overline{t}$.

By (a2), there is \bar{t}_1 such that $h_i(t) \geq \bar{t}$ for any $t \geq \bar{t}_1$, $i = 1, \ldots, s$. Note that $F(\bar{J}) \subset \underline{J}$ and $\underline{J} \subset Int(I_1)$. Define

$$\delta_1 := \min\left\{\min_{1 \le i \le s} \left(c_i - a_{i1}\right), \min_{1 \le i \le s} \left(b_{i1} - d_i\right)\right\},\tag{3.4}$$

which is positive by (3.3). We have for $t \geq \bar{t}_1$,

$$x'_{i}(t) \ge g_{i}(t) \left[\int_{H(t)} c_{i} d_{\tau} R_{i}(t,\tau) - a_{i1} \right] = g_{i}(t)(c_{i} - a_{i1}) \ge \delta_{1} g_{i}(t)$$

for any $x_i(t) \leq a_{i1}$ and conclude by (a4) that there is t_{i3} such that $x_i(t_{i3}) \in (a_{i1}, c_i)$. Moreover, $x_i(t) > a_{i1}$ for any $t \geq t_{i3}$. Assuming the contrary that $x_i(t) \in (a_{i1}, x_i(t_{i3})]$ for $t \in [t_{i3}, t_2^*)$ and $x_i(t_2^*) = a_{i1}$, we get for $t \in [t_{i3}, t_2^*)$, $f_i(X(t)) \geq c_i$

$$x_{i}'(t) \ge g_{i}(t) \left[\int_{H(t)} c_{i} d_{\tau} R_{i}(t,\tau) - x_{i}(t_{i3}) \right] = g_{i}(t)(c_{i} - x_{i}(t_{i3})) \ge g_{i}(t)(c_{i} - a_{i}) \ge \delta_{1}g_{i}(t) > 0,$$

which contradicts to the assumption that $x_i(t_2^*) = a_{i1} < x_i(t_{i3})$.

If $x_i(t) \ge b_{i1}$,

$$x'_{i}(t) \leq g_{i}(t) \left[\int_{H(t)} d_{i} \ d_{\tau} R_{i}(t,\tau) - b_{i1} \right] = r(t)(d_{i} - b_{i1}) \leq -\delta g_{i}(t).$$

Thus $x_i(t_{i4}) < b_{i1}$ for some t_{i4} , and similarly we get $x_i(t) < b_{i1}$ for any $t \ge t_{i4}$. Then, for $t \ge t_1$, where

$$t_1 = \max\left\{\max_{1 \le i \le s} t_{i3}, \max_{1 \le i \le s} t_{i4}\right\},\,$$

we have $x_i(t) \in (a_{i1}, b_{i1}), i = 1, ..., s$, or $X(t) \in Int(I_1), t \ge t_1$.

Since $x_i(t) \in (a_{i1}, b_{i1}), i = 1, ..., s, t \ge t_1$, we proceed to the next induction step from I_1 to I_2 .

By (a2), there exists $t_1^h > t_0$ such that $h(t) > t_1$ for any $t > t_1^h$. Then, we have an initial value problem with all initial values in $Int(I_1)$ and complete the induction step similarly, justifying that there is $t_2 > t_1$ such that $X(t) \in I_2$ for $t \ge t_2$, and also $h(t) > t_2$ for any $t > t_2^*$. Proceeding in the same manner from n to n+1, we prove that there is an increasing sequence of t_n such that $X(t) \in I_n$ for $t \ge t_n$. Since the intersection of I_n is z_* , this implies $\lim_{t \to +\infty} X(t) = z_*$.

Consider system (1.3) or (1.1) under the assumptions (a1)-(a5), as well as an additional assumption

(a6) $z_* = (x_1^*, \ldots, x_j^*, \ldots, x_s^*)$ is the only equilibrium in the domain D, and there exist $L_{ij} \ge 0$ such that, for any $(x_1, \ldots, x_s) \in D$, a.e.

$$\left| f_i(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_s) - f_i(x_1, \dots, x_{j-1}, x_j^*, x_{j+1}, \dots, x_s) \right|$$

$$\leq L_{ij} \left| x_j - x_j^* \right|, \quad i, j = 1, \dots, s, \quad L = (L_{ij})_{i,j=1}^s.$$

$$(3.5)$$

Note that (3.5) in (a6) is satisfied if f_i is globally Lipschitz, i.e. for any $(x_1, \ldots, x_j, \ldots, x_s) \in D$ and $(x_1, \ldots, y_j, \ldots, x_s) \in D$, a.e.,

$$|f_i(x_1,\ldots,x_{j-1},x_j,x_{j+1},\ldots,x_s) - f_i(x_1,\ldots,x_{j-1},y_j,x_{j+1},\ldots,x_s)| \le L_{ij} |x_j - y_j|$$

for i, j = 1, ..., s. In particular, if f_i are a.e. differentiable and

$$\left|\frac{\partial f_i}{\partial x_j}\right| \le L_{ij} \text{ a.e. }, \quad i, j = 1, \dots, s,$$

condition (a6) is satisfied.

We recall that a matrix $A = (a_{ij})_{i,j=1}^s$ is nonnegative if $a_{ij} \ge 0$ and positive if $a_{ij} > 0$, $i, j = 1, \ldots, s$. Let ||X|| be an arbitrary fixed norm of a column vector in \mathbb{R}^s , and ||A|| be the induced matrix norm. The classical definition of an M-matrix will be used. Following [11], we say that $A = (a_{ij})_{i,j=1}^s$ is a (non-singular) M-matrix if $a_{ij} \le 0$ for $i \ne j$ and A^{-1} is positive. By I we denote an $s \times s$ identity matrix.

There are many equivalent definitions of M-matrices, see [11] and also [9, Lemma 2.3].

Lemma 3.2 [11, p. 137,142, Exercise 2.9 of Chapter 6] A is an M-matrix if and only if $a_{ij} \leq 0, i \neq j$ and there exist positive numbers $\xi_i, i = 1, ..., s$ such that

$$\xi_i a_{ii} > \sum_{j \neq i} \xi_j |a_{ij}|, \ i = 1, \dots, s$$

Lemma 3.3 Let (a6) be satisfied, L be defined in (3.5), and I - L be an M-matrix. Then, there exist $\{a_{in}\}_{n=1}^{+\infty}$ and $\{b_{in}\}_{n=1}^{+\infty}$, $i = 1, \ldots, s$ such that

$$a_{in} < a_{in+1} < x_i^* < b_{in+1} < b_{in}, \quad n \in \mathbb{N}, \quad i = 1, \dots, s,$$

$$\lim_{n \to +\infty} a_{in} = \lim_{n \to +\infty} b_{in} = x_i^*, \quad i = 1, \dots, s,$$

and the domains

$$I_n = [a_{1n}, b_{1n}] \times [a_{2n}, b_{2n}] \cdots \times [a_{sn}, b_{sn}]$$

satisfy

$$I_1 \subset D, \quad F(I_n) \subset I_{n+1}, \quad n \in \mathbb{N}.$$

Proof. By definition of the nonnegative matrix L, all off-diagonal entries of I - L are non-positive. By Lemma 3.2, we have a finite set of $\xi_i > 0$, $i = 1, \ldots, s$ such that

$$\xi_i (1 - L_{ii}) > \sum_{j \neq i} \xi_j L_{ij}, \quad i = 1, \dots, s,$$

or

$$\alpha_i := \sum_{j \neq i} \frac{\xi_j}{\xi_i} L_{ij} + L_{ii} < 1, \quad i = 1, \dots, s.$$

Denote

$$\alpha = \max_{1 \le i \le s} \alpha_i \in (0, 1).$$
(3.6)

Choose for some c > 0,

$$a_{i1} = x_i^* - c\xi_i, \quad b_{i1} = x_i^* + c\xi_i, \quad i = 1, \dots, s$$

such that

$$I_1 = [a_{11}, b_{11}] \times [a_{21}, b_{21}] \cdots \times [a_{s1}, b_{s1}] \subset D$$

In particular, if

$$D = (a_{10}, b_{10}) \times (a_{20}, b_{20}) \times \dots \times (a_{s0}, b_{s0}),$$

we can take any positive c satisfying

$$c \le \min\left\{\min_{1\le j\le s} \frac{b_{j0} - x_j^*}{\xi_j}, \min_{1\le j\le s} \frac{x_j^* - a_{j0}}{\xi_j}\right\}.$$

We have

$$I_1 = [x_1^* - c\xi_1, x_1^* + c\xi_1] \times [x_2^* - c\xi_2, x_2^* + c\xi_2] \times \dots \times [x_s^* - c\xi_s, x_s^* + c\xi_s]$$
(3.7)

implying $|x_j - x_j^*| < c\xi_j$, j = 1, ..., s for $X \in I_1$ and $f_i(x_1^*, x_2^*, ..., x_s^*) = x_i^*$. By (3.5), once $X \in I_1$,

$$\begin{aligned} |f_i(X) - x_i^*| &\leq |f_i(x_1, x_2, \dots, x_{s-1}, x_s) - f_i(x_1^*, x_2, \dots, x_{s-1}, x_s)| \\ &+ |f_i(x_1^*, x_2, \dots, x_{s-1}, x_s) - f_i(x_1^*, x_2^*, \dots, x_{s-1}, x_s)| \\ &+ \dots + |f_i(x_1^*, x_2^*, \dots, x_{s-1}^*, x_s) - f_i(x_1^*, x_2^*, \dots, x_{s-1}^*, x_s^*)| \\ &\leq L_{ii} |x_i - x_i^*| + \sum_{j \neq i} L_{ij} |x_j - x_j^*| \\ &\leq L_{ii} c\xi_i + \sum_{j \neq i} L_{ij} c\xi_j \leq \alpha c\xi_i, \end{aligned}$$

where $\alpha \in (0, 1)$ is denoted in (3.6).

Recall (3.7) and denote for $n \in \mathbb{N}$,

$$I_{n+1} = [x_1^* - \alpha^n c\xi_1, x_1^* + \alpha^n c\xi_1] \times \dots \times [x_s^* - \alpha^n c\xi_s, x_s^* + \alpha^n c\xi_s].$$
(3.8)

We have justified $F(I_1) \subset I_2$, with I_2 defined in (3.8). Now let $X \in I_n$. Then,

$$\begin{aligned} |f_i(X) - x_i^*| &\leq |f_i(x_1, x_2, \dots, x_{s-1}, x_s) - f_i(x_1^*, x_2, \dots, x_{s-1}, x_s)| \\ &+ |f_i(x_1^*, x_2, \dots, x_{s-1}, x_s) - f_i(x_1^*, x_2^*, \dots, x_{s-1}, x_s)| \\ &+ \dots + |f_i(x_1^*, x_2^*, \dots, x_{s-1}^*, x_s) - f_i(x_1^*, x_2^*, \dots, x_{s-1}^*, x_s^*)| \\ &\leq L_{ii} |x_i - x_i^*| + \sum_{j \neq i} L_{ij} |x_j - x_j^*| \\ &\leq L_{ii} \alpha^{n-1} c\xi_i + \sum_{j \neq i} L_{ij} \alpha^{n-1} c\xi_j \leq \alpha \alpha^{n-1} c\xi_i = \alpha^n c\xi_i, \end{aligned}$$

so $F(X) \in I_{n+1}$, where I_{n+1} is defined in (3.8). F is an α -contraction, $\alpha \in (0, 1)$. Since

$$a_{in} = x_i^* - \alpha^{n-1} c \xi_i, \quad b_{in} = x_i^* + \alpha^{n-1} c \xi_i, \quad i = 1, \dots, s$$

we get $\lim_{n \to +\infty} a_{in} = \lim_{n \to +\infty} b_{in} = x_i^*$, i = 1, ..., s, which concludes the proof of the lemma. \Box

Lemma 3.3 and Theorem 3.1 immediately imply the following asymptotic stability result.

Theorem 3.4 Suppose (a1)-(a6) are satisfied. If I - L is an *M*-matrix, where *L* is defined in (3.5), then any solution of (1.3) with $X_0 \in D$ converges to z_* .

Proof. Let I - L be an *M*-matrix. By Lemma 3.3 and Definition 2.2, z_* is a strong attractor in *D* of difference system (2.6). Thus, the conditions of Theorem 3.1 are satisfied, and therefore any solution of (1.3),(2.1) with (3.1) being fulfilled converges to z_* .

4. Applications and Examples

Consider a particular case of s = 2. System (1.1) includes the model with variable delays

$$\frac{dx}{dt} = g_1(t) \left[f_1(y(h_1(t))) - x(t) \right],$$

$$\frac{dy}{dt} = g_2(t) \left[f_2(x(h_2(t))) - y(t) \right], \quad t \ge 0$$
(4.1)

and the integro-differential system

$$\frac{dx}{dt} = g_1(t) \left[\int_{h_1(t)}^t K_1(t,s) f_1(y(s)) \, ds - x(t) \right],$$

$$\frac{dy}{dt} = g_2(t) \left[\int_{h_2(t)}^t K_2(t,s) f_2(x(s)) \, ds - y(t) \right], \quad t \ge 0,$$
(4.2)

where for both (4.1) and (4.2), the functions h_i and g_i satisfy (a2) and (a4), respectively. For (4.2), in addition, a modification of (a3)

(a3*) $K_i(t,s) : \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{R}^+, i = 1, 2$ are locally integrable functions in both t and s satisfying $\int_{h_i(t)}^t K_i(t,s) \ ds \equiv 1, i = 1, 2$

is assumed to hold.

Further, for both (4.1) and (4.2), the functions f_i should satisfy

(a7) Both $f_1 : \mathbb{R}^+ \to \mathbb{R}^+$ and $f_2 : \mathbb{R}^+ \to \mathbb{R}^+$ are continuous strictly monotone increasing, $f_1(0) = f_2(0) = 0$ and

$$f_2(x) > f_1^{-1}(x), \ x \in (0, x^*), \ f_2(x) < f_1^{-1}(x), \ x \in (x^*, +\infty), \ y^* = f_1(x^*).$$
 (4.3)

Note that (4.3) implies that (x^*, y^*) is the unique equilibrium of systems (4.1) and (4.2).

Proposition 4.1 Let h_i , g_i and f_i satisfy (a2),(a4) and (a7), respectively, and, in the case of (4.2), (a3^{*}) hold. Then, all solutions of (4.1) and (4.2) with non-negative non-trivial in both x and y continuous initial conditions converge to (x^*, y^*) .

Proof. Define $D = (0, +\infty) \times (0, +\infty)$, $F = (f_1, f_2)^T$, then $F : D \to D$. Let us construct $I_n = [a_{1n}, b_{1n}] \times [a_{2n}, b_{2n}]$, where

$$a_{11} < a_{12} \cdots < a_{1n} < a_{1n+1} < \dots, \quad a_{21} < a_{22} \cdots < a_{2n} < a_{2n+1} < \dots, \tag{4.4}$$

$$\dots < b_{1n+1} < b_{1n} < \dots b_{12} < b_{11}, \quad \dots < b_{2n+1} < b_{2n} < \dots b_{22} < b_{21}, \tag{4.5}$$

and

$$\lim_{n \to +\infty} a_{1n} = \lim_{n \to +\infty} b_{1n} = x^*, \quad \lim_{n \to +\infty} a_{2n} = \lim_{n \to +\infty} b_{2n} = y^*, \tag{4.6}$$

which would imply that (x^*, y^*) is a strong attractor.

Since f_1 is monotone increasing, so is f_1^{-1} , also both f_2 and f_2^{-1} are monotone increasing. By (4.3) in (a6), $f_2(x) > f_1^{-1}(x)$ for $x \in (0, x^*)$. Denote $y = f_2(x)$, $x = f_2^{-1}(y)$. The function f_1 is also monotone increasing, thus, taking f_2 of both sides, we get

$$f_1(f_2(x)) > f_1(f_1^{-1}(x)) = x$$
, or $f_1(y) > f_2^{-1}(y)$, $y \in (0, y^*)$.

Similarly, considering $x > x^*$, or $y = f_2(x) > y^*$, we get $f_1(y) < f_2^{-1}(y)$ for $y \in (y^*, +\infty)$. Thus

$$f_1(y) > f_2^{-1}(y), y \in (0, y^*), f_2(y) < f_1^{-1}(y), y \in (y^*, +\infty).$$

Next, choose arbitrary initial left bounds $a_{11} \in (0, x^*)$ and $b_{11} \in (x^*, +\infty)$. For the left bound define $a_{21} = f_2(a_{11}), a_{12} = f_1(a_{21}), a_{22} = f_2(a_{12})$. By (4.3),

$$0 < f_1^{-1}(x) < f_2(x) < f_2(x^*) = y^*, \ x \in (0, x^*),$$

hence $f_2 : (0, x^*) \to (0, y^*)$. Recall that f_2 is monotone and $a_{11} \in (0, x^*)$, therefore $a_{21} = f_1(a_{11}) \in (0, y^*)$. In addition, for $x \in (0, x^*)$, (4.3) implies $f_1 : (0, y^*) \to (0, x^*)$ for monotone increasing f_1 . Therefore $a_{12} \in (0, x^*)$ and $a_{22} = f_2(a_{12}) \in (0, y^*)$. We have $a_{21}, a_{22} \in (0, y^*), a_{12} \in (0, x^*)$. Also,

$$a_{12} = f_1(a_{21}) = f_1(f_2(a_{11})) > f_1(f_1^{-1}(a_{11})) = a_{11},$$

$$a_{22} = f_2(a_{12}) = f_2(f_1(a_{21})) > f_2(f_2^{-1}(a_{21})) = a_{21}.$$

For an induction step, take

$$a_{1n+1} = f_1(a_{2n}), \quad a_{2n+1} = f_2(a_{1n+1}).$$
 (4.7)

From $a_{1n} \in (0, x^*)$, $a_{2n} \in (0, y^*)$ and monotonicity of f_1, f_2 we get $a_{1n+1} \in (0, x^*)$, $a_{2n+1} \in (0, y^*)$, as well as

$$a_{1n+1} = f_1(a_{2n}) = f_1(f_2(a_{1n})) > f_1(f_1^{-1}(a_{1n})) = a_{1n},$$

$$a_{22} = f_2(a_{1n+1}) = f_2(f_1(a_{2n})) > f_2(f_2^{-1}(a_{2n})) = a_{2n}.$$

Thus, (4.4) holds, and we have two monotone increasing sequences $\{a_{1n}\}$ and $\{a_{2n}\}$ bounded by x^* and y^* , respectively, from above. Hence both sequences have limits $\lim_{n \to +\infty} a_{1n} = d_1 \in (0, x^*]$, $\lim_{n \to +\infty} a_{2n} = d_2 \in (0, y^*]$. By (4.7) and continuity of $f_1, f_2, d_1 = f_1(d_2), d_2 = f_2(d_1)$, which implies $d_1 = x^*, d_2 = y^*$.

For the right bound we use

$$y^* = f_2(x^*) < f_2(x) < f_1^{-1}(x), \ x \in (x^*, +\infty),$$

 $f_2: (x^*, +\infty) \to (y^*, +\infty) \text{ and } f_2: (y^*, +\infty) \to (x^*, +\infty).$ Therefore, we get bounds for $b_{21} = f_2(b_{11}) \in (y^*, +\infty), \ b_{12} = f_1(a_{21}) \in (x^*, +\infty), \ b_{22} = f_1(b_{12}) \in (y^*, +\infty).$

The sequences of $b_{1n} \in (x^*, +\infty)$, $b_{2n} \in (y^*, +\infty)$ satisfying (4.5) are constructed similarly

$$b_{1n+1} = f_1(b_{2n}), \quad b_{2n+1} = f_2(b_{1n+1})$$

and the proof of (4.5) follows the same steps, as well as $\lim_{n \to +\infty} b_{1n} = x^*$, $\lim_{n \to +\infty} b_{2n} = y^*$. Therefore, (4.6) is satisfied, and (x^*, y^*) is a strong attractor. Thus, all the conditions of Theorem 3.1 are satisfied. The application of Theorem 3.1 concludes the proof. \Box

The statement of Proposition 4.1 is the main result of [7], and a two-dimensional cooperative system described in [7] is a particular case of the system considered in the present paper.

Example 4.2 Consider a particular case of (2.3)

$$\begin{aligned}
x' &= g_1(t) \left[\sqrt{y(h_1(t))} - x(t) \right], \\
y' &= g_2(t) \left[\sqrt{x(h_2(t))} - y(t) \right], \quad t \ge 0.
\end{aligned}$$
(4.8)

For (4.8), $f_1(y) = \sqrt{y}$, $f_1^{-1}(x) = x^2$, $f_2(x) = \sqrt{x}$, $f_2^{-1}(y) = y^2$, $f_1(1) = 1$, $f_2(1) = 1$ and

$$f_2(x) = \sqrt{x} > f_1^{-1}(x) = x^2, \quad x \in (0,1), \quad f_2(x) = \sqrt{x} < f_1^{-1}(x) = x^2, \quad x \in (1,+\infty).$$

Thus (4.3) holds, and Proposition 4.1 implies that any solution with nonnegative nontrivial initial conditions converges to (1,1).

Example 4.3 For the system

$$\begin{aligned}
x' &= g_1(t) \left[y^2(h_1(t)) - x(t) \right], \\
y' &= g_2(t) \left[\sqrt[4]{x(h_2(t))} - y(t) \right], \quad t \ge 0,
\end{aligned}$$
(4.9)

the functions $f_1(y) = y^2$ and $f_2(x) = \sqrt[4]{x}$ are continuous and monotone increasing on \mathbb{R}_+ , $f_1(1) = 1$, $f_2(1) = 1$. Also,

$$f_2(x) = \sqrt[4]{x} > f_1^{-1}(x) = \sqrt{x}, \quad x \in (0,1), \quad f_2(x) = \sqrt[4]{x} < f_1^{-1}(x) = \sqrt{x}, \quad x \in (1,+\infty).$$

Since (4.3) holds, by Proposition 4.1 any non-negative non-trivial in both x and y solution converges to (1, 1).

Example 4.4 For the BAM neural network without delays in the leakage terms

$$x'_{i} = g_{i}(t) \left[\left(\sum_{j=1}^{s} \alpha_{ij} x_{j}(h_{ij}(t)) \right)^{1/(2k_{i})} - x_{i}(t) \right],$$

where

$$\alpha_{ij} \ge 0, \quad \sum_{j=1}^{s} \alpha_{ij} = 1, \quad k_i \in \mathbb{N}, \quad i, j = 1, \dots s, \quad t \ge 0,$$

Theorem 3.4 implies that the equilibrium (1, 1, ..., 1) attracts all solutions with non-negative non-trivial continuous initial conditions.

Remark 4.5 In [9], a neural system which can be reduced to

$$\dot{x}_i(t) = \alpha_i(t) \left[-x_i(h_i(t)) + \sum_{j=1}^s F_{ij}(t, x_j(h_{ij}(t))) \right], \quad t \ge 0, \quad i = 1, \dots, s$$

was considered. If in the leakage terms $h_i(t) \equiv t$, i = 1, ..., s, the results of [9, Theorem 2.5] coincide with a particular case of Theorem 3.4 when delays are concentrated. Thus, compared to [9, Theorem 2.5], Theorem 3.4 considers more general distributed delays in non-leakage part but assumes a particular case of non-delayed leakage terms. Therefore, the results are independent.

Remark 4.6 According to [26, Theorem 3.1], all solutions of the system

$$\dot{x}_i(t) = -x_i(t) + \sum_{j=1}^s \alpha_{ij}(t) f_j(x_j(t-\tau_{ij})), \quad t > 0, \quad i = 1, \dots, s,$$

where

$$|f_j(u) - f_j(v)| \le L_j |u - v|, \ \forall u, v \in \mathbb{R}, \quad \max_{1 \le i \le s} \sum_{j=1}^s \alpha_{ij} L_j < 1,$$
 (4.10)

converge to the zero equilibrium (0, 0, ..., 0). Following the notation of the present paper, denote by L the matrix with the entries $L_{ij} = \alpha_{ij}L_j$. Thus zero is globally attractive, once a sum of the entries of each column is less than one. Note that Theorem 3.4 states attractivity of the zero equilibrium once the matrix I - L is an M-matrix. For example, let

$$L = \begin{bmatrix} \frac{1}{2} & 2\\ \frac{1}{16} & \frac{1}{2} \end{bmatrix},$$

then (4.10) is not satisfied, since the sum of the entries of the second column exceeds 1, but it is easy to check that I - L is an *M*-matrix by its form and positivity of the inverse matrix

$$I - L = \begin{bmatrix} \frac{1}{2} & -2\\ -\frac{1}{16} & \frac{1}{2} \end{bmatrix}, \quad (I - L)^{-1} = \begin{bmatrix} 4 & 16\\ \frac{1}{2} & 4 \end{bmatrix},$$

therefore Theorem 3.4 implies global attractivity of the zero equilibrium.

Next, consider the Nicholson-type system

$$\frac{dx_i}{dt} = g_i \left[\sum_{j \neq i} a_{ij} x_j + \sum_{k=1}^m \beta_{ik} x_i(\tau_{ik}(t)) e^{-x_i(\tau_{ik}(t))} - x_i(t) \right], \quad i = 1, \dots, s, \quad t \ge 0,$$
(4.11)

where $g_i > 0$, a_{ij} and β_{ik} are non-negative, while

$$\beta_i := \sum_{k=1}^m \beta_{ik} > 0,$$

and for some $\tau > 0$, $t - \tau_{ik}(t) \leq \tau$, $i = 1, \ldots, s$, $k = 1, \ldots, m$. Global attractivity conditions for (4.11) were obtained in [14, 19], see also references therein and [14] for a detailed history outline. A positive equilibrium for this system exists [14] once all the constants

$$\gamma_i := \frac{\beta_i}{1 - \sum_{j \neq i} a_{ij}} \tag{4.12}$$

satisfy $\gamma_i > 1, i = 1, \ldots, s$.

The unique positive equilibrium of (4.11) exists and is globally attractive [14, Theorems 2.5 and 3.3], once

$$1 < \gamma_i < e^2, \quad i = 1, \dots, s.$$
 (4.13)

Note that inequalities (4.13) imply $1 - \sum_{j \neq i} a_{ij} < \beta_i < e^2, i = 1, \dots, s.$

As an application of our results, consider the system

$$\frac{dx_i}{dt} = g_i(t) \left[\sum_{j \neq i} a_{ij} x_j(t) + \int_{h_i(t)}^t \beta_i x_i(\tau) e^{-x_i(\tau)} d_\tau r_i(t,\tau) - x_i(t) \right], \ i = 1, \dots, s, \ t \ge 0,$$
(4.14)

where for the functions g_i, r_i conditions (a2)-(a4) hold, $a_{ij} \ge 0, 1 < \beta_i \le e^2$.

In particular, (4.14) includes the system with several concentrated delays generalizing (4.11)

$$\frac{dx_i}{dt} = g_i(t) \left[\sum_{j \neq i} a_{ij} x_j(t) + \sum_{k=1}^{m_i} \beta_{ik} x_i(\tau_{ik}(t)) e^{-x_i(\tau_{ik}(t))} - x_i(t) \right], \quad i = 1, \dots, s, \quad t \ge 0,$$

where $\beta_i = \sum_{k=1}^{m_i} \beta_{ik} > 0, i = 1, \dots, s, g_i$ satisfy (a4), for τ_{ik} condition (a2) holds.

Assume that $x^* = (x_1^*, \ldots, x_s^*)$ is a unique positive equilibrium of (4.14). In particular, the fact that $\gamma_i > 1$, where γ_i are defined in (4.12), $i = 1, \ldots, s$ guarantees that such an equilibrium exists, similarly to systems with concentrated delays.

Denote

$$\alpha_{i} = \begin{cases} \max\{1 - \ln \beta_{i}, \beta_{i}e^{-2}\}, & 1 < \beta_{i} \le e, \\ \beta_{i}e^{-2}, & e < \beta_{i} < e^{2}. \end{cases}$$
(4.15)

Theorem 4.7 Let (a_2) - (a_4) hold and I - L be an M-matrix, where

$$L = (L_{ij})_{i,j=1}^{s}, \ L_{ij} = \begin{cases} a_{ij}, & j \neq i, \\ \alpha_{i}, & j = i, \end{cases}$$

and α_i are denoted in (4.15). Then all solutions of (4.14) with non-negative non-trivial initial conditions converge to x^* .

Proof. Denote

$$f_i(x_1,\ldots,x_s) = \sum_{j \neq i} a_{ij} x_j + \beta_i x_i e^{-x_i}.$$

To apply Theorem 3.4, we have to estimate the partial derivatives $\left|\frac{\partial f_i}{\partial x_j}\right|$. We have

$$\left|\frac{\partial f_i}{\partial x_j}\right| = \begin{cases} a_{ij}, & j \neq i, \\ \beta_i |1 - x_i| e^{-x_i}, & j = i. \end{cases}$$

The maximum of the function xe^{-x} is attained at x = 1 and equals 1/e. According to [13, Theorem 2.6], any positive solution of the equation

$$\frac{dy_i}{dt} = g_i(t) \left[\int_{h_i(t)}^t \beta_i y_i(\tau) e^{-y_i(\tau)} d_\tau r_i(t,\tau) - y_i(t) \right]$$
(4.16)

with $\beta_i > 1$ satisfying

$$\limsup_{t \to +\infty} y_i(t) \le \sup_{x \in [0, +\infty)} \beta_i x e^{-x} = \frac{\beta_i}{e},$$
$$\liminf_{t \to +\infty} y_i(t) \ge \min_{x \in (1, +\infty)} \beta_i x e^{-x} = x_i^0,$$

where

$$x_i^0 := \begin{cases} \ln \beta_i, & \beta_i \in [1, e), \\ \frac{\beta_i^2}{e} e^{-\frac{\beta_i}{e}}, & \beta_i \in [e, e^2). \end{cases}$$

Next, let x_i be a component of a solution in (4.14). Then, with the same initial conditions as in (4.16), since all components are positive, $x_i(t) \ge x_i^0$. Note that a similar result for concentrated delays was justified in [10, Theorem 2.3]. Hence it is sufficient to estimate $p_i(x_i) := \beta_i |1 - x_i| e^{-x_i}$ only on the interval $[x_i^0, +\infty)$. There are two cases: $x_i^0 \le 1$ corresponding to $\beta_i \in (1, e]$ and $x_i^0 > 1$ for $\beta_i \in (e, e^2)$.

corresponding to $\beta_i \in (1, e]$ and $x_i^0 > 1$ for $\beta_i \in (e, e^2)$. If $x_i^0 \le x_i < 1$ then $p_i = \beta_i (1 - x_i) e^{-x_i}$, $p'_i = -\beta_i (2 - x_i) e^{-x_i} < 0$. Hence $\max_{x_i^0 \le x_i < 1} p_i(x_i) = p_i(x_i^0) = \beta_i (1 - x_i^0) e^{-x_i^0} = 1 - \ln \beta_i.$

If $x_i^0 < 1$, $x_i > 1$ then $p_i = \beta_i (x_i - 1)e^{-x_i}$, $p'_i = \beta_i (2 - x_i)e^{-x_i}$ and $\max_{x_i > 1} p_i(x_i) = p_i(2) = \beta_i e^{-2}$.

If $x_i \ge x_i^0 > 1$ then $\max_{x_i \ge x_i^0 > 1} p_i(x_i) = p_i(2) = \beta_i e^{-2}$.

Theorem 3.4 implies that, since the matrix I-L is an *M*-matrix, x^* is a global attractor, which concludes the proof.

In particular, if $\beta_i > e$, the diagonal entries of I - L are positive, and the matrix is diagonally dominant

$$\beta_i e^{-2} < 1 - \sum_{i \neq j} a_{ij}, \quad i = 1, \dots, s,$$

the unique positive equilibrium of (4.14) is globally asymptotically stable, which generalizes the right inequality in (4.13) to the case of distributed delays and variable growth rates.

Corollary 4.8 Let n = 2, (a2)-(a4) hold and

$$e < \beta_1 < e^2, \ e < \beta_2 < e^2, \ a_{12}a_{21} < (1 - \alpha_1)(1 - \alpha_2),$$
 (4.17)

where α_i are introduced in (4.15). Then the positive equilibrium is globally attractive.

Proof. For n = 2, as $\alpha_i = \beta_i e^{-2}$, we have

$$I - A = \begin{pmatrix} 1 - \alpha_1 & -a_{12} \\ -a_{21} & 1 - \alpha_2 \end{pmatrix} = \begin{pmatrix} 1 - \beta_1 e^{-2} & -a_{12} \\ -a_{21} & 1 - \beta_2 e^{-2} \end{pmatrix}.$$

Thus I - A is an *M*-matrix if $\beta_i e^{-2} < 1$, i = 1, 2 and $a_{12}a_{21} < (1 - \alpha_1)(1 - \alpha_2)$, in particular, when (4.17) holds.

Example 4.9 Consider the system with h_1 , h_2 satisfying (a2), $r_1 > 0$, $r_2 > 0$,

$$\begin{aligned} x'(t) &= r_1 \left[0.5y(t) + 4x(h_1(t))e^{-x(h_1(t))} - x(t) \right], \\ y'(t) &= r_2 \left[0.2x(t) + 5y(h_2(t))e^{-y(h_2(t))} - y(t) \right], \quad t \ge 0. \end{aligned}$$
(4.18)

Obviously $\beta_1 = 4$ and $\beta_2 = 5$ are in (e, e^2) . Also,

$$a_{12}a_{21} = 0.1 < (1 - \beta_1 e^{-2})(1 - \beta_2 e^{-2}) \approx 0.148295$$

so (4.17) is satisfied, and the positive equilibrium is globally attractive.

Note that for n = 2, conditions (4.13) are equivalent to

$$1 - a_{12} < \beta_1 < (1 - a_{12})e^2, \quad 1 - a_{21} < \beta_2 < (1 - a_{21})e^2.$$
(4.19)

The right inequalities in (4.19) can be rewritten as

$$a_{12} < 1 - \beta_1 e^{-2}, \quad a_{21} < 1 - \beta_2 e^{-2},$$

where the first inequality is not satisfied since

$$a_{12} = 0.5 > 1 - \beta_1 e^{-2} \approx 0.45866.$$

Thus Corollary 4.8 establishes global attractivity of the positive equilibrium of (4.18), while (4.19) fails.

5. Discussion

General system (1.1) was motivated by neural networks but another common application is a compartment, or patch model of mathematical biology. For example, (4.14) is a particular type of a compartment model, where x_i is a population size in the *i*th patch, $a_{ij}(t)$ describes the relocation rate from the patch j to patch $i, i \neq j$, and Nicholson's growth rate. Assuming the logistic growth rate, we get for $K_i > 0$ being the carrying capacity of the *i*th patch, a model

$$\frac{dx_i}{dt} = g_i(t) \left[\sum_{j \neq i} a_{ij} x_j(t) + \int_{h_i(t)}^t \beta_i x_i(\tau) \left(1 - \frac{x_i(\tau)}{K_i} \right) d_\tau r_i(t,\tau) - x_i(t) \right], \ i = 1, \dots, s.$$
(5.1)

As possible extension of current research, another compartment model with the Mackey-Glass growth rate

$$\frac{dx_i}{dt} = r_i(t) \left[\sum_{j \neq i} a_{ij} x_j + \int_{h_i(t)}^t \beta_i \frac{x_i(\tau)}{1 + x_i^n(\tau)} \ d_\tau r_i(t,\tau) - x_i(t) \right], i = 1, \dots, s$$
(5.2)

can be explored under usual assumptions. It would be interesting to investigate existence, uniqueness and absolute attractivity of the positive equilibrium, and the dependency of this equilibrium on the parameters, as well as delay-dependent stability.

In addition to Nicholson-type system (4.14) studied in the present paper and proposed (5.1), (5.2), it is possible to consider Ricker-type model, for i = 1, ..., s,

$$\frac{dx_i}{dt} = g_i(t) \left[\int_{h_i(t)}^t \beta_i x_i(\tau) \exp\left\{ K_i - x_i(\tau) - \sum_{j \neq i} a_{ij} x_j(\tau) \right\} d_\tau r_i(t,\tau) - x_i(t) \right].$$
(5.3)

Global attractivity of a positive equilibrium for s = 2 and s = 3 was recently studied in [2], with explicit criteria obtained. It would be interesting to compare sufficient conditions under which the positive equilibrium of (5.3) attracts all positive solutions with these tests. In general, the strong attractivity is a stricter assumption that the fact that all solutions of a system of difference equations converge to a certain solution [27], so it is expected that global attractivity conditions for (5.3) may be more restrictive than the tests in [2].

Let us discuss whether we can replace a sequence of parallelepiped-type domains containing a fixed point z_* by any closed compact sets including z_* . Notice that a compact set on a line mentioned in the definition of a strong attractor in [26] can be reduced to a closed segment such that its interior contains z_* . Recall that every open set \mathbb{R} is a union of at most countable number of open disjoint intervals [29, p. 45, problems 22 and 29]. Hence a closed bounded set is a union of at most countable number of disjoint closed segments (some may consist of one point only). As segments are disjoint, only one of the segments includes z_* . Therefore at each stage we can consider only this segment. The fact that this segment has a non-empty interior, follows from (2.7). Thus, instead of a sequence of compact sets in [26], without loss of generality we can consider $I_n = [a_{1n}, b_{1n}] \times \cdots \times [a_{sn}, b_{sn}]$ as in Definition 2.2. Thus, our definition in fact coincides with the one in [26].

The main result of the present paper is the proof of global attractivity of nonautonomous equations with distributed and finite, not necessarily bounded, delays. One of the natural questions arising will be extension of the present results to equations with infinite, but exponentially decaying memory. Considering delay-dependent attractivity conditions for systems with distributed delays, similarly to the case of "small delays" in [13, 14], is another important question, once a cooperative system is not globally asymptotically stable for any delays.

The results of the present paper are concerned with non-autonomous systems. For relevant autonomous equations with distributed delays, it has been recently proved [3, 12] that, once we replace a distributed delay in an autonomous equation with its expected value, and the resulting delay equation is stable, so is the model with a distributed delay. It is an interesting and challenging problem to extend this result to autonomous systems with distributed delays.

Acknowledgments

The second author was partially supported by the NSERC grant RGPIN/05976-2015.

- A. N. Aliseyko, Lyapunov matrices for neutral time-delay systems with exponential kernel, Systems Control Lett. 131 (2019), 104497, 7 pp.
- [2] E. C. Balreira, S. Elaydi, R. Lus, Global stability of higher dimensional monotone maps, J. Difference Equ. Appl. 23 (2017), 2037-2071.
- [3] S. Bernard, J. Bélair, M.C. Mackey, Sufficient conditions for stability of linear differential equations with distributed delay, *Discrete Contin. Dyn. Syst. Ser.B* 1 (2001), 233-256.
- [4] L. Berezansky and E. Braverman, On nonoscillation and stability for systems of differential equations with a distributed delay, Automatica J. IFAC 48 (2012), 612-618.
- [5] L. Berezansky and E. Braverman, Stability of equations with a distributed delay, monotone production and nonlinear mortality, *Nonlinearity* 26 (2013), 2833-2849.
- [6] L. Berezansky and E. Braverman, On multistability of equations with a distributed delay, monotone production and the Allee effect, J. Math. Anal. Appl. 415 (2014), 873–888.
- [7] L. Berezansky and E. Braverman, On stability of cooperative and hereditary systems with a distributed delay, *Nonlinearity* 28 (2015), 1745–1760.
- [8] L. Berezansky and E. Braverman, On stability of delay equations with positive and negative coefficients with applications, Z. Anal. Anwend. 38 (2019), 157–189.
- [9] L. Berezansky, E. Braverman and L. Idels, New global exponential stability criteria for nonlinear delay differential systems with applications to BAM neural networks, *Appl. Math. Comput.* 243 (2014), 899–910.
- [10] L. Berezansky, L. Idels and L. Troib, Global dynamics of Nicholson-type delay systems with applications, Nonlinear Anal. Real World Appl. 12 (2011), 436-445.

- [11] A. Berman and R. Plemmons, Nonnegative Matrices in the Mathematical Sciences, Classics in Applied Mathematics, SIAM, Philadelphia, 1994.
- [12] S. Bernard and F. Crauste, Optimal linear stability condition for scalar differential equations with distributed delay, *Discrete Contin. Dyn. Syst. B* 20 (2015), 1855–1876.
- [13] E. Braverman and S. Zhukovskiy, Absolute and delay-dependent stability of equations with a distributed delay, *Discrete Contin. Dyn. Syst.* **32** (2012), 2041-2061.
- [14] D. Caetano and T. Faria, Stability and attractivity for Nicholson systems with time-dependent delays, Electron. J. Qual. Theory Differ. Equ. 2017, Paper No. 63, 19 pp.
- [15] S. A. Campbell and I. Ncube, Stability in a scalar differential equation with multiple, distributed time delays, J. Math. Anal. Appl. 450 (2017), 1104–1122.
- [16] Z. Cheng, Y. Wang and J. Cao, Stability and Hopf bifurcation of a neural network model with distributed delays and strong kernel, *Nonlinear Dynam.* 86 (2016), 323-335.
- [17] S. Esteves, E. Gökmen and J. J. Oliveira, Global exponential stability of nonautonomous neural network models with continuous distributed delays, *Appl. Math. Comput.* **219** (2013), 9296–9307.
- [18] T. Faria, Periodic solutions for a non-monotone family of delayed differential equations with applications to Nicholson systems, J. Differential Equations 263 (2017), 509–533.
- [19] T. Faria and G. Röst, Persistence, permanence and global stability for an n-dimensional Nicholson system, J. Dynam. Differential Equations 26 (2014), 723–744.
- [20] V. Y. Glizer, Uniform stabilizability of parameter-dependent systems with state and control delays by smooth-gain controls, J. Optim. Theory Appl. 183 (2019), 50–65.
- [21] S. A. Gourley, R. Liu and Y. Lou, Intra-specific competition and insect larval development: a model with time-dependent delay, Proc. Roy. Soc. Edinburgh Sect. A 147 (2017), 353–369.
- [22] K. Hattaf and N. Yousfi, A class of delayed viral infection models with general incidence rate and adaptive immune response. Int. J. Dyn. Control 4 (2016), no. 3, 254–265.
- [23] J.J. Hopfield, Neural networks with graded response have collective computation properties like those of two-state neurons, Proc. Natl. Acad. Sci. 81 (1984), 3088–3092.
- [24] S. Liu and E. Beretta, Competitive systems with stage structure of distributed-delay type, J. Math. Anal. Appl. 323 (2006), 331–343.
- [25] X. Liu and P. Stechlinski, Hybrid control of impulsive systems with distributed delays, Nonlinear Anal. Hybrid Syst. 11 (2014), 57–70.
- [26] E. Liz and A. Ruiz-Herrera, Attractivity, multistability, and bifurcation in delayed Hopfield's model with non-monotonic feedback, J. Differential Equations 255 (2013), no. 11, 4244–4266.
- [27] E. Liz and A. Ruiz-Herrera, Alfonso Addendum to "Attractivity, multistability, and bifurcation in delayed Hopfield's model with non-monotonic feedback", J. Differential Equations 257 (2014), 1307-1309.
- [28] Y. Muroya and T. Faria, Attractivity of saturated equilibria for Lotka-Volterra systems with infinite delays and feedback controls, *Discrete Contin. Dyn. Syst. Ser. B* 24 (2019), 3089-3114.
- [29] W. Rudin, Principles of Mathematical Analysis, third edition, International Series in Pure and Applied Mathematics, McGraw-Hill Book Co., New York-Auckland-Dsseldorf, 1976.
- [30] O. Solomon and E. Fridman, New stability conditions for systems with distributed delays, Automatica J. IFAC 49 (2013), 3467–3475.
- [31] X. Xu, L. Liu and G. Feng, Semi-global stabilization of linear systems with distributed infinite input delays and actuator saturations. Automatica J. IFAC 107 (2019), 398–405.
- [32] Y. Yang, L. Zou and S. Ruan, Global dynamics of a delayed within-host viral infection model with both virus-to-cell and cell-to-cell transmissions. Math. Biosci. 270 (2015), part B, 183–191.
- [33] Y. Yuan and J. Bélair, Stability and Hopf bifurcation analysis for functional differential equation with distributed delay, SIAM J. Appl. Dyn. Syst. 10 (2011), 551–581.