

# A NOTE ON LIE ALGEBRA COHOMOLOGY

MICHAEL J. LARSEN AND VALERY A. LUNTS

ABSTRACT. Given a finite dimensional Lie algebra  $L$  let  $I$  be the augmentation ideal in the universal enveloping algebra  $U(L)$ . We study the conditions on  $L$  under which the Ext-groups  $\text{Ext}(k, k)$  for the trivial  $L$ -module  $k$  are the same when computed in the category of all  $U(L)$ -modules or in the category of  $I$ -torsion  $U(L)$ -modules. We also prove that the Rees algebra  $\bigoplus_{n \geq 0} I^n$  is Noetherian if and only if  $L$  is nilpotent. An application to cohomology of equivariant sheaves is given.

## 1. INTRODUCTION

Let  $L$  be a finite dimensional Lie algebra over a field  $k$ . Consider the universal enveloping algebra  $U(L)$  with the augmentation ideal  $I \subset U(L)$ . Denote by  $U(L)\text{-mod}$  the category of finitely generated left  $U(L)$ -modules and by  $(U(L)\text{-mod})_I \subset U(L)\text{-mod}$  the Serre subcategory of  $I$ -torsion modules. We have the obvious functor

$$(1.1) \quad \Phi_L : D^b((U(L)\text{-mod})_I) \rightarrow D^b_I(U(L)\text{-mod})$$

where  $D^b_I(U(L)\text{-mod}) \subset D^b(U(L)\text{-mod})$  is the full triangulated subcategory consisting of complexes with  $I$ -torsion cohomology. In this paper we study the question:

**Question.** When is  $\Phi_L$  an equivalence?

The functor  $\Phi_L$  being an equivalence means that the Ext-groups  $\text{Ext}^i(k, k)$  for the trivial  $L$ -module  $k$  are the same in the categories  $U(L)\text{-mod}$  and  $(U(L)\text{-mod})_I$ .

We answer this question in Theorem 1.1 below.

Define inductively the decreasing sequence of ideals in  $L$ :

$$L_1 = L, \quad L_n = [L, L_{n-1}]$$

and put  $L_\infty = \bigcap_n L_n$ . This is an ideal in  $L$  such that the quotient Lie algebra  $L_{\text{nil}} := L/L_\infty$  is nilpotent. We have  $L_\infty = 0$  if and only if  $L$  is nilpotent.

For each  $i$  the cohomology  $H^i(L_\infty, k)$  is naturally an  $L_{\text{nil}}$ -module. Denote by  $H^{>0}(L_\infty, k)$  the positive degree cohomology.

---

ML was partially supported by NSF grant DMS-1702152. VL was partially supported by Laboratory of Mirror Symmetry NRU HSE, RF Government grant, ag. No. 14.641.31.0001.

**Theorem 1.1.** *The functor  $\Phi_L$  is an equivalence if and only if the  $L_{\text{nil}}$ -module  $H^{>0}(L_\infty, k)$  has no subquotients isomorphic to the trivial module  $k$ . For example  $\Phi_L$  is an equivalence if  $L$  is nilpotent.*

We find it natural to approach Theorem 1.1 by studying the graded Rees algebra

$$U(L)^* := \bigoplus_{n \geq 0} I^n = U(L) \oplus I \oplus I^2 \oplus \dots$$

It is easy to prove the following result.

**Proposition 1.2.** *If the algebra  $U(L)^*$  is graded left Noetherian, then the functor  $\Phi_L$  is an equivalence.*

It is, however, not necessary for  $U(L)^*$  to be graded left Noetherian in order for  $\Phi_L$  to be an equivalence.

The next theorem may be of independent interest.

**Theorem 1.3.** *The algebra  $U(L)^*$  is graded left Noetherian if and only if  $L$  is nilpotent.*

In the last section of the paper we mention an application of Theorem 1.1 to the cohomology of quasi-coherent sheaves which are equivariant with respect to a unipotent group.

In this paper we consider only left modules, but all the results are also valid (with the same proofs) for right modules.

We fix a field  $k$ . All Lie algebras are finite dimensional over  $k$ . All associative rings are unital.

*Acknowledgements.* We thank Grigory Papayanov for a useful discussion.

## 2. A CRITERION FOR EQUIVALENCE OF CATEGORIES

Let  $R$  be an associative left Noetherian ring with a 2-sided ideal  $I \subset R$ . Let  $M$  be a left  $R$ -module. An element  $m \in M$  is called  $I$ -torsion, if  $I^n m = 0$  for some  $n > 0$ . The collection of  $I$ -torsion elements in  $M$  is an  $R$ -submodule, which we denote by  $M_I$ . We say that  $M$  is torsion if  $M_I = M$ .

Let  $R\text{-mod}$  denote the abelian category of finitely generated left  $R$ -modules and let  $(R\text{-mod})_I \subset R\text{-mod}$  be its full Serre subcategory of  $I$ -torsion modules. Let  $C^b(R\text{-mod})$  (resp.  $C^b((R\text{-mod})_I)$ ) be the category of bounded complexes over  $R\text{-mod}$  (resp. over  $(R\text{-mod})_I$ ) and let  $C_I^b(R\text{-mod}) \subset C^b(R\text{-mod})$  be the full subcategory of complexes whose cohomology groups are torsion.

In the bounded derived category  $D^b(R\text{-mod})$  consider the full subcategory  $D_I^b(R\text{-mod})$  of complexes with torsion cohomology groups. We have the obvious functor

$$\Phi = \Phi_R : D^b((R\text{-mod})_I) \rightarrow D_I^b(R\text{-mod})$$

**Proposition 2.1.** *Assume that for every finitely generated left  $R$ -module  $M$  there exists a submodule  $N \subset M$  such that  $N_I = 0$  and  $M/N$  is  $I$ -torsion. Then the functor  $\Phi$  is an equivalence.*

*Proof.* Let  $A^\bullet$  be an object of  $C_I^b(R\text{-mod})$ . We claim there exists an object  $B^\bullet$  of  $C^b((R\text{-mod})_I)$  and a morphism of complexes  $f : A^\bullet \rightarrow B^\bullet$  which is a quasi-isomorphism. Indeed, let

$$A^\bullet = 0 \rightarrow A^i \xrightarrow{d^i} A^{i+1} \xrightarrow{d^{i+1}} \dots \xrightarrow{d^{n-1}} A^n \rightarrow 0$$

and let  $t$  be the lowest index such that  $A_I^t \neq A^t$ . By assumption there exists a submodule  $P \subset A^t$  such that  $P_I = 0$  and  $(A^t/P)_I = A^t/P$ . We claim that  $P \cap \ker d^t = 0$ . Indeed, since  $H^t(A^\bullet)$  and  $A^{t-1}$  are torsion, it follows that  $\ker d^t$  is torsion, so  $P \cap \ker d^t = 0$ . Therefore, the complex  $A^\bullet$  contains an acyclic subcomplex  $\tilde{P} := P \xrightarrow{\sim} d^t(P)$  and the components with index  $\leq t$  of the quotient complex  $A^\bullet/\tilde{P}$  are torsion. Iterating this process we find the required quasi-isomorphism  $f : A^\bullet \rightarrow B^\bullet$ . This shows that the functor  $\Phi$  is essentially surjective.

For complexes  $C^\bullet, D^\bullet$  representing objects in  $D^b((R\text{-mod})_I)$ , a morphism  $\Phi(D^\bullet) \rightarrow \Phi(C^\bullet)$  is represented by a diagram of complexes  $D^\bullet \rightarrow A^\bullet \xleftarrow{s} C^\bullet$ , where  $A^\bullet \in C_I^b(R\text{-mod})$  and  $s$  is a quasi-isomorphism. The fact that the functor  $\Phi$  is full and faithful now follows, since (as shown above) there exists a complex  $B^\bullet \in C^b((R\text{-mod})_I)$  and a morphism  $f : A^\bullet \rightarrow B^\bullet$  of complexes that is a quasi-isomorphism.  $\square$

Consider now the graded Rees algebra

$$R^* := \bigoplus_{n \geq 0} I^n = R \oplus I \oplus I^2 \oplus \dots$$

**Lemma 2.2.** *Assume that the algebra  $R^*$  is graded left Noetherian (i.e. every graded left ideal is finitely generated). Then the assumption of Proposition 2.1 holds: for any finitely generated left  $R$ -module  $M$  there exists a submodule  $N \subset M$  such that  $N_I = 0$  and  $M/N = (M/N)_I$*

*Proof.* Let  $M$  be a finitely generated  $R$ -module. If  $M = M_I$ , then we can take  $N = 0$ . So assume that  $M \neq M_I$ , i.e.  $I^s M \neq 0$  for all  $s > 0$ . Consider the graded finitely generated  $R^*$ -module

$$\tilde{M} = M \oplus IM \oplus I^2 M \oplus \dots$$

and its graded submodule

$$P = M_I \oplus (IM \cap M_I) \oplus (I^2 M \cap M_I) \oplus \dots$$

By our assumption,  $P$  is finitely generated; hence, there exists  $n > 0$  such that for all  $m > 0$

$$I^m(I^n M \cap M_I) = I^{n+m} M \cap M_I$$

As  $I^n M \cap M_I$  is finitely generated and  $I$ -torsion, it is annihilated by  $I^m$  for some  $m$ . Thus,

$$(I^{m+n} M)_I = I^{m+n} M \cap M_I = I^m(I^n M \cap M_I) = 0.$$

Putting  $N = I^{m+n} M$ , we have  $N_I = 0$  and  $(M/N)_I = M/N$ .  $\square$

**Corollary 2.3.** *Assume that the algebra  $R^*$  is graded left Noetherian. Then the functor  $\Phi$  is an equivalence.*

### 3. WHEN IS THE REES ALGEBRA OF A UNIVERSAL ENVELOPING ALGEBRA NOETHERIAN

Let  $L$  be a finite dimensional Lie algebra,  $U(L)$  its universal enveloping algebra and  $I \subset U(L)$  the augmentation ideal. As above, we consider the graded Rees algebra

$$U(L)^* = \bigoplus_{n \geq 0} I^n = U(L) \oplus I \oplus I^2 \oplus \dots$$

The main result of this section is the following theorem.

**Theorem 3.1.** *The algebra  $U(L)^*$  is graded left Noetherian if and only if the Lie algebra  $L$  is nilpotent.*

Before proving the theorem we formulate a useful corollary.

**Corollary 3.2.** *Let  $L$  be a nilpotent Lie algebra. Then the functor*

$$\Phi_{U(L)} : D^b((U(L)\text{-mod})_I) \rightarrow D_I^b(U(L)\text{-mod})$$

*is an equivalence.*

*Proof.* This follows from Theorem 3.1 and Corollary 2.3. □

*Proof.* The proof of the theorem will take several steps and will occupy the rest of the section.

**3.1. Proof of the “if” direction.** The universal enveloping algebra  $U(L)$  has a standard increasing filtration which induces a similar filtration in the Rees algebra  $U(L)^*$ . We will prove that if  $L$  is nilpotent, then the associated (double) graded algebra  $\text{gr } U(L)^*$  is commutative finitely generated, hence Noetherian.

So assume for now that the Lie algebra  $L$  is nilpotent.

We define  $L^m$  for positive integers  $m$  recursively;  $L^1 := L$ , and

$$L^d := \sum_{j=1}^{d-1} [L^j, L^{d-j}]$$

for  $m \geq 2$ . As  $L$  is nilpotent,  $L^d = \{0\}$  for all  $d$  sufficiently large. We choose an ordered basis  $\{e_1, \dots, e_n\}$  of  $L$  adapted to the decreasing filtration  $L^\bullet$  in the sense that there exists a non-decreasing sequence  $\nu_1 \leq \dots \leq \nu_n$  such that  $\{e_i \mid \nu_i \geq d\}$  spans  $L^d$  for all  $d$ . Thus, we can write

$$(3.1) \quad [e_i, e_j] = \sum_{\{k \mid \nu_k \geq \nu_i + \nu_j\}} c_k e_k.$$

Write for short  $U := U(L)$ . We recall the standard increasing filtration  $\{U_r\}_{r \geq 0}$  of  $U$  for which  $U_r$  is the span of all products  $x_1 x_2 \cdots x_r$ , where  $x_i \in L \subset U$ . For every multi-index  $a = (a_1, \dots, a_n)$  of non-negative integers,

we denote by  $e^a$  the monomial  $e_1^{a_1} \cdots e_n^{a_n}$ . Thus,  $e^a \in U_{|a|}$ , where  $|a| := a_1 + \cdots + a_n$ . For every  $x \in U$ , we write  $\deg x = 0$  if  $x \in U_0$  and  $\deg x = r$  if  $x \in U_r \setminus U_{r-1}$  for  $r \geq 1$ . We call this the *degree* of  $x$ .

The PBW theorem implies that for all  $r \in \mathbb{N}$  the image of the set

$$\{e^a : |a| = r\} \subset U_r$$

in  $U_r/U_{r-1}$  is a basis. Thus, the set of monomials  $e^a$  as  $a$  ranges over  $\mathbb{N}^n$  is a basis of  $U$ , which we call the *standard basis*, and  $\deg \sum_a c_a e^a$  is the largest value of  $|a|$  for which  $c_a \neq 0$ . For each  $r$ ,  $U_r$  is spanned by  $\{e^a : |a| \leq r\}$ . Moreover, for all  $x, y \in U$ ,

$$(3.2) \quad \deg(x + y) \leq \max(\deg x, \deg y).$$

By the PBW theorem, if  $x$  and  $y$  are non-zero elements of  $U$ , then

$$(3.3) \quad \deg xy = \deg x + \deg y.$$

Let  $I$  denote the augmentation ideal of  $U$ , or equivalently, the span of  $e^a$  for all  $a$  with  $|a| \geq 1$ .

**Lemma 3.3.** *For all positive integers  $m$ ,  $I^m$  has basis*

$$(3.4) \quad \{e^a \mid \sum_i a_i \nu_i \geq m\}.$$

*Proof.* This set is a subset of the standard basis, so it is linearly independent. For  $0 < k < d$ ,  $[L^{d-k}, L^k] \subset L^d$ , so by induction on  $d$ , every  $e_i \in L^d$  lies in  $I^d$ . Thus, the set (3.4) is contained in  $I^m$ .

We prove by induction that every element of  $I^m$  is a linear combination of elements of (3.4), the case  $m = 1$  being trivial. It suffices to prove that if  $i_0, \dots, i_k \in \{1, 2, \dots, n\}$ ,  $i_1 \leq i_2 \leq \cdots \leq i_k$ , and

$$\nu_{i_1} + \cdots + \nu_{i_k} \geq m - 1,$$

then  $e_{i_0} e_{i_1} \cdots e_{i_k}$  lies in the span of (3.4) for  $m$ . We prove more precisely that for any sequence  $i_0, \dots, i_k \in \{1, \dots, n\}$  which is non-decreasing with the possible exception of  $i_0$ , the product  $e_{i_0} e_{i_1} \cdots e_{i_k}$  is spanned by terms of the form  $e^a$  where  $|a| \leq k + 1$  and

$$(3.5) \quad \sum_i a_i \nu_i \geq \nu_{i_0} + \cdots + \nu_{i_k}.$$

We use double induction, first on  $k$  and then on  $i_0$ . The base case  $k = 0$  is trivial and for given  $k$  there is nothing to prove if  $i_0 \leq i_1$ , so the base case  $i_0 = 1$  is trivial. If  $i_0 > i_1$ , then writing

$$e_{i_0} e_{i_1} e_{i_2} \cdots e_{i_k} = e_{i_1} e_{i_0} e_{i_2} \cdots e_{i_k} + [e_{i_0}, e_{i_1}] e_{i_2} \cdots e_{i_k},$$

we need only prove the claim for both summands on the right hand side. By the induction hypothesis on  $k$ ,  $e_{i_0} e_{i_2} \cdots e_{i_k}$  is a linear combination of  $e^b$  for  $|b| \leq k$  and

$$\sum_i b_i \nu_i \geq \nu_{i_0} + \nu_{i_2} \cdots + \nu_{i_k}.$$

By the induction hypothesis on  $i_0, e_{i_1}$  times any such  $e^b$  is a linear combination of terms  $e^a$  with  $|a| \leq k+1$  and  $a$  satisfying (3.5).

By (3.1), we can write  $[e_{i_0}, e_{i_1}]$  as a linear combination of basis vectors  $e_j$  with  $\nu_j \geq \nu_{i_0} + \nu_{i_1}$ , so by the induction hypothesis on  $k$ ,  $[e_{i_0}, e_{i_1}]e_{i_2} \cdots e_{i_k}$  is a linear combination of terms of the form  $e^a$  where  $|a| \leq k$  and

$$\begin{aligned} \sum_i a_i \nu_i &\geq \nu_j + \nu_{i_2} + \cdots + \nu_{i_k} \\ &\geq \nu_{i_0} + \nu_{i_1} + \cdots + \nu_{i_k}. \end{aligned}$$

□

The filtration  $\{U_r\}$  of the universal enveloping algebra  $U = U(L)$  induces the filtration of the graded Rees algebra  $U(L)^*$ : we filter each summand  $I^m$  of  $U(L)^*$  by the induced filtration given by  $U_\bullet$ :

$$F_r U(L)^* := \bigoplus_{m=0}^{\infty} (I^m \cap U_r)$$

We get

$$\text{gr } U(L)^* = \bigoplus_{r,m} (I^m \cap U_r) / (I^m \cap U_{r-1})$$

is a (doubly) graded commutative algebra. We denote

$$U(L)^*_{r,m} = (I^m \cap U_r) / (I^m \cap U_{r-1})$$

so that  $\text{gr } U(L)^* = \bigoplus_{r,m} U(L)^*_{r,m}$ . By the PBW theorem,  $U_1 = L$ , so by Lemma 3.3,  $U(L)^*_{1,m}$  is naturally identified with  $L^m$ .

**Proposition 3.4.** *The graded algebra  $\text{gr } U(L)^*$  is a commutative graded algebra generated by  $U(L)^*_{1,m}$ ,  $m = 0, 1, \dots, s$ , where  $L^{s+1} = \{0\}$ .*

*Proof.* From Lemma 3.3, we see that  $I^m \cap U_r$  has basis

$$\{e^a \mid \sum_i a_i \nu_i \geq m, |a| \leq r\},$$

so  $U(L)^*_{r,m}$  has basis

$$\{e^a \mid \sum_i a_i \nu_i \geq m, |a| = r\}.$$

Notice that under the multiplication map

$$U(L)^*_{r,m} \otimes U(L)^*_{r',m'} \rightarrow U(L)^*_{r+r',m+m'}$$

we have  $e^a \cdot e^b = e^{a+b}$ . To prove that the classes  $U(L)^*_{1,m}$  generate, it suffices to note that the additive monoid

$$\{(a_1, \dots, a_n, r, m) \in \mathbb{N}^{n+2} \mid \sum_i a_i \nu_i \geq m, |a| = r\}$$

is generated by the set

$$\left\{ \underbrace{(0, \dots, 0)}_{i-1}, 1, \underbrace{(0, \dots, 0)}_{n-i}, 1, m \mid 1 \leq i \leq n, 0 \leq m \leq \nu_i \right\}$$

whose elements correspond to  $e_i \in L^m = U(L)_{1,m}^* \subset L$ .  $\square$

Now we recall a useful general result. Let  $R$  be an associative ring with increasing exhausting filtration

$$0 = F_{-1} \subset F_0 \subset F_1 \subset \dots = R$$

and consider the associated graded ring

$$\text{gr } R = \text{gr}_{F_\bullet} R = \bigoplus_{n \geq 0} F_n / F_{n-1}$$

If  $x \in F_n \setminus F_{n-1}$  we say that  $x$  has degree  $n$  and denote by  $\bar{x}$  its image in  $F_n / F_{n-1}$ .

**Lemma 3.5.** *Assume that  $\text{gr } R$  is graded left Noetherian. Then the ring  $R$  is left Noetherian.*

*Proof.* Let  $I \subset R$  be a left ideal. Then

$$\text{gr } I := \bigoplus_{n \geq 0} (I \cap F_n) / (I \cap F_{n-1})$$

is a graded left ideal in  $\text{gr } R$ . Let  $\bar{x}_1, \dots, \bar{x}_k$  be a set of homogeneous generators of the ideal  $\text{gr } I$ ; say  $\deg(\bar{x}_i) = n_i$ . Choose lifts  $x_1, \dots, x_k \in I$  of the  $\bar{x}_i$ 's and let  $(x_1, \dots, x_k) \subset I$  be the corresponding left ideal. We claim that  $(x_1, \dots, x_k) = I$ . Indeed, let  $x \in I$  be of degree  $d$ . If  $d = 0$ , then clearly  $x \in (x_1, \dots, x_k)$ . Otherwise, there exist  $r_i \in R$  of degree  $d - n_i$  such that

$$\bar{x} = \sum \bar{r}_i \bar{x}_i$$

It follows that  $x - \sum r_i x_i \in I \cap F_{d-1}$ , hence by induction on  $d$ ,  $x - \sum r_i x_i \in (x_1, \dots, x_k)$ .  $\square$

It follows from Proposition 3.4 and Lemma 3.5 that the Rees algebra  $U(L)^*$  is left Noetherian if the Lie algebra is nilpotent. This proves the “if” direction of Theorem 3.1.

**3.2. Proof of the “only if” direction.** For a Lie algebra  $L$  we consider the lower central series  $L_1 = L$ ,  $L_n = [L, L_{n-1}]$ . Thus

$$L = L_1 \supset L_2 \supset L_3 \supset \dots$$

is a nonincreasing sequence of ideals in  $L$ . We put

$$L_\infty := \bigcap_n L_n, \quad L_{\text{nil}} = L / L_\infty$$

The Lie algebra  $L_{\text{nil}}$  is nilpotent, and  $L$  is nilpotent if and only if  $L_\infty = 0$ .

We have the short exact sequence of Lie algebras

$$0 \rightarrow L_\infty \rightarrow L \rightarrow L_{\text{nil}} \rightarrow 0$$

This induces the surjection  $\theta : U(L) \rightarrow U(L_{\text{nil}})$  and  $\ker \theta$  is the ideal  $U(L)L_{\infty}U(L)$ . As before, let  $I \subset U(L)$  be the augmentation ideal.

**Lemma 3.6.** *We have the equality of ideals in  $U(L)$ :*

$$\bigcap_n I^n = \ker \theta$$

*Proof.* We have by construction  $L_n \subset I^n$  for all  $n$ , hence  $L_{\infty} \subset \bigcap_n I^n$  and so

$$\ker \theta = U(L)L_{\infty}U(L) \subset \bigcap_n I^n$$

To prove the opposite inclusion let  $\bar{I} \subset U(L_{\text{nil}})$  be the augmentation ideal, so that we have the surjection  $\theta : I \rightarrow \bar{I}$ . It suffices to prove that  $\bigcap_n \bar{I}^n = 0$ . This follows from Lemma 3.3.  $\square$

Assume that the Lie algebra  $L$  is not nilpotent, i.e.  $L_{\infty} \neq 0$ . Then Lemma 3.6 implies that  $L_{\infty} \subset \bigcap_n I^n$ . Fix  $0 \neq x \in L_{\infty}$  and consider the graded left ideal

$$J = \bigoplus_n U(L)x_n \subset U(L)^*$$

where  $x_n$  denotes the copy of  $x$  in  $I^n$ . We claim that  $J$  is not finitely generated. Assume, on the contrary, that

$$J = \sum_i U(L)^* f_i$$

for a finite number of homogeneous elements  $f_i \in U(L)x_{n_i}$ . Choose  $m > n_i$  for all  $i$ . We claim that

$$x_m \notin \sum_i U(L)^* f_i$$

Indeed, it suffices to notice that  $x \notin Ix$ : if  $x = fx$ , then  $f = 1$ , (since  $U(L)$  is a domain) and hence  $f \notin I$ . This completes the proof of Theorem 3.1.  $\square$

#### 4. MAIN THEOREM

Let  $L$  be a finite dimensional Lie algebra,  $U(L)$  its universal enveloping algebra,  $I \subset U(L)$  the augmentation ideal. As in Subsection 3.2, consider the ideal

$$L_{\infty} = \bigcap_n L_n \subset L$$

and the quotient nilpotent Lie algebra  $L_{\text{nil}} = L/L_{\infty}$ .

Each cohomology space  $H^i(L_{\infty}, k)$  is naturally a  $L_{\text{nil}}$ -module, so we have the Hochschild-Serre spectral sequence [HS]:

$$(4.1) \quad E_2^{pq} = H^p(L_{\text{nil}}, H^q(L_{\infty}, k)) \Rightarrow H^{p+q}(L, k).$$

**Theorem 4.1.** *Let  $L$  be a finite dimensional Lie algebra over a field  $k$ . The following conditions are equivalent:*

(1) *The natural functor*

$$\Phi_L : D^b((U(L)\text{-mod})_I) \rightarrow D^b_I(U(L)\text{-mod})$$

*is an equivalence.*

(2) *The natural map  $H^\bullet(L_{\text{nil}}, k) \rightarrow H^\bullet(L, k)$  is an isomorphism.*

(3) *The positive degree cohomology  $H^{>0}(L_\infty, k)$  considered as an  $L_{\text{nil}}$ -module has no subquotients isomorphic to the trivial module  $k$ .*

*Proof.* We first notice that the 3 conditions in the theorem hold in case  $L$  is nilpotent. Indeed, then  $L_\infty = 0$ , so (2) and (3) hold trivially. Also (1) holds by Corollary 3.2.

Let now  $L$  be general. As in section 3.2 we consider the short exact sequence of Lie algebras

$$0 \rightarrow L_\infty \rightarrow L \rightarrow L_{\text{nil}} \rightarrow 0$$

and the induced surjection  $\theta : U(L) \rightarrow U(L_{\text{nil}})$  with the kernel  $\ker \theta = \bigcap_n I^n$  (Lemma 3.6). Also, as in the proof of Lemma 3.6, denote by  $\bar{I}$  the augmentation ideal in  $U(L_{\text{nil}})$ . Lemma 3.6 implies that any  $U(L)$ -module  $M$  such that  $M = M_I$  is actually a  $U(L_{\text{nil}})$ -module (and  $M = M_{\bar{I}}$ ). Hence the functor of restriction of scalars

$$\theta_* : U(L_{\text{nil}})\text{-mod} \rightarrow U(L)\text{-mod}$$

induces the equivalence of categories

$$(U(L_{\text{nil}})\text{-mod})_{\bar{I}} \xrightarrow{\sim} (U(L)\text{-mod})_I$$

and therefore the equivalence of categories

$$(4.2) \quad D^b((U(L_{\text{nil}})\text{-mod})_{\bar{I}}) \xrightarrow{\sim} D^b((U(L)\text{-mod})_I)$$

We have the commutative diagram of functors

$$(4.3) \quad \begin{array}{ccc} D^b((U(L)\text{-mod})_I) & \xrightarrow{\Phi_L} & D^b_I(U(L)\text{-mod}) \\ \uparrow & & \uparrow \\ D^b((U(L_{\text{nil}})\text{-mod})_{\bar{I}}) & \xrightarrow{\Phi_{L_{\text{nil}}}} & D^b_{\bar{I}}(U(L_{\text{nil}})\text{-mod}) \end{array}$$

As explained above the left vertical arrow is an equivalence. Also  $\Phi_{L_{\text{nil}}}$  is an equivalence (Corollary 3.2). Hence  $\Phi_L$  is an equivalence if and only if the functor

$$(4.4) \quad \theta_* : D^b_{\bar{I}}(U(L_{\text{nil}})\text{-mod}) \rightarrow D^b_I(U(L)\text{-mod})$$

is an equivalence. Every finitely generated  $I$ -torsion  $U(L)$ -module (resp.  $\bar{I}$ -torsion  $U(L_{\text{nil}})$ -module) is a finite dimensional  $k$ -vector space on which  $U(L)$  (resp.  $U(L_{\text{nil}})$ ) acts nilpotently, so by Engel's theorem, it admits a stable flag with trivial 1-dimensional quotients. As triangulated categories,

therefore, both sides of (4.4) are generated by the trivial module  $k$ , and so the functor in (4.4) is an equivalence if and only if the natural map

$$(4.5) \quad \theta_* : \text{Ext}_{U(\mathcal{L}_{\text{nil}})}^\bullet(k, k) \rightarrow \text{Ext}_{U(\mathcal{L})}^\bullet(k, k)$$

is an isomorphism. This proves the equivalence of conditions (1) and (2) in the theorem. It remains to prove the equivalence of (2) and (3).

First we prove a lemma. Let  $\mathcal{L}$  be a nilpotent Lie algebra. By a theorem of Lie every simple  $\mathcal{L}$ -module  $M$  is one dimensional, hence it corresponds to an additive character

$$\chi_M : \mathcal{L}/[\mathcal{L}, \mathcal{L}] \rightarrow k$$

**Lemma 4.2.** *Let  $M$  be a simple nontrivial  $\mathcal{L}$ -module. Then*

$$H^\bullet(\mathcal{L}, M) = 0$$

*Proof.* The character  $\chi_M$  gives a short exact sequence of nilpotent Lie algebras

$$(4.6) \quad 0 \rightarrow \mathcal{L}' \rightarrow \mathcal{L} \xrightarrow{\chi_M} \bar{\mathcal{L}} \rightarrow 0$$

with  $\dim \bar{\mathcal{L}} = 1$ , where  $\mathcal{L}'$  acts trivially on  $M$ .

Recall [CE, (23.1)] the standard complex  $C^\bullet(\mathcal{L}', M)$ , which computes the cohomology  $H^\bullet(\mathcal{L}', M)$ . With our normalization, it is

$$(4.7) \quad 0 \rightarrow M = C^0(\mathcal{L}', M) \xrightarrow{\partial_0} C^1(\mathcal{L}', M) \xrightarrow{\partial_1} \dots,$$

where  $C^n(\mathcal{L}', M) := \text{Hom}_k(\bigwedge^n \mathcal{L}', M)$  and

$$\begin{aligned} \partial_n(f)(x_0 \wedge \dots \wedge x_n) &= \sum_{i=0}^n (-1)^i x_i f(x_0 \wedge \dots \wedge \hat{x}_i \wedge \dots \wedge x_n) \\ &+ \sum_{p < q} (-1)^{p+q} f([x_p, x_q] \wedge x_0 \wedge \dots \wedge \hat{x}_p \wedge \dots \wedge \hat{x}_q \wedge \dots \wedge x_n) \end{aligned}$$

The  $\mathcal{L}$ -action on  $\mathcal{L}'$  and  $M$  extends to an action on the complex  $C^\bullet(\mathcal{L}', M)$  by the formula

$$(4.8) \quad \begin{aligned} (xf)(x_1 \wedge \dots \wedge x_n) &= xf(x_1 \wedge \dots \wedge x_n) \\ &- \sum_{i=1}^n f(x_1 \wedge \dots \wedge x_{i-1} \wedge [x, x_i] \wedge x_{i+1} \wedge \dots \wedge x_n) \end{aligned}$$

This induces the  $\mathcal{L}$ -action on the cohomology  $H^\bullet(\mathcal{L}', M)$  which is trivial on  $\mathcal{L}'$  and hence gives the required  $\bar{\mathcal{L}}$ -action on  $H^\bullet(\mathcal{L}', M)$ .

We now take a closer look at the  $\mathcal{L}$ -action (4.8) on  $C^n(\mathcal{L}', M)$ . Since the  $\mathcal{L}$ -action on the Lie algebra  $\mathcal{L}'$  is nilpotent there exists a basis of  $\mathcal{L}'$  such that for every  $x \in \mathcal{L}$  the matrix of the operator  $[x, -]$  in this basis is strictly lower triangular. Then it follows from the formula (4.8) that there exists a basis for  $C^n(\mathcal{L}', M)$  such that the action of every  $x \in \mathcal{L}$  is given by a lower triangular matrix with all diagonal entries being  $\chi_M(x)$ .

Therefore any  $\bar{x} \in \bar{\mathcal{L}}$  acts on the cohomology  $H^\bullet(\mathcal{L}', M)$  by an operator whose characteristic polynomial is a power of  $(t - \chi_M(\bar{x}))$ . Since we assume that  $\chi_M(\bar{x}) \neq 0$  for  $\bar{x} \neq 0$ , it easily follows that

$$\text{Ext}_{U(\bar{\mathcal{L}})}^\bullet(k, H^\bullet(\mathcal{L}', M)) = H^\bullet(\bar{\mathcal{L}}, H^\bullet(\mathcal{L}', M)) = 0$$

Now the Hochschild-Serre spectral sequence

$$E_2^{pq} = H^p(\bar{\mathcal{L}}, H^q(\mathcal{L}', M)) \Rightarrow H^{p+q}(\mathcal{L}, M)$$

implies that  $H^\bullet(\mathcal{L}, M) = 0$ , which proves the lemma.  $\square$

We now return to the proof of the equivalence of conditions (2) and (3) in the theorem. The Hochschild-Serre spectral sequence (4.1) has  $E_2$  page

$$(4.9) \quad \begin{array}{ccccccc} E_2^{01} & & E_2^{11} & & \cdots & & \cdots \\ & \searrow & & \searrow & & & \\ E_2^{00} & & E_2^{10} & & E_2^{20} & & \cdots \end{array}$$

We have  $H^0(L_\infty, k) = k$  – the trivial  $L_{\text{nil}}$ -module and the graded space  $\text{Ext}_{U(L_{\text{nil}})}^\bullet(k, k)$  identifies naturally with the bottom row of this spectral sequence. The map  $\theta^* : \text{Ext}_{U(L_{\text{nil}})}^\bullet(k, k) \rightarrow \text{Ext}_{U(L)}^\bullet(k, k)$  then coincides with the projection

$$\text{Ext}_{U(L_{\text{nil}})}^\bullet(k, k) = H^\bullet(L_{\text{nil}}, H^0(L_\infty, k)) \rightarrow H^\bullet(L, k)$$

Notice that by a theorem of Lie every finite dimensional  $L_{\text{nil}}$ -module has a filtration with 1-dimensional subquotients.

Assume that the condition (3) holds, i.e. the  $L_{\text{nil}}$ -module  $H^{>0}(L_\infty, k)$  has no subquotients isomorphic to the trivial module  $k$ . Then all the 1-dimensional subquotients of the  $L_{\text{nil}}$ -module  $H^{>0}(L_\infty, k)$  are nontrivial. In this case it follows from Lemma 4.2 that only the bottom row of the spectral sequence (4.9) is nonzero. Therefore the natural map  $H^\bullet(L_{\text{nil}}, k) \rightarrow H^\bullet(L, k)$  is an isomorphism, i.e. the condition (2) of the theorem holds.

Assume, conversely, that condition (2) holds. Let  $d$  be the maximal integer such that  $H^d(L_{\text{nil}}, k) \neq 0$ . Again using Lie's theorem and Lemma 4.2 it follows that  $H^{>d}(L_{\text{nil}}, N) = 0$  for any finite dimensional  $L_{\text{nil}}$ -module  $N$ . If  $k$  is a subquotient of  $N$ , then by Engel's theorem, it is also a quotient of  $N$ . It follows, therefore, that if  $N$  has the trivial module as a subquotient, then  $H^d(L_{\text{nil}}, N) \neq 0$ . So if for some  $i > 0$  the  $L_{\text{nil}}$ -module  $H^i(L_\infty, k)$  has the trivial submodule  $k$  as a subquotient then  $E_2^{d,i} = H^d(L_{\text{nil}}, H^i(L_\infty, k))$  is nonzero, which means that it survives in  $H^{i+d}(L, k)$ . This is a contradiction and finishes the proof of the theorem.  $\square$

**4.1. Some examples.** (A) Consider the 2-dimensional Lie algebra with basis  $x, y$  and the relation  $[x, y] = y$ . This Lie algebra is solvable but not

nilpotent,  $L_\infty = ky$ . The standard complex, which computes the cohomology  $H^\bullet(L_\infty, k)$  has terms in degrees 0 and 1 and zero differential

$$k \xrightarrow{0} \text{Hom}_k(ky, k)$$

The element  $x \in L_{\text{nil}}$  acts on the space  $\text{Hom}_k(ky, k) = H^1(L_\infty, k)$  as minus the identity (see formula (4.8)), hence the condition (3) of Theorem 4.1 is satisfied.

(B) This is a generalization of example (A) above: assume that the  $L_{\text{nil}}$ -module  $\bigwedge^i L_\infty$  has no subquotients isomorphic to the trivial module  $k$  if  $i > 0$ . Then the condition (3) of Theorem 4.1 holds. For example this is the case when  $L$  is the Lie algebra of upper-triangular matrices. Then  $L_\infty$  is the ideal of strictly triangular matrices and  $L_{\text{nil}}$  is the abelian quotient.

(C) However, there exists solvable algebras  $L$  for which condition (3) does not hold. See Proposition 4.5 below.

(D) It may happen that condition (3) holds even though  $\bigwedge^{>0} L_\infty$  admits  $k$  as an  $L_{\text{nil}}$ -subquotient. See Proposition 4.6 below.

For the next two examples, we assume that  $k$  has characteristic zero.

(E) Assume that  $L_\infty \neq 0$  is semi-simple. If  $k = \mathbb{R}$  and  $L_\infty$  is compact, the cohomology  $H^\bullet(L_\infty, k)$  is isomorphic to the cohomology of any compact Lie group with the Lie algebra  $L_\infty$  [CE, Theorem 15.2], so  $H^{>0}(L_\infty, k) \neq 0$ . Since every compact semisimple Lie algebra has a compact real form,  $H^{>0}(L_\infty, k) \neq 0$  when  $k = \mathbb{C}$  and  $L_\infty$  is semisimple, the same statement follows for every Lie algebra over any field  $k$  of characteristic zero. However for any  $x \in L_{\text{nil}}$  the operator  $[x, -]$  on  $L_\infty$  is a derivation, so is inner. Therefore the action of  $L_{\text{nil}}$  on the cohomology is trivial and so the condition (3) of Theorem 4.1 fails.

(F) This is a generalization of example (E) above. We formulate it as a proposition.

**Proposition 4.3.** *Assume that the equivalent conditions of Theorem 4.1 are satisfied and the characteristic of  $k$  is zero. Then the algebra  $L_\infty$  is solvable.*

*Proof.* By the general theory the Lie algebra  $L_\infty$  has a maximal solvable ideal (the radical)  $L_\infty^{\text{rad}}$  such that

$$L_\infty^{\text{b}} := L_\infty / L_\infty^{\text{rad}}$$

is semi-simple. Put  $\mathfrak{g} := L_\infty^{\text{b}}$ . We need to prove that  $\mathfrak{g} = 0$ .

**Lemma 4.4.** *In the above notation the natural map*

$$H^\bullet(\mathfrak{g}, k) = \text{Ext}_{U(\mathfrak{g})}^\bullet(k, k) \rightarrow \text{Ext}_{U(L_\infty)}^\bullet(k, k) = H^\bullet(L_\infty, k)$$

*is injective.*

*Proof.* By the Levi theorem we know that the surjection of Lie algebras  $p : L_\infty \rightarrow \mathfrak{g}$  has a splitting  $s : \mathfrak{g} \rightarrow L_\infty$ . The corresponding surjection of universal enveloping algebras  $p : U(L_\infty) \rightarrow U(\mathfrak{g})$  induces the pair of adjoint

functors  $(Lp^*, p_*)$  – the extension and restriction of scalars between the derived categories  $D^b(U(L_\infty)\text{-mod})$  and  $D^b(U(\mathfrak{g})\text{-mod})$ , where  $A\text{-mod}$  is the category of (all) left modules over an associative ring  $A$ . It suffices to prove that the adjunction morphism of functors

$$\text{Id}_{D^b(U(\mathfrak{g})\text{-mod})} \rightarrow Lp^* \cdot p_*$$

has a left inverse, i.e.  $\text{Id}_{D^b(U(\mathfrak{g})\text{-mod})}$  is a direct summand of the functor  $Lp^* \cdot p_*$ .

Let  $S^\bullet$  be an object in  $D^b(U(\mathfrak{g})\text{-mod})$ . We may assume that  $S^\bullet$  consists of projective  $U(\mathfrak{g})$ -modules. Let us construct a special (functorial) projective resolution of  $p_*S^\bullet$ . The morphism  $s : \mathfrak{g} \rightarrow L_\infty$  gives a homomorphism  $s : U(\mathfrak{g}) \rightarrow U(L_\infty)$  such that  $p \cdot s = \text{id}$ . So we may consider  $U(L_\infty)$  as a (free) right  $U(\mathfrak{g})$ -module via the homomorphism  $s$ . Consider the obvious short exact sequence of complexes of  $U(L_\infty)$ -modules

$$0 \rightarrow K^\bullet \rightarrow U(L_\infty) \otimes_{U(\mathfrak{g})} S^\bullet \rightarrow p_*S^\bullet \rightarrow 0$$

Now we repeat this procedure with  $K^\bullet$  instead of  $S^\bullet$  (by first considering  $K^\bullet$  as a complex of  $U(\mathfrak{g})$ -modules via the map  $s$ ) and so on. Eventually we obtain the complex of (complexes of projective)  $U(L_\infty)$ -modules

$$P^\bullet := \cdots \xrightarrow{\partial_1} U(L_\infty) \otimes_{U(\mathfrak{g})} K^\bullet \xrightarrow{\partial_0} U(L_\infty) \otimes_{U(\mathfrak{g})} S^\bullet$$

which is a resolution of  $p_*S^\bullet$  and hence

$$Lp^* \cdot p_*S^\bullet = U(\mathfrak{g}) \otimes_{U(L_\infty)} P^\bullet$$

Note that by construction of  $P^\bullet$ , the map  $U(\mathfrak{g}) \otimes_{U(L_\infty)} \partial_0$  is zero. Hence  $U(\mathfrak{g}) \otimes_{U(L_\infty)} (U(L_\infty) \otimes_{U(\mathfrak{g})} S^\bullet) = S^\bullet$  is a direct summand of  $Lp^* \cdot p_*S^\bullet$ , which proves the lemma.  $\square$

Now the assertion of the proposition follows from the example (E) above. Indeed, if  $\mathfrak{g} \neq 0$ , then  $H^{>0}(\mathfrak{g}, k) \neq 0$ . As explained in example (E) the  $L_{\text{nil}}$  action on  $H^{>0}(\mathfrak{g}, k) \neq 0$  is trivial, so by the above lemma the space  $H^{>0}(L_\infty, k)$  contains a nonzero  $L_{\text{nil}}$ -submodule, which is trivial. This contradicts condition (3) of Theorem 4.1.  $\square$

**Proposition 4.5.** *There exist solvable Lie algebras  $L$  not satisfying the equivalent conditions of Theorem 4.1.*

*Proof.* For a nilpotent Lie algebra  $N$  acting on the trivial module  $k$ , the top differential,  $C^{\dim N - 1}(N, k) \rightarrow C^{\dim N}(N, k)$ , of the Chevalley-Eilenberg complex is always zero. Therefore, if  $L_\infty$  is nilpotent and  $L_{\text{nil}}$  acts trivially on  $\bigwedge^{\dim L_\infty} L_\infty$ , then condition (3) of Theorem 4.1 is violated.

Let  $L$  be the subalgebra of upper-triangular  $3 \times 3$  matrices such that the upper left and lower right entries are the same. Thus  $L_\infty$  consists of strictly upper triangular matrices, and its center  $Z$  consists of matrices which are zero except possibly in the upper right entry. The commutator map  $L_\infty/Z \times L_\infty/Z \rightarrow Z$  respects the action of  $L_{\text{nil}}$ , which is trivial on  $Z$ . This

implies that  $\bigwedge^2(L_\infty/Z) \cong k$  as  $L_{\text{nil}}$ -module, and therefore  $\bigwedge^3 L_\infty \cong k$  as  $L_{\text{nil}}$ -module.  $\square$

**Proposition 4.6.** *The conditions of Theorem 4.1 are strictly weaker than the condition that  $(\bigwedge^{>0} L_\infty^*)$  has a non-trivial  $L_{\text{nil}}$ -invariant subquotient.*

*Proof.* As  $H^{>0}(L_\infty, k)$  is a subquotient of  $\bigwedge^{>0} L_\infty^*$ , if the former has a non-trivial  $L_{\text{nil}}$ -invariant subquotient, the latter does as well.

We show that converse does not hold by exhibiting a case in which  $L_{\text{nil}}$  acts semisimply on  $\bigwedge^\bullet L_\infty^*$  and therefore on every  $L_{\text{nil}}$ -stable subquotient and for which

$$\dim(\bigwedge^\bullet L_\infty^*)^{L_{\text{nil}}} > 1 = \dim H^\bullet(L_\infty, k)^{L_{\text{nil}}}.$$

The free Lie algebra on two generators  $x$  and  $y$  admits a unique bigrading for which  $x$  and  $y$  have bidegree  $(1, 0)$  and  $(0, 1)$  respectively. Let  $M$  be the quotient of this algebra by the graded ideal generated by all elements of total degree  $\geq 4$  and also  $[[x, y], y]$ . Then  $M$  has basis:  $x, y, z, w$  of bidegree  $(1, 0)$ ,  $(0, 1)$ ,  $(1, 1)$ , and  $(2, 1)$  respectively, satisfying the following relations:

$$[x, y] = z, [x, z] = w, [x, w] = [y, z] = [y, w] = [z, w] = 0$$

(see [Bo, II, §2, no. 11, Théorème 1] and the computation of the Hall set for 2 generators given at the end of no. 10.)

We define  $t$  to be the derivation which acts on the bidegree  $(a, b)$  part of  $M$  by  $2a - 3b$ . Let  $L := M \oplus kt$  denote the semi-direct sum, so

$$[t, x] = 2x, [t, y] = -3y, [t, z] = -z, [t, w] = w.$$

We confirm that  $[L, L] = [L, M] = M$ , so  $L_\infty = M$ , and  $L_{\text{nil}}$  is the 1-dimensional algebra spanned by the class of  $t$ .

Next, we consider the Chevalley-Eilenberg complex of  $M$ . The underlying graded space is  $\bigwedge^\bullet M^*$ , which is spanned by wedge products of the dual basis  $x^*, y^*, z^*, w^*$  of  $M$ . The differential is given by

$$\delta(x^*) = 0, \delta(y^*) = 0, \delta(z^*) = y^* \wedge x^*, \delta(w^*) = z^* \wedge x^*.$$

The bigrading on  $M$  induces a bigrading on  $\bigwedge^\bullet M^*$ , and  $\delta$  preserves bidegree. The degree  $(3, 2)$ -part of  $\bigwedge^\bullet M^*$  is spanned by  $z^* \wedge w^*$  and  $x^* \wedge y^* \wedge w^*$ . As

$$\delta(z^* \wedge w^*) = -x^* \wedge y^* \wedge z^*,$$

the degree  $(3, 2)$ -part of  $H^*(L_\infty, k)$  is zero. On the other hand, the  $t$ -invariant part of  $H^*(L_\infty, k)$  is the sum of the  $(3n, 2n)$ -part over all integers  $n$ .

Now  $H^*(L_\infty, k)$  is a subquotient of  $\bigwedge^\bullet M^*$ , and the latter has non-trivial degree  $(3n, 2n)$ -part only for  $n = 0, 1$ . Thus,  $\dim(\bigwedge^\bullet L_\infty)^{L_{\text{nil}}} = 3$  but

$$\dim H^\bullet(L_\infty, k)^{L_{\text{nil}}} = 1.$$

$\square$

## 5. AN APPLICATION

Let  $k$  be an algebraically closed field of characteristic zero. Let  $V$  be a linear unipotent algebraic group over  $k$ ,  $L = \text{Lie } V$  the corresponding nilpotent Lie algebra.

Denote by  $V\text{-Mod}$  the abelian category of rational representations of  $V$ . Recall that an object of  $V\text{-mod}$  is by definition a  $V$ -module  $M$  which is a union of finite dimensional submodules  $M_i$ , such that the  $V$ -action on  $M_i$  comes from a homomorphism of  $k$ -algebraic groups  $V \rightarrow GL(M_i)$ . In particular every element of  $V$  acts on  $M_i$  via a unipotent operator.

Notice that we have a natural equivalence of abelian categories

$$\log : V\text{-Mod} \rightarrow (U(L)\text{-Mod})_I$$

where  $(U(L)\text{-Mod})_I$  is the abelian category of (all)  $U(L)$ -modules which are  $I$ -torsion.

This induces the equivalence of derived categories

$$(5.1) \quad \log : D^b(V\text{-Mod}) \rightarrow D^b((U(L)\text{-Mod})_I)$$

Recall that for  $M \in V\text{-mod}$  its cohomology is by definition

$$H_V^\bullet(M) := \text{Ext}_{V\text{-Mod}}^\bullet(k, M)$$

where  $k$  is the trivial rational  $V$ -module.

**Corollary 5.1.** *For any  $M \in V\text{-Mod}$  we have the isomorphism*

$$(5.2) \quad H_V^\bullet(M) \simeq H^\bullet(L, \log(M))$$

*In particular, the cohomology  $H_V^\bullet(M)$  can be computed using the standard complex for the Lie algebra  $L$ .*

*Proof.* The equivalence (5.1) implies the isomorphism

$$(5.3) \quad \text{Ext}_{V\text{-Mod}}^\bullet(k, M) = \text{Ext}_{(U(L)\text{-Mod})_I}^\bullet(k, \log(M))$$

The module  $M$  is a direct limit (union) of its finite dimensional submodules. The cohomology on both sides of (5.2) commutes with direct limits, hence we may assume that  $\dim_k M < \infty$  and so the  $\log(M) \in U(L)\text{-mod}$ . Using the standard methods one can show that

$$\text{Ext}_{(U(L)\text{-Mod})_I}^\bullet(k, \log(M)) = \text{Ext}_{(U(L)\text{-mod})_I}^\bullet(k, \log(M))$$

Finally, Corollary 3.2 implies the isomorphism

$$\text{Ext}_{(U(L)\text{-mod})_I}^\bullet(k, \log(M)) = \text{Ext}_{U(L)\text{-mod}}^\bullet(k, \log(M))$$

which proves the corollary.  $\square$

Let  $X$  be a  $k$ -scheme with an action of the group  $V$ . For a  $V$ -equivariant quasi-coherent sheaf  $F$ , its cohomology can be computed as

$$H_V^\bullet(X, F) = \text{Ext}_{V\text{-Mod}}^\bullet(k, \mathbb{R}\Gamma(X, F)),$$

and sometimes one wants to know that the Ext-space  $\text{Ext}_{V\text{-Mod}}^\bullet(k, -)$  can be computed using the standard complex for the Lie algebra  $L$  (by Corollary

5.1). This fact was used, for example, in the key computation on p. 8 of [Te].

#### REFERENCES

- [Bo] N. Bourbaki, *Éléments de mathématique*. Fasc. XXXVII. Groupes et algèbres de Lie. Chapitre II: Algèbres de Lie libres. Chapitre III: Groupes de Lie. *Actualités Scientifiques et Industrielles*, No. 1349. Hermann, Paris, 1972.
- [CE] C. Chevalley and S. Eilenberg, Theory of Lie groups and Lie algebras. *Trans. Amer. Math. Soc.* **63** (1948), 85–124.
- [HS] G. Hochschild and J-P. Serre, Cohomology of Lie algebras, *Annals of Math.* **57** (1953), 591–603.
- [Te] C. Teleman, The quantization conjecture revisited, *Annals of Math.* **152**, no. 1, (2000), 1–43.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405,  
U.S.A.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405,  
U.S.A.; HSE UNIVERSITY, RUSSIAN FEDERATION