

A Note on Residual Variables of an Affine Fibration

Prosenjit Das

*Department of Mathematics, Indian Institute of Space Science and Technology,
Valiamala P.O., Trivandrum 695 547, India
email: prosenjit.das@gmail.com*

Amartya K. Dutta

*Stat-Math Unit, Indian Statistical Institute,
203, B.T. Road, Kolkata 700 108, India
email: amartya@isical.ac.in*

Abstract

In a recent paper [El 13], M.E. Kahoui has shown that if R is a polynomial ring over \mathbb{C} , A an \mathbb{A}^3 -fibration over R , and W a residual variable of A then A is stably polynomial over $R[W]$. In this article we show that the above result holds over *any* Noetherian domain R provided the module of differentials $\Omega_R(A)$ of the affine fibration A (which is necessarily a projective A -module by a theorem of Asanuma) is a stably free A -module.

Keywords: Residual variable; Stably polynomial algebra; \mathbb{A}^n -fibrations, Module of differentials.

Mathematics Subject classifications (2010): Primary 14R25; Secondary 13B25, 13F20

1. Introduction

Let R denote a Noetherian integral domain. A finitely generated flat R -algebra A is said to be an \mathbb{A}^n -fibration over R if $A \otimes_R k(P)$ is a polynomial ring in n variables over $k(P)$ for every prime ideal P of R . The concept of residual variable was defined by S.M. Bhatwadekar and A.K. Dutta in [BD93] as an element W in the polynomial algebra $R[X, Y]$ for which $R[X, Y] \otimes_R k(P)$ is a polynomial algebra in one variable over $R[W] \otimes_R k(P)$ for every prime ideal P of R . The following statement is a part of their main result ([BD93], Theorem 3.1) on residual variables:

Theorem 1.1. *Let R be a Noetherian domain and W an element of $R[X, Y]$. Then the following are equivalent:*

- (1) W is a residual variable in $R[X, Y]$.
- (2) $R[X, Y]$ is a stably polynomial algebra over $R[W]$, i.e., $R[X, Y][Z_1, Z_2, \dots, Z_r] = R[W][T_0, T_1, \dots, T_r]$ for some indeterminates Z_i 's, T_i 's over R .

They also observed ([BD93], Remark 3.4) that an analogous result holds for a system of m algebraically independent elements W_1, W_2, \dots, W_m in the polynomial ring $R[X_1, X_2, \dots, X_{m+1}]$. Recently El Kahoui [El 13] has extended the concept of residual variable of a polynomial ring to that of an \mathbb{A}^n -fibration: he defines an element W of an \mathbb{A}^n -fibration A over a ring R to be a residual variable of A over R if $A \otimes_R k(P)$ is a polynomial algebra in $n - 1$ variables over $R[W] \otimes_R k(P)$ for each prime ideal P of R . He shows that when $n = 3$ and the base ring R is a polynomial algebra over \mathbb{C} , then the extended concept satisfies the following analogue of Theorem 1.1 ([El 13], Theorem 3.4):

Theorem 1.2. *Let R be a finite-dimensional polynomial algebra over \mathbb{C} , A an \mathbb{A}^3 -fibration over R and W an element of A . Then the following are equivalent:*

- (1) W is a residual variable of A over R .
- (2) $A[Z_1, Z_2, \dots, Z_r] = R[W][T_0, T_1, \dots, T_r, T_{r+1}]$ for some for some indeterminates Z_i 's, T_i 's over R .

Kahoui also observes in [El 13] that if the hypothesis “ R is a polynomial algebra over \mathbb{C} ” is weakened to “ R is any domain containing \mathbb{Q} ”, then the conclusion (2) implies (1) still holds in Theorem 1.2. He remarks ([El 13], Pg. 39) that it is not known whether the converse also holds. Example 3.14 of our paper shows that the converse does not hold in general even when R is a regular factorial affine domain. However, using the techniques of Bhatwadekar-Dutta, we shall show that the converse (i.e., (1) \implies (2) of Theorem 1.2) indeed holds over any Noetherian domain when the module of differentials $\Omega_R(A)$ is stably free over A . More generally, we shall prove (Corollary 3.6, Theorem 3.13, Proposition 3.16):

Main Theorem. Let R be a Noetherian domain and A an \mathbb{A}^n -fibration over R . Let $B = R[W_1, W_2, \dots, W_m]$ be a subring of A such that $A \otimes_R k(P)$ is a polynomial algebra in $n - m$ variables over $B \otimes_R k(P)$ for every prime ideal P of R . Then A is an \mathbb{A}^{n-m} -fibration over B . Moreover, the following statements are equivalent:

- (1) $\Omega_R(A)$ is stably free over A .
- (2) A is a stably polynomial algebra over B .

In [BD93], Bhatwadekar-Dutta also observed the following result ([BD93], Theorem 3.2):

Theorem 1.3. Let R be a Noetherian domain such that either R contains \mathbb{Q} or R is seminormal. Then an element W of $R[X, Y]$ is a residual variable of $R[X, Y]$ over R if and only if W is a variable of $R[X, Y]$.

We shall see that Theorem 1.3 holds if we replace the polynomial ring by an \mathbb{A}^2 -fibration whose module of differentials is stably free (Corollary 3.19).

2. Preliminaries

Throughout the article rings will be commutative with unity. For a ring R , $R^{[n]}$ will denote the polynomial ring in n variables over R . We shall use the notation $A = R^{[n]}$ to mean that A is isomorphic, as an R -algebra, to a polynomial ring in n variables over R .

For a prime ideal P of R , $k(P)$ will denote the residue field R_P/PR_P . A finitely generated flat R -algebra A is said to be an \mathbb{A}^n -fibration over R if $A \otimes_R k(P) = k(P)^{[n]}$ for every prime ideal P of R .

Note that several algebraic geometers use the term “ \mathbb{A}^n -fibration” to mean that $A/\mathfrak{m}A \cong (R/\mathfrak{m}R)^{[n]}$ for almost all maximal ideals \mathfrak{m} of R . We emphasise that we shall use it as defined by Sathaye in [Sat83], where the hypothesis is made on *all* fibre rings.

An R -algebra A is said to be stably polynomial algebra over an R -subalgebra B of A if there exist indeterminates Z_1, Z_2, \dots, Z_r over A and indeterminates T_1, T_2, \dots, T_s over B such that $A[Z_1, Z_2, \dots, Z_r] = B[T_1, T_2, \dots, T_s]$ (as B -algebras). We state below an elementary observation on stably polynomial algebras.

Lemma 2.1. Let A be a stably polynomial algebra over R . Then $\Omega_R(A)$ is a stably free A -module.

Proof. Set $D := A^{[m]} = R^{[n+m]}$ for some m, n and $M := \Omega_R(A) \oplus A^m$. Then $M \otimes_A D \cong (\Omega_R(A) \otimes_A D) \oplus D^m \cong \Omega_R(D) \cong D^{n+m}$ (cf. [Mat80], Example 26.J, Pg. 189). Thus $M \otimes_A D$ is a free D -module. Since A is a retract of $D = A^{[m]}$, it follows that M is a free A -module. Thus $\Omega_R(A)$ is stably free over A . \square

The following structure theorem on affine fibrations is due to T. Asanuma ([Asa87], Theorem 3.4):

Theorem 2.2. Let R be a Noetherian ring and A an \mathbb{A}^r -fibration over R . Then $\Omega_R(A)$ is a projective A -module of rank r and A is an R -subalgebra (up to an isomorphism) of a polynomial ring $R^{[m]}$ for some m such that $A^{[m]} = \text{Sym}_{R^{[m]}}(\Omega_R(A) \otimes_A R^{[m]})$ (as R -algebras). In particular, if $\Omega_R(A)$ is a stably free A -module, then A is a stably polynomial algebra over R .

We shall use the following result by Quillen-Suslin ([Qui76], [Sus76])

Theorem 2.3. If R is a PID, then any finitely generated projective $R^{[n]}$ -module is free.

We record the following result on cancellation by Hamann ([Ham75], Theorem 2.6 and Theorem 2.8).

Theorem 2.4. *Let R be a Noetherian ring such that either R contains \mathbb{Q} or R_{red} is seminormal. Then $R^{[1]}$ is R -invariant, i.e., if an R -algebra A is such that $A^{[m]} = R^{[m+1]}$, then $A = R^{[1]}$.*

The following result was first proved by Kambayashi-Miyayashi in ([KM78], Theorem 1). Since any rank-one projective module over a factorial domain is free, this result can now also be seen to follow from Theorem 2.2 and Theorem 2.4. A more general version of the result is given in ([Dut95], Theorem 3.4).

Theorem 2.5. *Let R be a Noetherian factorial domain and A an \mathbb{A}^1 -fibration over R . Then $A = R^{[1]}$.*

3. Main Theorem

Definition 3.1. Let R be a ring, A an R -algebra, $n \in \mathbb{N}$ and $\underline{W} := (W_1, W_2, \dots, W_m)$ an m -tuple of elements in A which are algebraically independent over R such that $A \otimes_R k(P) = (R[\underline{W}] \otimes_R k(P))^{[n-m]}$ for all $P \in \text{Spec}(R)$. We shall call such an m -tuple \underline{W} to be an m -tuple residual variable of A over R .

Remark 3.2. ([BD93], Example 4.1) provides an example of a residual variable in a polynomial ring which is not a variable. For an example of a residual variable in an affine fibration which is not a polynomial ring, see Example 3.14 or Remark 3.15.

We first observe an elementary result.

Lemma 3.3. *Let R be a ring, A an R -algebra, B_1, B_2 R -subalgebras of A and $B = B_1 \otimes_R B_2$. Suppose that $A \otimes_R k(P) = B \otimes_R k(P)^{[n]}$ for all $P \in \text{Spec}(R)$. Then $A \otimes_{B_2} k(Q) = B \otimes_{B_2} k(Q)^{[n]}$ for all $Q \in \text{Spec}(B_2)$.*

Proof. Fix $Q \in \text{Spec}(B_2)$ and set $P := Q \cap R \in \text{Spec}(R)$. Set $\overline{B}_2 := B_2 \otimes_R k(P)$. Then $B \otimes_{B_2} \overline{B}_2 = B \otimes_R k(P)$ and $A \otimes_{B_2} \overline{B}_2 = A \otimes_R k(P)$ so that

$$A \otimes_{B_2} k(Q) = A \otimes_{B_2} \overline{B}_2 \otimes_{\overline{B}_2} k(Q) = A \otimes_R k(P) \otimes_{\overline{B}_2} k(Q) = (B \otimes_R k(P) \otimes_{\overline{B}_2} k(Q))^{[n]} = (B \otimes_{B_2} \overline{B}_2 \otimes_{\overline{B}_2} k(Q))^{[n]} = (B \otimes_{B_2} k(Q))^{[n]}. \quad \square$$

As a consequence, we have

Remark 3.4. If $(\underline{U}, \underline{V}) := (U_1, U_2, \dots, U_s, V_1, V_2, \dots, V_t)$ is an $(s+t)$ -tuple residual variable of A over R , then \underline{V} is a t -tuple residual variable over $R[\underline{U}]$.

Next we record a result on flatness.

Lemma 3.5. *Let $R \subset B \subset A$ be Noetherian rings such that*

- (i) *A and B are flat over R .*
- (ii) *$A \otimes_R k(P)$ is flat over $B \otimes_R k(P)$ for all $P \in \text{Spec}(R)$.*

Then A is flat over B .

Proof. We shall show that A_Q is flat over $B_{Q \cap B}$ for all $Q \in \text{Spec}(A)$. Fix $Q \in \text{Spec}(A)$ and set $P' := Q \cap B \in \text{Spec}(B)$ and $P = P' \cap R \in \text{Spec}(R)$. Then we have local homomorphisms $R_P \rightarrow B_{P'} \rightarrow A_Q$. As $B_{P'}$ is flat over R_P , to show that A_Q is flat over $B_{P'}$, it is enough to show that $A_Q \otimes_{R_P} k(P)$ is flat over $B_{P'} \otimes_{R_P} k(P)$ (cf. [Mat80], 20.G, Pg. 152).

Since $A \otimes_R k(P)$ is flat over $B \otimes_R k(P)$ and $Q \cap R = P$, we see that $A_Q \otimes_{R_P} k(P)$ is flat over $B \otimes_R k(P)$ and hence $(A_Q \otimes_{R_P} k(P)) \otimes_B B_{P'}$ is flat over $(B \otimes_R k(P)) \otimes_B B_{P'}$. Now

$$(A_Q \otimes_{R_P} k(P)) \otimes_B B_{P'} = (A_Q \otimes_B B_{P'}) \otimes_{R_P} k(P) = (A_Q \otimes_{B_{P'}} B_{P'}) \otimes_{R_P} k(P) = A_Q \otimes_{R_P} k(P)$$

and

$$(B \otimes_R k(P)) \otimes_B B_{P'} = B_{P'} \otimes_R k(P) = B_{P'} \otimes_{R_P} k(P).$$

This shows that $A_Q \otimes_{R_P} k(P)$ is flat over $B_{P'} \otimes_{R_P} k(P)$ and hence A_Q is a flat $B_{P'}$ -algebra. Thus A is a flat B -algebra. \square

From Lemma 3.3 and Lemma 3.5 it follows that if \underline{W} is an m -tuple residual variable of an \mathbb{A}^n -fibration A over a ring R , then A is an \mathbb{A}^{n-m} -fibration over $R[\underline{W}]$. More generally, we have:

Corollary 3.6. *Let $R \subset B \subset A$ be Noetherian rings such that A is an \mathbb{A}^n -fibration over R and B an \mathbb{A}^m -fibration over R with $A \otimes_R k(P) = B \otimes_R k(P)^{[n-m]}$ for all $P \in \text{Spec}(R)$. Then A is an \mathbb{A}^{n-m} -fibration over B .*

Remark 3.7. Let A be an \mathbb{A}^3 -fibration over R and $W \in A$. It was shown in ([El 13], Proposition 3.2) that if $\mathbb{Q} \hookrightarrow R$ and A an \mathbb{A}^2 -fibration over $R[W]$, then W is a residual variable of A over R . The converse was also proved for the case R is a regular affine domain over \mathbb{C} ([El 13], Theorem 3.3). Corollary 3.6 shows that the converse holds for *any* Noetherian domain.

As a consequence of Corollary 3.6 and Theorem 2.5, we see that in an \mathbb{A}^{m+1} -fibration over a Noetherian factorial domain, any m -tuple residual variable is necessarily a variable.

Corollary 3.8. *Let R be a Noetherian factorial domain and A an \mathbb{A}^{m+1} -fibration over R . Then an m -tuple \underline{W} of A is an m -tuple residual variable of A over R if and only if $A = R[\underline{W}]^{[1]} = R^{[m+1]}$.*

Remark 3.9. Note that Corollary 3.8 need not hold for an m -tuple residual variable \underline{W} of an \mathbb{A}^n -fibration A over R when $n - m > 1$. Example 3.14 shows that A may not be even a stably polynomial algebra over $R[\underline{W}]$. The next result shows that if R is a polynomial algebra over a PID, then A happens to be a stably polynomial algebra over $R[\underline{W}]$.

Corollary 3.10. *Let R be a finite-dimensional polynomial algebra over a PID, A an \mathbb{A}^n -fibration over R and \underline{W} an m -tuple residual variable of A over R . Then A is a stably polynomial algebra over $R[\underline{W}]$.*

Proof. By Corollary 3.6, A is an \mathbb{A}^{n-m} -fibration over $R[\underline{W}]$ and hence by ([Asa87], Corollary 3.5), $A^{[r]} = \text{Sym}_{R[\underline{W}]}(M)$ for some $r \in \mathbb{N}$ where M is a finitely generated projective $R[\underline{W}]$ module of rank $n - m + r$. Since R is a polynomial algebra over a PID, by Theorem 2.3, we get that $A^{[r]} = R[\underline{W}]^{[n-m+r]}$, i.e., A is a stably polynomial algebra over $R[\underline{W}]$. \square

Remark 3.11.

- (1) Corollary 3.8 shows that if $n - m = 1$, then we have $A = R[\underline{W}]^{[1]}$ in Corollary 3.10. However, if $n - m > 1$, then an example of Asanuma ([Asa87], Theorem 5.1) shows that we need not have $A = R[\underline{W}]^{[n-m]}$ even in the case R is a PID, $m = 1$ and $n = 3$. In Asanuma's example, $\mathbb{Q} \not\hookrightarrow R$. When $\mathbb{Q} \hookrightarrow R$, it is not known whether, in Corollary 3.10, one can conclude that $A = R[\underline{W}]^{[n-m]}$ even in the case $m = 1$ and $n = 3$. For instance in ([BD94], Example 4.13), W is a residual variable in $A = R[X, Y, Z]$, where R is a DVR containing \mathbb{Q} , and it is not known whether $A = R[W]^{[2]}$.
- (2) The proof of Corollary 3.10 shows that the hypothesis that R is "a finite-dimensional polynomial algebra over a PID" can be replaced by the condition that R is "a regular ring with trivial Grothendieck group".

The following observation on module of differentials of an \mathbb{A}^n -fibration having residual variables is crucial for our main theorem.

Lemma 3.12. *Let R be a Noetherian ring and A an \mathbb{A}^{m+k} -fibration over R . If \underline{W} is an m -tuple residual variable of A over R , then $\Omega_R(A) = \Omega_{R[\underline{W}]}(A) \oplus A^m$. In particular, $\Omega_R(A)$ is a stably free A -module if and only if $\Omega_{R[\underline{W}]}(A)$ is a stably free A -module.*

Proof. By Corollary 3.6, A is an \mathbb{A}^k -fibration over $R[\underline{W}]$ and hence by Theorem 2.2, A is an $R[\underline{W}]$ -subalgebra of a polynomial algebra B over $R[\underline{W}]$ and $\Omega_{R[\underline{W}]}(A)$ is a projective A -module of rank k . Since $R \hookrightarrow R[\underline{W}] \hookrightarrow A \hookrightarrow B$, and since for any A -module M , every R -derivation $d : R[\underline{W}] \rightarrow M$, can be extended to an R -derivation $\tilde{d}|_A : A \rightarrow M$ where $\tilde{d}|_A$ is the restriction of an extension $\tilde{d} : B \rightarrow M$ of d , we have the following split short exact sequence ([Mat80], Theorem 57, p186):

$$0 \longrightarrow A \otimes_{R[\underline{W}]} \Omega_R(R[\underline{W}]) \longrightarrow \Omega_R(A) \longrightarrow \Omega_{R[\underline{W}]}(A) \longrightarrow 0$$

This shows that $\Omega_R(A) = \Omega_{R[\underline{W}]}(A) \oplus A \otimes_{R[\underline{W}]} \Omega_R(R[\underline{W}]) = \Omega_{R[\underline{W}]}(A) \oplus A^m$. \square

We now prove our main result.

Theorem 3.13. *Let R be a Noetherian ring and A an \mathbb{A}^n -fibration over R such that $\Omega_R(A)$ is a stably free A -module. Suppose \underline{W} is an m -tuple residual variable of A over R . Then A is a stably polynomial algebra over $R[\underline{W}]$; specifically, $A^{[\ell]} = R[\underline{W}]^{[n-m+\ell]}$ for some $\ell \in \mathbb{N}$.*

Proof. By Corollary 3.6, A is an \mathbb{A}^{n-m} -fibration over $R[\underline{W}]$ and hence, by Lemma 3.12, $\Omega_{R[\underline{W}]}(A)$ is a stably free A -module. Therefore, we get the result by Theorem 2.2. \square

The following example shows the necessity of the assumption “ $\Omega_R(A)$ is a stably free A -module” in Theorem 3.13 even when R is a regular factorial affine domain over the field of real numbers.

Example 3.14. Let $R = \mathbb{R}[X, Y, Z]/(X^2 + Y^2 + Z^2 - 1)$. It is well-known that $K_0(R)$, the Grothendieck group of R , is non-trivial (in fact, it is $\mathbb{Z} \oplus \mathbb{Z}/(2)$); in particular, there exists a finitely generated projective R -module M of rank 2 which is not stably free. Let W be an indeterminate over R and $A = \text{Sym}_R(M \oplus RW)$. Then $A_P = R_P[W]^{[2]}$ for all $P \in \text{Spec}(R)$ so that A is an \mathbb{A}^3 -fibration over R with W as a residual variable. If $A^{[\ell]} = R[W]^{[\ell+2]}$, then we would have $\text{Sym}_R(M \oplus RW \oplus R^\ell) \cong \text{Sym}_R(RW \oplus R^{\ell+2})$ and hence, by ([EH73], Lemma 1.3), we would have $M \oplus RW \oplus R^\ell \cong RW \oplus R^{\ell+2}$ contradicting that M is not stably free.

Remark 3.15. When R is a Noetherian factorial domain and $m = n - 1$ then Corollary 3.8 shows that the hypothesis “ $\Omega_R(A)$ is stably free” may be dropped from Theorem 3.13; in fact, in this case, $A = R[\underline{W}]^{[1]}$. But even over a (non-factorial) Dedekind domain and even for $n = 2$ and $m = 1$, an \mathbb{A}^2 -fibration need not be stably polynomial over $R[W]$ when W is a residual variable of A . For instance, choose a non-principal ideal I of the Dedekind domain R and set $A = \text{Sym}_R(I \oplus RW)$. As in Example 3.14, A is an \mathbb{A}^2 -fibration over R , W is a residual variable of A but A is not stably polynomial over $R[W]$.

The following result gives a converse of Theorem 3.13.

Proposition 3.16. *Let R be a Noetherian ring and A an \mathbb{A}^n -fibration over R . Suppose that there exists an m -tuple residual variable \underline{W} of A over R such that A is a stably polynomial algebra over $R[\underline{W}]$. Then $\Omega_R(A)$ is a stably free A -module.*

Proof. By Lemma 3.12, it suffices to show that $\Omega_{R[\underline{W}]}(A)$ is a stably free A -module. This follows from Lemma 2.1. \square

For convenience, we state below an easy result.

Lemma 3.17. *Let $R \hookrightarrow B \hookrightarrow A$ be integral domains such that A is an \mathbb{A}^n -fibration over B . Then $A \otimes_R k(P)$ is an \mathbb{A}^n -fibration over $B \otimes_R k(P)$ for every $P \in \text{Spec}(R)$. Moreover, if $B = R^{[m]}$, then $A \otimes_R k(P)$ is a stably polynomial algebra over $B \otimes_R k(P)$ for each $P \in \text{Spec}(R)$.*

Proof. Let $P \in \text{Spec}(R)$, $\overline{A} = A \otimes_R k(P)$ and $\overline{B} = B \otimes_R k(P)$. Since A is a finitely generated flat B -algebra, clearly \overline{A} is a finitely generated flat \overline{B} -algebra. Note that $\overline{A} = A \otimes_R k(P) = A \otimes_B B \otimes_R k(P) = A \otimes_B \overline{B}$. Let $\overline{Q} \in \text{Spec}(\overline{B})$ and let Q be the prime ideal of B for which $Q\overline{B} = \overline{Q}$. Then $Q \cap R = P$. Now $k(\overline{Q})$ is the field of fractions of B/Q and hence of $B_P/QB_P \cong \overline{B}/\overline{Q}$; thus $k(\overline{Q}) = k(Q)$. Hence, $\overline{A} \otimes_{\overline{B}} k(\overline{Q}) = A \otimes_B \overline{B} \otimes_{\overline{B}} k(\overline{Q}) = A \otimes_B k(Q) = k(Q)^{[m]} = k(\overline{Q})^{[m]}$. \square

Corollary 3.18. *Let R be a Noetherian domain and A an \mathbb{A}^{m+1} -fibration over R . Then an m -tuple \underline{W} of A is an m -tuple residual variable of A over R if and only if A is an \mathbb{A}^1 -fibration over $R[\underline{W}]$.*

Proof. Follows from Corollary 3.6, Lemma 3.17 and Corollary 3.8. \square

Finally, we give a condition for an m -tuple residual variable in an \mathbb{A}^{m+1} -fibration to be a variable.

Corollary 3.19. *Let R be a Noetherian domain and A an \mathbb{A}^{m+1} -fibration over R such that $\Omega_R(A)$ is a stably free A -module. Suppose that either R contains \mathbb{Q} or R is seminormal. Then for an m -tuple \underline{W} of A , the following are equivalent:*

- (I) \underline{W} is an m -tuple residual variable of A over R .
- (II) $A = R[\underline{W}]^{[1]} = R^{[m+1]}$.

Proof. Follows from Theorem 3.13 and Theorem 2.4. \square

4. Appendix

Theorem 3.13 can be slightly generalised when envisaged as a statement on “tensor product decomposition” discussed in [BW77]. Let $R \subset B \subset A$ be Noetherian rings with A an \mathbb{A}^n -fibration over R and also an \mathbb{A}^{n-m} -fibration over B . Theorem 3.13 shows that if $B = R^{[m]}$ and $\Omega_R(A)$ is stably free then $A^{[\ell]} = B^{[\ell]} \otimes_{R^{[\ell]}} C$, where $C = R^{[n-m+\ell]}$. One can see below (Proposition 4.2) that even when B is only an \mathbb{A}^m -fibration over R , it is stably a factor in a tensor product decomposition of A , even without the hypothesis that $\Omega_R(A)$ is stably free. We first note that the proof of Lemma 3.12 can be seen to yield the following general version:

Lemma 4.1. *Let $R \subset B \subset A$ be Noetherian rings such that A is an \mathbb{A}^n -fibration over R and B an \mathbb{A}^m -fibration over R with $A \otimes_R k(P) = B \otimes_R k(P)^{[n-m]}$ for all $P \in \text{Spec}(R)$. Then $\Omega_R(A) = \Omega_B(A) \oplus (\Omega_R(B) \otimes_B A)$.*

As a consequence we have

Proposition 4.2. *Under the hypothesis of Lemma 4.1, there exists $\ell \geq 0$ such that $A^{[\ell]} = C \otimes_{R^{[\ell]}} B^{[\ell]}$ for some $R^{[\ell]}$ -algebra C .*

Proof. Since A and B are affine fibrations over R , by Theorem 2.2, we can choose a sufficiently large positive integer ℓ such that

$$A^{[\ell]} = \text{Sym}_{R^{[\ell]}}(\Omega_R(A) \otimes_A R^{[\ell]}) \quad \text{and} \quad B^{[\ell]} = \text{Sym}_{R^{[\ell]}}(\Omega_R(B) \otimes_B R^{[\ell]}).$$

Therefore, by Lemma 4.1, we have

$$\begin{aligned} A^{[\ell]} &= \text{Sym}_{R^{[\ell]}}(\Omega_R(A) \otimes_A R^{[\ell]}) \\ &= \text{Sym}_{R^{[\ell]}}((\Omega_B(A) \oplus (\Omega_R(B) \otimes_B A)) \otimes_A R^{[\ell]}) \\ &= \text{Sym}_{R^{[\ell]}}(\Omega_B(A) \otimes_A R^{[\ell]}) \otimes_{R^{[\ell]}} \text{Sym}_{R^{[\ell]}}((\Omega_R(B) \otimes_B A) \otimes_A R^{[\ell]}) \\ &= \text{Sym}_{R^{[\ell]}}(\Omega_B(A) \otimes_A R^{[\ell]}) \otimes_{R^{[\ell]}} \text{Sym}_{R^{[\ell]}}(\Omega_R(B) \otimes_B R^{[\ell]}) \\ &= \text{Sym}_{R^{[\ell]}}(\Omega_B(A) \otimes_A R^{[\ell]}) \otimes_{R^{[\ell]}} B^{[\ell]} \\ &= C \otimes_{R^{[\ell]}} B^{[\ell]}, \end{aligned}$$

where $C = \text{Sym}_{R^{[\ell]}}(\Omega_B(A) \otimes_A R^{[\ell]})$. □

Acknowledgements: The authors thank Neena Gupta for carefully going through the draft and the referee for suggestions which have been incorporated in the Appendix.

References

- [Asa87] Teruo Asanuma, *Polynomial fibre rings of algebras over Noetherian rings*, Invent. Math. **87** (1987), no. 1, 101–127.
- [BD93] S. M. Bhatwadekar and Amartya K. Dutta, *On residual variables and stably polynomial algebras*, Comm. Algebra **21** (1993), no. 2, 635–645.
- [BD94] ———, *On affine fibrations*, Commutative algebra (Trieste, 1992), World Sci. Publ., River Edge, NJ, 1994, pp. 1–17.
- [BW77] H. Bass and D. Wright, *Localisation in the K-theory of invertible algebras*, J. Pure Appl. Algebra **9** (1976/77), no. 1, 89–105.
- [Dut95] Amartya K. Dutta, *On \mathbb{A}^1 -bundles of affine morphisms*, J. Math. Kyoto Univ. **35** (1995), no. 3, 377–385.
- [EH73] Paul Eakin and William Heinzer, *A cancellation problem for rings*, Conference on Commutative Algebra (Univ. Kansas, Lawrence, Kan., 1972), Springer, Berlin, 1973, pp. 61–77. Lecture Notes in Math., Vol. 311.
- [El 13] M’hammed El Kahoui, *On residual coordinates and stable coordinates of $R^{[3]}$* , Arch. Math. (Basel) **100** (2013), no. 1, 35–41.
- [Ham75] Eloise Hamann, *On the R-invariance of $R[X]$* , J. Algebra **35** (1975), 1–16.
- [KM78] T. Kambayashi and M. Miyanishi, *On flat fibrations by the affine line*, Illinois J. Math. **22** (1978), no. 4, 662–671.
- [Mat80] Hideyuki Matsumura, *Commutative algebra*, second ed., Mathematics Lecture Note Series, vol. 56, Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980.
- [Qui76] Daniel Quillen, *Projective modules over polynomial rings*, Invent. Math. **36** (1976), 167–171.
- [Sat83] A. Sathaye, *Polynomial ring in two variables over a DVR: a criterion*, Invent. Math. **74** (1983), no. 1, 159–168.
- [Sus76] A. A. Suslin, *Projective modules over polynomial rings are free*, Dokl. Akad. Nauk SSSR **229** (1976), no. 5, 1063–1066.