

BOUNDED RICCI CURVATURE AND POSITIVE SCALAR CURVATURE UNDER SINGULAR RICCI DE TURCK FLOW

KLAUS KRÖNCKE, TOBIAS MARXEN, AND BORIS VERTMAN

ABSTRACT. In this paper we consider a Ricci de Turck flow of spaces with isolated conical singularities, which preserves the conical structure along the flow. We establish that a given initial regularity of Ricci curvature is preserved along the flow. Moreover under additional assumptions, positivity of scalar curvature is preserved under such a flow, mirroring the standard property of Ricci flow on compact manifolds. The analytic difficulty is the a priori low regularity of scalar curvature at the conical tip along the flow, so that the maximum principle does not apply. We view this work as a first step toward studying positivity of the curvature operator along the singular Ricci flow.

CONTENTS

1. Introduction and statement of main results	1
2. Geometric preliminaries on conical manifolds	3
3. Weighted Hölder spaces on conical manifolds	4
4. Lichnerowicz Laplacian and tangential stability	6
5. Existence and regularity of singular Ricci de Turck flow	9
6. Evolution of curvatures under Ricci de Turck flow	10
7. Regularity of Ricci curvature along the Ricci de Turck flow	14
8. Positivity of scalar curvature along the Ricci de Turck flow	24
Appendix: Characterizing the spectral conditions	25
References	32

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Consider a compact smooth Riemannian manifold (M, g) without boundary. Its Ricci flow is a smooth family of metrics $(g(t))_{t \geq 0}$ such that

$$\partial_t g(t) = -2\text{Ric}(g(t)), \quad g(0) = g, \quad (1.1)$$

where $\text{Ric}(g(t))$ denotes the Ricci curvature tensor of $g(t)$. Due to diffeomorphism invariance of the Ricci tensor, this evolution equation fails to be strongly parabolic. One overcomes this problem by adding an additional term to the equation

Date: February 18, 2020.

2010 Mathematics Subject Classification. Primary 53C44; Secondary 53C25; 58J35.

Key words and phrases. Ricci flow, positive scalar curvature, conical singularities.

Partial support by DFG Priority Programme "Geometry at Infinity".

which brakes the diffeomorphism invariance. This leads to the equivalent and analytically more convenient Ricci de Turck flow

$$\partial_t g(t) = -2\text{Ric}(g(t)) + \mathcal{L}_{W(t)}g(t), \quad g(0) = g, \quad (1.2)$$

where $W(t)$ is the de Turck vector field defined in terms of the Christoffel symbols for the metrics $g(t)$ and a background metric h

$$W(t)^k = g(t)^{ij} (\Gamma_{ij}^k(g(t)) - \Gamma_{ij}^k(h)). \quad (1.3)$$

In the applications, the background metric h is usually taken as the initial metric g , or as its small Ricci flat perturbation. The de Turck vector field defines a one-parameter family of diffeomorphisms $(\phi(t))_{t \geq 0}$ and if $(g(t))_{t \geq 0}$ is a solution to the Ricci de Turck flow (1.2), then the pullback $(\phi(t)^*g(t))_{t \geq 0}$ is a solution to the Ricci flow (1.1).

On singular spaces, the de Turck vector field may point towards the singular stratum and thus lengths of the corresponding integral curves may not be bounded from below away from zero. Hence the one-parameter family of diffeomorphisms $(\phi(t))_{t \geq 0}$ may not exist for positive times. Therefore, in the singular setting (1.1) and (1.2) are generally not equivalent and we study the latter flow.

Our work establishes an interesting property of the flow, namely that a given initial regularity of the Ricci curvature is preserved along the flow.

Theorem 1.1. *The singular Ricci de Turck flow preserves the initial regularity of the Ricci curvature. In particular, if the initial metric has bounded Ricci curvature, the flow remains of bounded Ricci curvature as well.*

We are not able to deduce an analogous result for the scalar curvature. The reason is that the norm of the Ricci tensor appears as the reaction term in the evolution equation of the scalar curvature. Thus, unbounded Ricci curvature at the singularity pushes the scalar curvature to infinity after infinitesimal time. In contrast, the evolution equation on the Ricci curvature is tensorial which allows more flexibility. However, we are able to prove a different property of the scalar curvature along the Ricci flow which is well known in the smooth compact case.

Theorem 1.2. *An admissible Riemannian manifold with isolated conical singularities and positive scalar curvature admits a singular Ricci de Turck flow preserving the singular structure and under the additional assumption of strong tangential stability, positivity of scalar curvature along the flow.*

The present work is a continuation of a research program on singular Ricci flow, that preserves the initial singular structure, that has seen several recent advances. In the setting of surfaces with conical singularities, singular Ricci flow has been studied by Mazzeo, Rubinstein and Sesum in [MRS15] and Yin [Yin10]. The Yamabe flow, which coincides with the Ricci flow in the two-dimensional setting, has been studied in general dimension on spaces with edge singularities by Bahuaud and the third named author in [BAVe14] and [BAVe16]. In the setting of Kähler manifolds, Kähler-Ricci flow on spaces with edge singularities appears in the recent results on the Calabi-Yau conjecture on Fano manifolds, cf. Donaldson [DON12], Tian [Tia15], see also Jeffres, Mazzeo and Rubinstein [JMR16].

Kähler-Ricci flow in case of isolated conical singularities has been addressed by Chen and Wang [CHW15], Wang [WAN16], as well as Liu and Zhang [LiZh17].

Ricci flow on singular spaces of general dimension, without the Kähler condition, does not reduce to a scalar equation and has been studied in [VER16] by the third named author. Subsequently, the first and the third named authors in [KRVE19A, KRVE19B] established stability and studied solitons and Perelman entropies of this flow in the special case of isolated conical singularities. The present work is a continuation of this research program.

Let us point out that Ricci flow preserving the initial singular structure, is not the only possible way to evolve a singular metric. In fact, Giesen and Topping [GT10, GT11] construct a solution to the Ricci flow on surfaces with singularities, which becomes instantaneously complete. Alternatively, Simon [SIM13] constructs Ricci flow in dimension two and three that smoothes out the singularity.

2. GEOMETRIC PRELIMINARIES ON CONICAL MANIFOLDS

We begin with a definition of spaces with isolated conical singularities. We point out that part of our analysis in fact applies to spaces with non-isolated conical singularities, the so-called edges.

Definition 2.1. *Let M be the open interior of a compact smooth manifold \overline{M} with boundary $F := \partial M$. Let x be a boundary defining function and $\mathcal{C}(F)$ a tubular neighborhood of the boundary with open interior $\mathcal{C}(F) := (0, 1)_x \times F$. An incomplete Riemannian metric g on M with an isolated conical singularity is a smooth metric on M satisfying*

$$g \upharpoonright \mathcal{C}(F) = dx^2 + x^2 g_F + h =: \overline{g} + h,$$

where the higher order term h has the following asymptotics at $x = 0$. Let $\overline{g} = dx^2 + x^2 g_F$ denote the exact conical part of the metric g over $\mathcal{C}(F)$ and $\nabla_{\overline{g}}$ the corresponding Levi Civita connection. Then we require that for some $\gamma > 0$ and all integer $k \in \mathbb{N}_0$ the pointwise norm

$$|x^k \nabla_{\overline{g}}^k h|_{\overline{g}} = O(x^\gamma), \quad x \rightarrow 0. \quad (2.1)$$

Remark 2.2. *We emphasize here that we do not assume that the higher order term h is smooth up to $x = 0$ and do not restrict the order $\gamma > 0$ to be integer. In that sense the notion of conical singularities in the present discussion is more general than the classical notion of conical singularities where h is usually assumed to be smooth up to $x = 0$ with $\gamma = 1$. This minor generalization is necessary, since the Ricci de Turck flow, which will be introduced below, preserves a conical singularity only up to a higher order term h as above.*

We call (M, g) a manifold with an isolated conical singularity, or a *conical manifold* for short. The definition extends directly to conical manifolds with finitely many isolated conical singularities. Since the analytic arguments are local in nature, we may assume without loss of generality that M has a single conical singularity only.

Let $(z) = (z_1, \dots, z_n)$ be local coordinates on F , where $n = \dim F$. Then (x, z) are local coordinates on the conical neighborhood $\mathcal{C}(F) \subset M$. A b -vector field is by definition a smooth vector field on \overline{M} which is tangent to the boundary $\partial M = F$. The b -vector fields form a Lie algebra, denoted by \mathcal{V}_b . In the local coordinates (x, z) , the algebra \mathcal{V}_b is locally, near ∂M , generated by

$$\left\{ x \frac{\partial}{\partial x}, \partial_z = \left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n} \right) \right\},$$

with smooth coefficients on \overline{M} . The b -tangent bundle ${}^b\text{TM}$ is defined by requiring that the b -vector fields form a spanning set of sections for it, i.e. $\mathcal{V}_b = C^\infty(\overline{M}, {}^b\text{TM})$. The b -cotangent bundle ${}^bT^*M$ is the dual bundle and locally, near ∂M , generated by the one-forms

$$\left\{ \frac{dx}{x}, dz_1, \dots, dz_n \right\}. \quad (2.2)$$

Note that the differential form $\frac{dx}{x}$ is singular in the usual sense, but smooth as a section of ${}^bT^*M$. Extend the radial function x of the cone $\mathcal{C}(F)$ to a nowhere vanishing smooth function $x : \overline{M} \rightarrow [0, 2]$. Then we define the incomplete b -tangent space ${}^{ib}\text{TM}$ by asking that $C^\infty(\overline{M}, {}^{ib}\text{TM}) := x^{-1} C^\infty(\overline{M}, {}^b\text{TM})$. The dual bundle, the incomplete b -cotangent bundle ${}^{ib}T^*M$, is related to its complete counterpart ${}^bT^*M$ by $C^\infty(\overline{M}, {}^{ib}T^*M) = x C^\infty(\overline{M}, {}^bT^*M)$, and is locally generated by

$$\{dx, x dz_1, \dots, x dz_n\}. \quad (2.3)$$

3. WEIGHTED HÖLDER SPACES ON CONICAL MANIFOLDS

In this section we review definitions from [VER16, Section 1.3] in the case of isolated conical singularities. Let (M, g) be a manifold with isolated conical singularities.

Definition 3.1. Let $d_M(p, p')$ denote the distance between two points $p, p' \in M$ with respect to the conical metric g . In terms of the local coordinates (x, z) over the singular neighborhood $\mathcal{C}(F)$, the distance can be estimated up to a constant uniformly from above and below by the distance function of the model cone

$$d((x, z), (x', z')) = (|x - x'|^2 + (x + x')^2 |z - z'|^2)^{\frac{1}{2}}.$$

The Hölder space $C_{\text{ie}}^\alpha(M \times [0, T])$, $\alpha \in [0, 1)$, is by definition the set of functions $u(p, t)$ that are continuous on $\overline{M} \times [0, T]$ with finite α -th Hölder norm

$$\|u\|_\alpha := \|u\|_\infty + \sup \left(\frac{|u(p, t) - u(p', t')|}{d_M(p, p')^\alpha + |t - t'|^{\alpha/2}} \right) < \infty, \quad (3.1)$$

with supremum taken over all $p, p' \in M$ and $t, t' \in [0, T]$ with $p \neq p'$ and $t \neq t'$.¹

We now extend the notion of Hölder spaces to sections of the vector bundle $S := \text{Sym}^2({}^{ib}T^*M)$ of symmetric 2-tensors. Note that the Riemannian metric g induces a fibrewise inner product on S , which we also denote by g .

¹We can assume without loss of generality that the tuples (p, p') are always taken from within the same coordinate patch of a given atlas.

Definition 3.2. The Hölder space $\mathcal{C}_{ie}^\alpha(M \times [0, T], S)$ is by definition the set of all sections ω of S which are continuous on $\overline{M} \times [0, T]$, such that for any local orthonormal frame $\{s_j\}$ of S , the scalar functions $g(\omega, s_j)$ are in $\mathcal{C}_{ie}^\alpha(M \times [0, T])$.

The α -th Hölder norm of ω is defined using a partition of unity $\{\phi_j\}_{j \in J}$ subordinate to a cover of local trivializations of S , with a local orthonormal frame $\{s_{jk}\}$ over $\text{supp}(\phi_j)$ for each $j \in J$. We put

$$\|\omega\|_\alpha^{(\phi, s)} := \sum_{j \in J} \sum_k \|g(\phi_j \omega, s_{jk})\|_\alpha. \quad (3.2)$$

Different choices of $(\{\phi_j\}, \{s_{jk}\})$ lead to equivalent norms so that we may drop the upper index (ϕ, s) from notation. Next we come to weighted and higher order Hölder spaces of S -sections.

Definition 3.3. (1) The (hybrid) weighted Hölder space for $\gamma \in \mathbb{R}$ is

$$\mathcal{C}_{ie}^\alpha(M \times [0, T], S)_\gamma := x^\gamma \mathcal{C}_{ie}^\alpha(M \times [0, T], S) \cap x^{\gamma+\alpha} \mathcal{C}_{ie}^0(M \times [0, T], S)$$

with Hölder norm $\|\omega\|_{\alpha, \gamma} := \|x^{-\gamma} \omega\|_\alpha + \|x^{-\gamma-\alpha} \omega\|_\infty$.

(2) The weighted higher order Hölder spaces are defined by

$$\begin{aligned} \mathcal{C}_{ie}^{k, \alpha}(M \times [0, T], S)_\gamma &:= \{\omega \in \mathcal{C}_{ie}^\alpha \mid \{\mathcal{V}_b^j \circ (x^2 \partial_t)^l\} \omega \in \mathcal{C}_{ie}^\alpha \\ &\text{for all } j + 2l \leq k\}, \end{aligned}$$

for any $\gamma \in \mathbb{R}$ and $k \in \mathbb{N}$. For any $\gamma > -\alpha$ and $k \in \mathbb{N}$ we also define

$$\begin{aligned} \mathcal{C}_{ie}^{k, \alpha}(M \times [0, T], S)_\gamma^b &:= \{\omega \in \mathcal{C}_{ie}^\alpha \mid \{\mathcal{V}_b^j \circ (x^2 \partial_t)^l\} \omega \in \mathcal{C}_{ie}^\alpha \\ &\text{for all } j + 2l \leq k\}. \end{aligned}$$

The corresponding Hölder norms are defined using a finite cover of coordinate charts trivializing S_0 and a subordinate partition of unity $\{\phi_j\}_{j \in J}$. By a slight abuse of notation, we identify \mathcal{V}_b with its finite family of generators over each coordinate chart. Writing $\mathcal{D} := \{\mathcal{V}_b^j \circ (x^2 \partial_t)^l \mid j + 2l \leq k\}$ the Hölder norm on $\mathcal{C}_{ie}^{k, \alpha}(M \times [0, T], S)_\gamma$ is then given by

$$\|\omega\|_{k+\alpha, \gamma} = \sum_{j \in J} \sum_{X \in \mathcal{D}} \|X(\phi_j \omega)\|_{\alpha, \gamma} + \|\omega\|_{\alpha, \gamma}. \quad (3.3)$$

For the Hölder norm on $\mathcal{C}_{ie}^{k, \alpha}(M \times [0, T], S)_\gamma^b$ replace in (3.3) $\|\omega\|_{\alpha, \gamma}$ by $\|\omega\|_\alpha$.

The Hölder norms for different choices of coordinate charts, the subordinate partition of unity or vector fields \mathcal{V}_b are equivalent. Analogously we also consider time-independent Hölder spaces, which are denoted in the same way with $[0, T]$ deleted from notation above.

The vector bundle S decomposes into a direct sum of sub-bundles $S = S_0 \oplus S_1$, where the sub-bundle $S_0 = \text{Sym}_0^2(\text{ib} T^* M)$ is the space of trace-free (with respect to the fixed metric g) symmetric 2-tensors, and S_1 is the space of pure trace (with respect to the fixed metric g) symmetric 2-tensors. Definition 3.3 extends verbatim to subbundles S_0 and S_1 .

Remark 3.4. *The spaces presented here are slightly different from the spaces originally introduced in [VER16]. There, in case of S_1 -sections, higher order weighted Hölder spaces were defined in terms of $x^\gamma C_{ie}^\alpha$ instead of $C_{ie,\gamma}^\alpha$. Here, we present a more unified definition, which will become much more convenient below. The arguments of [VER16] still carry over to yield regularity statements in these unified Hölder spaces.*

Definition 3.5. *Let (M, g) be a compact conical manifold, $\gamma_0, \gamma_1 \in \mathbb{R}$. In order to simplify notation, we set $C_{ie}^{k,\alpha}(M \times [0, T], S)_\gamma^b := C_{ie}^{k,\alpha}(M \times [0, T], S)_\gamma^b$ for $\gamma \leq -\alpha$, with the same convention for S_1 as well. Then we define the following spaces.*

(1) *If (M, g) is not an orbifold, we set*

$$\mathcal{H}_{\gamma_0, \gamma_1}^{k, \alpha}(M \times [0, T], S) := C_{ie}^{k, \alpha}(M \times [0, T], S_0)_{\gamma_0} \oplus C_{ie}^{k, \alpha}(M \times [0, T], S_1)_{\gamma_1}^b.$$

(2) *If (M, g) is an orbifold, we set*

$$\mathcal{H}_{\gamma_0, \gamma_1}^{k, \alpha}(M \times [0, T], S) := C_{ie}^{k, \alpha}(M \times [0, T], S_0)_{\gamma_0}^b \oplus C_{ie}^{k, \alpha}(M \times [0, T], S_1)_{\gamma_1}^b.$$

Note that e.g. for the non-orbifold case, the different choice of spaces for the S_0 and S_1 components, simply ensures that the S_1 component is not (!) included in $x^\gamma C_{ie}^0$ for some positive weight γ . In case of orbifold singularities, this restriction is imposed on both the S_0 and S_1 components. We can now impose regularity assumptions on our initial data (M, g) . Definitions 3.2 and 3.3 extend naturally to associated bundles of ${}^{\text{ib}}\text{TM}$. Also we write $\mathcal{H}_\gamma^{k, \alpha} \equiv \mathcal{H}_{\gamma, \gamma}^{k, \alpha}$.

Definition 3.6. *Let $\alpha \in [0, 1)$, $k \in \mathbb{N}_0$ and $\gamma > 0$. A conical manifold (M, g) is (α, γ, k) Hölder regular if the following two conditions are satisfied:*

- (i) *The $(0, 4)$ curvature tensor $\text{Rm} \in C_{ie}^{k, \alpha}(M \times [0, T], \otimes^4 {}^{\text{ib}}\text{T}^*M)_{-2}$.*
- (ii) *The Ricci curvature tensor $\text{Ric} \in \mathcal{H}_{-2+\gamma}^{k, \alpha}(M \times [0, T], S)$.*

We continue under that assumption from now on.

Remark 3.7. *The asymptotics of the Ricci curvature tensor, as a section of S , is generically $O(x^{-2})$ as $x \rightarrow 0$. Hence (α, γ, k) Hölder regularity with $\gamma > 0$ in particular implies that the exact conical part $(\mathcal{C}(F), \bar{g})$ must be Ricci-flat. This is equivalent to (F^n, g_F) being Einstein with Einstein constant $n - 1$. Moreover, the weight γ corresponds to the weight in (2.1).*

4. LICHNEROWICZ LAPLACIAN AND TANGENTIAL STABILITY

Consider the right hand side $-2\text{Ric}(g(t)) + \mathcal{L}_{W(t)}g(t)$ of the Ricci de Turck flow equation (1.2). Write $W(t) = W(g(t), h)$ to indicate precisely what the de Turck vector field depends on. Choose $h = g$ as the background metric. Then, replacing $g(t) = g + s\omega$ for some symmetric 2-tensor ω , the linearization of right hand side of (1.2) is given by

$$\frac{d}{ds} \left(-2\text{Ric}(g + s\omega) + \mathcal{L}_{W(g+s\omega, g)}(g + s\omega) \right) |_{s=0} = -\Delta_L \omega, \quad (4.1)$$

where Δ_L is an elliptic operator, which is known as the Lichnerowicz Laplacian of g , acting on symmetric 2-tensors by

$$\Delta_L \omega_{ij} = \Delta \omega_{ij} - 2g^{pq} \text{Rm}_{qij}^r \omega_{rp} + g^{pq} R_{ip} \omega_{qj} + g^{pq} R_{jp} \omega_{iq},$$

where Δ denotes the rough Laplacian, and Rm_{qij}^r and R_{ij} denote the components of the $(1, 3)$ -Riemann curvature tensor and the Ricci tensor, respectively. Near the conical singularity Δ_L can be written as follows. We choose local coordinates (x, z) over the singular neighborhood $\mathcal{C}(F) = (0, 1)_x \times F$. Consider a decomposition of compactly supported smooth sections $C_0^\infty(\mathcal{C}(F), S \upharpoonright \mathcal{C}(F))$

$$\begin{aligned} C_0^\infty(\mathcal{C}(F), S \upharpoonright \mathcal{C}(F)) &\rightarrow C_0^\infty((0, 1), C^\infty(F) \times \Omega^1(F) \times \text{Sym}^2(T^*F)), \\ \omega &\mapsto (\omega(\partial_x, \partial_x), \omega(\partial_x, \cdot), \omega(\cdot, \cdot)), \end{aligned} \quad (4.2)$$

where $\Omega^1(F)$ denotes differential 1-forms on F . Under such a decomposition, the Lichnerowicz Laplace operator Δ_L associated to the singular Riemannian metric g attains the following form over $\mathcal{C}(F)$

$$\Delta_L = -\frac{\partial^2}{\partial x^2} - \frac{n}{x} \frac{\partial}{\partial x} + \frac{\square_L}{x^2} + \mathcal{O}, \quad (4.3)$$

where \square_L is a differential operator on $C^\infty(F) \times \Omega^1(F) \times \text{Sym}^2(T^*F)$, depending only on the exact conical part \bar{g} , and the higher order term is $\mathcal{O} \in x^{-2+\gamma} \mathcal{V}_b^2$ with Hölder regular coefficients.

4.1. Tangential stability.

We can now impose a central analytic condition, under which existence of singular Ricci de Turck flow has been established in [VER16].

Definition 4.1. (F, g_F) is called (strictly) *tangentially stable* if the tangential operator \square_L of the Lichnerowicz Laplacian on its cone restricted to tracefree tensors is non-negative (resp. strictly positive).

Tangential stability (in fact a smaller lower bound $\square_L \geq -((n-1)/2)^2$ would suffice) has a straightforward implication: for $\omega \in C_0^\infty(\mathcal{C}(F), S)$ we find

$$\begin{aligned} \Delta_L \omega &= \left(\frac{\partial}{\partial x} + \frac{1}{x} \left(\sqrt{\square_L + \left(\frac{n-1}{2} \right)^2} + \frac{n-1}{2} \right) \right)^t \\ &\quad \circ \left(\frac{\partial}{\partial x} + \frac{1}{x} \left(\sqrt{\square_L + \left(\frac{n-1}{2} \right)^2} + \frac{n-1}{2} \right) \right) \omega. \end{aligned} \quad (4.4)$$

Thus, tangential stability in particular implies that Δ_L , acting on compactly supported smooth sections, has a lower bound, i.e. there exists some $C \in \mathbb{R}$ such that

$$\Delta_L \upharpoonright C_0^\infty(M, S) \geq C. \quad (4.5)$$

For convenience of the reader we shall add here a complete characterization of tangential stability, obtained by the first and third named authors in [KRVE19A]. Recall that by Remark 3.7, the assumption of (α, γ, k) Hölder regularity implies that (F, g_F) is Einstein with Einstein constant $(n-1)$.

Theorem 4.2. *Let (F, g_F) , $n \geq 3$ be a compact Einstein manifold with constant $(n-1)$. We write Δ_E for its Einstein operator, and denote the Laplace Beltrami operator by Δ . Then (F, g_F) is tangentially stable if and only if $\text{Spec}(\Delta_E|_{\mathbb{T}}) \geq 0$ and $\text{Spec}(\Delta) \setminus$*

$\{0\} \cap (n, 2(n+1)) = \emptyset$. Similarly, (M, g) is strictly tangentially stable if and only if $\text{Spec}(\Delta_E|_{\text{TT}}) > 0$ and $\text{Spec}(\Delta) \setminus \{0\} \cap [n, 2(n+1)] = \emptyset$.

There are plenty of examples, where (strict) tangential stability is satisfied. Any spherical space form is tangentially stable. [KrVe19a] also provides a detailed list of strict tangentially stable Einstein manifolds that are symmetric spaces. Note that \mathbb{S}^n is tangentially stable but not strictly tangentially stable.

Theorem 4.3. *Let (F^n, g_F) , $n \geq 2$ be a closed Einstein manifold with constant $(n-1)$, which is a symmetric space of compact type. If it is a simple Lie group G , it is strictly tangentially stable if G is one of the following spaces:*

$$\text{Spin}(p) \ (p \geq 6, p \neq 7), \quad E_6, \quad E_7, \quad E_8, \quad F_4. \quad (4.6)$$

If (F, g_F) is a symmetric space of compact type G/K , it is strictly tangentially stable if it is one of the real Grassmannians

$$\frac{\text{SO}(p+q)}{\text{SO}(p) \times \text{SO}(q)} \quad p \geq 3, \quad q \geq 3, \quad (p, q) \neq (3, 3), \quad (4.7)$$

or one of the following spaces:

$$\begin{aligned} &E_6/[\text{Sp}(4)/\{\pm I\}], \quad E_6/\text{SU}(2) \cdot \text{SU}(6), \quad E_7/[\text{SU}(8)/\{\pm I\}] \\ &E_7/\text{SO}(12) \cdot \text{SU}(2), \quad E_8/\text{SO}(16), \quad E_8/E_7 \cdot \text{SU}(2), \quad F_4/\text{Sp}(3) \cdot \text{SU}(2). \end{aligned} \quad (4.8)$$

Remark 4.4. *In our original version of the theorem ([KrVe19a, Theorem 3.2]), the spaces $\text{SU}(2p)/\text{SO}(p)$, $p \geq 6$ also appear in the list. However, a small error occurred there. These space are only tangentially stable but not strictly tangentially stable. In fact, the condition*

$$\text{Spec}(\Delta) \cap (n, 2(n+1)) = \emptyset$$

for the Laplace Beltrami operator holds for these spaces, but $\Delta_E|_{\text{TT}}$ has a kernel, see e.g. [CAHe15, Table 2].

4.2. Self-adjoint extensions of the Lichnerowicz Laplacian.

We can now study the self-adjoint closed extensions of Δ_L acting on compactly supported sections $C_0^\infty(M, S)$. We write $L^2(M, S)$ for the completion of $C_0^\infty(M, S)$ with respect to the L^2 -norm $(\cdot, \cdot)_{L^2}$ defined by g . The maximal closed extension of Δ_L in $L^2(M, S)$ is defined by the following domain

$$\mathcal{D}(\Delta_{L, \max}) := \{\omega \in L^2(M, S) \mid \Delta_L \omega \in L^2(M, S)\}, \quad (4.9)$$

where $\Delta_L \omega$ is defined distributionally. The minimal closed extension of Δ_L in $L^2(M, S)$ is obtained as the domain of the graph closure of Δ_L acting on $C_0^\infty(M, S)$

$$\begin{aligned} \mathcal{D}(\Delta_{L, \min}) &:= \{\omega \in \mathcal{D}(\Delta_{L, \max}) \mid \exists (\omega_n)_{n \in \mathbb{N}} \subset C_0^\infty(M, S) : \\ &\quad \omega_n \xrightarrow{n \rightarrow \infty} \omega, \quad \Delta_L \omega_n \xrightarrow{n \rightarrow \infty} \Delta_L \omega \text{ in } L^2(M, S)\}. \end{aligned}$$

Let $(\lambda, \omega_\lambda)$ be the set of eigenvalues and corresponding eigentensors of the tangential operator \square_L . Assuming tangential stability, we have $\lambda \geq 0$, and hence we define

$$\nu(\lambda) := \sqrt{\lambda + \left(\frac{n-1}{2}\right)^2}. \quad (4.10)$$

Standard arguments, see e.g. [KLPo8], show that for each $\omega \in \mathcal{D}(\Delta_{L,\max})$ there exist constants $c_\lambda^\pm, \nu(\lambda) \in [0, 1)$, depending only on ω , such that ω admits a partial asymptotic expansion as $x \rightarrow 0$

$$\begin{aligned} \omega &= \sum_{\nu(\lambda)=0} \left(c_\lambda^+(\omega) x^{-\frac{(n-1)}{2}} + c_\lambda^-(\omega) x^{-\frac{(n-1)}{2}} \log(x) \right) \cdot \omega_\lambda \\ &+ \sum_{\nu(\lambda) \in (0,1)} \left(c_\lambda^+(\omega) x^{\nu(\lambda)-\frac{(n-1)}{2}} + c_\lambda^-(\omega) x^{-\nu(\lambda)-\frac{(n-1)}{2}} \right) \cdot \omega_\lambda \\ &+ \tilde{\omega}, \quad \tilde{\omega} \in \mathcal{D}(\Delta_{L,\min}). \end{aligned} \quad (4.11)$$

All self-adjoint extensions for Δ_L can be classified by algebraic conditions on the coefficients in the asymptotic expansion (4.11), see e.g. Kirsten, Loya and Park [KLPo8, Proposition 3.3]. The Friedrichs self-adjoint extension of Δ_L on $C_0^\infty(M, S) \subset L^2(M, S)$ is given by the domain

$$\mathcal{D}(\Delta_L) := \{\omega \in \mathcal{D}(\Delta_{L,\max}) \mid c_\lambda^-(\omega) = 0 \text{ for } \nu(\lambda) \in [0, 1)\}. \quad (4.12)$$

Note that if $n \geq 3$, then tangential stability implies that all $\nu(\lambda) \geq 1$. Hence the minimal and maximal domains coincide and we find

$$\mathcal{D}(\Delta_L) = \mathcal{D}(\Delta_{L,\max}) = \mathcal{D}(\Delta_{L,\min}), \quad \text{for } n \geq 3. \quad (4.13)$$

We close the section with an observation by Friedrichs and Stone, see Riesz and Nagy [RiNa90, Theorem on p. 330], cf. the corresponding statement in our previous work [KrVe19a, Proposition 2.2].

Proposition 4.5. *Assume that (M, g) is tangentially stable, so that by (4.5) the Lichnerowicz Laplacian Δ_L with domain $C_0^\infty(M, S)$ is bounded from below by a constant² $C \in \mathbb{R}$. Then the Friedrichs self-adjoint extension of the Lichnerowicz Laplacian Δ_L is bounded from below by C as well.*

5. EXISTENCE AND REGULARITY OF SINGULAR RICCI DE TURCK FLOW

The main result of [VER16, Theorem 4.1], see also [KrVe19a, Theorem 1.2], is existence and regularity of singular Ricci de Turck flow. Despite a slight difference of Hölder spaces used here and in [VER16], we still conclude from [VER16, Theorem 4.1] that the heat operator of the Friedrichs extension Δ_L maps

$$e^{-t\Delta_L} : \mathcal{H}_{-2+\gamma_0, -2+\gamma_1}^{k+2, \alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{\gamma_0, \gamma_1}^{k+2, \alpha}(M \times [0, T], S).$$

Therefore existence and regularity obtained in [VER16, Theorem 4.1], as well as in [KrVe19a, Theorem 1.2] for a different choice of a background metric, still hold in our spaces.

Theorem 5.1. *Let (M, g) be a conical manifold, which is tangentially stable and $(\alpha, k+1, \gamma)$ Hölder regular. In case the conical singularity is not orbifold, we assume strict tangential stability. Let the background metric be either equal to g or a conical Ricci flat metric, in which case g_0 is assumed to be a sufficiently small perturbation of \tilde{g} in $\mathcal{H}_{\gamma, \gamma}^{k+2, \alpha}(M, S)$. Then there exists some $T > 0$, such that the Ricci de Turck flow (1.2), starting at g admits a solution $g(\cdot) \in \mathcal{H}_{\gamma_0, \gamma_1}^{k+2, \alpha}(M \times [0, T], S)$ for some $\gamma_0, \gamma_1 \in (0, \gamma)$ sufficiently small.*

²The case of $C = 0$ is commonly referred to as linear stability in the literature.

Let us now explain in what sense the flow preserves the conical singularity. Given an admissible perturbation g of the conical metric g_0 , the pointwise trace of g with respect to g_0 , denoted as $\text{tr}_{g_0} g$ is by definition of admissibility an element of the Hölder space $\mathcal{C}_{\text{ie}}^{k,\alpha}(M, S_1)_{\gamma}^b$, restricting at $x = 0$ to a constant function $(\text{tr}_{g_0} g)(0) = u_0 > 0$. Setting $\tilde{x} := \sqrt{u_0} \cdot x$, the admissible perturbation $g = g_0 + h$ attains the form

$$g = d\tilde{x}^2 + \tilde{x}^2 g_F + \tilde{h},$$

where $|\tilde{h}|_g = O(x^\gamma)$ as $x \rightarrow 0$. Note that the leading part of the admissible perturbation g near the conical singularity differs from the leading part of the admissible metric g_0 only by scaling.

We now wish to specify conditions on the weights γ_0, γ_1 . Let us write \square'_L for the tangential operator of the Lichnerowicz Laplacian Δ_L acting on trace-free tensors S_0 . The tangential operator of Δ_L acting on the pure trace tensors S_1 is simply the Laplace Beltrami operator Δ_F of (F, g_F) . We set $u_0 := \min(\text{Spec } \square'_L \setminus \{0\}) > 0$ and $u_1 := \min(\text{Spec } \Delta_F \setminus \{0\}) > 0$. In the orbifold case we set $u := \min\{u_0, u_1\}$. Define ³

$$\mu_0 := \sqrt{u_0 + \frac{n-1}{2}} - \frac{n-1}{2}, \quad \mu_1 := \sqrt{u_1 + \frac{n-1}{2}} - \frac{n-1}{2}.$$

Then the admissible choice of weights (γ_0, γ_1) (in the orbifold case set $\gamma = \gamma_0 = \gamma_1$) and the Hölder exponent α is given by the following restrictions.

$$\begin{aligned} \text{i)} \quad & \gamma_0 \in (0, \mu_0), \quad \gamma_0 \leq 2\gamma_1, \quad \gamma_0 < \gamma, \\ \text{ii)} \quad & \gamma_1 \in (0, \mu_1), \quad \gamma_1 \leq \gamma_0, \quad \gamma_1 < \gamma, \\ \text{iii)} \quad & \alpha \in (0, \mu_0 - \gamma_0) \cap (0, \mu_1 - \gamma_1). \end{aligned} \tag{5.1}$$

Remark 5.2. Note that in case of $u_0, u_1 > n$, i.e. $\square_L > n$, and assuming additionally $\gamma > 1$, we may choose $\gamma_0, \gamma_1 > 1$ satisfying (5.1). This stronger condition is studied in the Appendix §8, where a list of examples is provided.

We close the section by pointing out regularity of the de Turck vector field. As before we may define weighted higher order Hölder spaces of the incomplete b-cotangent bundle ${}^{\text{ib}}T^*M$. By Theorem 5.1, cf. [VER16, §6]

$$W(t) \in \mathcal{C}_{\text{ie}}^{k+1,\alpha}(M \times [0, T], {}^{\text{ib}}T^*M)_{-1+\bar{\gamma}}, \tag{5.2}$$

where $\bar{\gamma} := \min\{\gamma_0, \gamma_1\}$, and $\bar{\gamma} = \gamma$ in the orbifold case. As explained in the previous Remark 5.2, assuming $\square_L > n$ we can choose $\bar{\gamma} > 1$, so that existence time of the integral curves of $W(t)$ is positive, uniformly bounded away from zero. In this case we may pass from the Ricci de Turck flow back to Ricci flow, which is generally not clear in the singular setting.

6. EVOLUTION OF CURVATURES UNDER RICCI DE TURCK FLOW

The results of this section are well-known in the setting of smooth compact manifolds where the Ricci flow is defined. Evolution equations are usually proven by studying evolution of curvatures under the Ricci flow, and then the

³cf. [VER16, Remark 2.5]

corresponding equations follow for the Ricci de Turck flow by applying the corresponding diffeomorphisms. In the singular setting there might be no globally well-defined diffeomorphism to go from the Ricci de Turck flow back to the Ricci flow, as the de Turck vector field may be pointing toward the singularity. Thus we need to establish the evolution of Ricci curvature under Ricci de Turck flow directly without passing back to the Ricci flow. The evolution of the scalar curvature is then a direct consequence.

Notation: Let $g(t)$, $t \in [0, T]$ be a Ricci de Turck flow of Riemannian metrics on M . We denote by ∇ , Δ , Δ_L and Γ the covariant derivative, the Laplace Beltrami operator, the Lichnerowicz Laplacian and the Christoffel symbols, respectively, defined with respect to the metric $g(t)$. We let $|\cdot|$ be the norm with respect to $g(t)$. We denote the Riemann curvature tensor by Rm , the Ricci tensor by Ric and the scalar curvature by R , and we use the following conventions:

$$Rm(X, Y)Z := \nabla_X(\nabla_Y Z) - \nabla_Y(\nabla_X Z) - \nabla_{[X, Y]}Z,$$

$$Rm\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right)\frac{\partial}{\partial x^k} = R_{ijk}^l \frac{\partial}{\partial x^l}.$$

Here and below we use the Einstein summation convention and sum over repeated upper and lower indices. We lower the upper index to the fourth slot $R_{ijkl} := g_{pl}R_{ijk}^p$. Then the Ricci tensor Ric and the scalar curvature R are given by

$$Ric_{jk} \equiv R_{jk} := R_{ijk}^i = g^{il}R_{ijkl}, \quad R := g^{jk}R_{jk}.$$

Theorem 6.1. *Let M be a smooth manifold. Let $g(t)$, $t \in [0, T]$ be a solution of the Ricci de Turck flow with initial metric g and reference metric h , i.e.*

$$\begin{cases} \frac{\partial}{\partial t}g_{ij}(x, t) = (-2R_{ij} + \nabla_i W_j + \nabla_j W_i)(x, t), & (x, t) \in M \times [0, T], \\ g_{ij}(x, 0) = g_{ij}(x), & x \in M, \end{cases} \quad (6.1)$$

where $W(t)^k = g(t)^{ij}(\Gamma_{ij}^k(g(t)) - \Gamma_{ij}^k(h))$ is the de Turck vector field. Then the Ricci tensor Ric evolves by

$$(\partial_t + \Delta_L)R_{jk} = \nabla_m R_{jk} W^m + R_{jm} \nabla_k W^m + R_{km} \nabla_j W^m. \quad (6.2)$$

Proof. By [CHKn04, Lemma 3.5] the evolution of the Ricci curvature is given by

$$\partial_t R_{jk} = \frac{1}{2}g^{pq}(\nabla_q \nabla_j a_{kp} + \nabla_q \nabla_k a_{jp} - \nabla_p \nabla_q a_{jk} - \nabla_j \nabla_k a_{pq}), \quad (6.3)$$

where the time-dependent $(0, 2)$ -tensor a is the time-derivative of the metric, $a_{ij} := \frac{\partial}{\partial t}g_{ij}$. Using the Ricci de Turck flow equation $\frac{\partial}{\partial t}g_{ij} = -2R_{ij} + \nabla_i W_j + \nabla_j W_i$ we obtain

$$\begin{aligned} \partial_t R_{jk} = & -g^{pq}(\nabla_q \nabla_j R_{kp} + \nabla_q \nabla_k R_{jp} - \nabla_p \nabla_q R_{jk} - \nabla_j \nabla_k R_{pq}) \\ & + \frac{1}{2}g^{pq}(\nabla_q \nabla_j \nabla_k W_p + \nabla_q \nabla_j \nabla_p W_k + \nabla_q \nabla_k \nabla_j W_p + \nabla_q \nabla_k \nabla_p W_j \\ & - \nabla_p \nabla_q \nabla_j W_k - \nabla_p \nabla_q \nabla_k W_j - \nabla_j \nabla_k \nabla_p W_q - \nabla_j \nabla_k \nabla_q W_p). \end{aligned} \quad (6.4)$$

The first line on the right-hand side is the time-derivative of the Ricci tensor when the metric evolves by Ricci flow. Since it is known [CHKn04, Lemma 6.9]

that under Ricci flow the time-derivative of the Ricci tensor equals the Lichnerowicz Laplacian of the Ricci tensor, i.e. $\partial_t R_{jk} = -\Delta_L R_{jk}$, it follows that

$$\Delta_L R_{jk} = g^{pq}(\nabla_q \nabla_j R_{kp} + \nabla_q \nabla_k R_{jp} - \nabla_p \nabla_q R_{jk} - \nabla_j \nabla_k R_{pq}). \quad (6.5)$$

To simplify the second and third line on the right-hand side, we compute several terms by commuting covariant derivatives. We have

$$\begin{aligned} \nabla_p \nabla_j \nabla_p W_k &= \nabla_p (\nabla_q \nabla_j W_k - R_{jqk}^m W_m) \\ &= \nabla_p \nabla_q \nabla_j W_k - \nabla_p R_{jqk}^m W_m - R_{jqk}^m \nabla_p W_m. \end{aligned} \quad (6.6)$$

Hence by exchanging j and k

$$\nabla_p \nabla_k \nabla_p W_j = \nabla_p \nabla_q \nabla_k W_j - \nabla_p R_{kqj}^m W_m - R_{kqj}^m \nabla_p W_m. \quad (6.7)$$

Furthermore,

$$\begin{aligned} \nabla_p \nabla_j \nabla_k W_q &= \nabla_j \nabla_p \nabla_k W_q - R_{pj k}^m \nabla_m W_q - R_{pj k}^m \nabla_k W_m \\ &= \nabla_j (\nabla_k \nabla_p W_q - R_{pkq}^m W_m) - R_{pj k}^m \nabla_m W_q - R_{pj k}^m \nabla_k W_m \\ &= \nabla_j \nabla_k \nabla_p W_q - \nabla_j R_{pkq}^m W_m - R_{pkq}^m \nabla_j W_m - R_{pj k}^m \nabla_m W_q - R_{pj k}^m \nabla_k W_m. \end{aligned} \quad (6.8)$$

By exchanging j and k and commuting covariant derivatives we have

$$\begin{aligned} \nabla_p \nabla_k \nabla_j W_q &= \nabla_k \nabla_j \nabla_p W_q - \nabla_k R_{pj q}^m W_m - R_{pj q}^m \nabla_k W_m - R_{pkj}^m \nabla_m W_q - R_{pkj}^m \nabla_j W_m \\ &= \nabla_j \nabla_k \nabla_p W_q - R_{kjp}^m \nabla_m W_q - R_{kjp}^m \nabla_p W_m \\ &\quad - \nabla_k R_{pj q}^m W_m - R_{pj q}^m \nabla_k W_m - R_{pkj}^m \nabla_m W_q - R_{pkj}^m \nabla_j W_m. \end{aligned} \quad (6.9)$$

Now plugging (6.5), (6.6), (6.7), (6.8), (6.9) into (6.4) we obtain

$$\begin{aligned} \partial_t R_{jk} &= -\Delta_L R_{jk} - \frac{1}{2} g^{pq} (\nabla_p R_{jqk}^m W_m + R_{jqk}^m \nabla_p W_m + \nabla_p R_{kqj}^m W_m + R_{kqj}^m \nabla_p W_m \\ &\quad + \nabla_j R_{pkq}^m W_m + R_{pkq}^m \nabla_j W_m + R_{pj k}^m \nabla_m W_q + R_{pj k}^m \nabla_k W_m \\ &\quad + R_{kjp}^m \nabla_m W_q + R_{kjp}^m \nabla_p W_m + \nabla_k R_{pj q}^m W_m + R_{pj q}^m \nabla_k W_m \\ &\quad + R_{pkj}^m \nabla_m W_q + R_{pkj}^m \nabla_j W_m). \end{aligned} \quad (6.10)$$

Simplifying and rearranging terms yields

$$\begin{aligned} \partial_t R_{jk} &= -\Delta_L R_{jk} - \frac{1}{2} g^{pq} \nabla_q W_m (R_{jpk}^m + R_{kpj}^m + R_{kjp}^m) - \frac{1}{2} g^{pq} \nabla_m W_q (R_{pj k}^m + R_{kjp}^m + R_{pkj}^m) \\ &\quad - \frac{1}{2} g^{pq} W_m (\nabla_p R_{jqk}^m + \nabla_p R_{kqj}^m + \nabla_j R_{pkq}^m + \nabla_k R_{pj q}^m) \\ &\quad - g^{pq} (R_{pkq}^m \nabla_j W_m + R_{pj q}^m \nabla_k W_m). \end{aligned} \quad (6.11)$$

The second line on the right-hand side is

$$\begin{aligned} & - \frac{1}{2} g^{pq} \nabla_q W_m (R_{jpk}^m + R_{kpj}^m + R_{kjp}^m) - \frac{1}{2} g^{pq} \nabla_m W_q (R_{pj k}^m + R_{kjp}^m + R_{pkj}^m) \quad (6.12) \\ &= - \frac{1}{2} \nabla^p W^m (R_{jpkm} + R_{kpjm} + R_{kjp m}) - \frac{1}{2} \nabla^m W^p (R_{pj km} + R_{kjp m} + R_{pkj m}) \\ &= - \frac{1}{2} \nabla^p W^m (R_{jpkm} + R_{kpjm} + R_{kjp m}) - \frac{1}{2} \nabla^p W^m (R_{mj kp} + R_{kj mp} + R_{mkjp}) \\ &= - \frac{1}{2} \nabla^p W^m (R_{jpkm} + R_{kpjm} + R_{kjp m} - R_{kpjm} - R_{kjp m} - R_{jpk m}) \\ &= 0, \end{aligned}$$

where in the penultimate line we used the symmetries of the curvature tensor. The fourth line on the right-hand side is given by

$$-g^{pq}(R_{pkq}^m \nabla_j W_m + R_{pjq}^m \nabla_k W_m) = R_{km} \nabla_j W^m + R_{jm} \nabla_k W^m. \quad (6.13)$$

To compute the third line on the right-hand side, we first observe that by the symmetries of the curvature tensor and the second Bianchi identity

$$\nabla_p R_{jqkm} = \nabla_p R_{kmjq} = -\nabla_m R_{pkjq} - \nabla_k R_{mpjq}. \quad (6.14)$$

Moreover, exchanging j and k yields

$$\nabla_p R_{kqjm} = -\nabla_m R_{pjkm} - \nabla_j R_{mpkq}. \quad (6.15)$$

Now using (6.14) and (6.15) we obtain for the third line on the right-hand side

$$\begin{aligned} & -\frac{1}{2}g^{pq}W_m(\nabla_p R_{jqk}^m + \nabla_p R_{kqj}^m + \nabla_j R_{pkq}^m + \nabla_k R_{pjq}^m) \\ &= -\frac{1}{2}g^{pq}W^m(-\nabla_m R_{pkjq} - \nabla_k R_{mpjq} - \nabla_m R_{pjkm} - \nabla_j R_{mpkq} + \nabla_j R_{pkqm} + \nabla_k R_{pjqm}) \\ &= -\frac{1}{2}W^m(-\nabla_m R_{kj} + \nabla_k R_{mj} - \nabla_m R_{jk} + \nabla_j R_{mk} - \nabla_j R_{km} - \nabla_k R_{jm}) \\ &= W^m \nabla_m R_{jk}. \end{aligned} \quad (6.16)$$

Plugging (6.12), (6.13) and (6.16) into (6.11) we finally obtain the claimed evolution equation of the Ricci tensor

$$\partial_t R_{jk} = -\Delta_L R_{jk} + \nabla_m R_{jk} W^m + R_{jm} \nabla_k W^m + R_{km} \nabla_j W^m.$$

□

Corollary 6.2. *Let M be a smooth manifold. Let $g(t)$, $t \in [0, T]$ be a solution of the Ricci de Turck flow with initial metric g and reference metric h , i.e.*

$$\begin{cases} \frac{\partial}{\partial t} g_{ij}(x, t) = (-2R_{ij} + \nabla_i W_j + \nabla_j W_i)(x, t), & (x, t) \in M \times [0, T], \\ g_{ij}(x, 0) = g_{ij}(x), & x \in M, \end{cases} \quad (6.17)$$

where $W(t)^k = g(t)^{ij}(\Gamma_{ij}^k(g(t)) - \Gamma_{ij}^k(h))$ is the de Turck vector field. Then the scalar curvature R evolves by

$$(\partial_t + \Delta)R = \langle W, \nabla R \rangle + 2|\text{Ric}|^2. \quad (6.18)$$

Proof. The scalar curvature is given by $R = g^{jk}R_{jk}$. Since

$$\partial_t g^{jk} = -g^{jp}g^{kq}\partial_t g_{pq} = -g^{jp}g^{kq}(-2R_{pq} + \nabla_p W_q + \nabla_q W_p)$$

and by the evolution equation for the Ricci tensor (6.2) we thus obtain

$$\begin{aligned} \partial_t R &= -g^{jp}g^{kq}(-2R_{pq} + \nabla_p W_q + \nabla_q W_p)R_{jk} \\ &\quad + g^{jk}(-\Delta_L R_{jk} + \nabla_m R_{jk} W^m + R_{jm} \nabla_k W^m + R_{km} \nabla_j W^m) \\ &= 2|\text{Ric}|^2 - 2g^{jp}g^{kq}\nabla_p W_q R_{jk} - g^{jk}\Delta_L R_{jk} + \nabla_m R W^m + 2g^{jk}R_{jm} \nabla_k W^m \\ &= -\Delta R + \nabla_m R W^m + 2|\text{Ric}|^2, \end{aligned}$$

where in the last step we used $g^{jk}\Delta_L R_{jk} = \Delta R$. □

Remark 6.3. *Note that in the proof we didn't use the special form of W , we just used that W is a (time-dependent) one-form.*

7. REGULARITY OF RICCI CURVATURE ALONG THE RICCI DE TURCK FLOW

Our aim in this section is to improve the a priori low regularity of the Ricci curvature along the singular Ricci de Turck flow, as noted in [VER16, Theorem 8.1]. Consider an $(\alpha, k+1, \gamma)$ -Hölder regular conical manifold (M, g) , satisfying tangential stability, with singular Ricci de Turck flow $g(\cdot) \in \mathcal{H}_{\gamma_0, \gamma_1}^{k, \alpha}(M \times [0, T], S)$, i.e. decomposing $g(t) = (1+u)g + \omega$ into trace and trace-free parts with respect to the initial metric g , we have

$$(\omega, u) \in \mathcal{C}_{\text{ie}}^{k+2, \alpha}(M \times [0, T], S_0)_{\gamma_0} \oplus \mathcal{C}_{\text{ie}}^{k+2, \alpha}(M \times [0, T], S_1)_{\gamma_1}^b.$$

Recall the following transformation rule for the Ricci curvature tensor under conformal transformations (setting $1+u = e^{2\phi}$ and noting that $\dim M = n+1$)

$$\begin{aligned} \text{Ric}((1+u)g) &= \text{Ric}(g) - (n-1)(\nabla\partial\phi - \partial\phi \cdot \partial\phi) \\ &\quad + (\Delta_L\phi - (n-1)\|\nabla\phi\|^2)g. \end{aligned} \tag{7.1}$$

From here we conclude that $\text{Ric}((1+u)g) - \text{R}(g) \in \mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{-2+\gamma_1}$. Now consider $\text{Ric}((1+u)g + \omega) - \text{Ric}((1+u)g)$, which is an intricate combination of α and ω , involving their second order $x^{-2} \mathcal{V}^2$ derivatives. Hence that difference lies in $\mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{-2+\bar{\gamma}}$ with $\bar{\gamma} := \min\{\gamma_0, \gamma_1\}$. We conclude

$$\begin{aligned} \text{Ric}(g) &\in \mathcal{C}_{\text{ie}}^{k+1, \alpha}(M \times [0, T], S)_{-2+\gamma}, \\ \text{Ric}(g(t)) &\in \mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{-2+\bar{\gamma}}, \quad t > 0. \end{aligned} \tag{7.2}$$

In particular, e.g. if $\mu_0, \mu_1 \leq 2 \leq \gamma$, then $\bar{\gamma} \leq 2 \leq \gamma$. In that case, the initial Ricci curvature $\text{Ric}(g)$ is bounded as a section of S , while for positive times $\text{Ric}(g(t))$ is singular as a section of S . In this section we improve this low regularity result.

7.1. Expansion of the Lichnerowicz Laplacian. Below we fix the following notation: A superscript “ \sim ” indicates that the quantity is taken with respect to the initial metric $g(0)$. For example, $\widetilde{\nabla}$ and $\widetilde{\Delta}_L$ refer to the covariant derivative and the Lichnerowicz Laplacian with respect to $g(0)$. Otherwise the quantities are defined with respect to the flow $g(t)$.

Theorem 7.1. *Consider an $(\alpha, k+1, \gamma)$ -Hölder regular conical manifold (M, g) , satisfying tangential stability, with singular Ricci de Turck flow $g(\cdot) \in \mathcal{H}_{\gamma_0, \gamma_1}^{k, \alpha}(M \times [0, T], S)$, i.e. decomposing $g(t) = \alpha g + \omega$ into trace and trace-free parts with respect to the initial metric g . Then there exist*

$$\begin{aligned} A &\in \mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{\bar{\gamma}}, \\ B &\in \mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{-1+\bar{\gamma}}, \\ C &\in \mathcal{C}_{\text{ie}}^{k, \alpha}(M \times [0, T], S)_{-2+\bar{\gamma}}, \end{aligned}$$

such that for any (time-dependent) symmetric 2-tensor c

$$\Delta_L c = \alpha^{-1} \widetilde{\Delta}_L c + (A \cdot (x^{-1} \mathcal{V}_b)^2 + B \cdot x^{-1} \mathcal{V}_b + C) c. \tag{7.3}$$

Proof. Consider the rough Laplacian $\Delta = -g^{ij}\nabla_i\nabla_j$ first. We start by writing out Δ in local coordinates and later on turn to the Lichnerowicz Laplacian Δ_L . We compute for any symmetric 2-tensor c

$$\begin{aligned} (\Delta c)_{kl} &= -g^{ij}\nabla_i\nabla_j c_{kl} \\ &= -g^{ij}(\partial_i\partial_j c_{kl} - \partial_i\Gamma_{jk}^m c_{ml} - \Gamma_{jk}^m\partial_i c_{ml} - \partial_i\Gamma_{jl}^m c_{km} - \Gamma_{jl}^m\partial_i c_{km} \\ &\quad - \Gamma_{ij}^m\partial_m c_{kl} + \Gamma_{ij}^m\Gamma_{mk}^p c_{pl} + \Gamma_{ij}^m\Gamma_{ml}^p c_{kp} - \Gamma_{ik}^m\partial_j c_{ml} + \Gamma_{ik}^m\Gamma_{jm}^p c_{pl} \\ &\quad + \Gamma_{ik}^m\Gamma_{jl}^p c_{mp} - \Gamma_{il}^m\partial_j c_{km} + \Gamma_{jk}^m\Gamma_{il}^p c_{mp} + \Gamma_{il}^m\Gamma_{jm}^p c_{kp}). \end{aligned} \quad (7.4)$$

Our goal is to obtain an expansion for each term on the right-hand side of (7.4). The metric $g(t)$ has the form $g_{ij} = a\tilde{g}_{ij} + \omega_{ij}$, with inverse (cf. [VER16, p. 28])

$$g^{ij} = a^{-1}\tilde{g}^{ij} - a^{-2}\tilde{g}^{il}\tilde{g}^{jp}\omega_{pl} + a^{-2}g^{jl}\tilde{g}^{ir}\tilde{g}^{pq}\omega_{lp}\omega_{rq}. \quad (7.5)$$

Hence the first term on the right-hand side of (7.4) is given by

$$-g^{ij}\partial_i\partial_j c_{kl} = -a^{-1}\tilde{g}^{ij}\partial_i\partial_j c_{kl} - (-a^{-2}\tilde{g}^{is}\tilde{g}^{jp}\omega_{ps} + a^{-2}g^{js}\tilde{g}^{ir}\tilde{g}^{pq}\omega_{sp}\omega_{rq})\partial_i\partial_j c_{kl}.$$

Now we study the asymptotics of the last expression at the conical singularity. This is analogous to the discussion in [VER16, p. 28-29]. Consider the local coordinates (z_0, \dots, z_n) of $\mathcal{C}(F)$ with $z_0 = x$ and (z_1, \dots, z_n) being local coordinates of F . An upper index $i = 0$ does not contribute any singular factor of x due to the structure of the inverse g^{-1} . A lower index $i = 0$ indicates a differentiation by $\partial_x \in x^{-1}\mathcal{V}_b$. Hence an index $i = 0$ (as a combination of a lower and upper index) contributes $x^{-1}\mathcal{V}_b$ up to a term of type A .

Similarly, an upper index $i > 0$ contributes a singular factor x^{-1} due to the structure of the inverse g^{-1} . A lower index $i > 0$ indicates a differentiation by $\partial_{z_j} \in \mathcal{V}_b$. Hence an index $i > 0$ (as a combination of a lower and upper index) contributes $x^{-1}\mathcal{V}_b$ up to a term of type A . Hence in total we find

$$-g^{ij}\partial_i\partial_j c_{kl} = -a^{-1}\tilde{g}^{ij}\partial_i\partial_j c_{kl} + A \cdot (x^{-1}\mathcal{V}_b)^2 c_{kl}, \quad (7.6)$$

for some $A \in C_{ie}^{k,\alpha}(M \times [0, T], S)_{\overline{\gamma}}$. In order to study the remaining terms in (7.4), involving Christoffel symbols, we note that the Christoffel symbols Γ_{ij}^k with respect to the metric $g(t)$ are related to the Christoffel symbols $\tilde{\Gamma}_{ij}^k$ of the initial metric \tilde{g} as follows

$$\begin{aligned} \Gamma_{ij}^k &= \tilde{\Gamma}_{ij}^k + \frac{1}{2}a^{-1}\tilde{g}^{km}(\partial_i a \tilde{g}_{jm} + \partial_j a \tilde{g}_{im} - \partial_m a \tilde{g}_{ij} + \partial_i \omega_{jm} \\ &\quad + \partial_j \omega_{im} - \partial_m \omega_{ij}) + \frac{1}{2}(-a^{-2}\tilde{g}^{kl}\tilde{g}^{mp}\omega_{pl} + a^{-2}g^{ml}\tilde{g}^{kr}\tilde{g}^{pq}\omega_{lp}\omega_{rq}) \\ &\quad \times (a(\partial_i \tilde{g}_{jm} + \partial_j \tilde{g}_{im} - \partial_m \tilde{g}_{ij}) + \partial_i a \tilde{g}_{jm} + \partial_j a \tilde{g}_{im} - \partial_m a \tilde{g}_{ij} \\ &\quad + \partial_i \omega_{jm} + \partial_j \omega_{im} - \partial_m \omega_{ij}) \end{aligned} \quad (7.7)$$

This allows us to conclude by counting upper and lower indices as above that for any $\{Y_\ell\}_\ell$ with $Y_0 := \partial_x$ and $Y_i := x^{-1}\partial_{z_i}$ for $i = 1, \dots, n$, we have

$$(\Delta c)(Y_k, Y_l) = a^{-1}(\tilde{\Delta} c)(Y_k, Y_l) + (A \cdot (x^{-1}\mathcal{V}_b)^2 + B \cdot x^{-1}\mathcal{V}_b + C) c, \quad (7.8)$$

for some terms A, B, C as in the statement of the theorem. Now we turn to the Lichnerowicz Laplacian. Recall that it is given by

$$\Delta_L c_{kl} = \Delta c_{kl} - 2g^{pq} R_{qkl}^r c_{rp} + g^{pq} R_{kp} c_{ql} + g^{pq} R_{lp} c_{kq}.$$

To deal with the extra terms involving the Riemann curvature tensor and the Ricci curvature on the right-hand side, we note the following formula for the components of the curvature tensor in local coordinates

$$R_{ijk}^l = \partial_i \Gamma_{jk}^l - \partial_j \Gamma_{ik}^l + \Gamma_{jk}^p \Gamma_{ip}^l - \Gamma_{ik}^p \Gamma_{jp}^l, \quad (7.9)$$

which also leads to a formula for the components of the Ricci tensor

$$R_{jk} = R_{ijk}^i = \partial_i \Gamma_{jk}^i - \partial_j \Gamma_{ik}^i + \Gamma_{jk}^p \Gamma_{ip}^i - \Gamma_{ik}^p \Gamma_{jp}^i. \quad (7.10)$$

These extra terms can be treated using (7.7), which leads to the claimed expansion for the Lichnerowicz Laplacian. \square

Corollary 7.2. *Under the conditions of Theorem 7.1 we conclude*

$$(\partial_t + \alpha^{-1} \widetilde{\Delta}_L) \text{Ric} =: X(\text{Ric}) \in \mathcal{C}_{ie}^{k-1, \alpha}(M \times [0, T], S)_{-4+2\gamma}. \quad (7.11)$$

Proof. Recall the evolution (6.2) of Ricci curvature along the Ricci de Turck flow

$$(\partial_t + \Delta_L) \text{Ric}_{jk} = \nabla_m \text{Ric}_{jk} W^m + \text{Ric}_{jm} \nabla_k W^m + \text{Ric}_{km} \nabla_j W^m, \quad (7.12)$$

By (5.2) and (7.2) we conclude

$$(\partial_t + \Delta_L) \text{Ric} \in \mathcal{C}_{ie}^{k-1, \alpha}(M \times [0, T], S)_{-4+2\gamma}.$$

The statement now follows from Theorem 7.1 and (7.2). \square

7.2. Mapping properties of the heat operator for $\widetilde{\Delta}_L$. Let us simplify notation by writing $\mathcal{H}_\gamma^{k, \alpha} \equiv \mathcal{H}_{\gamma, \gamma}^{k, \alpha}$. The heat operator for $\widetilde{\Delta}_L$ has been studied in [VER16, Theorem 3.1], which asserts that for $n \geq 3$, any $\gamma \in (0, 2 + \min\{\mu_0, \mu_1\})$ and $\alpha \in (0, 1)$ sufficiently small, the following is a bounded mapping

$$e^{-t\widetilde{\Delta}_L} : \mathcal{H}_{-4+\gamma}^{k, \alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2, \alpha}(M \times [0, T], S). \quad (7.13)$$

The following mapping properties are needed later in this section, and follow by similar arguments as in [VER16, Theorem 3.1] and (7.13).

Lemma 7.3. *Consider smooth functions $\rho \in C^\infty(\overline{M})$ smooth up to the boundary, $\eta \in C_0^\infty(M)$ smooth with compact support away from the conical singularity. Let $\nu \in C_0^\infty(M, TM)$ be a smooth 1-form and $\omega \in C_0^\infty(M, {}^{\text{ib}}T^*M \otimes {}^{\text{ib}}T^*M)$ a smooth $(0, 2)$ -tensor, both with compact support away from the conical singularity. Let H be the heat operator of the Lichnerowicz Laplacian Δ_L . Then*

$$\begin{aligned} E_0 &:= \eta H \rho : \mathcal{H}_{-4+\gamma}^{k, \alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2, \alpha}(M \times [0, T], S), \\ \widetilde{E}_0 &:= \nu \otimes H \rho : \mathcal{H}_{-4+\gamma}^{k, \alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2, \alpha}(M \times [0, T], {}^{\text{ib}}T^*M \otimes S), \\ \widetilde{\widetilde{E}}_0 &:= \omega \otimes H \rho : \mathcal{H}_{-4+\gamma}^{k, \alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2, \alpha}(M \times [0, T], {}^{\text{ib}}T^*M^{\otimes 2} \otimes S), \end{aligned}$$

with $\|E_0\| \rightarrow 0$, $\|\widetilde{E}_0\| \rightarrow 0$ and $\|\widetilde{\widetilde{E}}_0\| \rightarrow 0$ as $T \rightarrow 0$.

The next result is a straightforward consequence of Lemma 7.3.

Corollary 7.4. *In the notation of Lemma 7.3 we find*

$$\begin{aligned} F_0 &:= \nu \otimes \nabla H\rho : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-3+\gamma}^{k+1,\alpha}(M \times [0, T], {}^{\text{ib}}T^*M^{\otimes 2} \otimes S), \\ G_0 &:= g^{ij}\nu_i \nabla_j H\rho : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-4+\gamma}^{k+1,\alpha}(M \times [0, T], S), \end{aligned}$$

with $\|F_0\| \rightarrow 0, \|G_0\| \rightarrow 0$ as $T \rightarrow 0$.

Proof. By Lemma 7.3 we have

$$\nabla(\nu \otimes H\rho) : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S) \rightarrow \mathcal{H}_{-3+\gamma}^{k+1,\alpha}(M \times [0, T], {}^{\text{ib}}T^*M^{\otimes 2} \otimes S),$$

with $\|\nabla(\nu \otimes H\rho)\| \rightarrow 0$ as $T \rightarrow 0$. On the other hand

$$\nabla(\nu \otimes H\rho) = \nabla\nu \otimes H\rho + \nu \otimes \nabla H\rho.$$

Hence the mapping properties of F_0 follow by applying Lemma 7.3 to $\nabla\nu \otimes H\rho$. The mapping properties of G_0 follow from the ones of F_0 , as G_0 is simply F_0 composed with a contraction. \square

7.3. Heat operator parametrix for $\alpha^{-1}\widetilde{\Delta}_L$. Our goal in this section is the existence of inverses \mathcal{Q} and \mathcal{R} for the parabolic operator $P := \partial_t + \alpha^{-1}\widetilde{\Delta}_L$, where α is the pure trace part of $g(t)$ with respect to the initial metric g and is positive uniformly bounded away from zero for short time. These parametrices are constructed out of the heat operator for the Friedrichs extension $\widetilde{\Delta}_L$, using (7.13), Lemma 7.3 and Corollary 7.4. Our main result in this subsection is as follows.

Theorem 7.5. *Let $n \geq 3$. Consider any $\gamma \in (0, 2 + \min\{\mu_0, \mu_1\})$ and $\alpha \in (0, 1)$ sufficiently small, such that (7.13) holds. Then there exists $T_0 > 0$ sufficiently small and a bounded linear map*

$$\mathcal{Q} : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M \times [0, T_0], S),$$

such that if $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$, then $u = \mathcal{Q}f$ solves the initial value problem

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)u = f, \quad u(\cdot, 0) = 0.$$

The proof of that theorem will occupy the rest of this subsection. Before we proceed, let us note an immediate consequence.

Theorem 7.6. *Let $n \geq 3$. Consider any $\gamma \in (0, 2 + \min\{\mu_0, \mu_1\})$ and $\alpha \in (0, 1)$ sufficiently small, such that (7.13) holds. Then there exists $T_0 > 0$ sufficiently small and a bounded linear map*

$$\mathcal{R} : \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M, S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M \times [0, T_0], S),$$

such that if $u_0 \in \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M, S)$, then $u = \mathcal{R}u_0$ solves the initial value problem

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)u = 0, \quad u(\cdot, 0) = u_0.$$

Proof. We have $\widetilde{\Delta}_L u_0 \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M, S)$ and thus $\alpha^{-1}\widetilde{\Delta}_L u_0 \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$. We set $\mathcal{R}u_0 := u_0 + \mathcal{Q}(\alpha^{-1}\widetilde{\Delta}_L u_0)$, which yields the desired solution to the initial value problem above. \square

7.3.1. Construction of a boundary parametrrix.

We now begin with the proof of Theorem 7.5. The proof idea is to construct boundary and interior parametrices, and glue them together to an approximate solution to $(\partial_t + a^{-1}\widetilde{\Delta}_L)$. Provided the error is sufficiently "nice", we can obtain a solution \mathcal{Q} by a Neumann series argument. We follow the analytic path outlined in the work by the third named author jointly with Bahuaud [BAVe16].

We start by introducing the notion of a *reference covering* which is a special case of a covering considered in [BAVe16]. Let $U = [0, 1]_x \times (-1, 1]_z^n$ be a model half-cube. For each $p \in \partial M$, there is a coordinate chart $\Phi_p : U \rightarrow W_p$ onto an open neighborhood $W_p \subset \mathcal{C}(F) \subset \overline{M}$, centered around p_i . Due to compactness of ∂M , there are finitely many such charts $\{W_i, \Phi_{p_i}\}_{i=1}^N$ covering $\mathcal{C}(F)$. Together with the open set $W_0 = \overline{M} \setminus ([0, \frac{1}{2}) \times F)$ we obtain an open cover of \overline{M} that we call a reference covering.

Let $\sigma : \mathbb{R}^+ \rightarrow [0, 1]$ be a smooth cutoff function with $\sigma(s) = 1$ for $s \leq 1/4$ and $\sigma(s) = 0$ for $s \geq 1/2$. Denote by $(x, z) \in \Phi_{p_i}^{-1}(W_i)$ the local coordinates on the coordinate chart W_i , centered around $p_i \in \partial M$. Then we define

$$\varphi_i(\Phi_{p_i}(x, z)) := \sigma(x)\sigma(\|z\|), \quad \psi_i(\Phi_{p_i}(x, z)) := \sigma\left(\frac{x}{2}\right)\sigma\left(\frac{\|z\|}{2}\right).$$

Notice that $\varphi_i, \psi_i \in C^\infty(\overline{M})$ are both identically 1 near $(0, 0)$, and $\psi_i \equiv 1$ on $\text{supp } \varphi_i$. Furthermore, $\varphi_i, \psi_i \in C_{ie}^\alpha(M)$, since they are constant near the cone points. These functions are illustrated in the radial direction in Figure 1.




FIGURE 1. The cutoff functions φ_i, ψ_i .

Next, for $\varepsilon \in (0, 1)$ to be specified later, and any $p = z_0 \in W_i \cap \partial M$ we set

$$\tilde{\varphi}_{i,p}(x, z) := \varphi_i\left(\frac{x}{\varepsilon}, z - z_0\right), \quad \tilde{\psi}_{i,p}(x, z) := \psi_i\left(\frac{x}{\varepsilon}, z - z_0\right).$$

This defines smooth cutoff functions $\tilde{\varphi}_{i,p}, \tilde{\psi}_{i,p} \in C^\infty(\overline{M})$ which again lie in $C_{ie}^{k,\alpha}(M)$, with the Hölder norms bounded by $\varepsilon^{-\alpha}\|\varphi_i\|_{C_{ie}^{k,\alpha}(M)}$ and $\varepsilon^{-\alpha}\|\psi_i\|_{C_{ie}^{k,\alpha}(M)}$, respectively

$$\|\tilde{\varphi}_{i,p}\|_{C_{ie}^{k,\alpha}(M)} \leq \varepsilon^{-\alpha}\|\varphi_i\|_{C_{ie}^{k,\alpha}(M)}, \quad \|\tilde{\psi}_{i,p}\|_{C_{ie}^{k,\alpha}(M)} \leq \varepsilon^{-\alpha}\|\psi_i\|_{C_{ie}^{k,\alpha}(M)}. \quad (7.14)$$

Let L_i be the lattice of points in the coordinate chart W_i of the form $\{(0, w) \mid w \in \mathbb{Z}^n\}$. By construction, every point on ∂M lies in the support of at most a fixed number (independent of ε) of functions $\{\tilde{\psi}_{i,p} \mid i = 1, \dots, N, p \in L_i\}$. Furthermore,

let $F_\varepsilon = F_\varepsilon(x)$ be a cutoff function with $\text{supp } F_\varepsilon \subset \{x \geq \varepsilon/4\} \subset M$, which is identically 1 on the set $\{x \geq \varepsilon\}$. Let $G_\varepsilon(x) = F_\varepsilon(\frac{x}{2})$. For any $p \in L_i$ we normalize

$$\varphi_{i,p} := \frac{\tilde{\varphi}_{i,p}}{G_\varepsilon + \sum_k \sum_{q \in L_k} \tilde{\varphi}_{k,q}}, \quad \psi_{i,p} := \frac{\tilde{\psi}_{i,p}}{F_\varepsilon + \sum_k \sum_{q \in L_k} \tilde{\varphi}_{k,q}}. \quad (7.15)$$

By (7.14), these functions again lie in $C_{ie}^\alpha(M)$, with

$$\begin{aligned} \|\varphi_{i,p}\|_{C_{ie}^{k,\alpha}(M)} &\leq \text{const} \cdot \varepsilon^{-\alpha} \|\varphi_i\|_{C_{ie}^{k,\alpha}(M)} \leq \text{const} \cdot \varepsilon^{-\alpha}, \\ \|\psi_{i,p}\|_{C_{ie}^{k,\alpha}(M)} &\leq \text{const} \cdot \varepsilon^{-\alpha} \|\psi_i\|_{C_{ie}^{k,\alpha}(M)} \leq \text{const} \cdot \varepsilon^{-\alpha}. \end{aligned} \quad (7.16)$$

Finally note that due to normalization, for any $0 < \varepsilon < 1$, the functions

$$\Phi := \sum_i \sum_{p \in L_i} \varphi_{i,p}, \quad \Psi := \sum_i \sum_{p \in L_i} \psi_{i,p}$$

are smooth cutoff functions in $C^\infty(\overline{M})$ which are identically 1 in a neighborhood of the cone singularity. We can now introduce an approximate boundary parametrix. Consider any $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$ and denote by H_p the heat kernel of $\alpha(p, 0)^{-1} \widetilde{\Delta}_L$ for any $p \in \partial M$ (note this is simply a rescaling of $\widetilde{\Delta}_L$ by a positive constant). Then we define our boundary parametrix by

$$Q_b f := \sum_{j=1}^N \sum_{p \in L_j} \psi_{j,p} H_p[\varphi_{j,p} f]. \quad (7.17)$$

Lemma 7.7. *The solution $u_p := \psi_{j,p} H_p[\varphi_{j,p} f]$ satisfies*

$$(\partial_t + \alpha^{-1} \widetilde{\Delta}_L) u_p = \varphi_{j,p} f + E_{j,p}^0 f + E_{j,p}^1 f,$$

where the operators $E_{j,p}^0$ and $E_{j,p}^1$ are as follows.

- (i) $E_{j,p}^0$ is a bounded linear map on $\mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S)$, and there exists a constant $C > 0$ independent of j, p , such that for $T/\varepsilon^2 < 1$

$$\|E_{j,p}^0 f\| \leq C(\varepsilon^\gamma + T^{\alpha/2}) \|\varphi_{j,p} f\|,$$

- (ii) $E_{j,p}^1$ is a bounded linear map on $\mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S)$, with operator norm satisfying $\|E_{j,p}^1\| \rightarrow 0$ as $T \rightarrow 0^+$.

Proof. We simplify notation by omitting the subscripts on φ, ψ and E^0, E^1 . Then

$$\begin{aligned} (\partial_t + \alpha^{-1} \widetilde{\Delta}_L) u_p &= (\partial_t + \alpha^{-1} \widetilde{\Delta}_L)(\psi H_p[\varphi f]) \\ &= \psi \partial_t H_p[\varphi f] + \alpha^{-1} (\widetilde{\Delta} \psi \cdot H_p[\varphi f] + 2\tilde{g}^{ij} \widetilde{\nabla}_i \psi \widetilde{\nabla}_j H_p[\varphi f] + \psi \widetilde{\Delta}_L H_p[\varphi f]) \\ &= \psi (\partial_t + \alpha^{-1}(p, 0) \widetilde{\Delta}_L) H_p[\varphi f] + \psi (\alpha^{-1} - \alpha^{-1}(p, 0)) \widetilde{\Delta}_L H_p[\varphi f] \\ &\quad + \alpha^{-1} \widetilde{\Delta} \psi \cdot H_p[\varphi f] + \alpha^{-1} 2\tilde{g}^{ij} \widetilde{\nabla}_i \psi \widetilde{\nabla}_j H_p[\varphi f] =: \psi \varphi f + E^0 f + E^1 f, \end{aligned}$$

where the operators E^0 and E^1 are explicitly given by

$$\begin{aligned} E^0 f &:= \psi (\alpha^{-1} - \alpha^{-1}(p, 0)) \widetilde{\Delta}_L H_p[\varphi f], \\ E^1 f &:= \alpha^{-1} \widetilde{\Delta} \psi \cdot H_p[\varphi f] + 2\alpha^{-1} \tilde{g}^{ij} \widetilde{\nabla}_i \psi \widetilde{\nabla}_j H_p[\varphi f]. \end{aligned}$$

⁴Recall that $g(p, t) = \alpha(p, t)g(p) \oplus \omega(p, t)$.

Since $\psi \equiv 1$ on $\text{supp } \varphi$, any derivative of ψ vanishes on a neighborhood of the conical singularity. Hence the claimed mapping properties of E^1 follow from Lemma 7.3 and Corollary 7.4. What is left is establishing the mapping properties of E^0 . Writing out definitions of the various Hölder norms, we find

$$\|E^0 f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \leq C \|\psi(a^{-1} - a^{-1}(p, 0))\|_{C_{ie}^{k,\alpha}} \cdot \|\widetilde{\Delta}_L H_p[\varphi f]\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}}. \quad (7.18)$$

We write $p = (0, z_0)$. For any (x, z) in the support of ψ we obtain

$$\begin{aligned} |a^{-1}(x, z, t) - a^{-1}(0, z_0, 0)| &\leq |x|^\gamma \|x^{-\gamma}(\partial_x a^{-1})\|_\infty + t^{\frac{\alpha}{2}} \|a^{-1}\|_\alpha \\ &\leq |x|^\gamma \|x^{-\gamma}(\partial_x a)\|_\infty + t^{\frac{\alpha}{2}} \|a\|_\alpha \\ &\leq C(d_M(p, q)^\gamma + t^{\frac{\alpha}{2}}) \|a\|_{C_{ie}^{1,\alpha}(M \times [0, T], S_1)}^\flat \\ &\leq C(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) \|a\|_{C_{ie}^{1,\alpha}(M \times [0, T], S_1)}^\flat, \end{aligned}$$

where we used that $a(0, \cdot, t)$ is constant in z , and $\partial_x a^{-1} = -\partial_x a \cdot a^{-2}$. Recall the notation \mathcal{D} of Definition 3.3. Then for any partial differential operator $X \in \mathcal{D}$ of order at least 1, we compute similar to above for any $(x, z) \in \text{supp } \psi$

$$\begin{aligned} |X(a^{-1}(x, z, t) - a^{-1}(0, z_0, 0))| &= |Xa^{-1}(x, z, t)| \leq |x|^\gamma \|x^{-\gamma} Xa^{-1}\|_\infty \\ &\leq C\varepsilon^\gamma \|a\|_{C_{ie}^{1,\alpha}(M \times [0, T], S_1)}^\flat \leq C(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) \|a\|_{C_{ie}^{1,\alpha}(M \times [0, T], S_1)}^\flat. \end{aligned}$$

Analogous estimates hold for Hölder differences. Noting (7.16), we conclude

$$\|\psi(a^{-1} - a^{-1}(p, 0))\|_{C_{ie}^{k,\alpha}} \leq C\varepsilon^{-\alpha}(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) \|a\|_{C_{ie}^{k+1,\alpha}(M \times [0, T], S_1)}^\flat. \quad (7.19)$$

Now we estimate $\widetilde{\Delta}_L H_p[\varphi f]$. We first note that it vanishes identically at $t = 0$. This is obtained from the following mapping properties of the heat operator

$$\begin{aligned} H_p \circ \varphi : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S) &\rightarrow \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M \times [0, T], S), \\ H_p \circ \varphi : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S) &\rightarrow t^{\frac{\alpha}{2}} \mathcal{H}_{-2+\gamma}^{k+2,0}(M \times [0, T], S), \end{aligned}$$

where the first mapping property is due to (7.13) and the second is obtained by similar arguments, converting the lesser target regularity into a time weight. Hence $\widetilde{\Delta}_L H_p[\varphi f] \in \mathcal{H}_{-4+\gamma}^{k,0}$ vanishes identically at $t = 0$ and hence

$$\|\widetilde{\Delta}_L H_p[\varphi f]\|_{\mathcal{H}_{-4+\gamma}^{k,0}} = \|\widetilde{\Delta}_L H_p[\varphi f] - \widetilde{\Delta}_L H_p[\varphi f](t = 0)\|_{\mathcal{H}_{-4+\gamma}^{k,0}} \quad (7.20)$$

$$\leq CT^{\frac{\alpha}{2}} \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \leq C(\varepsilon^\alpha + T^{\frac{\alpha}{2}}) \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}}. \quad (7.21)$$

Analogous estimates hold for Hölder differences, if we take supremums over $\text{supp } \psi$. Hence overall we arrive at the estimate

$$\|\widetilde{\Delta}_L H_p[\varphi f]\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}(\text{supp } \psi \times [0, T], S)} \leq C(\varepsilon^\alpha + T^{\frac{\alpha}{2}}) \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}}. \quad (7.22)$$

Plugging the estimates (7.19) and (7.22) into (7.18), we conclude for $T/\varepsilon^2 < 1$

$$\begin{aligned} \|E^0 f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} &\leq C(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) \varepsilon^{-\alpha} (\varepsilon^\alpha + T^{\frac{\alpha}{2}}) \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \\ &\leq C(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) (1 + (T/\varepsilon^2)^{\frac{\alpha}{2}}) \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \\ &\leq C(\varepsilon^\gamma + T^{\frac{\alpha}{2}}) \|\varphi f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}}. \end{aligned}$$

This finishes the proof. \square

Proposition 7.8. *Let $\alpha \in (0, \min\{\gamma, 1\})$. For every $\delta > 0$, there exists $\varepsilon > 0$ and $T_0 > 0$ sufficiently small, such that the heat parametrix defined in (7.17)*

$$Q_b : \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S) \rightarrow \mathcal{H}_{-2+\gamma}^{k+2,\alpha}(M \times [0, T_0], S),$$

is a bounded linear map, solving

$$(\partial_t + a^{-1}\widetilde{\Delta}_L)(Q_b f) = \Phi f + E^0 f + E^1 f,$$

where E^0 and E^1 are bounded linear maps on $\mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$ with operator norms $\|E^0\| < \delta$ and $\|E^1\| \rightarrow 0$ as $T_0 \rightarrow 0^+$.

Proof. By Lemma 7.7 we compute for any $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$

$$(\partial_t + a^{-1}\widetilde{\Delta}_L)(Q_b f) = \sum_{j=1}^N \sum_{p \in L_j} (\partial_t + a^{-1}\widetilde{\Delta}_L)\psi_{j,p} H_p[\varphi_{j,p} f] = \Phi f + E^0 f + E^1 f,$$

where we have defined for $i = 0, 1$

$$E^i f := \sum_{j=1}^N \sum_{p \in L_j} E_{j,p}^i f.$$

The operators $E_{j,p}^i$, and hence also both E^0 and E^1 , are bounded operators on $\mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$ by Lemma 7.7. It remains to estimate their operator norms as $T_0 \rightarrow 0$. The fact that $\|E^1\| \rightarrow 0$ as $T_0 \rightarrow 0^+$ is a direct consequence of the second statement in Lemma 7.7. For the estimate of $\|E^0\|$ we argue as follows. We choose $\varepsilon > 0$ and $T_0 > 0$ with $T_0 < \varepsilon^2$. Then by Lemma 7.7 and (7.16) we have

$$\begin{aligned} \|E_{j,p}^0 f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} &\leq C(\varepsilon^\gamma + T_0^{\alpha/2}) \|\varphi_{j,p} f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \\ &\leq C\varepsilon^{-\alpha}(\varepsilon^\gamma + T_0^{\alpha/2}) \|f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \\ &= C(\varepsilon^{\gamma-\alpha} + \varepsilon^{-\alpha} T_0^{\alpha/2}) \|f\|_{\mathcal{H}_{-4+\gamma}^{k,\alpha}} \end{aligned}$$

Now set $\delta' := \frac{\delta}{\sum_j |L_j| N} > 0$ for a given $\delta > 0$. Then fix $\varepsilon > 0$ sufficiently small $C\varepsilon^{\gamma-\alpha} < \delta'/2$ (note that $\gamma - \alpha > 0$). For the fixed $\varepsilon > 0$ choose $T_0 > 0$ sufficiently small, such that $C\varepsilon^{-\alpha} T_0^{\alpha/2} < \delta'/2$. These choices yield $\|E^0\| < \delta$, and the proof is finished. \square

7.3.2. Construction of an interior parametrix. Next we construct an approximate interior parametrix. This construction is analogous to the one in [BAVe16], but for the convenience of the reader we repeat it here.

Recall that the radial function of the cone $\chi : \mathcal{C}(F) \rightarrow (0, 1)$ is extended smoothly to a nowhere vanishing function $\chi \in C^\infty(M)$. We assume that $\chi \geq 1$ outside of the singular neighborhood $\mathcal{C}(F)$. For $\varepsilon > 0$ small enough, $Y_\varepsilon := \{\chi \geq \varepsilon/2\} \subset M$ is a manifold with smooth boundary $\{\varepsilon/2\} \times F$. Let \bar{Y} denote the double of Y_ε . Since the Riemannian metric on \bar{Y} need not be smooth, we smoothen the metric in a small collar neighborhood of $\{\varepsilon/2\} \times F$, such that the metrics on \bar{Y} and M coincide over $Y_{2\varepsilon}$.

Since $\Phi \in C^\infty(\bar{M})$ is by construction identically 1 in $\{\chi < \varepsilon\}$, the function $1 - \Phi$ defines a smooth cutoff function on the closed double \bar{Y} , which is again denoted

by $1 - \Phi$. Let \bar{P} denote the extension of $P = \partial_t + \alpha^{-1}\widetilde{\Delta}_L$ to a uniformly parabolic operator on \bar{Y} . Note that $\mathcal{H}_{-4+\gamma}^{k,\alpha} \upharpoonright Y_\varepsilon \equiv \mathcal{C}_{ie}^{k,\alpha}(Y_\varepsilon \times [0, T_0], S)$. Let $\tilde{Q}_i : \mathcal{C}_{ie}^{k,\alpha}(Y_\varepsilon \times [0, T_0], S) \rightarrow \mathcal{C}_{ie}^{k+2,\alpha}(Y_\varepsilon \times [0, T_0], S)$ be the solution operator of the inhomogeneous Cauchy problem

$$\bar{P}u = (1 - \Phi)f, \quad u(\cdot, 0) = 0.$$

Also let χ be any smooth cutoff function which is identically 1 on $\text{supp}(1 - \Phi)$. Then we define the interior parametrix for any $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}$ by

$$Q_i f = \chi \tilde{Q}_i [(1 - \Phi)f].$$

7.3.3. Construction of the parametrix. From the approximate boundary and interior parametrices we obtain an approximate parametrix by

$$Qf := Q_b f + Q_i f$$

for $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$. From there we get a parametrix as follows.

Proof of Proposition 7.5. By Proposition 7.8 we conclude

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)Qf = \Phi f + E^0 f + E^1 f + (1 - \Phi)f + E^2 f =: f + Ef,$$

where $E^2 f := [\chi, \alpha^{-1}\widetilde{\Delta}_L]\tilde{Q}_i[(1 - \Phi)f]$. Now similarly as in Lemma 7.3 and Corollary 7.4 we find that $\|E^2\| \rightarrow 0$ as $T_0 \rightarrow 0$. Hence we can choose T_0 sufficiently small, such that the error term $Ef := E^0 f + E^1 f + E^2 f$ has operator norm less than 1. Then $I + E$, as an operator on $\mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T_0], S)$, is invertible, and we set

$$\mathcal{Q} := Q(I + E)^{-1}.$$

□

7.4. Extension of parametrices to the full time interval. In this subsection we extend the existence results of Theorems 7.5 and 7.6 from the shorter time interval $[0, T_0]$ to the full time interval $[0, T]$. By Theorem 7.5, for any $f \in \mathcal{H}_{-4+\gamma}^{k,\alpha}(M \times [0, T], S)$ there exists a $T_0 \in (0, T]$ and a solution $u = \mathcal{Q}f \in \mathcal{H}_{-2+\gamma}^{k,\alpha}(M \times [0, T_0], S)$ to the parabolic initial value problem

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)u = f, \quad u(t = 0) = 0.$$

If $T_0 < T$, we consider the homogeneous Cauchy problem

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)v_1 = 0, \quad v_1(t = 0) = u(t = T_0),$$

where the initial data $u(t = T_0) \in \mathcal{H}_{-2+\gamma}^{k,\alpha}(M, S)$. By Theorem 7.6, the solution to this problem, $v_1 = \mathcal{R}v_1(t = 0)$, exists on the time interval $[0, T_0]$ independent of the initial value $u(t = T_0)$. We may use Theorem 7.5 to solve

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)u_1 = f(p, t + T_0), \quad u_1(0) = 0,$$

on the interval $[T_0, 2T_0]$, and then the function

$$\tilde{u}(p, t) = \begin{cases} u(p, t) & \text{for } 0 \leq t \leq T_0 \\ u_1(p, t - T_0) + v_1(p, t - T_0) & \text{for } T_0 < t \leq 2T_0 \end{cases}$$

extends u past T_0 . This process continues until $nT_0 > T$, and produces a solution u in $\mathcal{H}_{-2+\gamma}^{k,\alpha}(M \times [0, T], S)$. Thus Theorem 7.5 and hence also Theorem 7.6 hold on the full time interval $[0, T]$.

Remark 7.9. *To see that the solution produced in this way indeed has the claimed regularity (in particular including Hölder regularity in time), we observe that instead of piecing together solutions on the time intervals $[0, T_0]$, $[T_0, 2T_0]$ etc. we could have instead chosen overlapping intervals $[0, T_0]$, $[T_0 - \varepsilon, 2T_0 - \varepsilon]$ etc. with a small $\varepsilon > 0$. Then these solutions agree on the overlaps by a uniqueness argument analogous to the proof of Theorem 7.10, based on the fact that the Friedrichs self adjoint extension $\widetilde{\Delta}_L$ is bounded from below.*

7.5. Regularity of Ricci curvature along the flow. We continue under the previously fixed notation where the upper script “ \sim ” indicates that the quantity is taken with respect to the initial metric. The following result proves our main Theorem 1.1.

Theorem 7.10. *Let (M, g) be a tangentially stable conical manifold of dimension at least 4, with $(\alpha, \gamma, k+1)$ Hölder regular geometry. Let $g(t)$, $t \in [0, T]$ be the solution of the Ricci de Turck flow with initial metric g and reference metric as in Theorem 5.1. Then, assuming $\gamma \in (0, 2 + \min\{\mu_0, \mu_1\})$, if $\widetilde{\text{Ric}} \in \mathcal{H}_{-2+\gamma}^{k,\alpha}(M \times [0, T], S)$, then $\text{Ric} \in \mathcal{H}_{-2+\gamma}^{k,\alpha}(M \times [0, T], S)$.*

Proof. By Proposition 7.2 we know

$$\begin{aligned} (\partial_t + \alpha^{-1}\widetilde{\Delta}_L)\text{Ric} &=: X(\text{Ric}) \in \mathcal{C}_{\text{ie}}^{k-1,\alpha}(M \times [0, T], S)_{-4+2\gamma} \\ &\subseteq \mathcal{H}_{-4+2\gamma}^{k,\alpha}(M \times [0, T], S). \end{aligned}$$

Consider the parametrix \mathcal{Q} of Theorem 7.5 and the parametrix \mathcal{R} of Theorem 7.6. We set $\gamma' := \min\{\gamma, 2\gamma\}$ and define

$$\text{Ric}' := \mathcal{Q}(X(\text{Ric})) + \mathcal{R}(\widetilde{\text{Ric}}) \in \mathcal{H}_{-2+\gamma'}^{k,\alpha}(M \times [0, T], S), \quad (7.23)$$

which is a solution of the parabolic initial value problem

$$(\partial_t + \alpha^{-1}\widetilde{\Delta}_L)\text{Ric}' = X(\text{Ric}), \quad \text{Ric}(t=0) = \widetilde{\text{Ric}}. \quad (7.24)$$

Due to tangential stability, the Friedrichs self adjoint extension $\widetilde{\Delta}_L$ is bounded from below. For $n \geq 3$ (note that we assume that $\dim M$ is at least 4) we find that $u := \text{Ric}' - \text{Ric} \in \mathcal{D}(\widetilde{\Delta}_L)$ and hence we may compute (note that $\alpha > 0$ and hence $\alpha^{-1}\widetilde{\Delta}_L \geq C$ for some lower bound $C \in \mathbb{R}$)

$$\partial_t \|u\|_{L^2(M, g_0)}^2 = -(\alpha^{-1}\widetilde{\Delta}_L u, u)_{L^2(M, g_0)} \leq C \|u\|_{L^2(M, g_0)}^2.$$

This differential inequality leads to the estimate

$$\|u\|_{L^2(M, g_0)}^2 \leq e^{Ct} \|u(0)\|_{L^2(M, g_0)}^2.$$

Since $u(t=0) \equiv 0$, we find that $u \equiv 0$. Hence we still conclude, despite having no maximum principle at hand

$$\text{Ric} \equiv \text{Ric}' \in \mathcal{H}_{-2+\gamma'}^{k,\alpha}(M \times [0, T], S).$$

We iterate the argument, improving the weight as long as $\gamma' < \gamma$. This proves the theorem. \square

8. POSITIVITY OF SCALAR CURVATURE ALONG THE RICCI DE TURCK FLOW

We can now prove our second main result on positivity of scalar curvature along the singular Ricci de Turck flow. At this final step, we will need a stronger tangential stability hypothesis. We impose, cf. Remark 5.2, the following additional

Assumption 8.1. We assume strong tangential stability: $u_0, u_1 > n$, i.e. $\square_L > n$, so that we may choose $\gamma_0, \gamma_1 \geq 1$ satisfying (5.1). This stronger condition is studied in the Appendix §8, where a list of examples is provided. Amongst the symmetric spaces of compact type, only

$$E_8, \quad E_7/[SU(8)/\{\pm I\}], \quad E_8/SO(16), \quad E_8/E_7 \cdot SU(2)$$

satisfy the strong tangential stability condition⁵.

The following theorem proves our second main result, Theorem 1.2.

Theorem 8.1. *Let (M, g) be a strong tangentially stable conical manifold of dimension at least 4, with $(\alpha, \gamma, k+1)$ Hölder regular geometry, where we assume $\gamma > 3$. Let $g(t), t \in [0, T]$ be the solution of the Ricci de Turck flow with initial metric g and reference metric as in Theorem 5.1. Assume that $R_g \geq 0$. Then $R_{g(t)} \geq 0$ for all $t \in [0, T]$. Furthermore, if R_g is positive at some point in the interior M , then $R_{g(t)}$ is positive in the interior M for all $t \in (0, T]$.*

Proof. By Theorem 7.10, $\text{Ric} \in \mathcal{H}_{-2+\gamma'}^{k, \alpha}(M \times [0, T], S)$, where γ' is any weight smaller than $\{\gamma, 2 + \mu_0, 2 + \mu_1\}$. Due to Assumption 8.1 of strong tangential stability, we find in the evolution equation (6.18) for the scalar curvature along the flow

$$\partial_t R + \Delta R = \langle W, \nabla R \rangle + 2|\text{Ric}|^2 \in \mathcal{C}_{\text{ie}}^\alpha(M \times [0, T]). \quad (8.1)$$

We want to express the Laplace Beltrami operator Δ in terms of $a^{-1}\tilde{\Delta}$, where $\tilde{\Delta}$ denotes the Laplace Beltrami operator with respect to $\tilde{g} \equiv g(0)$. Consider first the Laplace Beltrami operator for $\hat{g} = a\tilde{g}$,

$$\begin{aligned} \Delta_{\hat{g}} &= -\frac{1}{\det \hat{g}} \partial_i (\sqrt{\det \hat{g}} \cdot \hat{g}^{ij} \partial_j) = a^{-1} \tilde{\Delta} - \left(\frac{m}{2} - 1\right) \cdot \partial_i a \cdot a^{-2} \cdot \tilde{g}^{ij} \partial_j \\ &= a^{-1} \tilde{\Delta} - \{x^{-1} \mathcal{V}_b a, a\} x^{-1} \mathcal{V}_b, \end{aligned}$$

where $\{x^{-1} \mathcal{V}_b a, a\}$ refers to a linear combination of monomials consisting of the terms in the brackets. If we take into account higher order terms with $g = a\tilde{g} \oplus \omega$, we obtain in the same notation and higher regularity of R

$$\begin{aligned} \Delta R - a^{-1} \tilde{\Delta} R &= \{x^{-1} \mathcal{V}_b a, a\} x^{-1} \mathcal{V}_b R + \{x^{-1} \mathcal{V}_b \omega, x^{-1} \omega, \omega\} x^{-1} \mathcal{V}_b R \\ &\quad + \{a, \omega\} x^{-1} \mathcal{V}_b^2 R \in \mathcal{C}_{\text{ie}}^\alpha(M \times [0, T]). \end{aligned} \quad (8.2)$$

⁵We hope to lift that restriction in the forthcoming work.

The relation above is similar to Theorem 7.1, where we note that the Lichnerowicz Laplacian acting on the pure trace component S_1 coincides with the Laplace Beltrami operator. Combining (8.1) and (8.2), we conclude

$$\partial_t R + \alpha^{-1} \tilde{\Delta} R = P \in \mathcal{C}_{ie}^\alpha(M \times [0, T]). \quad (8.3)$$

By [BAVE16, Proposition 4.1, 4.6] there exists a solution $R' \in \mathcal{C}_{ie}^\alpha(M \times [0, T])$ of (8.3), for a given initial value $R(0) \in \mathcal{C}_{ie}^\alpha(M)$, such that $\Delta R' \in \mathcal{C}_{ie}^\alpha(M \times [0, T])$ again. Exactly as in Theorem 7.10, we find $R = R'$. Hence the maximum principle obtained in [BAVE16, Theorem 3.1] applies to R and, denoting by $R_{\min}(t)$ the minimum of the scalar curvature at time t , which is attained at $p_{\min}(t) \in \bar{M}$, we conclude

$$\begin{aligned} \partial_t R_{\min}(t) &\geq \langle W(p_{\min}(t)), \nabla R_{\min}(t) \rangle + 2|\text{Ric}(p_{\min}(t))|^2 \\ &\geq \langle W(p_{\min}(t)), \nabla R_{\min}(t) \rangle \end{aligned}$$

Now, if $p_{\min}(t)$ lies in the open interior of M , then $\nabla R_{\min}(t) = 0$. If $p_{\min}(t)$ lies at the conical singularity, we argue as follows: note that $\langle W, \nabla R \rangle \in \mathcal{C}_{ie}^\alpha(M \times [0, T])_{\gamma'-3}$. Since by strong tangential stability, we can choose $\gamma' > 3$, we find that $\langle W, \nabla R \rangle$ is vanishing at the conical singularity and hence

$$\partial_t R_{\min}(t) \geq \langle W(p_{\min}(t)), \nabla R_{\min}(t) \rangle = 0.$$

This implies that $R \geq 0$ for all $t \in [0, T]$. Now the last statement follows from the strong maximum principle [KLoo8, Theorem A.5]. \square

APPENDIX: CHARACTERIZING THE SPECTRAL CONDITIONS

In this appendix, we aim to characterize strong tangential stability in terms of eigenvalues of geometric operators on the cross-section of a cone. Note that this condition is only used in the last part of our argument, proving Theorem 1.1. It is not used in our first main result Theorem 1.2 on higher regularity of Ricci curvature.

Note that the operator \square_L can be entirely be described as an operator on the cross section (F^n, g_F) of the cone. Therefore in this section, all scalar products and geometric operators are taken with respect to (F^n, g_F) . The operator Δ_E denotes the Einstein operator on symmetric two-tensors over F . It is given by

$$\Delta_E \omega_{ij} = \Delta \omega_{ij} - 2g^{pq} \text{Rm}_{qij}^r \omega_{rp},$$

where Δ is the rough Laplacian. We write Δ for the Laplace Beltrami operator on F . Moreover, Π denotes the space of symmetric two-tensors which are trace-free and divergence-free at each point.

Theorem 8.2. *Let (F^n, g_F) , $n \geq 3$ be a compact Einstein manifold with constant $n - 1$. Then (F, g_F) is strongly tangentially stable if and only if we have the conditions*

$$\begin{aligned} \text{Spec}(\Delta_E|_{\Pi}) &> n, \\ \text{Spec}(\Delta_{\Omega^1(F) \cap \ker(\text{div})}) &> n + \sqrt{n^2 + 2n + 1}, \end{aligned}$$

and if for all positive eigenvalues λ of the Laplace-Beltrami operator on functions satisfies

$$\begin{aligned} & n(\lambda - 3n + 2)(\lambda + 4 - n)n(\lambda + n + 2) \\ & - 8n(n - 1)(\lambda - n)(\lambda + n + 2) - 8\lambda n(n + 1)(\lambda - 3n + 2) > 0. \end{aligned}$$

Proof. We first recall from the discussion before Remark 5.2 that strong tangential stability is equivalent to the two estimates

$$u_0 = \min(\text{Spec } \square'_L \setminus \{0\}) > n, \quad u_1 = \min(\text{Spec } \Delta_F \setminus \{0\}) > n.$$

However, the condition $u_1 > n$ holds for any Einstein metric except the sphere, where we have equality [OBA62]. For the rest of the proof, it thus suffices to consider the bundle $S_0 \upharpoonright F$. We use the same methodology as in [KrVe19A] which builds up on a decomposition of symmetric 2-tensors established in [KrÖ17]. We use the notation in [KrÖ17, Section 2] and the calculations in Section 3 of the same paper where we remove all terms containing radial derivatives in order to obtain expressions for the tangential operator. More precisely, we write

$$\begin{aligned} \{h_i\} & \text{ - basis of } L^2(TT), \quad \Delta_E h_i = \kappa_i h_i, \quad V_{1,i} := \langle r^2 h_i \rangle, \\ \{\omega_i\} & \text{ - basis of coclosed sections } L^2(T^*F), \quad \Delta \omega_i = \mu_i \omega_i, \\ V_{3,i} & := \langle r^2 \delta^* \omega_i \rangle \oplus \langle dr \odot r \omega_i \rangle, \\ \{v_i\} & \text{ - basis of } L^2(F), \quad \Delta v_i = \lambda_i v_i, \\ V_{4,i} & := \langle r^2 (n \nabla^2 v_i + \Delta v_i g) \rangle \oplus \langle dr \odot r \nabla v_i \rangle \\ & \oplus \langle v_i (r^2 g - n dr \otimes dr) \rangle. \end{aligned} \tag{8.4}$$

Here, Δ in $\Delta \omega_i$ denotes the connection Laplacian, while Δ in Δv_i denotes the Laplace Beltrami operator. The spaces $V_{1,i}, V_{3,i}, V_{4,i}$, with $L^2(0, 1)$ coefficients, span all trace-free sections $L^2(S_0 \upharpoonright F)$ over F , and are invariant under the action of the Lichnerowicz Laplacian. In [KrÖ17, Section 2], there is also a notion for the spaces $V_{2,i} := \langle v_i (r^2 g + n dr \otimes dr) \rangle$. But these spaces span the full trace sections $L^2(S_1 \upharpoonright F)$, whose discussion is not relevant here. At first, if $\tilde{h} = r^2 h_i \in V_{1,i}$,

$$(\square_L \tilde{h}, \tilde{h})_{L^2} = \kappa_i \|\tilde{h}\|_{L^2}^2$$

such that $\square_L > n$ on $V_{1,i}$ for all i if and only if $\kappa_i > n$ for all eigenvalues of the Einstein operator on TT -tensors are positive (non-negative).

Let now $\tilde{h} \in V_{3,i}$ so that it is of the form $\tilde{h} = \tilde{h}_1 + \tilde{h}_2 = \varphi r^2 \delta^* \omega_i + \psi dr \odot r \omega_i$. In this case, we have the scalar products

$$\begin{aligned} (\square_L \tilde{h}_1, \tilde{h}_1)_{L^2} &= \frac{\varphi^2}{2} (\mu_i - (n - 1))^2, \\ (\square_L \tilde{h}_2, \tilde{h}_2)_{L^2} &= \psi^2 [2\mu_i + (2n + 6)], \\ (\square_L \tilde{h}_1, \tilde{h}_2)_{L^2} &= -2(\mu_i - (n - 1))\psi\varphi. \end{aligned}$$

Taking $r^2 \delta^* \omega_i$ and $dr \odot r \omega_i$ as a basis, \square_L respects the subspace and acts as 2×2 -matrix

$$\begin{pmatrix} \frac{1}{2}(\mu_i - (n - 1))^2 & -2(\mu_i - (n - 1)) \\ -2(\mu_i - (n - 1)) & 2\mu_i + (2n + 6) \end{pmatrix}.$$

Because

$$\|\tilde{h}_1\|_{L^2}^2 = \frac{1}{2}(\mu_i - (n-1)) \cdot |\varphi|^2, \quad \|\tilde{h}_2\|_{L^2}^2 = 2|\psi|^2,$$

the operator $\square_L - n \cdot \text{id} > 0$ is represented by

$$A := \begin{pmatrix} \frac{1}{2}(\mu_i - (n-1))^2 - \frac{n}{2}(\mu_i - (n-1)) & -2(\mu_i - (n-1)) \\ -2(\mu_i - (n-1)) & 2\mu_i + 6 \end{pmatrix}.$$

The matrix A is positive definite if and only if the matrix

$$B := \begin{pmatrix} \frac{1}{2}(\mu_i - (2n-1)) & -2(\mu_i - (n-1)) \\ -2 & 2\mu_i + 6 \end{pmatrix}$$

is positive definite because A is obtained from B by multiplying the first column by $\mu_i - (n-1)$. By computing principal minors, this holds if

$$\det(B) = \frac{1}{2}(\mu_i - (2n-1))(2\mu_i + 6) - 4(\mu_i - (n-1)) > 0.$$

This in turn holds if

$$\mu_i > n + \sqrt{n^2 + 2n + 1}.$$

Therefore, $\square_L > n$ on the spaces $V_{3,i}$ if and only if $\Delta > n + \sqrt{n^2 + 2n + 1}$ on coclosed sections $L^2(T^*F)$. It remains to consider the case $\tilde{h} \in V_{4,i}$, so that it is of the form

$$\tilde{h} = \tilde{h}_1 + \tilde{h}_2 + \tilde{h}_3 = \varphi r^2(n\nabla^2 v_i + \Delta v_i g) + \psi dr \odot r\nabla v_i + \mathcal{X}v_i(r^2 g - n dr \otimes dr).$$

This case is the most delicate one. We have the scalar products

$$\begin{aligned} (\square_L \tilde{h}_1, \tilde{h}_1) &= n(n-1)\lambda_i(\lambda_i - n)(\lambda_i - 2(n-1))\varphi^2, \\ (\square_L \tilde{h}_2, \tilde{h}_2) &= [2\lambda_i(\lambda_i - (n-1)) + (2n+6)\lambda_i]\psi^2, \\ (\square_L \tilde{h}_3, \tilde{h}_3) &= [n\{(n+1)\lambda_i - 2(n+1)\} + 2n^2(n+3)]\mathcal{X}^2, \\ (\square_L \tilde{h}_1, \tilde{h}_2) &= -4(n-1)\lambda_i(\lambda_i - n)\psi\varphi, \\ (\square_L \tilde{h}_2, \tilde{h}_3) &= 4(n+1)\lambda_i\psi\mathcal{X}, \\ (\square_L \tilde{h}_1, \tilde{h}_3) &= 0 \end{aligned}$$

and the norms

$$\begin{aligned} \|\tilde{h}_1\|_{L^2}^2 &= n(n-1)\lambda_i(\lambda_i - n)\varphi^2, \\ \|\tilde{h}_2\|_{L^2}^2 &= 2\varphi^2\lambda_i, \\ \|\tilde{h}_3\|_{L^2}^2 &= (n+1)n. \end{aligned}$$

Consider $(\square_L - n \cdot \text{id})$. It acts as a matrix $A = (a_{ij})_{1 \leq i, j \leq 3}$, whose coefficients are given by

$$\begin{aligned} a_{11} &= n(n-1)\lambda_i(\lambda_i - n)[\lambda_i - 2(n-1) - n], \\ a_{22} &= 2\lambda_i[\lambda_i - (n-1) + 3], \\ a_{33} &= n\{(n+1)\lambda_i - 2(n-1) - n(n+1) + 2n(n+3)\}, \\ a_{12} &= a_{21} = -4(n-1)\lambda_i(\lambda_i - n), \\ a_{23} &= a_{32} = 4(n+1)\lambda_i, \\ a_{13} &= a_{31} = 0. \end{aligned}$$

In order to prove positivity of this matrix, we consider its principal minors A_{33} (which is the lower right entry), A_{23} (the lower right 2×2 -matrix) and A (the whole matrix). At first,

$$A_{33} = n\{(n+1)\lambda_i + n^2 + 3n + 2\} > 0.$$

Observe that in the case $\lambda_i = 0$, $\tilde{h}_1 \equiv 0$ and $\tilde{h}_2 \equiv 0$, so that $V_{4i} = \text{span}\{\tilde{h}_3\}$ and hence, $(\square_L - n \cdot \text{id})$ acts as $A_{33} > 0$. Therefore, we may from now on assume that $\lambda_i > 0$, which means that actually $\lambda_i \geq n$ (due to eigenvalue estimates for Einstein manifolds, see e.g. [OBA62]) with $\lambda_i = n$ only for \mathbb{S}^n . By considering the matrix

$$\begin{pmatrix} 2(\lambda_i + 4 - n) & 4(n+1)\lambda_i \\ 4(n+1) & n\{(n+1)\lambda_i + n^2 + 3n + 2\} \end{pmatrix},$$

from which one recovers A_{23} by multiplying the first column by λ_i , we see that

$$\begin{aligned} \frac{\det A_{23}}{\lambda_i} &= 2(\lambda_i + 4 - n) \cdot n \cdot [(n+1)\lambda_i + n^2 + 3n + 2] - 16(n+1)^2\lambda_i \\ &= 2n(n+1)\lambda_i^2 - 4(n+1)(n+4)\lambda_i - 2n(n-4)(n^2 + 3n + 2), \end{aligned}$$

which is positive if

$$\lambda_i > \frac{n+4}{n} + \sqrt{\frac{n+4}{n} + (n-4)(n+2)}.$$

Here, the right hand side is smaller than n if $n \geq 4$ such that this condition holds anyway. Before we compute the full determinant of A , we remark that in the case $\lambda_i = n$, the tensor \tilde{h}_1 is vanishing so that in this case, the matrix A describing \square_L on $V_{4,i}$ reduces to the matrix A_{23} which just has been considered. Therefore, there is nothing more to prove in this case and we may assume $\lambda = \lambda_i > n$ from now on. To compute the full determinant of A , we first consider the matrix

$$\begin{pmatrix} n[\lambda - 2(n-1) - n] & -2(n-1)(\lambda - n) & 0 \\ -4 & \lambda - (n-1) + 3 & 4\lambda \\ 0 & 2(n+1) & n[\lambda + 2(n+1) - n] \end{pmatrix}$$

from which we recover A by multiplying the three columns by $(n-1)\lambda(\lambda - n)$ and $2\lambda, (n+1)$, respectively. We get

$$\begin{aligned} [(n-1)\lambda(\lambda - n)2\lambda \cdot (n+1)]^{-1} \det A &= n(\lambda - 3n + 2)(\lambda + 4 - n)n(\lambda + n + 2) \\ &\quad - 8n(n-1)(\lambda - n)(\lambda + n + 2) - 8\lambda n(n+1)(\lambda - 3n + 2) \end{aligned}$$

which finishes the proof. \square

Theorem 8.3. *Amongst the symmetric spaces of compact type, only*

$$E_8, \quad E_7/[SU(8)/\{\pm I\}], \quad E_8/SO(16), \quad E_8/E_7 \cdot SU(2)$$

are strongly tangentially stable.

Proof. We merge and analyse the Tables 2 and 3 in [Krö17] and the Tables 1 and 2 in [CAHe15]. In the tables below, Λ denotes the smallest nonzero eigenvalue of the Laplace-Beltrami operator divided by the Einstein constant $n - 1$ and Θ denotes the smallest eigenvalue of the Lichnerowicz Laplacian Δ_L on symmetric 2-tensors with $\int_F \text{tr} h \, dV = 0$ divided by the same constant $n - 1$.

To check that the above mentioned spaces satisfy the condition of Theorem 8.2, we check on one hand that the estimate for λ in this theorem holds for $(n - 1) \cdot \Lambda$. On the other hand, we have the relations $\Delta_L = \Delta_E + 2(n - 1)\text{id}$, $\delta^* \circ \Delta_H = \Delta_L \circ \delta^*$ and $\Delta_H = \Delta + (n - 1)\text{id}$. Therefore, the condition $\text{Spec}(\Delta_E|_{\mathbb{T}\mathbb{T}}) > n$ holds, if $(n - 1)\Theta - 2(n - 1) > n$ and the condition $\text{Spec}(\Delta_{\Omega^1(F) \cap \ker(\text{div})}) > n + \sqrt{n^2 + 2n + 1}$ holds if $(n - 1)\Theta > 2n - 1 + \sqrt{n^2 + 2n + 1}$.

To show that all the other examples do not satisfy $\square_L > n$, we proceed as follows: We check that $(n - 1)\Lambda$ satisfies

$$\Lambda \leq \frac{n+4}{n} + \sqrt{\frac{n+4}{n} + (n-4)(n+2)}$$

or

$$\begin{aligned} & n(\Lambda - 3n + 2)(\Lambda + 4 - n)n(\Lambda + n + 2) \\ & - 8n(n - 1)(\Lambda - n)(\Lambda + n + 2) - 8\Lambda n(n + 1)(\Lambda - 3n + 2) \leq 0. \end{aligned}$$

This holds for example, if $\Lambda \leq 3$. Because $\Delta_L(f \cdot g_F) = \Delta f \cdot g_F$ and $\Delta_L \circ \nabla^2 = \nabla^2 \circ \Delta$, we clearly have $\Theta \leq \Lambda$. Because of the decomposition

$$C^\infty(\text{Sym}^2(T^*F)) = C^\infty(F) \cdot g_F \oplus \nabla^2(C^\infty(F)) \oplus \delta^*(\Omega^1(F) \cap \ker(\text{div})) \oplus \mathbb{T}\mathbb{T},$$

(c.f. [Krö17]) Θ is attained on

$$\delta^*(\Omega^1(F) \cap \ker(\text{div})) \oplus \mathbb{T}\mathbb{T}$$

if $\Theta < \Lambda$. In this case, we check if Θ satisfies $(n - 1)\Theta - 2(n - 1) \leq n$ and $(n - 1)\Theta < 2n - 1 + \sqrt{n^2 + 2n + 1}$ (both conditions are satisfied if $\Theta \leq 3$). Due to the relations above, this implies that either $\text{Spec}(\Delta_E|_{\mathbb{T}\mathbb{T}}) > n$ or $\text{Spec}(\Delta_{\Omega^1(F) \cap \ker(\text{div})}) > n + \sqrt{n^2 + 2n + 1}$ fails so that $\square_L > n$ fails. Therefore, the condition $\text{Spec}(\Delta_E|_{\mathbb{T}\mathbb{T}}) > n$ holds, if $(n - 1)\Theta - 2(n - 1) > n$ and the condition $\text{Spec}(\Delta_{\Omega^1(F) \cap \ker(\text{div})}) > n + \sqrt{n^2 + 2n + 1}$ holds if $(n - 1)\Theta > 2n - 1 + \sqrt{n^2 + 2n + 1}$.

To finish the proof, one just has to go through all the values of Θ and Λ in the tables two below. Table 1 lists the simple Lie groups while Table 2 lists all other symmetric spaces of non-group type. Strong tangential stability is abbreviated by STS. As was already said, for the spaces mentioned in the statement of the theorem, one manually checks the estimates above to verify that the conditions of Theorem 8.2 are satisfied. In all the other cases, one finds $\Theta \leq 3$ except in the

case $E_7/SO(12) \cdot SU(2)$ where one manually checks the condition on Λ mentioned above. This finishes the proof of the theorem. \square

type	G	dim(G)	Λ	Θ	STS
A_p	$SU(p+1), p \geq 2$	$p^2 - 1$	$\frac{2p(p+2)}{(p+1)^2}$	$\frac{2p(p+2)}{(p+1)^2}$	no
B_n	$Spin(5)$	10	$\frac{5}{3}$	$\frac{4}{3}$	no
	$Spin(7)$	21	$\frac{21}{10}$	$\frac{12}{5}$	no
	$Spin(2p+1), n \geq 4$	$2p(p+1)$	$\frac{4p}{2p-1}$	$\frac{4p}{2p-1}$	no
C_p	$Sp(p), p \geq 3$	$p(2p+1)$	$\frac{2p+1}{p+1}$	$\frac{4p-1}{2(p+1)}$	no
D_p	$Spin(2p), p \geq 3$	$p(2p+1)$	$\frac{2p-1}{p-1}$	$\frac{2p-1}{p-1}$	no
E_6	E_6	156	$\frac{26}{9}$	$\frac{17}{6}$	no
E_7	E_7	266	$\frac{19}{6}$	3	no
E_8	E_8	496	4	$\frac{47}{15}$	yes
F_4	F_4	52	$\frac{8}{3}$	$\frac{8}{3}$	no
G_2	G_2	14	2	2	no

TABLE 1. Conditions of Theorem 8.2 for simple Lie groups

type	G/K	dim(G/K)	Λ	Θ	STS
A I	$SU(p)/SO(p), 5 \geq p \geq 3$	$\frac{(p-1)(p+2)}{2}$	$\frac{2(p-1)(p+2)}{p^2}$	2	no
	$SU(p)/SO(p), p \geq 6$	$\frac{(p-1)(p+2)}{2}$	$\frac{2(p-1)(p+2)}{p^2}$	2	no
A II	$SU(4)/Sp(2) = S^5$	5	$\frac{5}{4}$	3	no
	$SU(2p)/Sp(p), p \geq 3$	$2p^2 - p - 1$	$\frac{(2p+1)(p-1)}{p^2}$	2	no
A III	$\frac{U(p+1)}{U(p) \times U(1)} = \mathbb{CP}^p$	$2p$	2	2	no
	$\frac{U(p+q)}{U(q) \times U(p)}, q \geq p \geq 2$	$2pq$	2	2	no
B I	$\frac{SO(5)}{SO(3) \times SO(2)}$	6	2	$\frac{4}{3}$	no
	$\frac{SO(2p+3)}{SO(2p+1) \times SO(2)}, p \geq 2$	$4p+2$	2	$\frac{8}{2p+1}$	no
	$\frac{SO(7)}{SO(4) \times SO(3)}$	12	$\frac{12}{5}$	$\frac{8}{5}$	no
	$\frac{SO(2p+3)}{SO(3) \times SO(2p)}, p \geq 3$	$6p$	$\frac{4p+6}{2p+1}$	$\frac{8}{2q+1}$	no
	$\frac{SO(2q+2p+1)}{SO(2q+1) \times SO(2p)}, p, q \geq 2$	$2n(2m+1)$	$\frac{4m+4n+2}{2m+2n-1}$	$\frac{8}{2p+2q-1}$	no
B II	$\frac{SO(2p+1)}{SO(2p)} = \mathbb{S}^{2p}, p \geq 1$	$2p$	$\frac{2p}{2p-1}$	$\frac{4p+2}{2p-1}$	no

type	G/K	dim(G/K)	Λ	Θ	STS
C I	$\mathrm{Sp}(p)/\mathrm{U}(p), p \geq 3$	$p(p+1)$	2	$\frac{2p}{p+1}$	no
C II	$\frac{\mathrm{Sp}(2)}{\mathrm{Sp}(1) \times \mathrm{Sp}(1)} = \mathbb{S}^4$	4	$\frac{4}{3}$	$\frac{10}{3}$	no
	$\frac{\mathrm{Sp}(p+1)}{\mathrm{Sp}(p) \times \mathrm{Sp}(1)} = \mathbb{H}\mathbb{P}^p, p \geq 2$	$4p$	$\frac{2(p+1)}{p+2}$	$\frac{2(p+1)}{p+2}$	no
	$\frac{\mathrm{Sp}(p+q)}{\mathrm{Sp}(q) \times \mathrm{Sp}(p)}, q \geq p \geq 2$	$4pq$	$\frac{2(p+q)}{p+q+1}$	$\frac{2(p+q)}{p+q+1}$	no
D I	$\frac{\mathrm{SO}(8)}{\mathrm{SO}(5) \times \mathrm{SO}(3)}$	15	$\frac{5}{2}$	$\frac{5}{2}$	no
	$\frac{\mathrm{SO}(2p+2)}{\mathrm{SO}(2p) \times \mathrm{SO}(2)}, p \geq 3$	$4p$	2	2	no
	$\frac{\mathrm{SO}(2p)}{\mathrm{SO}(p) \times \mathrm{SO}(p)}, p \geq 4$	p^2	$\frac{2p}{p-1}$	$\frac{2p}{p-1}$	no
	$\frac{\mathrm{SO}(2p+2)}{\mathrm{SO}(p+2) \times \mathrm{SO}(p)}, p \geq 4$	$p(p+2)$	$\frac{2p+2}{p}$	$\frac{2p+2}{p}$	no
	$\frac{\mathrm{SO}(2p)}{\mathrm{SO}(2p-q) \times \mathrm{SO}(q)}, p-2 \geq q \geq 3$	$(2p-q)q$	$\frac{2p}{p-1}$	$\frac{2p}{p-1}$	no
D II	$\frac{\mathrm{SO}(2p+2)}{\mathrm{SO}(2p+1)} = \mathbb{S}^{2p+1}, p \geq 3$	$2p+1$	$\frac{2p+1}{2p}$	$\frac{2(p+1)}{p}$	no
D III	$\mathrm{SO}(2p)/\mathrm{U}(p), p \geq 5$	$p(p-1)$	2	2	no
E I	$\mathrm{E}_6/[\mathrm{Sp}(4)/\{\pm I\}]$	42	$\frac{28}{9}$	3	no
E II	$\mathrm{E}_6/\mathrm{SU}(2) \cdot \mathrm{SU}(6)$	40	3	3	no
E III	$\mathrm{E}_6/\mathrm{SO}(10) \cdot \mathrm{SO}(2)$	32	2	2	no
E IV	$\mathrm{E}_6/\mathrm{F}_4$	26	$\frac{13}{9}$	$\frac{13}{9}$	no
E V	$\mathrm{E}_7/[\mathrm{SU}(8)/\{\pm I\}]$	70	$\frac{10}{3}$	$\frac{28}{9}$	yes
E VI	$\mathrm{E}_7/\mathrm{SO}(12) \cdot \mathrm{SU}(2)$	64	$\frac{28}{9}$	$\frac{28}{9}$	no
E VII	$\mathrm{E}_7/\mathrm{E}_6 \cdot \mathrm{SO}(2)$	54	2	2	no
E VIII	$\mathrm{E}_8/\mathrm{SO}(16)$	128	$\frac{62}{15}$	$\frac{16}{5}$	yes
E IX	$\mathrm{E}_8/\mathrm{E}_7 \cdot \mathrm{SU}(2)$	112	$\frac{16}{5}$	$\frac{16}{5}$	yes
F I	$\mathrm{F}_4/\mathrm{Sp}(3) \cdot \mathrm{SU}(2)$	28	$\frac{26}{9}$	$\frac{26}{9}$	no
F II	$\mathrm{F}_4/\mathrm{Spin}(9)$	16	$\frac{4}{3}$	$\frac{4}{3}$	no
G	$\mathrm{G}_2/\mathrm{SO}(4)$	8	$\frac{7}{3}$	$\frac{7}{3}$	no

TABLE 2. Conditions of Theorem 8.2 for symmetric spaces of non-group type

REFERENCES

- [AkBoo2] K. Akutagawa, B. Botvinnik, *The Relative Yamabe Invariant*, *Comm. Anal. Geom.* **10** (2002), no. 5, 935–969
- [BAVe14] Eric Bahuaud and Boris Vertman, *Yamabe flow on manifolds with edges*, *Math. Nachr.* **287** (2014), no. 23, 127–159
- [BAVe16] Eric Bahuaud and Boris Vertman, *Long time existence of the edge Yamabe flow*, *J. Math. Soc. Japan.* **71** (2019), no. 2, 651–688.
- [CAHe15] Huai-Dong Cao and Chenxu He, *Linear Stability of Perelmans ν -entropy on Symmetric spaces of compact type*, *J. Reine Angew. Math.* **709** (2015), no. 1, 229–246
- [CHWa15] Xiuxiong Chen and Yuanqi Wang, *Bessel functions, heat kernel and the conical Kähler - Ricci flow*, *J. Funct. Anal.* **269**, no. 2 (2015) 551–632
- [CHKn04] B. Chow, D. Knopf, *The Ricci Flow: An Introduction*, *Mathematical Surveys and Monographs*, Vol. 110. American Mathematical Society, Providence, RI, 2004
- [DON12] Simon K. Donaldson, *Kähler metrics with cone singularities along a divisor*, *Essays in Math. and Appl.*, p.49-79, Springer Heidelberg (2012)
- [GiTo10] Gregor Giesen and Peter M. Topping, *Ricci flow of negatively curved incomplete surfaces*, *Calc. Var. Partial Differ. Equ.* **38** (2010), no. 3-4, 357–367
- [GiTo11] ———, *Existence of Ricci flows of incomplete surfaces*, *Comm. Partial Differential Equations* **36** (2011), no. 10, 1860–1880
- [JMR16] Thalia Jeffres, Rafe Mazzeo, and Yanir Rubinstein, *Kähler-Einstein metrics with edge singularities*, *Ann. of Math.* **183** (2016), no. 1, 95–176
- [KLPO8] Klaus Kirsten, Paul Loya and Jinsung Park, *Functional determinants for general self-adjoint extensions of Laplace-type operators resulting from the generalized cone*, *Manuscripta Math.* **125** (2008), no. 1, 95–126.
- [KLoo8] B. Kleiner, J. Lott, *Notes on Perelman’s papers*, *Geometry and Topology* **12** (2008) 2587–2858
- [KRö17] K. Kröncke, *Stable and unstable Einstein warped products*, *Trans. Amer. Math. Soc.* **369** (2017) no. 9, 6537–6563
- [KRVe19A] Klaus Kröncke and Boris Vertman, *Stability of Ricci de Turck flow on Singular Spaces*, *Calc. Var. Partial Differ. Equ.* **58** (2019), no. 2, 74
- [KRVe19B] ———, *Perelman’s Entropies for Manifolds with conical Singularities*, preprint on arXiv:1902.02097 [math.DG] (2019)
- [LiZh17] Jiawei Liu and Xi Zhang, *Conical Kähler-Ricci flows on Fano manifolds*, *Adv. Math.* **307** (2017), 1324–1371
- [MRS15] Rafe Mazzeo, Yanir Rubinstein, and Natasha Sesum, *Ricci flow on surfaces with conic singularities*, *Anal. PDE* **8** (2015), no. 4, 839–882
- [OBA62] M. Obata, *Certain conditions for a Riemannian manifold to be isometric with a sphere*, *J. Math. Soc. Japan* **14** (1962), 333–340
- [SIM13] Miles Simon, *Local smoothing results for the Ricci flow in dimensions two and three*, *Geom. Topol.* **17** (2013), no. 4, 2263–2287
- [RiNA90] Frigyes Riesz and Béla Sz.-Nagy, *Functional analysis*. Translated from the second French edition by Leo F. Boron. Reprint of the 1955 original. Dover Books on Advanced Mathematics. Dover Publications, Inc., New York, 1990
- [Tia15] Gang Tian, *K-stability and Kähler-Einstein metrics*, *Comm. Pure Appl. Math.* **68** (2015), no. 7, 1085–1156
- [TOPPO6] P. Topping, *Lectures on the Ricci flow*, <http://homepages.warwick.ac.uk/maseq/RFnotes.html> (2006)
- [VER16] B. Vertman, *Ricci flow on singular manifolds*, arXiv:1603.06545v3 [math.DG] (2016)
- [WAN16] Yuanqi Wang, *Smooth approximations of the conical Kähler - Ricci flows*, *Math. Ann.* **365** (2016), no. 1-2, 835–856
- [YIN10] Hao Yin, *Ricci flow on surfaces with conical singularities*, *J. Geom. Anal.* **20** (2010), no. 4, 970–995.

UNIVERSITY HAMBURG, GERMANY

E-mail address: klaus.kroencke@uni-hamburg.de

UNIVERSITY OLDENBURG, GERMANY

E-mail address: tobias.marxen@uni-oldenburg.de

UNIVERSITY OLDENBURG, GERMANY

E-mail address: boris.vertman@uni-oldenburg.de