

ON THE NULLSTELLENSATZ FOR C-HOLOMORPHIC FUNCTIONS WITH ALGEBRAIC GRAPHS

ADAM BIAŁOŻYT, MACIEJ P. DENKOWSKI, PIOTR TWORZEWSKI

ABSTRACT. C-holomorphic functions defined on algebraic sets and having algebraic graphs can be considered as a complex counterpart of *regular* functions introduced recently in real geometry. This note is a part of our study on the subject; we prove herein some effective Nullstellensätze.

1. INTRODUCTION

This article is a second one, after [BDTW], of a series devoted to continuous functions defined on a given algebraic subset of a complex finite-dimensional vector space M and having algebraic graphs, i.e. the class of c-holomorphic functions with algebraic graphs. They can be seen as a kind of complex counterpart of the recently introduced *regular* functions [FHMM], [Kol] in connection with [K].

To simplify the notation, we will consider $M = \mathbb{C}^m$ (anyway, everything here is invariant under linear isomorphisms). In this paper we study some basic Nullstellensätze in this class. Dealing with such functions requires the use of purely geometric methods.

For the convenience of the reader we recall some basic facts. Let $A \subset \Omega$ be an analytic subset of an open set $\Omega \subset \mathbb{C}^m$. R. Remmert generalized the notion of holomorphic mapping onto sets having singularities in a more convenient way (from the geometric point of view) than the usual notion of *weakly holomorphic functions* (i.e. functions defined and holomorphic on $\text{Reg}A$ and locally bounded on A), namely:

Definition 1.1. (cf. [R]) A mapping $f: A \rightarrow \mathbb{C}^n$ is called *c-holomorphic* if it is continuous and the restriction of f to the subset of regular points $\text{Reg}A$ is holomorphic. We denote by $\mathcal{O}_c(A, \mathbb{C}^n)$ the vector space of c-holomorphic mappings, and by $\mathcal{O}_c(A)$ the ring of c-holomorphic functions.

The corner stone of all our further considerations is the following theorem:

Theorem 1.2 ([Wh] 4.5Q). *A mapping $f: A \rightarrow \mathbb{C}^n$ defined on an analytic set $A \subset \mathbb{C}^m$ is c-holomorphic iff it is continuous and its graph $\Gamma_f := \{(x, f(x)) \mid x \in A\}$ is an analytic subset of $\Omega \times \mathbb{C}^n$.*

Date: January 28th 2020.

1991 Mathematics Subject Classification. 32B15, 32A17, 32A22.

Key words and phrases. Complex analytic and algebraic sets, c-holomorphic functions, Nullstellensatz.

For a more detailed list of basic properties of c-holomorphic mappings see [Wh], [D1] and [BDTW]. It is easy to see that for $f \in \mathcal{O}_c(A, \mathbb{C}^n)$, the graph Γ_f is irreducible iff A is irreducible.

Finally, we recall some notions we will be using. If $\Gamma \subset \mathbb{C}^n$ is algebraic of pure dimension k , then $\deg \Gamma = \#(L \cap \Gamma)$ for any $L \subset \mathbb{C}^n$ affine subspace of dimension $n - k$ transversal to Γ and such that $L_\infty \cap \bar{\Gamma} = \emptyset$, where $\bar{\Gamma}$ is the projective closure and L_∞ denotes the points of L at infinity (see [L] VII.§7 and [Ch]). The condition $L_\infty \cap \bar{\Gamma} = \emptyset$ is equivalent both to $L_\infty \cap \Gamma_\infty = \emptyset$ and to the inclusion

$$\Gamma \subset \{u + v \in L' + L \mid \|v\| \leq \text{const.}(1 + \|u\|)\}$$

where L' is any k -dimensional affine subspace such that $L' + L = \mathbb{C}^n$. Moreover, for any $(n - k)$ -dimensional affine subspace L cutting A in a zero-dimensional set, $\#(L \cap \Gamma) \leq \deg \Gamma$.

We write $G'_{n-k}(\mathbb{C}^n)$ for the set of affine hyperplanes of dimension $n - k$ and we have

$$\deg \Gamma = \max\{\#(L \cap \Gamma) \mid L \in G'_{n-k}(\mathbb{C}^n): \dim(L \cap \Gamma) = 0\}.$$

A natural subclass of c-holomorphic functions that we can expect will behave like polynomials or regular functions is formed by those c-holomorphic function whose graph is algebraic.

Definition 1.3. We will call *c-algebraic* any continuous function $f: A \rightarrow \mathbb{C}$ having an algebraic graph, where $A \subset \mathbb{C}^m$ is a fixed analytic set. We will denote by

$$\mathcal{O}_c^a(A) = \{f \in \mathcal{O}_c(A) \mid \Gamma_f \text{ is algebraic}\}$$

the ring of such functions and write $f = (f_1, \dots, f_n) \in \mathcal{O}_c^a(A, \mathbb{C}^n)$ whenever all the components f_j are c-algebraic which is clearly equivalent to f having an algebraic graph.

Remark 1.4. Of course, by the Chevalley-Remmert Theorem

$$\mathcal{O}_c^a(A) \neq \emptyset \Rightarrow A \text{ is algebraic.}$$

Example 1.5. The eternal example of a c-algebraic but non-regular function is $f(x, y) = y/x$ on $A: y^2 = x^3$ extended at the origin of \mathbb{C}^2 by putting $f(0, 0) = 0$.

As observed in [BDTW] each c-algebraic function on an algebraic pure dimensional set is the restriction of a regular function (this is the link to [FHMM]).

From now on we will assume that $A \subset \mathbb{C}^m$ is a pure k -dimensional algebraic set of degree $d := \deg A$. Obviously, we shall assume also $k \geq 1$ unless something else is stated.

Let $\|\cdot\|$ denote one of the usual norms in \mathbb{C}^m . In [BDTW] we studied the *growth exponent* at infinity that generalizes the notion of degree of a polynomial for $f \in \mathcal{O}_c^a(A)$:

$$\mathcal{B}(f) := \inf\{s \geq 0 \mid |f(x)| \leq \text{const.}\|x\|^s, x \in A: \|x\| \gg 1\}$$

where the defining inequality is equivalent to

$$(*) \quad |f(x)| \leq \text{const} \cdot (1 + \|x\|)^s, \quad x \in A$$

and this characterises $\mathcal{O}_c^a(A)$ in $\mathcal{O}_c(A)$, cf. [BDTW] Lemma 3.1. In the polynomial case, i.e. $A = \mathbb{C}^k$, we have $\mathcal{B}(f) = \deg f$, for c -algebraic functions are polynomials by the Serre Graph Theorem.

2. CHARACTERISTIC POLYNOMIALS

In the setting of c -algebraic functions we are obliged to make do more with the geometric structure than the algebraic one which is a basic hindrance when trying to transpose the results known, for instance, for polynomial dominating mappings to the c -holomorphic algebraic case.

We consider now the following situation:

Let $A \subset \mathbb{C}^m$ be algebraic of pure dimension $k > 0$ and suppose $f \in \mathcal{O}_c^a(A, \mathbb{C}^k)$ is a proper mapping.

Since Γ_f is algebraic with proper projection onto \mathbb{C}^k , the cardinality of the fibre $\#f^{-1}(y)$ is constant for the generic point $y \in \mathbb{C}^k$. Following [BDTW] we denote this number by $d(f)$ and call it the *geometric degree* of f just as in the polynomial case. We call *critical* for f any point $y \in \mathbb{C}^k$ for which $\#f^{-1}(w) \neq d(f)$. In that case one has actually $\#f^{-1}(w) < d(f)$. Obviously $d(f) \leq \deg \Gamma_f$ (cf. [L]).

In the sequel we shall use intensively the notion of characteristic polynomial relative to a proper mapping f . For any $g \in \mathcal{O}_c^a(A)$ let us introduce the *characteristic polynomial* of g relative to f : for any $y \in \mathbb{C}^k$ not critical for f we put

$$(\#) \quad P_g(y, t) := \prod_{x \in f^{-1}(y)} (t - g(x)) = t^{d(f)} + a_1(y)t^{d(f)-1} + \dots + a_{d(f)}(y)$$

extending the coefficients through the critical locus of f thanks to the Riemann Extension Theorem (they are continuous; see below their form). Therefore $P_g \in \mathcal{O}(\mathbb{C}^k)[t]$.

Proposition 2.1. *In the introduced setting, P_g is a pure-bred polynomial, i.e. $P_g \in \mathbb{C}[y_1, \dots, y_k][t]$.*

Proof. This follows directly from the expressions for the coefficients:

$$a_j(y) = (-1)^j \sum_{1 \leq \iota_1 < \dots < \iota_j \leq d(f)} g(x^{(\iota_1)}) \cdot \dots \cdot g(x^{(\iota_j)}),$$

where $f^{-1}(y) = \{x^{(1)}, \dots, x^{(d(f))}\}$ consists of $d(f)$ points.

Since $g \in \mathcal{O}_c^a(A)$, there is $|g(x)| \leq C_1(1 + \|x\|^r)$ for $x \in A$ with some constants $C_1, r > 0$ (cf. (*) and [BDTW] Lemma 3.1). By assumption, Γ_f has proper projection onto \mathbb{C}^k and so by the Rudin-Sadullaev Theorem (see [L]),

$$\Gamma_f \subset \{(x, y) \in \mathbb{C}^m \times \mathbb{C}^k \mid \|x\| \leq C_2(1 + \|y\|^s)\}$$

for some constants $C_2, s > 0$. Therefore, for any $x \in A$, $\|x\| \leq C_2(1 + \|f(x)\|)^s$. We obtain thus

$$|g(x)| \leq C_1(1 + C_2^r(1 + \|f(x)\|)^{rs}) \leq C_1 2 \max\{1, C_2^r\}(1 + \|f(x)\|)^{rs}.$$

This means in particular that for any y not critical for f , and for all j ,

$$|a_j(y)| \leq \text{const.}(1 + \|y\|)^{rsj},$$

since $y = f(x^{(j)})$. By continuity this inequality can be extended to the whole of \mathbb{C}^k and so by Liouville's Theorem $a_j \in \mathbb{C}[y_1, \dots, y_k]$ for all j . \square

Remark 2.2. In the proof above we may put $r = \mathcal{B}(g)$. On the other hand, by [TW1], we can also take $s = \deg \Gamma_f - d(f) + 1$. Therefore, we obtain

$$\deg a_j \leq j\mathcal{B}(g)(\deg \Gamma_f - d(f) + 1).$$

The following Lemma of Płoski is crucial when using the characteristic polynomial:

Lemma 2.3 ([P1] lemma (2.1)). *$P(x, t) = t^d + a_1(x)t^{d-1} + \dots + a_d(x)$ be a polynomial with $a_j \in \mathbb{C}[x_1, \dots, x_k]$. Then $\delta(P) := \max_{j=1}^d (\deg a_j/j)$ is the minimal exponent $q > 0$ such that*

$$\{(x, t) \in \mathbb{C}^k \times \mathbb{C} \mid P(x, t) = 0, |x| \geq R\} \subset \{(x, t) \in \mathbb{C}^k \times \mathbb{C} \mid |t| \leq C|x|^q\}$$

for some $R, C > 0$.

We present a simple application of the characteristic polynomial in the polynomial case. Let $f: \mathbb{C}^m \rightarrow \mathbb{C}$ be a polynomial of degree d and such that the complex gradient (with respect to $\partial/\partial z_j$) $\nabla f: \mathbb{C}^m \rightarrow \mathbb{C}^m$ is proper with multiplicity μ ⁽¹⁾. Let $P(y, t)$ be the characteristic polynomial of f with respect to ∇f . By Proposition 2.1 it is a polynomial $P \in \mathbb{C}[y_1, \dots, y_m, t]$ of the form

$$P(y, t) = \prod_{x \in \nabla f^{-1}(y)} (t - f(x)) = t^\mu + a_1(y)t^{\mu-1} + \dots + a_\mu(y).$$

Let us write $D = \deg \Gamma_{\nabla f}$. Obviously, $D \geq \mu$.

Proposition 2.4. *In the setting introduced above,*

$$|f(x)|^\Theta \leq \text{const.} \|\nabla f(x)\|, \quad \|x\| \gg 1$$

where

$$\Theta = \frac{1}{d(D - \mu + 1)} \in \left(0, \frac{1}{d}\right].$$

Proof. This follows essentially from Lemma and Remark 2.2, but we may also give a direct argument. Clearly, $f(x)$ is a root of $P(\nabla f(x), \cdot)$. Therefore,

$$\|f(x)\| \leq 2 \max_{j=1}^\mu |a_j(\nabla f(x))|^{\frac{1}{j}}.$$

¹Which is equivalent to saying that the fibres are finite and the sum of the local multiplicities always gives μ .

By Remark 2.2 we have

$$|a_j(\nabla f(x))|^{\frac{1}{j}} \leq \text{const} \cdot \|\nabla f(x)\|^{d(D-\mu+1)}$$

whence the result. \square

3. NULLSTELLENSÄTZE AND DEGREE OF CYCLES OF ZEROES

We shall deal first with the 0-dimensional case, i.e. we assume that $f = (f_1, \dots, f_n)$ is a proper c -algebraic map on a set $A \subset \mathbb{C}^m$ of pure dimension $k > 0$ (it follows that $n \geq k$). We assume that $f^{-1}(0) \neq \emptyset$. Then the intersection of the graph Γ_f with $\mathbb{C}^m \times \{0\}^n$ is isolated but possibly improper ⁽²⁾. We may compute the improper intersection indices

$$i_a(f) := i(\Gamma_f \cdot (\mathbb{C}^m \times \{0\}^n); a), \quad a \in f^{-1}(0)$$

along [ATW] (see also [D3]).

The properness of f implies by the Chevalley-Remmert Theorem that $f(A)$ is algebraic. If it happens to be *irreducible*, then we can define the *generalized geometric degree* $d(f)$ to be the degree (sheet number) of the branched covering (cf. [Ch]) $f|_{A \setminus f^{-1}(\text{Sng}f(A))}$ over the connected manifold $\text{Reg}f(A)$.

Example 3.1. It should be pointed out that although $d(f)$ corresponds to the cardinality of the generic nonempty fibre of f , it may not be the greatest cardinality of the fibre: if A denotes the graph of $t \mapsto (t^2 - 1, t(t^2 - 1))$ in \mathbb{C}^3 and f is the projection onto the last two coordinates, then $d(f) = 1$, whereas $\#f^{-1}(0) = 2$.

For $n = k$, which means that the intersection is proper and f is surjective, the numbers $i_a(f)$ coincide with the usual local geometric multiplicities $m_a(f)$ ⁽³⁾ and the Stoll formula yields $d(f) = \sum_{a \in f^{-1}(0)} m_a(f)$. Thanks to a Spodzieja-type reduction as in [Sp], we obtain the following general c -algebraic counterpart of a result of E. Cygan [Cg]:

Theorem 3.2. *In the setting introduced above, let $g \in \mathcal{O}_c^a(A)$ be such that $g^{-1}(0) \supset f^{-1}(0)$. Then, if $f(A)$ is irreducible, there are n functions $h_j \in \mathcal{O}_c^a(A)$ such that*

$$g^{d(f) \deg f(A)} = \sum_{j=1}^n h_j f_j \quad \text{on the whole of } A.$$

Moreover,

$$d(f) \deg f(A) \geq \sum_{a \in f^{-1}(0)} i_a(f)$$

²Recall that the intersection $X \cap Y$ of two pure-dimensional analytic sets X, Y in a manifold M is called proper, if at any $a \in X \cap Y$ it has the minimal possible dimension $\dim X + \dim Y - \dim M$.

³I.e. $m_a(f) = \limsup_{v \rightarrow f(a)} \#(U \cap f^{-1}(v))$ where U is a neighbourhood of a such that $\overline{U} \cap f^{-1}(f(a)) = \{a\}$ (the result is the same for all U small enough); see [Ch].

and equality holds, provided $k = n$ (in which case $\deg f(A) = 1$).

In the proof we will need the following refinement of [D3] Theorem 2.3:

Proposition 3.3. *Let $A \subset \mathbb{C}^m \times D$ be a pure k -dimensional analytic set with proper projection $p|_A$ where $p: \mathbb{C}^m \times D \ni (x, y) \mapsto y \in D$ and $D \subset \mathbb{C}^n$ is a domain. Assume that $0 \in X := p(A)$ is irreducible and let μ denote the multiplicity of the branched covering $p|_{A \setminus p^{-1}(\text{Sng} X)}$ over the connected manifold $\text{Reg} X$. Then*

$$\mu \cdot \deg_0 X = \sum_{a \in p^{-1}(0) \cap A} i(A \cdot \text{Ker } p; a),$$

where $\deg_0 X$ stands for the local degree (Lelong number) of X at the origin.

Remark 3.4. Asearlier we are dealing here with a possibly improper isolated intersection $A \cap \text{Ker } p$ and we are using [ATW] to compute the intersection multiplicities which we may denote as earlier $i_a(p|_A) := i(A \cdot \text{Ker } p; a)$.

Proof of Proposition 3.3. We can find a neighbourhood U of $0 \in \mathbb{C}^n$ and pairwise disjoint neighbourhood V_1, \dots, V_ν ($\nu \leq \mu$) of the points in the fibre $p^{-1}(0) \cap A$ such that $U \cap X = \bigcup_{j=1}^r X_j$ corresponds to the decomposition of the germ $(X, 0)$ into irreducible components and $p^{-1}(U) = \bigcup_{i=1}^\nu V_i \cap A$. We regroup the sets

$$W_j := \bigcup_{s=1}^{t_j} V_{i_s} \cap A$$

according to their projection, i.e. to obtain W_j we take all the sets $V_{i_s} \cap A$, $s = 1, \dots, t_j$, for which $p(V_i \cap A) = X_j$. We get as much sets W_j as there are components X_j and $p|_{W_j}: W_j \rightarrow X_j$ is proper and becomes a μ -sheeted branched covering over $\text{Reg} X_j$.

Moreover, the points of $p^{-1}(0) \cap A$ form groups $\{a_{i_1}, \dots, a_{i_{t_j}}\} = p^{-1}(0) \cap W_j$ and we may compute the *regular multiplicity* (cf.[D3])

$$\tilde{m}_{a_{i_s}}(p|_A) = \limsup_{\text{Reg} \ni y \rightarrow 0} \#p^{-1}(0) \cap A \cap V_{i_s} = \tilde{m}_{a_{i_s}}(p|_{W_j}),$$

provided V_{i_s} is small enough (which, of course, we may assume with no harm for the generality). Then, clearly, $\mu = \sum_{s=1}^{t_j} \tilde{m}_{a_{i_s}}(p|_A)$. Finally, by [D3] Theorem 2.3,

$$\tilde{m}_{a_{i_s}}(p|_{W_j}) \cdot \deg_0 X_j = i_a(p|_{W_j})$$

and $i_{a_{i_s}}(p|_{W_j}) = i_{a_{i_s}}(p|_A)$. Therefore,

$$\begin{aligned} \sum_{i=1}^\nu i_{a_i}(p|_A) &= \sum_{j=1}^r \sum_{s=1}^{t_j} (\tilde{m}_{a_{i_s}}(p|_{W_j}) \cdot \deg_0 X_j) = \\ &= \sum_{j=1}^r \mu \cdot \deg_0 X_j = \mu \cdot \deg_0 X \end{aligned}$$

as required. □

On the Nullstellensatz for c-holomorphic functions with algebraic graphs

Proof of Theorem 3.2. First, let us deal with the proper intersection case, i.e. let us assume that $k = n$. Let P_g be the characteristic polynomial of g relative to f as in (#). From the definition we have clearly $P_g(f(x), g(x)) = 0$ for $x \in A$, which means

$$g(x)^{d(f)} = -a_1(f(x))g(x)^{d(f)-1} + \dots - a_{d(f)}(f(x)).$$

Now, for any j , $a_j \in \mathbb{C}[y_1, \dots, y_k]$ and since $g = 0$ on $f^{-1}(0)$, it follows from the expression of a_j (see the proof of Proposition 2.1) that $a_j(0) = 0$ for any j . Therefore $a_j(y) = \sum_{\iota=1}^k a_{j,\iota}(y)y_\iota$ with $a_{j,\iota} \in \mathbb{C}[y_1, \dots, y_k]$ and the assertion follows.

Now, assume that $k > n$ and write $X := f(A)$. Then X is an irreducible k -dimensional algebraic subset of \mathbb{C}^n and for the generic epimorphism $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^k$ we have $d(\pi|_X) = \deg X$. Consider the proper c-algebraic map $\tilde{f} := \pi \circ f$. From the first part of the proof we get

$$g^{d(\tilde{f})} = \sum_{j=1}^k h_j \tilde{f}_j,$$

but $\tilde{f}_j = (\pi \circ f)_j = (\pi_j \circ f)$ and $\pi_j(y) = \sum_{i=1}^n \alpha_{ji} y_i$ which yields the assertion sought after, provided we show $d(\tilde{f}) = d(f) \cdot \deg X$. But for the generic $x \in \mathbb{C}^k$, the fibre $\pi^{-1}(x)$ lies entirely in $\text{Reg} A$ and does not meet the critical set of the $d(f)$ -sheeted branched covering $f|_{A \setminus f^{-1}(\text{Sng} X)}$ over the connected manifold $\text{Reg} X$, whence it follows that $d(\tilde{f}) = d(f) \cdot \deg X$.

To prove the second part of the assertion we may assume that the coordinates in \mathbb{C}^n are chosen so that π is the projection onto the first k of them. Put $T := \mathbb{C}^m \times \{0\}^k$. Then, by the Stoll Formula

$$d(\tilde{f}) = d(p|_{\Gamma_{\tilde{f}}}) = \sum_{a \in T \cap \Gamma_{\tilde{f}}} i(T \cdot \Gamma_{\tilde{f}}; a)$$

where now the intersection multiplicities $i(T \cdot \Gamma_{\tilde{f}}; a)$ are computed along Draper, as we are dealing with an isolated proper intersection. In particular $d(p|_{\Gamma_{\tilde{f}}}) \geq \sum_{b \in f^{-1}(0)} i(T \cdot \Gamma_{\tilde{f}}; (b, 0))$ where $0 \in \mathbb{C}^k$.

Observe that $(\text{id}_{\mathbb{C}^m} \times \pi)|_{\Gamma_f} = \Gamma_{\tilde{f}}$ and this is a one to one correspondence. Put $N := \mathbb{C}^m \times \{0\}^k \times \mathbb{C}^{n-k}$. It is classical in proper intersection theory that $i(\Gamma_{\tilde{f}} \cdot T; (b, 0)) = m_{(b,0)}(\tau|_{\Gamma_{\tilde{f}}})$ for the linear projection τ such that $\text{Ker } \tau = T$. Similarly, $i(\Gamma_f \cdot N; (b, 0)) = m_{(b,0)}(p|_{\Gamma_f})$ (here $0 \in \mathbb{C}^n$) for the linear projection p with $\text{Ker } p = N$. Now, if $z \in \mathbb{C}^k$, then the fibres $\tau^{-1}(z) \cap \Gamma_{\tilde{f}} \cap (U \times V)$ and $p^{-1}(z) \cap \Gamma_f \cap (U \times V \times W)$ are bijective by $\text{id}_{\mathbb{C}^m} \times \pi$, where $b \in U \subset \mathbb{C}^m$, $0 \in V \subset \mathbb{C}^k$, $0 \in W \subset \mathbb{C}^{n-k}$ are sufficiently small neighbourhoods. This implies that $m_{(b,0)}(\tau|_{\Gamma_{\tilde{f}}}) = m_{(b,0)}(p|_{\Gamma_f})$ and we are done. \square

Remark 3.5. Example 3.3 from [D2] shows that the coefficients h_j may well be strictly c-algebraic, i.e. having no holomorphic extension onto a neighbourhood of A in \mathbb{C}^m (even locally).

Theorem 3.2 can be further generalized in two directions. The first one is the c-algebraic counterpart of Lemma 1.1 from [PT] (compare also [D1] for the c-holomorphic version which was much more straightforward) and its important consequence.

Proposition 3.6. *Let $f = (f_1, \dots, f_k) \in \mathcal{O}_c^a(A, \mathbb{C}^k)$ be proper on the pure k -dimensional set $A \subset \mathbb{C}^m$, with geometric degree $d(f)$. Fix $\ell \in \{1, \dots, k\}$ and assume that $g \in \mathcal{O}_c^a(A)$ satisfies $g^{-1}(0) \supset \bigcap_{j=1}^{\ell} f_j^{-1}(0)$. Then we can find ℓ functions $h_j \in \mathcal{O}_c^a(A)$ for which*

$$g^{d(f)} = \sum_{j=1}^{\ell} h_j f_j, \quad \text{on } A.$$

Proof. By the Chevalley-Remmert Theorem, the set

$$X := \{(f(x), g(x)) \in \mathbb{C}^k \times \mathbb{C} \mid x \in A\}$$

is algebraic as the proper ⁽⁴⁾ projection of the graph of (f, g) and it has pure dimension k . It is obvious that the characteristic polynomial P_g (having the form $(\#)$) of g relative to f yields $X = P_g^{-1}(0)$. Moreover,

$$X \cap (\{0\}^{\ell} \times \mathbb{C}^{k-\ell} \times \mathbb{C}) = \{0\}^{\ell} \times \mathbb{C}^{k-\ell} \times \{0\} \Rightarrow a_j|_{\{0\}^{\ell} \times \mathbb{C}^{k-\ell}} \equiv 0,$$

for all indices j (which is easy to check starting from the last one, i.e. a_d). This allows us to write $a_j(w) = y_1 a_{j,1}(w) + \dots + y_{\ell} a_{j,\ell}(w)$ where $w = (y, z) \in \mathbb{C}^{\ell} \times \mathbb{C}^{k-\ell}$ and $a_{j,s}$ are polynomials. Eventually, the identity $P_g(f(x), g(x)) \equiv 0$ gives the result. \square

We are able now to generalize this to the case of a set-theoretical complete intersection for strictly regular sequences in connection with [FPT], [D2] and [PT]. Suppose that $f: A \rightarrow \mathbb{C}^n$ is c-algebraic, A has pure dimension $k > 0$ and $f^{-1}(0)$ is pure $(k - n)$ -dimensional. This means exactly that the intersection $\Gamma_f \cap (\mathbb{C}^m \times \{0\}^k)$ is proper (i.e. it is a set-theoretical complete intersection). In such a case we may consider the *algebraic effective cycle of zeroes* of f :

$$(\star) \quad Z_f := \Gamma_f \cdot (\mathbb{C}^k \times \{0\}^k) = \sum_{j=1}^r i(\Gamma_f \cdot (\mathbb{C}^k \times \{0\}^k); V_j) V_j,$$

where $f^{-1}(0) = \bigcup_{j=1}^r V_j$ is the decomposition into irreducible components and $i(\Gamma_f \cdot (\mathbb{C}^k \times \{0\}^k); V_j)$ is the intersection multiplicity along V_j computed following Draper [Dr] (see also [Ch]).

Since all V_j are algebraic we may define the *degree of the cycle* Z_f to be the number

$$\deg Z_f = \sum_{j=1}^r i(\Gamma_f \cdot (\mathbb{C}^k \times \{0\}^k); V_j) \cdot \deg V_j.$$

⁴This is simple topological fact: given three locally compact topological spaces X, Y, Z and two continuous functions $\varphi: X \rightarrow Y$ and $\psi: X \rightarrow Z$ such that ψ is in addition proper, the function $(\varphi, \psi): X \rightarrow Y \times Z$ is proper.

Note that for $k = n$ we clearly obtain $\deg Z_f = d(f)$.

Following [FPT] we introduce the notion of a strictly regular sequence:

Definition 3.7. A map $f \in \mathcal{O}_c^a(A, \mathbb{C}^n)$ where $A \subset \mathbb{C}^m$ has pure dimension $k > n$, is called *strictly regular*, if there are $k - n$ affine forms $L_j: \mathbb{C}^m \rightarrow \mathbb{C}$ such that $(f, L_1|_A, \dots, L_{k-n}|_A): A \rightarrow \mathbb{C}^k$ is proper.

Remark 3.8. If $f \in \mathcal{O}_c^a(A, \mathbb{C}^n)$ is strictly regular, then $f^{-1}(0)$ has pure dimension $k - n$ (cf. [D2]) and so Z_f is well-defined.

Theorem 3.9. Assume $f \in \mathcal{O}_c^a(A, \mathbb{C}^n)$ is strictly regular. Then for any $g \in \mathcal{O}_c^a(A)$ such that $g^{-1}(0) \supset f^{-1}(0)$ there are n functions $h_j \in \mathcal{O}_c^a(A)$ yielding

$$g^{\deg Z_f} = \sum_{j=1}^n h_j f_j \quad \text{on the whole of } A.$$

Proof. Let $L = (L_1, \dots, L_{k-n})$ be the affine mapping from the definition of strict regularity. Then $\varphi := (f, L|_A) \in \mathcal{O}_c^a(A, \mathbb{C}^k)$ is a proper mapping. We may assume that coordinates in \mathbb{C}^m are chosen in such a way that L is linear. Observe that $\varphi^{-1}(u, v) = f^{-1}(u) \cap L^{-1}(v)$, for $(u, v) \in \mathbb{C}^n \times \mathbb{C}^{k-n}$. Consider $\psi(\zeta, u) := (u, L(\zeta)) \in \mathbb{C}^n \times \mathbb{C}^{k-n}$, for $(\zeta, u) \in \mathbb{C}^m \times \mathbb{C}^n$. It is easy to see that $\psi|_{\Gamma_f}$ is proper: if $K \subset \mathbb{C}^k$ is a compact set, then $\psi^{-1}(K) \cap \Gamma_f = \{(\zeta, f(\zeta)) \mid \zeta \in \varphi^{-1}(K)\}$ is clearly compact.

Then $d(\varphi) = d(\psi|_{\Gamma_f})$, because $\psi^{-1}(u, v) = f^{-1}(u) \cap L^{-1}(v)$. The intersection $\Gamma_f \cap \text{Ker } \psi$ is isolated and proper and, obviously, $d(\psi|_{\Gamma_f}) = \deg(\Gamma_f \cdot \text{Ker } \psi)$. Since $\text{Ker } \psi \subset \mathbb{C}^m \times \{0\}^n$, we can apply [TW2] Theorem 2.2:

$$\begin{aligned} \Gamma_f \cdot_{\mathbb{C}^{m+n}} \text{Ker } \psi &= (\Gamma_f \cdot_{\mathbb{C}^{m+n}} (\mathbb{C}^m \times \{0\}^n)) \cdot_{\mathbb{C}^m} \text{Ker } \psi = \\ &= Z_f \cdot \text{Ker } L. \end{aligned}$$

Therefore, in view of (\star) ,

$$d(\varphi) = \deg(Z_f \cdot \text{Ker } L) = \sum_{j=1}^r i(\Gamma_f \cdot (\mathbb{C}^k \times \{0\}^k); V_j) \deg(V_j \cdot \text{Ker } L),$$

where $\deg(V_j \cdot \text{Ker } L) = \sum_{a \in V_j \cap \text{Ker } L} m_a(\pi^L|_{V_j})$ with the local geometric multiplicities $m_a(\pi^L|_{V_j})$ of the projection π^L along $\text{Ker } L$. Note that we have necessarily $\deg(V_j \cdot \text{Ker } L) \leq \deg V_j$ whence $d(\varphi) \leq \deg Z_f$. Now, Proposition 3.6 gives

$$g^{d(\varphi)} = \sum_{j=1}^n g_j f_j, \quad \text{on } A,$$

with some $g_j \in \mathcal{O}_c^a(A)$. Multiplying both sides by $g^{\deg Z_f - d(\varphi)}$ and putting $h_j := g^{\deg Z_f - d(\varphi)} \cdot g_j$ we get the result sought for. \square

REFERENCES

- [ATW] R. Achilles, P. Tworzewski, T. Winiarski, *On improper isolated intersection in complex analytic geometry*, Ann. Polon. Math. LI (1990), 21-36;
- [BDTW] A. Białożyty, M. P. Denkowski, P. Tworzewski, T. Wawak, *On the growth exponent of c -holomorphic functions with algebraic graphs*, arXiv:1406.5293, submitted 2020;
- [Cg] E. Cygan, *Nullstellensatz and cycles of zeroes of holomorphic mappings*, Ann. Polon. Math. 78 no.2 (2002), 181-191;
- [Ch] E. M. Chirka, *Complex Analytic Sets*, Kluwer Acad. Publ. 1989;
- [D1] M. P. Denkowski, *The Lojasiewicz exponent of c -holomorphic mappings*, Ann. Polon. Math. LXXXVII.1 (2005), pp. 63-81;
- [D2] M. P. Denkowski, *A note on the Nullstellensatz for c -holomorphic functions*, Ann. Polon. Math. XC.3 (2007), 219-228;
- [D3] M. P. Denkowski, *Multiplicity and the pull-back problem*, Manuscripta Math. 149 (2016), 83-91;
- [Dr] R. N. Draper, *Intersection theory in analytic geometry*, Math. Ann. 180 (1969), 175-204;
- [FPT] A. Fabiano, A. Płoski, P. Tworzewski, *Effective Nullstellensatz for strictly regular sequences*, Univ. Jagell. Acta Math. 38 (2000), 163-167;
- [FHMM] G. Fichou, J. Huisman, F. Mangolte, J.-P. Monnier, *Fonctions régulières*, arXiv 1112.3800 [math.AG] (2011);
- [Kol] J. Kollár, *Continuous rational functions on real and p -adic varieties*, arXiv 1101.3737 [math.AG] (2011);
- [K] W. Kucharz, *Rational maps in real algebraic geometry*, Adv. Geom. 9 (4) (2009), 517-539;
- [L] S. Lojasiewicz, *Introduction to Complex Analytic Geometry*, Birkhäuser, Basel 1991;
- [P1] A. Płoski, *On the growth of proper polynomial mappings*, Ann. Polon. Math. XLV.3 (1985), 297-309;
- [P2] A. Płoski, *Multiplicity and the Lojasiewicz exponent*, Singularities, Banach Center Publ. vol. 20 PWN Warsaw (1988), 353-364;
- [PT] A. Płoski, P. Tworzewski, *Effective Nullstellensatz on analytic and algebraic varieties*, Bull. Pol. Acad. Sci. Math. vol. 46 nr 1 (1998), 31-38;
- [R] R. Remmert, *Projektionen analytischer Mengen*, Math. Ann. 130 (1956), 410-441;
- [Sp] S. Spodzieja, *The Lojasiewicz exponent at infinity for overdetermined polynomial mappings*, Ann. Pol. Math. LXXVIII.1 (2002), 1-10;
- [TW1] P. Tworzewski, T. Winiarski, *Analytic sets with proper projections*, Journ. Reine Angew. Math. 337 (1982), 68-76;
- [TW2] P. Tworzewski, T. Winiarski, *Cycles of zeroes of holomorphic mappings*, Bull. Pol. Acad. Sci. Math. vol. 37 (1986), 95-101;
- [Wh] H. Whitney, *Complex Analytic Varieties*, Addison-Wesley Publ. Co. 1972.

JAGIELLONIAN UNIVERSITY, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE,
INSTITUTE OF MATHEMATICS, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

E-mail address: adam.bialozyt@doctoral.uj.edu.pl

E-mail address: maciej.denkowski@uj.edu.pl

E-mail address: piotr.tworzewski@uj.edu.pl