

FIBRATION THEOREMS À LA MILNOR FOR DIFFERENTIABLE MAPS WITH NON-ISOLATED SINGULARITIES

JOSÉ LUIS CISNEROS-MOLINA, AURÉLIO MENEGON, JOSÉ SEADE,
AND JAWAD SNOUSSI

ABSTRACT. We prove fibration theorems à la Milnor for C^ℓ real maps with non isolated critical values. We study the situation for maps with linear discriminant, and prove that the concept of d -regularity is the key point for the existence of a Milnor fibration on the sphere. We also explain how one can modify the target space by homeomorphisms to linearize a general discriminant. Whenever the composed map is d -regular one has fibration on the sphere. Plenty of examples are discussed along the text, particularly the interesting family of functions $(f, g): \mathbb{R}^n \rightarrow \mathbb{R}^2$ of the type

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^p, \sum_{i=1}^n b_i x_i^q \right),$$

where $a_i, b_i \in \mathbb{R}$ are constants in generic position and $p, q \geq 2$ are integers.

1. INTRODUCTION

We study the topological behaviour of the non-critical levels of differentiable functions near a critical point.

As motivation, let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a real analytic map, $n \geq k \geq 2$, with $0 \in \mathbb{R}^n$ an isolated critical point and $\dim(f^{-1}(0)) > 0$. By the implicit function theorem, for every sufficiently small sphere $\mathbb{S}_\epsilon^{n-1}$ around 0, one has the following *transversality property*: there exists

Date: February 18, 2020.

2010 Mathematics Subject Classification. Primary 14D06, 14B05, 14J17, 32S55, 58K05, 58K15.

Key words and phrases. Milnor fibration, d -regularity, linear discriminant, sub-analytic.

The first author is Regular Associate of the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. Supported by CONACYT 253506.

The second author had partial support from CNPq (Brazil).

The third author was supported by CONACYT 282937 and UNAM-DGAPA-PAPIIT IN110517.

The last author was supported by UNAM-DGAPA-PAPIIT 113817.

$\delta > 0$ such that for all $t \in \mathbb{R}^k$ with $\|t\| \leq \delta$ the fiber $f^{-1}(t)$ meets $\mathbb{S}_\epsilon^{n-1}$ transversally. Hence, by the Relative Ehresmann Fibration Theorem (see [16, p. 23]), one has a locally trivial fibration restricting f to the “tube”:

$$(1) \quad f: (\mathbb{B}_\epsilon^n \cap (f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \{0\}))) \longrightarrow \mathbb{S}_\delta^{k-1} \setminus \{0\},$$

where \mathbb{B}_ϵ^n is the closed ball in \mathbb{R}^n of radius ϵ around the origin. In his now classical book [21] Milnor proves (see [21, Theorem 11.2] or [20, Theorem 2]) that (1) gives rise to a locally trivial fibration

$$(2) \quad \phi: \mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(0) \rightarrow \mathbb{S}^{k-1},$$

where \mathbb{S}^{k-1} is the sphere of radius 1 around $0 \in \mathbb{R}^k$. Milnor proves this constructing an integrable non-zero vector field on $\mathbb{B}_\epsilon \setminus f^{-1}(0)$ which is transverse to the fibres of f and to the spheres centered at 0 contained in \mathbb{B}_ϵ . The integrable curves of this vector field carry diffeomorphically the tube to the complement of $f^{-1}(\mathbb{B}_\delta^k) \cap \mathbb{S}_\epsilon^{n-1}$ in $\mathbb{S}_\epsilon^{n-1}$ keeping its boundary fixed.

Milnor points out in his book that this theorem has two main weaknesses: The condition for an analytic map $(\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$, to have an isolated critical point is very stringent, and even when this is satisfied, the projection map ϕ may not always be taken as the natural map $f/\|f\|$ (see [21, p. 99]).

The above discussion was extended in [25, Theorem 1.3] to functions f with an isolated critical value, a more general but still stringent setting. The authors proved¹ that if f has Thom’s a_f property, which implies the *transversality property* mentioned above, one also has fibration (1) on the tube, known as Milnor-Lê fibration, and using Milnor’s vector field one also has an equivalent fibration (2) on the sphere.

The question of whether we can take the projection ϕ of (2) to be the natural one, $\phi = f/\|f\|$ was answered in [10] for analytic functions with an isolated critical value, by introducing the concept of d -regularity (see also [12]): The map f is d -regular if and only if fibration (1) is equivalent to a fibration (2) where the projection is given by $\phi = f/\|f\|$.

The d -regularity condition actually springs from [9] and is defined by means of a canonical pencil as follows: For every line $0 \in \ell \subset \mathbb{R}^k$ consider the set

$$X_\ell = \{x \in \mathbb{R}^n \mid f(x) \in \ell\}.$$

This is a pencil of real analytic varieties intersecting at $f^{-1}(0)$ and smooth away from it. The map f is said to be *d -regular at 0* if there

¹It was proved in the more general context where the domain of f is a real analytic variety with an isolated singularity at 0.

exists $\epsilon_0 > 0$ such that every $X_\ell \setminus V$ is transverse to every sphere centred at 0 and contained in \mathbb{B}_{ϵ_0} , whenever the intersection is not empty.

Several works have been published in the last two decades studying Milnor fibrations for real analytic maps with either an isolated critical point or value, see for instance [28, 24, 27, 4, 25, 5, 10, 22, 1, 3, 13, 2] or the survey articles [6, 11, 12, 26].

In this paper we envisage the general case of functions (analytic or not) with arbitrary critical set (cf. [7, 19]). Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$, $n > k \geq 2$ with a critical point at 0, of class C^ℓ , $\ell \geq 1$, and $\dim(f^{-1}(0)) > 0$. Our first result is (Theorem 2.7) that f has the transversality property if and only if it admits local Milnor-Lê fibrations in tubes over the complement of the discriminant Δ_f .

Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be an analytic map with critical point at $0 \in \mathbb{R}^n$ and consider small balls $\mathbb{B}_\epsilon^n \subset \mathbb{R}^n$ and $\mathbb{B}_\delta^k \subset \mathbb{R}^k$ with $0 < \delta \ll \epsilon$. Let Δ_ϵ be the *extended discriminant* of f with respect to \mathbb{B}_ϵ^n , which consists of the images by f of critical points of f in the interior of \mathbb{B}_ϵ^n together with the images of the critical points of the restriction of f to the boundary sphere $\mathbb{S}_\epsilon^{n-1}$ (see Section 2). Following [23, §IV.4.4], in [7, Corollary 2.2] was proved that the restriction of f to the tube

$$(3) \quad f: \mathbb{B}_\epsilon \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon) \rightarrow \mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon,$$

is a locally trivial fibration, where $\mathbb{S}_\delta^{k-1} = \partial\mathbb{B}_\delta^k$. Hence, in this general setting there is always a fibration on the tube. Moreover, if f satisfies the *transversality property*, there is no contribution to the extended discriminant Δ_ϵ by points on the sphere $\mathbb{S}_\epsilon^{n-1}$, the extended discriminant is just the discriminant of f and the fibration on the tube essentially does not depend on ϵ . We remark that in [19] there is a simple criterium to decide when a map has the transversality condition. However, Milnor's vector field on $\mathbb{B}_\epsilon^n \setminus f^{-1}(\Delta_\epsilon)$ not necessarily works to inflate the tube to the sphere, since in general, $\mathbb{B}_\epsilon^n \setminus f^{-1}(\Delta_\epsilon)$ is not invariant under its flow.

In this article we extend the discussion of existence of fibrations (in the tube and in the sphere) for differentiable functions of class C^ℓ , $\ell \geq 1$, with possibly non-isolated critical value. The fibration on the tube always exists, since [7, Corollary 2.2] extends without problem to this setting. Concerning the fibration on the sphere, we answer the question below:

Question 1.1. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be an analytic map with critical point at $0 \in \mathbb{R}^n$ and arbitrary critical set. Is there a condition on f which ensures that there exists a fibration on the sphere?

The answer is d -regularity. We show that if for a map $(\mathbb{R}^n, 0) \xrightarrow{f} (\mathbb{R}^k, 0)$, $n \geq k \geq 2$, there exists a local homeomorphism $(\mathbb{R}^k, 0) \xrightarrow{h} (\mathbb{R}^k, 0)$ such that the preimage of the discriminant $h^{-1}(\Delta_f)$ is linear and the composition $f_h = h^{-1} \circ f$ is d -regular, then the map $f_h/\|f_h\|$ is the projection of a fibration on the sphere minus the preimage of Δ_f (see Theorem 5.2 for the precise statement). In this case, we say that f is d_h -regular. Every d -regular map is d_h -regular for h the identity map. In the analytic case, this fibration is equivalent to fibration (3) on the tube.

Notice that the discriminant Δ_f can have real codimension 1 and therefore its complement splits into finitely many connected components, say S_1, \dots, S_r . The topology of the fibers $f^{-1}(t)$ can change for values in different S_i . This happens for instance for the map germs

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^2, \sum_{i=1}^n b_i x_i^2 \right),$$

where $a_i, b_i \in \mathbb{R}$ are constants in generic position, studied by López de Medrano in [17]. Notice that each sector determines an analytic family of smooth manifolds that degenerate to the singular fiber over 0. If we take an arc joining two adjacent sectors we get a family of smooth compact manifolds that transform into other smooth compact manifolds as we cross the discriminant. It would be very interesting to determine how the topology changes as we move from one sector to another. This can clearly be seen in the example given in Subsection 5.2 of a non-analytic d_h -regular map.

Remark 1.2. Throughout this paper, we will assume that f is *locally surjective*, that is, the image of f contains an open neighbourhood of the origin in \mathbb{R}^p , and we shall not mention it all the time. Nevertheless, it is easy to see that in the general case the same results hold if one intersects the bases of the locally trivial fibrations with their image. This choice is to avoid a heavy notation.

Remark 1.3. The results of this article extend to analytic map-germs $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ if we consider f in the class of *nice analytic map-germs* defined in [2, Definition 2.2] for which the discriminant is a well-defined set-germ at 0, so it does not depend on the radius ϵ .

The article is organized as follows. In section 2, we give the fibration on the tube (Proposition 2.1), we define the transversality property and present Theorem 2.7 mentioned above. In section 3, we define when f has linear discriminant and we exhibit examples. Then we extend

the concept of d -regularity and the corresponding fibration theorems to this case (Theorem 3.9 and Theorem 3.12). In section 4, we define conic homeomorphisms on the target of f and d_h -regularity, and we give examples of both concepts. Finally in section 5, for an arbitrary f , we discuss how using a conic homeomorphism one can linearize the discriminant of f . Then one can see if the linearized map is d -regular, in this case we get the main fibration theorem (Theorem 5.2) and we apply it to the case of maps with isolated critical value and the transversality property (Theorem 5.4). We give several examples, including the family mentioned in the abstract and an example of a non-analytic map. We include Appendix A where we prove Lemma A.1, which says that the composition of two C^ℓ -locally trivial fibrations between smooth manifolds is a C^ℓ -locally trivial fibration. It is used in the proof of Theorem 3.9.

2. FIBRATION ON THE TUBE

Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$, $n > k \geq 2$, be a map of class C^ℓ with $\ell \geq 1$ and a critical point at 0. Following [7, §2] we see that there always exists a fibration on a tube.

Assume that f is locally surjective (see Remark 1.2). Equip \mathbb{R}^n with a Whitney stratification adapted to $V = f^{-1}(0)$, and let $\mathbb{B}_{\epsilon_0}^n$ be a closed ball in \mathbb{R}^n , centred at 0, of sufficiently small radius $\epsilon_0 > 0$, so that every sphere in this ball, centred at 0, meets transversely every stratum of V . We call ϵ_0 a *Milnor radius* for f (at 0). In what follows, for $0 < \epsilon < \epsilon_0$ we shall consider the restriction f_ϵ of f to the closed ball $\mathbb{B}_\epsilon^n \subset \mathbb{B}_{\epsilon_0}^n$.

Denote by Σ_ϵ the intersection of the critical set of f with the ball \mathbb{B}_ϵ and set $\Delta_\epsilon := f_\epsilon(\Sigma_\epsilon)$, which we call the *discriminant* of f_ϵ . The discriminant Δ_ϵ is a subanalytic set and it may depend on the choice of the radius ϵ , as showed in [2].

Also denote by $\Sigma_\epsilon(\mathbb{S}_\epsilon^{n-1})$ the set of critical points in $\mathbb{S}_\epsilon^{n-1}$ of the restriction $f_\epsilon|_{\mathbb{S}_\epsilon^{n-1}}$. Set $\hat{\Sigma}_\epsilon := \Sigma_\epsilon \cup \Sigma_\epsilon(\mathbb{S}_\epsilon^{n-1})$ and denote by $\hat{\Delta}_\epsilon := f(\hat{\Sigma}_\epsilon)$ which we call the *extended discriminant*² of f .

Let \mathbb{B}_δ^k be an open ball in \mathbb{R}^k centred at 0 of radius $0 < \delta \ll \epsilon$.

Proposition 2.1 ([7, Corollary 2.2]). *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$ be a map of class C^ℓ with $\ell \geq 1$. Then the restrictions:*

$$(4) \quad f_\epsilon|: \mathbb{B}_\epsilon^n \cap f_\epsilon^{-1}(\mathring{\mathbb{B}}_\delta^k \setminus \hat{\Delta}_\epsilon) \rightarrow \mathring{\mathbb{B}}_\delta^k \setminus \hat{\Delta}_\epsilon$$

²In [23, §IV.4.4] $\hat{\Sigma}_\epsilon$ is called the *apparent contour at the source* and Δ_ϵ is called the *apparent contour at the target* or just *apparent contour*.

and

$$(5) \quad f_\epsilon|: \mathbb{B}_\epsilon^n \cap f_\epsilon^{-1}(\mathbb{S}_\delta^{k-1} \setminus \hat{\Delta}_\epsilon) \rightarrow \mathbb{S}_\delta^{k-1} \setminus \hat{\Delta}_\epsilon$$

are locally trivial fibrations.

If f is analytic, then the fibrations above are smooth.

We can also consider a further condition of f which ensures that the fibrations of Proposition 2.1 essentially does not depend on ϵ .

Definition 2.2. Let $0 < \epsilon < \epsilon_0$, we say that f has the *transversality property in the ball* \mathbb{B}_ϵ^n if there exist $0 < \delta \ll \epsilon$ such that for every $y \in \mathbb{B}_\delta^k \setminus \Delta_\epsilon$ the fibre $f^{-1}(y)$ is transverse to the sphere $\mathbb{S}_\epsilon^{n-1}$. Since we are assuming f locally surjective we can take $\mathbb{B}_\delta^k \subset \text{Im}(f_\epsilon)$.

Remark 2.3. In the case that f is not locally surjective for the transversality property we need to ask that for every $y \in (\mathbb{B}_\delta^k \setminus \Delta_\epsilon) \cap \text{Im}(f_\epsilon)$ the fibre $f^{-1}(y)$ meets $\mathbb{S}_\epsilon^{n-1}$ transversely. But if we consider that an empty fibre intersects transversely the sphere $\mathbb{S}_\epsilon^{n-1}$ we can state in general the transversality property as above.

Remark 2.4. If we work with a nice analytic map-germ $f: (U, 0) \rightarrow (\mathbb{R}^k, 0)$ (see Remark 1.3), the discriminant does not depend on the radius ϵ and we just denote it by Δ . In this case we need the following stronger *transversality property*. We say that f satisfies the *strong transversality property* if for every $0 < \epsilon < \epsilon_0$ there exists a real number $\delta = \delta(\epsilon)$ such that $0 < \delta \ll \epsilon$ and for every $y \in \mathbb{B}_\delta^k \setminus \Delta$ the fibre $f^{-1}(y)$ meets $\mathbb{S}_\epsilon^{n-1}$ transversely. It is clear that if f satisfies the strong transversality property, then for every $0 < \epsilon < \epsilon_0$, f has the transversality property in the ball \mathbb{B}_ϵ^n .

Remark 2.5. If f is analytic and satisfies the Thom a_f -property then it has the strong transversality property (compare with [10, Remark 5.7]).

Theorem 2.6. Consider $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$, $n > k \geq 2$ a nice map-germ with a critical point at 0 and $\dim(f^{-1}(0)) > 0$. Then f has the strong transversality property if and only if there exists $\epsilon_0 > 0$ such that for every $0 < \epsilon < \epsilon_0$ and every $0 < \delta \ll \epsilon$ there is no contribution to the extended discriminant $\hat{\Delta}_\epsilon$ coming from the intersection $\mathbb{S}_\epsilon^n \cap f^{-1}(\mathbb{B}_\delta^k)$, that is, $\Sigma_\epsilon = \hat{\Sigma}_\epsilon$, and $\hat{\Delta}_\epsilon \cap \mathbb{B}_\delta^k = \Delta_\epsilon \cap \mathbb{B}_\delta^k = f(\Sigma_f(\mathring{\mathbb{B}}_\epsilon)) \cap \mathbb{B}_\delta^k$ is the discriminant of f_ϵ restricted to \mathbb{B}_δ^k . Therefore, fibrations (4) and (5) essentially do not depend on ϵ .

So we have the following

Theorem 2.7. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$, $n > k \geq 2$ be a map of class C^ℓ , $\ell \geq 1$ with a critical point at 0 and $\dim(f^{-1}(0)) > 0$. Then

the ball \mathbb{B}_ϵ^n has the transversality property if and only if it admits local Milnor-Lê fibrations in tubes over the complement of the discriminant Δ_ϵ .

3. THE d -REGULARITY FOR REAL ANALYTIC MAPS WITH LINEAR DISCRIMINANT

In this section, we extend the concept of d -regularity to real differentiable maps with linear discriminant. The main idea is that d -regularity can be defined as transversality of preimages of lines, different from those lying in the discriminant, with sufficiently small spheres. Then we show that in this context, d -regularity guarantees a fibration on the sphere.

First, let us make the definition of linear discriminant precise:

Definition 3.1. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ , $\ell \geq 1$ with a critical point at 0. Let ϵ_0 be a Milnor radius for f . Let $0 < \epsilon < \epsilon_0$, we say that f has *linear discriminant in the ball \mathbb{B}_ϵ^n* if Δ_ϵ is a union of line-segments with one endpoint at $0 \in \mathbb{R}^k$. We say that $\eta > 0$ is a *linearity radius* for Δ_ϵ if each of these line-segments intersect \mathbb{S}_η^{k-1} , that is, if

$$\Delta_\epsilon \cap \mathbb{B}_\eta^k = \text{Cone}(\Delta_\epsilon \cap \mathbb{S}_\eta^{k-1}).$$

Remark 3.2. The case when f has $0 \in \mathbb{R}^k$ as isolated critical value is considered to have linear discriminant with $\Delta_\epsilon \cap \mathbb{S}_\eta^{k-1} = \emptyset$.

Remark 3.3. Let $f: (\mathbb{B}_{\epsilon_0}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$ be a locally surjective analytic map. Let $0 < \epsilon < \epsilon_0$ and suppose f has the transversality property in the ball \mathbb{B}_ϵ^n and that f_ϵ has linear discriminant in \mathbb{B}_ϵ^n . Notice that in this case we can take the linearity radius to be the δ in the definition of the transversality property.

Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with $\ell \geq 1$ with linear discriminant in the ball \mathbb{B}_ϵ^n and consider a linearity radius $\eta > 0$ for f . Set:

$$\mathcal{A}_\eta := \Delta_\epsilon \cap \mathbb{S}_\eta^{k-1}.$$

Let $\pi: \mathbb{S}_\eta^{k-1} \rightarrow \mathbb{S}^{k-1}$ be the projection onto the unit sphere \mathbb{S}^{k-1} and set $\mathcal{A} = \pi(\mathcal{A}_\eta)$. For each point $\theta \in \mathbb{S}_\eta^{k-1}$, let $\mathcal{L}_\theta \subset \mathbb{R}^k$ be the open ray in \mathbb{R}^k from the origin that contains the point θ . Set:

$$E_\theta := f^{-1}(\mathcal{L}_\theta).$$

Notice that E_θ is a manifold of class C^ℓ for any θ in $\mathbb{S}_\eta^{k-1} \setminus \mathcal{A}_\eta$. We have:

Definition 3.4. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with $\ell \geq 1$ and linear discriminant. We say that f is *d-regular (in the ball \mathbb{B}_ϵ^n)* if E_θ intersects the sphere $\mathbb{S}_{\epsilon'}^{n-1}$ transversely in \mathbb{R}^n , for every ϵ' with $0 < \epsilon' \leq \epsilon$ and for every $\theta \in \mathbb{S}_\eta^{k-1} \setminus \mathcal{A}_\eta$.

Example 3.5. Consider the real analytic map $f: (\mathbb{R}^4, 0) \rightarrow (\mathbb{R}^3, 0)$ given by:

$$f(x, y, z, w) := (x^2 - y^2z, y, w).$$

Its critical set is the plane $\{x = y = 0\}$ in \mathbb{R}^4 , and its discriminant is the axis $\{u_1 = u_2 = 0\}$ in \mathbb{R}^3 , which is linear. One can check that f is not *d-regular*.

A class of examples.

Example 3.6. In [17] S. López de Medrano studied the topology of real analytic maps $(f, g): \mathbb{R}^n \rightarrow \mathbb{R}^2$ such that both f and g are homogeneous quadratic polynomials. For (f, g) satisfying some generic hypothesis, he completely described the topology of $V(f, g) = f^{-1}(0) \cap g^{-1}(0)$ in terms of the coefficients of f and g .

Let us restrict to the diagonal case, that is, when (f, g) has the form:

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^2, \sum_{i=1}^n b_i x_i^2 \right),$$

where a_i and b_i are real constants in generic position. This means that the origin is in the convex hull of the points (a_i, b_i) (which guarantees that the link of $V(f)$ is non-empty) and that no two of the points (a_i, b_i) are linearly dependent, that is, $a_i b_j \neq a_j b_i$, for any $i \neq j$ (*Weak Hyperbolicity Hypothesis*).

A simple calculation shows that the set Σ of critical points of (f, g) coincides with the coordinate axis of \mathbb{R}^n and the discriminant $\Delta(f, g)$ is the union of the n line-segments in \mathbb{R}^2 joining the origin and the points $\lambda_i := (a_i, b_i)$. Hence (f, g) has linear discriminant.

We are going to show that (f, g) is *d-regular*. First, notice that any ray \mathcal{L}_θ is given by one of the following forms:

$\{u + \alpha v = 0\} \cap \{v > 0\}$, $\{u + \alpha v = 0\} \cap \{v < 0\}$, $\{v = 0\} \cap \{u > 0\}$ or $\{v = 0\} \cap \{u < 0\}$ for some $\alpha \in \mathbb{R}$. Hence any E_θ has one of the following forms:

$\{f + \alpha g = 0\} \cap \{g > 0\}$, $\{f + \alpha g = 0\} \cap \{g < 0\}$, $\{g = 0\} \cap \{f > 0\}$ or $\{g = 0\} \cap \{f < 0\}$ for some $\alpha \in \mathbb{R}$.

In any case, all the equations and inequations are homogeneous, so it is easy to verify that E_θ intersects any sphere centered at the origin transversally.

The following example generalizes López de Medrano's maps:

Example 3.7. Let \mathbb{K} be either \mathbb{R} or \mathbb{C} . Let $(f, g) : \mathbb{K}^n \rightarrow \mathbb{K}^2$ be a \mathbb{K} -analytic map of the form:

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^p, \sum_{i=1}^n b_i x_i^p \right),$$

where $a_i, b_i \in \mathbb{K}$ are constants in generic position (as in Example 3.6) and $p \geq 2$ is an integer.

As before, one has that the critical set Σ of (f, g) is given by the coordinate axis of \mathbb{K}^n and the discriminant Δ is linear. Moreover, the same argument of the example above shows that (f, g) is d -regular.

Let us show that the link Z of $V = (f, g)^{-1}(0)$ is non-empty, as before. In order to deal simultaneously with both the real and the complex case, set $c = 1$ if $\mathbb{K} = \mathbb{R}$ and $c = 2$ if $\mathbb{K} = \mathbb{C}$.

If p is odd, consider the homeomorphism:

$$\begin{aligned} \phi_1 : \quad \mathbb{K}^n &\longrightarrow \mathbb{K}^n \\ (x_1, \dots, x_n) &\longmapsto (x_1^p, \dots, x_n^p) \end{aligned} .$$

Then (f, g) equals the composition $(f^1, g^1) \circ \phi_1$, where (f^1, g^1) is the submersion given by:

$$(f^1, g^1)(x) := \left(\sum_{i=1}^n a_i x_i, \sum_{i=1}^n b_i x_i \right),$$

Hence V is homeomorphic to a $(n-2)$ - \mathbb{K} -dimensional plane in \mathbb{K}^n and Z is homeomorphic to the sphere $\mathbb{S}^{c(n-2)-1}$.

If $p = 2r$ for some odd integer $r \geq 3$, consider the homeomorphism:

$$\begin{aligned} \phi_2 : \quad \mathbb{K}^n &\longrightarrow \mathbb{K}^n \\ (x_1, \dots, x_n) &\longmapsto (x_1^r, \dots, x_n^r) \end{aligned} .$$

So $(f, g) = (f^2, g^2) \circ \phi_2$, where $(f^2, g^2) := (\sum_{i=1}^n a_i x_i^2, \sum_{i=1}^n b_i x_i^2)$ is López de Medrano's map as in Example 3.6. Therefore Z is homeomorphic to the link of (f^2, g^2) , which is non-empty.

Finally, if $p = 2r$ for some even integer $r \geq 2$, consider the map:

$$\begin{aligned} \phi_3 : \quad \mathbb{K}^n &\longrightarrow \mathbb{K}^n \\ (x_1, \dots, x_n) &\longmapsto (x_1^r, \dots, x_n^r) \end{aligned} ,$$

which is not a homeomorphism. Once again we have that $(f, g) = (f^2, g^2) \circ \phi_3$. Hence $V = (\phi_3)^{-1}(\hat{V})$, where $\hat{V} := (f^2, g^2)^{-1}(0)$ has dimension greater than zero. So the dimension of V is bigger than zero.

Now we give a characterization of d -regularity for maps with linear discriminant, based on some definitions and preliminary results from [10, §3].

Let $0 < \epsilon < \epsilon_0$ be such that f has linear discriminant and it is d -regular in the ball \mathbb{B}_ϵ^n . Set $W := f^{-1}(\Delta_\epsilon)$ and consider the maps $\Phi: \mathbb{B}_\epsilon^n \setminus W \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ and $\mathfrak{F}: \mathbb{B}_\epsilon^n \setminus W \rightarrow \mathbb{R}^k \setminus \Delta_\epsilon$ given respectively by

$$\Phi(x) = \frac{f(x)}{\|f(x)\|}, \quad \text{and} \quad \mathfrak{F} = \|x\|\Phi(x).$$

Notice that given $y \in \mathcal{L}_\theta$ with $\theta \in \mathbb{S}_\eta^{k-1} \setminus \mathcal{A}_\eta$ the fibre $\mathfrak{F}^{-1}(y)$ is the intersection of E_θ with the sphere of radius $\|y\|$ centred at 0. Each E_θ is a union of fibres of \mathfrak{F} , just as it is a union of fibres of f , so we have

$$(6) \quad E_\theta = f^{-1}(\mathcal{L}_\theta) = \mathfrak{F}^{-1}(\mathcal{L}_\theta).$$

The map \mathfrak{F} is called the *spherefication* of f .

The following proposition is a straightforward generalization of [10, Proposition 3.2].

Proposition 3.8. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with $\ell \geq 1$, with linear discriminant. Let ϵ_0 be a Milnor radius for f and let $0 < \epsilon < \epsilon_0$. The following conditions are equivalent*

- The map f is d -regular in the ball \mathbb{B}_ϵ^n .
- For each sphere $\mathbb{S}_{\epsilon'}^{n-1}$ in \mathbb{R}^n centred at 0 of radius $\epsilon' < \epsilon$, the restriction map $\mathfrak{F}_{\epsilon'}: \mathbb{S}_{\epsilon'}^{n-1} \setminus W \rightarrow \mathbb{S}_{\epsilon'}^{k-1} \setminus \Delta_\epsilon$ of \mathfrak{F} is a submersion.
- The spherefication map \mathfrak{F} is a submersion.
- The map $\phi = \frac{f}{\|f\|}: \mathbb{S}_{\epsilon'}^{n-1} \setminus W \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ is a submersion for every sphere $\mathbb{S}_{\epsilon'}^{n-1}$ with $\epsilon' < \epsilon$.

We can now state the fibration theorem:

Theorem 3.9. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$ be a map of class C^ℓ with $\ell \geq 1$. Let ϵ_0 be a Milnor radius for f and let $0 < \epsilon < \epsilon_0$. Suppose f has linear discriminant and the transversality property in the ball \mathbb{B}_ϵ^n . If f is d -regular in the ball \mathbb{B}_ϵ^n , then the restriction of Φ given by*

$$(7) \quad \phi = \Phi|: \mathbb{S}_\epsilon^{n-1} \setminus W \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$$

is a (differentiable) locally trivial fibration over its image, where $W := f^{-1}(\Delta_\epsilon)$. If f is analytic then the fibration ϕ is smooth.

Proof. Suppose that f is surjective. The general case is analogous. Remember from Remark 3.3 that we can take $\delta > 0$ for the transversality property as the linearity radius of f . Set

$$\mathcal{M} := \mathbb{S}_\epsilon^{n-1} \setminus W.$$

Notice that \mathcal{M} is an open submanifold of $\mathbb{S}_\epsilon^{n-1}$ since Δ_ϵ is closed in \mathbb{R}^k . Consider the following decomposition

$$\mathcal{M} = (\mathcal{M} \cap f^{-1}(\mathbb{B}_\delta^k)) \cup (\mathcal{M} \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)),$$

where $\mathring{\mathbb{B}}_\delta^k$ is the interior of the closed ball \mathbb{B}_δ^k . Both pieces are submanifolds-with-boundary of \mathcal{M} of dimension $n - 1$ and their intersection is the common boundary submanifold of dimension $n - 2$

$$\mathbb{S}_\epsilon^{n-1} \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon) = (\mathcal{M} \cap f^{-1}(\mathbb{B}_\delta^k)) \cap (\mathcal{M} \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)).$$

We are going to show that the restriction of ϕ to each of these components is a (differentiable) fibre bundle, so they can be glued into a global fibre bundle.

The restriction of f given by $f_1: \mathcal{M} \cap f^{-1}(\mathbb{B}_\delta^k) \rightarrow \mathbb{B}_\delta^k \setminus \Delta_\epsilon$ is proper since $\mathbb{S}_\epsilon^{n-1} \cap f^{-1}(\mathbb{B}_\delta^k)$ is compact, and since f has the transversality property in the ball \mathbb{B}_ϵ^n it is a submersion, and by Ehresmann fibration theorem it is a (differentiable) fibre bundle. Now consider the radial projection $\tilde{\pi}: \mathbb{B}_\delta^k \setminus \Delta_{f,\eta} \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ which is a (trivial and smooth) fibre bundle. The restriction

$$\phi_1: (\mathcal{M} \cap f^{-1}(\mathbb{B}_\delta^k)) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$$

of ϕ is given by the composition $\tilde{\pi} \circ f_1$. By Lemma A.1 the composition $\phi_1 = \tilde{\pi} \circ f_1$ is a C^ℓ -locally trivial fibration.

So now we just have to show that the restriction:

$$\phi_2: \mathcal{M} \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$$

is a C^ℓ -fibration. We have that ϕ_2 is proper since $\mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)$ is compact.

Since f is d -regular, by Proposition 3.8 the map $\phi: \mathbb{S}_\epsilon^{n-1} \setminus W \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ has no critical points. So ϕ_2 is a submersion restricted to the interior $\mathcal{M} \setminus f^{-1}(\mathbb{B}_\delta^k)$ of $\mathcal{M} \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)$. Since ϕ_1 and ϕ_2 coincide on the boundary $\mathcal{M} \cap f^{-1}(\mathbb{S}_\delta^{k-1})$ and we already saw that ϕ_1 restricted to this boundary is a submersion. The result follows from the Ehresmann fibration theorem for manifolds with boundary. \square

Remark 3.10. In the case of f with isolated critical value and transversality property, we have $\Delta_\epsilon = \{0\}$ and $\mathcal{A} = \emptyset$ and Theorem 3.9 in this case gives another proof of the existence of the fibration on the sphere $\phi = \frac{f}{\|f\|}: \mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(0) \rightarrow \mathbb{S}^{k-1}$ without ‘‘inflating’’ the fibration (5) on the tube (see [10, Theorem 5.3 (2)]).

3.1. The analytic case: equivalence of fibrations. In this subsection we consider $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$, an analytic map with linear discriminant and the transversality property in the ball \mathbb{B}_ϵ^n .

Consider the locally trivial fibration (5) of Proposition 2.1. Composing it with the radial projection $\tilde{\pi}: \mathbb{B}_\delta^k \setminus \Delta_{f,\eta} \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ we get an equivalent locally trivial fibration

$$\tilde{f} := \tilde{\pi} \circ f|: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_{f,\eta}) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}.$$

We want to show that this locally trivial fibration is equivalent to the locally trivial fibration (7), provided that f is d -regular. For this, we use the following generalization of a characterization of d -regularity given in [8, Theorem 3.7] to the case of f with linear discriminant.

Theorem 3.11 ([10, Lemma 5.2]). *The map f is d -regular in the ball \mathbb{B}_ϵ^n , if and only if there exists an analytic vector field \tilde{w} on $\mathring{\mathbb{B}}_\epsilon^n \setminus V$ which has the following properties:*

- (1) *It is radial, i.e., it is transverse to all spheres in $\mathring{\mathbb{B}}_\epsilon^n$ centred at 0.*
- (2) *It is transverse to all the tubes $f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon)$.*
- (3) *It is tangent to each E_θ for $\theta \notin \mathcal{A}_\eta$, whenever it is not empty.*

Proof. Same as the proof of [8, Theorem 3.7] but replacing $\mathring{\mathbb{B}}_\epsilon^n \setminus V$ by $\mathring{\mathbb{B}}_\epsilon^n \setminus f^{-1}(\Delta_\epsilon)$, see [8, Remark 3.9]. \square

Now we can use the vector field \tilde{w} of Theorem 3.11 to give the equivalence of fibrations.

Theorem 3.12. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ with $n \geq k \geq 2$ be a map of class C^ℓ with $\ell \geq 1$. Let ϵ_0 be a Milnor radius for f and let $0 < \epsilon < \epsilon_0$. Suppose f has linear discriminant, has the transversality property and that it is d -regular in the ball \mathbb{B}_ϵ^n . Then the fibre bundles:*

$$\tilde{f} := \pi_\delta \circ f_\epsilon|: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \mathcal{A}_\delta) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$$

and

$$\phi: \mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(\Delta) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$$

are equivalent, where $\pi_\delta: \mathbb{S}_\delta^{k-1} \rightarrow \mathbb{S}^{k-1}$ is the radial projection.

Proof. In the proof of Theorem 3.9 we saw that the restriction of ϕ given by $\phi|: (\mathbb{S}_\epsilon^{n-1} \setminus W) \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ is a fibre bundle. The flow associated to the vector field \tilde{w} of Theorema 3.11 defines in the usual way a diffeomorphism τ between $\mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon)$ and $(\mathbb{S}_\epsilon^{n-1} \setminus W) \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)$: for a point $x \in \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon)$ follow the solution of \tilde{w} that passes through x till it meets $\mathbb{S}_\epsilon^{n-1}$ at some point \hat{x} . This point exists and is unique because \tilde{w} satisfies conditions (1) and (2) in

Theorem 3.11. Define $\tau(x) = \hat{x}$. By condition (3) in Theorem 3.11 the solutions of \tilde{w} lie in an E_θ , then we have that $\tilde{f}(x) := \Phi(x) = \Phi(\hat{x}) =: \phi(\hat{x})$. Therefore, the diffeomorphism $\tau: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon) \rightarrow (\mathbb{S}_\epsilon^{n-1} \setminus W) \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k)$ gives an equivalence of fibre bundles

$$\begin{array}{ccc} \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_f^\eta) & \xrightarrow{\tau} & (\mathbb{S}_\epsilon^{n-1} \setminus W) \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k) \\ & \searrow \tilde{f} & \swarrow \phi \\ & \mathbb{S}^{k-1} \setminus \mathcal{A} & \end{array}$$

We saw in the proof of Theorem 3.9 that the locally trivial fibration $\phi|: (\mathbb{S}_\epsilon^{n-1} \setminus W) \setminus f^{-1}(\mathring{\mathbb{B}}_\delta^k) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$ can be extended to the locally trivial fibration $\phi: \mathbb{S}_\epsilon^{n-1} \setminus W \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}$. \square

Now we want to apply Theorem 3.12 in Example 3.7. First, let us prove the following two propositions:

Proposition 3.13. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ . If $V(f) := f^{-1}(0)$ is not contained in the critical set Σ_f of f , then f is surjective.*

Proof. By hypothesis, for any $\epsilon > 0$ there exists $x \in V(f) \cap \mathbb{B}_\epsilon^n$ such that $x \notin \Sigma_f$. So, up to a change of coordinates, f is a projection on an open neighborhood W_x of x in \mathbb{B}_ϵ^n . Hence $f(W_x)$ is an open neighborhood of 0 in \mathbb{R}^k . \square

Proposition 3.14. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a real analytic map. If $V(f) = \{0\}$ or if $\Sigma_f(\mathring{\mathbb{B}}_\epsilon) \cap V(f) \subset \{0\}$, then f has the transversality property.*

Proof. Suppose that $V(f) = \{0\}$ and fix $\epsilon > 0$. By continuity of f there exists $\delta > 0$ sufficiently small such that $f^{-1}(t) \subset \mathbb{B}_\epsilon^n$, for any $t \in \mathbb{B}_\delta^k$. So f has the transversality property.

Now suppose that $\dim V(f) > 0$. Since f is real analytic, there exists $\epsilon_0 > 0$ sufficiently small such that $V(f)$ intersects the sphere $\mathbb{S}_\epsilon^{n-1}$ transversally in \mathbb{R}^n , for any ϵ with $0 < \epsilon \leq \epsilon_0$.

Fix ϵ and consider the restriction $f|$ of f to $\mathbb{S}_\epsilon^{n-1}$. Its critical points are the critical points of f that are in $\mathbb{S}_\epsilon^{n-1}$ and the regular points x of f that are in $\mathbb{S}_\epsilon^{n-1}$ such that $f^{-1}(f(x))$ intersect $\mathbb{S}_\epsilon^{n-1}$ not transversally at x in \mathbb{R}^n .

Since $\Sigma_f(\mathring{\mathbb{B}}_\epsilon) \cap V(f) \subset \{0\}$, it follows that any $x \in V(f) \cap \mathbb{S}_\epsilon^{n-1}$ is a regular point of f . Up to a change of coordinates, f is a projection on an open neighborhood of x in \mathbb{B}_ϵ^n , therefore the fibres of f near V are transverse to $\mathbb{S}_\epsilon^{n-1}$. So there exists an open neighborhood W of

$V(f) \cap \mathbb{S}_\epsilon^{n-1}$ in $\mathbb{S}_\epsilon^{n-1}$ such that any $y \in W$ is a regular point of $f|_W$. So we just have to take $\delta = \delta(\epsilon)$ sufficiently small such that $f^{-1}(\mathbb{B}_\delta^k) \cap \mathbb{S}_\epsilon^{n-1}$ is contained in W . \square

Corollary 3.15. *Any map $(f, g) : (\mathbb{K}^n, 0) \rightarrow (\mathbb{K}^2, 0)$ as in Example 3.7 satisfies all the hypothesis of Theorem 3.12, hence it induces equivalent fibrations on the tube and the sphere.*

Question 3.16. Are fibrations (5) and (7) also equivalent when f , with linear discriminant, transversality property and d -regularity, is only of class C^ℓ with $\ell \geq 1$?

4. CONIC HOMEOMORPHISMS AND d_h -REGULARITY

In this section, we want to extend the concept of d -regularity, allowing some maps to become d -regular after a homeomorphism on the target space.

Recall that given $\eta > 0$, for each point $\theta \in \mathbb{S}_\eta^{k-1}$ the set $\mathcal{L}_\theta \subset \mathbb{R}^k$ is the open segment of line that starts in the origin and ends at the point θ (but not containing these two points).

Definition 4.1. Let $h : (\mathbb{R}^k, 0) \rightarrow (\mathbb{R}^k, 0)$ be a homeomorphism. Suppose that there exists $\eta > 0$ sufficiently small, such that the restriction of h to the ball \mathbb{B}_η^k

$$(8) \quad h_\eta : \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k,$$

with $\mathcal{B}_\eta^k := h(\mathbb{B}_\eta^k)$, satisfies the following:

- (i) For each $\theta \in \mathbb{S}_\eta^{k-1}$ the image $h_\eta(\mathcal{L}_\theta)$ is a path in \mathbb{R}^k of class C^ℓ with $\ell \geq 1$;
- (ii) The inverse map h^{-1} of h is of class C^ℓ with $\ell \geq 1$ outside the origin;
- (iii) The map h^{-1} is a submersion outside the origin.

We say that the restriction (8) is a *conic homeomorphism*.

The identity map is obviously a conic homeomorphism. Given a conic homeomorphism $h_\eta : \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$, set:

$$\varphi_{h,\theta} := h(\mathcal{L}_\theta),$$

for each $\theta \in \mathbb{S}_\eta^{k-1}$.

Example 4.2. Consider the map $h(u, v) = (u, v^3)$, whose inverse is given by $h^{-1}(u, v) = (u, \sqrt[3]{v})$. Clearly, h is a differentiable homeomorphism, although its inverse h^{-1} is not differentiable at the u -axis. It is easy to check that h^{-1} is a conic homeomorphism.

Example 4.3. Now consider the map $h: (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ given by:

$$h(u, v) = \begin{cases} (\sqrt{u}, \frac{v}{2}), & \text{if } u \geq 0; \\ (-\sqrt{-u}, \frac{v}{2}), & \text{if } u < 0. \end{cases}$$

whose inverse is given by

$$h^{-1}(u, v) = \begin{cases} (u^2, 2v), & \text{if } u \geq 0; \\ (-u^2, 2v), & \text{if } u < 0. \end{cases}$$

One can check that h is a conic homeomorphism and the inverse map h^{-1} is not.

If $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ is a conic homeomorphism, then each point $x \in \mathcal{B}_\eta^k$ belongs to some smooth path $\varphi_{h, \theta(x)}$. Precisely, $\theta(x) = \frac{h^{-1}(x)}{\|h^{-1}(x)\|}$.

So if we set

$$\mathcal{S}_\eta^{k-1} := h(\mathbb{S}_\eta^{k-1}),$$

the conic homeomorphism h induces a continuous and surjective map

$$\xi_h : \mathcal{B}_\eta^k \setminus \{0\} \longrightarrow \mathcal{S}_\eta^{k-1} \\ x \longmapsto h\left(\eta \frac{h^{-1}(x)}{\|h^{-1}(x)\|}\right).$$

That is, $\xi_h(x)$ is the point where the smooth curve $\varphi_{h, \theta(x)}$ that contains x intersects \mathcal{S}_η^{k-1} . In other words, $\xi_h(x)$ sends each smooth curve $\varphi_{h, \theta}$ to the point $h(\theta) \in \mathcal{S}_\eta^{k-1}$.

So for each θ in \mathbb{S}_η^{k-1} we have that

$$h(\mathcal{L}_\theta) := \varphi_{h, \theta} = \xi_h^{-1}(h(\theta)).$$

Now, given a map $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ and a conic homeomorphism $h: (\mathbb{R}^k, 0) \rightarrow (\mathbb{R}^k, 0)$, we define the map

$$f_h := h^{-1} \circ f,$$

which we call a *conic modification* of f :

$$\begin{array}{ccc} & (\mathbb{R}^n, 0) & \\ & \swarrow f_h & \downarrow f \\ (\mathbb{R}^k, 0) & \xrightarrow{h} & (\mathbb{R}^k, 0) \end{array}.$$

Once a representative $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ of h is fixed, for each θ in \mathbb{S}_η^{k-1} we set

$$E_{f, h, \theta} := f^{-1}(\varphi_{h, \theta}).$$

Remark 4.4. For each $\theta \in \mathbb{S}_\eta^{k-1}$ we have that

$$E_{f,h,\theta} = (f_h)^{-1}(\mathcal{L}_\theta) = \left(\eta \frac{f_h}{\|f_h\|} \right)^{-1}(\theta) = E_{f_h, id, \theta}.$$

One should also notice that $E_{f,h,\theta}$ may be the empty set for some (or for any) $\theta \in \mathbb{S}_\eta^{k-1}$, since f is not necessarily surjective.

Definition 4.5. We say that a map $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ and a conic homeomorphism $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ are *compatible* in a subset $\mathcal{C} \subset \mathbb{S}_\eta^{k-1}$ if $E_{f,h,\theta}$ is a differentiable manifold, for each $\theta \in \mathcal{C}$. We say that f and h are *compatible* if f and h are compatible in \mathbb{S}_η^{k-1} .

Remark 4.6. Observe that if f has an isolated critical value, then any conic homeomorphism h is compatible with f , since each $E_{f,h,\theta}$ is the inverse image by f of a path $\varphi_{h,\theta}$ of class C^ℓ that contains no critical value.

Now we can define d -regularity up to homomorphism:

Definition 4.7 (d_h -regularity). Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with $\ell \geq 1$. Let ϵ_0 be a Milnor radius for f and let $0 < \epsilon < \epsilon_0$. Let $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ be a conic homeomorphism compatible with f in $\mathcal{C} \subset \mathbb{S}_\eta^{k-1}$ such that $f(\mathbb{B}_{\epsilon_0}^n) \subset \mathring{\mathcal{B}}_\eta^k$. We say that f is d_h -regular relative to \mathcal{C} in the ball \mathbb{B}_ϵ^n if for any $0 < \epsilon' \leq \epsilon$ the sphere $\mathbb{S}_{\epsilon'}^{n-1}$ intersects $E_{f,h,\theta}$ transversally, whenever such intersection is not empty for each $\theta \in \mathcal{C}$. If $\mathcal{C} = \mathbb{S}_\eta^{k-1}$, we just say that f is d_h -regular in the ball \mathbb{B}_ϵ^n .

Since by Remark 4.4 we have that $E_{f,h,\theta} = E_{f_h, id, \theta}$, it follows:

Proposition 4.8. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a C^ℓ -map with $\ell \geq 1$ and let $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ be a conic homeomorphism.

- (i) The map f is compatible with h in \mathcal{C} if and only if the conic modification $f_h := h^{-1} \circ f$ is compatible with the identity map in \mathcal{C} .
- (ii) Suppose that f and h are compatible in $\mathcal{C} \subset \mathbb{S}_\eta^{k-1}$. Then f is d_h -regular relative to \mathcal{C} in the ball \mathbb{B}_ϵ^n if and only if f_h is d -regular relative to \mathcal{C} in the ball \mathbb{B}_ϵ^n .

Hence the definition of d_h -regularity above generalizes the definition of d -regularity, in the sense that a C^ℓ -map with linear discriminant (as in section 2) is d -regular if and only if it is d_{id} -regular relative to $\mathcal{C} := \mathbb{S}_\eta^{k-1} \setminus \mathcal{A}_\eta$.

Example 4.9. Let $f: (\mathbb{R}^3, 0) \rightarrow (\mathbb{R}^2, 0)$ be the real analytic map given by

$$f(x, y, z) := (x^2z + y^3, x).$$

As pointed out in Example 3.9 of [12], the map f has an isolated critical value, but along the line $L := \{x - z = 0, y = 0\}$, the spaces $E_{f, id, \theta}$ are not transversal to the corresponding spheres. So f is not d -regular.

Now consider the conic homeomorphisms $h(u, v) = (u, v^3)$ and its inverse $h^{-1}(u, v) = (u, \sqrt[3]{v})$ of Example 4.2. By Remark 4.6 we have that f is compatible with both h and h^{-1} .

Let us show that f is $d_{(h^{-1})}$ -regular:

$$f_{(h^{-1})}(x, y, z) = (x^2 z + y^3, x^3),$$

it is an analytic map, whose critical set is the plane $\{x = 0\}$, and its discriminant is the line $\{v = 0\}$ in \mathbb{R}^2 . Using [12, Proposition 3.8] one can check that $f_{(h^{-1})}$ is d -regular and hence f is $d_{(h^{-1})}$ -regular.

This leads us to the following:

Question 4.10. Is there always a conic modification of a real analytic map f with isolated critical value, which is d -regular?

5. THE d_h -REGULARITY AND FIBRATION THEOREMS

In this section we study the possibility to modify the discriminant of a map by homeomorphism, making it linear. Then one can check if the new map is d -regular, and in that case we establish a fibration theorem. This process applies in particular to maps with isolated critical value and the transversality property.

Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$, with $n \geq k \geq 2$, be a map of class C^ℓ . Let ϵ_0 be a Milnor radius for f , let $0 < \epsilon < \epsilon_0$ and let Δ_ϵ be the discriminant of f_ϵ .

Definition 5.1. We say that a conic homeomorphism $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ is a linearization for f_ϵ if we have that

$$h^{-1}(\Delta_\epsilon \cap \mathcal{B}_\eta^k) = \text{Cone}(h^{-1}(\Delta_\epsilon \cap \partial \mathcal{B}_\eta^k)).$$

Note that the definition above includes the case where f_ϵ has an isolated critical value.

Recall the notation of Section 4. Given a linearization h for a map f of class C^ℓ , we have the following paths of class C^ℓ

$$\varphi_{h, \theta} = h(\mathcal{L}_\theta)$$

and the induced map

$$\xi_h: \mathbb{B}_\eta^k \setminus \{0\} \rightarrow \mathbb{S}_\eta^{k-1},$$

where $\xi_h(x)$ is the point where the closure of the path $\varphi_{h, \theta}$ that contains x intersects \mathbb{S}_η^{k-1} . So $\varphi_{h, \theta} = (\xi_h)^{-1}(\theta)$.

Also, given a linearization h for f_ϵ , we set

$$\mathcal{A}_{h,\eta} := h^{-1}(\Delta_\epsilon \cap \partial \mathcal{B}_\eta^k) = h^{-1}(\Delta_\epsilon) \cap \mathbb{S}_\eta^{k-1}.$$

Finally, for each θ in \mathbb{S}_η^{k-1} we have

$$E_{f,h,\theta} := f^{-1}(\varphi_{h,\theta}),$$

which is a manifold of class C^ℓ if $\theta \notin \mathcal{A}_{h,\eta}$. Recall that

$$E_{f,h,\theta} = (\xi_h \circ f)^{-1}(\theta) = f^{-1}(h(\mathcal{L}_\theta)).$$

So any linearization h for a map f_ϵ of class C^ℓ is compatible with f_ϵ in $\mathbb{S}_\eta^{k-1} \setminus \mathcal{A}_{h,\eta}$.

Once again, let $\pi: \mathbb{S}_\eta^{k-1} \rightarrow \mathbb{S}^{k-1}$ be the projection onto the unit sphere \mathbb{S}^{k-1} and set $\mathcal{A}_h = \pi(\mathcal{A}_{h,\eta})$. Recall the map of class C^ℓ

$$\phi_{f,h}: \mathbb{S}_\epsilon^{n-1} \setminus K_f \rightarrow \mathbb{S}^{k-1}$$

given by

$$(\phi_{f,h})(x) = \frac{f_h(x)}{\|f_h(x)\|}.$$

As before, using Proposition 4.8 together with Theorem 3.9 and Theorem 3.12, we have:

Theorem 5.2. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with the transversality property in the ball \mathbb{B}_ϵ^n , and suppose it admits a linearization $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ making f d_h -regular in the ball \mathbb{B}_ϵ^n . Then the restriction*

$$(\phi_{f,h})|: \mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(\Delta_f) \rightarrow \mathbb{S}^{k-1} \setminus \mathcal{A}_h$$

is a C^ℓ -locally trivial fibration over its image.

If f is analytic, by Theorem 3.12 we have the following corollary.

Corollary 5.3. *In the case that f is real analytic and with the transversality property in the ball \mathbb{B}_ϵ^n , the fibration given in Theorem 5.2 is equivalent to the locally trivial fibration (5) on the tube:*

$$f_h|: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon) \rightarrow \mathbb{S}_\delta^{k-1} \setminus \Delta_\epsilon.$$

5.1. Isolated critical value with transversality property. Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ , with isolated critical value. Let ϵ_0 be a Milnor radius for f , let $0 < \epsilon < \epsilon_0$ and suppose f has the transversality property in the ball \mathbb{B}_ϵ^n . In this case, one can use Milnor's vector field ([21, Theorem 11.2]) to give a fibration on the sphere, but without knowing what is the projection map. Since maps with linear discriminant include the case of isolated critical value, we can apply Theorem 5.2 to this case getting a sharper result:

Theorem 5.4. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with an isolated critical value and the transversality property in the ball \mathbb{B}_ϵ^n . If it is d_h -regular for some conic homeomorphism $h: \mathbb{B}_\eta^k \rightarrow \mathcal{B}_\eta^k$ of class C^ℓ , then the map*

$$\phi_{f,h} := (\pi \circ f_h)|: \mathbb{S}_\epsilon^{n-1} \setminus f^{-1}(0) \rightarrow \mathbb{S}^{k-1},$$

is a C^ℓ -locally trivial fibration over its image.

It is an immediate corollary of Theorem 5.2, together with the fact that the map $f_h := h^{-1} \circ f$ induces a submersion of class C^ℓ outside $f^{-1}(0)$.

In the case of f analytic with isolated critical value and transversality property in the ball \mathbb{B}_ϵ^n , we can now say more on equivalence of fibrations. By Theorem 5.4 and Theorem 3.12 the two fibrations:

$$(9) \quad \phi_{f,h}: \mathbb{S}_\epsilon^{n-1} \setminus K_f \rightarrow \mathbb{S}^{k-1}$$

and

$$(10) \quad f_h: \mathbb{B}_\epsilon^n \cap f_h^{-1}(\mathbb{S}_{\delta'}^{k-1}) \rightarrow \mathbb{S}_{\delta'}^{k-1}$$

are equivalent. This last one is clearly equivalent to the fiber bundle

$$(11) \quad f: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathcal{S}_{\delta'}^{k-1}) \rightarrow \mathcal{S}_{\delta'}^{k-1}$$

Let us show that fibration (11) is equivalent to the fibration on the tube:

$$(12) \quad f: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathbb{S}_{\delta''}^{k-1}) \rightarrow \mathbb{S}_{\delta''}^{k-1}$$

over any sphere sufficiently small such that $\mathbb{S}_{\delta''}^{k-1} \subset \mathbb{B}_\delta^k$. This is true even when f is of class C^ℓ .

Proposition 5.5. *Let $f: (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^k, 0)$ be a map of class C^ℓ with an isolated critical value and the transversality property in the ball \mathbb{B}_ϵ^n . Consider the fibration (4) on the solid tube*

$$(13) \quad f|: \mathbb{B}_\epsilon^n \cap f^{-1}(\mathring{\mathbb{B}}_\delta^k \setminus \{0\}) \rightarrow (\mathring{\mathbb{B}}_\delta^k \setminus \{0\}).$$

We have that $\pi_{k-1}(\mathring{\mathbb{B}}_\delta^k \setminus \{0\}) \cong \mathbb{Z}$. Let $\tilde{g}, \tilde{h}: \mathbb{S}^{k-1} \rightarrow \mathring{\mathbb{B}}_\delta^k \setminus \{0\}$ be two continuous embeddings such that both represent the generator of $\pi_{k-1}(\mathring{\mathbb{B}}_\delta^k \setminus \{0\})$. Then the restrictions

$$(14) \quad f|: \mathbb{B}_\epsilon^n \cap f^{-1}(g(\mathbb{S}^{k-1})) \rightarrow g(\mathbb{S}^{k-1}),$$

and

$$(15) \quad f|: \mathbb{B}_\epsilon^n \cap f^{-1}(h(\mathbb{S}^{k-1})) \rightarrow h(\mathbb{S}^{k-1})$$

are topologically equivalent fibre bundles.

Proof. For paracompact spaces locally trivial fibrations are homotopy invariant, that is, pull-backs over homotopic maps give equivalent fibre bundles (see for instance [15, §4 Theorem 9.8] plus a reduction to the principal bundle case). Since \tilde{g} and \tilde{h} are homotopic, the pull-backs of the fibre bundle (13) by \tilde{g} and \tilde{h} are equivalent. Since \tilde{g} and \tilde{h} are embeddings those pull-backs are equivalent to the restrictions (14) and (15). \square

Corollary 5.6. *The fibre bundles (11) and (12) are equivalent.*

Proof. Take \tilde{g} as the projection of the unit sphere onto the sphere $\mathbb{S}_{\delta''}^{k-1}$ and $\tilde{h} = h \circ \pi_{\delta'}$ where $\pi_{\delta'}$ is the projection of the unit sphere onto the sphere $\mathbb{S}_{\delta'}^{k-1}$. Then \tilde{g} and \tilde{h} satisfy the conditions of Proposition 5.5. \square

In particular, we have:

Corollary 5.7. *For f real analytic, the fiber bundle $\phi_{f,h}$ of Proposition 5.4 does not depend on the choice of the conic homeomorphism h (as long as f is d_h -regular), up to topological equivalence of fiber bundles.*

This means that we are not going to find different fibrations on the sphere by looking for different conic homeomorphisms h such that f is d_h -regular.

Example 5.8. Consider the real analytic map $f: (\mathbb{R}^3, 0) \rightarrow (\mathbb{R}^2, 0)$ given by

$$f(x, y, z) := (x + y, x^2 + y^2 + z^3).$$

Its critical locus Σ_f is the line $\{x = y, z = 0\}$ and its discriminant Δ_f is the parabola $\{v = \frac{u^2}{2}\}$.

Since $\Sigma_f(\hat{\mathbb{B}}_\epsilon) \cap V_f = \{0\}$, it follows from Propositions 3.13 and 3.14 that f is surjective and has the transversality property.

The germ of conic homeomorphism $h: (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ given by

$$h(u, v) = \begin{cases} (\sqrt{u}, \frac{v}{2}), & \text{if } u \geq 0, \\ (-\sqrt{-u}, \frac{v}{2}), & \text{if } u < 0, \end{cases}$$

whose inverse is given by

$$h^{-1}(u, v) = \begin{cases} (u^2, 2v), & \text{if } u \geq 0, \\ (-u^2, 2v), & \text{if } u < 0, \end{cases}$$

is compatible with f in

$$\mathcal{C} := \mathbb{S}_\eta^1 \setminus \{\theta_{\pi/4}, \theta_{3\pi/4}\},$$

and h^{-1} sends the parabola $\Delta_f = \{v = \frac{u^2}{2}\}$ to the set $L := \mathcal{L}_{\pi/4} \cup \mathcal{L}_{3\pi/4} \cup \{0\}$. So h is a linearization for f .

Moreover, the conic modification $f_h := h^{-1} \circ f$, given by

$$f_h(x, y, z) := \begin{cases} ((x+y)^2, 2(x^2+y^2+z^3)), & \text{if } x+y \geq 0, \\ (-(x+y)^2, 2(x^2+y^2+z^3)), & \text{if } x+y < 0. \end{cases}$$

is d -regular relative to \mathcal{C} . Therefore f is d_h -regular.

Hence, since $f^{-1}(\Delta_f) = \{(x-y)^2 + 2z^3 = 0\}$, it follows by Theorem 3.12 that the map

$$\phi_{f,h} = \frac{f_h}{\|f_h\|} : \mathbb{S}_\epsilon^2 \setminus \{(x-y)^2 + 2z^3 = 0\} \rightarrow \mathbb{S}_1^1 \setminus \{\theta_{\pi/4}, \theta_{3\pi/4}\}$$

is a topological locally trivial fibration (equivalent to the fibration in the tube associated to f_h).

The following example generalizes Example 3.7:

Example 5.9. Let $(f, g) : \mathbb{R}^n \rightarrow \mathbb{R}^2$ be a real analytic map of the form

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^p, \sum_{i=1}^n b_i x_i^q \right),$$

with $p, q \geq 2$ integers, such that the points $\lambda_i = (a_i, b_i)$ satisfy the Weak Hyperbolicity Hypothesis (that is, no two of them are linearly dependent). Example 3.7 concerns the case $p = q$.

An easy calculation shows that the critical set Σ of (f, g) is linear and that the discriminant Δ is the union of the parametrized curves $(a_i t^p, b_i t^q)$ in \mathbb{R}^2 (with $t \in \mathbb{R}$), for each $i = 1, \dots, n$. So it follows from Proposition 3.14 that (f, g) has the transversality property.

Moreover, if $V(f, g) := (f, g)^{-1}(0)$ has dimension bigger than zero, it follows from Propositions 3.13 that (f, g) is surjective. If $V(f, g) = \{0\}$, then (f, g) may be not surjective. Recall from Example 3.7 that if $p = q$, then $V(f, g) = \{0\}$ if and only if the origin is not in the convex hull of the points $\lambda_i = (a_i, b_i)$.

We must consider four cases:

- (1) Suppose that both p and q are odd:

Consider the germ of conic homeomorphism $h(u, v) = (u^{1/q}, v^{1/p})$, whose inverse is given by $h^{-1}(u, v) = (u^q, v^p)$. It is a linearization for the map (f, g) , since for each $i = 1, \dots, n$ we have

$$h^{-1}(a_i t^p, b_i t^q) = (a_i^q t^{pq}, b_i^p t^{pq}).$$

Let us show that (f, g) is d_h -regular. The conic modification $(f, g)_h := h^{-1} \circ (f, g)$ is given by

$$(f, g)_h = \left(\left(\sum_{i=1}^n a_i x_i^p \right)^q, \left(\sum_{i=1}^n b_i x_i^q \right)^p \right).$$

So each $E_{(f,g)_h, id, \theta}$ has one of the following forms:

$$\{f^q + \alpha g^p = 0\} \cap \{g > 0\}$$

or

$$\{f^q + \alpha g^p = 0\} \cap \{g < 0\}$$

or

$$\{g = 0\} \cap \{f > 0\}$$

or

$$\{g = 0\} \cap \{f < 0\},$$

for some $\alpha \in \mathbb{R}$. The first one corresponds to $\theta \in (0, \pi/2]$ if $\alpha \leq 0$ and to $\theta \in (\pi/2, \pi)$ if $\alpha > 0$. The second one corresponds to $\theta \in (\pi, 3\pi/2]$ if $\alpha \leq 0$ and to $\theta \in (3\pi/2, 2\pi)$ if $\alpha > 0$. The third one corresponds to $\theta = 0$ and the fourth one corresponds to $\theta = \pi$.

Since all the equalities above deal with homogeneous polynomials, it is easy to see that each $E_{(f,g)_h, id, \theta}$ intersects any sphere in \mathbb{R}^n transversally.

(2) Suppose that p is even and q is odd:

Consider the germ of conic homeomorphism $h : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$ given by

$$h(u, v) = \begin{cases} (u^{1/q}, v^{1/p}), & \text{if } v \geq 0, \\ (u^{1/q}, -(-v)^{1/p}), & \text{if } v < 0, \end{cases}$$

whose inverse is given by:

$$h^{-1}(u, v) = \begin{cases} (u^q, v^p), & \text{if } v \geq 0, \\ (u^q, -v^p), & \text{if } v < 0. \end{cases}$$

It is a linearization for (f, g) , since for each $i = 1, \dots, n$ we have:

$$h^{-1}(a_i t^p, b_i t^q) = \begin{cases} (a_i^q t^{pq}, b_i^p t^{pq}), & \text{if } t \geq 0, \\ (a_i^q t^{pq}, -b_i^p t^{pq}), & \text{if } t < 0, \end{cases} \quad \text{if } b_i \geq 0,$$

and

$$h^{-1}(a_i t^p, b_i t^q) = \begin{cases} (a_i^q t^{pq}, -b_i^p t^{pq}), & \text{if } t \geq 0, \\ (a_i^q t^{pq}, b_i^p t^{pq}), & \text{if } t < 0. \end{cases} \quad \text{if } b_i < 0.$$

The conic modification $(f, g)_h$ is given by

$$(f, g)_h(x) = \begin{cases} (f(x)^q, g(x)^p), & \text{if } g(x) \geq 0, \\ (f(x)^q, -g(x)^p), & \text{if } g(x) < 0. \end{cases}$$

Proceeding as in Case (1), one can see that (f, g) is d_h -regular.

- (3) The case when p is odd and q is even is analogous, considering the linearization given by

$$h(u, v) = \begin{cases} (u^{1/q}, v^{1/p}), & \text{if } u \geq 0, \\ (-(-u)^{1/q}, v^{1/p}), & \text{if } u < 0. \end{cases}$$

- (4) The case when both p and q are even is also analogous, considering the linearization given by

$$h(u, v) = \begin{cases} (u^{1/q}, v^{1/p}), & \text{if } u \geq 0 \text{ and } v \geq 0, \\ (-(-u)^{1/q}, v^{1/p}), & \text{if } u < 0 \text{ and } v \geq 0, \\ (u^{1/q}, -(-v)^{1/p}), & \text{if } u \geq 0 \text{ and } v < 0, \\ (-(-u)^{1/q}, -(-v)^{1/p}), & \text{if } u < 0 \text{ and } v < 0, \end{cases}$$

So we have proved:

Theorem 5.10. *Let $(f, g): \mathbb{R}^n \rightarrow \mathbb{R}^2$ be a real analytic map of the form*

$$(f, g) = \left(\sum_{i=1}^n a_i x_i^p, \sum_{i=1}^n b_i x_i^q \right),$$

where $a_i, b_i \in \mathbb{R}$ are constants in generic position and $p, q \geq 2$ are integers, as in Example 5.9. Then there exist ϵ and δ sufficiently small, with $0 < \delta \ll \epsilon$, such that the restriction

$$(f, g)|: (f, g)^{-1}(\mathbb{D}_\delta^2 \setminus \Delta) \cap \mathbb{B}_\epsilon^n \rightarrow \mathbb{D}_\delta^2 \setminus \Delta$$

is a smooth locally trivial fibration over its image, where the discriminant Δ of (f, g) is given by the parametrized curves $t \mapsto (a_i t^p, b_i t^q)$ in \mathbb{R}^2 , for each $i = 1, \dots, n$.

Moreover, we have an equivalent smooth locally trivial fibration

$$\phi: \mathbb{S}_\epsilon^{n-1} \setminus ((f, g)^{-1}(\Delta) \cap \mathbb{S}_\epsilon^{n-1}) \rightarrow (\mathbb{S}_1^1 \setminus \mathcal{A}_h) \cap \text{Im}(\phi)$$

where

$$\phi = \frac{h^{-1} \circ f}{\|h^{-1} \circ f\|}$$

for the corresponding homeomorphism h as in Example 5.9, and $\mathcal{A}_h := h^{-1}(\Delta) \cap \mathbb{S}_1^1$.

If $V(f, g) \neq \{0\}$, then the maps above are surjective.

5.2. An example of a non-analytic d_h -regular map. Consider the real function $\varsigma: \mathbb{R} \rightarrow \mathbb{R}_+$ given by:

$$\varsigma(t) := \begin{cases} e^{-1/t} & \text{if } t > 0; \\ 0 & \text{if } t \leq 0. \end{cases}$$

It is a classic example of a function that is smooth and non-analytic.

Now define $\alpha: \mathbb{R}^n \rightarrow \mathbb{R}$ by $\alpha(x) = 1 - \|x - \bar{1}\|^2$ where $\bar{1} := (1, 0, \dots, 0)$.

So the function $f: \mathbb{R}^n \rightarrow \mathbb{R}_+$ given by $f(x) := \varsigma(\alpha(x))$, is smooth and non-analytic at the origin.

Notice that:

- $f(x) = 0$ if and only if $\|x - \bar{1}\| \geq 1$;
- $f(x) = t$ for some $t > 0$ if and only if $t \leq e^{-1}$ and

$$\|x - \bar{1}\|^2 = \frac{1}{\ln t} + 1.$$

So we have that:

- (i) $\text{Im}(f) = [0, e^{-1}]$;
- (ii) $V(f) = \mathbb{R}^n \setminus \mathring{\mathbb{B}}^n(\bar{1}; 1)$, where $\mathring{\mathbb{B}}^n(\bar{1}; 1)$ is the open ball of radius 1 around the point $\bar{1}$;
- (iii) $f^{-1}(t) = \mathbb{S}^{n-1}\left(\bar{1}; \sqrt{\frac{1}{\ln t} + 1}\right)$, where $\mathbb{S}^{n-1}\left(\bar{1}; \sqrt{\frac{1}{\ln t} + 1}\right)$ is the $(n-1)$ -sphere around $\bar{1}$ of radius $\sqrt{\frac{1}{\ln t} + 1}$, for any $0 < t < e^{-1}$;
- (iv) $f^{-1}(e^{-1}) = \{\bar{1}\}$.

The gradient of α is given by

$$\nabla\alpha(x) = -2(x - \bar{1}).$$

The derivative of ς is given by

$$\varsigma'(t) = \begin{cases} \frac{1}{t^2}e^{-\frac{1}{t}} & t > 0 \\ 0 & t \leq 0. \end{cases}$$

So by the chain rule, the gradient vector of f at a point $x \in \mathbb{R}^n$ is given by:

$$\nabla f(x) = \zeta'(\alpha(x))\nabla\alpha(x) = \begin{cases} -\frac{2}{\alpha(x)^2}f(x)(x - \bar{1}), & \text{if } \|x - \bar{1}\| < 1, \\ \bar{0}, & \text{if } \|x - \bar{1}\| \geq 1, \end{cases}$$

where $\bar{0} := (0, 0, \dots, 0)$.

Hence the critical set and the discriminant of f are given by:

$$\Sigma_f = V(f) \cup \{\bar{1}\}, \quad \Delta_f = \{0, e^{-1}\}.$$

Now we consider a function g analogous to the function f . Define $\beta: \mathbb{R}^n \rightarrow \mathbb{R}$ by $\beta(x) = 4 - \|x - \bar{2}\|^2$ where $\bar{2} := (2, 0, \dots, 0)$. Define $g: \mathbb{R}^n \rightarrow \mathbb{R}_+$ by $g(x) := \zeta(\beta(x))$.

In this case we have:

- (1) $\text{Im}(g) = [0, e^{-\frac{1}{4}}]$;
- (2) $V(g) = \mathbb{R}^2 \setminus \mathring{\mathbb{B}}^n(\bar{2}; 2)$;
- (3) $g^{-1}(t) = \mathbb{S}^{n-1}\left(\bar{2}; \sqrt{\frac{1}{\ln t} + 4}\right)$, for any $0 < t < e^{-\frac{1}{4}}$;
- (4) $g^{-1}(e^{-\frac{1}{4}}) = \{\bar{2}\}$.

Doing a computation analogous to that for f , we get that the critical set and the discriminant of g are given by:

$$\Sigma_g = V(g) \cup \{\bar{2}\}, \quad \Delta_g = \{0, e^{-\frac{1}{4}}\}.$$

Finally, set the map $\Psi: \mathbb{R}^n \rightarrow \mathbb{R}^2$ given by $\Psi := (f, g)$. It is a smooth map that is not analytic at the origin.

We have that:

- (a) $\text{Im}(\Psi) \subset [0, e^{-1}] \times [0, e^{-\frac{1}{4}}]$;
- (b) $V(\Psi) = V(g) = \mathbb{R}^n \setminus \mathring{\mathbb{B}}^n(\bar{2}; 2)$, since $V(g) \subset V(f)$;
- (c) $\Psi^{-1}(t_1, t_2) = \mathbb{S}^{n-1}\left(\bar{1}; \sqrt{\frac{1}{\ln t_1} + 1}\right) \cap \mathbb{S}^{n-1}\left(\bar{2}; \sqrt{\frac{1}{\ln t_2} + 4}\right)$, for any $t_1 \neq 0$ and $t_2 \neq 0$. Notice that if the two spheres $f^{-1}(t_1)$ and $g^{-1}(t_2)$ are transverse, the intersection is either homeomorphic to a sphere \mathbb{S}^{n-2} or empty. If they are tangent, the intersection is a point;
- (d) $\Psi^{-1}(0, t_2) = \left(\mathbb{R}^n \setminus \mathring{\mathbb{B}}^n(\bar{1}; 1)\right) \cap \mathbb{S}^{n-1}\left(\bar{2}; \sqrt{\frac{1}{\ln t_2} + 4}\right)$, for any $t_2 \neq 0$. Notice that this is homeomorphic to a ball \mathbb{B}^{n-1} , except when $t_2 = e^{-\frac{1}{4}}$ we get the point $\{\bar{2}\}$.
- (e) $\Psi^{-1}(t_1, 0) = \mathbb{S}^{n-1}\left(\bar{1}; \sqrt{\frac{1}{\ln t_1} + 1}\right) \cap \left(\mathbb{R}^n \setminus \mathring{\mathbb{B}}^n(\bar{2}; 2)\right)$, for any $t_1 \neq 0$. Notice that this is the empty set.

The Jacobian matrix of Ψ at a point $x = (x_1, \dots, x_n)$ is given by the following matrix

$$\begin{bmatrix} -\frac{2}{\alpha(x)^2}f(x)(x_1 - 1) & -\frac{2}{\alpha(x)^2}f(x)x_2 & \dots & -\frac{2}{\alpha(x)^2}f(x)x_n \\ -\frac{2}{\beta(x)^2}g(x)(x_1 - 2) & -\frac{2}{\beta(x)^2}g(x)x_2 & \dots & -\frac{2}{\beta(x)^2}g(x)x_n \end{bmatrix}.$$

So we have that the critical set and the discriminant are given by

$$\Sigma_\Psi = V(f) \cup \{x_2 = \dots = x_n = 0\}, \quad \Delta_\Psi = \{(0, t_2) \mid 0 \leq t_2 \leq e^{-1/4}\} \cup \mathcal{C},$$

where \mathcal{C} is the curve in \mathbb{R}^2 given by:

$$\mathcal{C}(s) := \begin{cases} \left(e^{-\frac{1}{s(2-s)}}, e^{-\frac{1}{s(4-s)}} \right) & \text{if } s > 0; \\ (0, 0) & \text{if } s \leq 0. \end{cases}$$

In particular, Ψ **does not have** linear discriminant (see Figure 1).




FIGURE 1. The blue closed curve is the discriminant.

Remark 5.11. The discriminant Δ_Ψ divides the plane in two connected components. Here we can see the phenomenon described in the Introduction: by (c), over points inside the discriminant, the fibres are spheres \mathbb{S}^{n-2} , while over points outside the discriminant, the fibres are empty.

We claim that Ψ has the transversality property, so there is no further contribution to the discriminant by critical points of Ψ restricted to the sphere $\mathbb{S}_\epsilon^{n-1}$. Let \mathbb{B}_ϵ the closed n -ball of small radius $\epsilon > 0$ centred at the origin $\bar{0} \in \mathbb{R}^n$. Consider the $(n-1)$ -sphere $\mathbb{S}^{n-1}(\bar{1}; 1)$ of radius 1 centred at $\bar{1}$. Firstly, we want to find the equation of the $(n-2)$ -sphere

which is the intersection of the $(n - 1)$ -spheres $\mathbb{S}_\epsilon^{n-1}$ and $\mathbb{S}^{n-1}(\bar{1}; 1)$ which respectively have the equations

$$(16) \quad x_1^2 + x_2^2 + \cdots + x_n^2 = \epsilon^2$$

$$(17) \quad (x_1 - 1)^2 + x_2^2 + \cdots + x_n^2 = 1.$$

Getting x_2^2 from (16) and substituting in (17) we get that $x_1 = \frac{\epsilon^2}{2}$, so the intersection is the $(n - 2)$ -sphere with equation

$$(18) \quad x_2^2 + \cdots + x_n^2 = \epsilon^2 - \frac{\epsilon^4}{4}.$$

Now we want to compute the radius r of the $(n - 1)$ -sphere $\mathbb{S}^{n-1}(\bar{2}; r)$ with equation

$$(19) \quad (x_1 - 2)^2 + x_2^2 + \cdots + x_n^2 = r^2$$

which intersects the hyperplane $x_1 = \frac{\epsilon^2}{2}$ on the $(n - 2)$ -sphere given by (18). Substituting $x_1 = \frac{\epsilon^2}{2}$ and (18) in (19) we obtain that $r^2 = 4 - \epsilon^2$. The image of the $(n - 1)$ -sphere $\mathbb{S}^{n-1}(\bar{2}; r)$ under g is $e^{-\frac{1}{r(4-r)}}$. Any $(n - 1)$ -sphere $\mathbb{S}^{n-1}(\bar{2}; r')$ of radius $r' > r > 0$ intersects any $(n - 1)$ -sphere $\mathbb{S}^{n-1}(\bar{1}; r'')$ with $0 < r'' \leq 1$ in either, an $(n - 2)$ -sphere **contained** in the interior of the n -ball \mathbb{B}_ϵ^n or the empty set. Taking $\delta \leq e^{-\frac{1}{r(4-r)}}$ we get that the fibre $\Psi^{-1}(t_1, t_2)$ with $(t_1, t_2) \in \mathbb{B}_\delta^k \setminus \Delta_\Psi$ is either, contained in the interior of the n -ball \mathbb{B}_ϵ^n or the empty set, hence Ψ has the transversality property.

Consider the homeomorphism $h: (0, 1) \times (0, 1) \rightarrow (0, e^{-1}) \times (0, e^{-\frac{1}{3}})$ given by

$$h(u, v) := \left(e^{\frac{1}{u(u-2)}}, e^{\frac{1}{v(v-4)}} \right),$$

with inverse

$$h^{-1}(u, v) := \left(1 - \sqrt{1 + \frac{1}{\ln u}}, 2 - \sqrt{4 + \frac{1}{\ln v}} \right).$$

For $\eta < e^{-1}$ the restriction of h to $\mathbb{B}_\eta^2 \cap ((0, 1) \times (0, 1))$ is a conic homeomorphism that gives a linearization for Ψ , since h takes the segment $\mathcal{L}_{\frac{\pi}{2}}$ to itself and the segment $\mathcal{L}_{\frac{\pi}{4}}$ onto the curve \mathcal{C} (see the small rectangle in Figure 1).

Set $E_\theta := (h^{-1} \circ \Psi)^{-1}(\mathcal{L}_\theta)$ for $\theta \in (0, \frac{\pi}{2}]$. For any $\theta \in (\pi/4, \pi/2)$, one can check that E_θ is a manifold homeomorphic to the cylinder $\mathbb{S}^{n-2} \times (0, 1)$ that intersects the sphere $\mathbb{S}_{\epsilon'}$ transversally, for any $\epsilon' \leq \epsilon$,

with ϵ small enough as above, and for $\theta \in [0, \pi/4)$ we have that E_θ is empty. Moreover:

$$E_{\frac{\pi}{4}} = \{x_2 = \cdots = x_n = 0\}$$

and $E_{\frac{\pi}{2}}$ is a manifold homeomorphic to a disk \mathbb{D}^n that intersects the sphere $\mathbb{S}_{\epsilon'}$ transversally, for any $\epsilon' \leq \epsilon$. Hence Ψ is d_h -regular. Alternatively, one can check this by using Proposition 3.8 of [3] for the composition $h^{-1} \circ \Psi$.

APPENDIX A. PROOF OF LEMMA A.1

In this section we use an extension of Ehresmann Fibration Theorem proved by Wolf in [29] to prove the following:

Lemma A.1. *Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be C^ℓ -locally trivial fibrations with $1 \leq l \leq \infty$, between smooth manifolds possibly with boundary. Then $g \circ f: X \rightarrow Z$ is a C^ℓ -locally trivial fibration.*

Analogous results are given by Ekedahl [14] and McKay [18, Corollary 7].

We follow Section 2 of [29] to give the necessary definitions to enunciate Wolf's theorem. In [29] the results are stated for smooth manifolds and smooth maps between them. Here we also deal with smooth (C^∞) manifolds but the maps may be only of class C^ℓ for $1 \leq l \leq \infty$.

Let $\varphi: E \rightarrow B$ be a submersion of class C^ℓ with $1 \leq l \leq \infty$. Since φ is a submersion, for any $b \in B$ the fibre $\varphi^{-1}(b)$ is a submanifold of E of dimension $\dim E - \dim B$.

Given $x \in E$, the *vertical space* V_x at x is the subspace of $T_x E$ defined by

$$V_x = \{v \in T_x E \mid D_x \varphi(v) = 0\},$$

that is, the space tangent to the fibre $\varphi^{-1}(\varphi(x))$. One has that $\dim V_x = \dim E - \dim B$. The *vertical distribution* is $\mathcal{V} = \{V_x\}_{x \in E}$. An *Ehresmann connection* for φ is a smooth distribution $\mathcal{H} = \{H_x\}_{x \in E}$ on E that is complementary to \mathcal{V} , i.e., $T_x E = V_x \oplus H_x$ for every $x \in E$. So $D_x \varphi$ restricts to a linear isomorphism from H_x onto $T_{\varphi(x)} B$. The space H_x is the *horizontal space* at x . Notice that using a Riemannian metric on E it is always possible to construct an Ehresmann connection taking the orthogonal complement of the vertical distribution.

Fix an Ehresmann connection \mathcal{H} of $\varphi: E \rightarrow B$. A tangent vector $v \in T_x E$ is *horizontal* (respectively, *vertical*) if $v \in H_x$ (respectively, $v \in V_x$); a sectionally C^ℓ -curve in E with $1 \leq l \leq \infty$ is *horizontal*

(respectively, *vertical*) if each of its tangent vectors is *horizontal* (respectively, *vertical*). We make the convention that all sectionally C^ℓ -curves are parametrised so as to be regular (nowhere vanishing tangent vector) on each smooth arc.

Let $\alpha(t)$, $t \in [0, 1]$, be a sectionally C^ℓ -curve in $\varphi(E) \subset B$. Given $x \in \varphi^{-1}(\alpha(0))$, there is at most one sectionally C^ℓ -horizontal curve $\alpha_x(t)$, $t \in [0, 1]$, in E such that:

- (a) $\alpha_x(0) = x$, and
- (b) $\varphi \circ \alpha_x = \alpha$.

If it exists, α_x is called the *horizontal lift* of α to x . If α_x exists for every $x \in \varphi^{-1}(\alpha(0))$, then we say that α has *horizontal lifts*. In such case, the *translation of the fibres along α* is the map

$$\begin{aligned} \rho_\alpha: \varphi^{-1}(\alpha(0)) &\rightarrow \varphi^{-1}(\alpha(1)), \\ x &\mapsto \alpha_x(1), \end{aligned}$$

which is differentiable by Lemma 2.2 of [29].

Theorem A.2 (Corollary 2.5 of [29]). *Let $\varphi: E \rightarrow B$ be a submersion of class C^ℓ with $1 \leq \ell \leq \infty$, where E and B are paracompact and B connected.³ Then the following statements are equivalent:*

- (i) $\varphi: E \rightarrow \varphi(E)$ is a locally trivial differentiable fibre bundle.
- (ii) There exists an Ehresmann connection for φ , relative to which every sectionally C^ℓ -curve in $\varphi(E)$ has horizontal lifts.
- (iii) If \mathcal{H} is an Ehresmann connection for φ , then every sectionally C^ℓ -curve in $\varphi(E)$ has horizontal lifts relative to \mathcal{H} .

It is easy to extend Theorem A.2 when E is a manifold with boundary ∂E asking that the restriction $\phi|_{\partial E}: \partial E \rightarrow B$ is also a submersion and applying Theorem A.2 to the restriction of φ to the interior of E and to the restriction of φ to the boundary ∂E .

Proof of Lemma A.1. Since g is a C^ℓ -locally trivial fibration, by Theorem A.2 there exists an Ehresmann connection \mathcal{H}^g for g , relative to which every sectionally C^ℓ -curve in $g(Y) \subset Z$ has horizontal lifts. For any $y \in Y$ we have $T_y Y = V_y^g \oplus H_y^g$ where V_y^g and H_y^g are respectively the vertical and horizontal subspaces of $T_y Y$. Recall that V_y^g is the tangent space of the fibre $g^{-1}(g(y))$ at y and that H_y^g projects isomorphically onto $T_{g(y)} Z$ under $D_y g$.

Analogously, there exists an Ehresmann connection \mathcal{H}^f for f , relative to which every sectionally C^ℓ -curve in $f(X) \subset Y$ has horizontal

³The hypothesis in [29] of E being connected is not used in the proof and it works without it.

lifts. For any $x \in X$ we have $T_x X = V_x^f \oplus H_x^f$ where V_x^f and H_x^f are respectively the vertical and horizontal subspaces of $T_x X$. Recall that V_x^f is the tangent space of the fibre $f^{-1}(f(x))$ at x and that H_x^f projects isomorphically onto $T_{f(x)} Y = V_{f(x)}^g \oplus H_{f(x)}^g$ under $D_x f$. This isomorphism induces a direct sum decomposition $H_x^f = \tilde{H}_x^f \oplus H_x^{g \circ f}$, where \tilde{H}_x^f and $H_x^{g \circ f}$ correspond respectively to $V_{f(x)}^g$ and $H_{f(x)}^g$. Hence we have $T_x X = V_x^f \oplus \tilde{H}_x^f \oplus H_x^{g \circ f}$. Set $V_x^{g \circ f} = V_x^f \oplus \tilde{H}_x^f$, then we have $T_x X = V_x^{g \circ f} \oplus H_x^{g \circ f}$ and we claim that $V_x^{g \circ f}$ is the vertical space of $g \circ f$ at x and that the distribution $\mathcal{H}^{g \circ f} = \{H_x^{g \circ f}\}_{x \in X}$ is an Ehresmann connection for $g \circ f$. Firstly, it is easy to see that $H_x^{g \circ f}$ is mapped isomorphically onto $T_{g(f(x))} Z$ under $D_x(g \circ f)$

$$D_{f(x)} g(D_x f(H_x^{g \circ f})) = D_{f(x)} g(H_{f(x)}^g) = T_{g(f(x))} Z.$$

To see that $V_x^{g \circ f}$ is the vertical space of $g \circ f$ at x we need to check two cases: 1) if $v \in V_x^f$ we have that $D_x f(v) = 0$, then $D_{f(x)} g(D_x f(v)) = D_{f(x)} g(0) = 0$, 2) if $v \in \tilde{H}_x^f$ then $D_x f(v) \in V_{f(x)}^g$ and $D_{f(x)} g(D_x f(v)) = 0$.

Let $z \in (g \circ f)(X) \subset Z$, $x \in (g \circ f)^{-1}(z)$ and $y = f(x) \in g^{-1}(z) \subset Y$. Let $\alpha: I \rightarrow Z$ be a sectionally C^ℓ -curve in $(g \circ f)(X) \subset Z$ with $\alpha(0) = z$ and let $\alpha_y: I \rightarrow f(X) \subset Y$ be its horizontal lift relative to \mathcal{H}^g , so we have that $\alpha_y(0) = y$ and $g \circ \alpha_y = \alpha$. Now let $\alpha_x: I \rightarrow X$ be the horizontal lift of α_y relative to \mathcal{H}^f , so we have that $\alpha_x(0) = x$ and $f \circ \alpha_x = \alpha_y$. Thus we have $g \circ f \circ \alpha_x = g \circ \alpha_y = \alpha$, so α_x is a lift of α by $g \circ f$. To conclude the proof we need to check that α_x is a horizontal lift relative to the Ehresmann connection $\mathcal{H}^{g \circ f}$. Since $\alpha_x: I \rightarrow X$ is the horizontal lift of α_y relative to \mathcal{H}^f we have that $\alpha'_x(t) \in H_{\alpha_x(t)}^f = \tilde{H}_{\alpha_x(t)}^f \oplus H_{\alpha_x(t)}^{g \circ f}$ for every $t \in I$. We claim that $\alpha'_x(t) \in H_{\alpha_x(t)}^{g \circ f}$, suppose this is not true, that $\alpha'_x(t) \in \tilde{H}_{\alpha_x(t)}^f$, then $D_{\alpha_x(t)} f(\alpha'_x(t)) = \alpha'_y(t) \in V_{f(\alpha_x(t))}^g$, but this contradicts the fact that α_y is a horizontal lift of α relative to the Ehresmann connection \mathcal{H}^g . \square

REFERENCES

- [1] R. N. Araújo dos Santos. Equivalence of real Milnor fibrations for quasi-homogeneous singularities. *Rocky Mountain J. Math.*, 42(2):439–449, 2012. [3](#)
- [2] Raimundo N. Araújo dos Santos, Maico F. Ribeiro, and Mihai Tibăr. Fibrations of highly singular map germs. *Bull. Sci. Math.*, 155:92–111, 2019. [3](#), [4](#), [5](#)
- [3] Javier Fernández de Bobadilla and Aurélio Menegon. The boundary of the Milnor fibre of complex and real analytic non-isolated singularities. *Geom. Dedicata*, 173:143–162, 2014. [3](#)

- [4] Arnaud Bodin and Anne Pichon. Meromorphic functions, bifurcation sets and fibred links. *Math. Res. Lett.*, 14(3):413–422, 2007. [3](#)
- [5] José Luis Cisneros-Molina. Join theorem for polar weighted homogeneous singularities. In *Singularities II*, volume 475 of *Contemp. Math.*, pages 43–59. Amer. Math. Soc., Providence, RI, 2008. [3](#)
- [6] José Luis Cisneros-Molina and Raimundo N. Araújo dos Santos. About the existence of Milnor fibrations. In *Real and complex singularities*, volume 380 of *London Math. Soc. Lecture Note Ser.*, pages 82–103. Cambridge Univ. Press, Cambridge, 2010. [3](#)
- [7] José Luis Cisneros-Molina, Nivaldo G. Grulha Jr., and José Seade. On the topology of real analytic maps. *International Journal of Mathematics*, 25(7), 2014. [3](#), [5](#)
- [8] José Luis Cisneros-Molina and Aurélio Menegon-Neto. Equivalence of Milnor and Milnor-Lê fibrations for real analytic maps. To appear in *International Journal of Mathematics*, 2019. [12](#)
- [9] José Luis Cisneros-Molina, José Seade, and Jawad Snoussi. Refinements of Milnor’s fibration theorem for complex singularities. *Adv. Math.*, 222(3):937–970, 2009. [2](#)
- [10] José Luis Cisneros-Molina, José Seade, and Jawad Snoussi. Milnor fibrations and d -regularity for real analytic singularities. *Internat. J. Math.*, 21(4):419–434, 2010. [2](#), [3](#), [6](#), [10](#), [11](#), [12](#)
- [11] José Luis Cisneros-Molina, José Seade, and Jawad Snoussi. Milnor Fibrations for Real and Complex Singularities. In José Ignacio Cogolludo-Agustín and Eriko Hironaka, editors, *Topology of Algebraic Varieties and Singularities*, volume 538 of *Contemporary Mathematics*, pages 345–362. American Mathematical Society, 2011. [3](#)
- [12] José Luis Cisneros-Molina, José Seade, and Jawad Snoussi. Milnor fibrations and the concept of d -regularity for analytic map germs. In *Real and complex singularities*, volume 569 of *Contemp. Math.*, pages 1–28. Amer. Math. Soc., Providence, RI, 2012. [2](#), [3](#), [17](#)
- [13] Nicolas Dutertre and Raimundo N. Araújo dos Santos. Topology of real Milnor fibrations for non-isolated singularities. *Int. Math. Res. Not. IMRN*, (16):4849–4866, 2016. [3](#)
- [14] [Torsten Ekedahl](#). Is the composition of two bundle projections necessarily a bundle projection? MathOverflow. URL:<http://mathoverflow.net/q/74722>. [28](#)
- [15] Dale Husemoller. *Fibre Bundles*. Graduate Texts in Mathematics 20. Springer-Verlag, third edition, 1993. [20](#)
- [16] Klaus Lamotke. The topology of complex projective varieties after S. Lefschetz. *Topology*, 20(1):15–51, 1981. [2](#)
- [17] Santiago López de Medrano. Singularities of homogeneous quadratic mappings. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM*, 108(1):95–112, 2014. [4](#), [8](#)
- [18] Benjamin McKay. Morphisms of Cartan Connections. arXiv:0802.1473, 2010. [28](#)
- [19] Aurélio Menegon Neto and José Seade. On the Lê-Milnor fibration for real analytic maps. *Math. Nachr.*, 2016. Early view version. [3](#)
- [20] John Milnor. On isolated singularities of hypersurfaces. Preprint, June 1966. [2](#)

- [21] John Milnor. *Singular points of complex hypersurfaces*. Annals of Mathematics Studies, No. 61. Princeton University Press, Princeton, N.J., 1968. [2](#), [18](#)
- [22] Mutsuo Oka. Non-degenerate mixed functions. *Kodai Math. J.*, 33(1):1–62, 2010. [3](#)
- [23] Frédéric Pham. *Singularities of integrals*. Universitext. Springer London, 2011. [3](#), [5](#)
- [24] Anne Pichon. Real analytic germs $f\bar{g}$ and open-book decompositions of the 3-sphere. *Internat. J. Math.*, 16(1):1–12, 2005. [3](#)
- [25] Anne Pichon and José Seade. Fibred multilinks and singularities $f\bar{g}$. *Math. Ann.*, 342(3):487–514, 2008. [2](#), [3](#)
- [26] José Seade. On Milnor’s fibration theorem and its offspring after 50 years. *Bull. Amer. Math. Soc.*, 56:281–348, 2019. [3](#)
- [27] Maria Aparecida Soares Ruas and Raimundo Nonato Araújo dos Santos. Real Milnor fibrations and (c)-regularity. *Manuscripta Math.*, 117(2):207–218, 2005. [3](#)
- [28] Maria Aparecida Soares Ruas, José Seade, and Alberto Verjovsky. On real singularities with a Milnor fibration. In A. Libgober and M. Tibar, editors, *Trends in singularities*, Trends Math., pages 191–213. Birkhäuser, Basel, 2002. [3](#)
- [29] Joseph A. Wolf. Differentiable fibre spaces and mappings compatible with Riemannian metrics. *Michigan Math. J.*, 11:65–70, 1964. [28](#), [29](#)

INSTITUTO DE MATEMÁTICAS, UNIDAD CUERNAVACA, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, AVENIDA UNIVERSIDAD S/N, COLONIA LOMAS DE CHAMILPA, CUERNAVACA, MORELOS, MEXICO.

E-mail address: jlcisneros@im.unam.mx

UNIVERSIDADE FEDERAL DA PARAÍBA, DEPARTAMENTO DE MATEMÁTICA, CEP 58051-900, JOÃO PESSOA - PB, BRAZIL

E-mail address: aurelio@mat.ufpb.br

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, ÁREA DE LA INVESTIGACIÓN CIENTÍFICA, CIRCUITO EXTERIOR, CIUDAD UNIVERSITARIA, COYOACÁN, 04510, CIUDAD DE MÉXICO, MEXICO.

E-mail address: jseade@im.unam.mx

INSTITUTO DE MATEMÁTICAS, UNIDAD CUERNAVACA, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, AVENIDA UNIVERSIDAD S/N, COLONIA LOMAS DE CHAMILPA, CUERNAVACA, MORELOS, MEXICO.

E-mail address: jsnoussi@im.unam.mx