

INDEX ONE MINIMAL SURFACES IN POSITIVELY CURVED 3-MANIFOLDS

ANTONIO ROS

ABSTRACT. We construct a Riemannian metric of positive sectional curvature on the 3-dimensional projective space with a two-sided closed embedded minimal surface of genus 3, index 1 and nullity 0.

Mathematics Subject Classification: 53A10, 58E12, 57M50.

1. INTRODUCTION

The study of the index of minimal surfaces in Riemannian three manifolds is a topic of interest in different contexts and in particular in Min-max theory for the area functional. If M is a compact orientable Riemannian three-manifold with positive Ricci curvature, then the index of closed two-sided minimal surfaces cannot be 0 and Ketover, Marques and Neves [9] prove that admits a closed embedded two-sided minimal surface Σ of index 1 realizing the Heegaard genus of M such that its area give the width of the ambient metric. Hamilton, [7], showed that such a M is diffeomorphic to a spherical space form and therefore $Heegaard\ genus(M) \leq 2$. On the other hand, from the second variation formula of area it follows that the genus of an index 1 minimal surface Σ in M is ≤ 3 . This bound holds true even for $Ric \geq 0$, Ros [18], and it is known to be sharp for the cubic flat 3-torus, Ross [20]. The geometric significance of the option $genus(\Sigma) = 3$ is not clear and its relevance has been noted by Schoen and Neves [14].

In the case of constant curvature, Viana [22] proved that there is a positive c such that, if $\Sigma \subset M^3$ is a closed 2-sided embedded minimal surface of index 1 and genus 3 in a spherical space form M , then the order of the fundamental group of M is bounded $|\pi_1(M)| \leq c$.

In this paper, we construct, for the first time, a closed minimal surface of genus 3 and index 1 in a positively curved ambient space. We prove the following:

This work is supported in part by the IMAG–Maria de Maeztu grant CEX2020-001105-M / AEI / 10.13039/501100011033, MICINN grant PID2020-117868GB-I00 and Junta de Andalucía grant P18-FR4049.

Theorem 1. *There is a Riemannian metric g of positive sectional curvature on the real projective space RP^3 and a compact orientable embedded minimal surface Σ of genus 3 in (RP^3, g) with index 1 and nullity 0.*

As a consequence of the property of RP^3 and the result of [22], the question above about the behaviour of index one minimal surfaces could be stated as follows:

Problem 1. *Which compact orientable three-manifolds M admit a Riemannian metric g with positive Ricci curvature and a closed two-sided surface $\Sigma \subset M$ of genus 3 such that Σ is an index one minimal surface in (M, g) ? Are there only finitely many of them?*

We outline the proof of the theorem. Consider the *Euclidean projective space* $RP^3(1/2)$ given as a compact 3-manifold diffeomorphic to usual projective space $RP^3 \simeq S^3/\pm$ with a singular flat Riemannian metric. The singular set is a 1-net with six edges and four vertices, Figure 2. It is the quotient space of the cubic 3-torus $T^3(1)$ under the action of a finite group of rigid motions, see [3] and Section 3 below. By using International Crystallographic Notation [1], the group is named $I222$ and it is generated by the 2-fold screw motions around the principal axes of the cube of side $1/2$ that apply a perpendicular face into the opposite one,

$$RP^3(1/2) = T^3(1)/I222.$$

The proof of Theorem 1 depends on the fact that the flat projective space can be approximated by smooth Riemannian metrics on RP^3 with positive sectional curvature in terms of the Gromov-Hausdorff distance, the approximation being in the C^∞ topology outside of any neighborhood of the singular net.

In a second step we construct an index one minimal surface in the flat projective space. To do that we start with the Schwarz P minimal surface, Figure 3. Marty Ross [20] showed that this surface has index one in the flat 3-torus and that result plays an important role in the history of the second variation of area for minimal surfaces. If we think of $RP^3(1/2)$ as a twisted kind of the flat 3-torus, then the surface we use is a twisted version of the Schwarz P minimal surface. By combining Ross' theorem and the symmetry properties of Schwarz' surface, we show that the new surface has index 1 and nullity 0.

The above ingredients allow us to prove Theorem 1 by applying an implicit function theorem argument.

Remark 1. Note that the Riemannian metrics of Theorem 1 admit a second closed minimal surface of index 1 and genus 1 (equal to the Heegaard genus of RP^3), [9].

2. PRELIMINARIES

Let Σ be an orientable closed minimal surface immersed in an orientable Riemannian 3-manifold M . We denote by $Isom(\Sigma)$ the group of transformations on Σ that come from isometries of the ambient space M .

The second variation formula of area is given by the *index form* $Q(-, -)$ associated to the *Jacobi operator* $L = \Delta + Ric(N) + |A|^2$, where Δ is the Laplacian on Σ , $Ric(N)$ is the Ricci curvature of M along the unit normal vector N of the surface and $|A|^2$ is the square of the norm of the second fundamental A of the immersion. If $\varphi \in C^2(\Sigma)$ is an smooth function, then

$$Q(\varphi, \varphi) = \int_{\Sigma} |\nabla \varphi|^2 - (Ric(N) + |A|^2) \varphi^2 = - \int_{\Sigma} \varphi L \varphi \, dA.$$

The eigenvalues of L are noted as $\lambda_0 < \lambda_1 \cdots < \lambda_k \rightarrow \infty$, each one with finite multiplicity m_k . The eigenspace $V_k(\Sigma)$ is the m_k -dimensional space of λ_k -eigenfunctions given by the functions u such that

$$Lu + \lambda_k u = 0 \text{ on } \Sigma.$$

The eigenvalue λ_0 is of multiplicity one and the eigenspace is generated by a positive function φ_0 . In particular, φ_0 is invariant under any isometry in $Isom(\Sigma)$. The *Jacobi functions* are the eigenfunctions with an eigenvalue equal to 0 (if any) and the *nullity* of Σ is the multiplicity of the 0-eigenvalue. The immersion is said to be *stable* when $\lambda_0 \geq 0$ and the *index* of Σ is the number of negative eigenvalues (counted with multiplicities).

We will consider embedded non totally geodesic two-sided minimal surfaces $\Sigma \subset M$ in a compact orientable ambient space with nonnegative Ricci curvature. From the second variation formula we get $\lambda_0 < 0$ and so, Σ is unstable. We focus on the *index one* case (or equivalently $\lambda_1 \geq 0$).

We will also consider the second variation formula for fixed boundary. Let $D \subset \Sigma$ be a domain with piecewise smooth boundary ∂D . The eigenvalues of L will be denoted as $\mu_1 < \mu_2 < \cdots$ and the corresponding eigenfunctions satisfy the equation

$$Lv + \mu_k v = 0 \text{ on } D \text{ and } v = 0 \text{ on } \partial D.$$

The first eigenfunction φ_1 is positive and if $\mu_1 > 0$ then we say that D is *strictly stable* (for the fixed boundary problem).

Let M be a compact orientable 3-manifold and $\Sigma \subset M$ a closed two-sided surface. The surface is said to be a *Heegaard surface* if it separates M in two handle bodies $M - \Sigma = \Omega \cup \Omega'$. The *Heegaard genus* of M is the minimal genus of its Heegaard surfaces. We say that the Heegaard surface is *strongly irreducible* if any two proper essential discs $\Delta \subset \Omega$ and $\Delta' \subset \Omega'$ at different sides of Σ have nondisjoint boundaries. If M has positive Ricci curvature $Ric > 0$ and Σ is a minimal surface, then Lawson [11] proved that Σ is a Heegaard surface. The same result holds for non totally geodesic minimal surfaces in a flat 3-manifold, Meeks [13].

3. THE CRYSTALLOGRAPHIC GROUPS $I222$ AND $Immm$

We consider the solid 3-cube of side $a > 0$

$$C(a) = \{(x, y, z) \in \mathbb{R}^3 / 0 \leq x, y, z \leq a\}.$$

Its principal axes are the straight lines passing through the center of the cube $(a/2, a/2, a/2)$ and parallel to the coordinate axes. If we identify opposite faces of $C(a)$ by the translations

$$(x, y, 0) \mapsto (x, y, a) \quad (x, 0, z) \mapsto (x, a, z) \quad (0, y, z) \mapsto (a, y, z)$$

we obtain the *cubic* 3-torus $T^3(a)$ with the flat Riemannian metric. Equivalently, the flat torus is obtained as the quotient of the Euclidean 3-space by the lattice Γ_a generated by the vectors $(a, 0, 0)$, $(0, a, 0)$ and $(0, 0, a)$, $T^3(a) = \mathbb{R}^3 / \Gamma_a$ and $C(a)$ is a fundamental region of the torus.

We will consider in particular the cases $a = 1$ and $a = 1/2$.

We have the 8:1 covering map

$$\Pi : T^3(1) \longrightarrow T^3(1/2).$$

We will also consider the tiling of $T^3(1)$ by eight cubes of edge $1/2$,

$$(1) \quad T^3(1) = \bigcup_{\alpha_i=0,1/2} C(1/2) + (\alpha_1, \alpha_2, \alpha_3).$$

In this case $C(1/2)$ is a fundamental region of that tiling, see Figure 1 (a).

Throughout this paper, by a *crystallographic group* \mathcal{G} we mean a finite subgroup of the isometry group of the cubic 3-torus $T^3(1)$, $\mathcal{G} \subset Isom(T^3(1))$, see the Bilbao Crystallographic Server [1]. Following [1], to describe rigid motions of $T^3(1)$ we will simply write the image of




FIGURE 1.

a general point (x, y, z) of \mathbb{R}^3 , the quotient by the lattice must be understood. The torus $T^3(1)$ is the quotient by the translations $(x + 1, y, z)$, $(x, y + 1, z)$, $(x, y, z + 1)$.

Let us consider the three order-2 screw motions whose screw axes are the principal axes of $C(1/2)$, one of which transforms the face $x = 0$ into the opposite one $x = 1/2$, and the same for the two other directions, Figure 1 (b),

$$(2) \quad \begin{cases} (-x + 1/2, -y + 1/2, z + 1/2) \\ (-x + 1/2, y + 1/2, -z + 1/2) \\ (x + 1/2, -y + 1/2, -z + 1/2). \end{cases}$$

The screw-motions (2) generate the group $I222$ of order 8 which, viewed as a transformation group on $T^3(1)$, is given by

$$(3) \quad I222 = \begin{cases} (0, 0, 0) + (1/2, 1/2, 1/2) + \\ (x, y, z) \quad (-x, -y, z) \quad (-x, y, -z) \quad (x, -y, -z). \end{cases}$$

The rigid motions of the group consists of the transformations in the second row plus any of the translations given in the first row. Its elements are as follows

- the identity map,
- the three screw motions given by equation (2),
- three axial symmetries around the edges of $C(1/2)$ (composition of two screw motions). Each one of these symmetries has four parallel axes in the torus $T^3(1)$.
- The central translation $(x + 1/2, y + 1/2, z + 1/2)$ (composition of the three screw motions).

If we add the mirror symmetry $(x, y, -z)$ to the screw motions (2), then the generated crystallographic group $Immm$ is given by the sixteen transformations of $T^3(1)$, see [1],

$$(4) \quad Immm = \left\{ \begin{array}{ll} (0, 0, 0) + & (1/2, 1/2, 1/2) + \\ (\pm x, \pm y, \pm z). \end{array} \right.$$

The group $Immm$ is formed by eight orientation preserving motions $Immm^+ = I222$ and eight orientation reversing ones, $Immm^-$. So $I222$ is a subgroup of $Immm$, the quotient is the cyclic group of order two $Immm/I222 = \mathbb{Z}_2$ and $Immm^-$ is a coset of that quotient.

The rigid motions of $Immm^-$ are described as follows:

- Two central symmetries:
 - one centered at the eight vertices of the cube $C(1/2) \subset T^3(1)$,
 - and the other centered at the middle points of the eight cubes of the tiling (1).
- Three planar reflections at the faces of the cube $C(1/2)$. The mirror planes of each one of these reflections are given by
 - $x = 0$ and $x = 1/2$,
 - $y = 0$ and $y = 1/2$ and
 - $z = 0$ and $z = 1/2$.
- Three glide reflections (no fixed points).

Note that the reflections $(-x, y, z)$ and $(1 - x, y, z)$ are different in \mathbb{R}^3 but they coincide if viewed as transformations of the 3-torus $T^3(1)$.

Another system of generators of $Immm$ is given by the following rigid motions of \mathbb{R}^3 :

$$(5) \quad \left\{ \begin{array}{l} \text{six mirror symmetries with respect to the faces of } C(1/2) \\ \text{and the central symmetry with respect to its center.} \end{array} \right.$$

3.1. The Euclidean projective space. If the crystallographic group $\mathcal{G} \subset Isom(T^3(1))$ consists of orientation preserving rigid motions, then the map $T^3(1) \rightarrow T(1)^3/\mathcal{G}$ is called an (orientable) *Euclidean orbifold*. The quotient space $T(1)^3/\mathcal{G}$ admits the structure of a closed orientable 3-manifold, called the *underlying space*, endowed with a flat riemannian metric with singularities. The *singular set* is the 1-cycle given by the image of fixed points of elements of $\mathcal{G} - \{Identity\}$ and the quotient map $T^3 \rightarrow T^3(1)/\mathcal{G}$ is a branched riemannian covering. For a presentation of Euclidean orbifolds see Dunbar [3].

The real projective space RP^3 is the manifold obtained by identifying antipodal points of the boundary sphere of the Euclidean 3-ball.

Following a related point of view, let us introduce the *Euclidean projective space* $RP^3(1/2)$ the (Alexandrov) metric space obtained from the region $C(1/2)$ by identifying each face of the cube with the opposite one by using the equivalent relation

$$(6) \quad \sim = \begin{cases} (x, y, 0) \sim (1/2 - x, 1/2 - y, 1/2) \\ (x, 0, z) \sim (1/2 - x, 1/2, 1/2 - z) \\ (0, y, z) \sim (1/2, 1/2 - y, 1/2 - z) \end{cases} \quad RP^3(1/2) = C(1/2)/\sim,$$

see Figure 1 (b). From (2) and (6) we have that the Euclidean projective space is the *orbifold* space given as the quotient

$$(7) \quad RP^3(1/2) = T^3(1)/I222, \quad \Pi^* : T^3(1) \longrightarrow RP^3(1/2).$$

The projection map Π^* is 8 to 1 and $RP^3(1/2)$ is a flat riemannian 3-manifold with singularities. The singular set is the image by Π^* of the edges of the cube $C(1/2) \subset T^3(1)$.




FIGURE 2. Picture of the Euclidean orbifold $T^3(1)/I222$ appearing in the PhD thesis of Dunbar [3], p 51. The underlying space is the projective space given as an Euclidean 3-ball with antipodal points identified. The 1-cycle inside the ball represents the singular set of the projective manifold.

4. THE SCHWARZ' P MINIMAL SURFACE

The *Schwarz' P minimal surface* $\Sigma \subset T^3(1/2)$ is a closed orientable minimal surface of genus 3 embedded in the cubic 3-torus, see Figure 3, which consists of 24 congruent skew minimal quadrilaterals. It contains 12 closed straight lines parallel to the plane coordinates forming a net with 8 triple vertices and 12 double ones.

The Gauss map $N : \Sigma \longrightarrow S^2(1)$ is a conformal map of degree 2 whose branch values are the eight vertices of the cube inscribed in




FIGURE 3. The fundamental piece S of Schwarz' P minimal surface in $C(1/2)$ has genus 0 and is orthogonal to the boundary of the cube along six congruent convex curves. So, it is a free boundary minimal surface in the cube and identifying the opposite faces we obtain the Schwarz P minimal surface in the 3-torus $\Sigma \subset T^3(1/2)$.

the unit sphere $a(\pm 1, \pm 1, \pm 1)$ with $a = 1/\sqrt{3}$. In particular, Σ is a hyperelliptic Riemann surface.

The isometry group of the P surface is $Isom(\Sigma) = Im\bar{3}m$ of order 96, and it is obtained by composition of the following rigid motions:

- a) the translations $(1/2, 0, 0), (0, 1/2, 0), (0, 0, 1/2)$ generating the simple cubic lattice,
- b) the 48 isometries of the group of the cube $m\bar{3}m$ and
- c) the body centered translation $(x + 1/4, y + 1/4, z + 1/4)$.

The subgroup of *side preserving* isometries of Σ , i.e. that preserve the unit normal vector N or, equivalently, the components of $T^3(1/2) - \Sigma$, is $Isom(\Sigma)^+ = Pm\bar{3}m$ of order 48 and is generated by the transformations in a) and b).

The body centered translation c) interchanges these components and belongs to the class of side reversing isometries $Isom(\Sigma)^-$. The groups under consideration verify the relationship

$$(8) \quad I222 \subset Immm \subset Pm\bar{3}m \subset Im\bar{3}m.$$

Ross showed the following interesting property of the Schwarz P minimal surface.

Theorem 2 (Ross [20]). *The Schwarz' P minimal surface $\Sigma \subset T^3(1/2)$ has index 1 and nullity 3.*

In particular, the only Jacobi functions on Σ are the linear functions of its Gauss map. Some other classical periodic minimal surfaces have index one too, see Ejiri and Shoda [4]. The known proofs depend (at some moment) of explicit calculations. For more about index one minimal surfaces in flat 3-manifolds see [17, 16, 18].

4.1. The Square Catenoid. The planar 2-tori $z = 1/8$ and $z = 3/8$ divide $T^3(1/2)$ in two connected components. The part of Σ in each one of these components is called the *Square Catenoid*. It is an annulus bounded by a pair of squares lying in the above horizontal planes and at intermediate heights the horizontal sections are convex curves, see Figure 4,

$$Cat = \Sigma \cap \{1/8 \leq z \leq 3/8\}, \quad Cat' = \Sigma \cap \{3/8 \leq z \leq 5/8\}.$$

Moreover, both annuli are congruent and the centered traslation by the vector $(1/4, 1/4, 1/4)$ transforms one into the other,

$$Cat + (1/4, 1/4, 1/4) = Cat' \quad \text{and} \quad Cat \cup Cat' = \Sigma.$$




FIGURE 4. The Square Catenoid is a minimal annulus bounded by two parallel squares of side $\sqrt{2}/4$ which differ by the vertical translation $(0, 0, 1/4)$. It decomposes into twelve congruent minimal quadrilaterals and the Schwarz' P surface is the union of two square catenoids.

Although we will not give a proof of Theorem 2, we show the following consequence, which in fact takes part of the inside arguments of [20] and is equivalent to the theorem itself.

Corollary 3. *The Square Catenoid is strictly stable (for the fixed boundary problem).*

Proof. Let $v : Cat \rightarrow \mathbb{R}$ be the first eigenfunction of the Square Catenoid

$$Lv + \mu_1 v = 0 \text{ on } Cat \quad \text{and} \quad v = 0 \text{ at } \partial Cat$$

We know that v is positive in the interior of Cat and if we extend v as 0 on Cat' we get a nonzero piecewise smooth function $v : \Sigma \rightarrow \mathbb{R}$. If we repeat the same procedure by changing the role of Cat and Cat' we obtain a second piecewise function v' on Σ with support equal to Cat' such that

$$\int_{\Sigma} vv' = 0 \quad Q(v, v') = 0.$$

As v and v' are linearly independent, if we denote by U the plane generated by v and v' , then for any $u \in U$ we have

$$(9) \quad Q(u, u) = \mu_1 \int_{\Sigma} u^2.$$

Reasoning by contradiction, assume that $\mu_1 \leq 0$. Let us take $u \in U - \{0\}$ orthogonal to the λ_0 -eigenfunction φ_0 on Σ , i. e.

$$\int_{\Sigma} u\varphi_0 = 0.$$

As Σ has index 1, from (9) we conclude that $\mu_1 = 0$ and so u is a Jacobi function. The nullity of Σ is equal to 3 and so it follows that u is linear function of the Gauss map N . At the boundary of Cat we have $u = 0$. This implies that u is identically zero and this contradiction gives $\mu_1 > 0$. Therefore the Square Catenoid is strictly stable. \square

4.2. The Twisted P minimal surface. Let us start with the covering $\Pi : T^3(1) \rightarrow T^3(1/2)$ in (7) and the Schwarz' P minimal surface $\Sigma \subset T^3(1/2)$. We consider the pullback image and the restricted 8:1 covering map

$$\Sigma^{-1} = \Pi^{-1}(\Sigma) \subset T^3(1) \quad \Pi : \Sigma^{-1} \rightarrow \Sigma.$$

The closed minimal surface Σ^{-1} has genus 17, see Figure 5.

From now on, the transformation groups $I222$ and $Immm$ and the coset $Immm^-$, will be viewed as acting either on the unit 3-torus or on the surface $\Sigma^{-1} \subset T^3(1)$, depending on the case.

We define the *twisted P minimal surface* Σ^* in the flat projective space $RP^3(1/2)$ by taking the quotient by the group $I222$,

$$\Sigma^* = \Sigma^{-1}/I222 \quad RP^3(1/2) = T^3(1)/I222 \quad \Sigma^* \subset RP^3(1/2).$$

The quotient surface Σ^* is a compact, orientable embedded minimal surface with $genus(\Sigma^*) = 3$ that does not meet the singular set of $RP^3(1/2)$. We will also consider the 8:1 covering maps

$$\Pi^* : T^3(1) \rightarrow RP^3(1/2) \quad \Pi^* : \Sigma^{-1} \rightarrow \Sigma^*$$

with covering transformations group $I222$. Note that Π^* is a branched covering between the 3-torus and the real projective space and when restricted to the surfaces Σ^{-1} and Σ^* , it defines a (unbranched) covering map.




FIGURE 5. The fundamental domain of the surface Σ^{-1} on the cube of side 1 is formed by eight copies of the fundamental piece S of the Schwarz surface Σ .




FIGURE 6. For later use it will be of interest to retain the set of straight lines in the Schwarz P surface $\Sigma^{-1} \subset T^3(1)$. Each of these lines is parallel to one of the coordinate planes and the figure shows the straight lines (in red) at height $z = 1/8$ in the cube $C(1)$.

Another way of thinking about the twisted surface is as follows. Identifying the boundary of the cube $C(1/2)$ by the translations vectors $(1/2, 0, 0)$, $(0, 1/2, 0)$, $(0, 0, 1/2)$, the fundamental piece S gives the closed *Schwarz' P minimal surface* in the cubic 3-torus $\Sigma \subset T^3(1/2)$, see Figure 3. However, if we identify the opposite faces of the cube by

using the screw motions in (6) we obtain the *twisted P minimal surface* of genus 3 in the flat projective space $\Sigma^* \subset RP^3(1/2)$.

In this subsection we will show one of the central results of the paper: The twisted Schwarz P surface has index 1 and nullity 0, Theorem 4. If the fundamental piece S of the Schwarz P minimal surface in Figure 3 had index 1 for the free boundary problem in the cube $C(1/2)$, then Theorem 4 will follow as a direct consequence. But it is known that free boundary minimal surfaces of the cube with six boundary components cannot have index 1, see Theor. 5 in Ros and Vergasta [19]. This illustrates the need for a more elaborate argument.

A Continuous function on the twisted P surface Σ^* will be thought of as one on the pullback surface Σ^{-1} , $\Pi^* : \Sigma^{-1} \rightarrow \Sigma^*$, invariant under the group $I222$

$$u : \Sigma^{-1} \rightarrow \mathbb{R}, \quad \text{with } u \circ \phi = u \quad \forall \phi \in I222.$$

We say that a function u of this type is *odd* (with respect to the pair $I222 \subset Immm$) if $u \circ \phi = -u$, $\forall \phi \in Immm^-$. In the case u is invariant under of all the rigid motions of $Immm$, then we say that u is *even*. As $Immm/I222 = \mathbb{Z}_2$, then the space of continuous $I222$ -invariant functions on Σ^{-1} decomposes as a direct sum

$$\{u \in C^0(\Sigma^{-1}) / u \circ \phi = u, \forall \phi \in I222\} = Odd \oplus Even.$$

On the minimal surface $\Sigma^* \subset RP^3(1/2)$ we have the Jacobi operator $L = \Delta + |A|^2$, the eigenvalues $\lambda_k^* = \lambda_k(\Sigma^*)$ and the eigenspaces $V_k(\Sigma^*)$. In this section we will see these objects over Σ^{-1} . The eigenspaces $V_k(\Sigma^*) = \{u : \Sigma^* \rightarrow \mathbb{R} / Lu + \lambda_k^* u = 0\}$ will be identified with the space of λ_k^* -eigenfunctions on Σ^{-1} which are invariant under the transformations of $I222$,

$$(10) \quad W_k = \{u : \Sigma^{-1} \rightarrow \mathbb{R} / Lu + \lambda_k^* u = 0, \quad u \circ \phi = u \quad \forall \phi \in I222\}.$$

The minimal surface Σ^{-1} could have other eigenvalues that do not admit $I222$ -invariant eigenfunctions. These are not considered here. However, the smallest eigenvalue λ_0^* of Σ^* is also the smallest eigenvalue of Σ and Σ^{-1} , $\lambda_0^* = \lambda_0 < 0$. This is because, in all the cases, the first eigenfunction is invariant under all the isometries of the surface. In particular, the λ_0 -eigenspace on Σ^{-1} is generated by a positive even eigenfunction $\varphi_0 : \Sigma^{-1} \rightarrow \mathbb{R}$. Moreover, the decomposition of W_k into *Odd* and *Even* functions holds on the eigenspaces W_k , for any k , too.

From theorem 2 we have $\lambda_1 = \lambda_1(\Sigma) = 0$. Now we will prove that $\lambda_1^* = \lambda_1(\Sigma^*) > 0$.

Theorem 4. *The Twisted P minimal surface $\Sigma^* \subset RP^3(1/2)$ has index 1 and nullity 0.*

Proof. The first eigenvalue λ_0^* is negative and the eigenspace of W_0 is generated by a positive function $\varphi_0 \in W_0$ invariant under all the isometries of the surface Σ^{-1} .

The space W_1 consists of λ_1^* -eigenfunctions of Σ^{-1} invariant under the group $I222$ and the statement of the theorem means that $\lambda_1^* > 0$.

Reasoning by contradiction, suppose that $\lambda_1^* \leq 0$ and then we will obtain the two *Claims* bellow.

Claim 1. *Every function $u \in W_1$ is invariant under the group $Immm$, i.e. u is even.*

In fact, if $u \neq 0$ is odd then u vanishes at the fixed points of all the isometries $\phi \in Immm^-$. Therefore u is equal to zero at the six Jordan curves in $\Sigma \cap \partial C(1/2)$, as it are the fixed points of the three mirror symmetries of the group $Immm$, see Figure 3. After identifying opposite sides of the cube, we obtain a continuous piecewise smooth function \tilde{u} on the Schwarz P surface Σ defined as u restricted to $\Sigma \cap C(1/2) = \Sigma^* \cap C(1/2)$. Moreover, u and φ_0 are orthogonal on Σ^* , so we have

$$\int_{\Sigma} \tilde{u} \varphi_0 = \int_{\Sigma^*} u \varphi_0 = 0, \quad \text{and} \quad Q_{\Sigma}(\tilde{u}, \tilde{u}) = Q_{\Sigma^*}(u, u) \leq 0.$$

From Theorem 2 we conclude that $\lambda_1^* = 0$ and u is a linear function of the Gauss map of the Schwarz P minimal surface. But none of these linear functions is well defined on the twisted P surface Σ^* and this contradiction proves the claim.

The axial symmetries of the group $I222$ are those whose axes are the edges of the cube $C(1/2)$. Hence every eigenfunction $u \in W_1$ is invariant under these involutions.

Now we will show that if $\lambda_1^* \leq 0$ then u is also invariant under other types of axial symmetries: Those given by the straight lines contained in the Schwarz P surface, see Figure 3.

Claim 2. *The functions in W_1 are invariant under the axial symmetries whose axes are contained in Σ^{-1} .*

To prove claim 2, suppose on the contrary that there is $u \in W_1 - \{0\}$ anti-invariant with respect to a horizontal straight line $\mathcal{L} \subset T^3(1)$ contained in $\Sigma \cap \{z = 1/8\}$ (a red line in Figure 6). In particular, u vanishes along \mathcal{L} . Claim 1 says that u is invariant under $Immm$ and therefore u is symmetric with respect to the faces of the cube $C(1/2)$. It

follows that u vanishes at all the straight lines in the Schwarz minimal surface that are at that same height than \mathcal{L} : all the red lines in Figure 6.

The function u is also invariant by the central symmetry at $(1/4, 1/4, 1/4) \in C(1/2)$ (recall this motion belongs to the group $Immm$) and, therefore, the function u vanishes at the horizontal lines in Σ^{-1} at height $z = 3/8$, too. Therefore u gives an eigenfunction for the fixed boundary problem of the Square Catenoid and Corollary 3 contradicts that $\lambda_1^* \leq 0$. That proves the claim.

Now we continue with the proof of Theorem 4. Recall that, reasoning by contradiction, we are assuming that $\lambda_1^* \leq 0$. Consider two straight lines $\mathcal{L}, \mathcal{L}' \subset \Sigma^{-1}$ in the Schwarz surface such that $\mathcal{L} \subset \{z = 1/8\}$ and $\mathcal{L}' = \mathcal{L} + (0, 0, 1/4) \subset \{z = 3/8\}$. From *Claim 2*, any function $u \in W_1$ is invariant under the axial symmetries with respect to \mathcal{L} and \mathcal{L}' . So, u is invariant with respect to the translation $(0, 0, 1/2)$ (the composition of the two reflections above).

In the same way, we have that u is invariant under the translations $(1/2, 0, 0), (0, 1/2, 0)$ and therefore it is well-defined on the Schwarz surface $\Sigma \subset T^3(1/2)$. From Theorem 2 we have that $\lambda_1^* = 0$ and u is a linear function of the Gauss map of the Schwarz P surface, $u = \langle N, a \rangle$ for some $a \in \mathbb{R}^3 - 0$.

Let $C = \{z = 0\} \cap S$ be the convex curve at the bottom of ∂S , Figure 3. As the normal vector N is horizontal along C and u is well-defined in both Σ and Σ^* , it follows that $u = 0$ on C . Then a is a vertical vector. Repeating the argument for the directions $(1, 0, 0)$ and $(0, 1, 0)$ we get $a = 0$ and this contradiction concludes the proof. \square

5. THE MAIN RESULT

The flat projective space is the quotient of the 3-torus $T^3(1)$ by the group $I222$ or, equivalently, by identifying the points of the boundary of the cube by the order-two screw motions in (6)

$$RP^3(1/2) = T^3(1)/I222 = C(1/2)/\sim.$$

To construct an approximation of the *singular* flat projective space $RP^3(1/2)$ by nonsingular Riemannian manifolds, we consider the hyperplane $\mathbb{R}^3 = \{(x, y, z, t) \in \mathbb{R}^4 / t = 0\}$, $\varepsilon > 0$ and the following figures:

- the cubes $C(1/2)$ and $C(1/2 + \pi\varepsilon)$ in \mathbb{R}^3 and the homothety $h : C(1/2) \longrightarrow C(1/2 + \pi\varepsilon)$ of ratio $1 + 2\pi\varepsilon$.




FIGURE 7. Two-dimensional version of the main construction in the proof of Theorem 1. The objective is to obtain two blue domains which, after identification, are isometric from the intrinsic point of view. One of them lies in the *flat projective plane* and the other one in the usual projective plane with a nonnegatively curved metric. Left: We have a blue domain inside the square in the plane meeting its boundary orthogonally. The sides of the square are identified by the glide motions whose axes are the axes of the figure. The generated plane crystallographic group is pgg and the quotient orbifold T^2/pgg is the *flat projective plane* with two singular points (corresponding to the vertices of the square). Right: The planar square has been deformed into a centrally symmetric, smooth convex surface that meets the plane $z = 0$ orthogonally. The central part remains horizontal and near the edges of the square (outside of a neighborhood of its vertices) it has been folded into cylindrical pieces. Around the vertices we can deform the initial domain into any convex shape. By identifying antipodal points of the boundary, we get a blue domain in the usual projective plane with a metric of nonnegative curvature. The two constructed blue domains become isometric, as we wanted. Note that the gluing along the edges is of class C^1 at the level of the immersed convex surface, but it is of class C^∞ if we just consider the induced flat Riemannian surface structure.

- convex ball $B_\varepsilon = \{p \in \mathbb{R}^4 / \text{dist}(p, C(1/2)) \leq \varepsilon\}$ in \mathbb{R}^4 and the top part of its boundary $E_\varepsilon = \partial B_\varepsilon \cap \{t \geq 0\}$.

E_ε is a C^1 convex graph over $\{p \in \mathbb{R}^3 / \text{dist}(p, C(1/2)) \leq \varepsilon\} \subset \mathbb{R}^3$, its boundary meets orthogonally the hyperplane $t = 0$ and it consists of the union of the following pieces:

- 1) A flat part parallel to the cube, $C(1/2) + (0, 0, 0, \varepsilon)$.

- 2) A part made by quarter of cylinders of the type $S^1(\varepsilon) \times \mathbb{R}^2$ following the faces of the cube. Each one of these pieces is isometric (as a Riemannian manifold) to a region of \mathbb{R}^3 .
- 3) Hypercylindrical pieces $\{(x, y, z) \in S^2(\varepsilon) / x, y, z \geq 0\} \times \mathbb{R}$ along the edges of the cube.
- 4) Spherical caps as $\{(x, y, z, t) \in S^3(\varepsilon) / x, y, z, t \geq 0\}$ centered at the vertices of the cube.

Lemma 5. *There are two domains $D \subset C(1/2)$ and $D_\varepsilon \subset E_\varepsilon$ and a dilation map $\phi : D \longrightarrow D_\varepsilon$ satisfying the following:*

- i) *D is equal to $C(1/2)$ minus a small neighborhood of the edges of the cube.*
- ii) *The domain D_ε is the union of the pieces 1) and 2) above. From the intrinsic point of view, D_ε is the piecewise smooth C^∞ flat Riemannian manifold.*
- iii) *The ratio of the dilation $\phi : D \longrightarrow D_\varepsilon$ is $1 + 2\pi\varepsilon$.*

Proof. The union along the boundary of the halfspace $\{(x, y, z, \varepsilon) \in \mathbb{R}^4 / z \geq 0\}$ and the semi cylinder $\mathbb{R}^2 \times \{(z, t) \in S^1(\varepsilon) / z \leq 0\}$ give a hypersurface of class C^1 in \mathbb{R}^4 . However, from the Riemannian point of view, the union is a C^∞ manifold isometric to an open subset of \mathbb{R}^3 .

As a consequence, the union of the pieces of type 1) and 2) is a C^∞ flat Riemannian manifold D_ε contained in E_ε . Furthermore we have:

- A domain $D'_\varepsilon \subset C(1/2 + \pi\varepsilon)$ and a Riemannian isometry $f : D'_\varepsilon \longrightarrow D_\varepsilon$.
- the pullback homothetic image $D = h^{-1}(D'_\varepsilon)$, $h : D \longrightarrow D'_\varepsilon$, and
- the C^∞ -dilation map $\phi = h \circ f$ of ratio $1 + 2\pi\varepsilon$ between flat manifolds

$$\phi : D \longrightarrow D_\varepsilon \quad D \subset C(1/2) \quad \text{and} \quad D_\varepsilon \subset E_\varepsilon.$$

The transformation ϕ is the composition of a Euclidean homothety and a Riemannian isometry. \square

Proof of Theorem 1. Since ∂B_ε is formed by pieces of hyperplanes and hypercylinders of the type 1), 2), 3) y 4) above, it is a convex hypersurface of class C^1 but not C^2 .

A compact convex body in $\Omega \subset \mathbb{R}^4$ can be approximated by a sequence of smooth convex bodies Ω_n such that the principal curvatures of $\partial\Omega_n$ are positive and tend to the principal curvatures of $\partial\Omega$ in the

sense of the surface area measure, Gruber [5] p 133, Weil [23]. For the case of abstract positively curved polyhedral 3-manifolds we can also use the alternative smoothing result in [12].

Therefore, in our case, we have that the convex hypersurface ∂B_ε can be approximated in the C^1 topology by (a centrally symmetric) strictly convex compact hypersurface S_ε of class C^∞ . Moreover, by Lemma 5, the approximation can be assumed of class C^∞ over D_ε .

For small $\varepsilon > 0$, if we identify antipodal points of the boundary we obtain two metric copies of the 3-dimensional projective space

$$RP_\varepsilon^3 = E_\varepsilon / \sim \quad \text{and} \quad \overline{RP}_\varepsilon^3 = \overline{E}_\varepsilon / \sim$$

that can be taken arbitrarily close in the C^1 -topology and satisfy the following:

i) RP_ε^3 contains an smooth flat domain $U_\varepsilon = D_\varepsilon / \sim$ obtained from D_ε by identifying antipodal points of ∂D_ε at level $t = 0$. Furthermore this domain contains itself an dilated image of the twisted P minimal surface $\Sigma_\varepsilon^* \subset U_\varepsilon$.

ii) $\overline{RP}_\varepsilon^3$ is a C^∞ Riemannian manifold with positive sectional curvature containing a domain \overline{U}_ε as close as we want of U_ε in the C^∞ topology.

From Theorem 4, the twisted P minimal surface Σ_ε^* has index 1 and nullity 0 and, therefore, the theorem follows from the implicit function theorem. \square

The existence of index one closed embedded minimal surfaces in spaces of positive Ricci curvature follows from Ketover, Marques and Neves [9]. There are minimal genus Heegaard surfaces, $genus(\Sigma) \leq 2$. D. Ketover, Y. Liokumovich and A. Song [8] obtain index one minimal surfaces in compact Riemannian 3-manifolds, without curvature assumptions, isotopic to a given strongly irreducible Heegaard splitting. These results follow Almgren-Pitts [15] and Simon-Smith [21] approaches. The surfaces can also been constructed by using Min-max theory for Allen-Cahn equation, see Guaraco [6], Chodosh and Mantoulidis [2] and references therein.

Although the Schwarz P surface Σ in $T^3(1/2)$ is a Heegaard splitting of minimum genus, the min-max construction in this space does not give the P surface but a multiplicity 2 horizontal 2-torus. In the flat projective space $RP^3(1/2)$, the Heegaard surface Σ^* is not even a minimum genus.

Another remarkable topological property of these surfaces is the following. Let S a Heegaard surface in a closed orientable 3-manifold M and $\Omega, \Omega' \subset M$ be the two complementary handlebodies. The Heegaard splitting is said to be *strongly irreducible* if any pair of essential discs $\Delta \subset \Omega$ and $\Delta' \subset \Omega'$ intersect at their boundary. In our situation, neither the Heegaard splitting given by the Schwarz P surface $\Sigma \subset T^3(1/2)$ nor the one given by the twisted P surface $\Sigma^* \subset RP^3(1/2)$ are strongly irreducible. In fact, if we consider the *inside/outside* regions determined by the P surface in the cube $C(1/2)$, Figure 3, then the disc Δ given by $\{z = 0\} \cap \text{Inside}$ is an essential disc and $\{z = 1/4\} \cap \text{Outside}$ consists in four quadrants that after identifications, in both $T^3(1/2)$ and $RP^3(1/2)$, give a second essential disc Δ' at the other side, the boundary curves being disjoint.

As the strong irreducibility implies the minimal genus property, see e.g. [10], the *index 1/genus 3* Schoen' question [14] could also be approached, instead of as in Problem 1, as follows:

Problem 2. *Let (M, g) a compact orientable Riemannian 3-manifolds with positive Ricci curvature and $\Sigma \subset M$ be a closed two-sided minimal surface of index one and genus 3. Hence Σ is a Heegaard splitting, Lawson [11], but can it be strongly irreducible?*

REFERENCES

- [1] M. I. Aroyo, J. M. Perez-Mato, D. Orobengoa, E. Tasci, G. de la Flor, A. Kirov, *Crystallography online: Bilbao Crystallographic Server*, Bulg. Chem. Commun. 43 (2) (2011), 183–197. <http://www.cryst.ehu.es>
- [2] O. Chodosh and C. Mantoulidis, *Minimal surfaces and the Allen-Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates*, Ann. of Math. 191 no. 1 (2020), 213–328, MR4045964, Zbl 1431.49045.
- [3] W. D. Dunbar, *Fibered Orbifolds and crystallographic groups*, PhD thesis, Advisor: W. P. Thurston, Princeton Univ, 1981, MR2631317.
- [4] N. Ejiri and T. Shoda, *The Morse index of a triply periodic minimal surface*, Differential Geom. Appl. 58 (2018), 177–201, MR3777753, Zbl 1388.53012.
- [5] P. M. Gruber, *Approximation of convex bodies*, Convexity and its applications, ed. P. M. Gruber and M. Wills, Birkhauser Verlag, (1983), 131–162, Zbl0519.52005.
- [6] M. Guaraco, *Min-max for phase transitions and the existence of embedded minimal hypersurfaces* J. Differential Geom. 108(1) (2018), 91–133, MR3743704, Zbl1387.49060.
- [7] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. 17 (2) (1982), 255–306, MR0664497, Zbl 0504.53034.

- [8] D. Ketover, Y. Liokumovich and A. Song, *On the existence of minimal Heegaard surfaces*, preprint, arXiv:1911.07161.
- [9] D. Ketover, F. C. Marques and A. Neves, *Catenoid estimate and its geometric applications*, J. Differential Geom. 115, no 1 (2020), 1–26, MR4081930, Zbl1439.53064.
- [10] T. Kobayashi and Y. Rieck, *Manifolds admitting both strongly irreducible and weakly reducible minimal genus Heegaard splittings*, Comm. Anal. Geom. 17 (2009), 637–649, MR2601347, Zbl 1439.53064.
- [11] H. B. Lawson, *The unknottedness of minimal embeddings*, Invent. Math. 11 (1970), 183–187, MR0287447, Zbl 0205.52002.
- [12] N. Lebedeva, V. Matveev, A. Petrunin and V. Shevchishin, *Smoothing 3-dimensional polyhedral spaces*, Electron. Res. Announc. Math. Sci. 22 (2015), 12–19, MR3361404, Zbl 1346.53040.
- [13] W. H. Meeks, *Lectures on Plateau’s problem*, IMPA, Rio de Janeiro, Brazil, 1978.
- [14] A. Neves, *New applications of Min-max Theory*, Proceedings of International Congress of Mathematics (2014), 939–957, MR3728646, Zbl 1314.00104.
- [15] J. Pitts, *Existence and Regularity of minimal surfaces on Riemannian manifolds*, Mathematical Notes 27, Princeton University Press, Princeton, 1981, MR0626027, Zbl 0462.58003.
- [16] M. Ritoré, *Index one minimal surfaces in flat three-space forms*, Indiana Univ. Math. J. 46 (1997), 1137–1153, MR1631568, Zbl 0922.53022.
- [17] M. Ritoré and A. Ros, *The spaces of index one minimal surfaces and stable constant mean curvature surfaces embedded in flat three manifolds*, Trans. Amer. Math. Soc. 348 (1996), 391–410, MR1322955, Zbl 0867.53007.
- [18] A. Ros, *One-sided complete stable minimal surfaces*, J. Differential Geom. 74, no. 1 (2006), 69–92, MR 2260928, Zbl 1110.53009.
- [19] A. Ros and E. Vergasta, *Stability for hypersurfaces of constant mean curvature with free boundary*, Geom. Dedicata, 56 (1995), 19–33, MR1338315, Zbl0912.53009.
- [20] M. Ross, *Schwarz’ P and D surfaces are stable*, Differential Geom. Appl. 2, no. 2 (1992), 179–195, MR1245555, Zbl 0747.53010.
- [21] F. Smith, *On the existence of embedded minimal two spheres in the three sphere, endowed with an arbitrary riemannian metric*. PhD thesis, Advisor: L. Simon, University of Melbourne, 1982.
- [22] C. Viana, *Index one minimal surfaces in spherical space forms*, Adv. Math. 360 (2020), 106900, 18 pp, MR4031123, Zbl 1430.53069.
- [23] W. Weil, *Ein Approximationssatz für konvexe Körper*, Manuscripta Math. 8 (1973) 335–362, MR0467623, Zbl 0251.52007.

Antonio Ros, aros@ugr.es,

Department of Geometry and Topology and Institute of Mathematics (IMAG),
University of Granada, 18071 Granada, Spain.