

Comment on “Path integral action of a particle with the generalized uncertainty principle and correspondence with noncommutativity”

DaeKil Park^{1,2*} and Eylee Jung¹,

¹*Department of Electronic Engineering,*

Kyungnam University, Changwon 631-701, Korea

²*Department of Physics, Kyungnam University, Changwon 631-701, Korea*

Abstract

Recently in [Phys. Rev. D 99 (2019) 104010] the non-relativistic Feynman propagator for harmonic oscillator system is presented when the generalized uncertainty principle is employed. In this short comment it is shown that the expression is incorrect. We also derive the correct expression of it.

* dkpark@kyungnam.ac.kr

Recently, S. Gangopadhyay and S. Bhattacharyya[1] have derived the non-relativistic Feynman propagators for free and harmonic oscillator systems when the generalized uncertainty principle (GUP) is employed and have discussed on their correspondence with noncommutativity. The GUP[2] they used can be summarized as a modified Heisenberg algebra

$$[Q_i, P_j] = i\hbar(\delta_{ij} + \beta\delta_{ij}P^2 + 2\beta P_i P_j), \quad (1)$$

where β is a GUP parameter, which has a dimension $(\text{momentum})^{-2}$. The modified Heisenberg algebra can be readily represented up to first order of β as $Q_i = q_i, P_i = p_i(1 + \beta p^2)$, where $\{p_i, q_i\}$ satisfies the usual Heisenberg algebra $[q_i, p_j] = i\hbar\delta_{ij}$.

The authors of Ref.[1] have considered the harmonic oscillator system, whose Hamiltonian is

$$\hat{H} = \frac{1}{2m}P^2 + \frac{1}{2}m\omega^2X^2 = \frac{p^2}{2m} + \frac{\beta}{m}p^4 + \frac{1}{2}m\omega^2x^2 + \mathcal{O}(\beta^2). \quad (2)$$

Without any explicit explanation they presented the following Feynman propagator (see Eq. (41) of Ref. [1]) of this system in a form:

$$\begin{aligned} & \langle q_f, t_f | q_0, t_0 \rangle \\ &= \sqrt{\frac{m\omega}{2\pi i\hbar \sin \omega T}} \left[1 + \frac{3i\beta\hbar m}{T} - 6\beta m^2 \left(\frac{q_f - q_0}{T} \right)^2 - \frac{3}{4}\beta m\hbar\omega^2 T \cot \omega T + \mathcal{O}(\beta^2) \right] e^{\frac{i}{\hbar}S_{cl}}, \end{aligned} \quad (3)$$

where $T = t_f - t_0$ and S_{cl} is a classical action. The only comment the authors presented is that the $\omega \rightarrow 0$ limit of Eq. (3) is

$$\begin{aligned} K_F[q_f, t_f : q_0, t_0] &= \sqrt{\frac{m}{2\pi i\hbar T}} \left(1 + \frac{3i\beta\hbar m}{T} - \frac{6\beta m^2(q_0 - q_f)^2}{T^2} + \mathcal{O}(\beta^2) \right) \\ &\quad \times \exp \left[\frac{im}{2\hbar T} (q_0 - q_f)^2 \left\{ 1 - 2\beta m^2 \left(\frac{q_0 - q_f}{T} \right)^2 \right\} \right], \end{aligned} \quad (4)$$

which is Feynman propagator for free particle. Since Eq. (3) is one of the main results of Ref. [1] and it can be used in other GUP-related issues, it is worthwhile to check the validity of Eq. (3) more carefully. Unfortunately, it is incorrect although it approaches to a correct $\omega \rightarrow 0$ limit. As we will show, this should be changed into

$$\langle q_f, t_f | q_0, t_0 \rangle = \sqrt{\frac{m\omega}{2\pi i\hbar \sin \omega T}} [1 + \beta f(q_0, q_f : T) + \mathcal{O}(\beta^2)] e^{\frac{i}{\hbar}(S_0 + \beta S_1)}, \quad (5)$$

where

$$\begin{aligned}
S_0 &= \frac{m\omega}{2 \sin \omega T} [(q_0^2 + q_f^2) \cos \omega T - 2q_0 q_f] \\
S_1 &= -\frac{m^3 \omega^3}{32 \sin^4 \omega T} \left[\{12\omega T + 8 \sin 2\omega T + \sin 4\omega T\} (q_0^4 + q_f^4) \right. \\
&\quad \left. - 4 \{12\omega T \cos \omega T + 11 \sin \omega T + 3 \sin 3\omega T\} q_0 q_f (q_0^2 + q_f^2) \right. \\
&\quad \left. + 12 \{4\omega T + 2\omega T \cos 2\omega T + 5 \sin 2\omega T\} q_0^2 q_f^2 \right] \\
f(q_0, q_f : T) &= \frac{3i\hbar m\omega}{8 \sin^2 \omega T} (2\omega T + 5 \sin \omega T \cos \omega T + \omega T \cos 2\omega T) \\
&\quad - \frac{3m^2 \omega^2}{8 \sin^3 \omega T} \left[2\omega T \{3 \cos \omega T (q_0^2 + q_f^2) - 2(2 + \cos 2\omega T) q_0 q_f\} \right. \\
&\quad \left. + 10 \sin \omega T (q_0^2 + q_f^2 - 2q_0 q_f \cos \omega T) - 6 \sin^3 \omega T (q_0^2 + q_f^2) \right].
\end{aligned} \tag{6}$$

Of course, $S_0 + \beta S_1$ is a classical action. It is straightforward to show that the $\omega \rightarrow 0$ limit of Eq. (5) also goes to $K_F[q_f, t_f : q_0, t_0]$.

In order to show Eq. (5) explicitly we note that the Feynman propagator $\langle q_f, t_f | q_0, t_0 \rangle$ can be derived from Schrödinger equation as

$$\langle q_f, t_f | q_0, t_0 \rangle = \sum_n \psi_n(q_f) \psi_n^*(q_0) e^{-(i/\hbar)E_n(t_f - t_0)}, \tag{7}$$

where $\psi_n(q)$ and E_n are n^{th} -order eigenfunction and eigenvalue of Schrödinger equation. The Schrödinger equation for the harmonic oscillator system is given by

$$\left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + \frac{\beta \hbar^4}{m} \frac{\partial^4}{\partial x^4} + \frac{1}{2} m \omega^2 x^2 + \mathcal{O}(\beta^2) \right] \psi_n(x) = E_n \psi_n(x). \tag{8}$$

If we treat the GUP term $\frac{\beta \hbar^4}{m} \frac{\partial^4}{\partial x^4}$ as small perturbation, one can derive $\psi_n(x)$ and E_n in a form:

$$\begin{aligned}
\psi_n(x) &= \phi_n(x) + (\beta m \hbar \omega) \left[\frac{(2n+3)\sqrt{(n+1)(n+2)}}{4} \phi_{n+2}(x) - \frac{(2n-1)\sqrt{n(n-1)}}{4} \phi_{n-2}(x) \right. \\
&\quad \left. + \frac{\sqrt{n(n-1)(n-2)(n-3)}}{16} \phi_{n-4}(x) - \frac{\sqrt{(n+1)(n+2)(n+3)(n+4)}}{16} \phi_{n+4}(x) \right] + \mathcal{O}(\beta^2) \\
E_n &= \left(n + \frac{1}{2} \right) \hbar \omega \left[1 + \frac{3(2n^2 + 2n + 1)}{2(2n + 1)} (\beta m \hbar \omega) \right] + \mathcal{O}(\beta^2),
\end{aligned} \tag{9}$$

where $n = 0, 1, 2, \dots$ and

$$\phi_n(x) = \frac{1}{\sqrt{2^n n!}} \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} H_n \left(\sqrt{\frac{m\omega}{\hbar}} x \right) \exp \left[-\frac{m\omega}{2\hbar} x^2 \right]. \quad (10)$$

In Eq. (10) $H_n(z)$ is a n^{th} -order Hermite polynomial. We assume $\phi_m(z) = 0$ for $m < 0$.

Inserting Eq. (9) into Eq. (7) one can express $\langle q_f, t_f | q_0, t_0 \rangle$ as

$$\langle q_f, t_f | q_0, t_0 \rangle = J + (\beta m \hbar \omega)(K_1 + K_2) + \mathcal{O}(\beta^2) \quad (11)$$

where

$$\begin{aligned} J &= \sum_{n=0}^{\infty} \phi_n(q_f) \phi_n(q_0) \exp \left[-\frac{i}{\hbar} \left(n + \frac{1}{2} \right) \hbar \omega T \left\{ 1 + \frac{3(2n^2 + 2n + 1)}{2(2n + 1)} (\beta m \hbar \omega) \right\} \right] \\ K_1 &= \left[\sum_{n=0}^{\infty} \frac{(2n+3)\sqrt{(n+1)(n+2)}}{4} [\phi_n(q_f) \phi_{n+2}(q_0) + \phi_n(q_0) \phi_{n+2}(q_f)] \right. \\ &\quad \left. - \sum_{n=2}^{\infty} \frac{(2n-1)\sqrt{n(n-1)}}{4} [\phi_n(q_f) \phi_{n-2}(q_0) + \phi_n(q_0) \phi_{n-2}(q_f)] \right] \exp \left[-\frac{i}{\hbar} \left(n + \frac{1}{2} \right) \hbar \omega T \right] \\ K_2 &= \left[\sum_{n=4}^{\infty} \frac{\sqrt{n(n-1)(n-2)(n-3)}}{16} [\phi_n(q_f) \phi_{n-4}(q_0) + \phi_n(q_0) \phi_{n-4}(q_f)] \right. \\ &\quad \left. - \sum_{n=0}^{\infty} \frac{\sqrt{(n+1)(n+2)(n+3)(n+4)}}{16} [\phi_n(q_f) \phi_{n+4}(q_0) + \phi_n(q_0) \phi_{n+4}(q_f)] \right] \times \exp \left[-\frac{i}{\hbar} \left(n + \frac{1}{2} \right) \hbar \omega T \right]. \end{aligned} \quad (12)$$

Using the extended Mehler's formula[3]

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{t^k}{k!} H_{k+m}(x) H_{k+n}(y) &= (1 - 4t^2)^{-(m+n+1)/2} \exp \left[\frac{4txy - 4t^2(x^2 + y^2)}{1 - 4t^2} \right] \\ &\quad \times \sum_{k=0}^{\min(m,n)} 2^{2k} k! \binom{m}{k} \binom{n}{k} t^k H_{m-k} \left(\frac{x - 2ty}{\sqrt{1 - 4t^2}} \right) H_{n-k} \left(\frac{y - 2tx}{\sqrt{1 - 4t^2}} \right), \end{aligned} \quad (13)$$

one can show

$$\begin{aligned} J &= \sqrt{\frac{m\omega}{\pi\hbar}} e^{-\frac{i}{2}\omega T} \left[\left(1 - \frac{3i}{4} (\beta m \hbar \omega^2 T) \right) + \frac{3i}{2} (\beta m \hbar \omega^2 T) \frac{\partial^2}{\partial \mu^2} + \frac{3}{2} (\beta m \hbar \omega^2 T) \frac{\partial}{\partial \mu} \right] F(\mu) \Big|_{\mu=\omega T} \\ K_1 &= -\frac{1}{2} \sqrt{\frac{m\omega}{2\pi i \hbar}} e^{-\frac{3}{2}i\omega T} \sin \omega T \left(2i \frac{\partial}{\partial \mu} + 3 \right) G(\mu) \Big|_{\mu=\omega T}, \end{aligned} \quad (14)$$

where

$$\begin{aligned} F(\mu) &= \frac{e^{\frac{i}{2}\mu}}{\sqrt{2i \sin \mu}} \exp \left[\frac{im\omega}{2\hbar \sin \mu} \{ (q_0^2 + q_f^2) \cos \mu - 2q_0 q_f \} \right] \\ G(\mu) &= \frac{\sqrt{2i} e^{i\mu}}{\sin \mu} \left[\frac{im\omega}{\hbar \sin \mu} \{ (q_0^2 + q_f^2) \cos \mu - 2q_0 q_f \} + 1 \right] F(\mu). \end{aligned} \quad (15)$$

Computing Eq. (14) explicitly, one can show

$$J = \sqrt{\frac{m\omega}{2\pi i\hbar \sin \omega T}} e^{\frac{i}{\hbar} S_0} \tilde{J} \quad K_1 = \sqrt{\frac{m\omega}{2\pi i\hbar \sin \omega T}} e^{\frac{i}{\hbar} S_0} \tilde{K}_1, \quad (16)$$

where

$$\begin{aligned} \tilde{J} &= 1 - \frac{3i\beta m\omega^2 T}{8\hbar \sin^4 \omega T} \left[-3i\hbar m\omega(q_0^2 + q_f^2) \sin 2\omega T + m^2\omega^2(q_0^2 + q_f^2 - 2q_0q_f \cos \omega T)^2 \right. \\ &\quad \left. + 4i\hbar m\omega \sin \omega T (2 + \cos 2\omega T) q_0q_f - \hbar^2 \sin^2 \omega T (2 + \cos 2\omega T) \right] \\ \tilde{K}_1 &= -\frac{i}{8\hbar^2 \sin^3 \omega T} \left[-4m^2\omega^2 q_0q_f (q_0^2 + q_f^2)(3 + \cos 2\omega T) + 3\hbar^2(\cos 3\omega T - \cos \omega T) \right. \\ &\quad \left. + 4m\omega \cos \omega T \{m\omega(q_0^4 + 6q_0^2q_f^2 + q_f^4) + 12i\hbar q_0q_f \sin \omega T\} - 3i\hbar m\omega(q_0^2 + q_f^2)(5 \sin \omega T + \sin 3\omega T) \right]. \end{aligned} \quad (17)$$

Using Eq. (13) again and $H_4(z) = 16z^4 - 48z^2 + 12$, one can show again $K_2 = \sqrt{\frac{m\omega}{2\pi i\hbar \sin \omega T}} e^{\frac{i}{\hbar} S_0} \tilde{K}_2$, where

$$\begin{aligned} \tilde{K}_2 &= -\frac{i \cos \omega T}{16\hbar^2 \sin^3 \omega T} \left[12m^2\omega^2 q_0^2 q_f^2 - 3\hbar^2(1 - \cos 2\omega T) + 2m\omega \left\{ m\omega \cos 2\omega T (q_0^4 + q_f^4) \right. \right. \\ &\quad \left. \left. - 4m\omega q_0q_f (q_0^2 + q_f^2) \cos \omega T - 6i\hbar \sin \omega T \{(q_0^2 + q_f^2) \cos \omega T - 2q_0q_f\} \right\} \right]. \end{aligned}$$

Inserting J , K_1 , and K_2 into Eq. (11), it is possible to show that the Feynman propagator becomes Eq. (5).

- [1] S. Gangopadhyay and S. Bhattacharyya, *Path integral action of a particle with the generalized uncertainty principle and correspondence with noncommutativity*, Phys. Rev. **D 99** (2019) 104010 [arXiv:1901.03411 (quant-ph)].
- [2] A. Kempf, G. Mangano, and R. B. Mann, *Hilbert Space Representation of the Minimal Length Uncertainty Relation*, Phys. Rev. **D 52** (1995) 1108 [hep-th/9412167].
- [3] A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev, *Integrals and Series vol. 2*, (Gordon and Breach Science Publishers, 1983, New York).