arXiv:2002.09379v1 [math.AG] 21 Feb 2020

1

A Note on Dieudonné Theory over Perfectoid Rings

Timo Henkel¹

Abstract. For a perfectoid ring R and a natural number n we investigate the essential image of the category of truncated by n Barsotti-Tate groups under the anti-equivalence between commutative, finite, locally free, R-group schemes of p-power order and torsion Breuil-Kisin-Fargues modules over R. We describe the associated semi-liner algebra data and show as a consequence that every BT_n -group over R is the p^n -torsion of some BT-group.

Introduction

For k being a perfect field of characteristic p, classical Dieudonné theory provides a semi-linear algebra description of truncated by n Barsotti-Tate groups. This can be used to show that every such group arises as the p^n torsion subgroup of some Barsotti-Tate group. Lau extended these results to perfect rings of characteristic p in [5], reproving unpublished results of Gabber by a different method. Recently, in [1], Anschütz and Le Bras generalized the existing instances of Dieudonné theory to so called Prismatic Dieudonné Theory, including results on Dieudonné theory for perfectoid rings, which were partially proven by Lau before in [6]. In this note, we use the results of [1] to classify BT_n -groups over perfectoid rings in terms of semi-linear algebra and investigate the consequences for lifting properties of BT_n -groups to BT-groups in this setting.

The semi-linear algebra objects associated with BT_n -groups are the following: A BK_n-module over a perfectoid ring $R = W(S)/\xi$ is a triple (M, φ, ψ) , where M is a finite projective $W_n(S)$ -module, $\varphi \colon M^\sigma \to M$ and $\psi \colon M \to M^\sigma$ are $W_n(S)$ -linear maps such that $\varphi \circ \psi = \xi$ and $\psi \circ \varphi = \xi$. In the case n = 1 one additionally demands that $\operatorname{coker}(\varphi)$ is finite projective as an S/ξ -module and that the induced sequence $M/\xi \xrightarrow{\overline{\psi}} M^\sigma/\xi \xrightarrow{\overline{\varphi}} M/\xi$ is exact. The category of BK_n-modules over R, denoted by BK_n(R), also admits a fibrewise characterization: A torsion Breuil-Kisin-Fargues module over Ris a BK_n-module if and only if it is killed by p^n and all fibres are truncated Dieudonné modules of level n. Combining this fact with the fibrewise description of BT_n-groups and the properties of the functor of [1], which classifies commutative finite locally free p-groups, we show the following result:

¹Department of Mathematics, Technische Universität Darmstadt, Schloßgartenstr. 7, 64289 Darmstadt, Germany, thenkel@mathematik.tu-darmstadt.de

Theorem 1. Let R be a perfectoid ring. For all $n \in \mathbb{N}$ there is an antiequivalence of categories

$$\operatorname{BT}_{n}(R) \cong \operatorname{BK}_{n}(R).$$

We apply lifting arguments to obtain the following corollary:

Theorem 2. Let R be a perfectoid ring and $n \in \mathbb{N}$. The truncation functor

$$\operatorname{BT}(R) \to \operatorname{BT}_{\operatorname{n}}(R)$$

is essentially surjective, i.e. every BT_n -group over R arises as the p^n -torsion subgroup of some Barsotti-Tate group over R.

In the first chapter we recall the definition and general facts about perfectoid rings. Over such a ring R we briefly recall the relevant categorical anti-equivalence \mathbb{M} of [1] which is the starting point of our investigation (cf. 1.9). In the next chapter we turn attention to the category of BT₁-groups and BK₁-modules. After proofing the fibrewise characterization (cf. 2.14) we establish Theorem 1 for n = 1 (cf. 2.15). In the last chapter, we explain analogous results for BT_n-groups (cf. 3.8) and finally show that, with this description at hand, we can use lifting of normal representations to proof Theorem 2 (cf 3.14).

Acknowledgements: The author thanks E. Lau and T. Wedhorn for help-ful discussions.

Contents

1	Prerequisits		3
	1.1	Generalities on perfectoid rings	3
	1.2	Short review of classification results	4
2	BT_1 -groups and Dieudonné spaces		6
	2.1	Truncated by 1 Barsotti-Tate groups	6
	2.2	Modules associated with BT_1 -groups $\ldots \ldots \ldots \ldots \ldots$	8
3	Generalization to BT_n and lifting properties		
	3.1	Modules associated with BT_n -groups $\ldots \ldots \ldots \ldots \ldots \ldots$	11
	3.2	Normal representations for BK_n -modules $\ldots \ldots \ldots \ldots \ldots$	13

Notations and assumptions:

- · Once and for all we fix some prime number p.
- \cdot All rings are commutative with 1.
- For a ring R with pR = 0 we denote by W(R) the ring of p-typical Witt vectors of R and for $n \in \mathbb{N}$ by $W_n(R)$ the ring of p-typical Witt vectors truncated by n.
- For a ring R with pR = 0 we denote the Frobenius morphism $x \mapsto x^p$ by σ . By abuse of notation we also denote the induced Frobenius morphism on W(S) and on $W_n(S)$ by σ .

1 Prerequisits

We recall some general facts about perfectoid rings and briefly state the antiequivalences obtained in [6] (§9 and §10), [1] (§ 4 and §5.1) and [7] (Theorem 17.5.2) which characterize Barsotti-Tate groups and *p*-groups over perfectoid rings in terms of semilinear algebra.

1.1 Generalities on perfectoid rings

We use the definition of perfectoid rings as quotients of perfect prisms as used in [6].

- **Definition 1.1.** 1. A ring R with pR = 0 is called *perfect* if the Frobenius morphism $\sigma: R \to R, x \mapsto x^p$ is bijective.
 - 2. For a perfect ring S an element $\xi = (\xi_0, \xi_1, ...) \in W(S)$ is called *distin*guished if $\xi_1 \in S^{\times}$ and S is ξ_0 -adically complete.
 - 3. A ring R is called *perfectoid* if there is an isomorphism $R \cong W(S)/\xi$, where S is a perfect ring and $\xi \in S$ is a distinguished element.
- **Remark 1.2.** 1. A ring R is perfected if and only if it is π -adically complete for some element $\pi \in R$ such that π^p divides p, the Frobenius map $\varphi \colon R/p \to R/p$ is surjective and the kernel of $\theta \colon W(R^{\flat}) \to R$ is principal, say generated by an element ξ (here θ denotes the unique lift of the natural surjection $R^{\flat} := \lim_{\epsilon \to \sigma} R/p \to R/p$). In this case we obtain an isomorphism $W(R^{\flat})/\xi \to R$ which is induced by θ and whenever we have a presentation of a perfectoid ring $R = W(S)/\xi$ we may identify S with R^{\flat} (cf. [6] Remark 8.6).

- 2. The notion of a perfectoid ring defined above is sometimes called *integral perfectoid ring* to distinguish from the notion of a *perfectoid Tate Huber ring*, which plays the central role in the theory of perfectoid spaces. However, given a perfectoid Tate Huber pair (R, R^+) , the ring of integral elements R^+ is a perfectoid ring in the sense defined above.
- 3. For a perfectoid ring $R = W(S)/\xi$ we have $W_n(S)/\xi = R/p^n$ for all $n \in \mathbb{N}$ and the elements p and $\xi \in W(S)$ are nonzerodivisors. The latter implies that R is p-torsion free if and only if $W_n(S)$ is ξ -torsion free for all $n \in \mathbb{N}$. Indeed, the W(S)-module

$$\{(x,y) \in W(S)^2 \mid \xi x = p^n y\} / \{(p^n z, \xi z) \mid z \in W(S)\}$$

is isomorphic to both $\{x \in W_n(S) \mid \xi x = 0\}$ and $\{y \in R \mid p^n y = 0\}$ via $(x, y) \mapsto x$ and $(x, y) \mapsto y$ respectively.

- 4. Every perfectoid ring $R = W(S)/\xi$ is *p*-adically complete. In particular, $Max(R) = Max(R/p) = Max(S/\xi_0) = Max(S)$.
- **Example 1.3.** 1. Every perfect ring S may be written as W(S)/p and is hence a perfectoid ring. In particular, every perfect field of characteristic p is a perfectoid ring.
 - 2. Let *C* be an algebraically closed complete non-archimedean extension of \mathbb{Q}_p with ring of integers \mathcal{O}_C . Choosing a compatible system of *p*power roots of *p* gives an element p^{\flat} of \mathcal{O}_C^{\flat} such that \mathcal{O}_C^{\flat} is p^{\flat} -adically complete. We obtain an isomorphism $W(\mathcal{O}_C^{\flat})/([p^{\flat}] - p) \cong \mathcal{O}_C$ which depicts \mathcal{O}_C as a perfectoid ring.

1.2 Short review of classification results

Let $R = W(S)/\xi$ be a perfectoid ring. We denote the category of *p*-divisible groups over R by BT(R).

- **Definition 1.4.** 1. A (minuscule) Breuil-Kisin-Fargues module for R is a triple (M, φ, ψ) , where M is a finite projective W(S)-module and $\varphi: M^{\sigma} \to M$ and $\psi: M \to M^{\sigma}$ are W(S)-linear maps such that $\psi \circ \varphi = \xi$ and $\varphi \circ \psi = \xi$.
 - 2. We denote the category of Breuil-Kisin-Fargues modules for R by BK(R) (the morphisms are W(S)-linear maps which are compatible with the φ 's; the compatibility with ψ 's is then automatic as explained below).

Note that since $\xi \in W(S)$ is a nonzerodivisor, the maps φ and ψ are injective and hence ψ is determined by φ (and vice versa) via the formula $\psi(x) = \varphi^{-1}(\xi x)$. We only add the datum to draw the connection to other semi linear algebra data as defined below.

Theorem 1.5. There is an exact, compatible with base change, anti-equivalence

$$BT(R) \cong BK(R),$$
$$G \mapsto \mathbb{M}(G),$$

admitting an exact, compatible with base change, inverse.

Proof. [7] Theorem 17.5.2. resp. [1] Corollary 4.3.8.

Definition 1.6. Let $BT_{tor}(R)$ denote the category of commutative, finite, locally free *R*-group schemes which are killed by a power of *p*.

Note that the category $BT_{tor}(R)$ is denoted by pGR(Spec R) in [6]. We justify our notation by Theorem 1.9 and Remark 1.10 below.

The following definition is taken from [6] Definition 10.5. Such objects are introduced as *torsion prismatic Dieudonné modules* in [1] Definition 5.1.1.

- **Definition 1.7.** 1. A torsion Breuil-Kisin-Fargues module for R is a triple $M = (M, \varphi, \psi)$, where M is a finitely presented W(S)-module of projective dimension ≤ 1 , killed by a power of p, and $M^{\sigma} \xrightarrow{\varphi} M, M \xrightarrow{\psi} M^{\sigma}$ are W(S)-linear maps such that $\varphi \circ \psi = \xi$ and $\psi \circ \varphi = \xi$.
 - 2. We denote the category of torsion Breuil-Kisin-Fargues modules for R by $BK_{tor}(R)$ (the morphisms are W(S)-linear maps which are compatible with the φ 's and the ψ 's).

Remark 1.8. If R is p-torsion free (equivalently $W_n(S)$ is ξ -torsion free for all $n \in \mathbb{N}$) for any object $M = (M, \varphi, \psi)$ of $BK_{tor}(R)$ the module M is ξ torsion free and hence φ and ψ are injective. So φ determines $\psi \colon M \to M^{\sigma}$ via the formula $\psi(x) = \varphi^{-1}(\xi x)$ (and vice versa). Indeed, assume $p^n M = 0$ then M fits into a short exact sequence

$$0 \to P_1 \to P_0 \to M \to 0,$$

where P_1 and P_0 are finite projective W(S)-modules s.t. $p^n P_0 \subseteq P_1$. We obtain an injection

$$p^n P_0/p^n P_1 \rightarrow P_1/p^n P_1$$

of $W_n(S)$ -modules, where the latter is projective and hence $M = p^n P_0/p^n P_1$ is torsion free. But $\xi \in W_n(S)$ is a nonzerodivisor by assumption.

By abuse of notation, we use the same symbol \mathbb{M} for the anti-equivalences in Theorem 1.5 and Theorem 1.9 below.

Theorem 1.9. There is an exact, compatible with base change, anti-equivalence

$$BT_{tor}(R) \cong BK_{tor}(R),$$
$$G \mapsto \mathbb{M}(G),$$

admitting an exact, compatible with base change, inverse.

Proof. [1] Theorem 5.1.4.

- **Remark 1.10.** 1. Zariski-locally on $\operatorname{Spec}(R)$ every $G \in \operatorname{BT}_{\operatorname{tor}}(R)$ can be written as the kernel of an isogeny of *p*-divisible groups over R, so Zariski-locally, $G = \ker(H_1 \to H_2)$ (cf. [2] Theorem 3.1.1). The functor of the theorem above can Zariski-locally on R be defined as $\mathbb{M}(G) :=$ $\operatorname{coker}(\mathbb{M}(H_1) \to \mathbb{M}(H_2))$ using the anti-equivalence of Theorem 1.5 (cf. proof of [6] Theorem 10.12 and proof of [1] Theorem 5.1.4).
 - By [1] Proposition 4.3.2 and the explicit description of M given in Theorem 5.1.4, the anti-equivalence M extends crystalline Dieudonné Theory of [2].

2 BT₁-groups and Dieudonné spaces

The goal of this section is to determine the essential image of the full subcategory $BT_1(R)$ of $BT_{tor}(R)$, defined below, for a perfectoid ring R under the anti-equivalence $BT_{tor}(R) \cong BK_{tor}(R)$. Fix a perfectoid ring $R = W(S)/\xi$.

2.1 Truncated by 1 Barsotti-Tate groups

Definition 2.1. Let $BT_1(R)$ denote the full subcategory of $BT_{tor}(R)$ consisting of objects G such that G is annihilated by p and for $G_{\circ} := G \otimes_R R/p$ the sequence

$$G_{\circ} \xrightarrow{F} G_{\circ}^{(p)} \xrightarrow{V} G_{\circ}$$

is exact (*F* resp. *V* denote the Frobenius resp. Verschiebung of G_{\circ}). Objects of $BT_1(R)$ are called *truncated by* 1 *Barsotti-Tate groups over* R or BT_1 -*groups over* R.

Definition 2.2. Let G = Spec(A) be a BT₁-group over R.

- 1. The locally constant function h: $\operatorname{Spec}(R/p) \to \mathbb{N}_0$ which is uniquely determined by $s \mapsto \log_p(\dim_{\kappa(s)}(A \otimes_R \kappa(s)))$, for $s \in \operatorname{Max}(R/p)$ with residue field $\kappa(s)$, is said to be the *height of G*.
- 2. The locally constant function dim: $\operatorname{Spec}(R/p) \to \mathbb{N}_0$ which is uniquely determined by $s \mapsto \dim_{\kappa(s)}(\operatorname{Lie}(G \otimes_R \kappa(s)))$, for $s \in \operatorname{Max}(R/p)$ with residue field $\kappa(s)$, is said to be the *dimension of* G.

Example 2.3. Let G be a p-divisible group over R of constant height h and dimension d, and let G[p] denote its group of p-torsion. Then G[p] is a BT_1 -group of height h and dimension d. As a corollary of the lifting results of §3, we will see that every object of $BT_1(R)$ is of this form.

Remark 2.4. 1. Let G be some BT_1 -group over R with $M = \operatorname{M}(G)$ the associated $\operatorname{BK}_{\operatorname{tor}}$ -module under the anti-equivalence of Theorem 1.9. Since M commutes with base change and extends the Dieudonné Theory of [2], given some point $s \in \operatorname{Max}(S)$ with residue field κ (which is perfect and of characteristic p), the base change $M_s = (M_s := M \otimes_S \kappa, \varphi_s := \varphi \otimes \operatorname{id}_{\kappa}, \psi_s := \psi \otimes \operatorname{id}_{\kappa})$ of M to κ is a Dieudonné space over κ , i.e. M_s is a finite dimensional κ -vector space such that $\psi_s \circ \varphi_s = 0$, $\varphi_s \circ \psi_s = 0$ and the sequence

$$M_s \xrightarrow{\psi_s} M_s^{\sigma} \xrightarrow{\varphi_s} M_s$$

is exact (note that this implies that $M_s^{\sigma} \xrightarrow{\varphi_s} M_s \xrightarrow{\psi_s} M_s^{\sigma}$ is exact as well).

- 2. The *height* of a Dieudonné space (M, φ, ψ) over a perfect field k of characteristic p is defined to be the k-dimension of M. The k-dimension of coker (φ) is called *dimension of* M.
- 3. For a perfect field k of characteristic p we denote the category of Dieudonné spaces over k by $BK_1(k)$. The functor $BT_1(k) \to BK_1(k)$ which is induced by M is a height and dimension preserving antiequivalence, since the Dieudonné Theory of [2] §3 provides this result and M extends this theory.

The following lemma gives a fibrewise characterization of BT_1 -groups. This will lead to a description of the essential image of $BT_1(R)$ under the anti-equivalence of Theorem 1.9.

Lemma 2.5. Let $G \in BT_{tor}(R)$ such that G is killed by p. Then $G \in BT_1(R)$ if and only if $G \otimes_R \kappa(s) \in BT_1(\kappa(s))$ for all $s \in Max(R/p)$. *Proof.* Noting that Max(R) = Max(R/p) this is a corollary of the following general assertion on finite locally free group schemes.

Lemma 2.6. Let X be an affine scheme. A sequence of finite locally free group schemes

$$G' \xrightarrow{\varphi} G \xrightarrow{\psi} G''$$

over X is exact if and only if $\psi \circ \varphi = 0$ and for every closed point $x \in X$ the sequence $G'_x \to G_x \to G''_x$ is exact.

Proof. [4] Proposition 1.1.

2.2 Modules associated with BT₁-groups

The fibrewise characterization of $\operatorname{BT}_1(R)$ hints that its essential image consists of such $M \in \operatorname{BK}_{\operatorname{tor}}(R)$ which satisfy pM = 0 and whose fibres are Dieudonné spaces for all closed points of $\operatorname{Spec}(S)$. The goal of this section is to show that this essential image is also given via the following global definition.

Definition 2.7. Let $BK_1(R)$ be the full subcategory of $BK_{tor}(R)$ whose objects are given by triples $M = (M, \varphi, \psi)$, such that

- (a) M is killed by p (and hence is a finite projective S-module by Lemma 2.8),
- (b) $\operatorname{coker}(\varphi)$ is finite projective as an S/ξ_0 -module,
- (c) the induced sequence

$$M/\xi_0 \xrightarrow{\overline{\psi}} M^\sigma/\xi_0 \xrightarrow{\overline{\varphi}} M/\xi_0$$

is exact.

Objects of $BK_1(R)$ are also called BK_1 -modules over R. The lemma below ensures that the underlying S-module of a BK_1 -module is in fact finite projective as indicated in (a):

Lemma 2.8. Let M be a finite W(S)-module which is killed by p. Then M is of projective dimension 1 as a W(S)-module if and only if it is finite projective as an S-module.

Proof. [3] Lemma 7.8.

- **Remark 2.9.** 1. If R = k is a perfect field of characteristic p, we can write k = W(k)/p. Hence S = k and $\xi_0 = 0$ and in this case and we recover the definition of Dieudonné spaces over k.
 - 2. More generally, if R = S is a perfect ring, a BK₁-module is the same as a *truncated Dieudonné module of level* 1 *over* R as defined in [5] Definition 6.1.
 - 3. If R is p-torsion free, an object of BK_{tor} which is killed by p automatically satisfies property (c) of the definition above.
- **Example 2.10.** 1. Let $\mathcal{O}_C = W(\mathcal{O}_C^{\flat})/([p^{\flat}] p)$ be the ring of integers of some algebraically closed complete non archimedean extension of \mathbb{Q}_p with tilt \mathcal{O}_C^{\flat} . In this case we have $\xi_0 = p^{\flat}$ and a BK_{tor}-module over \mathcal{O}_C which is killed by p is uniquely determined by a pair (M, φ) , where M is a finite free \mathcal{O}_C^{\flat} -module and $\varphi \colon M^{\sigma} \to M$ is a linear map whose cokernel is killed by p^{\flat} .
 - 2. For every $a \in \mathcal{O}_{C}^{\flat}$ such that $|p^{\flat}| \leq |a| \leq 1$ the cokernel of the linear map $(\mathcal{O}_{C}^{\flat})^{\sigma} \to \mathcal{O}_{C}^{\flat}, x \mapsto a\sigma^{-1}(x)$ is killed by p^{\flat} . However, its cokernel is finite projective as a $\mathcal{O}_{C}^{\flat}/p^{\flat}\mathcal{O}_{C}^{\flat}$ -module if and only if it is isomorphic to $(\mathcal{O}_{C}^{\flat}/p^{\flat}\mathcal{O}_{C}^{\flat})^{n}$ as an $\mathcal{O}_{C}^{\flat}/p^{\flat}\mathcal{O}_{C}^{\flat}$ -module for some $n \in \mathbb{N}$ because $\mathcal{O}_{C}^{\flat}/p^{\flat}\mathcal{O}_{C}^{\flat}$ is local. But this is satisfied if and only if $|a| \in \{|p^{\flat}|, 1\}$ since multiplication with a is not equal to 0 on $(\mathcal{O}_{C}^{\flat}/p^{\flat}\mathcal{O}_{C}^{\flat})^{n}$ if $|a| > |p^{\flat}|$. Hence, up to isomorphism, we only obtain two different BK₁-modules in this case. One associated with the BT₁-group $\mathbb{Z}/p\mathbb{Z}_{\mathcal{O}_{C}}$ and one associated with $\mu_{p,\mathcal{O}_{C}}$. This observation matches our expectation that every BK₁-module over \mathcal{O}_{C} can be lifted to some BK-module over \mathcal{O}_{C} . Those are by the anti-equivalence of Theorem 1.5 given by the BT-groups $\mathbb{Q}_{p}/\mathbb{Z}_{p}$ and $\mu_{p^{\infty}}$.

We also have a notion of height and dimension for BK₁-modules:

Definition 2.11. Let $M = (M, \varphi, \psi) \in BK_1(R)$.

- 1. The locally constant function h: $\operatorname{Spec}(S/\xi_0) \to \mathbb{Z}$ uniquely determined by $s \mapsto \dim_{\kappa(s)}(M \otimes_S \kappa(s))$, for $s \in \operatorname{Max}(S/\xi_0)$ with residue field $\kappa(s)$, is said to be the *height of* M.
- 2. The locally constant function dim: $\operatorname{Spec}(S/\xi_0) \to \mathbb{Z}$ uniquely determined by $s \mapsto \dim_{\kappa(s)}(\operatorname{coker}(\varphi) \otimes_{S/\xi_0} \kappa(s))$, for $s \in \operatorname{Max}(S/\xi_0)$ with residue field $\kappa(s)$, is said to be the *dimension of* M.

The following lemma is the analogue of Lemma 2.6 providing a fibrewise characterization of $BT_1(R)$:

Lemma 2.12. Let A be a ring and

$$M^{\bullet} = (M' \xrightarrow{\varphi} M \xrightarrow{\psi} M' \xrightarrow{\varphi} M)$$

be a sequence of finite projective A-modules such that $\varphi \circ \psi = 0$ and $\psi \circ \varphi = 0$. The following assertions are equivalent:

- 1. $M^{\bullet} \otimes_A \kappa(x)$ is exact for all $x \in Max(A)$,
- 2. M^{\bullet} is exact and coker(φ) is a finite projective A-module.
- 3. $M \xrightarrow{\psi} M' \xrightarrow{\varphi} M$ is exact and $\operatorname{coker}(\varphi)$ is a finite projective A-module.

In this case $\operatorname{im}(\psi)$ and $\operatorname{ker}(\varphi)$ are direct summands of M', $\operatorname{im}(\varphi)$ and $\operatorname{ker}(\psi)$ are direct summands of M and $\operatorname{coker}(\psi)$ is a finite projective A-modules as well.

Proof. Obviously, 2. implies 3. If $M \xrightarrow{\psi} M' \xrightarrow{\varphi} M$ is exact and $\operatorname{coker}(\varphi)$ is finite projective, $\operatorname{im}(\varphi)$, $\operatorname{ker}(\varphi)$, $\operatorname{coker}(\psi)$, $\operatorname{im}(\psi)$ and $\operatorname{ker}(\psi)$ satisfy the additional assertion of the lemma. Hence 1. follows. So lets assume 1. and consider the sequence

$$0 \to \operatorname{im}(\varphi) \to M \to \operatorname{coker}(\varphi) \to 0.$$

To show that $\operatorname{coker}(\varphi)$ is a projective *A*-module, we establish that $\operatorname{im}(\varphi) \otimes_A \kappa(x) \to M \otimes_A \kappa(x)$ is injective for each $x \in \operatorname{Max}(A)$. By assumption, we have an exact sequence of $\kappa(x)$ -vector spaces

$$M \otimes_A \kappa(x) \xrightarrow{\psi_x} M' \otimes_A \kappa(x) \xrightarrow{\varphi_x} M \otimes_A \kappa(x).$$

Here φ_x factors as

$$M' \otimes_A \kappa(x) \xrightarrow{\pi} \operatorname{im}(\varphi) \otimes_A \kappa(x) \to M \otimes_A \kappa(x).$$

Now let $s \in \operatorname{im}(\varphi) \otimes_A \kappa(x)$ which is mapped to 0. By surjectivity of π we can choose $y \in M' \otimes_A \kappa(x)$ with $\pi(y) = s$. Then $\varphi_x(y) = 0$, hence we find $z \in M \otimes_A \kappa(x)$ with $\psi_x(z) = y$. Therefore $s = \pi(\psi_x(z)) = 0$ as desired. Note that we can do the same calculations for ψ instead of φ to obtain projectivity of $\operatorname{coker}(\psi)$. Consequently, $\operatorname{im}(\psi)$, $\operatorname{ker}(\psi)$, $\operatorname{im}(\varphi)$ and $\operatorname{ker}(\varphi)$ satisfy the additional assertion of the lemma. The exactness of M^{\bullet} then follows by applying Nakayama's lemma.

11

Remark 2.13. Let $R = W(S)/\xi \to R' = W(S')/\xi'$ be a morphism of perfectoid rings. Then the image of ξ under the induced morphism $W(S) \to W(S')$ is a distinguished element of W(S') and we can identify $R' = W(S')/\xi$. By the previous lemma we obtain a base change functor $BT_1(R) \to BT_1(R')$.

Proposition 2.14. Let $M = (M, \varphi, \psi) \in BK_{tor}(R)$ such that pM = 0. Then $M \in BK_1(R)$ if and only if $M \otimes_S \kappa(s) \in BK_1(\kappa(s))$ for all $s \in Max(S/\xi_0)$. Moreover in this case the sequence

$$M^{\sigma}/\xi_0 \xrightarrow{\overline{\varphi}} M/\xi_0 \xrightarrow{\overline{\psi}} M^{\sigma}/\xi_0$$

is exact as well.

Proof. This follows from Lemma 2.12.

Proposition 2.15. The anti-equivalence \mathbb{M} of Theorem 1.9 induces a height and dimension preserving anti-equivalence

$$\operatorname{BT}_1(R) \cong \operatorname{BK}_1(R).$$

Proof. Let G be a BT_{tor} -group over R and $M = \operatorname{M}(G)$ be the associated BK_{tor} -module over R. Since M is exact and has an exact inverse, G is annihilated by p if and only if M is annihilated by p. Since M and an inverse are compatible with base change and induce height and dimension preserving anti-equivalences $\operatorname{BT}_1(k) \cong \operatorname{BK}_1(k)$ for every perfect field k, the proposition follows from Lemma 2.5 and Lemma 2.14.

3 Generalization to BT_n and lifting properties

We still fix a perfectoid ring $R = W(S)/\xi$. Similar to the previous section, we determine the essential image of BT_n -groups under the anti-equivalence \mathbb{M} of Theorem 1.9. At the end we apply a lifting argument to the obtained semilinear algebra data to show that over R every BT_n -group arises as the p^n -torsion of some BT-group.

3.1 Modules associated with BT_n -groups

Definition 3.1. For $n \geq 2$ we denote by $\operatorname{BT}_n(R)$ the full subcategory of $\operatorname{BT}_{\operatorname{tor}}(R)$ whose objects are those groups G which are annihilated by p^n and such that the sequence

$$G \xrightarrow{\cdot p^{n-1}} G \xrightarrow{\cdot p} G$$

is exact.

As in the case for n = 1, we have a fibrewise description of the category $BK_n(R)$:

Lemma 3.2. Let $G \in BT_{tor}(R)$ and $n \in \mathbb{N}$ such that G is annihilated by p^n . Then G is an object of $BT_n(R)$ if and only if $G \otimes_R \kappa(s)$ is an object of $BT_n(\kappa(s))$ for all $s \in Max(R/p)$.

Proof. Note that Max(R) = Max(R/p) and use Lemma 2.6.

Remark 3.3. Let $n \ge m$ be two natural numbers and $G \in BT_n(R)$. The p^m -torsion subgroup $G[p^m]$ of G is an object of $BT_m(R)$.

Now we define the full subcategory of $BK_{tor}(R)$ which will be shown to correspond to $BT_n(R)$ under the anti-equivalence of Theorem 1.9.

Definition 3.4. For $n \geq 2$ we denote by $BK_n(R)$ the full subcategory of $BK_{tor}(R)$ consisting of objects $M = (M, \varphi, \psi)$ such that M is killed by p^n and finite projective as a $W_n(S)$ -module.

Objects of $BK_n(R)$ are also called BK_n -modules over R and if R = S is perfect, those are the same as *truncated Dieudonné modules of rank n over* R as defined in [5] Definition 6.1.

If R = S is perfect, so in particular if R is a perfect field of characteristic p, we have the following classification result:

Proposition 3.5. Assume that R is perfect. Then the anti-equivalence \mathbb{M} : $\mathrm{BT}_{\mathrm{tor}}(R) \to \mathrm{BK}_{\mathrm{tor}}(R)$ of Theorem 1.9 induces an anti-equivalence

$$BT_n(R) \cong BK_n(R)$$

for all $n \geq 1$.

Proof. Since \mathbb{M} extends the Dieudonné theory of [2], this is the content of [5] Theorem 6.4.

Remark 3.6. Let $R = W(S)/\xi \to R' = W(S')/\xi$ be a morphism of perfectoid rings. We obtain a base change functor $BK_n(R) \to BK_n(R')$.

As in the n = 1 case, we also have the following fibrewise description:

Lemma 3.7. Let $M = (M, \varphi, \psi) \in BK_{tor}(R)$ such that $p^n M = 0$. Then $M \in BK_n(R)$ if and only if $M \otimes_{W_n(S)} W_n(\kappa(s)) \in BK_n(\kappa(s))$ for all $s \in Max(R/p)$.

3.2 Normal representations for BK_n-modules

Proof. One direction is clear. Conversely, assume that all fibres of $M = (M, \varphi, \psi)$ are BK_n -modules. We have to show that M is a finite projective $W_n(S)$ -module. The assumption $M \otimes_{W_n(S)} W_n(\kappa(s)) \in BT_n(\kappa(s))$ implies that $M/p \otimes_S \kappa(s) = (M \otimes_{W_n(S)} W_n(\kappa(s)))/p$ is an object of $BT_1(\kappa(s))$ for all $s \in Max(R/p)$ by [5] Lemma 6.2. This shows that $M/p \otimes_S \kappa(s)$ is an object of $BK_1(\kappa(s))$. Consequently, $M/p \in BK_1(R)$ by Lemma 2.14 and M/p is a finite projective S-module. By the local flatness criterion we are reduced to show that

$$\operatorname{Tor}_{1}^{W_{n}(S)}(S,M) = 0.$$

This is the case if and only if the sequence

$$M \xrightarrow{\cdot p^{n-1}} M \xrightarrow{\cdot p} M$$

is exact. But since the anti-equivalence \mathbb{M} (of Theorem 1.9) and an inverse are exact and commute with base change, this is equivalent to

$$M \otimes_{W_n(S)} W_n(\kappa(s)) \xrightarrow{:p^{n-1}} M \otimes_{W_n(S)} W_n(\kappa(s)) \xrightarrow{:p} M \otimes_{W_n(S)} W_n(\kappa(s))$$

being exact for all $s \in Max(W_n(S))$ by Lemma 2.6. This is true because $M \otimes_{W_n(S)} W_n(\kappa(s))$ is a finite projective $W_n(\kappa(s))$ -module by assumption. \Box

The fibrewise description then leads to

Proposition 3.8. The anti-equivalence \mathbb{M} : $\mathrm{BT}_{\mathrm{tor}}(R) \cong \mathrm{BK}_{\mathrm{tor}}(R)$ of Theorem 1.9, induces an anti-equivalence $\mathrm{BT}_{\mathrm{n}}(R) \cong \mathrm{BK}_{\mathrm{n}}(R)$ for all $n \geq 1$.

Proof. This follows formally from Lemma 3.2 and Lemma 3.7 using the exactness and base change properties of the anti-equivalence \mathbb{M} and the classification in the perfect case as stated in Proposition 3.5.

3.2 Normal representations for BK_n-modules

The classification of $\mathrm{BT}_n(R)$ via BK_n -modules at hand, we show that any BT_n -group can be lifted to a BT-group. We fix some $n \in \mathbb{N} \cup \{\infty\}$ and use the notation $W_{\infty}(R) := W(R)$, $\mathrm{BK}_{\infty}(R) := \mathrm{BK}(R)$ and $\mathrm{BT}_{\infty} := \mathrm{BT}(R)$ for a fixed perfectoid ring $R = W(S)/\xi$.

Definition 3.9. A normal representation for a BK_n-module (M, φ, ψ) over R is a triple (L, P, Φ) where $L \subseteq M$ and $P \subseteq M$ are direct summands with $M = L \oplus P$ and $\Phi: M \to M^{\sigma}$ is a $W_n(S)$ -linear bijection, such that, with respect to the decompositions $M = L \oplus P$ and $M^{\sigma} = \Phi(L) \oplus \Phi(P)$, we have $\psi = \alpha \oplus \xi\beta$ and $\varphi = \xi\alpha^{-1} \oplus \beta^{-1}$, where $\alpha = \Phi|_L: L \cong \Phi(L)$ and $\beta = \Phi|_P: P \cong \Phi(P)$.

Remark 3.10. Every triple (L, P, Φ) , where P and L are finite projective $W_n(S)$ -modules and $\Phi: P \oplus L \cong (P \oplus L)^{\sigma}$ is an isomorphism, determines a unique BK_n-module over R, which we call the BK_n-module associated with (L, P, Φ) .

The key ingredient to lifting BK_n -modules is the fact that every BK_n -module admits a normal representation. We need the following version of Nakayama's lemma for the proof.

Lemma 3.11. Let A be a ring and let $g: M \to N$ be a morphism of Amodules with M finitely generated and N finite projective. In both of the following cases g is an isomorphism:

- 1. $g \otimes_A \kappa(x) \colon M \otimes_A \kappa(x) \to N \otimes_A \kappa(x)$ is an isomorphism for all closed points $x \in \text{Spec}(A)$.
- 2. $g \otimes_A A/I: M/IM \to N/IN$ is an isomorphism for some ideal $I \subseteq A$ contained in the Jacobson radical of A.

Proof. 2. follows from 1. which in turn is [1] Lemma 4.5.3.

Lemma 3.12. Every BK_n -module (M, φ, ψ) over R admits a normal representation.

Proof. Let $M = (M, \varphi, \psi) \in BK_n(R)$ and $I := \ker(W_n(S) \to S/\xi_0) = (\xi, p)$ which is contained in the Jacobson radical of $W_n(S)$. The pair $(W_n(S), I)$ even is Henselian since the pairs $(W_n(S), (p))$ and $(S, (\xi))$ have this property. Consider $\overline{M} = (\overline{M}, \overline{\varphi}, \overline{\psi})$, the base change of M to S/ξ_0 . Choose submodules $\overline{L} \subseteq \overline{M}$ and $\overline{Q} \subseteq \overline{M}^{\sigma}$ such that $\overline{M} = \overline{L} \oplus \operatorname{im}(\overline{\varphi})$ and $\overline{M}^{\sigma} = \operatorname{im}(\overline{\psi}) \oplus \overline{Q}$. We have isomorphisms of S/ξ_0 -modules $\overline{\alpha} \colon \overline{L} \to \operatorname{im}(\overline{\psi})$ and $\overline{\beta} \colon \operatorname{im}(\overline{\varphi}) \to \overline{Q}$ which are induced by $\overline{\psi}$ and $\overline{\varphi}^{-1}$ respectively. We claim that we can find a direct summand $Q \subseteq M^{\sigma}$ such that $Q/IQ = \overline{Q}$. Indeed, let Q' be any finite projective $W_n(S)$ -module such that $Q'/IQ' = \overline{Q}$, which exists since the pair $(W_n(S), I)$ is Henselian. We can lift the maps $\overline{Q} \to \overline{M}^{\sigma} \to \overline{Q}$ to maps $Q' \to M^{\sigma} \to Q'$ whose composition is an isomorphism by Nakayama. Then $Q \subseteq M^{\sigma}$, the image of the injection $Q' \to M^{\sigma}$, does the trick. The same argument shows that we can find a direct summand $L \subseteq M$ such that $L/IL = \overline{L}$. Consider the commutative diagram

$$Q \xrightarrow{\varphi} \varphi(Q) \xleftarrow{M} M$$
$$\downarrow / I \qquad \downarrow / I \qquad \downarrow / I$$
$$\overline{Q} \xrightarrow{\overline{\beta}^{-1}} \operatorname{im}(\overline{\varphi}) \xleftarrow{M}.$$

Since $\overline{\beta}^{-1}$ is bijective, we see that $\varphi(Q)/(I\varphi(Q)) = \operatorname{im}(\overline{\varphi})$. Now consider the map $\varphi(Q) \oplus L \to M$. This is an isomorphism modulo I and hence itself an isomorphism by Lemma 3.11. In particular, $\varphi(Q)$ is a finite projective $W_n(S)$ -module. Moreover, $\varphi|_Q \colon Q \to \varphi(Q)$ is an isomorphism since it is an isomorphism modulo I. We set $\beta := (\varphi|_Q)^{-1} \colon \varphi(Q) \to Q$. The same arguments show that $Q \oplus \psi(L) = M^{\sigma}$ and that $\alpha := \psi|_L \colon L \to \psi(L)$ is bijective. Now define $\Phi \colon M = L \oplus \varphi(Q) \to \psi(L) \oplus Q = M^{\sigma}$ to be $\alpha \oplus \beta$. By construction, the triple $(L, \varphi(Q), \Phi)$ then is a normal representation for M.

Proposition 3.13. For $n \ge m$ $(m, n \in \mathbb{N} \cup \{\infty\})$ the truncation functor $BK_n(R) \to BK_m(R)$ is essentially surjective.

Proof. Let M be a BK_m-module over R with normal decomposition (L, P, Φ) . We choose finite projective $W_n(S)$ -modules \tilde{L} and \tilde{P} such that $\tilde{L}/p^m = L$ and $\tilde{P}/p^m = P$. We set $\tilde{M} := \tilde{L} \oplus \tilde{P}$. Moreover, we can lift the map $\tilde{M} \to M \xrightarrow{\Phi} M^{\sigma}$ along $\tilde{M}^{\sigma} \to M^{\sigma}$ to obtain an isomorphism $\tilde{\Phi} \colon \tilde{M} \to \tilde{M}^{\sigma}$. Let $\tilde{M} = (\tilde{M}, \tilde{\varphi}, \tilde{\psi})$ be the BK_n-module over R associated with $(\tilde{L}, \tilde{P}, \tilde{\Phi})$. Then $\tilde{M}/p^m = M$ as desired.

Now setting $n = \infty$ and using Proposition 3.8 and Theorem 1.5, we obtain

Theorem 3.14. Every BT_n -group G over R is of the form $H[p^n]$ for some BT-group H over R.

References

- J. Anschütz, A.-C. Le Bras: Prismatic Dieudonné Theory, preprint (2019), arXiv:1907.10525.
- [2] P. Berthelot, L. Breen, W. Messing: *Théorie de Dieudonné cristalline II*. Lecture Notes in Math. 930, Springer Verlag, 1982.
- B. Bhatt, P. Scholze: Projectivity of the Witt Vector affine Grassmanian. Invent. Math.209(2017), no. 2, 329423.
- [4] A.J. de Jong: Finite locally free group schemes in characteristic p and Dieudonné modules. Invent. Math., 114(1): 89-137, 1993.
- [5] E. Lau: Smoothness of the truncated display functor. J. Amer. Math. Soc. 26 (2013), no. 1, 129-165.
- [6] E. Lau: Dieudonné theory for semiperfect rings and perfectoid rings. Compos. Math.154(2018), no. 9, 1974-2004.