

New Lower Bounds for Tverberg Partitions with Tolerance in the Plane

Sergey Bereg*

Mohammadreza Haghpanah*

February 25, 2020

Abstract

Let P be a set n points in a d -dimensional space. Tverberg's theorem says that, if n is at least $(k-1)(d+1)+1$, then P can be partitioned into k sets whose convex hulls intersect. Partitions with this property are called *Tverberg partitions*. A partition has tolerance t if the partition remains a Tverberg partition after removal of any set of t points from P . Tolerant Tverberg partitions exist in any dimension provided that n is sufficiently large. Let $N(d, k, t)$ be the smallest value of n such that tolerant Tverberg partitions exist for any set of n points in \mathbb{R}^d . Only few exact values of $N(d, k, t)$ are known.

In this paper we establish a new tight bound for $N(2, 2, 2)$. We also prove many new lower bounds on $N(2, k, t)$ for $k \geq 2$ and $t \geq 1$.

1 Introduction

The classical Tverberg's theorem [18] states that a sufficiently large set of points in \mathbb{R}^d can be partitioned into subsets such that their convex hulls intersect. For the history and recent advances around Tverberg's theorem, we refer the reader to [3, 4, 6, 13].

Theorem 1 (Tverberg [18]). *Any set P of at least $(k-1)(d+1)+1$ points in \mathbb{R}^d can be partitioned into k sets P_1, P_2, \dots, P_k whose convex hulls intersect, i.e.*

$$\bigcap_{i=1}^k \text{conv}(P_i) \neq \emptyset. \quad (1)$$

In 1972, David Larman proved the first tolerant version of Tverberg's theorem: any set of $2d+3$ points in \mathbb{R}^d can be partitioned into two sets A and B such that their convex hulls intersect with tolerance one, i.e., for any point $x \in \mathbb{R}^d$, $\text{conv}(A \setminus \{x\}) \cap \text{conv}(B \setminus \{x\}) \neq \emptyset$. García-Colín [8] generalized it for any tolerance. Soberón and Strausz [17] generalized it further for partitions into many sets. The general problem can be stated as follows.

Problem (Tverberg partitions with tolerance). Given positive integers d, k, t , find the smallest positive integer $N(d, k, t)$ such that any set P of $N(d, k, t)$ points in \mathbb{R}^d can be partitioned into k subsets P_1, P_2, \dots, P_k such that for any set $Y \subset P$ of at most t points

$$\bigcap_{i=1}^k \text{conv}(P_i \setminus Y) \neq \emptyset. \quad (2)$$

We call a partition of P into k subsets P_1, P_2, \dots, P_k *t-tolerant* if condition (2) holds for any set Y of at most t points. Several upper bounds for $N(d, k, t)$ are known [10, 12, 14, 17]. Some lower bounds for $N(d, k, t)$ are known [9, 2, 15].

*Department of Computer Science, University of Texas at Dallas. This research is supported in part by NSF award CCF-1718994.

Theorem 2 (Ramírez-Alfonsín [2]). *For any $d \geq 4$*

$$N(d, 2, 1) \geq \left\lceil \frac{5d}{3} \right\rceil + 3. \quad (3)$$

Theorem 3 (García-Colín and Larman [9]).

$$N(d, 2, t) \geq 2d + t + 1. \quad (4)$$

Theorem 4 (Soberón [15]).

$$N(d, k, t) \geq k(t + \lfloor d/2 \rfloor + 1). \quad (5)$$

In contrast to Tverberg's theorem, most bounds are not known to be tight. The only known tight bounds are the following. Larman [12] proved $N(d, 2, 1) = 2d + 3$ for $d \leq 3$. Forge, Las Vergnas and Schuchert [7] proved $N(4, 2, 1) = 11$. For all $k \geq 2, t \geq 1$ and dimension one, $N(1, k, t) = k(t + 2) - 1$ by Mulzer and Stein [14]. In this paper we establish a new tight bound for $N(2, 2, 2)$.

Theorem 5. $N(2, 2, 2) = 10$.

We also improve the bound (5) for the plane.

Theorem 6. *Let c be a positive integer. For any integers $k \geq 2c$ and $t \geq c$*

$$N(2, k, t) \geq k(t + 2) + c. \quad (6)$$

Remark. The bound of Theorem 6 can be stated without parameter c

$$N(2, k, t) \geq k(t + 2) + \min(t, \lfloor k/2 \rfloor). \quad (7)$$

If $d = 2$ and $k = 2$ then Equation (6) provides a lower bound $N(2, 2, t) \geq 2(t + 2) + 1$. We further improve the bound for $k = 2$ and $t \geq 3$.

Theorem 7. *For any $t \geq 3$, $N(2, 2, t) \geq 2t + 6$.*

Several upper bounds for $N(d, k, t)$ are known [8, 10, 16]. For example, Soberón [16] proved $N(d, k, t) = kt + O(\sqrt{t})$ for fixed k and d . Therefore, it is interesting to find lower and upper bounds for $P(d, k, t) = N(d, k, t) - kt$. Theorems 5, 6, and 7 provide new lower bounds for $P(d, k, t)$ for $d = 2$.

Remark. Recently, we improved some lower bounds using computer programs [5]. For example, $N(2, 2, t) \geq 2t + 7$ for $5 \leq t \leq 10$.

The rest of the paper is organized as follows. Section 2 introduces some observations and lemmas that will be used in our proofs. Theorems 6 and 7 are proven in Section 3. Theorem 5 is proven in Section 4. Finally, we discuss the results and open problems in Section 5.

2 Preliminaries

We start with a simple observation.

Observation 8. *Every part in a t -tolerant partition has size at least $t + 1$.*

The following lemma is simple, yet very useful in proving that a partition is not t -tolerant.

Lemma 9. *Let P_1, P_2, \dots, P_k be a partition of a finite set P in \mathbb{R}^d and let X be a subset of a set P_i such that*

$$\text{conv}(X) \cap \text{conv}(P \setminus X) = \emptyset \quad (8)$$

and $|X| = m, |P_i| = t + m$. Then the partition of P is not t -tolerant.

Proof. Let $Y = P_i \setminus X$. Then $|Y| = t$. Therefore

$$\bigcap_{j=1}^k \text{conv}(P_j \setminus Y) = \emptyset \quad (9)$$

by Equation 8 since one of the sets in the intersection is X and every other set is a subset of $P \setminus X$. \square

Two special cases of Lemma 9.

Lemma 10. *A partition P_1, P_2, \dots, P_k of a finite set P in \mathbb{R}^d is not t -tolerant if one of the following conditions holds.*

- (1) *A set P_j contains a vertex of $\text{conv}(P)$ and $|P_j| = t + 1$.*
- (2) *A set P_j contains two points p_i, p_{i+1} such that p_i, p_{i+1} is an edge of $\text{conv}(P)$ and $|P_j| = t + 2$.*

Proof. The first claim follows from Lemma 9 by setting $X = \{p_i\}$ where $p_i \in P_j$ is a vertex of $\text{conv}(P)$. The second claim follows from Lemma 9 by setting $X = \{p_i, p_{i+1}\}$. \square

3 Proofs of Theorems 6 and 7

First, we prove Theorem 6.

Proof of Theorem 6. Construct a set P of $k(t+2) + c - 1$ points in the plane as follows. Let Q be a regular $(k(t+2) - 1)$ -gon with center at the origin O . Place $k(t+2) - 1$ points $q_1, \dots, q_{k(t+2)-1}$ at the vertices of Q and c points p_1, \dots, p_c close to the origin. More specifically, we assume that $|p_1|, \dots, |p_c| < d$ where d is the distance from the origin to the line $p_1 p_k$. Let $V = \{q_1, \dots, q_{k(t+2)-1}\}$. We assume that the vertices of Q are sorted in clockwise order, see Figure 1.




Figure 1: A point set for Theorem 6.

We show that any partition of P into P_1, P_2, \dots, P_k is not t -tolerant. By Observation 8, we assume $|P_i| \geq t + 1$ for all i . If $|P_i| = t + 1$ for some i , then P_i contains a point q_j since $t + 1 > c$. The partition is not t -tolerant by Lemma 10(1).

It remains to consider the case where all sets in the partition have size at least $t + 2$. At most $c - 1$ sets in the partition may have size $\geq t + 3$. Since $k \geq 2c$ there exists a set P_i of size $t + 2$ such that $P_i \subset V$. Since $|V| = k(t+2) - 1$, there exist two points q_a and q_{a+j} with $j \leq k - 1$ such that the vertices between q_a and q_{a+j} are in $Q \setminus P_i$ (we assume here that $q_x = q_y$ if $x \equiv y \pmod{k(t+2) - 1}$). There is a set $P_m, m \neq i$ such that

none of these vertices are in P_m . Therefore the partition is not t -tolerant since $\text{conv}\{q_a, q_b\} \cap \text{conv}(P_m) = \emptyset$. Note that the distance from the origin to any point in $\{p_1, \dots, p_c\}$ is smaller than the distance from the origin to the line $q_a q_b$. \square

Next we prove Theorem 7.




Figure 2: A point set for Theorem 7.

Proof of Theorem 7. Construct a set P of $2t + 5$ points in the plane as follows. First, place $2t + 2$ points in convex position $P' = \{p_1, p_2, \dots, p_{2t+2}\}$ where points are in clockwise order. Then place three points $Q = \{q_1, q_2, q_3\}$ inside the convex hull of P' such that point $q_i, i = 1, 2, 3$ is close to edge $p_{2i-1}p_{2i}$ (it is required that q_i is inside triangles $\Delta p_{2i-2}p_{2i-1}p_{2i}$ and $\Delta p_{2i-1}p_{2i}p_{2i+1}$), see Figure 2. We show that any partition of P into P_1, P_2 is not t -tolerant.

Without loss of generality, we assume $|P_1| \leq |P_2|$. Therefore, $|P_1| \leq t + 2$. By Observation 8, we assume $|P_1| \geq t + 1$. Suppose if $|P_1| = t + 1$. Since $t \geq 3$, P_1 contains a point of P' . The partition is not t -tolerant by Lemma 10(1).

It remains to consider the case where $|P_1| = t + 2$. If set P_1 does not contain a point q_i , then $P_1 \subset P'$ and $|P_1| \geq |P'|/2$. Then P_1 contains two consecutive points of P' . The partition is not t -tolerant by Lemma 10(2). Therefore $P_1 \cap Q \neq \emptyset$.

Suppose $q_i \in P_1$ for some i . If $p_{2i-1} \in P_1$ or $p_{2i} \in P_1$ then the partition is not t -tolerant using $X = \{q_i, p_{2i-1}\}$ or $X = \{q_i, p_{2i}\}$, respectively, and Lemma 9. Therefore, we assume that both points p_{2i-1} and p_{2i} are in P_2 .

Let $m = |P_1 \cap \{q_1, q_2, q_3\}|$. Then $m \in \{1, 2, 3\}$. By the previous argument, set P_2 contains at least m pairs of consecutive points of P' (the pair p_{2i-1}, p_{2i} for each point $q_i \in P_1$). Note that $|P' \cap P_1| = t + 2 - m$ and $|P' \cap P_2| = t + m$. We will show that set P_1 contains a pair of consecutive points of P' . Then, the partition is not t -tolerant by Lemma 10(2).

If $m = 1$, then $|P' \cap P_1| = |P' \cap P_2| = t + 1$ and set P_2 contains a pair of consecutive points of P' . Then set P_1 contains a pair of consecutive points of P' .

Suppose $m = 2$, say $q_i, q_j \in P_1$ where $i < j$. Then set P_2 contains $p_{2i-1}, p_{2i}, p_{2j-1}$, and p_{2j} . There are two intervals $p_{2i+1}, \dots, p_{2j-2}$ and $p_{2j+1}, \dots, p_{2i-2}$ in P' containing points of P_1 . Note that each interval contains an even number of points. If set P_1 does not contain a pair of consecutive points of P' , then set P_1 contains at most half of points in each interval. Then $|P' \cap P_1| < t$. This contradiction implies that set P_1 contains a pair of consecutive points of P' .

Finally, if $m = 3$, then there exists only one interval $p_7, p_8, \dots, p_{2t+2}$ and set P_1 must contain a pair of consecutive points of P' ; otherwise $|P' \cap P_1| < t + 1$. The theorem follows. \square

4 Proof of Theorem 5

We found a special configuration of nine points to prove a lower bound for Theorem 5. For this, we checked point configurations of all order types for $n = 9$. Order types, introduced by Goodman and Pollack [11], are useful for characterizing the combinatorial properties of point configurations. An order type of a set of points p_1, p_2, \dots, p_n in general position in the plane is defined using a mapping that assigns each triple of integers i, j, k with $1 \leq i < j < k \leq n$ the orientation (either clockwise or counter-clockwise) of the triple p_i, p_j, p_k . Two point sets P_1 and P_2 have the same *order type* if there is a bijection π from P_1 to P_2 preserving this map, i.e., for any three distinct points a, b, c in P_1 , the orientation of a, b, c and the orientation of $\pi(a), \pi(b), \pi(c)$ are the same. Aichholzer *et al.* [1] established that there are 158,817 order types for $n = 9$. We developed a program for testing each of these point sets whether it admits a 2-tolerant partition into two sets. The proof of the tolerance of a configuration found by the program turns out to be not simple even using the tools from Section 2.




Figure 3: A set S of nine points for Lemma 11.

Lemma 11. *There exists a set of nine points in the plane that does not admit a 2-tolerant partition into two sets.*




Figure 4: Case 2a. The points of A are red and the points of B are black. The shaded polygons are the convex hulls of $A - C$ and $B - C$.

Proof. Consider a set S of nine points shown in Fig.3 where $S = P \cup Q$, $P = \{p_1, \dots, p_6\}$ and $Q = \{q_1, q_2, q_3\}$. We will prove that any partition of $S = A \cup B$ is not 2-tolerant, i.e. there exists a set $C = \{a, b\} \subseteq S$ such

that

$$\text{conv}(A \setminus C) \cap \text{conv}(B \setminus C) = \emptyset.$$

We can assume that $|A| < |B|$. By Observation 8, we assume $|A| \geq 3$. Suppose that $|A| = 3$. If $A = Q$, then take $C = \{p_1, p_2\}$. If $A \neq Q$, then the partition of S is not 2-tolerant by Lemma 10(1).

It remains to analyze partitions with $|A| = 4$. By Lemma 10(2), we assume that A does not contain two consecutive points p_i, p_{i+1} (assuming $p_7 = p_1$). For example, this implies that A contains at least one point of Q . Consider the following cases depending on $|A \cap P|$.

Case 1. Set A contains exactly one point of P , say p_i . Then $Q \subset A$. One of the sets $\{p_1, p_2, p_3\}$, $\{p_3, p_4, p_5\}$, $\{p_4, p_5, p_6\}$, say set X , does not contain p_i . The partition is not 2-tolerant by Lemma 9 using X .




Figure 5: (a) Case 2b. (b) Case 2c.

Case 2. Set A contains exactly two points of P , i.e. $A = \{p_i, p_j, q_c, q_d\}$ for some $1 \leq i < j \leq 6$ and $1 \leq c < d \leq 3$.

Case 2a. Suppose $i = 1$. By our assumption $j \notin \{2, 6\}$. We can assume that $j \neq 3$ using $X = \{p_4, p_5, p_6\}$ and Lemma 9. So, $j \in \{4, 5\}$.

We also can assume that $q_1 \in B$ using $X = \{p_1, q_1\}$ and Lemma 9. Therefore $\{q_2, q_3\} \subset A$. If $j = 4$ (so $A = \{p_1, p_4, q_2, q_3\}$) then take $C = \{p_1, p_3\}$, see Fig.4(a). If $j = 5$ (so $A = \{p_1, p_5, q_2, q_3\}$) then take $C = \{p_6, q_2\}$, see Fig.4(b).




Figure 6: Case 3a.

Case 2b. Suppose $i = 2$. Then $j \geq 4$. We can assume $j \neq 4$ using $X = \{p_1, p_5, p_6\}$ and Lemma 9. We can assume $j \neq 6$ using $X = \{p_3, p_4, p_5\}$ and Lemma 9. Thus, $j = 5$. We can assume $q_1 \in A$ using $X = \{p_1, p_6, q_1\}$ and Lemma 9.

If $q_3 \in A$ then $A = \{p_2, p_5, q_1, q_3\}$ we can use $X = \{p_3, p_4, q_2\}$ and Lemma 9. If $q_2 \in A$ then $A = \{p_2, p_5, q_1, q_2\}$ and we can use $C = \{p_1, p_5\}$, see Fig.5(a).

Case 2c. Suppose $i, j \in \{3, 4, 5, 6\}$. We can assume $q_1, q_2 \in A$ using $X = \{p_1, p_2, x\}$ for $x \in \{q_1, q_2\}$ and Lemma 9. Since p_i and p_{i+1} cannot be in A together, there are three options. If $p_3, p_5 \in A$, take $C = \{p_4, q_1\}$, see Fig.5(b). If $p_4, p_6 \in A$ then we can use $X = \{p_1, p_2, p_3\}$ and Lemma 9. If $p_3, p_6 \in A$ then we can use $X = \{p_3, q_2\}$ and Lemma 9.

Case 3. Set A contains exactly three points of P , i.e. $\{p_1, p_3, p_5\} \subset A$ or $\{p_2, p_4, p_6\} \subset A$.

Case 3a. Suppose $\{p_1, p_3, p_5\} \subset A$. If $q_1 \in A$ then we can use $X = \{p_1, q_1\}$ and Lemma 9. If $q_2 \in A$, take $C = \{p_2, p_5\}$, see Fig.6(a). If $q_3 \in A$, take $C = \{p_3, p_6\}$, see Fig.6(b).

Case 3b. Suppose $\{p_2, p_4, p_6\} \subset A$. If $q_1 \in A$, take $C = \{p_1, p_4\}$, see Fig.7(a). If $q_2 \in A$, take $C = \{p_3, p_6\}$, see Fig.7(b). If $q_3 \in A$, take $C = \{p_2, p_5\}$, see Fig.7(c). The lemma follows. \square




Figure 7: Case 3b.

Corollary 12. $N(2, 2, 2) \geq 10$.

Soberón and Strausz [17] proved an upper bound $N(d, k, t) \leq (k - 1)(t + 1)(d + 1) + 1$ which implies $N(2, 2, 2) \leq 10$. Therefore $N(2, 2, 2) = 10$.

5 Further Discussion and Open Problems

Improving the bounds for Tverberg partitions with tolerance is an interesting problem. There are many triples (d, k, t) with a gap between known lower and upper bounds. Finding sharp bounds for $N(d, k, t)$ is a challenging problem. For example, Larman [12] conjectured that $N(d, 2, 1) = 2d + 3$ for any $d \geq 1$. The conjecture is open for $d \geq 5$. For dimension $d = 5$, the best known upper bound is $N(5, 2, 1) \leq 13$ by Larman [12] and the best known lower bound is $N(5, 2, 1) \geq 12$ by García-Colín and Larman [9].

Another possibility for a new sharp bound is for $d = 2, k = 2$ and tolerance $t = 3$. The lower bound is $N(2, 2, 3) \geq 12$ by Theorem 7. The best upper bound is $N(2, 2, 3) \leq 13$ by Soberón and Strausz [17]. Based on our computer experiments, we conjecture that $N(2, 2, 3) = 12$. One way to improve the upper bound for $d = 2, k = 2$ and $t = 3$ is to study the following core lemma in the proof of the upper bound for $N(d, k, t)$ [17].

Lemma 13. *Let $p \geq 1$ and $r \geq 0$ be integers, $S \subset \mathbb{R}^p$ be a set of $n = p(r + 1) + 1$ points a_1, a_2, \dots, a_n and G be a group with $|G| \leq p$. If there is an action of G in a set $S' \subset \mathbb{R}^p$ which is compatible with $S \subset S'$, then for each a_i there is a $g_i \in G$ such that the set $\{g_1 a_1, g_2 a_2, \dots, g_n a_n\}$ captures the origin with tolerance r .*

For $d = 2, k = 2$ and tolerance 3, Lemma 13 uses $n = 13$ points since $p = (k - 1)(d + 1) = 3$. Is this bound for n sharp?

5.1 Order types and tolerant Tverberg partitions

Our proof of Theorem 5 uses the set of nine points shown in Figure 3. It was computed by a program that we developed for testing the tolerance of a given point set. We have applied this program to 158,817 order types for $n = 9$ which are provided by Aichholzer *et al.* [1]. Our program found that 155,115 point sets admit a 2-tolerant partition into two sets. Only 3,702 point sets do not admit a 2-tolerant partition into two sets and therefore, can be used as examples for proving the lower bound $N(2, 2, 2) \geq 10$.

We analyzed these 3,702 order types and found that their convex hulls have sizes between three and six. We count the number of different order types for each size of the convex hull, see Table 1. Figures 8, 9, and 10 show some order types with 5, 4, and 3 points on the convex hull, respectively. It follows from our computer experiments that any set of nine points in the plane with at least seven points on the convex hull admits a 2-tolerant partition into two sets. It is interesting to investigate how the size of convex hull of a point set affects lower bounds of $N(d, k, t)$.

h	$h = 3$	$h = 4$	$h = 5$	$h = 6$	$h = 7$	$h = 8$	$h = 9$
N_h	1303	1554	769	76	0	0	0

Table 1: 3,702 order types of nine points that do not admit a 2-tolerant partition into two sets. The point sets are partitioned using the size of the convex hull h . The size of each group is N_h .

References

- [1] O. Aichholzer, F. Aurenhammer, and H. Krasser. Enumerating order types for small point sets with applications. *Order*, 19:265–281, 2002.
- [2] J. L. R. Alfonsín. Lawrence oriented matroids and a problem of McMullen on projective equivalences of polytopes. *Eur. J. Comb.*, 22(5):723–731, 2001.
- [3] I. Bárány, P. V. M. Blagojevic, and G. M. Ziegler. Tverberg’s theorem at 50: extensions and counterexamples. *Notices Amer. Math. Soc.*, 63(7):732–739.




Figure 8: Two point sets of nine points each with five points on the convex hull. The point sets have different order types. Both point sets do not admit a 2-tolerant partition into two sets.




Figure 9: Two point sets of nine points each with four points on the convex hull. The point sets have different order types. Both point sets do not admit a 2-tolerant partition into two sets.

- [4] I. Bárány and P. Soberón. Tverbergs theorem is 50 years old: a survey. *Bulletin of the American Mathematical Society*, 55(4):459–492, 2018.
- [5] S. Bereg and M. Haghpanah. Algorithms for Radon partitions with tolerance. In *6th Internat. Conf. on Algorithms and Discrete Applied Mathematics*, pages 476–487, 2020.
- [6] J. De Loera, X. Goaoc, F. Meunier, and N. Mustafa. The discrete yet ubiquitous theorems of carathéodory, helly, sperner, tucker, and tverberg. *Bulletin of the American Mathematical Society*, 56(3):415–511, 2019.
- [7] D. Forge, M. Las Vergnas, and P. Schuchert. 10 points in dimension 4 not projectively equivalent to the vertices of a convex polytope. *European Journal of Combinatorics*, 22(5):705–708, 2001.
- [8] N. García-Colín. *Applying Tverberg type theorems to geometric problems*. PhD thesis, University College of London, 2007.




Figure 10: Two point sets of nine points each with three points on the convex hull. The point sets have different order types. Both point sets do not admit a 2-tolerant partition into two sets.

- [9] N. García-Colín and D. G. Larman. Projective equivalences of k -neighbourly polytopes. *Graphs and Combinatorics*, 31(5):1403–1422, 2015.
- [10] N. García-Colín, M. Raggi, and E. Roldán-Pensado. A note on the tolerant Tverberg theorem. *Discrete & Computational Geometry*, 58(3):746–754, 2017.
- [11] J. E. Goodman and R. Pollack. Multidimensional sorting. *SIAM J. Comput.*, 12(3):484–507, Aug. 1983.
- [12] D. G. Larman. On sets projectively equivalent to the vertices of a convex polytope. *Bulletin of the London Mathematical Society*, 4(1):6–12, mar 1972.
- [13] J. Matoušek. *Lectures on discrete geometry*, volume 212. Springer, 2002.
- [14] W. Mulzer and Y. Stein. Algorithms for tolerant Tverberg partitions. *Int. J. Comput. Geometry Appl.*, 24(4):261–274, 2014.
- [15] P. Soberón. Equal coefficients and tolerance in coloured Tverberg partitions. *Combinatorica*, 35(2):235–252, 2015.
- [16] P. Soberón. Robust Tverberg and colourful Carathéodory results via random choice. *Combinatorics, Probability and Computing*, 27(3):427–440, 2018.
- [17] P. Soberón and R. Strausz. A generalisation of Tverberg’s theorem. *Discrete & Computational Geometry*, 47(3):455–460, 2012.
- [18] H. Tverberg. A generalization of Radon’s theorem. *J. London Math. Soc.*, 41:123–128, 1966.