

WHAT DETERMINES A VARIETY?

JÁNOS KOLLÁR

ABSTRACT. We show that, over a field of characteristic 0, a normal, projective variety of dimension at least 4 is uniquely determined by its underlying topological space. The proof builds on previous work of Lieblich and Olsson.

CONTENTS

1. Introduction	1
2. Conjectures	7
Noether-Lefschetz theorem over countable fields	7
Independence of intersection points	8
Sections with few zeros	8
3. Set-theoretic complete intersection property	9
4. Mordell-Weil fields	13
5. Reducible scip subsets	16
6. Projective spaces	20
7. Sections and their zero sets	23
8. Linear similarity	27
9. Linkage of divisors and residue fields	31
10. Minimally restrictive linking and transversality	35
11. Degrees of curves and divisors	38
12. Topological pencils	42
13. Complete intersections	45
14. Picard group, class group and Albanese variety	48
15. Commutative algebraic groups	54
16. Special fields	57
References	63

1. INTRODUCTION

By definition, a scheme X is a topological space—which we denote by $|X|$ —and a sheaf of rings \mathcal{O}_X on the open subsets of $|X|$.

As a continuation of [LO19, KLOS20], we study the following natural but seldom considered questions.

- How to read off properties of X from $|X|$?
- Does $|X|$ alone determine X ?

In some cases, the answer is clearly negative.

- If C is an irreducible curve over an infinite field k , then $|C|$ has 1 generic point and $|k|$ closed points, so $|C|$ depends only on the cardinality of k .

- For any variety X , the seminormalization map $X^{\text{sn}} \rightarrow X$ is a homeomorphism, so we should clearly restrict attention to seminormal varieties. The same applies to the weak normalization in positive characteristic.
- Let K/L be a finite field extension and X an irreducible, normal K -variety. We can also view it as an L -variety X_L , and then $|X| = |X_L|$. Note that X_L has at least $\deg(K/L)$ geometric irreducible components, we can thus avoid this problem by working only with geometrically irreducible varieties.
- Purely inseparable morphisms in positive characteristic give homeomorphisms between quite dissimilar varieties.
- A more unexpected example is constructed in [WK81]: $|\mathbb{P}_{\mathbb{F}_p}^2| \sim |\mathbb{P}_{\mathbb{F}_q}^2|$ for any 2 primes p, q . In general it seems that surfaces over finite fields and their algebraic extensions give many similar examples (27).
- Let S, T be real algebraic surfaces birational to $\mathbb{P}_{\mathbb{R}}^2$ and $\Phi : S(\mathbb{R}) \sim T(\mathbb{R})$ a homeomorphism in the classical topology. By [KM09], Φ can be approximated by maps that are homeomorphisms both in the classical and the Zariski topology.

Despite these examples, it is possible that the above are exceptional instances and the answer is positive in almost all other cases.

In Sections 3–6 we study curves $C \subset X$ such that every finite subset $Z \subset C$ is obtained (or almost obtained) as $C \cap D$ for some divisor $D \subset X$. It turns out that this depends mostly on the base field k . Studying a similar question for the union of a curve and a divisor allows us to distinguish projective spaces from all other varieties.

Theorem 1. *Let L be a field of characteristic 0 and K an arbitrary field. Let Y_L be a normal, projective, geometrically irreducible variety of dimension $n \geq 2$ over L such that $|Y_L|$ is homeomorphic to $|\mathbb{P}_K^n|$. Then $K \cong L$ and $Y_L \cong \mathbb{P}_L^n$.*

The linkage of divisors is investigated in Sections 7–12. This leads to an answer to the basic problem in characteristic 0, save in low dimensions. (We comment on the dimension restrictions in (12).)

Theorem 2. *Let K, L be fields of characteristic 0 and X_K, Y_L normal, projective, geometrically irreducible varieties over K (resp. L). Let $\Phi : |X_K| \sim |Y_L|$ be a homeomorphism. Assume that*

$$(2.1) \text{ either } \dim X \geq 4,$$

$$(2.2) \text{ or } \dim X \geq 3 \text{ and } K, L \text{ are finitely generated field extensions of } \mathbb{Q}.$$

Then Φ is the composite of a field isomorphism $\phi : K \cong L$ and an algebraic isomorphism of L -varieties $X_L^\phi \cong Y_L$.

For completeness, let us state the most general form of (2) that could be true.

Conjecture 3. *Let K, L be fields and X_K, Y_L seminormal, geometrically irreducible varieties over K (resp. L). Let $\Phi : |X_K| \sim |Y_L|$ be a homeomorphism. Assume that $\text{char } L = 0$ and $\dim X_L \geq 2$.*

Then Φ is the composite of a field isomorphism $\phi : K \cong L$ and an algebraic isomorphism of L -varieties $X_L^\phi \cong Y_L$.

It is possible that modifications of the current methods will handle the seminormal and proper case. Our approach has many difficulties with open varieties; though some cases are treated in [LO19, KLOS20].

Ancillary results are collected in Sections 13–16. These are mostly known but hard to find references for the forms that we need. The notion of weakly Hilbertian fields introduced in Section 16 is new and may be of independent interest.

4 (Terminology and notation). For ease of reference, here is a list of new or non-standard terminology and notation that we use.

Albanese variety, classical $\mathbf{Alb}(X)$, à la Grothendieck $\mathbf{Alb}^{\text{gr}}(X)$ (128).

Absolutely scip, variant of scip (32).

Ample-ci, complete intersection of ample divisors (117).

Ample-sci, set-theoretic complete intersection of ample divisors (117).

Anti-Mordell-Weil field, (152).

Bertini-Hilbert dimension of a field, (63).

$\text{BH}(k)$, the Bertini-Hilbert dimension; it is either 1 or 2 (63).

Detects linear similarity, (65).

Field-locally thin subset, (71).

Generically scip, variant of scip (19).

Linked, L-linked, (75).

Linking is free, (76).

Linking on W_2 determines linking on W_1 , (87).

Linking is minimally restrictive, (87).

Locally finite field, algebraic extension of a finite field (150).

Mordell-Weil field, (151).

Non-Cartier center, (136).

Scip, set-theoretic complete intersection property.

For irreducible subsets (19) and for reducible subsets (36).

Scip with finite defect, variant of scip (29).

Thin subset, (71).

Stably dense, (64).

Topological pencil, t-pencil, (105).

Weakly Hilbertian field, (156).

$\Gamma^B(X, L)$, sections of (powers of) L of that vanish exactly along B (55).

$\mathcal{R}_V^B(X, L)$, restriction of $\Gamma^B(X, L)$ to $V \subset X$ (57).

$\mathcal{R}_k^A \mathbb{G}_m$, Weil restriction of the multiplicative group (145).

$\rho(X)$, Picard number using Cartier divisors.

$\rho^{\text{cl}}(X)$, Picard number (or class rank) using non-Cartier divisors (124).

$\Sigma(X)$, $\{x \in X : \dim \mathcal{O}_{x, X} = 0, \text{ or } = 1 \text{ and not regular, or not } S_2.\}$ (54.3).

\sim , linear equivalence of divisors D_1, D_2 .

$\sim_{\mathbb{Q}}$, \mathbb{Q} -linear equivalence: $mD_1 \sim mD_2$ for some $m \neq 0$.

\sim_s , linear similarity: $m_1D_1 \sim m_2D_2$ for some $m_1, m_2 \neq 0$ (64).

\sim_{sa} , linear similarity + D_1, D_2 ample and irreducible (8).

Introduction to Sections 3–6

Let X be a normal, projective K -variety over some field K and $C \subset X$ an irreducible curve. In Sections 3–4 we study which finite subsets of C are obtained as the intersection of C with some divisor; a condition that depends only on the topology of the pair $|C| \subset |X|$.

Somewhat surprisingly, at the most basic level the answer is governed by the base field K . More precisely, by the qualitative behavior of the group of K -points

of Abelian varieties over K . There are 3 classes of fields K for which the ‘size’ of $A(K)$ is about the same for every Abelian variety over K .

- (Finite fields) For these $A(K)$ is finite. More generally, if K is *locally finite* then $A(K)$ is a torsion group.
- (Number fields) For these $A(K)$ is finitely generated by the Mordell-Weil theorem. More generally, the same holds for fields that are finitely generated over a prime field [LN59].
- (Geometric fields) For these $A(K)$ has infinite rank. This holds for example if K is algebraically closed, except for $\overline{\mathbb{F}}_p$. These are called anti-Mordell-Weil fields [IL19]; see (152) for the main examples.

Roughly speaking, our results show how to read off closely related properties of K from the topology of $|X|$ in the first 2 cases and to recognize rational curves on X in the third case.

The conclusions are most complete for locally finite fields.

Theorem 5. *Let X be an irreducible, quasi-projective variety of dimension ≥ 2 over a perfect field K . The following are equivalent.*

- (5.1) *For every irreducible curve $C \subset X$ and every finite, closed subset $P \subset C$, there is a divisor $D \subset X$ such that $C \cap D = P$ (as sets).*
- (5.2) *K is locally finite.*
- (5.3) *$A(K)$ is a torsion group for every commutative algebraic group A over K .*

We have a more complicated characterization of the Mordell-Weil case.

Theorem 6. *Let X be an irreducible, quasi-projective variety of dimension ≥ 2 over a field K . The following are equivalent.*

- (6.1) *For every irreducible curve $C \subset X$ there is a finite, closed subset $\Sigma \subset C$ such that for every finite, closed subset $P \subset C$, there is a divisor $D \subset X$ such that $P \subset C \cap D \subset P \cup \Sigma$ (as sets).*
- (6.2) *$A(K)$ has finite \mathbb{Q} -rank for every Abelian variety A over K .*

In the geometric cases our considerations yield a Zariski-topological characterization of rational curves. The complete statement in (34) needs several definitions, so here we state it somewhat informally.

Proposition 7. *Let X be an irreducible, quasi-projective variety of dimension $n \geq 2$ over an anti-Mordell-Weil field and $C \subset X$ an irreducible, geometrically reduced curve. One can decide using only the topology of the pair $|C| \subset |X|$ whether C is rational or not.*

The last result is especially useful if X contains many rational curves, for example for $X = \mathbb{P}^n$. However, in Sections 5–6 we get better results by observing that, from the topological point of view,

$$(\text{hyperplane}) \cup (\text{line}) \subset \mathbb{P}^n$$

is a very unusual configuration. This leads to the proof of (1).

Introduction to Sections 7–12

A program towards showing that $|X|$ determines X was started in [LO19]. As an intermediate step, [LO19] proposed to look for additional data that, together with $|X|$, determine X . A candidate is *linear equivalence* of divisors, denoted

by $D_1 \sim D_2$. In (8) we introduce a variant, called *linear similarity of ample divisors* and denoted by \sim_{sa} . Then we show first that $|X|$ determines \sim_{sa} , then that $(|X|, \sim_{\text{sa}})$ determines \sim and finally [LO19] proves that $(|X|, \sim)$ determines X . Symbolically:

$$|X| \xrightarrow{(9)} (|X|, \sim_{\text{sa}}) \xrightarrow{(10)} (|X|, \sim) \xrightarrow{[\text{LO19}]} X.$$

Definition 8 (Linear similarity of ample divisors). Let X be a normal variety and $\text{PDiv}(X)$ the set of prime divisors on X . We define a relation on $\text{PDiv}(X) \times \text{PDiv}(X)$ by declaring that $D_1 \sim_{\text{sa}} D_2$ iff

- (8.1) D_1, D_2 are \mathbb{Q} -Cartier, ample and
- (8.2) $m_1 D_1 \sim m_2 D_2$ for some $m_1, m_2 > 0$.

Note that $\text{rank}_{\mathbb{Q}} \text{Cl}(X) = 1$ iff $D_1 \sim_{\text{sa}} D_2$ for any 2 prime divisors on X . In these cases the relation \sim_{sa} carries no extra information.

The first step of the proof finds \sim_{sa} .

Proposition 9. *Let X be a normal, projective variety of dimension ≥ 3 over a field k . Assume that k is not locally finite. Then $|X|$ determines \sim_{sa} .*

The proof is actually a quite short argument in Section 8, which is surprising since \sim and \sim_{sa} seem very closely related at first sight. We show how to recognize

- (9.1) irreducible, ample \mathbb{Q} -Cartier divisors (67),
- (9.2) linear similarity of irreducible, ample \mathbb{Q} -Cartier divisors (68), and
- (9.3) irreducible \mathbb{Q} -Cartier divisors (69).

Then a longer argument shows that once we know $|X|$ and \sim_{sa} , then we also know \sim .

Theorem 10. *Let X be a normal, projective, geometrically irreducible variety over a field k of characteristic 0. Assume that*

- (10.1) *either $\dim X \geq 4$,*
- (10.2) *or $\dim X \geq 3$ and k is a finitely generated field extension of \mathbb{Q} .*

Then $|X|$ and \sim_{sa} together determine \sim . (See (116) for a more general version.)

This is longer; we step-by-step recognize the following objects/properties.

- (10.1) k -points (82).
- (10.2) Isomorphism of residue fields of closed points (83).
- (10.3) Isomorphism of reduced, 0-dimensional subschemes (84).
- (10.4) Transversality of 0-dimensional intersections of subvarieties (91).
- (10.5) Two irreducible curves having the same degree (95).
- (10.6) Two irreducible divisors having the same degree (96).
- (10.7) \mathbb{Q} -linear equivalence of ample divisors (97.5).

We can now use (107) to recognize algebraic pencils of divisors.

The remaining problem is to decide which members of a topological pencil are members of the corresponding algebraic pencil; this is the *true membership* problem [KLOS20, 5.3.5] As in [KLOS20, Chap.5], instead of a complete solution, we only deal with ‘well behaved’ linear systems. We find sufficient conditions for

- (10.8) linearity of a pencil (109),
- (10.9) membership in a pencil (113) and

(10.10) linear equivalence of reduced divisors (115).

Then, as in [KLOS20, 5.3.8], we see that linear equivalences between reduced divisors generate the full linear equivalence relation. The next result then completes the proof of (2).

Theorem 11. [LO19, Main Thm.] *Let X be a normal, projective, geometrically irreducible variety of dimension ≥ 2 over a field. Then $|X|$ and \sim together determine X . \square*

The main technical tool for all this is the study of linkage of divisors.

12 (Linkage of divisors). Let X be a normal, projective, irreducible k -variety and L an ample line bundle on X . Let $Z_1, Z_2 \subset X$ be closed, irreducible subvarieties such that $\dim(Z_1 \cap Z_2) = 0$. We consider the following

Linkage problem 12.1. We say that $s_i \in H^0(X, L^{m_i})$ with zero sets $H_i := (s_i = 0)$ are L -linked on $Z_1 \cup Z_2$ if there is an $s \in H^0(X, L^r)$ with zero set $H := (s = 0)$ such that

$$\text{Supp}(H_1 \cap Z_1) \cup \text{Supp}(H_2 \cap Z_2) = \text{Supp}((Z_1 \cup Z_2) \cap H).$$

Note that if the Picard number of X is 1, then this is clearly a question involving only the underlying topology $|X|$. In fact, by (9), this is almost always a question about $|X|$.

Sufficient condition 12.2. If $s_1^{r_1}|_{Z_1 \cap Z_2} = c \cdot s_2^{r_2}|_{Z_1 \cap Z_2}$ for some nonzero $c \in k^\times$ and $r_1, r_2 \in \mathbb{N}$, then the $H_i := (s_i = 0)$ are L -linked on $Z_1 \cup Z_2$.

Next note that $s_i|_{Z_1 \cap Z_2}$ can be an arbitrary element of $H^0(Z_1 \cap Z_2, L|_{Z_1 \cap Z_2}) \cong H^0(Z_1 \cap Z_2, \mathcal{O}_{Z_1 \cap Z_2})$ for L sufficiently ample. Thus we obtain the following.

Proposition 12.3. If the sufficient condition (12.2) is necessary and any two H_1, H_2 are L -linked, then $H^0(Z_1 \cap Z_2, \mathcal{O}_{Z_1 \cap Z_2}) \cong k$. That is, $Z_1 \cap Z_2$ is a reduced k -point of X .

This gives us a topological way to identify k -points and also check whether an intersection is transverse or not.

When is condition (12.2) necessary? 12.4. We know that $\text{Supp}(s_i|_{Z_i} = 0) = \text{Supp}(s|_{Z_i} = 0)$. Thus if

- (a) the Z_i are normal, geometrically irreducible, and
- (b) the $\text{Supp}(s_i|_{Z_i} = 0)$ are irreducible, then

$$s_i^{m_i}|_{Z_i} = c_i \cdot s^{n_i}|_{Z_i} \tag{12.4.c}$$

for some nonzero $c_i \in k^\times$ and $m_i, n_i \in \mathbb{N}$. Thus we conclude that there is a constant $c \in k^\times$ such that

$$s_1^{m_1 n_2}|_{Z_1 \cap Z_2} = c \cdot s_2^{m_2 n_1}|_{Z_1 \cap Z_2}, \tag{12.4.d}$$

as needed.

It remains to deal with the conditions (12.4.a–b). It turns out that normality can be avoided, this is worked out in Section 7. Instead of geometric irreducibility, the key condition is geometric connectedness, which is guaranteed if the Z_i are set-theoretic complete intersections of ample divisors. Thus the troublesome condition is (12.4.b).

Irreducibility of $H \cap Z$ 12.5. Let $Z \subset X$ be an irreducible subvariety and $H \subset X$ a general ample divisor. When is $H \cap Z$ irreducible?

Bertini's theorem says that this holds if $\dim Z \geq 2$. Since we also need $\dim(Z_1 \cap Z_2) = 0$, we must have $\dim X \geq 4$. For the best results we also need $Z_1 \cap Z_2$ to be a single point, which usually can be arranged only if $\dim X \geq 5$. This is the case when the methods work best.

There are 2 ways to lower the dimension. First, it turns out that we get almost everything if one of the Z_i has dimension ≥ 2 , the other can be a curve. Thus we get all results if $\dim X \geq 4$, as in (2.1).

If k is finitely generated over \mathbb{Q} , then a theorem of Hilbert guarantees that $H \cap Z$ is irreducible for most ample divisors H even if $\dim Z = 1$. This property defines Hilbertian fields (153), and, for such fields, we can work with varieties of dimension ≥ 3 ; leading to (2.2).

In order to give unified treatments, we introduce the Bertini-Hilbert dimension of fields in (63).

Introduction to Sections 13–16

These sections provide background material, most of which is known but may be hard to find in the form that we need. Section 13 summarizes properties of complete intersections and various Bertini-type theorems.

The theory of Picard group, Picard variety and Albanese variety is recalled in Section 14. The literature is much less complete about the class group and its scheme version, which does not even seem to have a name.

Basic results on commutative algebraic groups and the multiplicative group of Artin algebras are studied in Section 15.

Section 16 recalls the definitions and basic properties of locally finite, Mordell-Weil, anti-Mordell-Weil and Hilbertian fields. We introduce the new notion of weakly Hilbertian fields (156).

Acknowledgments. My interest in this subject was started by a lecture of M. Olsson about [LO19] at MSRI. After that, a series of discussions with M. Lieblich and M. Olsson led to further results [KLOS20]. These had a major influence on this work.

I received help from M. Larsen with algebraic tori, B. Poonen with Bertini and Noether-Lefschetz theorems, J. Silverman with Néron's theorem, K. Smith with discrete valuation rings and C. Voisin with Noether-Lefschetz theorems; I am grateful to all of them.

Partial financial support was provided by the NSF under grant number DMS-1901855.

2. CONJECTURES

We discuss several conjectures that came up while studying the above questions, but which are of independent interest.

Noether-Lefschetz theorem over countable fields.

The following is probably old, [Ter85] attributes a version of it to T. Shioda.

Conjecture 13. *Let k be a field that is not locally finite, X a normal projective k -variety of dimension ≥ 3 and H an ample Cartier divisor. Then, for $m \gg 1$ and for 'most' k -divisors $D \in |mH|(k)$, the restriction maps*

$$\mathrm{Pic}(X) \rightarrow \mathrm{Pic}(D) \quad \text{and} \quad \mathrm{Cl}(X) \rightarrow \mathrm{Cl}(D) \quad \text{are isomorphisms.}$$

The traditional statement of the Noether-Lefschetz theorem says that the conclusion holds outside a countable union of proper, closed subvarieties of $|mH|$; see [Gro68] for the Picard group and [RS06, RS09, Ji20] for the class group. This gives a positive answer to the conjecture whenever k is uncountable.

For countable algebraically closed fields a positive answer is given in [And96, MP12, Amb18, Chr18]. See also [Ter85] for similar results over \mathbb{Q} for complete intersections in \mathbb{P}^n .

Independence of intersection points.

To start with an example, let $E \subset \mathbb{P}^2$ be an elliptic curve, and $L \subset \mathbb{P}^2$ a very general line intersecting E at 3 points p_1, p_2, p_3 . Let $[H] \in \text{Pic}(E)$ denote the hyperplane class. One can see that $m_1 p_1 + m_2 p_2 + m_3 p_3 \sim nH$ holds only for $m_1 = m_2 = m_3 = n$. This is easy over \mathbb{C} , gets quite a bit harder over \mathbb{Q} if we want the line to be also defined over \mathbb{Q} . The following conjecture says that a similar claim holds for all smooth curves.

Conjecture 14. *Let k be a field that is not locally finite. Let C be a smooth, projective curve of genus ≥ 1 over k and L a very ample line bundle on C . For a section $s \in H^0(C, L)$ write $\{p_i(s) : i \in I\}$ (resp. $\{\bar{p}_i(s) : i \in \bar{I}\}$) for the closed points (resp. \bar{k} -points) of $(s = 0)$. Then, for ‘most’ sections, we have injections*

$$\begin{aligned} \bigoplus_{i \in I} \mathbb{Z}[p_i(s)] &\hookrightarrow \text{Pic}(C) \quad (\text{weak form}), \\ \bigoplus_{i \in \bar{I}} \mathbb{Z}[\bar{p}_i(s)] &\hookrightarrow \text{Pic}(C_{\bar{k}}) \quad (\text{strong form}). \end{aligned}$$

It is not clear what ‘most’ should mean. It is possible that this holds outside a field-locally thin set (71), but some heuristics suggest otherwise.

For the proof of (80) we would need the following stronger variant. If true, it would allow us to prove (2) for 3-folds as well.

Conjecture 15. *Using the notation of (14), let $A \subsetneq \mathbf{Pic}^\circ(C)$ be an Abelian subvariety and $\Gamma \subset \text{Pic}(C)$ a finitely generated subgroup that contains $[L]$. Then, for ‘most’ sections, we have injections*

$$\begin{aligned} \bigoplus_{i \in I} \mathbb{Z}[p_i(s)] / \sum_{i \in I} [p_i(s)] &\hookrightarrow \text{Pic}(C) / \langle A(k), \Gamma \rangle \quad (\text{weak form}), \\ \bigoplus_{i \in \bar{I}} \mathbb{Z}[\bar{p}_i(s)] / \sum_{i \in \bar{I}} [\bar{p}_i(s)] &\hookrightarrow \text{Pic}(C_{\bar{k}}) / \langle A(\bar{k}), \Gamma \rangle \quad (\text{strong form}). \end{aligned}$$

Sections with few zeros.

The next 2 conjectures posit that, for ‘most’ ample line bundles, every section has many zeros.

Conjecture 16. *Let K be an algebraically closed field other than $\bar{\mathbb{F}}_p$. Let C be a smooth, projective curve over K . Then, for ‘most’ ample line bundles L , every section of L^m has at least $g(C)$ zeros for every $m \geq 1$.*

Line bundles of degree d that have a section with fewer than g zeros form a closed subset of dimension $g - 1$ of $\mathbf{Pic}^d(C)$ obtained as the image of the maps

$$\phi_{\mathbf{m}} : C^{g-1} \rightarrow \mathbf{Pic}^d(C) \quad \text{given by} \quad (c_1, \dots, c_{g-1}) \mapsto \mathcal{O}_C(\sum_i m_i [c_i]),$$

where $\mathbf{m} := (m_1, \dots, m_{g-1})$ such that $\sum m_i = d$. Thus (16) is true if K is uncountable. The most interesting open case is probably \mathbb{Q} . By (162), there is a curve C and a line bundle L over \mathbb{Q} , such that every section of L^m has at least 2 zeros for every $m \geq 1$.

We prove the nodal rational curve cases of (16) in (164).

Thinking of the curve C as a subvariety of its Jacobian leads to the following stronger form.

Conjecture 17. *Let K be an algebraically closed field other than $\overline{\mathbb{F}}_p$. Let A be an Abelian variety over K and $Z_i \subset A$ subvarieties such that $\sum_i \dim Z_i < \dim A$. Then, for ‘most’ $p \in A(K)$, the equation*

$$n[p] = \sum_i m_i [z_i] \quad n, m_i \in \mathbb{Z}, z_i \in Z_i(K),$$

has only the trivial solution $n = m_i = 0$.

Next we give an example with only one Z_i where this holds.

Example 18. Let k be any field. Assume that $A = B \times E$ where B is a simple Abelian variety, E an elliptic curve, and we have only one $Z = Z_1 \subset A$ of dimension $\leq \dim A - 2$. Assume also that Z does not contain any translate of E .

Let $\pi : A \rightarrow B$ be the coordinate projection. If $p \in E(k)$, $z \in Z(k)$ and $n[p] = m[z]$, then $m[\pi(z)] = 0$, that is, $\pi(z)$ is a torsion point in $\pi(Z)$. By [Zha98] there are only finitely many such, so there are only finitely many $\{z_j \in Z : j \in J\}$ for which there is an $m_j > 0$ such that $m_j [z_j] \in E$.

Thus if $p \in E(k)$ then $n[p] = m[z]$ has a nontrivial solution iff p is in the saturation of $m_j [z_j]$ for some $j \in J$.

If $\text{rank}_{\mathbb{Q}} E(k) \geq 2$, then finitely many subgroups of \mathbb{Q} -rank 1 do not cover $E(k)$.

To get such Jacobian examples, fix an elliptic curve E over $\overline{\mathbb{Q}}$ and let C be a sufficiently general member of a very ample linear system on $E \times \mathbb{P}^1$. Then, by [Koc18, 1.6], $\text{Jac}(C)$ is isogeneous to the product of E and of a simple Abelian variety B .

3. SET-THEORETIC COMPLETE INTERSECTION PROPERTY

Definition 19. Let X be an irreducible scheme and $Z \subset X$ a closed, irreducible subset. We say that Z has the *set-theoretic complete intersection property*—or that Z is *scip*—if the following holds.

(19.1) Let $D_Z \subset Z$ be a closed subset of pure codimension 1. Then there is an effective divisor $D_X \subset X$ such that $\text{Supp}(D_X \cap Z) = D_Z$.

In some cases only ‘nice’ subvarieties $D_Z \subset Z$ are set-theoretic complete intersections. It is usually hard to formulate this in general, but the next variant allows us to ignore finitely many ‘bad’ points of Z .

We say that Z is *generically scip* if there is a finite (not necessarily closed) subset $\Sigma_Z \subset Z$ such that the following holds.

(19.2) Let $D_Z \subset Z$ be a closed subset of pure codimension 1 that is disjoint from Σ_Z . Then, for every finite (not necessarily closed) subset $\Sigma_X \subset X \setminus D_Z$, there is an effective divisor $D_X \subset X$ that is disjoint from Σ_X and such that $\text{Supp}(D_X \cap Z) = D_Z$.

As a simple example, the quadric cone $Q \subset \mathbb{P}^4$ is not scip (over \mathbb{C}) but it is generically scip with $\Sigma_Z = \{\text{vertex}\}$ and Σ_X arbitrary.

The introduction of Σ_Z means that we do not have to worry about some very singular points on Z . This is especially clear on curves, where we may assume that Σ_Z contains all singular points.

The introduction of Σ_X at first seems to make finding D_X harder. However, if X is normal and Σ_X contains all non-Cartier centers of X (136), then D_X is a Cartier

divisor. Thus we can usually work with the Picard group of X (for which there are solid references), rather than the class group (for which modern references seem to be lacking).

It is clear that these notions depend only on the topological pair $|Z| \subset |X|$.

At the beginning we study the case when Z is an irreducible curve, but later we need to understand many cases when Z is reducible, and not even pure dimensional.

We check in (28) that being generically scip is invariant under purely inseparable morphisms and purely inseparable base field extensions. Thus, in order to save considerable trouble with non-reduced group schemes, we usually work over perfect base fields.

Lemma 20. *Let X be a normal, quasi-projective variety over a perfect field k and $C \subset X$ an integral, generically scip curve. Then*

$$\text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(C)] \quad \text{is a torsion group.}$$

Proof. We may assume that $\Sigma_Z \supset \text{Sing } C$ and $\Sigma_Z \cup \Sigma_X$ contains all non-Cartier centers of X (136). Let $p \in C \setminus \Sigma_Z$ be a point. By assumption there is an effective, Cartier divisor D_p such that $\text{Supp}(D_p \cap C) = \{p\}$. We do not know the intersection multiplicity at p , so we can only say that $\mathcal{O}_X(D_p)|_C \cong \mathcal{O}_C(m[p])$ for some $m > 0$. (Here we use that p is a regular point.) That is, $\mathcal{O}_C(p)$ is torsion in $\text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(C)]$. Since the $\mathcal{O}_C(p)$ generate $\text{Pic}(C)$, we are done. \square

The rest of Sections 3–4 is essentially devoted to trying to understand the converse of (20). Let us see first that the direct converse does not hold.

Example 21. Let $C \subset \mathbb{P}^2$ be a smooth cubic over a number field. By the Mordell-Weil theorem $\text{Pic}(C)$ is finitely generated. Choose points $\{p_i \in C : i \in I\}$ such that $\mathcal{O}_C(1)$ and the $\mathcal{O}_C(p_i)$ form a basis of $\text{Pic}(C) \otimes \mathbb{Q}$.

Let X be obtained by blowing up the points $p_i \in C \subset \mathbb{P}^2$. Let $E_i \subset X$ be the exceptional curves and let $C_X \subset X$ denote the birational transform of C . Note that $\text{Pic}(X)$ is spanned by the E_i and the pull-back of $\mathcal{O}_{\mathbb{P}^2}(1)$. Thus $\text{Pic}(X) \rightarrow \text{Pic}(C_X)$ is an injection with torsion cokernel.

Claim 21.1. $C_X \subset X$ is not generically scip.

Proof. Choose $n_i > 0$ and let $p \in C_X$ be a closed point such that $[p] \sim \sum n_i [p_i]$. Assume that $\{p\} = \text{Supp}(C_X \cap D)$ for some effective divisor $D \subset X$. Then

$$\mathcal{O}_X(D)|_{C_X} \cong \mathcal{O}_{C_X}(m[p]) \cong \mathcal{O}_{C_X}(\sum mn_i [p_i])$$

for some $m > 0$. Since $\text{Pic}(X) \rightarrow \text{Pic}(C_X)$ is an injection, this implies that $D \sim \sum mn_i [E_i]$. But then $D = \sum mn_i [E_i]$ and so $C_X \cap D = \{p_i : i \in I\}$. \square

The following is a partial converse of (20).

Lemma 22. *Let X be a projective variety over a field k and $C \subset X$ a reduced, irreducible curve. Assume that*

$$\text{coker}[\text{Pic}^\circ(X) \rightarrow \text{Pic}^\circ(C)] \quad \text{is a torsion group.}$$

Then C is scip.

Proof. Let L be an ample line bundle such that $H^1(X, L^m \otimes T \otimes I_C) = 0$ for every $m \geq 1$ and every $T \in \text{Pic}^\circ(X)$, where $I_C \subset \mathcal{O}_X$ denotes the ideal sheaf of C . Then $H^0(X, L^m \otimes T) \rightarrow H^0(C, L^m \otimes T|_C)$ is surjective for every $m \geq 1$. Set $d = \deg_C L$.

For a point $p \in C$ let P be a Cartier divisor on C whose support is p and set $r = \deg P$. (We can take $P = p$ if p is a regular point.) Then $L_C^r(-dP) \in \text{Pic}^\circ(C)$, thus there is an $m \geq 1$ and $T \in \text{Pic}^\circ(X)$ such that

$$L_C^{mr}(-mdP) \cong T^{-1}|_C.$$

This gives a section $s_C \in H^0(C, L^m \otimes T|_C)$ whose divisor is mdP . It lifts to a section $s \in H^0(X, L^m \otimes T)$ and $D := (s = 0)$ works. \square

The next example shows that C can be scip even if $\text{coker}[\text{Pic}^\circ(X) \rightarrow \text{Pic}^\circ(C)]$ is non-torsion.

Example 23. Again let $C \subset \mathbb{P}^2$ be a smooth cubic over a number field. Choose a finite subset $L_i \in \text{Pic}^\circ(C)$, closed under inverse, that generates a full rank subgroup.

Choosing general sections in each $L_i^{-1}(3)$, their zero sets $P_i \subset C$ are irreducible and distinct. Let $S_i \rightarrow \mathbb{P}^2$ denote the blow-up of P_i and $C_i \subset S_i$ the birational transform of C . Then $\mathcal{O}_{S_i}(C_i)$ is a nef line bundle on S_i and $\mathcal{O}_{S_i}(C_i)|_{C_i} \cong L_i$.

Finally consider the diagonal embedding $C \subset \prod_i C_i \subset \prod_i S_i =: X$.

Claim 23.1. $C \subset X$ is scip.

Proof. The key point is that $\{T|_C : T \in \text{Pic}(X) \text{ and } T \text{ is nef}\} \subset \text{Pic}^\circ(C)$ contains a full rank subgroup of $\text{Pic}^\circ(C)$. By Fujita's vanishing theorem [Laz04, 1.4.35], there is an ample line bundle L such that $H^1(X, L^m \otimes T \otimes I_C) = 0$ for every $m \geq 1$ and every nef T . The rest of the argument in the proof of (22) works.

24 (Cokernel of $\text{Pic}(X) \rightarrow \text{Pic}(Y)$). (See (123) for definitions and notation involving the Picard group.)

If X is a proper scheme then $\text{Pic}(X)$ is an extension of $\text{NS}(X)$ by $\text{Pic}^\circ(X)$. While $\text{NS}(X)$ is always a finitely generated abelian group, $\text{Pic}^\circ(X)$ can be trivial or very large, depending on the ground field and X . However, $\mathbf{Pic}(X)$ is an algebraic group and $\mathbf{Pic}(X)(k)/\text{Pic}^\circ(X)$ is torsion. Thus, if $p : Y \rightarrow X$ is a morphism, we aim to understand $p^* : \text{Pic}(X) \rightarrow \text{Pic}(Y)$, in terms of

$$p^* : \mathbf{Pic}^\circ(X) \rightarrow \mathbf{Pic}^\circ(Y) \quad \text{and} \quad p^* : \text{NS}(X) \rightarrow \text{NS}(Y). \quad (24.1)$$

We have better theoretical control of these maps since the first is a map of Abelian varieties and the second a map of finitely generated abelian groups.

Proposition 25. *Let $p : Y \rightarrow X$ be a morphism of proper k -schemes. Then*

$$\begin{aligned} \text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(Y)] &\geq \\ &\geq \text{rank}_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^\circ(X) \rightarrow \mathbf{Pic}^\circ(Y)](k) - \text{rank}_{\mathbb{Q}} \text{NS}(X). \end{aligned}$$

Proof. Let $\text{Pic}^*(X) \subset \text{Pic}(X)$ denote the preimage of $\text{Pic}^\circ(Y)$. Then $\text{Pic}^\circ(X) \subset \text{Pic}^*(X)$ and the quotient is a subgroup of $\text{NS}(X)$. Thus we see that

$$\begin{aligned} &\text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(Y)] \\ &\geq \text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}^*(X) \rightarrow \text{Pic}^\circ(Y)] \\ &\geq \text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}^\circ(X) \rightarrow \text{Pic}^\circ(Y)] - \text{rank}_{\mathbb{Q}} \text{NS}(X) \\ &= \text{rank}_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^\circ(X) \rightarrow \mathbf{Pic}^\circ(Y)](k) - \text{rank}_{\mathbb{Q}} \text{NS}(X), \end{aligned}$$

where the last equality holds since the maps $\text{Pic}^\circ(Z) \otimes \mathbb{Q} \rightarrow \mathbf{Pic}^\circ(Z)(k) \otimes \mathbb{Q}$ are isomorphisms for proper k -schemes and $A \mapsto A(k) \otimes \mathbb{Q}$ is an exact functor of commutative group varieties (144). \square

The first application is a characterization of locally finite fields.

Theorem 26. *Let X be an irreducible, quasi-projective variety of dimension ≥ 2 over a perfect field k . The following are equivalent.*

(26.1) *k is locally finite.*

(26.2) *Every irreducible curve $C \subset X$ is scip.*

(26.3) *Every irreducible curve $C \subset X$ is generically scip.*

Proof. Assume first that k is locally finite and let $\bar{X} \supset X$ be a compactification. Let $C \subset X$ be an irreducible curve with closure \bar{C} . If k is locally finite then $\mathbf{Pic}^\circ(\bar{C})(k)$ is torsion by (141.1), hence \bar{C} is scip by (22) and so is $C \subset X$. (26.2) \Rightarrow (26.3) is clear.

It remains to prove that (26.3) \Rightarrow (26.1). Note that if (26.3) holds for X then it holds for every open subset of it, we may thus assume that X is normal (or even smooth). Let $\bar{X} \supset X$ be a normal compactification.

If k is not locally finite then let $\bar{C} \subset \bar{X}^{\text{ns}}$ be an irreducible curve with at least 1 node contained in X . Note that $\mathbf{Pic}^\circ(\bar{X})$ is an Abelian variety (123) and $\mathbf{Pic}^\circ(\bar{C})$ contains a k -torus (142). Thus $\text{coker}[\mathbf{Pic}^\circ(\bar{X}) \rightarrow \mathbf{Pic}^\circ(\bar{C})]$ contains a k -torus, hence its \mathbb{Q} -rank is infinite (141). Thus $\text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(\bar{X}) \rightarrow \text{Pic}(\bar{C})] = \infty$ by (25) and so \bar{C} is not generically scip by (20).

Assume that the normalization of \bar{C} has m_∞ points lying over $\bar{C} \setminus X$. Then the kernel of the restriction map $\text{Pic}(\bar{C}) \rightarrow \text{Pic}(C)$ has rank $\leq m_\infty$, thus we still have $\text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(C)] = \infty$, hence C is not generically scip by (20). \square

We can use this to strengthen the results of [WK81].

Proposition 27. *Let S be a normal, projective surface over a field k . Then $|S|$ is homeomorphic to $\mathbb{P}_{\mathbb{F}_2}^2$ if and only if*

(27.1) *k is locally finite, and*

(27.2) *any 2 curves in S have a non-empty intersection.*

Proof. By (26) every irreducible curve in $\mathbb{P}_{\mathbb{F}_2}^2$ is scip, hence the same holds for S , so k is locally finite by (26). Any 2 curves in $\mathbb{P}_{\mathbb{F}_2}^2$ have a non-empty intersection, hence the same holds for S .

The converse, which is the combination of the next 2 claims, is essentially proved in [WK81].

Claim 27.3. Up to homeomorphism there is a unique countable, noetherian, 2-dimensional topological space X with the following properties.

(a) Every curve $C \subset X$ contains infinitely many points.

(b) Any 2 irreducible curves $C_1, C_2 \subset X$ have a non-empty intersection.

(c) Given any 1-dimensional, closed $D \subset X$ and a 0-dimensional, closed $P \subset D$ such that $P \cap D_i \neq \emptyset$ for every irreducible component $D_i \subset D$, there is an irreducible curve $C \subset X$ such that $D \cap C = P$.

Proof. Assume that we have 2 such X, X' . Choose well orderings of their points and curves Z_1, Z_2, \dots and Z'_1, Z'_2, \dots . Assume that we have a subset $I \subset \mathbb{N}$ and an injection $\phi_I : I \rightarrow \mathbb{N}$ such that ϕ_I induces a homeomorphism

$$\{Z_i : i \in I\} \longrightarrow \{Z'_{\phi_I(i)} : i \in I\}.$$

Set $j := \inf\{\mathbb{N} \setminus I\}$. We extend ϕ_I to $J := I \cup \{j\}$ as follows.

If Z_j is a point on a Z_i (resp. on none of them), we pick $Z'_{\phi_J(j)}$ to be any new point on $Z'_{\phi_I(i)}$ (resp. on none of them). If Z_j is a curve, then first we pick images of all the new points in $Z_j \cap Z_i$ for every $i \in I$ and then use (c) to pick $Z'_{\phi_J(j)}$.

Continuing this way would only give an injection $\phi_\infty : \mathbb{N} \rightarrow \mathbb{N}$. We fix this by alternating the construction between ϕ_I and ϕ_I^{-1} . \square

Claim 27.4. Let K be locally finite field and S a normal, projective surface over K . Then $|S|$ satisfies (27.3.a–c) iff any 2 irreducible curves $C_1, C_2 \subset S$ have a non-empty intersection.

Proof. We need to prove (27.c). First we blow up P and normalize to get $S_1 \rightarrow S$. Repeatedly blowing up points over P we get $S_r \rightarrow S$ such that the intersection matrix of D_r (the birational transform of D) is negative definite. By [Art62], $D_r \subset S_r$ can be contracted to get $\pi : S_r \rightarrow T$. By [CP16], there is an irreducible hypersurface section $C_T \subset T$ that is disjoint from $\pi(D_r)$. Let $C \subset S$ be its birational transform. \square

Lemma 28. *Let $p : X' \rightarrow X$ be a morphism between normal, projective varieties. Let $C' \subset X'$ be an irreducible curve. Set $C := p(C')$ and assume that $k(C')/k(C)$ is purely inseparable.*

Then C generically scip $\Rightarrow C'$ generically scip.

Proof. Let $\Sigma_X \subset X$ be a finite subset that contains all non-Cartier centers and such that $C' \setminus p^{-1}(\Sigma_X) \rightarrow C \setminus \Sigma_X$ is a bijection.

Pick any $q' \in C' \setminus p^{-1}(\Sigma_X)$ and set $q = p(q')$. There is a divisor $D(q)$ such that $C \cap D(q) = \{q\}$. Then $D(q)$ is Cartier, hence its pull-back gives a divisor $D(q')$ such that $C' \cap D(q') = \{q'\}$. \square

4. MORDELL-WEIL FIELDS

The Mordell-Weil theorem says that if A is an Abelian variety over a number field k then $A(k)$ is a finitely generated group. Our results are not sensitive to torsion in $A(k)$, this is why we need the concept of \mathbb{Q} -Mordell-Weil fields where $\text{rank}_{\mathbb{Q}} A(k)$ is always finite; see (151).

\mathbb{Q} -Mordell-Weil fields have a nice characterization involving complete intersections on curves.

Definition 29. Let X be a scheme and $C \subset X$ a curve.

(29.1) C is *scip with defect* $\Sigma \subset C$ if, for every closed, finite subset $P \subset C$, there is an effective divisor $D = D(C, P) \subset X$ such that $P \subset \text{Supp}(D \cap C) \subset P \cup \Sigma$.

(29.2) C is *scip with finite defect* if it is scip with defect Σ for some finite subset $\Sigma \subset C$.

It is clear that these depend only on the topological pair $|C| \subset |X|$.

Note that being scip with finite defect is invariant under birational maps that are isomorphisms at the generic point of C , we just need to add to Σ all the indeterminacy points that lie on C .

Let X, Y be irreducible varieties and $\pi : Y \rightarrow X$ a dominant, finite morphism. Let $C_X \subset X$ be a curve with reduced preimage $C_Y \subset Y$. If C_Y is scip with finite defect Σ_Y then C_X is scip with finite defect $\Sigma_X := \pi(\Sigma_Y)$.

Thus most questions about these notions can be reduced to normal, projective varieties.

Lemma 30. *Let X be a normal, projective variety over a perfect field k and $C \subset X$ an irreducible curve with normalization $\pi : \bar{C} \rightarrow C$. Then C is scip with finite defect iff*

$$\text{rank}_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^{\circ}(X) \rightarrow \mathbf{Jac}(\bar{C})](k) < \infty. \quad (30.1)$$

Proof. Note first that the difference between

$$\begin{aligned} & \text{rank}_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^{\circ}(X) \rightarrow \mathbf{Jac}(\bar{C})](k) \quad \text{and} \\ & \text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(\bar{C})] \end{aligned}$$

is at most $\text{rank}_{\mathbb{Q}} \text{NS}(X)$.

Assume that C is scip with finite defect. We may enlarge Σ_X to achieve that $C \setminus \Sigma_X$ is smooth and every Weil divisor on X not containing Σ_X is Cartier along $C \setminus \Sigma_X$ (139). Let the defect set be $\Sigma \subset C \setminus \Sigma_X$ and $\bar{\Sigma} \subset \bar{C}$ its preimage. Let $\Gamma \subset \text{Pic}(\bar{C})$ be the subgroup generated by all $\bar{q} \in \bar{\Sigma}$.

Pick any closed point $p \in C \setminus \Sigma_X$. By assumption we have an effective Cartier divisor D_p such that $p \in \text{Supp}(D_p \cap C) \subset \{p\} \cup \Sigma$. This shows that, for some $m > 0$,

$$m[p] \in \langle \Gamma, \text{im}[\text{Pic}(X) \rightarrow \text{Pic}(\bar{C})] \rangle.$$

Since these $\{[p] : p \in C \setminus \Sigma\}$ generate $\text{Pic}(\bar{C})$, we get that

$$\text{rank}_{\mathbb{Q}} \text{coker}[\text{Pic}(X) \rightarrow \text{Pic}(\bar{C})] \leq \text{rank}_{\mathbb{Q}} \Gamma.$$

Conversely, assume that (30.1) holds. Curve singularities can be resolved, thus we may assume that C is smooth. Then there is a finite subset $\{F_i : i \in I\} \subset \text{Jac}(\bar{C})$, closed under inverses, that generates $\text{coker}[\text{Pic}^{\circ}(X) \rightarrow \text{Jac}(\bar{C})]$ modulo torsion. Fix a point $p_0 \in C \setminus \Sigma$ and an ample line bundle L on X such that $\deg_C L = c \deg p_0$ for some $c > 0$. Set $d_0 = \deg p_0$ and choose r_1 such that $r_1[p_0] + F_i \sim G_i$, where the G_i are effective and disjoint from Σ_X .

Now pick any $p \in C \setminus \Sigma_X$ and set $d = \deg p$. The line bundle $\pi^* L^r (rc[p_0] + d[p_0] - d_0[p])$ is in $\text{Jac}(\bar{C})$. So, by assumption, there are nonnegative m_i and $T \in \text{Pic}^{\circ}(X)$ such that

$$\pi^* L^r (rc[p_0] + d[p_0] - d_0[p]) \sim_{\mathbb{Q}} \mathcal{O}_{\bar{C}}(\sum_i m_i F_i) \otimes T^{-1}.$$

We can rewrite this as

$$\pi^*(L^r \otimes T) \sim_{\mathbb{Q}} \mathcal{O}_{\bar{C}}(d_0[p] + (rc - d - r_1 \sum m_i)[p_0] + \sum_i m_i G_i)$$

Every section of $\pi^*(L^r \otimes T)|_C$ lifts back to a section of $L^r \otimes T$ for large enough r . Thus C is scip with defect $\Sigma = \text{Supp}(p_0 + \sum G_i)$. \square

Theorem 31. *Let X be an irreducible, quasi-projective variety of dimension ≥ 2 over a perfect field k . The following are equivalent.*

(31.1) *k is \mathbb{Q} -Mordell-Weil.*

(31.2) *Every irreducible curve $C \subset X$ is scip with finite defect.*

Proof. As we observed in (29), it is enough to prove (31.1) \Rightarrow (31.2) for normal, projective varieties. If k is \mathbb{Q} -Mordell-Weil then (31.2) holds for these by (30).

Conversely, if every irreducible curve $C \subset X$ is scip with finite defect then the same holds for the smooth locus $X^{\text{ns}} \subset X$. Let $X' \supset X^{\text{ns}}$ be a normal compactification. If $C' \subset X'$ has nonempty intersection with X^{ns} then C' is also scip with finite defect.

Now let A be an Abelian variety over k . By (35) there is an irreducible, projective curve $C' \subset X'$ and a \mathbb{Q} -injection

$$A \hookrightarrow_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^{\circ}(X') \rightarrow \mathbf{Jac}(\bar{C}')].$$

By (30) $\text{coker}[\mathbf{Pic}^{\circ}(X') \rightarrow \mathbf{Jac}(\bar{C}')](k)$ has finite \mathbb{Q} -rank, and so does $A(k)$. Thus k is \mathbb{Q} -Mordell-Weil. \square

By (30), being scip with finite defect depends on the interaction of $\mathbf{Pic}^{\circ}(X)$ and $\mathbf{Jac}(\bar{C})$. The following definition is designed to get rid of the influence of $\mathbf{Pic}^{\circ}(X)$.

Definition 32. Let X be a scheme, $C \subset X$ an irreducible curve. We say that C is *absolutely scip with finite defect* if the following holds.

(32.1) Let $C' \neq C$ be any irreducible curve. Then there are finite subsets $\Sigma \subset C$ and $\Sigma' \subset C'$ such that for every finite subset $P \subset C$ there is an effective divisor $D \subset X$ such that $P \subset \text{Supp}(D \cap (C \cup C')) \subset P \cup \Sigma \cup \Sigma'$.

Note that P is a subset of C only. This has the following effect.

Let $\Gamma' \subset \text{Jac}(\bar{C}')$ be the subgroup generated by the preimages of Σ' . Let $\Gamma'_X \subset \text{Pic}(X)$ be the preimage of Γ' under $\text{Pic}(X) \rightarrow \text{Jac}(\bar{C}')$. Then the class of D has to be in Γ'_X .

If C' is general ample curve then the kernel of $\text{Pic}(X) \rightarrow \text{Jac}(\bar{C}')$ is torsion, thus Γ'_X is a finitely generated group.

Now when we run the proof of (30) for $C \subset X$, instead of the whole $\text{Pic}(X)$, we have only Γ'_X to choose D from. The condition (30.1) now becomes

$$\text{rank}_{\mathbb{Q}} \text{coker}[\Gamma'_X \rightarrow \text{Pic}(\bar{C})] < \infty.$$

Since Γ'_X a finitely generated, this holds iff $\text{rank}_{\mathbb{Q}} \text{Pic}(\bar{C}) < \infty$, and we get the following.

Proposition 33. *Let X be a normal, projective variety of dimension ≥ 2 over a perfect field k and $C \subset X$ an irreducible curve. Then C is absolutely scip with finite defect iff $\text{Pic}(\bar{C})$ has finite \mathbb{Q} -rank.* \square

This is especially useful over fields where the opposite of the Mordell-Weil theorem happens, these are the *anti-Mordell-Weil* fields (152). For varieties over such fields we can recognize rational curves using only their set-theoretic intersection properties.

Putting together (33) with (152) gives the topological characterization of rational curves.

Corollary 34. *Let k be a perfect, anti-Mordell-Weil field, X an irreducible, quasi-projective k -variety of dimension ≥ 2 and $C \subset X$ an irreducible curve. Then C is absolutely scip with finite defect iff every irreducible component of $C_{\bar{k}}$ is rational.* \square

Lemma 35 (Curves with large Jacobians). *Let X be a geometrically normal, projective variety of dimension ≥ 2 and A an Abelian variety over k . Then there is an irreducible, projective curve $C \subset X^{\text{ns}}$ such that there is a \mathbb{Q} -injection (that is, with finite kernel)*

$$A \hookrightarrow_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^{\circ}(X) \rightarrow \mathbf{Jac}(\bar{C})].$$

Proof. Let $\bar{C} \subset A^{\vee} \times X$ be a curve that is a general, irreducible, complete intersection of ample divisors; we use [Poo08] in case k is finite. Let $C \subset X$ be the

image of the second coordinate projection $\pi : \bar{C} \rightarrow C$. Then $C \subset X^{\text{ns}}$ and \bar{C} is the normalization of C . (In fact, $\bar{C} \cong C$ if $\dim X \geq 3$ and k is infinite.) By (131) the natural map

$$A \times \mathbf{Pic}^\circ(X) = \mathbf{Pic}^\circ(A^\vee) \times \mathbf{Pic}^\circ(X) \rightarrow \mathbf{Jac}(\bar{C})$$

is \mathbb{Q} -injective, hence we have a \mathbb{Q} -injection $A \hookrightarrow_{\mathbb{Q}} \text{coker}[\mathbf{Pic}^\circ(X) \rightarrow \mathbf{Jac}(\bar{C})]$. \square

5. REDUCIBLE SCIP SUBSETS

Definition 36. Let X be a scheme and $Z \subset X$ a closed subset. We say that Z is *scip* if the following holds.

(36.1) Let $D_Z \subset Z$ be a closed subset of pure codimension 1 that has nonempty intersection with every irreducible component of Z . Then there is an effective divisor $D_X \subset X$ such that $\text{Supp}(D_X \cap Z) = D_Z$.

We say that Z is *generically scip* if the following holds.

(36.2) There is a (not necessarily closed) finite subset $\Sigma_Z \subset Z$ such that, if D_Z in (36.1) is disjoint from Σ_Z , then, for every (not necessarily closed) finite subset $\Sigma_X \subset X \setminus D_Z$, we can find $D_X \subset X$ as in (36.1) that is also disjoint from Σ_X .

It is clear that these depend only on the topological pair $|Z| \subset |X|$. Also, if (36.2) holds for some Σ_Z then it also holds for every larger $\Sigma'_Z \supset \Sigma_Z$. We usually just take the union $\Sigma := \Sigma_Z \cup \Sigma_X$ large enough.

If Z is scip (resp. generically scip) then any union of its irreducible components is also scip (resp. generically scip).

Example 37. In \mathbb{P}^n with coordinates x_0, \dots, x_n , set $L_1 = (x_1 = \dots = x_i = 0)$ and $L_2 = (x_{i+1} = \dots = x_n = 0)$. We claim that $L_1 \cup L_2$ is generically scip. Indeed, given divisors $D_{Z_i} \subset L_i$ not containing $L_1 \cap L_2 = (1:0:\dots:0)$, they can be given by equations

$$D_{Z_1} = (g_1(x_0, x_{i+1}, \dots, x_n) = 0) \quad \text{and} \quad D_{Z_2} = (g_2(x_0, x_1, \dots, x_i) = 0).$$

We may assume that $g_i(1, 0, \dots, 0) = 1$. Then

$$D_X := (g_1^{\deg g_2} + g_2^{\deg g_1} = x_0^{\deg g_1 \deg g_2})$$

satisfies $\text{Supp}(D_X \cap (L_1 \cup L_2)) = D_{Z_1} \cup D_{Z_2}$.

The following strong converse also illustrates the big difference between fields of characteristic 0, fields of positive characteristic and subfields of $\bar{\mathbb{F}}_p$.

Claim 37.1. Let k be a field, $D \subset \mathbb{P}_k^n$ an irreducible divisor and $C \not\subset D$ an irreducible, ample-sci (117) curve. Then $C \cup D$ is scip (resp. generically scip) iff one of the following holds.

- (a) $\text{char } k = 0$, C is a line and D is a hyperplane.
- (b) $\text{char } k > 0$, $\text{Supp}(C \cap D)$ is a single k^{ins} -point and C, D are both scip (resp. generically scip).
- (c) k is locally finite and D is scip (resp. generically scip).

Proof. Assume first that $\text{char } k = 0$. By (40) $(C \cdot D) = 1$, so $\deg C = 1$ and $\deg D = 1$. If $\text{char } k > 0$ but k is not locally finite, then $\text{Supp}(C \cap D)$ is a single k^{ins} -point by (39.1). If $C \cup D$ is scip (resp. generically scip) then C and D are both scip (resp. generically scip). This shows that the conditions of (b) are necessary. Their sufficiency also follows from (39). \square

Note that if C is smooth and rational, then it is scip. If $n \geq 4$ and D is smooth then it is scip. Let $D \subset \mathbb{P}^n$ be a smooth hypersurface and $n \geq 3$. If $\deg D \leq n$ then there are lots of smooth rational curves that meet D in 1 point only. If $\deg D \geq n + 2$ then there should be few such curves, but there are examples of arbitrary large degree.

Next we prove a general result about reducible scip subschemes.

Notation 38. For a k -scheme Y we use $k[Y] := H^0(Y, \mathcal{O}_Y)$ to denote the ring of regular functions. If Y is normal and proper then $k[Y] = k$ iff Y is geometrically integral.

If Y is reduced then $k(Y)$ denotes the ring of rational functions. Y is irreducible iff $k(Y)$ is a field. If Y_i are the irreducible components of Y then $k(Y) \cong \bigoplus_i k(Y_i)$.

Proposition 39. *Let X be a normal, projective k -variety such that $\rho(X) = 1$. Let $Z, W \subset X$ be reduced, irreducible subvarieties such that $Z \cap W$ is 0-dimensional. Assume that k is not locally finite. Then $Z \cup W$ is generically scip (36) iff the following hold.*

- (39.1) Z and W are generically scip,
- (39.2) $Z \cap W$ is irreducible,
- (39.3) either $k[\text{red}(Z \cap W)]/k[W]$ or $k[\text{red}(Z \cap W)]/k[Z]$ is purely inseparable, and
- (39.4) if $\text{char } k = 0$ then $Z \cap W$ is reduced.

Proof. Assume first that $Z \cup W$ is generically scip. Choose any Σ that contains $\Sigma(Z \cup W)$ (54.3) and the non-Cartier centers of X (136). Let L be an ample line bundle on X such that $H^0(X, L) \rightarrow H^0(Z \cap W, L_{Z \cap W})$ is surjective. Choose sections $s_Z, s_W \in H^0(X, L)$ that are nowhere zero on Σ . Write $(s_Z|_Z = 0) = \sum_i a_i A_i$ and $(s_W|_W = 0) = \sum_j b_j B_j$. By assumption, for every i, j there is a divisor $D_{ij} \subset X$ such that $D_{ij}|_{Z \cup W} = c_{ij} A_i + d_{ij} B_j$ for some $c_{ij}, d_{ij} > 0$. The D_{ij} are Cartier since they are disjoint from Σ .

For each j a suitable positive linear combination of the D_{ij} gives a divisor D_j such that $D_j|_Z$ is a multiple of $(s_Z|_Z = 0)$ and $D_j|_W$ is a multiple of B_j . Then we can take a suitable positive linear combination D of the D_j such that $D|_Z$ is a multiple of $(s_Z|_Z = 0)$ and $D|_W$ is a multiple of $(s_W|_W = 0)$.

Since $\rho(X) = 1$, after passing to a suitable power we may assume that

$$D = (s = 0) \quad \text{for some } s \in H^0(X, L^m) \quad \text{and } m > 0. \quad (39.5)$$

As we check in (54.4), this implies that

$$\begin{aligned} s|_Z &= u_Z s_Z^m|_Z && \text{for some } u_Z \in k[Z]^\times \quad \text{and} \\ s|_W &= u_W s_W^m|_W && \text{for some } u_W \in k[W]^\times, \end{aligned} \quad (39.6)$$

and hence

$$(s_Z/s_W)^m|_{Z \cap W} = u_W|_{Z \cap W} \cdot u_Z^{-1}|_{Z \cap W} \in \text{im}[k[W]^\times \times k[Z]^\times \rightarrow k[Z \cap W]^\times]. \quad (39.7)$$

We can arrange s_Z/s_W to be an arbitrary element of $k[Z \cap W]^\times$, hence we conclude that

$$k[Z \cap W]^\times / k[W]^\times \times k[Z]^\times \quad \text{is a torsion group.} \quad (39.8)$$

Now (149) shows that either $k[\text{red}(Z \cap W)]/k[W]$ or $k[\text{red}(Z \cap W)]/k[Z]$ is purely inseparable, proving (39.3). Since Z, W are irreducible, $k[Z]$ and $k[W]$ are finite field extensions of k . Thus $k[\text{red}(Z \cap W)]$ is a finite field extension of k , hence $Z \cap W$ is irreducible. Finally (39.4) follows from (147).

Conversely, assume that (39.1–4) hold and let A_Z and B_W be effective divisors on Z and W that are disjoint from Σ . By (39.1) they are both restrictions of Cartier divisors from X . Since $\rho(X) = 1$, there is a power L^m and sections

$$\sigma_Z \in H^0(Z, L^m|_Z) \quad \text{and} \quad \sigma_W \in H^0(W, L^m|_W).$$

As in (39.4–5) we see that suitable powers of σ_Z^r and σ_W^r can be glued to a section of $\sigma_{Z \cup W} \in H^0(Z \cap W, L^{mr}|_{Z \cup W})$ iff

$$\sigma_Z \sigma_W^{-1} \in k[Z \cap W]^\times / k[W]^\times \times k[Z]^\times \quad \text{is torsion.} \quad (39.8)$$

This is guaranteed by (39.3–4) using (149).

Once $A_Z \cup B_W$ is defined as the zero set of a section $\sigma_{Z \cup W}$, we can lift (a possibly higher power of) it to a section $\sigma_X \in H^0(X, L^N)$, and $D_X := \text{Supp}(\sigma_X = 0)$ shows that $Z \cup W$ is generically scip. \square

Corollary 40. *Let X be a smooth, projective k -variety such that $\rho(X) = 1$ and $\text{char } k = 0$. Let $C \subset X$ be a geometrically connected curve and $D \subset X$ a divisor. If $C \cup D$ is generically scip then $(C \cdot D) = 1$. \square*

Looking at the above proof shows that there should be even fewer generically scip reducible subsets if $\rho(X) > 1$, but for now we have the following slightly weaker result.

Proposition 41. *Let X be a normal, projective k -variety where k is not locally finite. Let $Z, W \subset X$ be reduced, irreducible subvarieties such that $Z \cap W$ is 0-dimensional. Assume that $Z \cup W$ is generically scip (36). Then*

(41.1) $Z \cap W$ is irreducible,

(41.2) either $k[\text{red}(Z \cap W)]/k[W]$ or $k[\text{red}(Z \cap W)]/k[Z]$ is purely inseparable, and

(41.3) if $\text{char } k = 0$ then

$$\dim_k \ker [k[Z \cap W] \rightarrow k[\text{red}(Z \cap W)]] \leq \frac{\rho(X) - 1}{\deg[k : \mathbb{Q}]}.$$

Proof. Choose Σ to contain $\Sigma(Z \cup W)$ and the non-Cartier centers of X (136). As in (125), let $\text{WDiv}(X, \Sigma)$ denote the group of Weil divisors whose support is disjoint from Σ . These are all Cartier by our choice of Σ . We get restriction maps $r_Z : \text{WDiv}(X, \Sigma) \rightarrow \text{WDiv}(Z, \Sigma)$ and $r_W : \text{WDiv}(X, \Sigma) \rightarrow \text{WDiv}(W, \Sigma)$. These descend to maps of the Picard groups $\bar{r}_Z : \text{Pic}(X) \rightarrow \text{Pic}(Z)$ and $\bar{r}_W : \text{Pic}(X) \rightarrow \text{Pic}(W)$, which do not depend on Σ . Set $K_Z(X) := \ker \bar{r}_Z$, $K_W(X) := \ker \bar{r}_W$ and $K_{ZW}(X) = K_Z(X) \cap K_W(X)$.

As in (125), the kernels of \bar{r}_Z and \bar{r}_W define closed subgroups $\mathbf{K}_Z(X) \subset \mathbf{Pic}(X)$ and $\mathbf{K}_W(X) \subset \mathbf{Pic}(X)$. Their intersection is denoted by $\mathbf{K}_{ZW}(X)$.

Let B be a divisor in $\text{WDiv}(X, \Sigma)$ whose class $[B]$ lies in $K_{ZW}(X)$. Then $B|_Z = (s_Z)$, where s_Z is unique up to $k[Z]^\times$, and $B|_W = (s_W)$, where s_W is unique up to $k[W]^\times$; here we use that $\Sigma \supset \Sigma(Z \cup W)$ and (54). Restricting both to $Z \cap W$ we get

$$s_Z|_{Z \cap W} \cdot s_W^{-1}|_{Z \cap W} \in k[Z \cap W]^\times / (k[W]^\times \times k[Z]^\times),$$

which defines a homomorphism

$$K_{ZW}(X) \rightarrow k[Z \cap W]^\times / (k[W]^\times \times k[Z]^\times).$$

As in (145), we get a homomorphism of algebraic groups

$$\partial_{ZW} : \mathbf{K}_{ZW}(X) \rightarrow (\mathcal{R}_k^{Z \cap W} \mathbb{G}_m) / (\mathcal{R}_k^W \mathbb{G}_m \times \mathcal{R}_k^Z \mathbb{G}_m).$$

Note that $\mathbf{K}_{ZW}(X, \Sigma) \cap \mathbf{Pic}^\circ(X)$ is an Abelian variety (128.7), hence a positive dimensional subgroup of it has no morphisms to a linear algebraic group. Set

$$\mathrm{NS}_{ZW}(X) := K_{ZW}(X) / (K_{ZW}(X) \cap \mathbf{Pic}^\circ(X)(k)).$$

Thus ∂_{ZW} factors through

$$\mathrm{NS}_{ZW}(X) \rightarrow (\mathcal{R}_k^{Z \cap W} \mathbb{G}_m) / (\mathcal{R}_k^W \mathbb{G}_m \times \mathcal{R}_k^Z \mathbb{G}_m).$$

Let $\Gamma_{ZW}(X)$ denote its image.

By the above, $\Gamma_{ZW}(X)$ is the image of a subgroup of $\ker[\mathrm{NS}(X) \rightarrow \mathrm{NS}(Z)]$ (modulo torsion). All we need from this is that

$$\mathrm{rank}_{\mathbb{Q}} \Gamma_{ZW}(X) \leq \rho(X) - 1.$$

Now we start to follow the proof of (39). The departure from it happens at (39.5), where now σ is not a section of L^m , but of some $L^m(B)$ for some $B \in K_{ZW}(X)$. Thus we conclude that

$$(s/t)^m|_{Z \cap W} = u_W|_{Z \cap W} \cdot u_Z^{-1}|_{Z \cap W} \cdot \gamma \in \mathrm{im}[k[W]^\times \times k[Z]^\times \rightarrow k[Z \cap W]^\times], \quad (41.3)$$

for some $\gamma \in \Gamma_{ZW}(X)$. We can arrange s/t to be an arbitrary element of $k[Z \cap W]^\times$, hence we conclude that

$$k[Z \cap W]^\times / k[W]^\times \times k[Z]^\times \times \Gamma_{ZW}(X) \quad \text{is a torsion group.} \quad (41.4)$$

Now we use (149) to get that $Z \cap W$ is irreducible and (147) implies (41.2). Finally we get that

$$\mathrm{rank}_{\mathbb{Q}} \ker[k[Z \cap W] \rightarrow k[\mathrm{red}(Z \cap W)]] \leq \mathrm{rank}_{\mathbb{Q}} \Gamma_{ZW}(X) \leq \rho(X) - 1.$$

Thus (41.3) follows from (141.3). \square

In characteristic 0 we can reformulate the bound of (41) as follows.

Corollary 42. *Let X be a normal, projective variety over a field k of characteristic 0. Let $Z, W \subset X$ be reduced, irreducible subvarieties such that $Z \cap W$ is 0-dimensional. Assume that $Z \cup W$ is generically scip (36). Then $Z \cap W$ is irreducible and*

$$\dim_k k[Z \cap W] \leq \max\{\dim_k k[Z], \dim_k k[W]\} + \frac{\rho(X) - 1}{\deg[k : \mathbb{Q}]}. \quad \square$$

Example 43. Combining the ideas of (37) with (39) we get a method to recognize k -points. The assumptions are restrictive, but this gives the first indication that one can get detailed scheme-theoretic information from the topology. However, scip and generically scip turn out to be too restrictive in general; searching for a more flexible variant lead to the notion of linkage in Section 9.

Claim 43.1. Let X be a smooth, projective k -variety of dimension ≥ 7 such that $\rho(X) = 1$. Then $p \in X$ is a k -point iff there are 3-dimensional, set-theoretic complete intersections $Z, W \subset X$ such that

$$(43.1) \quad \mathrm{Supp}(Z \cap W) = \{p\} \text{ and}$$

$$(43.2) \quad Z \cup W \text{ is generically scip.}$$

Proof. Assume that $p \in X$ is a k -point and let $Z, W \subset X$ be 3-dimensional, smooth, complete intersections such that $Z \cap W = \{p\}$ as schemes. Lefschetz theorem tells us that if $D_Z \subset Z$ is any divisor then (some multiple of) it is a complete intersection. Arguing as in (37) we see that $Z \cup W$ is generically scip.

Conversely, $k[Z] = k[W] = k$ since Z, W are set-theoretic complete intersections (119.1), thus (39) says that $p \in X(k)$. \square

Note that the bound $\dim X \geq 7$ can be improved to $\dim X \geq 5$ if the Noether-Lefschetz theorem applies over k ; see (13) for such cases.

6. PROJECTIVE SPACES

We study the scip property for the union of a curve and of a divisor. As we observed in Section 5, this happens very rarely, and it leads to the following stronger version of (1)

Theorem 44. *Let L be a field of characteristic 0 and K an arbitrary field. Let Y_L be a normal, projective, geometrically irreducible L -variety of dimension $n \geq 2$ and $\Phi : |\mathbb{P}_K^n| \sim |Y_L|$ a homeomorphism. Then*

$$(44.1) \quad Y_L \cong \mathbb{P}_L^n,$$

$$(44.2) \quad K \cong L, \text{ and}$$

$$(44.3) \quad \Phi \text{ is the composite of a field isomorphism } \phi : K \cong L \text{ and of an automorphism of } \mathbb{P}_K^n.$$

We start with an easy to prove but interesting special case of (44).

45 (Proof of (44) when $Y_L \cong \mathbb{P}_L^n$). Let $H \subset \mathbb{P}_K^n$ be a hyperplane and $\ell \subset \mathbb{P}_K^n$ a line not contained in H . Then $\ell \cup H$ is scip by (36.2) hence so is $\Phi(\ell) \cup \Phi(H) \subset \mathbb{P}_L^n$. So $\Phi(H) \subset \mathbb{P}_L^n$ is a hyperplane by (37.1.a). By taking intersections, we see that Φ gives an isomorphism of the projective geometries $K\mathbb{P}^n$ and $L\mathbb{P}^n$. By the Veblen-Young theorem [VY08] this is induced by a field isomorphism $\phi : K \cong L$.

Composing Φ with the natural isomorphism induced by ϕ^{-1} , we get a homeomorphism $\Psi : |\mathbb{P}_K^n| \rightarrow |\mathbb{P}_K^n|$ that is the identity on K -points. It remains to show that it is the identity on all points. Let $C \subset \mathbb{P}_K^n$ be a K -rational curve. It has infinitely many K -points and these are fixed by Ψ . Thus $C \cap \Psi(C)$ is infinite, hence $C = \Psi(C)$. However, we do not yet know that $\Psi|_C$ is the identity.

Let $p \in \mathbb{P}_K^n$ be a closed point. Assume that there are K -rational curves $C_\lambda \subset \mathbb{P}_K^n$ such that $\{p\} = \cap C_\lambda$. Then $\{\Psi(p)\} = \cap \Psi(C_\lambda) = \cap C_\lambda = \{p\}$, as needed.

It remains to construct such curves C_λ . For this we can work in an affine chart $p \in \mathbb{A}_K^n \subset \mathbb{P}_K^n$ with coordinates x_i . Note that $K(p)/K$ is a finite, separable extension, hence can be generated by a single element $z_p \in K(p)$. We can thus write $x_i(p) = h_i(z_p)$ for some $h_i \in K[t]$.

Let $g(t) \in K[t]$ be the minimal polynomial of z_p ; we can then identify z_p with a root of g in \bar{K} .

For $i \in \{1, \dots, n\}$ and $a \in K$ let $C_{i,a}$ be the image of

$$\tau_{i,a} : t \mapsto (h_1(t), h_2(t), \dots, h_n(t)) + ag(t)e_i,$$

where e_i is the i th standard basis vector.

The $C_{i,a}$ are K -rational curves, hence stabilized by Φ .

Claim 45.1. $\cap_{i,a} C_{i,a} = \{p\}$.

Proof. First, the $\tau_{i,a}$ all map $z_p \in \mathbb{A}^1(\bar{K})$ to p , so $p \in \cap_{i,a} C_{i,a}$. To see the converse, assume that $p \neq q \in \cap_{i,a} C_{i,a}$. After permuting the coordinates we may assume that $p_n \neq q_n$. If $q = \tau_{1,a}(z')$ then $h_n(z') = q_n$. The equation $h_n(*) = q_n$ has finitely many solutions z'_j and they are all different from z_p . Then, for all but finitely many $a \in K$, $h_1(z'_i) + ag(z'_i) \neq q_i$ for every i . Thus $q \notin C_{1,a}$. \square

In positive characteristic the above proof and (39) give the following.

Claim 45.2. Let K, L be perfect fields that are not locally finite, $n \geq 2$ and $\Phi : |\mathbb{P}_K^n| \sim |\mathbb{P}_L^n|$ a homeomorphism. Then Φ induces a bijection $\mathbb{P}^n(K) \leftrightarrow \mathbb{P}^n(L)$. \square

For the proof of (44), the key step is the following.

Lemma 46. *Using the notation and assumptions of (44), let $|H|$ denote the linear system of all hyperplanes in \mathbb{P}_K^n . Then $\{\Phi(H) : H \in |H|(K)\}$ is a bounded family of divisors on Y_L .*

47 (Proof of (44) assuming (46)). First note that K is not locally finite by (26).

Set $Z := \Phi^{-1}(\text{Sing}(Y_L))$. Let $|D| \subset |H|$ be a pencil of hyperplanes whose base locus is not contained in Z , and $\{D_\lambda : \lambda \in \Lambda\}$ the corresponding t-pencil (105). Thus $\{\Phi(D_\lambda) : \lambda \in \Lambda\}$ is a t-pencil on Y_L .

There are infinitely many hyperplanes among the $\{D_\lambda : \lambda \in \Lambda\}$ and their Φ -images form a bounded family of divisors by (46). Thus $\{\Phi(D_\lambda) : \lambda \in \Lambda\}$ is algebraic (107), linear (109) and the images of the K -hyperplanes are true members (113). Thus, by (48), the $\{\Phi(H) : H \in |H|(K)\}$ span an n -dimensional linear system $|H|^Y$, which is basepoint-free since already the $\{\Phi(H) : H \in |H|(K)\}$ have no point in common. So $|H|^Y$ gives a morphism $g : Y \rightarrow \mathbb{P}_L^n$. Since any hyperplane H has nonempty intersection with every curve, the same holds for $\Phi(H)$, so $g : Y \rightarrow \mathbb{P}_L^n$ is finite and $|H|^Y$ is ample.

Therefore members of $|H|^Y$ are geometrically connected, and so are the Φ -images of the lines, since they are set-theoretic complete intersections of members of $|H|^Y$ (119.1). By (39) this shows that $\text{Supp}(\Phi(\ell) \cap \Phi(H))$ is an L -point whenever it is a smooth point of Y_L . We also obtain this point as $\Phi(H_1) \cap \dots \cap \Phi(H_n)$, or as a fiber of $g : Y \rightarrow \mathbb{P}_L^n$. Since $\text{char } L = 0$, general fibers of g are reduced. Thus $g : Y \rightarrow \mathbb{P}_L^n$ is finite and of degree 1, hence an isomorphism. The rest now follows from (45). \square

The following lemma, which essentially says that pencils determine higher dimensional linear systems, is longer to state than to prove.

Lemma 48. *Let Y be a normal, projective variety over a field L . Let K be an infinite field and $e_0, \dots, e_n \in K\mathbb{P}^n$ independent points. Assume that we have a map*

$$\Phi : K\mathbb{P}^n \rightarrow (\text{effective Weil divisors on } Y)$$

with the following property.

(48.1) *For $r = 1, \dots, n$ there are Zariski open subsets $\emptyset \neq U_{r-1} \subset \langle e_0, \dots, e_{r-1} \rangle$ such that, for every $p \in U_{r-1}$, the divisors $\{\Phi(q) : q \in \langle p, e_r \rangle\}$ are L -members of a linear pencil on Y .*

Then there is a Zariski open subset $W \subset K\mathbb{P}^n$ such that the divisors

$$\{\Phi(q) : q \in W\}$$

are L -members of a linear system of dimension $\leq n$ on Y . \square

49 (Proof of (46) when $\dim Y = 2$). Let ℓ, ℓ' be 2 lines in \mathbb{P}_K^2 . We aim to show that $(\Phi(\ell) \cdot \Phi(\ell'))$ is bounded. If $\Phi(\ell) \cap \Phi(\ell')$ is a smooth point of Y , then

$$(\Phi(\ell) \cdot \Phi(\ell')) = \dim_k k[\Phi(\ell) \cap \Phi(\ell')],$$

and, by (42), the latter is bounded by

$$\max\{\dim_k k[\Phi(\ell)], \dim_k k[\Phi(\ell')]\} + \frac{\rho(Y) - 1}{\deg[L : \mathbb{Q}]}.$$

If $C \subset Y$ is any integral curve then $\dim_k k[C]$ is the number of connected components of $C_{\bar{k}}$. By (53) then either $\dim_k k[C] < \rho(Y_{\bar{k}})$, or C is contained in members of a basepoint-free pencil. In the latter case it is disjoint from other members of the pencil, which does not happen for $\Phi(\ell)$.

Thus we conclude that $(\Phi(\ell) \cdot \Phi(\ell'))$ is bounded, except possibly for lines that meet at a singular point of Y .

We check in (50) that almost all $\Phi(\ell)$ have positive self-intersection number. Since $\Phi(\ell)$ has positive intersection number with every other curve, we see that almost all $\Phi(\ell)$ are ample. A general such $\Phi(\ell)$ is contained in the smooth locus of Y , hence it has bounded intersection number with every other $\Phi(\ell')$. Hence the $\Phi(\ell)$ form a bounded family. \square

Lemma 50. *Let S be a normal, projective surface and $\{C_i : i \in I\}$ an infinite collection of irreducible curves on S such that the intersection numbers $\{(C_{i_1} \cdot C_{i_2}) : i_1 \neq i_2 \in I\}$ are positive and bounded from above. Then $(C_i^2) > 0$ for almost all $i \in I$.*

Proof. Set $d_1 := \sup\{(C_{i_1} \cdot C_{i_2}) : i_1 \neq i_2 \in I\}$. First we claim that the self-intersections $(C_i^2) : i \in I$ are bounded from below by some $-d_2 < 0$. If not then for every r there is a subset $J_r \subset I$ of r elements such that $(C_j^2) < -rd_1$ for every $j \in J_r$. Then the intersection matrix of the curves $\{C_j : j \in J_r\}$ is negative definite. This is only possible for $r < \rho^{\text{cl}}(S)$ (124).

Note that the local intersection numbers at a point $s \in S$ are bounded from below by some $\beta(s) > 0$ (the inverse of the determinant of the intersection matrix of the exceptional curves on a resolution). So there is a $\beta > 0$ such that $(C_{i_1} \cdot C_{i_2}) \geq \beta$ for every $i_1 \neq i_2 \in I$.

Let $J \subset I$ be any finite subset of $> 1 + d_2/\beta$ elements. Then $D := \sum_{j \in J} C_j$ is nef and big. Since $(D \cdot C_i) \leq d_1|J|$ for every $i \in I \setminus J$, these C_i are contained in a family of curves parametrized by a k -scheme of finite type W by (51). The family of curves parametrized by W contains only finitely many curves with negative self-intersection, since these are isolated points of W . Also, W contains only finitely many basepoint-free pencils, and $\{C_i : i \in I\}$ can contain at most one member of each such pencil since $(C_{i_1} \cdot C_{i_2}) > 0$ for $i_1 \neq i_2 \in I$. Thus $(C_i^2) > 0$ for all but finitely many $i \in I$. \square

Lemma 51. *Let S be a normal, projective surface and M a big \mathbb{Q} -divisor on S . Then for every m , the set of curves*

$$\{C : C \text{ is irreducible and } (M \cdot C) \leq m\}$$

is a bounded family.

Proof. Note that on a normal surface the intersection numbers make sense even for divisors that are not \mathbb{Q} -Cartier. After a resolution and pull-back, we may assume that S is smooth. By Kodaira's lemma [KM98, 2.60] we can write $M \sim_{\mathbb{Q}} A + E$ where A is ample and E is effective. Thus either $C \subset E$ or $(A \cdot C) \leq m$. \square

52 (Proof of (46) when $\dim Y \geq 3$). In this case the $\Phi(H)$ are ample by (67), hence geometrically connected, and so are the Φ -images of the lines since they are set-theoretic complete intersections of members of $|H|^Y$. Thus

$$\dim_k [\Phi(\ell) \cap \Phi(H)] \leq 1 + \frac{\rho(Y) - 1}{\deg[k : \mathbb{Q}]} \quad (52.1)$$

by (42). If $\Phi(\ell) \cap \Phi(H)$ is a smooth point of Y , then

$$(\Phi(\ell) \cdot \Phi(H)) = \dim_k [\Phi(\ell) \cap \Phi(H)].$$

So the $\Phi(H)$ have bounded intersection number with a curve that is an intersection of ample divisors, hence they form a bounded family. \square

The following result is proved in [BPS16], sharpening earlier versions of [Tot00, Per06].

Theorem 53. *Let X be a normal, projective, irreducible k -variety and $\{D_i : i \in I\}$ pairwise disjoint divisors. Then*

$$(53.1) \text{ either } |I| \leq \rho^{\text{cl}}(X) - 1,$$

$$(53.2) \text{ or all the } D_i \text{ are contained in members of a basepoint-free pencil.} \quad \square$$

7. SECTIONS AND THEIR ZERO SETS

We discuss foundational results about sections and their zero sets that are needed in our study of linkage.

54. Let X be a normal, geometrically integral, proper k -variety, L a line bundle on X and $s_1, s_2 \in H^0(X, L)$ sections of L with corresponding divisors ($s_i = 0$). Then $(s_1 = 0) = (s_2 = 0)$ iff $s_1 = s_2 \cdot c$ for some $c \in k^\times$. Our aim is to relax normality as much as possible and still keep the conclusion.

Let Y be a reduced scheme, L a line bundle on Y and $s \in H^0(Y, L)$ a section. It has *scheme-theoretic zeros* ($s = 0$) and *divisor-theoretic zeros*; the latter is the Weil divisor $\sum_{\eta} \text{length}_{k(\eta)}(L/s\mathcal{O}_Y)[\bar{\eta}]$, where the summation is over all codimension 1 points of Y . The scheme-theoretic zeros determine the divisor-theoretic zeros, but the converse does not always hold.

We consider 2 genericity conditions.

$$(54.1) \text{ Every generic point of } \text{Supp}(s = 0) \text{ is a regular, codimension 1 point of } Y.$$

$$(54.2) \text{ } Y \text{ is } S_2 \text{ along } \text{Supp}(s = 0).$$

If (54.1) holds then $(s = 0)$ is well-defined as a Weil divisor that is generically Cartier. If (54.2) holds then the zero set $(s = 0)$ has no embedded points. Thus if (54.1–2) both hold then the divisor-theoretic zeros determine the scheme-theoretic zeros. In these cases we do not distinguish the 2 concepts and simply talk about the *zero set*, and denote it by $(s = 0)$.

Both of these conditions hold, except at some special points.

Definition 54.3. For a reduced scheme Y , let $\Sigma(Y) \subset Y$ denote the set of points $y \in Y$ such that $\mathcal{O}_{y,Y}$ is either of dimension 0, or of dimension 1 but not regular, or not S_2 .

If Y is of finite type then $\Sigma(Y)$ is finite. Note that s satisfies the conditions (54.1–2) iff $(s = 0)$ is disjoint from $\Sigma(Y)$.

The usual correspondence between divisors and sections holds outside $\Sigma(Y)$.

Claim 54.4. Let Y be a reduced scheme, L a line bundle on Y and $s_1, s_2 \in H^0(Y, L)$ sections that do not vanish at any point of $\Sigma(Y)$. Then $(s_1 = 0) = (s_2 = 0)$ iff $s_1 = s_2 \cdot u$ for some $H^0(Y, \mathcal{O}_Y)^\times$. \square

Example 54.5. Let $Y \subset \mathbb{P}_k^4$ be the union of $(x_1 = x_2 = 0)$ and of $(x_3 = x_4 = 0)$. Note that $(1:0:0:0)$ is a non- S_2 point and $H^0(Y, \mathcal{O}_Y) = k$.

Consider $s(a, c) = ax_1 + cx_3 \in H^0(Y, \mathcal{O}_Y(1))$. Note that its divisor ($s(a, c) = 0$) is independent of $a, c \in k^\times$. However, $s(a, c) = s(a', c') \cdot u$ for some $H^0(Y, \mathcal{O}_Y)^\times$ iff $a/c = a'/c'$.

Definition 55. Let Y be a reduced scheme, $B \subsetneq Y$ a closed subset and L a line bundle on Y . For $m > 0$ set

$$\Gamma^B(Y, L, m) := \{s \in H^0(Y, L^m) : \text{Supp}(s = 0) = B\}. \quad (55.1)$$

We define analogously $\Gamma^{cB}(Y, L, m)$,

$$\Gamma^B(Y, L) := \bigoplus_m \Gamma^B(Y, L^m) \quad \text{and} \quad \Gamma^{cB}(Y, L) := \bigoplus_m \Gamma^{cB}(Y, L, m). \quad (55.2)$$

These are all unions of $k[Y]^\times$ -orbits (38).

Note that, in view of (64) and (68), $\Gamma^B(Y, L)$, is a very natural object to consider for us.

Lemma 56. *Let Y be a reduced, projective scheme, $B \subset Y$ a closed subset that is disjoint from $\Sigma(Y)$, and L a line bundle on Y .*

(56.1) *If B is irreducible then $\Gamma^B(Y, L, m)$ consists of at most one $k[Y]^\times$ -orbit.*

(56.2) *$\Gamma^{cB}(Y, L, m)/k[Y]^\times$ is finite.*

(56.3) *$\Gamma^{cB}(Y, L)/k[Y]^\times$ is a sub-semigroup of a finitely generated group.*

Proof. The first claim follows from (54.4). To see the other 2, let $B_i \subset B$ be the irreducible, divisorial components. By (54.4) $\Gamma^{cB}(Y, L)/k[Y]^\times \subset \bigoplus_i \mathbb{Z}B_i$, proving (56.3). If $s \in \Gamma^{cB}(Y, L, m)$ and $(s = 0) = \sum_i m_i B_i$ then, computing the degrees (with respect to some ample divisor) gives that $\sum_i m_i \deg B_i = \deg D$, hence $m_i \leq \deg D$ for every i and (56.2) holds. \square

Next we look at the evaluation of a section of a line bundle L at a point or at a 0-dimensional subscheme V . The twist is that we can not distinguish 2 sections if their zero sets have the same support, and we also can not distinguish various powers of L from each other. Thus for us the outcome of evaluation is not a single element of $H^0(V, \mathcal{O}_V) \otimes L$, but a subsemigroup of $H^0(V, \mathcal{O}_V) \otimes \bigoplus_m L^m$. Our aim is then to understand when this subsemigroup is small.

Definition 57. Let X be a normal, projective, irreducible k -variety, L a line bundle and D an effective divisor on X . Let $W \subset X$ be a closed, integral subvariety and $V \subset W \setminus D$ a 0-dimensional subscheme. Set

$$\begin{aligned} \mathcal{R}_V^W(D, L, m) &:= \text{im}[\Gamma^{W \cap D}(W, L|_W, m) \rightarrow H^0(V, L^m|_V)] \quad \text{and} \\ \mathcal{R}_V^W(D, L) &:= \bigoplus_{m > 0} \mathcal{R}_V^W(D, L, m). \end{aligned}$$

Note that $\mathcal{R}_V^W(D, L)$ is a semigroup that is closed under multiplication by $k[W]^\times = H^0(W, \mathcal{O}_W)^\times$ and, if $D \cap \Sigma(W) = \emptyset$, then $\mathcal{R}_V^W(D, L)/k[W]^\times$ is a subsemigroup of a finitely generated group by (56.3).

The following elementary observations turn out to be crucial.

Proposition 58. *Using the notation and assumptions of (57), assume also that W is irreducible, $D := (s = 0)$ for some $s \in H^0(X, L^r)$, $D \cap W$ is irreducible and disjoint from $\Sigma(W)$. Then*

(58.1) *$\mathcal{R}_V^W(D, L) \cong \langle s \rangle_{\mathbb{Q}} \cdot k[W]^\times$, where $\langle s \rangle_{\mathbb{Q}}$ denotes the saturation of the sub-semigroup generated by powers of s .*

(58.2) $\mathcal{R}_V^W(D, L)$ depends only on D, L and V (but not on W) in the following cases

(58.2.a) $k[W] = k$,

(58.2.b) $k[W]/k$ is Galois and V is irreducible, or

(58.2.c) $k[W] = k[\text{red } V]$ and it is separable over k .

Proof. The first assertion follows from (56.1). Indeed, $s|_W$ is the unique section of $L^r|_W$ (up to $k[W]^\times$) that defines $\text{Supp}(D \cap W)$. Therefore

$$\mathcal{R}_V^W(D, L, rm) = s^m|_W \cdot k[W]^\times|_V = s^m|_V \cdot k[W]^\times|_V. \quad (58.3)$$

For other values, $\mathcal{R}_V^W(D, L, m')$ is either empty or consists of a single $k[W]^\times$ -orbit. We can write an element of it as $(s|_W)^{m'/r}$, though it is well defined only up to an r th root of unity.

If (58.2.a) holds then $k[W]^\times = k^\times$ and the k^\times -action on $H^0(V, L^m|_V)$ is independent of W .

In general the image of the restriction map $\sigma : k[W] \rightarrow k[V]$ depends on W , but not much. If $\text{red } V = \{p_i : i \in I\}$ then we have finitely many choices for each embedding $k[W] \hookrightarrow k(p_i)$, giving finitely many possibilities for

$$\text{red}(\sigma) : k[W] \rightarrow \bigoplus_i k(p_i) = k[\text{red } V].$$

(58.2.a–c) are the cases when there is a unique $k[W] \rightarrow \bigoplus_i k(p_i)$. If $k[W]/k$ is a separable extension, then $\text{red}(\sigma)$ has a unique lifting to $\sigma : k[W] \rightarrow k[V]$ by (146.3). \square

Note that $\mathcal{R}_V^W(D, L)$ is nonempty only if D is the support of some section of L^m for some m . Thus (58) is meaningful only if there are sections with irreducible support. For most applications we need many such sections. This leads us to the following definition.

Bertini-Hilbert dimension

59. Let X be a projective variety and L an ample line bundle on X . We are looking for sections $s \in H^0(X, L)$ that satisfy 3 properties.

(59.1) The zero set ($s = 0$) is irreducible.

(59.2) The values of s at some points $x_i \in X$ are specified. More generally, given a 0-dimensional subscheme $Z \subset X$, we would like to specify $s|_Z$.

(59.3) The zero set ($s = 0$) avoids a finite set of points $\Sigma \subset X$.

To formalize these, let X be a scheme over a field k , $Z \subset X$ a subscheme, L a line bundle on X and $s_Z \in H^0(Z, L|_Z)$. Set

$$H^0(X, L, s_Z) := \{s \in H^0(X, L) : s|_Z = cs_Z \text{ for some } c \in H^0(X, \mathcal{O}_X)\}. \quad (59.4)$$

This is a vector subspace of $H^0(X, L)$. If X is integral then for the corresponding linear systems we use the notation $|L, s_Z| \subset |L|$. For a finite subset $\Sigma \subset X$, let

$$|L, s_Z, \Sigma^c| := \{D \in |L, s_Z| : D \cap \Sigma = \emptyset\} \quad (59.5)$$

denote the subset of those divisors that are disjoint from Σ . Finally we use

$$|L, s_Z, \Sigma^c|^{\text{irr}} := \{D \in |L, s_Z, \Sigma^c| : D \text{ is irreducible.}\} \quad (59.6)$$

For our purposes we are free to replace L by L^m . Thus $H^0(X, L^m) \rightarrow H^0(Z, L^m|_Z)$ is surjective and $|L^m, s_Z^m|$ is very ample on $X \setminus Z$ for $m \gg 1$. Hence conditions

(59.2–3) are easy to satisfy and the key question is the irreducibility condition (59.1). Next we discuss 3 cases when we can guarantee irreducibility.

The optimal situation is when $\dim X \geq 2$.

Lemma 60. *Let X be an irreducible, projective variety of dimension ≥ 2 over an infinite field k . Let $\Sigma \subset X$ be a finite subset and $Z \subset X$ a finite subscheme. Let L be an ample line bundle on X and $s_Z \in H^0(Z, L|_Z)^\times$. Then,*

$$(60.1) \quad |L^m, s_Z^m, \Sigma^c|^{\text{irr}} \text{ contains an open and dense subset of } |L^m, s_Z^m| \text{ for } m \gg 1.$$

Proof. $|L^m, s_Z^m|$ is very ample on $X \setminus Z$, hence this follows from the usual Bertini theorems (118). \square

Next we consider Hilbertian fields (153). Here $|L^m, s_Z^m, \Sigma^c|^{\text{irr}}$ need not be open, but it is still quite large. By (60), we need to pay attention only to curves.

Lemma 61. *Let C be an irreducible, projective curve over a Hilbertian field k . Let $\Sigma \subset C$ be a finite subset and $Z \subset C$ a finite subscheme. Let L be an ample line bundle on C and $s_Z \in H^0(Z, L|_Z)^\times$. Then,*

$$(61.1) \quad |L^m, s_Z^m, \Sigma^c|^{\text{irr}} \text{ contains the complement of a thin subset (71) of } |L^m, s_Z^m| \text{ for } m \gg 1.$$

Proof. As before, $|L^m, s_Z^m|$ is very ample on $X \setminus Z$, hence this follows from a basic property of Hilbertian fields (153). \square

Analyzing the proofs in Sections 9–10 shows that a weaker version of (61.1) is sufficient. We only need $|L^m, s_Z^m, \Sigma^c|^{\text{irr}}$ to be nonempty for some $m > 0$. This led to the definition of *weakly Hilbertian* fields (156). The following is essentially their definition; we state it as a lemma to emphasize the similarity to (61).

Lemma 62. *Let C be an irreducible, projective curve over a weakly Hilbertian field k . Let $\Sigma \subset C$ be a finite subset and $Z \subset C$ a finite subscheme. Let L be an ample line bundle on C . Then*

$$(62.1) \quad |L^m, s_Z^m, \Sigma^c|^{\text{irr}} \text{ is nonempty for some } m > 0. \quad \square$$

Note that although we ask for only 1 irreducible divisor, by enlarging Σ we see that we get infinitely many. In fact, the sets

$$|L^m, s_Z^m, \Sigma^c|^{\text{irr}} \subset |L^m, s_Z^m|$$

seem to be quite large, though we do not have a precise way of stating what this means.

For most of the proofs we need to know the smallest dimension where linear systems are guaranteed to have many irreducible members. This leads to the following definition.

Definition 63. Let k be a field that is not locally finite. We define the *Bertini-Hilbert dimension* of k —denoted by $\text{BH}(k)$ —by setting

$$(63.1) \quad \text{BH}(k) = 1 \text{ if } k \text{ is weakly Hilbertian (156), and}$$

$$(63.2) \quad \text{BH}(k) = 2 \text{ otherwise.}$$

In view of (60), the distinction is only about curves. If k is Hilbertian then $\text{BH}(k) = 1$ by (61).

We leave the definition open for locally finite fields. If k is locally finite and L is an ample line bundle on an irreducible curve C , then every smooth point $p \in C$ is

the co-support of some section of some L^m . This would suggest that $\text{BH}(k)$ should be 1, but in some applications setting $\text{BH}(\mathbb{F}_q) = 2$ or even $\text{BH}(\mathbb{F}_q) = \infty$ would seem the right choice.

8. LINEAR SIMILARITY

Definition 64. Let X be a normal variety. Two divisors D_1, D_2 are *linearly similar* if there are nonzero integers m_1, m_2 such that $m_1 D_1 \sim m_2 D_2$. We denote it by $D_1 \sim_s D_2$.

If $\text{rank}_{\mathbb{Q}} \text{Cl}(X) = 1$ then any 2 effective divisors are linearly similar. Thus this notion is nontrivial only if $\text{rank}_{\mathbb{Q}} \text{Cl}(X) > 1$.

The set of all effective divisors linearly similar to a fixed divisor D is naturally an infinite union of linear systems, we denote it by $|\mathbb{Q}D|$.

Let $|\mathbb{Q}D|^{\text{irr}} \subset |\mathbb{Q}D|$ be the subset parametrizing irreducible (but not necessarily reduced) divisors.

Some of the linear systems $|D'| \subset |\mathbb{Q}D|$ may be small and behave exceptionally. So let us call a subset $W \subset |\mathbb{Q}D|$ *stably dense* if $W \cap |mD|$ is dense in $|mD|$ for $m \gg 1$. Note that $|\mathbb{Q}D|^{\text{irr}}$ need not be dense in $|\mathbb{Q}D|$, but if D is ample and $\dim X \geq 2$ (more generally, if D is mobile and has Kodaira dimension ≥ 2) then, by (118.3), $|\mathbb{Q}D|^{\text{irr}}$ is stably dense in $|\mathbb{Q}D|$.

If $\dim X = 1$ then $|\mathbb{Q}D|^{\text{irr}}$ is frequently empty. This presents a serious technical difficulty in our treatment. However, if $\deg D > 0$ then $|\mathbb{Q}D|^{\text{irr}}$ is stably dense in $|\mathbb{Q}D|$ provided $\text{BH}(k) = 1$.

Let $H^0(X, \mathcal{O}_X(\mathbb{Q}D))$ denote the Cox ring of $\mathbb{Q}[D]$, that is, the direct sum of all $H^0(X, \mathcal{O}_X(D'))$ where we pick one D' from each linear system in $|\mathbb{Q}D|$.

If $L = \mathcal{O}_X(D)$ is a line bundle, then a power of $H^0(X, \mathcal{O}_X(\mathbb{Q}D))$ is contained in $H^0(X, \oplus_m L^m)$. It is sometimes convenient to work with $H^0(X, \oplus_m L^m)$.

65 (Restriction and linear similarity). Let X be a variety, $Z \subset X$ a subvariety and D_1, D_2 effective divisors on X . If $D_1 \sim_s D_2$ then (aside from some problems that appear for non-Cartier divisors), $D_1|_Z \sim_s D_2|_Z$.

For us the main interest will be the converse: if $D_1|_Z \sim_s D_2|_Z$, when can we conclude that $D_1 \sim_s D_2$?

Let D be an irreducible divisor. We say that a subvariety $Z \subset X$ *detects* linear similarity to D if the following holds.

(65.1) Let D' be an effective divisor such that $\text{Supp}(D \cap Z) = \text{Supp}(D' \cap Z)$. Then $D' \sim_s D$.

It is not always easy to see when this happens, but the following is quite useful.

Criterion 65.2. Assume that $Z \cap \text{Sing } X$ has codimension ≥ 2 in Z , the kernel of $\text{Cl}(X) \rightarrow \text{Pic}(Z \setminus \text{Sing } X)$ is torsion, D is disjoint from $\Sigma(Z)$ (54.3) and $D \cap Z$ is irreducible. Then Z detects linear similarity to D .

Proof. If $Z \cap \text{Sing } X$ has codimension ≥ 2 in Z then we have a restriction map from rank 1 reflexive sheaves on X (that are locally free along $\Sigma(Z)$) to rank 1 reflexive sheaves on Z (that are locally free along $\Sigma(Z)$) and such a rank 1 reflexive sheaf on Z is determined by the divisors of its sections (54.4). \square

An easy argument (66) shows that this criterion applies if Z is a general, ample, complete intersection subvariety of dimension ≥ 2 .

The kernel of $\mathrm{Cl}(X) \rightarrow \mathrm{Cl}(C)$ is also torsion for many ample, complete intersection curves $C \subset X$, but $D \cap C$ is essentially never irreducible if the base field is algebraically closed. There are 2 ways to go around this problem.

If k is Hilbertian (153) then $D \cap C$ is irreducible for many choices of C .

For arbitrary fields, we need the following.

Criterion 65.3. Let $C \subset X^{\mathrm{ns}}$ be a smooth, projective curve. Assume that the kernel of $\mathrm{Cl}(X) \rightarrow \mathrm{Pic}(C)$ is torsion and the following holds.

(a) Let $D \cap C = \{p_1, \dots, p_r\}$. Then the points p_1, \dots, p_{r-1} are linearly independent over $\mathrm{im}[\mathrm{Cl}(X) \rightarrow \mathrm{Pic}(C)]$. More precisely,

$$\langle p_1, \dots, p_r \rangle \cap \mathrm{im}[\mathrm{Cl}(X) \rightarrow \mathrm{Pic}(C)] = \mathbb{Z}[D|_C].$$

Then C detects linear similarity to D .

Proof. Let D' be another effective divisor such that $\mathrm{Supp}(D' \cap C) = \mathrm{Supp}(D \cap C)$. Note that both D, D' are Cartier along C . Thus $D|_C = \sum d_i [p_i]$ and $D'|_C = \sum d'_i [p_i]$. By (65.3.a) $\sum d'_i [p_i] = m_1 \sum d_i [p_i]$ for some m_1 , hence $D' - m_1 D$ is in the kernel of $\mathrm{Cl}(X) \rightarrow \mathrm{Pic}(C)$. Thus $m_2(D' - m_1 D) \sim 0$ for some $m_2 > 0$. \square

We see in (74) that the condition (65.3.a) holds for many ample-ci curves (117) and use this in (74) to show that if k is not locally finite, then many ample-ci curves detect linear similarity to a divisor.

Lemma 66. *Let X be a normal, projective variety over a field k and D_1, \dots, D_r irreducible divisors on X . Then linear similarity to the D_i is detected by*

(66.1) *general, ample, complete intersections of dimension ≥ 2 , and*

(66.2) *a dense subset of complete intersection curves if k is Hilbertian.*

Proof. Let $Z \subset X$ be a general, ample, complete intersection surface. Then $\mathrm{Cl}(X) \rightarrow \mathrm{Cl}(Z)$ is an injection by (120) and, if $\dim Z \geq 2$, then $Z \cap D_i$ is irreducible and reduced for every i by Bertini's theorem (118.3).

The injectivity of $\mathrm{Cl}(X) \rightarrow \mathrm{Cl}(C)$ also holds for a dense subset of complete intersection curves $C \subset X$ (74), but for irreducibility we need k to be weakly Hilbertian. So (65.3) applies in both cases. \square

We prove a much stronger form of (66.2) in (74), but first we derive some consequences of (66) and (74).

Lemma 67. *Let X be a normal, projective variety over a field k . Assume that*

(67.1) *either k is not locally finite and $\dim X \geq 2$,*

(67.2) *or k is locally finite and $\dim X \geq 3$.*

Then an irreducible divisor H is \mathbb{Q} -Cartier and ample iff the following holds.

(67.3) *For every divisor $D \subset X$ and closed points $p, q \in X \setminus D$, there is a divisor*

$H(p, q) \subset X$ such that

(67.3.a) *$H \cap D = H(p, q) \cap D$,*

(67.3.b) *$p \notin H(p, q)$ and*

(67.3.c) *$q \in H(p, q)$.*

Proof. If H is \mathbb{Q} -Cartier and ample then the restriction map

$$H^0(X, \mathcal{O}_X(mH)) \twoheadrightarrow H^0(D, \mathcal{O}_D(mH|_D)) + \mathcal{O}_X(mH) \otimes k(p) + \mathcal{O}_X(mH) \otimes k(q)$$

is surjective for some $m > 0$. We can thus find a section $s(p, q) \in H^0(X, \mathcal{O}_X(mH))$ as needed.

Conversely, by (66) and (74) we can choose an ample divisor $D \subset X$ that detects linear similarity to H . Then assumption (67.3.a) guarantees that $H(p, q) \sim_s H$. Assumption (67.3.b) implies that H is \mathbb{Q} -Cartier at p . Since p, q are arbitrary points (if we also vary D), H is \mathbb{Q} -Cartier and a multiple of it separates points. \square

Together with (67), the next result proves (9).

Lemma 68. *Let X be a normal, projective variety over a field k and H_1, H_2 irreducible, \mathbb{Q} -Cartier, ample divisors. Assume that*

(68.1) *either k is not locally finite and $\dim X \geq 3$,*

(68.2) *or k is locally finite and $\dim X \geq 5$.*

Then the following are equivalent.

(68.3) $H_1 \sim_s H_2$.

(68.4) $|\mathbb{Q}H_1|^{\text{irr}} = |\mathbb{Q}H_2|^{\text{irr}}$.

(68.5) *Let $Z_1, Z_2 \subset X$ be 2 disjoint, irreducible subvarieties of dimension ≥ 2 if k is locally finite and ≥ 1 otherwise. Then there is a \mathbb{Q} -Cartier, ample divisor H' such that $\text{Supp}(H' \cap Z_i) = \text{Supp}(H_i \cap Z_i)$ for $i = 1, 2$.*

Proof. (68.3) \Leftrightarrow (68.4) is clear. If (68.3) holds then choose $m_1, m_2 \gg 1$ such that $m_1 H_1 \sim m_2 H_2$ and

$$H^0(X, \mathcal{O}_X(m_1 H_1)) \twoheadrightarrow H^0(Z_1, \mathcal{O}_X(m_1 H_1)|_{Z_1}) + H^0(Z_2, \mathcal{O}_X(m_2 H_2)|_{Z_2})$$

is surjective. We can then find $H' \in |m_1 H_1| = |m_2 H_2|$ whose restriction to Z_i is $m_i H_i|_{Z_i}$.

Finally assume (68.5). By (66) and (74) we can choose both Z, W normal, disjoint and such that they detect linear similarity to the H_i . Then we have the chain of linear similarities

$$H_1 \stackrel{\text{(by } Z_1)}{\sim_s} H' \stackrel{\text{(by } Z_2)}{\sim_s} H_2. \quad \square$$

69 (Variants for reducible divisors). With X as in (68), let B_1, B_2 be effective divisors. The above proof shows that (68.5) is equivalent to the following.

(69.1) There are \mathbb{Q} -Cartier, ample, irreducible divisors H_1, H_2 such that $\text{Supp } H_i = \text{Supp } B_i$ and $H_1 \sim_s H_2$.

We can thus recognize irreducible \mathbb{Q} -Cartier divisors using the following criterion.

Claim 69.2. An irreducible divisor $D \subset X$ is \mathbb{Q} -Cartier iff there are irreducible, \mathbb{Q} -Cartier, ample divisors A_1, A_2 such that the above criterion holds for $B_1 := D + A_1$ and $B_2 := A_2$. \square

Remark 70. Using (67) and (68) we get our first topological invariance claims. Namely, let X_K, Y_L be normal, projective varieties such that $|X| \sim |Y|$. Assume that either L is not locally finite and $\dim Y \geq 3$, or $\dim Y \geq 5$. Then

(70.1) If X is \mathbb{Q} -factorial then so is Y .

(70.2) If $\text{rank}_{\mathbb{Q}} \text{Cl}(X) = 1$ then $\text{rank}_{\mathbb{Q}} \text{Cl}(Y) = 1$.

Note that by (27), $\mathbb{P}_{\mathbb{F}_p}^2$ is homeomorphic to smooth surfaces with arbitrary large Picard number, so some restriction on the dimension is necessary in (68).

Néron's theorem and consequences

Definition 71. Let X be an irreducible variety. Following [Ser89], a subset $T \subset X(k)$ is called *thin* if there is a morphism $\pi : Y \rightarrow X$ such that $T \subset \pi(Y(k))$ and there is no rational section $\sigma : X \dashrightarrow Y$. This notion is most interesting for finitely generated, infinite fields. For such fields, $\mathbb{A}^1(k) \subset \mathbb{A}^1(k)$ is not thin; this is essentially due to Hilbert.

A rather typical example to keep in mind is the following. The map $\mathbb{A}^1 \rightarrow \mathbb{A}^1$ given by $x \rightarrow x^2$ shows that the set of all squares is a thin subset of $\mathbb{A}^1(k)$.

We also need a version of this for arbitrary fields K . Let us say that a subset $T \subset X(K)$ is *field-locally thin* if for every finitely generated subfield $k \subset K$, the intersection $T \cap X(k)$ is thin.

Theorem 72. [Nér52b, Thm.6] *Let k be a finitely generated, infinite field. Let $U \subset \mathbb{P}_k^1$ be an open subset and $\pi : T_U \rightarrow U$ a smooth, projective morphism of relative dimension 1. Then there is a dense set $N(T_U) \subset U(k)$, such that the restriction map*

$$\mathrm{Pic}(T_U) \rightarrow \mathrm{Pic}(T_u) \quad \text{is injective for all } u \in N(T_U).$$

Moreover, $N(T_U)$ contains the complement of a thin set. \square

A stronger version is proved in [Sil83, Thm.C], though it applies only to number fields and finite extensions of $\mathbb{F}_p(t)$.

Corollary 73. *Let K be a field that is not locally finite. Let S be a normal, projective surface over K and $|C| = \{C_u : u \in \mathbb{P}^1\}$ a mobile, linear pencil of curves with basepoints $\{p_1, \dots, p_r\}$. Assume that a general C_u is smooth and S is smooth along it. Let $\{B_j : j \in J\}$ be the irreducible components of the reducible members of $|C|$, plus one of the irreducible members. Let m_{ij} be the intersection multiplicity of B_j with a general C_u at p_i ; this is independent of u .*

Then there is a dense set $N(S, |C|) \subset \mathbb{P}^1(K)$ such that, for $u \in N(S, |C|)$, all the linear relations among

$$[p_1(u)], \dots, [p_r(u)] \in \frac{\mathrm{Pic}(C_u)}{\mathrm{im}[\mathrm{Cl}(S) \rightarrow \mathrm{Pic}(C_u)]}$$

are generated by $\sum_i m_{ij} [p_i(u)] = 0$ for all $j \in J$.

Moreover, $N(S, |C|)$ contains the complement of a field-locally thin set.

Proof. Note that the point p_i is contained in every C_u ; the notation $[p_i(u)]$ indicates that we take its class in $\mathrm{Pic}(C_u)$, which depends on u .

The restriction of B_j to C_u is $\sum_i m_{ij} [p_i(u)]$, so we do need to have the equations $\sum_i m_{ij} [p_i(u)] = 0$. The interesting part is to show that there are no other relations.

Let T be the normalization of the closure of the graph of $|C| : S \dashrightarrow \mathbb{P}^1$. The projection $\pi_1 : T \rightarrow S$ is birational, with exceptional curves $E_i \subset T$ sitting over p_i . Let $B_j^T \subset T$ denote the birational transform of B_j . Note that

$$B_j^T \sim \pi_1^* B_j - \sum_i m_{ij} E_i.$$

The second projection $\pi_2 : T \rightarrow \mathbb{P}^1$ is generically smooth and the irreducible components of its fibers are exactly the B_j^T .

Let $U \subset \mathbb{P}^1$ be the largest open set over which π_2 is smooth. By restriction we get $T_U \rightarrow U$. The Picard group of T_U is then

$$\mathrm{Pic}(T_U) = \mathrm{Cl}(T) / \langle B_j^T : j \in J \rangle.$$

Choose now a finitely generated subfield $k \subset K$ such that $S, |C|$, the p_i and the B_j are defined over k .

Note that $\text{Cl}(T_k) = \pi_1^* \text{Cl}(S_k) + \sum_i [E_i]$, and killing $\pi_1^* \text{Cl}(S_k)$ gives a natural surjection

$$\text{Cl}(T_k)/\pi_1^* \text{Cl}(S_k) \longrightarrow \langle E_i : i \in I \rangle / \langle \sum_i m_{ij} [p_i] : j \in J \rangle.$$

Thus all the linear relations among $[E_1], \dots, [E_r] \in \text{Cl}(T_k)/\pi_1^* \text{Cl}(S_k)$ are generated by $\sum_i m_{ij} [E_i] = 0$ for all $j \in J$.

We now apply (72) to get $N(T_U) \subset \mathbb{P}^1(k)$ such that, for $u \in N(T_U)$, all the linear relations among

$$[p_1(u)], \dots, [p_r(u)] \in \text{Pic}(C_u)/\text{im}[\text{Cl}(S_k) \rightarrow \text{Pic}(C_u)]$$

are generated by $\sum_i m_{ij} [p_i(u)] = 0$ for all $j \in J$.

This is not exactly what we want since $\text{Cl}(S_k)$ may be much bigger than $\text{Cl}(S_k)$. However, if we have a linear relation

$$\sum_i \lambda_i [p_i(u)] \sim [L_K] \quad \text{where} \quad L_K \in \text{im}[\text{Cl}(S_k) \rightarrow \mathbf{Pic}(C_u)](K),$$

then in fact $L_K \in \text{im}[\text{Cl}(S_k) \rightarrow \mathbf{Pic}(C_u)](k)$, hence a power of L_K is in $\text{im}[\text{Cl}(S_k) \rightarrow \text{Pic}(C_u)]$ (144). Letting k vary now proves our claim. \square

Theorem 74. *Let k be a field that is not locally finite. Let X be a normal, projective variety of dimension $n \geq 2$ over k , $\{D_i : i = 1, \dots, r\}$ irreducible Weil divisors and H_1, \dots, H_{n-1} ample divisors. Then, for $m_1 \gg, m_2, \dots, m_r \gg 1$, there is a dense set*

$$U \subset |m_1 H_1|(k) \times \dots \times |m_{n-1} H_{n-1}|(k)$$

such that, for $u \in U$, the corresponding complete intersection curve C_u detects linear similarity to each D_i .

Proof. By (66), there is a Zariski open

$$U_2 \subset |m_2 H_2| \times \dots \times |m_{n-1} H_{n-1}|$$

such that $\text{Cl}(X) \rightarrow \text{Cl}(H_2 \cap \dots \cap H_{n-1})$ is an injection for $(H_2, \dots, H_{n-1}) \in U_2$ and the $D_i \cap H_2 \cap \dots \cap H_{n-1}$ are irreducible. This reduces us to the case $n = 2$.

Thus from now on we have a normal, projective surface X over k , $\{D_i : i = 1, \dots, r\}$ irreducible Weil divisors on X and an ample divisor H on X .

Now choose a pencil $|C| \subset |mH|$ such that

$$(74.1) \quad D_0 + \dots + D_r \in |C| \text{ for some irreducible curve } D_0,$$

$$(74.2) \quad \text{all other members of } |C| \text{ are irreducible,}$$

$$(74.3) \quad \text{the general member of } |C| \text{ is smooth and } X \text{ is smooth along it.}$$

Applying (73) to it we get a dense set of $|C|(k)$ where the requirements hold. \square

Remark. Most likely one can choose U such that it contains the complement of a field-locally thin set.

9. LINKAGE OF DIVISORS AND RESIDUE FIELDS

Definition 75. Let Y be a reduced, projective scheme (not necessarily pure dimensional) and L an ample line bundle on Y . Two effective divisors $D_1, D_2 \subset Y$ are called (*topologically, directly*) L -linked if there is a section $s \in H^0(Y, L^m)$ such that $\text{Supp}(s = 0) = \text{Supp } D_1 \cup \text{Supp } D_2$.

Next we look at an unusual special case of linkage.

Definition 76. Let X be a normal, projective, irreducible k -variety and $Z, W \subset X$ closed subsets. Let L be an ample line bundle on X . Two effective divisors $H_Z, H_W \in |\mathbb{Q}L|$ are (topologically, directly) L -linked on $Z \cup W$ if $H_Z \cap Z$ and $H_W \cap W$ are L -linked on $Z \cup W$. That is,

$$(76.1) \text{ there is a section } s \in H^0(X, L^m) \text{ such that } (s = 0) \cap Z = H_Z \cap Z \text{ and } (s = 0) \cap W = H_W \cap W \text{ (as sets).}$$

Note that (75) would suggest working with a section $s' \in H^0(Z \cup W, L^m|_{Z \cup W})$, but a power of s' lifts to a section on X , so the 2 versions are equivalent.

As we see below, this notion is not interesting if $Z \cap W = \emptyset$ and it has various problems if $\dim(Z \cap W) \geq 1$. Thus we focus on the case when $\dim(Z \cap W) = 0$.

A key observation is that linkage carries very significant information about

- residue fields of $Z \cap W$ in every characteristic, and
- the structure of $Z \cap W$ in characteristic 0.

Proposition 77. *Let X be a normal, projective, irreducible k -variety and $Z, W \subset X$ closed subsets such that $\dim(Z \cap W) = 0$. Let L be an ample line bundle on X and $H_Z, H_W \in |\mathbb{Q}L|$. Then H_Z, H_W are L -linked on $Z \cup W$ iff (using (57))*

$$\mathcal{R}_{Z \cap W}^Z(H_Z, L) \cap \mathcal{R}_{Z \cap W}^W(H_W, L) \neq \emptyset.$$

Proof. Assume that $H = (s = 0)$ gives the L -linkage for some $s \in H^0(X, L^m)$. Then $s|_Z \in \Gamma^{Z \cap H_Z}(Z, L|_Z, m)$ and $s|_W \in \Gamma^{W \cap H_W}(W, L|_W, m)$ have the same restriction to $Z \cap W$.

Conversely, if $s_Z \in H^0(Z, L^m|_Z)$ and $s_W \in H^0(W, L^m|_W)$ have the same image in $H^0(Z \cap W, L^m|_{Z \cap W})$, then they glue to a section $s_{Z \cup W} \in H^0(Z \cup W, L^m|_{Z \cup W})$, and then $s_{Z \cup W}^{m'}$ lifts to a section of $H^0(X, L^{m'm})$ for some $m' > 0$. \square

78. The conditions in (77) give the strongest restriction if

$$\mathcal{R}_{Z \cap W}^Z(H_Z, L)/k^\times \quad \text{and} \quad \mathcal{R}_{Z \cap W}^W(H_W, L)/k^\times \quad (78.1)$$

both have \mathbb{Q} -rank 1. However, in general these objects are essentially extensions of a finite rank semigroup by $k[Z]^\times$ (resp. $k[W]^\times$). Criteria for (78.1) are given in (55.2). This case can be used to identify the k -points of X (82).

We get further interesting consequences if we relax these restrictions. The general situation seems rather complicated. In our applications it is advantageous to work with a non-symmetric situation:

$$(78.2) \quad H^0(Z, \mathcal{O}_Z) = k, \text{ and}$$

$$(78.3) \quad \mathcal{R}_{Z \cap W}^W(H_W, L)/k[W]^\times \text{ has } \mathbb{Q}\text{-rank } 1.$$

Note that (78.2) holds if Z is geometrically connected and reduced. In applications we achieve this by choosing Z to be ample-ci (117).

We saw in (58) that (78.3) holds if $\dim W \geq \text{BH}(k)$ (with some additional mild genericity conditions).

Next we study the case when linking is always possible.

Definition 79 (Free linking). Let X be a normal, projective, irreducible k -variety and L an ample line bundle on X . Let $Z, W \subset X$ be closed, integral subvarieties such that $\dim(Z \cap W) = 0$.

We say that L -linking is free on $Z \cup W$ iff two divisors $H_Z, H_W \in |\mathcal{O}_X(L)|$ are L -linked on $Z \cup W$ whenever they are disjoint from $\Sigma(Z \cup W)$ (54.3).

In the rest of the section we discuss various cases when the topological notion of free linking makes it possible to obtain information about the residue fields of closed points. The key is the following.

Conjecture 80. *Let X be a normal, projective, irreducible k -variety and L an ample line bundle on X . Let $Z, W \subset X$ be closed, integral, positive dimensional subvarieties such that $\dim(Z \cap W) = 0$. The following are equivalent.*

(80.1) L -linking is free on $Z \cup W$.

(80.2) $k[Z \cap W]^\times / k[Z]^\times \cdot k[W]^\times$ is a torsion group.

(80.3) One of the following holds.

(80.3.a) $\text{char } k = 0$, $Z \cap W$ is reduced, and
either $k[Z \cap W] = k[Z]$ or $k[Z \cap W] = k[W]$.

(80.3.b) $\text{char } k > 0$, and either $k[\text{red}(Z \cap W)]/k[Z]$ or $k[\text{red}(Z \cap W)]/k[W]$ is purely inseparable.

(80.3.c) k is locally finite.

Comments. Note that (80.2) \Rightarrow (80.1) holds by (77) and the equivalence of (80.2) and (80.3) is proved in (149).

Next we show that (80.1) \Rightarrow (80.3) holds if W is geometrically connected and $\dim W \geq \text{BH}(k)$. A careful study of the proof shows that the first assumption is not necessary, and the validity of (15) would imply that (80.1) \Rightarrow (80.2) holds unconditionally.

Proposition 81. *Let X be a normal, projective, geometrically irreducible k -variety and L an ample line bundle on X . Let $Z, W \subset X$ be closed, integral, positive dimensional subvarieties such that $\dim(Z \cap W) = 0$. Assume that W is geometrically connected and $\dim W \geq \text{BH}(k)$.*

Then (80.1–4) are equivalent.

Proof. As we noted, we need to show that (80.1) \Rightarrow (80.3). Thus assume (80.1). First choose $s_Z \in H^0(X, L^m)$ such that $\text{Supp}(s_Z = 0) = \text{Supp } H_Z$ is disjoint from $\Sigma(Z \cup W)$.

By (60) and (62), for every $s_{Z \cap W} \in k[Z \cap W]^\times$, there is an $n > 0$ such that $s_{Z \cap W}^n$ extends to $s_W \in H^0(X, L^n)$ and $W \cap H_W$ is irreducible, where $H_W := \text{Supp}(s_W = 0)$.

If H_Z, H_W are L -linked, then there is an $s \in H^0(X, L^m)$ as in (76). (We can use the same m , if we pass to suitable powers of s, s_Z, s_W .)

By (56), there are $u_Z \in k[Z]^\times, u_W \in k[W]^\times = k^\times$, a finitely generated subgroup $\Gamma_Z \subset k(Z)^\times, \gamma_Z \in \Gamma_Z$ and a natural number r such that

$$s_W^r = s^r|_W \cdot u_W \quad \text{and} \quad s_Z^r = s^r|_Z \cdot u_Z \cdot \gamma_Z. \quad (81.1)$$

Therefore

$$\begin{aligned} s_W^r|_{Z \cap W} \cdot s_Z^{-r}|_{Z \cap W} &= u_W|_{Z \cap W} \cdot u_Z^{-1}|_{Z \cap W} \cdot \gamma_Z^{-1}|_{Z \cap W} \\ &\in k^\times \cdot k[Z]^\times \cdot \Gamma_Z|_{Z \cap W} = k[Z]^\times \cdot \Gamma_Z|_{Z \cap W}. \end{aligned} \quad (81.2)$$

Next note that $s_W|_{Z \cap W} = s_{Z \cap W}^n$ where $s_{Z \cap W}$ is arbitrary. Thus $s_{Z \cap W}^n s_Z^{-1}|_{Z \cap W}$ is an arbitrary element of $k[Z \cap W]^\times$ (up to n -torsion). Therefore we get that

$$k[Z \cap W]^\times / (k[Z]^\times \cdot \Gamma_Z|_{Z \cap W}) \quad \text{is torsion.} \quad (81.3)$$

Thus we obtain that

$$k[Z \cap W]^\times / k[Z]^\times \quad \text{has finite } \mathbb{Q}\text{-rank.} \quad (81.4)$$

By (147) we are in one of 4 cases.

(81.5) k is locally finite; giving (80.4.c).

(81.6) $\text{char } k > 0$ and $k[Z] \hookrightarrow k[\text{red}(Z \cap W)]$ is a purely inseparable extension; giving (80.4.b).

(81.7) $\text{char } k = 0$ and $k[Z] \cong k[Z \cap W]$; giving (80.4.a).

(81.8) $\deg(k/\mathbb{Q}) < \infty$.

In the latter case k is Hilbertian. Once k is Hilbertian, at the beginning of the proof we can choose $Z \cap H_Z$ to be irreducible; in which case $\Gamma_Z = \{1\}$ by (56.1). Thus (81.3) becomes

$$k[Z \cap W]^\times / k[Z]^\times \quad \text{is torsion,} \quad (81.9)$$

and (147) implies that $Z \cap W$ is reduced. \square

Using (81) we get a topological way of recognizing k -points.

Corollary 82. *Let k be a perfect field that is not locally finite, and X a normal, projective, geometrically irreducible k -variety of dimension $> 1 + \text{BH}(k)$. Let L be an ample line bundle and $p \in X$ a closed point. Assume that either $\text{char } k > 0$ or p is a smooth point of X . The following are equivalent.*

(82.1) p is a k -point.

(82.2) There are integral, ample-sci (117) subvarieties Z, W such that

(82.2.a) $\dim Z = 1, \dim W = \text{BH}(k)$,

(82.2.b) $\text{Supp}(Z \cap W) = \{p\}$ and

(82.2.c) L -linking is free on $Z \cup W$.

Proof. (82.2) \Rightarrow (82.1) follows from (81). Conversely, we can take Z, W to be general complete intersections of ample divisors containing p . \square

Remark 82.3. If $\text{char } k = 0$ and (82.2) holds then $Z \cap W$ is a k -point, even if X is singular there. However, for a singular k -point it may not be possible to find Z, W such that $Z \cap W = \{p\}$ (as schemes). Thus the method does not yet provide a topological way of identifying singular k -points if $\text{char } k = 0$.

Corollary 83. *Let k be a perfect field that is not locally finite, and X a normal, projective, geometrically irreducible k -variety of dimension $> 1 + \text{BH}(k)$. Let L be an ample line bundle and $p, q \in X$ closed points. Assume that either $\text{char } k > 0$ or p is a smooth point. The following are equivalent.*

(83.1) There is a k -embedding $k(p) \hookrightarrow k(q)$.

(83.2) There are irreducible subvarieties Z, W such that

(83.2.a) $\dim Z = 1, \dim W = \text{BH}(X)$,

(83.2.b) $\text{Supp}(Z \cap W) = \{p\}$,

(83.2.c) $q \in Z$,

(83.2.d) W is geometrically connected, and

(83.2.e) L -linking is free on $Z \cup W$.

Proof. If (83.2) holds then $k(p) \cong k[Z]$ by (81) and (83.2.c) gives an embedding $k[Z] \hookrightarrow k(q)$.

Conversely, given $k(p) \hookrightarrow k(q)$, the required Z is constructed in (122) and then choose W to be a general complete intersection containing p . \square

Reversing the role of p, q we obtain a criterion to decide whether $k(p) \cong k(q)$. Note, however, that we get no information about $\deg(k(p)/k)$. Nonetheless, we have enough information to obtain the following.

84 (Isomorphism of 0-cycles from $|X|$ and \sim_{sa}). Let k be a perfect field that is not locally finite, and X a normal, projective, irreducible k -variety of dimension $> 1 + \text{BH}(k)$. Let $Z_1, Z_2 \subset X$ be reduced 0-dimensional subschemes. Assume that either $\text{char } k > 0$ or $Z_1, Z_2 \subset X^{\text{ns}}$.

We can then decide, using only $(|X|, \sim_{\text{sa}})$, whether Z_1, Z_2 are isomorphic as k -schemes. \square

85 (Imperfect fields). If k is an imperfect field, we can apply the above results to k^{ins} . This results in the following obvious changes in the statements.

In (82.1) we characterize k^{ins} -points.

In (83.1) we characterize k -embeddings $k^{\text{ins}}(p) \hookrightarrow k^{\text{ins}}(q)$.

In (84) we characterize isomorphisms $Z_1 \times_k k^{\text{ins}} \cong Z_2 \times_k k^{\text{ins}}$.

Remark 86. Let X a normal, projective, geometrically irreducible k -variety of dimension $> 1 + \text{BH}(k)$. Then $\text{char } k > 0$ iff the following holds.

(86.1) There is an integral curve $C \subset X$ and a point $p \in C$, such that L -linking is free on $C \cup W$ for every ample-sci subvariety W of dimension $\text{BH}(k)$, for which $\text{Supp}(C \cap W) = \{p\}$.

Indeed, if $\text{char } k = 0$ then for any $p \in C$ we can choose W such that $C \cap W$ is non-reduced, and then L -linking is not free on $C \cup W$ by (81).

Conversely, we use (122) to get $p \in C$ such that $k(p)^{\text{ins}} = k[C]^{\text{ins}}$, and then (81) and (85) show that L -linking is free on $C \cup W$ if $\text{char } k > 0$.

10. MINIMALLY RESTRICTIVE LINKING AND TRANSVERSALITY

Definition 87. Let X be a normal, projective, irreducible k -variety and $Z, W_1, W_2 \subset X$ closed, irreducible, geometrically connected subvarieties such that $\dim(Z \cap W_i) = 0$. We say that L -linking on W_2 determines L -linking on W_1 if the following holds.

(87.1) Let $H_Z, H_W \in |\mathbb{Q}L|^{\text{irr}}$ be effective divisors disjoint from the $\Sigma(Z \cup W_i)$ such that $W_2 \cap H_W$ is irreducible. Then

$$\left(\begin{array}{c} H_Z, H_W \text{ are} \\ \text{linked on } Z \cup W_2 \end{array} \right) \Rightarrow \left(\begin{array}{c} H_Z, H_W \text{ are} \\ \text{linked on } Z \cup W_1 \end{array} \right).$$

In applying this notion we always assume that $\dim W_i \geq \text{BH}(k)$, hence the above conditions are not empty.

We say that L -linking is *minimally restrictive* on W_1 , if L -linking on W_2 determines L -linking on W_1 , whenever $\text{Supp}(Z \cap W_1) = \text{Supp}(Z \cap W_2)$.

The key result—and rationale for the definition—is the following.

Proposition 88. *Let k be a field of characteristic 0, X a normal, projective, irreducible k -variety and L an ample line bundle on X . Let $Z, W_1, W_2 \subset X$ be closed, integral, geometrically connected subvarieties such that $\dim Z \geq 1$, $\dim W_i \geq \text{BH}(k)$ and $\dim(Z \cap W_i) = 0$. The following are equivalent.*

(88.1) $Z \cap W_1 \subset Z \cap W_2$ as schemes.

(88.2) L -linking on W_2 determines L -linking on W_1 .

Proof. Pick $H_Z = (s_Z = 0)$ and $H_W = (s_W = 0)$. By assumption $W_2 \cap H_W$ is irreducible and disjoint from $\Sigma(Z \cup W_2)$. Thus, by (58.1),

$$\mathcal{R}_{Z \cap W_2}^{W_2}(H_Z, L) = \langle s_W|_{Z \cap W_2} \rangle_{\mathbb{Q}} \cdot k[W_2]^{\times} = \langle s_W|_{Z \cap W_2} \rangle \cdot k^{\times}.$$

So, by (77), H_Z, H_W are linked on $Z \cup W_2$ iff, for some $r > 0$,

$$s_W^r|_{Z \cap W_2} \in \mathcal{R}_{Z \cap W_2}^Z(H_Z, L) \subset H^0(Z \cap W_2, \oplus_m L^m|_{Z \cap W_2}). \quad (88.3)$$

Note that (88.1) holds iff there is a natural surjection

$$H^0(Z \cap W_2, \oplus_m L^m|_{Z \cap W_2}) \rightarrow H^0(Z \cap W_1, \oplus_m L^m|_{Z \cap W_1}).$$

Thus (88.3) implies that

$$s_W^r|_{Z \cap W_1} \in \mathcal{R}_{Z \cap W_1}^Z(H_Z, L) \subset H^0(Z \cap W_1, \oplus_m L^m|_{Z \cap W_1}), \quad (88.4)$$

proving (88.2).

To see the converse, let N be the kernel of

$$\rho : H^0(Z \cap W_1, \mathcal{O}_{Z \cap W_1})^{\times} \rightarrow H^0(Z \cap W_1 \cap W_2, \mathcal{O}_{Z \cap W_1 \cap W_2})^{\times}.$$

It is a direct sum of a commutative, unipotent group over k and of the $k(p_i)^{\times}$ for every $p_i \in W_1 \setminus W_2$. N is positive dimensional iff $Z \cap W_1 \not\subset Z \cap W_2$. We distinguish 2 cases, depending on whether $\text{rank}_{\mathbb{Q}} N(k) = \infty$ or not.

Next choose any $\sigma_2 \in H^0(Z \cap W_2, L|_{Z \cap W_2})^{\times}$. If $\text{rank}_{\mathbb{Q}} N(k) = \infty$ then the restriction of σ_2 to $Z \cap W_1 \cap W_2$ can be lifted in 2 different ways to

$$\sigma_Z, \sigma_W \in H^0(Z \cap W_1, L|_{Z \cap W_1})^{\times}$$

such that $\sigma_Z \sigma_W^{-1}$ is non-torsion in

$$H^0(Z \cap W_1, \oplus_m L^m|_{Z \cap W_1})^{\times} / \mathcal{R}_{Z \cap W_1}^Z(H_Z, L).$$

We can now glue σ_2, σ_W to a section of $H^0(Z \cap (W_1 \cup W_2), L|_{Z \cap (W_1 \cup W_2)})^{\times}$ and then lift (some power of) it to $s_W \in H^0(X, L)$ such that both $W_i \cap (s_W = 0)$ are irreducible and disjoint from $\Sigma(Z \cup W_1 \cup W_2)$. Similarly, we can glue σ_2, σ_Z to a section of $H^0(Z \cap (W_1 \cup W_2), L|_{Z \cap (W_1 \cup W_2)})^{\times}$ and then lift (some power of) it to $s_Z \in H^0(X, L)$ such that $Z \cap (s_Z = 0)$ is irreducible and disjoint from $\Sigma(Z \cup W_1 \cup W_2)$.

By construction, $s_Z|_{Z \cap W_2} = s_W|_{Z \cap W_2}$, hence $(s_Z = 0)$ and $(s_W = 0)$ are L -linked on $Z \cup W_2$, but $s_Z|_{Z \cap W_1}$ and $s_W|_{Z \cap W_1}$ are multiplicatively independent, hence $(s_Z = 0)$ and $(s_W = 0)$ are not L -linked on $Z \cup W_2$.

We are left with the case when $\text{rank}_{\mathbb{Q}} N(k) < \infty$. In this case $\deg(k/\mathbb{Q}) < \infty$ by (141), hence k is Hilbertian (153). We can thus choose s_Z such that $Z \cap H_Z$ is irreducible and disjoint from $\Sigma(Z \cup W_2)$. So, by (58.1),

$$\mathcal{R}_{Z \cap W_2}^Z(H_Z, L) = \langle s_Z|_{Z \cap W_2} \rangle_{\mathbb{Q}} \cdot k[Z]^{\times} = \langle s_Z|_{Z \cap W_2} \rangle_{\mathbb{Q}} \cdot k^{\times}.$$

This implies that $\mathcal{R}_{Z \cap W_2}^Z(H_Z, L)$ has trivial intersection with N . We can thus again choose $\sigma_Z, \sigma_W \in H^0(Z \cap W_1, L|_{Z \cap W_1})^{\times}$ such that $\sigma_Z \sigma_W^{-1}$ is non-torsion, and complete the proof as before. \square

Corollary 89. *Let k be a field of characteristic 0, X a normal, projective, irreducible k -variety and L an ample line bundle. Let $Z, W \subset X$ be closed, integral, geometrically connected subvarieties such that $\dim(Z \cap W) = 0$. Assume that $\dim X > \dim Z + \text{BH}(k)$*

Then L -linking is minimally restrictive on $Z \cup W$ iff $Z \cap W$ is reduced.

Proof. Set $W_1 := W$ in (88) and let W_2 run through all ample-sci subvarieties of dimension $\text{BH}(k)$ (117) that intersect Z exactly along $Z \cap W_1$. Then apply (90). \square

Lemma 90. *Let X be a projective k -variety, $Z \subset X$ a subscheme of codimension $> r$ and $P \subset Z$ a reduced, finite subscheme. Let $\mathcal{W}(Z, P)$ be the set of all irreducible, r -dimensional, ample-sci (117) subvarieties $W \subset X$ for which $\text{Supp}(Z \cap W) = P$. Then*

$$\bigcap_{W \in \mathcal{W}(Z, P)} W = P \quad (\text{scheme theoretically}). \quad \square$$

The following consequence of (89) allows us to understand intersection multiplicities topologically.

91 (Determining transversality from $|X|$ and \sim_{sa}). Let k be a field of characteristic 0, X a normal, projective, geometrically irreducible k -variety of dimension $> 1 + \text{BH}(k)$. Let $H \subset X$ be an irreducible, ample divisor and $C \subset X$ an irreducible, geometrically connected curve. Assume that $C \cap H \subset X^{\text{ns}}$. The following are equivalent.

(91.1) All intersections of $C \cap H$ are transversal.

(91.2) $C \cap H$ is reduced.

(91.3) L -linking is minimally restrictive on $C \cup H$ for some ample line bundle L .

Indeed, (91.2) \Leftrightarrow (91.3) follows from (89) and (91.1) \Leftrightarrow (91.2) is a basic property of intersection multiplicities; see for example [Ful98, 8.2]. \square

Note that there are several weaknesses of the current form of the above equivalences.

First, we do not yet know how to decide which are the smooth points of X . We usually go around this by saying that some assertion holds outside some codimension ≥ 2 subset.

Second, we also do not yet know how to decide whether a curve C is geometrically connected or not. However, if C is ample-sci (117), then C is geometrically connected (117.1).

The above arguments also show the following.

Corollary 92. *Let k be a field of characteristic 0, X a k -variety, $Z \subset X$ an irreducible, geometrically connected subvariety of codimension $r > \text{BH}(k)$ and $p \in Z$ a closed point such that X is smooth at p . Then Z is smooth at p iff there is an irreducible, ample-sci subvariety $W \subset X$ of dimension r such that $p \in \text{Supp}(Z \cap W)$ and L -linking is minimally restrictive on W . \square*

Interchanging the roles of Z, W gives the following dual version.

Corollary 93. *Let k be a field of characteristic 0, X a k -variety, $W \subset X$ an irreducible, geometrically connected subvariety of dimension $r > \text{BH}(k)$ and $p \in W$ a closed point such that X is smooth at p . Then W is smooth at p iff there is an irreducible, ample, complete intersection subvariety $Z \subset X$ of codimension r such that $p \in \text{Supp}(Z \cap W)$ and H -linking is minimally restrictive on W . \square*

94. The argument in (88) also applies in positive characteristic, except that then the kernel of

$$H^0(Z \cap W, \mathcal{O}_{Z \cap W})^\times \rightarrow H^0(\text{red}(Z \cap W), \mathcal{O}_{\text{red}(Z \cap W)})^\times$$

is p -power torsion. Thus multiplicative independence is not changed as we pass from $Z \cap W$ to $\text{red}(Z \cap W)$. We get that, if k is not locally finite, then the following are equivalent.

$$(94.1) \text{ Supp}(Z \cap W_1) \subset \text{Supp}(Z \cap W_2).$$

$$(94.2) \text{ } L\text{-linking on } W_1 \text{ determines } L\text{-linking on } W_2.$$

Thus, while L -linking carries scheme-theoretic information in characteristic 0, it does not in positive characteristic.

11. DEGREES OF CURVES AND DIVISORS

Next we try to determine intersection numbers $(C \cdot H)$ of curves and divisors. First we use (91) to check that all intersections are transversal. If the field is algebraically closed, then $(C \cdot H) = \#(C \cap H)$, and we are done.

If the field is not algebraically closed, we would need to compute $\deg k(p)/k$ for the intersection points. This we can not do, but, by (84), we can determine when $\deg k(p_1)/k = \deg k(p_2)/k$ for 2 points. The following result says that, although we are not able to compute $(C \cdot H)$, we can decide whether $(C_1 \cdot H) = (C_2 \cdot H)$ for 2 curves. This is enough for many applications.

Theorem 95. *Let X be a projective variety over an infinite field, H an irreducible, ample, Cartier divisor, $\{C_i : i \in I\}$ finitely many irreducible, geometrically reduced curves. The following are equivalent.*

$$(95.1) (C_i \cdot H) \text{ is independent of } i.$$

$$(95.2) \text{ There is an irreducible divisor } G \sim_{\text{sa}} H \text{ such that the (scheme theoretic) intersections } \{C_i \cap G : i \in I\} \text{ are reduced and isomorphic to each other.}$$

Moreover, we can choose G to be disjoint from any finite subset $\Sigma \subset X$.

Proof. Assume that (95.2) holds and $G \sim mH$. Then $m(C_i \cdot H) = \deg_k(C_i \cap G)$, proving (95.1). To see the converse, let C denote the union of the C_i . Set $L := \mathcal{O}_X(H)|_C$. Choose $m \gg 1$ such that $H^0(X, \mathcal{O}_X(mH)) \rightarrow H^0(C, \mathcal{O}_C(mH|_C))$ is surjective and there is a section $s \in H^0(C, \mathcal{O}_C(mH|_C))$ as in (99). Then $G := (s = 0)$ works. \square

Corollary 96. *Let X be a normal, projective variety over an infinite field, H an irreducible, ample, Cartier divisor, $\{D_i : i \in I\}$ irreducible, geometrically reduced divisors and $B \subset X$ a closed subset of dimension $\leq n - 2$ containing $\text{Sing } X$. The following are equivalent.*

$$(96.1) (D_i \cdot H^{n-1}) \text{ is independent of } i.$$

$$(96.2) \text{ There are irreducible, } H\text{-sci curves } A \text{ (117) that are disjoint from } B \text{ and such that the (scheme theoretic) intersections } A \cap D_i \text{ are reduced and isomorphic to each other.}$$

Proof. As before, (96.2) \Rightarrow (96.1) is clear. For the converse, we look for A contained in a general complete intersection surface $S \subset X$. This reduces us to the special case when $\dim X = 2$. Then the D_i are curves, so (96) follows from (95). \square

The following observation allows us to describe \mathbb{Q} -linear equivalence topologically.

97 (Determining numerical equivalence from $|X|$ and \sim_{sa}). We state 4 pairs of results. In each of them the first is the information we seek, the second shows how it can be obtained using (95–96) and the previous ones. We assume that X is a projective variety over a field of characteristic 0. The latter is necessary in order to use (89). We use several claims about intersections of curves and divisors; these are recalled in (98).

Claim 97.1. Let C_1, C_2 be irreducible curves, not contained in $\text{Sing } X$. Then

- $(C_1 \cdot H) = (C_2 \cdot H)$ iff
- for every finite subset $\Sigma \subset C_1 \cup C_2$ there is an irreducible divisor $G \sim_s H$ disjoint from Σ , such that
 - (a) H -linking is minimally restrictive on $C_i \cup G$ for $i = 1, 2$, and
 - (b) $C_1 \cap G$ and $C_2 \cap G$ are isomorphic as k -schemes. □

Claim 97.2. Let C_1, C_2 be irreducible curves, not contained in $\text{Sing } X$. Then

- $C_1 \equiv C_2$ iff
- $(C_1 \cdot H) = (C_2 \cdot H)$ for every ample divisor H . □

Claim 97.3. Let D_1, D_2 be irreducible, geometrically connected divisors on X and H an ample divisor. Then

- $(D_1 \cdot H^{n-1}) = (D_2 \cdot H^{n-1})$ iff
- For every codimension ≥ 2 subset $B \subset X$ there are irreducible, H -sci curves A that are disjoint from B and such that
 - (a) H -linking is minimally restrictive on $A \cup D_i$ for $i = 1, 2$, and
 - (b) $A \cap D_1$ and $A \cap D_2$ are isomorphic as k -schemes. □

Claim 97.4. Let D_1, D_2 be irreducible, geometrically connected divisors on X . Then

- $D_1 \equiv D_2$ iff
- $(D_1 \cdot H^{n-1}) = (D_2 \cdot H^{n-1})$ for every ample divisor H . □

Claim 97.5. Let H_1, H_2 be irreducible, ample divisors. Then

- $H_1 \sim_{\mathbb{Q}} H_2$ iff
- $H_1 \sim_{\text{sa}} H_2$ and $(H_1 \cdot H_2^{n-1}) = (H_2 \cdot H_1^{n-1})$. □

Lemma 98. *Let X be a normal, projective variety.*

(98.1) *Let C_1, C_2 be 1-cycles. Then $C_1 \equiv C_2$ iff $(C_1 \cdot H) = (C_2 \cdot H)$ for every ample Cartier divisor H .*

(98.2) *Let D_1, D_2 be \mathbb{Q} -Cartier divisors. Then $D_1 \equiv D_2$ iff $(D_1 \cdot A) = (D_2 \cdot A)$ for every irreducible, ample-sci curve (117).*

Proof. We use the theory of cones and divisors; see for example [Laz04, 1.4.C]. Let $N^1(X) = \text{NS}(X) \otimes \mathbb{Q}$ be the Néron-Severi space (the space of \mathbb{Q} -Cartier \mathbb{Q} -divisors modulo algebraic equivalence) and $N_1(X)$ the space of curves with \mathbb{Q} -coefficients, modulo algebraic equivalence. By Kleiman's theorem [Laz04, 1.4.23] they are dual to each other, hence they have the same dimension. Ample divisors span $N^1(X)$, thus (98.1) holds. For (98.2) it remains to show that irreducible, ample-sci curves span $N_1(X)$.

First we claim that if H is ample then $H^{n-2} : N^1(X) \rightarrow N_1(X)$ is an isomorphism. This is a special case of the Hard Lefschetz theorem [Laz04, 3.1.39], which is usually stated for smooth varieties. Since $N^1(X)$ and $N_1(X)$ have the same dimension, it is enough to show that H^{n-2} gives an injection. This follows from the Grothendieck-Lefschetz hyperplane theorem: if $S \subset X$ is a general H -complete intersection surface then $\text{Pic}(X) \rightarrow \text{Pic}(S)$ is an injection.

Finally, let D be any Cartier divisor. Then $H + \epsilon D$ is ample for $0 < \epsilon \ll 1$ and

$$H^{n-2}D \in \langle (H^i D^{n-1-i}) : i = 0, \dots, n-1 \rangle = \langle (H + \epsilon D)^{n-1} : 0 < \epsilon \ll 1 \rangle. \quad \square$$

Zero cycles on curves

Let L be a very ample line bundle on a reduced, projective curve over an algebraically closed field. The zero set of a general section of L consist of $\deg_C L$ distinct points. However, if we work over a non-closed field k , then the zero set of a general section $s \in H^0(C, L)$ is a union of points of the form $\text{Spec}_k k_i$ for some field extensions k_i/k , that depend on the choice of the section in a rather unpredictable way. We may thus aim to find sections $s \in H^0(C, L)$ whose zero set is arithmetically simple. If k is Hilbertian, we can chose the zero set to be irreducible. Another direction would be to find zero sets that consist of low degree points. This is, however, impossible already for genus 1 curves over \mathbb{Q} .

The next result says that, for any finite set of curves C_i and line bundles L_i , a uniformly nice choice of sections is possible.

Theorem 99. *Let C be a geometrically reduced, projective curve over a field k with irreducible components $\{C_i : i \in I\}$. Let L be an ample line bundle on C and $\Sigma \subset C$ a finite set. Then there is an $m > 0$, a separable field extension K/k and a section $s \in H^0(C, L^m)$ such that*

(99.1) $(s = 0)$ is disjoint from $\Sigma \cup \text{Sing } C$, and

(99.2) $C_i \cap (s = 0)$ is isomorphic to the disjoint union of $(m / \deg(K/k)) \cdot \deg_{C_i} L$ copies of $\text{Spec}_k K$ for every i .

Proof. For m_1 large enough there is a separable morphism $\pi : C \rightarrow \mathbb{P}^1$ such that $L^{m_1} \cong \pi^* \mathcal{O}_{\mathbb{P}^1}(1)$. By (102) there is a separable point $p \in \mathbb{P}^1$ that is disjoint from $\pi(\Sigma \cup \text{Sing } C)$ and such that $\pi^{-1}(p)$ is a reduced, disjoint union of copies of p . Let $s' \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(m_2))$ be a defining equation of p . Then $s := \pi^* s' \in H^0(C, L^{m_1 m_2})$ has the required properties. \square

Corollary 100. *Let X be a projective variety over a field k , L an ample line bundle, $\{C_i : i \in I\}$ a finite set of geometrically reduced curves and $\Sigma \subset X$ a finite subset. Then there is an $m > 0$, a section $s \in H^0(X, L^m)$ and a separable field extension K/k such that*

(100.1) $(s = 0)$ is disjoint from $\Sigma \cup \text{Sing}(\cup_i C_i)$, and

(100.2) $C_i \cap (s = 0)$ is isomorphic to the disjoint union of $(m / \deg(K/k)) \cdot \deg_{C_i} L$ copies of $\text{Spec}_k K$ for every i . \square

101. Another consequence of (99) is the following. Let X be a projective variety over a field k . Then every element of the Chow group of 0-cycles of degree 0 has a representative of the form

$$\sum_i ([p_i] - [q_i]) \quad \text{where} \quad k(p_i) \cong k(q_i).$$

The following can be viewed as a generalization of a special case of Chebotarev's density theorem: there are infinitely many completely split primes in any separable field extension.

Proposition 102. [Poo01] *Let C be a geometrically reduced k -curve and $\pi : C \rightarrow \mathbb{P}^1$ a quasi-finite, separable morphism. Then there are infinitely many separable points $p_j \in \mathbb{P}^1$ such that $\pi^{-1}(p_j)$ is a reduced, disjoint union of copies of p_j for every j .*

Proof. Let C_i be the irreducible components of C . Let D be the normalization of \mathbb{P}^1 in the Galois closure of a composite of the $k(C_i)/k(\mathbb{P}^1)$. If p_j works for $\sigma : D \rightarrow \mathbb{P}^1$ and π is étale over p_j then p_j works for $C \rightarrow \mathbb{P}^1$.

If k is infinite, then a general pencil of very ample divisors gives a separable morphism $\rho : D \rightarrow \mathbf{P}^1 := \mathbb{P}^1$ such that

$$(\rho, \sigma) : D \rightarrow D' \subset \mathbf{P}^1 \times \mathbb{P}^1 \quad (102.1)$$

is birational onto its image. (We use the notation \mathbf{P}^1 to distinguish the 2 factors.) Let $S \subset D$ be the union of the preimage of $\text{Sing } D'$, the ramification locus of σ and the ramification locus of ρ .

Pick any $c \in \mathbf{P}^1(k) \setminus \rho(S)$. Let $p_D \in \rho^{-1}(c)$ be any closed point, p'_D its image in D' and $p := \sigma(p_D) \in \mathbb{P}^1$. Then

$$k(p_D) = k(p'_D) = k(c) \otimes_k k(p) = k(p).$$

Since D/\mathbb{P}^1 is Galois, the same holds for all points in $\sigma^{-1}(p)$.

If k is finite, choose $q = p^r$ such that D decomposes into m irreducible components that are geometrically irreducible. Then D has about mq points in \mathbb{F}_q . All these map to \mathbb{F}_q points in \mathbb{P}^1 . We show that for most of them, their image is not defined over a subfield of \mathbb{F}_q . All subfields of \mathbb{F}_q have at most \sqrt{q} elements and the number of maximal ones equals the number of prime divisors of r , so there are at most $\log_2 r$ of them. Thus at most $\log_2 r \cdot \sqrt{q}$ points of $\mathbb{P}^1(\mathbb{F}_q)$ are in a smaller subfield and these have at most $\deg \pi \cdot \log_2 r \cdot \sqrt{q}$ preimages in D . So for $q \gg 1$, almost all \mathbb{F}_q points of D map to points of \mathbb{P}^1 whose residue field is \mathbb{F}_q . \square

103 (Variants of (102)). The following versions are also useful.

Claim 103.1. Let X, Y be geometrically reduced k -schemes and $\pi : X \rightarrow Y$ a quasi-finite, separable morphism. Then there is a Zariski dense set of separable points $p_j \in Y$ such that $\pi^{-1}(p_j)$ is a reduced, disjoint union of copies of p_j .

Proof. We can replace Y by a general curve $B \subset Y$. The curve case is reduced to (102) by composing with a quasi-finite, separable morphism $B \rightarrow \mathbb{P}^1$. \square

Claim 103.2. Let X, Y be geometrically reduced k -schemes and $\pi : X \rightarrow Y$ a quasi-finite, separable morphism. Then there are infinitely many irreducible divisors $D_j \subset Y$ such that $\pi^{-1}(D_j)$ is a reduced union of divisors D_j^i , such that, each $D_j^i \rightarrow D_j$ is birational.

Proof. As in (102) and in (103.1), we may assume that $Y = \mathbb{P}^n$ and $X \rightarrow \mathbb{P}^n$ is Galois. If k is infinite, the proof works as in (102), but we replace (102.1) by

$$(\rho, \sigma) : X \rightarrow X' \subset \mathbf{P}^1 \times \mathbb{P}^n.$$

We can also reduce (103.2) to (103.1) by choosing a coordinate projection $p : \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-1}$ and work with the generic fibers of $Y \dashrightarrow \mathbb{P}^{n-1}$ and $X \dashrightarrow \mathbb{P}^{n-1}$. \square

12. TOPOLOGICAL PENCILS

Definition 104 (Pencils). Let k be a field and X an integral k -variety. Let C be an integral curve and $p : X \dashrightarrow C$ a dominant map with indeterminacy locus $B \subset X$. We care about the map p up to birational equivalence, so we may as well assume that C is nonsingular and projective.

For a point $c \in C(\bar{k})$ let $D_c \subset X$ denote the closure of $p^{-1}(c)$. In traditional terminology (see, for example, [Zar41])

$$|D|_{\text{alg}} := \{D_c : c \in C(\bar{k})\}$$

is an (*algebraic*) *pencil of divisors* parametrized by the curve C . We use $|D|_{\text{alg}}$ to emphasize that C can be a non-rational curve. We refer to the map $p : X \dashrightarrow C$ itself as an (*algebraic*) *pencil*. The pencil is called *linear* if $C \cong \mathbb{P}_k^1$. These are usually denoted by $|D|$.

If p factors as $X \dashrightarrow C' \dashrightarrow C$ then $X \dashrightarrow C'$ determines another pencil $|D'|_{\text{alg}}$ and members of $|D|_{\text{alg}}$ are certain unions of members of $|D'|_{\text{alg}}$. If $C' \dashrightarrow C$ is not birational, we say that $|D|_{\text{alg}}$ is *composite with* $|D'|_{\text{alg}}$.

A theorem of Bertini—which seems to have been first fully proved by [vdW37]—states that the following are equivalent.

(104.1) Almost all members of $|D|_{\text{alg}}$ are irreducible and generically reduced.

(104.2) $|D|_{\text{alg}}$ is not composite with any other pencil.

(104.3) $k(C)$ is algebraically closed in $k(X)$.

As in [Zar41], we call such a pencil *non-composite*.

Every pencil is composite with a unique non-composite pencil; nowadays this is pretty clear using Stein factorization.

Note. The definition of linear system frequently allows fixed components; those without fixed components are called *mobile*. In this terminology, our pencils are the *mobile pencils*.

Definition 105 (Topological pencil). Let X be a projective, geometrically normal k -variety. A *t-pencil* is a collection of effective divisors $\{D_\lambda : \lambda \in \Lambda\}$ such that

(105.1) Every closed point of X is contained in some D_λ .

(105.2) Almost all of the D_λ are irreducible.

(105.3) There is a closed subset $B \subset X$ of codimension ≥ 2 such that $D_\lambda \cap D_\mu \subset B$ for every $\lambda \neq \mu \in \Lambda$. The smallest such B is called the *base locus*.

(105.4) Each $D_\lambda \setminus B$ is connected.

We call a *t-pencil* *ample* if almost all members are ample \mathbb{Q} -Cartier divisors.

Example 106. Let X be a projective, geometrically normal k -variety and $|D|_{\text{alg}}$ a non-composite algebraic pencil with parameter curve C and base locus B .

For $c \in C(\bar{k})$ let $\{D_{c,i} : i = 1, \dots, r_c\}$ be the connected components of $\text{red } D_c \setminus B$. As σ runs through all embeddings $k(c) \hookrightarrow \bar{k}$, we get k^{ins} -divisors $\cup_\sigma D_{c,i}^\sigma$. The set of all $D_{c,i}$ forms a *t-pencil* with base locus B . We denote it by $|D|^t$ and write its members as $\{D_\lambda : \lambda \in \Lambda\}$. These are the *algebraic t-pencils*.

Note that $|D|^t$ is ample iff the members of $|D|_{\text{alg}}$ are ample \mathbb{Q} -Cartier divisors.

The following is proved in [KLOS20, 5.2.8].

Proposition 107. *Let X be a normal, projective variety of dimension n over an infinite field and H an ample divisor. A *t-pencil* $\{D_\lambda : \lambda \in \Lambda\}$ is algebraic iff there*

is an infinite subset $\Lambda^* \subset \Lambda$ such that the intersection numbers $(H^{n-1} \cdot D_\lambda)$ are independent of $\lambda \in \Lambda^*$.

Proof. Let $p : X \dashrightarrow C$ be an algebraic pencil. C has infinitely many points of degree d for some $d > 0$, these give infinitely many divisors of the same degree in the corresponding t-pencil.

The converse is proved in [KLOS20, 5.2.8]. \square

Combining (107) with (96) we get the following.

Theorem 108. *Let X be a normal, projective, geometrically integral variety of dimension $> 1 + \text{BH}(k)$ over a field k of characteristic 0. Then algebraicity of ample t-pencils is a property of $(|X|, \sim_{\text{sa}})$.* \square

Remark 108.1. We use ampleness only to guarantee that almost all members are geometrically connected; a little more work should establish the theorem for non-ample t-pencils as well. If the pencil has a basepoint, then most likely almost all members are geometrically connected, hence we can use (97.3) to decide when they have the same degree. Basepoint-free t-pencils are always algebraic by (53).

The following is a linearity criterion for algebraic t-pencils.

Lemma 109 (Linearity test). *Let X be a normal, projective variety over a perfect field k . Let $|D|_{\text{alg}}$ be a non-composite pencil with parameter curve C and corresponding topological pencil $|D|^t$.*

(109.1) *If $|D|^t$ has a smooth basepoint then $g(C) = 0$.*

(109.2) *If $g(C) = 0$ and there is a geometrically irreducible subvariety $W \subset \text{Bs } |D|^t$ such that some $D_\lambda \in |D|^t$ is generically smooth along W , then $C \cong \mathbb{P}^1$.*

Proof. The first claim is classical, see for example [Kol96, VI.1.9] for a more general assertion.

If $g(C) = 0$ and C has a k -point then $C \cong \mathbb{P}^1$. Every member of $|D|^t$ is obtained as $D_\lambda = \cup_\sigma D_{c,i}^\sigma$ (106). If $D_{c,i}$ contains W then so do its conjugates, so D_λ is singular along W iff $k(c) \neq k$. \square

The following is a very simple topological formula computing the intersection number of a curve with a pencil of divisors.

Lemma 110. *Let X be a normal, projective variety over a field k of characteristic 0, $|D|_{\text{alg}}$ a pencil with parameter curve C and corresponding topological pencil $|D|^t$. Let $A \subset X$ be an irreducible curve disjoint from $\text{Bs } |D|^t$. Then*

$$(A \cdot |D|_{\text{alg}}) = \max\{\#|A \cap D_\lambda| : D_\lambda \in |D|^t\}.$$

Proof. The inequality \geq is clear. Conversely, the pencil is given by a map $X \dashrightarrow C$ that induces a dominant morphism $\pi : A \rightarrow C$. We need to find a $c \in C(\bar{k})$ such that π is étale over c and $k(a_i) = k(c)$ for every $a_i \in \pi^{-1}(c)$. This is possible by (102). \square

True members

Definition 111. Let X be a normal, projective, geometrically integral k -variety and $|D|_{\text{alg}}$ a non-composite algebraic pencil with corresponding t-pencil $|D|^t = \{D_\lambda : \lambda \in \Lambda\}$. D_λ is called a *true member* of $|D|^t$ if there is a k -member $D_c \in |D|_{\text{alg}}$ such that $D_\lambda = \text{red } D_c = D_c$.

If $|D|_{\text{alg}}$ is a linear pencil with parameter curve $C \cong \mathbb{P}^1$, then D_c is a true member for all but finitely many $c \in \mathbb{P}^1(k)$, but in general there may not be any true members.

Definition 112. Let X be a normal, projective, geometrically integral k -variety over an infinite field and L an ample line bundle on X . Let $|D|^t$ be a linear, algebraic t -pencil. Define its L -degree as the smallest number in the set $\{\deg_L D_\lambda : \lambda \in \Lambda\}$ that is taken infinitely many times. We denote it by $\deg_L |D|^t$.

Note that $\deg_L |D|^t = \deg_L |D|$. (For non-linear pencils the definition makes sense but frequently gives a multiple of $\deg_L |D|$.)

Lemma 113 (True membership test). *Let X be a normal, projective variety over a perfect field k and $|D|^t = \{D_\lambda : \lambda \in \Lambda\}$ a linear, algebraic t -pencil.*

(113.1) *Almost all of the $\{D_\mu : \deg_H D_\mu = \deg_L |D|^t\}$ are true k -members of $|D|^t$.*

(113.2) *If such a D_μ is generically smooth along a geometrically irreducible subvariety $W \subset \text{Bs } |D|^t$, then it is a true member of $|D|^t$.*

Proof. The first claim is clear. As in (109), $D_\lambda = \cup_\sigma D_{c,i}^\sigma$ (106) is smooth along W iff $c \in C(k)$. If this holds then $\deg_L D_\lambda \leq \deg_L D_c = \deg_L |D|$ and equality holds iff $D_\lambda = D_c$, that is, iff D_λ is a true member of $|D|^t$. \square

The following example shows that in general there can be false members that are linearly equivalent to true members.

Example 114. Start with the pencil on \mathbb{A}^2

$$|(u^2 + v^2 + u)(v^2 + u), (u^2 + v^2 + v)(u^2 + v)|.$$

Its general member is a quartic with a node at the origin, but it has 2 members that split into conics.

Next make a change of variables $u = x + iy, v = x - iy$. The resulting pencil $|D|$ is still defined over \mathbb{Q} but now it has a conjugate pair of reducible members. Thus we obtain that

$$\begin{aligned} & ((x + iy)^2 + (x - iy)^2 + (x - iy))((x + iy)^2 + (x - iy)^2 + (x + iy)) = \\ & (2x^2 + 2y^2 + x - iy)(2x^2 + 2y^2 + x + iy) = \\ & (2x^2 + 2y^2 + x)^2 + y^2 \quad \text{and} \\ & ((x - iy)^2 + (x + iy))((x + iy)^2 + (x - iy)) = \\ & (x^2 - y^2 + x + i(y - 2xy))(x^2 - y^2 + x + i(y - 2xy)) = \\ & (x^2 - y^2 + x)^2 + (y - 2xy)^2 \end{aligned}$$

both give degree 4 false members of $|D|^t$.

Theorem 115. *Let X be a normal, projective variety over an infinite, perfect field k . Two reduced divisors A_1, A_2 are linearly equivalent iff the following holds.*

There is a codimension 2 subset $\Sigma \subset X$ such that for every irreducible curve $C \not\subset \Sigma \cup A_1 \cup A_2$ there are irreducible divisors A', H, H' and t -pencils $|D_1|^t$ and $|D_2|^t$, such that, for $i = 1, 2$,

(115.1) H, H' are \mathbb{Q} -Cartier and ample,

(115.2) $|D_i|^t$ are algebraic,

(115.3) $A_i + A' + H$ and H' are reduced members of $|D_i|^t$,

(115.4) $(A_i \cup A' \cup H) \setminus (\Sigma \cup H')$ are connected,

(115.5) $C \subset \text{Bs } |D_i|^t$, and

(115.6) $A_i + A' + H$ and H' are generically smooth along C .

Proof. If A_1, A_2 are linearly equivalent then first choose $\Sigma = \text{Sing } X$ and A' such that $A_i + A'$ are reduced and \mathbb{Q} -Cartier. Then choose H sufficiently ample, generically smooth along C and such that $|A_1 + A' + H| = |A_2 + A' + H|$ is very ample and has an irreducible member H' that is generically smooth along C and such that the $(A_i \cup A' \cup H) \setminus (\Sigma \cup H')$ are connected. Set $|D_i| := |A_i + A' + H, H'|$. Then $A_i + A' + H$ and H' are reduced members of $|D_i|^t$ and the rest is clear.

Conversely, choose C to be geometrically irreducible such that X is generically smooth along C . Then the $|D_i|^t$ are linear pencils by (109). Furthermore, $A_i + A' + H$ and H' are true members of $|D_i|^t$ by (113). Thus $A_1 + A' + H \sim H' \sim A_2 + A' + H$ by (115.3), hence $A_1 \sim A_2$. \square

Theorem 116. *Let X be a normal, projective variety of dimension $> 1 + \text{BH}(k)$ over a field k of characteristic 0. Then linear equivalence of divisors is determined by $(|X|, \sim_{\text{sa}})$.*

Proof. Let $A = \sum a_i A_i$ be an effective divisor and fix an ample divisor H . For every i there is an m_i such that $|A_i + m_i H|$ is mobile. Let A_i^* be the sum of a_i distinct, irreducible members of $|A_i + m_i H|$. Then $A + (\sum a_i m_i) H \sim \sum A_i^*$ and $\sum A_i^*$ is reduced. Thus, as in [KLOS20, 5.3.8], we see that linear equivalence of all divisors is determined by linear equivalence of reduced divisors.

Thus it remains to show that linear equivalence of reduced divisors is determined by $(|X|, \sim_{\text{sa}})$. For this we need to check that the conditions (115.1–6) are determined by $(|X|, \sim_{\text{sa}})$.

For (115.1) we use (67) and for (115.2) we need (108). Conditions (115.3–5) are topological. Finally for (115.6) we can use (93). \square

13. COMPLETE INTERSECTIONS

In this section we collect various results on complete intersections that were used earlier.

117 (Complete intersections). Let X be an irreducible variety. A subscheme $Z \subset X$ of codimension r is a *complete intersection* (resp. *set-theoretic complete intersection*) if there are effective Cartier divisors D_1, \dots, D_r such that $Z = D_1 \cap \dots \cap D_r$ scheme theoretically (resp. $\text{Supp } Z = \text{Supp}(D_1 \cap \dots \cap D_r)$).

If the D_i are ample, we call Z an ample (set-theoretic) complete intersection, usually abbreviated as *ample-ci* resp. *ample-sci*.

If H is a Cartier divisor and $D_i \in |m_i H|$ for every i , then we say that Z is a *complete H -intersection*. These we usually abbreviate as *H -ci*, the set-theoretic versions as *H -sci*.

Ample complete intersections inherit many properties of a variety, but the strongest results are for general complete intersections; that is, when the $D_i \in |m_i H|$ are sufficiently general.

118 (General complete intersections). We say that a *general complete intersection has property \mathcal{P}* if the following holds.

(118.1) Let X be a projective variety and H_i ample divisors on X . Then for $m_i \gg 1$ there is an open, dense subset $U \subset \prod_i |m_i H_i|$ such that if $D_i \in |m_i H_i|(\bar{k})$ and $(D_1, \dots, D_r) \in U$ then $Z := D_1 \cap \dots \cap D_r$ satisfies \mathcal{P} .

There is a long list of such properties \mathcal{P} ; below is a list of those that we use. They usually go by the name *Bertini theorems*. Most of these are proved in [Har77, II.8.18, III.10.9], but for (118.5) one needs to use that a general hypersurface section of an S_2 scheme is also S_2 ; see [Gro60, IV.12.1.6].

- (118.2) Given finitely many irreducible subvarieties $W_j \subset X$, we have $\dim(Z \cap W_j) = \dim W_j - r$ (or the intersection is empty).
- (118.3) Given finitely many irreducible subvarieties $W_j \subset X$ of dimension $\geq r + 1$, the $Z \cap W_j$ are irreducible.
- (118.4) Given finitely many locally closed, smooth subvarieties $W_j \subset X$, the $Z \cap W_j$ are smooth.
- (118.5) Given finitely many locally closed, normal subvarieties $W_j \subset X$, the $Z \cap W_j$ are normal.

In many applications, we need complete intersections that are special in some respects but general in some others. Here we deal with local conditions.

- (118.6) A set LC of *local conditions* consist of finitely many smooth points $p_j \in X$ with maximal ideals $m_j \subset \mathcal{O}_{p_j, X}$, natural numbers n_j and regular sequences $g_{j1}, \dots, g_{jr} \in \mathcal{O}_{p_j, X}$.

For example, we can specify whether $p_i \in Z$ or that Z be smooth at p_i with given tangent space.

A codimension r complete intersection $Z = D_1 \cap \dots \cap D_r$ satisfies LC if g_{ji} is a local equation for D_i at p_j , modulo $m_j^{n_j}$.

For every i the local conditions define a linear subspace $(|m_i H_i|, \text{LC}) \subset |m_i H_i|$ (which may be empty). The complete intersections that satisfy LC form an open, dense subset of

$$\prod_i (|m_i H_i|, \text{LC}) \subset \prod_i |m_i H_i|.$$

The following combines the local conditions with (118.2–5).

- (118.7) Let X be a projective variety, H_i ample divisors on X , LC a set of local conditions at the points $\{p_j\}$ and \mathcal{P} any of the properties (118.2–5). Then for $m_i \gg 1$ there is an open, dense subset $U \subset \prod_i (|m_i H_i|, \text{LC}_i)$ such that if $(D_1, \dots, D_r) \in U(\bar{k})$ then
 - (a) $Z := D_1 \cap \dots \cap D_r$ satisfies \mathcal{P} on $X \setminus \{p_j\}$, and
 - (b) $(I_{D_i}, m_j^{n_j}) = (g_{ji}, m_j^{n_j})$ for every j . □

Finite fields 118.8. If k is finite, then a dense open subset of \mathbb{A}^n may be disjoint from $\mathbb{A}^n(k)$. Nonetheless, the results of [Poo08, CP16] say that the open sets U above have k -points.

Our proofs have other problems with finite fields, so we will be able to make only very limited use of these cases.

119 (Connectedness). Let Z be a scheme. Connectedness and irreducibility of Z depends only on the topological space $|Z|$, but geometric connectedness and geometric irreducibility can not be determined using $|Z|$ only.

We frequently need to guarantee that certain schemes are geometrically connected. The next criterion can be proved by repeatedly using [Har77, II.7.8]; see also [Har62].

Claim 119.1. Let X be a normal, projective, geometrically irreducible variety and $Z \subset X$ a positive dimensional, ample-sci. Then Z is geometrically connected. □

Note that a proper k -scheme Y is geometrically connected iff $H^0(Y, \mathcal{O}_Y)$ is a local, Artin k -algebra such that $H^0(Y, \mathcal{O}_Y)/\sqrt{0}$ is a purely inseparable field extension of k . We can thus restate (119.1) as follows.

Claim 119.2. Let X be a normal, projective, geometrically irreducible variety and $Z \subset X$ a positive dimensional, reduced, ample-sci. Then $k[Z]/k[X]$ is purely inseparable. \square

We also use the following variant of the Lefschetz hyperplane theorem. For the Picard variety, this is proved in [Gro68]. For normal varieties in characteristic 0, the class group version is proved in [RS06, RS09]. For positive characteristic see [Ji20].

Theorem 120. *Let X be a geometrically normal, projective variety and $|H|$ an ample linear system on X . Let $Z \subset X$ be a normal, ≥ 2 dimensional, complete H -intersection. Then the restriction map*

$$\mathrm{Cl}(X) \hookrightarrow \mathrm{Cl}(Z) \quad \text{is injective.}$$

121 (Disjointness of conjugates). Let K/k be a finite, separable field extension and $A \in \mathrm{Mat}_{r \times n}(K)$ an $r \times n$ matrix. Then $A\mathbf{x} = 0$ defines a linear subspace $L_A \subset K^n$, usually of codimension r . Let $\sigma : K \hookrightarrow \bar{k}$ be a k -embedding, it defines A^σ and $L_A^\sigma \subset K^n$. We aim to show that usually $L_A \cap L_A^\sigma$ has codimension $2r$.

After choosing a k -basis e_i for K , we can write $A = \sum_i e_i A_i$ where $A_i \in \mathrm{Mat}_{r \times n}(k)$.

Claim 121.1. There is a Zariski open subset $U \subset \bigoplus_i \mathrm{Mat}_{r \times n}(k)$ such that if $A \in U$ then $\mathrm{codim}(L_A \cap L_A^\sigma) = \min\{2r, n\}$ for all non-identity k -embeddings $\sigma : K \hookrightarrow \bar{k}$.

Proof. Let us work out the critical case when $n = 2r$. We use $x_1, \dots, x_r, y_1, \dots, y_r$ as variables. Write the equations as $A\mathbf{x} + B\mathbf{y} = 0$ where $A, B \in \mathrm{Mat}_{r \times r}(K)$. The conjugate equations are $A^\sigma \mathbf{x} + B^\sigma \mathbf{y} = 0$. We need to show that

$$\mathrm{rank} \begin{pmatrix} A & B \\ A^\sigma & B^\sigma \end{pmatrix} = 2r$$

is a non-empty, Zariski open condition. It is clearly an open condition, so need to find one example with maximal rank. Take $A, C \in \mathrm{Mat}_{r \times r}(k)$ and $B = \alpha C$. By row reduction

$$\begin{pmatrix} A & B \\ A^\sigma & B^\sigma \end{pmatrix} = \begin{pmatrix} A & \alpha C \\ A & \alpha^\sigma C \end{pmatrix} \rightsquigarrow \begin{pmatrix} A & \alpha C \\ 0 & (\alpha^\sigma - \alpha)C \end{pmatrix}$$

whose determinant is $(\alpha^\sigma - \alpha)^r \det A \det C$. \square

The intrinsic way of passing from $\mathrm{Mat}_{r \times n}(K)$ to $\bigoplus_i \mathrm{Mat}_{r \times n}(k)$ is the *Weil restriction*, denoted by $\mathcal{R}_k^K(\cdot)$; see, for instance [BLR90, Sec.7.6]. We can thus globalize (121.1) first to projective spaces and then to their subvarieties as follows.

Claim 121.2. Let X be a k -variety of pure dimension n and $|M_1|, \dots, |M_r|$ basepoint-free linear systems. Let K/k be a finite, separable field extension. Then there is a dense, Zariski open subset

$$U \subset \mathcal{R}_k^K |M_1| \times \cdots \times \mathcal{R}_k^K |M_r|,$$

such that, if $(D_1, \dots, D_r) \in U$, then

$$\mathrm{codim}_X(D_1 \cap \cdots \cap D_r \cap D_1^\sigma \cap \cdots \cap D_r^\sigma) = \min\{2r, n+1\},$$

for all non-identity k -embeddings $\sigma : K \hookrightarrow \bar{k}$. \square

Lemma 122. *Let k be an infinite field and X a normal, projective, irreducible k -variety of dimension $n > 2r$. Let $p, q \in X$ be closed points such that there are embeddings $k \subset k(p) \subset k(q)^{\text{ins}} \subset \bar{k}$.*

Then there is an irreducible, r -dimensional k -variety $W \subset X$ such that

(122.1) $p, q \in W$ and

(122.2) $k(p)/k[W]$ is purely inseparable.

Furthermore, if p is a smooth, separable point of X then we can also assume that

(122.3) p is a smooth point of W .

Proof. Let $k \subset K_p \subset k(p)$ and $k \subset K_q \subset k(q)$ be maximal separable subextensions. After base change to K_p , we have a degree 1 point \bar{p} lying over p and a degree $= \deg(K_q/K_p)$ point \bar{q} lying over q . Let $\{\sigma\}$ be the set of all k -embeddings $\sigma : K_p \hookrightarrow \bar{k}$. Thus \bar{p} and \bar{q} each have $\deg(K_p : k)$ conjugates over k and these are disjoint from each other.

Next take a general ample-ci variety (117) $W_1 \subset X_{K_p}$ that contains \bar{p} and \bar{q} . By (121.2) the W_1^σ are disjoint from each other. Thus their union $W_{K_p} = \cup_\sigma W_1^\sigma$ descends to a k -subvariety $W \subset X$ with the required properties. \square

14. PICARD GROUP, CLASS GROUP AND ALBANESE VARIETY

For the Picard group and Picard scheme, [Gro62, Lects.V-VI], [Mum66, Sec.19], [Mum70] or [BLR90] contain proofs and details; for these we just fix our notation. Modern references for the class group and Albanese variety are harder to find; about these we give more details.

123 (Picard group of a normal variety). The group of line bundles on a scheme X is the *Picard group* of X , denoted by $\text{Pic}(X)$. If X is proper then $\text{Pic}^\circ(X) \subset \text{Pic}(X)$ denotes the subgroup of divisors that are algebraically equivalent to 0. The quotient $\text{NS}(X) := \text{Pic}(X)/\text{Pic}^\circ(X)$ is the *Néron-Severi* group of X . It is a finitely generated abelian group.

If X is proper over a field k with algebraic closure \bar{k} then $\text{Pic}(X_{\bar{k}})$ has a natural k -scheme structure, denoted by $\mathbf{Pic}(X)$. The identity component is denoted by $\mathbf{Pic}^\circ(X)$, it is a commutative algebraic group. If X is geometrically normal and $\text{char } k = 0$ then $\mathbf{Pic}^\circ(X)$ is an Abelian variety. If X is geometrically normal and k is perfect then $\text{red } \mathbf{Pic}^\circ(X)$ is an Abelian variety. The non-reduced structure of $\mathbf{Pic}^\circ(X)$ will play no role in our questions.

There is a natural inclusion $\text{Pic}(X) \hookrightarrow \mathbf{Pic}(X)(k)$ which is an isomorphism if X has a k -point. In general the quotient $\mathbf{Pic}(X)(k)/\text{Pic}(X)$ is a torsion group.

Claim 123.1. For Abelian varieties, $A \mapsto \mathbf{Pic}^\circ(A)$ is a duality.

Proof. Over \mathbb{C} , this is a special case of the Appell-Humbert theorem [Mum70, pp.21-22]. In general see [Mum70, Sec.13]. \square

124 (Class group of a normal variety). For a normal and proper k -variety X , let $\text{Cl}(X)$ denote the group of Weil divisors modulo linear equivalence. It is also isomorphic to the group of rank 1 reflexive sheaves, where the product is the double dual of the tensor product.

Let $\mathrm{Cl}^\circ(X) \subset \mathrm{Cl}(X)$ be the subgroup of divisors that are algebraically equivalent to 0. I call the quotient $\mathrm{NS}^{\mathrm{cl}}(X) := \mathrm{Cl}(X)/\mathrm{Cl}^\circ(X)$ the *Néron-Severi class group*¹ of X , and its \mathbb{Q} -rank the *class rank* of X , denoted by $\rho^{\mathrm{cl}}(X)$.

Note that we have natural inclusions

$$\mathrm{Pic}^\circ(X) \subset \mathrm{Cl}^\circ(X) \quad \text{and} \quad \mathrm{NS}(X) \subset \mathrm{NS}^{\mathrm{cl}}(X), \quad (124.1)$$

that are isomorphisms iff every Weil divisor is Cartier, for example when X is smooth.

Basic results about these groups are the following.

Claim 124.2. Let $p : Y \rightarrow X$ be a birational morphism of normal, proper varieties over a perfect field k . Then

- (a) $p_* : \mathrm{Cl}^\circ(Y) \rightarrow \mathrm{Cl}^\circ(X)$ is an isomorphism and
- (b) $p_* : \mathrm{NS}^{\mathrm{cl}}(Y) \rightarrow \mathrm{NS}^{\mathrm{cl}}(X)$ is onto.

Claim 124.3. Let X be a normal, proper variety over a perfect field k . Then there is normal, proper variety Y and a birational morphism $p : Y \rightarrow X$ such that $\mathrm{Cl}^\circ(Y_K) = \mathrm{Pic}^\circ(Y_K)$ for every $K \supset k$.

It is quickest to prove these by using the Albanese variety, see (128.5–6). As a consequence, we can define the scheme structure of Cl° by

$$\mathrm{Cl}^\circ(X) \cong \mathbf{Cl}^\circ(Y) \cong \mathrm{red} \mathbf{Pic}^\circ(Y). \quad (124.4)$$

In the complex case these results go back to Picard [Pic95] and Severi [Sev06], but the most complete references may be the papers of Matsusaka [Mat52] and of Néron [Nér52a]; see also [Kol18, Sec.3] for some discussions.

More recent results on various aspects of the class group of singular varieties are discussed in [BVS93, BVRs09, RS09].

Definition 125. Let X be a normal, proper k -variety and $\Sigma \subset X$ a subset. Let $\mathrm{WDiv}(X, \Sigma) \subset \mathrm{WDiv}(X)$ and $\mathrm{Cl}(X, \Sigma) \subset \mathrm{Cl}(X)$ denote the subgroup of those Weil divisors that are Cartier at every point $x \in \Sigma$. (The concept and notation are not standard.)

Note that $\mathrm{Cl}(X, \Sigma)$ is isomorphic to the group of those rank 1 reflexive sheaves that are locally free at every point $x \in \Sigma$.

We see in (126) that $\mathrm{Cl}(X_{\bar{k}}, \Sigma_{\bar{k}})$ is naturally identified with a closed k -subgroup $\mathbf{Cl}(X, \Sigma) \subset \mathbf{Cl}(X)$. We denote its identity component by $\mathbf{Cl}^\circ(X, \Sigma)$. Note that in general $\mathbf{Cl}(X, \Sigma) \cap \mathbf{Cl}^\circ(X)$ may be disconnected.

The quotient $\mathrm{NS}^{\mathrm{cl}}(X, \Sigma) := \mathbf{Cl}(X, \Sigma)/\mathbf{Cl}^\circ(X, \Sigma)$ is finitely generated.

Lemma 126. *Let X be a normal, proper variety over a perfect field k and $\Sigma \subset X$ an arbitrary subset. Then there is a closed, algebraic k -subgroup $\mathbf{Cl}(X, \Sigma) \subset \mathbf{Cl}(X)$ such that $\mathrm{Cl}(X_{\bar{k}}, \Sigma_{\bar{k}}) = \mathbf{Cl}(X, \Sigma)(\bar{k})$.*

Proof. Assume first that $\Sigma = \{x\}$ is a closed point and there is a universal family \mathbf{L} on $X \times \mathbf{Cl}^\circ(X)$ that is flat over $\mathbf{Cl}^\circ(X)$. The set of points $V \subset X \times \mathbf{Cl}^\circ(X)$ where \mathbf{L} is not locally free is closed. Since $\mathbf{Cl}(X, \{x\})$ is the complement of the image of $V \cap (\{x\} \times \mathbf{Cl}^\circ(X))$, it is constructible. It is also a subgroup and a constructible subgroup is closed.

¹The literature seems inconsistent. Frequently this is called the Néron-Severi group.

In general such an \mathbf{L} does not exist, but we check in (127) that a flat universal family exists after a finite field extension and a constructible subdivision $\tau : \coprod_j W_j \rightarrow \mathbf{CI}^\circ(X)$. The argument above then shows that $\mathbf{CI}(X, \{x\})$ is constructible, hence closed as before.

If Σ is any set of closed points then $\mathbf{CI}(X, \Sigma) = \bigcap_{x \in \Sigma} \mathbf{CI}(X, \{x\})$.

If η is a non-closed point, then $\mathbf{CI}(X, \{\eta\})$ is the union of all $\mathbf{CI}(X, \Sigma_U)$, where U runs through all open subsets of $\bar{\eta}$ and Σ_U denotes the set of closed points of U . By the Noetherian property, $\mathbf{CI}(X, \{\eta\}) = \mathbf{CI}(X, \Sigma_U)$ for some U .

Finally, $\mathbf{CI}(X, \Sigma) = \bigcap_{x \in \Sigma} \mathbf{CI}(X, \{x\})$ holds for any set of points Σ . \square

Lemma 127. *Let X be a normal, proper variety over an algebraically closed field K . There is a locally closed decomposition $\tau : \coprod_j W_j \rightarrow \mathbf{CI}^\circ(X)$ such for every j there is a universal family \mathbf{L}_j on $X \times W_j$ that is flat over W_j and whose fiber over $w \in W_j$ is the reflexive sheaf corresponding to $\tau(w) \in \mathbf{CI}^\circ(X)$.*

Proof. By (124.3), there is a proper, birational morphism from a normal variety $p : Y \rightarrow X$ such that $\mathbf{CI}^\circ(X) = \mathbf{Pic}^\circ(Y)$. Let \mathbf{L} be the universal line bundle over $Y \times \mathbf{Pic}^\circ(Y)$. Pushing it forward we get a rank 1 sheaf

$$\mathbf{L}_X := (\pi_* \mathbf{L})^{[**]} \quad \text{over} \quad X \times \mathbf{CI}^\circ(X).$$

In general \mathbf{L}_X is not flat over $\mathbf{CI}^\circ(X)$. However, by generic flatness, \mathbf{L}_X is flat with reflexive fibers over a dense, open subset $W_1 \subset \mathbf{CI}^\circ(X)$. Repeating this with $\mathbf{CI}^\circ(X) \setminus W_1$ we get the required locally closed decomposition. \square

Albanese variety

128 (Albanese variety). Let X be a proper, normal variety over a perfect field k . There are 2 different notions of the *Albanese variety* of X in the literature. In [Gro62, VI.3.3] it is the target of the universal *morphism* from X to an Abelian torsor; that is, a principal homogeneous space under an Abelian variety. We denote this by

$$\text{alb}_X^{\text{gr}} : X \rightarrow \mathbf{Alb}^{\text{gr}}(X). \quad (128.1)$$

Pull-back by alb_X^{gr} gives an isomorphism

$$\mathbf{Pic}^\circ(\mathbf{Alb}^{\text{gr}}(X)) \cong \text{red } \mathbf{Pic}^\circ(X). \quad (128.2)$$

If X has a k -point then $\mathbf{Alb}^{\text{gr}}(X)$ an Abelian variety. $\mathbf{Alb}^{\text{gr}}(X)$ is a birational invariant for smooth, proper varieties, but not a birational invariant for normal varieties.

In the older literature, for example [Mat52, Ser59], the Albanese map is the universal *rational map* from X to an Abelian torsor, called the *classical Albanese variety*

$$\text{alb}_X : X \dashrightarrow \mathbf{Alb}(X). \quad (128.3)$$

If X has a smooth k -point then so does $\mathbf{Alb}(X)$ and then it is an Abelian variety.

$\mathbf{Alb}(X)$ is a birational invariant of X (for normal, proper varieties) and the two versions coincide if X is smooth. Therefore, if $X' \rightarrow X$ is a resolution then $\mathbf{Alb}(X) = \mathbf{Alb}(X') = \mathbf{Alb}^{\text{gr}}(X')$. In any case, we get a morphism over the smooth locus

$$\text{alb}_X : X^{\text{sm}} \rightarrow \mathbf{Alb}(X). \quad (128.4)$$

Let X' be the normalization of the closure of the graph of alb_X . Then we have a commutative diagram

$$\begin{array}{ccc} & X' & \\ p \swarrow & & \searrow \text{alb}_{X'} \\ X & \xrightarrow{\text{alb}_X} & \mathbf{Alb}(X) = \mathbf{Alb}(X'), \end{array} \quad (128.5)$$

where $\text{alb}_{X'}$ is a morphism. In particular (128.2) gives that

$$\mathbf{Cl}^\circ(X') = \text{red } \mathbf{Pic}^\circ(X') \cong \mathbf{Pic}^\circ(\mathbf{Alb}(X')). \quad (128.6)$$

Therefore

$$\mathbf{Cl}^\circ(X) \cong \mathbf{Pic}^\circ(\mathbf{Alb}(X)). \quad (128.7)$$

Let $p : Y \dashrightarrow X$ be a map of normal varieties. As long as $p(Y)$ is not contained in the singular locus of X , the composite $\mathbf{Alb}_X \circ p : Y \dashrightarrow \mathbf{Alb}(X)$ is defined, hence we get a morphism

$$\text{alb}_p : \mathbf{Alb}(Y) \rightarrow \mathbf{Alb}(X). \quad (128.8)$$

129. Let $p : Y \rightarrow X$ be a morphism of normal varieties. Let

$$\text{alb}_{Y \setminus X}^{\text{gr}} : X \rightarrow \mathbf{Alb}^{\text{gr}}(Y \setminus X) \quad (129.1)$$

denote the universal morphism from X to an Abelian torsor that maps every irreducible component of Y to a point. Thus we get

$$\mathbf{Alb}^{\text{gr}}(Y) \rightarrow \mathbf{Alb}^{\text{gr}}(X) \rightarrow \mathbf{Alb}^{\text{gr}}(Y \setminus X). \quad (129.2)$$

We claim that the dual sequence

$$0 \rightarrow \mathbf{Pic}^\circ(\mathbf{Alb}^{\text{gr}}(Y \setminus X)) \rightarrow \mathbf{Pic}^\circ(X) \rightarrow \mathbf{Pic}^\circ(Y) \quad (129.3)$$

is exact. To see this, let K denote the kernel of $\mathbf{Pic}^\circ(X) \rightarrow \mathbf{Pic}^\circ(Y)$. It is clear that $\mathbf{Pic}^\circ(\mathbf{Alb}^{\text{gr}}(Y \setminus X)) \subset K$. To see the converse, we may assume that X has a k -point. By (123.1) we get an exact sequence

$$\mathbf{Alb}^{\text{gr}}(Y) \rightarrow \mathbf{Alb}^{\text{gr}}(X) \rightarrow \mathbf{Pic}^\circ(K) \rightarrow 0.$$

The resulting $Y \rightarrow \mathbf{Pic}^\circ(K)$ maps every irreducible component of Y to a point, so it factors through $\mathbf{Alb}^{\text{gr}}(Y \setminus X)$. By duality we get $K \rightarrow \mathbf{Pic}^\circ(\mathbf{Alb}^{\text{gr}}(Y \setminus X))$. \square

We would like to know when alb_p is dominant. Lefschetz theory suggests that this should hold if $p(Y)$ is ample-ci (117). We will need two generalizations of this.

Lemma 130. *Let X, Y be normal, projective varieties and $p : Y \rightarrow X$ a morphism. Assume that $p(Y)$ has nonempty intersection with every nonzero divisor in X and alb_X is a morphism along $p(Y)$. Then $\text{alb}_p : \mathbf{Alb}(Y) \rightarrow \mathbf{Alb}(X)$ is surjective.*

Proof. If alb_p is not surjective then the quotient $\mathbf{Alb}(X)/\text{alb}_p(\mathbf{Alb}(Y))$ is positive dimensional. Hence there is a nonzero, effective divisor $D \subset \mathbf{Alb}(X)/\text{alb}_p(\mathbf{Alb}(Y))$ whose pull-back to $\mathbf{Alb}(X)$ is disjoint from $\text{alb}_p(\mathbf{Alb}(Y))$. Then its pull-back to X is a divisor which is disjoint from $p(Y)$. \square

Corollary 131. *Let X be a normal, projective variety and $C \subset X^{\text{ns}}$ an irreducible ample-sci curve (117). Then $\mathbf{Cl}^\circ(X) \rightarrow \mathbf{Jac}(C)$ is injective modulo torsion.* \square

Lemma 132. *Let X be a normal, proper variety. Then there is a finite subset $\Sigma \subset X$ such that the following holds.*

Let $Y \subset X$ be an irreducible divisor that is disjoint from Σ . Assume that Y has nonempty intersection with every nonzero divisor in X . Then $\mathbf{Alb}(\bar{Y}) \rightarrow \mathbf{Alb}(X)$ is surjective.

Proof. Consider the normalization of the closure of the graph of alb_X

$$X \xleftarrow{\pi} X' \xrightarrow{\text{alb}_X} \mathbf{Alb}(X).$$

Let $E'_i \subset X'$ be the π -exceptional divisors. Choose Σ to contain the generic point of each $\pi(E'_i)$ and every non-Cartier center (136).

Then Y is a Cartier divisor and $\pi^{-1}(Y) = \pi_*^{-1}(Y)$. Therefore, if $D' \subset X'$ is a divisor then

$$\pi(\pi_*^{-1}(Y) \cap D') = \pi(\pi^{-1}(Y) \cap D') = Y \cap \pi(D') \neq \emptyset.$$

Let \bar{Y}' denote the normalization of $\pi_*^{-1}(Y)$. Then $\mathbf{Alb}(\bar{Y}') \rightarrow \mathbf{Alb}(X)$ is surjective by (130) and $\mathbf{Alb}(\bar{Y}') \cong \mathbf{Alb}(\bar{Y})$. \square

Definition 133 (Partial Albanese variety). Let X be a proper, normal variety over a perfect field k and $\Sigma \subset X$ a subset.

Define the *Albanese map* of (X, Σ) as the universal rational map from X to an Abelian torsor, that is a morphism along Σ

$$\text{alb}_{X, \Sigma} : X \dashrightarrow \mathbf{Alb}(X, \Sigma). \quad (133.1)$$

If $\Sigma \subset X^{\text{sm}}$ then $\mathbf{Alb}(X, \Sigma) = \mathbf{Alb}(X)$. In general $\mathbf{Alb}(X, \Sigma)$ is a quotient of $\mathbf{Alb}(X)$.

Theorem 134. *Let X be a normal, proper variety over a perfect field K and $\Sigma \subset X$ a subset. Then pull-back by $\text{alb}_{X, \Sigma}$ gives an isomorphism*

$$\text{alb}_{X, \Sigma}^* \mathbf{Pic}^\circ(\mathbf{Alb}(X, \Sigma)) \cong \mathbf{Cl}^\circ(X, \Sigma). \quad (134.1)$$

Proof. Consider $\text{alb}_{X, \Sigma} : X \dashrightarrow \mathbf{Alb}(X, \Sigma)$. By assumption it is a morphism along Σ , thus the pull-back of a line bundle on $\mathbf{Alb}(X, \Sigma)$ is locally free along Σ . That is,

$$\text{alb}_{X, \Sigma}^* \mathbf{Pic}^\circ(\mathbf{Alb}(X, \Sigma)) \subset \mathbf{Cl}^\circ(X, \Sigma). \quad (134.2)$$

For the converse, assume first that $\Sigma = \{x\}$ is a closed point. As in (128.5) we have

$$\begin{array}{ccc} & X' & \\ p \swarrow & & \searrow \text{alb}_{X'} \\ X & \xrightarrow{\text{alb}_X} & \mathbf{Alb}(X) = \mathbf{Alb}(X'), \end{array}$$

where $\text{alb}_{X'}$ is a morphism. Let Y' be the normalization of $p^{-1}(x)$. Since $X' \rightarrow \mathbf{Alb}^{\text{gr}}(Y' \setminus X')$ contracts every irreducible component of Y' to a point, the composite $X \dashrightarrow X' \rightarrow \mathbf{Alb}^{\text{gr}}(Y' \setminus X')$ is a morphism at x by Zariski's main theorem. This gives $\mathbf{Alb}(X, \{x\}) \rightarrow \mathbf{Alb}^{\text{gr}}(Y' \setminus X')$. Thus we get a commutative diagram

$$\begin{array}{ccc} X' & \rightarrow & \mathbf{Alb}^{\text{gr}}(Y' \setminus X') \\ \downarrow & & \uparrow \\ X & \rightarrow & \mathbf{Alb}(X, \{x\}). \end{array} \quad (134.3)$$

If $L \in \mathbf{Cl}^\circ(X, \{x\})(\bar{k})$ then its pull-back to X' is trivial along Y' , hence it is obtained as the pull-back of a line bundle on $\mathbf{Alb}^{\text{gr}}(Y' \setminus X')$. Factoring through $\mathbf{Alb}(X, \{x\})$ shows that $\mathbf{Cl}^\circ(X, \{x\}) \subset \text{alb}_{X, \{x\}}^* \mathbf{Pic}^\circ(\mathbf{Alb}(X, \{x\}))$.

The same argument works for any finite number of closed points. If Σ is an infinite set of closed point then, by the Noetherian property, $\mathbf{CI}^\circ(X, \Sigma) = \mathbf{CI}^\circ(X, \Sigma')$ for every large enough finite subset $\Sigma' \subset \Sigma$.

Finally assume that y is a non-closed point. Then $\mathbf{CI}^\circ(X, \{y\})$ is the union of all $\mathbf{CI}^\circ(X, \Sigma_U)$ where Σ_U is the set of all closed points in some open subset $U \subset \overline{\{y\}}$. By the Noetherian property, we have equality $\mathbf{CI}^\circ(X, \{y\}) = \mathbf{CI}^\circ(X, \Sigma_U)$ for some fixed U . \square

Corollary 135. *Let X be a normal, projective variety over a perfect field and $Z \subset X$ a closed, reduced subscheme with generic points g_Z . Then there is a normal, projective variety X' , a birational morphism $p : X' \rightarrow X$ and a closed, reduced subscheme $Z' \subset X'$ with generic points $g_{Z'}$ such that*

(135.1) p is a local isomorphism at all generic points of Z' ,

(135.2) $Z = p(Z')$,

(135.3) $\text{alb}_{X', g_{Z'}}$ is a morphism along Z' and

(135.4) $\mathbf{CI}^\circ(X, g_Z) = \mathbf{CI}^\circ(X', g_{Z'}) = \mathbf{CI}^\circ(X', Z')$.

If either $\dim Z = 1$ or the characteristic is 0, we can also achieve that

(135.5) Z' is smooth,

Proof. We can take X' to be the normalization of the closure of the graph of alb_{X, g_Z} . Then we can resolve the singularities of Z' if desired. \square

Non-Cartier centers

Definition 136 (Non-Cartier centers). Let X be a reduced scheme and D an effective Weil divisor. There is a unique largest open subscheme $X_D^{\text{car}} \subset X$, called the *Cartier locus* of D such that the restriction of D to X_D^{car} is Cartier. The complement $X \setminus X_D^{\text{car}}$ is the *non-Cartier locus* of D . A point $x \in X$ is a *non-Cartier center* of X if there is a Weil divisor D such that x is the generic point of an irreducible component of the non-Cartier locus of D .

For example, let $X = (xy = 0) \subset \mathbb{A}_{xyz}^3$ and set $D_c := (x = z - c = 0)$. Its non-Cartier locus is the point $(x = y = z - c = 0)$. Thus every closed point of the z -axis is a non-Cartier center of X . The generic point of the z -axis is also a non-Cartier center of X for the divisor $(x = y = 0)$.

In direct analogy one can define the notions of \mathbb{Q} -Cartier locus and non- \mathbb{Q} -Cartier center.

The next result of [BGS11, 6.7] shows that the situation is quite different for normal varieties. (Note that [BGS11] works over an algebraically closed field, but this is not necessary.)

Theorem 137. *A geometrically normal variety has only finitely many non-Cartier or non- \mathbb{Q} -Cartier centers.*

Proof. We may assume that X is proper and irreducible. Let $U \subset X$ be an open subset such that X has only finitely many non- (\mathbb{Q}) -Cartier or centers in U . We show that there is a strictly larger open subset $U \subsetneq U' \subset X$ such that X has only finitely many non- (\mathbb{Q}) -Cartier centers in U' . We can start with the smooth locus $U = X^{\text{sm}}$, since it is disjoint from every non- (\mathbb{Q}) -Cartier center. Noetherian induction then gives that X has only finitely many non- (\mathbb{Q}) -Cartier centers.

Let $Z \subset X \setminus U$ be an irreducible component. By (138) there is a dense, open subset $Z^0 \subset Z$ such that if a Weil divisor D is (\mathbb{Q}) -Cartier at the generic point $g_Z \in Z$ then it is (\mathbb{Q}) -Cartier along Z^0 . We may assume that Z^0 is disjoint from every other irreducible component of $X \setminus U$. Then $U' := U \cup Z^0$ is open in X and g_Z is the only possible new non- (\mathbb{Q}) -Cartier center in U' . \square

Lemma 138. *Let X be a normal, proper variety over an algebraically closed field and $Z \subset X$ an irreducible subvariety. Then there is a dense, open subset $Z^0 \subset Z$ such that the following holds.*

Let D be a Weil divisor that is (\mathbb{Q}) -Cartier at the generic point $g_Z \in Z$. Then it is (\mathbb{Q}) -Cartier everywhere along Z^0 .

Proof. As in (125), let $\mathbf{Cl}(X, g_Z) \subset \mathbf{Cl}(X)$ be the subgroup of those divisors that are Cartier at the generic point of Z and $\mathbf{Cl}^\circ(X, g_Z) \subset \mathbf{Cl}^\circ(X)$ the identity component.

As we noted in (125), the quotient $\mathbf{Cl}(X, g_Z)/\mathbf{Cl}^\circ(X, g_Z)$ is finitely generated; say by the divisors D_i . There is a dense, open subset $Z_1^0 \subset Z$ such that every D_i is Cartier along Z_1^0 , hence the same holds for every linear combination of the D_i .

Next we show that there is a dense, open subset $Z_2^0 \subset Z$ such that every divisor in $\mathbf{Cl}^\circ(X, g_Z)$ is Cartier along Z_2^0 . Consider the Albanese map $\mathrm{alb}_{X,Z} : X \dashrightarrow \mathbf{Alb}(X, Z)$. By (133) it is defined at g_Z , hence on a dense, open subset $Z_2^0 \subset Z$. By (134), $\mathbf{Cl}^\circ(X, g_Z)$ is the pull-back of $\mathbf{Pic}^\circ(\mathbf{Alb}(X, g_Z))$, hence every member of $\mathbf{Cl}^\circ(X, g_Z)$ is locally free along Z_2^0 . Finally $Z^0 = Z_1^0 \cap Z_2^0$ is the dense, open subset that we need.

If D is \mathbb{Q} -Cartier at g_Z then mD is Cartier at g_Z for some $m > 0$, hence D is \mathbb{Q} -Cartier along Z^0 by the previous results. \square

Definition 139 (Relative factorial locus). Let X be a normal variety over a perfect field and $Z \subset X$ a closed subset. Let $\{w_i \in X : i \in I\}$ be those non-Cartier centers of X whose closure does not contain any irreducible component of Z . Then

$$\mathrm{Fact}(Z \subset X) := Z \setminus \cup_i \bar{w}_i \quad (139.1)$$

is the unique largest dense open subset $Z^\circ \subset Z$ such that if a Weil divisor D is Cartier at some point of each irreducible component of Z then it is Cartier everywhere along Z° . We call $\mathrm{Fact}(Z \subset X)$ the *relative factorial locus* of X along Z . The factorial locus of X along X is the largest open subscheme $X^\circ \subset X$ whose local rings are factorial.

15. COMMUTATIVE ALGEBRAIC GROUPS

140 (Structure of commutative algebraic groups). Let A denote a commutative algebraic group over a perfect field k , $A^\circ \subset A$ the identity component and $A^{\mathrm{lin}} \subset A$ the largest connected linear algebraic subgroup. Then A/A° is a finite abelian group and A°/A^{lin} is an Abelian variety.

Let $A^{\mathrm{unip}} \subset A^{\mathrm{lin}}$ be the largest unipotent subgroup and $A^{\mathrm{tor}} \subset A^{\mathrm{lin}}$ the largest semisimple subgroup. Then A^{tor} is a *torus* (that is, isomorphic to \mathbb{G}_m^r over k^{sep} for some r) and $A^{\mathrm{lin}} = A^{\mathrm{unip}} + A^{\mathrm{tor}}$.

A is called *semi-Abelian* if $A^{\mathrm{unip}} = 0$.

Let $A^{\mathrm{prop}} \subset A$ denote the largest proper, connected subgroup. Then $A^{\mathrm{prop}} \cap A^{\mathrm{lin}}$ is finite but usually $A^{\mathrm{prop}} + A^{\mathrm{lin}}$ does not equal A .

See [Bor91], [Mil17, Chap.8] or [Bri17b] for details and proofs.

141 (\mathbb{Q} -rank). For Abelian varieties, the \mathbb{Q} -rank of $A(k)$ is a subtle invariant of A and k ; see for example (151) and (152). By contrast the \mathbb{Q} -rank of a linear algebraic group is easy to compute.

(141.1) $G(k)$ is torsion for every algebraic group G over a locally finite field k .

(141.2) $U(k)$ is p^∞ -torsion for every unipotent algebraic group U over a field k of characteristic $p > 0$.

(141.3) $\text{rank}_{\mathbb{Q}} U(k) = \dim U \cdot \deg(k/\mathbb{Q})$ for every unipotent algebraic group U over a field k of characteristic 0.

(141.4) (Larsen) $\text{rank}_{\mathbb{Q}} T(k) = \infty$ for every positive dimensional torus over a field that is not locally finite.

Of these only the last claim is nontrivial. More precise versions follow from weak approximation, but here is a shorter argument.

T is defined over a finitely generated subfield, so we may as well assume that k is either a number field or the field of functions of a geometrically integral curve C over a subfield $k_0 \subset k$. Over a dense, open, regular subset $U \subset C$ we have a torus $T_U \rightarrow U$.

Assume to the contrary that $t_1, \dots, t_s \in T(k)$ generate a maximal rank subgroup. We can view the t_i as rational sections of $T_U \rightarrow U$. After further shrinking U we may assume that they are all regular sections. If $t \in T(k)$ then $t^n \in \langle t_1, \dots, t_s \rangle$ for some $n > 0$. Thus t is a rational section that is also finite over U , hence a regular section. Thus every rational section of $T_U \rightarrow U$ is regular.

Next we show that this is not the case. T is isomorphic to \mathbb{G}_m^r over k^{sep} , hence there is a finite, separable field extension K/k such that $T_K \cong \mathbb{G}_m^r$. (Such a K is called a *splitting field* of T .) After further shrinking U we may assume that we have a finite morphism $\pi : V \rightarrow U$ such that $T_V \cong V \times \mathbb{G}_m^r$. Let now $p \in U$ be a point with preimages $p_1, \dots, p_r \in V$ such that $k(p_i) \cong k(p)$ for every i and π is étale over p . In the geometric cases this is possible by (103.1), for number fields this follows from the Chebotarev density theorem.

Then T_V has a rational section s_V that has a pole along p_1 but regular at p_2, \dots, p_r . Then $\text{norm}_{K/k}(s_V)$ is a rational section of T_U with a pole at p , a contradiction. \square

142 (Jacobians of curves). Let C be a proper scheme of dimension 1 over a field k . Then $\mathbf{Pic}^\circ(C)$ is called the *Jacobian* or *generalized Jacobian* of C and denoted by $\mathbf{Jac}(C)$.

Let $C^{\text{wn}} \rightarrow C$ denote the *weak normalization* and $\bar{C} \rightarrow C^{\text{wn}} \rightarrow C$ the *normalization*. The pull-back maps give

$$\mathbf{Jac}(C) \rightarrow \mathbf{Jac}(C^{\text{wn}}) \rightarrow \mathbf{Jac}(\bar{C}).$$

The kernel of $\mathbf{Jac}(C) \rightarrow \mathbf{Jac}(C^{\text{wn}})$ is $\mathbf{Jac}(C)^{\text{unip}}$ and the kernel of $\mathbf{Jac}(C) \rightarrow \mathbf{Jac}(\bar{C})$ is $\mathbf{Jac}(C)^{\text{lin}}$. Thus (141) gives the following.

(142.1) If k is locally finite then $\mathbf{Jac}(C)(k)$ is torsion.

(142.2) If $\text{char } k > 0$ and C is geometrically integral, then $\mathbf{Jac}(C)(\bar{k})$ is torsion iff C is rational and $C^{\text{wn}} = \bar{C}$.

(142.3) If $\text{char } k = 0$ then $\mathbf{Jac}(C)(\bar{k})$ is torsion iff $h^1(C, \mathcal{O}_C) = 0$. This implies that every irreducible component of $C_{\bar{k}}$ is smooth and rational.

Definition 143. Let P be a 0-cycle on A and $P_{\bar{k}} = \cup_i m_i [p_i]$. Set

$$\mathrm{tr}_A P := \sum_i m_i [p_i] \quad (\text{summation in } A.) \quad (143.1)$$

Note that $\mathrm{tr}_A P \in A(k^{\mathrm{ins}})$. If A is the additive group \mathbb{G}_a then this is the usual trace, but for the multiplicative group \mathbb{G}_m this is the norm. Since we usually use additive notation, trace seems a better choice. For $Z \subset A$ set

$$\mathrm{tr}_A Z := \{\mathrm{tr}_A P : P \text{ is a 0-cycle on } Z\}. \quad (143.2)$$

Let C be a smooth, projective curve. There is a natural embedding $j : C \hookrightarrow \mathbf{Jac}_1(C) \subset \mathbf{Pic}(C)$. If P is a 0-cycle on C then

$$\mathrm{tr}_{\mathbf{Pic}(C)}(j(P)) = [\mathcal{O}_C(P)] \in \mathrm{Jac}_{\mathrm{deg} P}(C). \quad (143.3)$$

The following is a restatement of [Bri17a, 4.9].

Lemma 144. $A \mapsto A(k) \otimes \mathbb{Q}$ is an exact functor on the category of commutative algebraic groups.

Proof. The only nontrivial claim is that if $g : A \rightarrow B$ is a dominant morphism then $g(k) : A(k) \rightarrow B(k)$ is surjective modulo torsion. To see this pick $b \in B(k)$ and let P be a 0-cycle on the fiber A_b . Then $\mathrm{tr}_A P \in A(k)$ and $g(\mathrm{tr}_A P) = \mathrm{deg} P \cdot b$. \square

The multiplicative group of Artin algebras

Definition 145. Let A be an Artin algebra. The multiplicative group of its invertible elements is denoted by A^\times .

If A is a k -algebra, it is frequently convenient to view A^\times as an algebraic group over k ; this is called *Weil restriction*. We denote it by $\mathcal{R}_k^A \mathbb{G}_m$. That is,

$$(\mathcal{R}_k^A \mathbb{G}_m)(B) = (A \otimes_k B)^\times \quad \text{for a } k\text{-algebra } B.$$

Note that $\dim \mathcal{R}_k^A \mathbb{G}_m = \dim_k A$.

For example, if K/k is a field extension of degree n , choose a basis $e_i \in K$. As a variety, $\mathcal{R}_k^K \mathbb{G}_m$ is $\mathbb{A}^n \setminus (\mathrm{norm}_{K/k}(\sum x_i e_i) = 0)$.

146. Let (A, m) be a semi-local, Artin k algebra and $K = A/m$. There is an exact sequence

$$1 \rightarrow U \rightarrow A^\times \rightarrow K^\times \rightarrow 1, \quad (146.1)$$

where the map $a \mapsto 1 + a$ identifies m with U . Note that $a \mapsto 1 + a$ is a group isomorphism if $m^2 = 0$ but not otherwise. In characteristic 0 one can correct this by taking $a \mapsto \exp(a)$. We will think of (146.1) as the k -points of an exact sequence of algebraic k -groups

$$1 \rightarrow U \rightarrow \mathcal{R}_k^A \mathbb{G}_m \rightarrow \mathcal{R}_k^K \mathbb{G}_m \rightarrow 1, \quad (146.2)$$

where U is a unipotent group. In positive characteristic the algebraic groups $(m, +)$ and (U, \cdot) need not be isomorphic.

We also use the following variant of Hensel's lemma.

Claim 146.3. Let $k \subset k' \subset K$ be a subfield that is separable over k . Then there is a unique lifting $k' \rightarrow A$. \square

Combining the above with the previous discussion on algebraic groups yield the following lemmas.

Lemma 147. Let k be a field and $A \rightarrow B$ a homomorphism of Artin k -algebras. Then $\mathrm{coker}[A^\times \rightarrow B^\times]$ is torsion iff one of the following holds.

(147.1) $A \rightarrow B$ is surjective.

(147.2) $\text{char } k > 0$ and $B/\sqrt{0}$ is purely inseparable over $A/\sqrt{0}$.

(147.3) k is locally finite.

Moreover, $\text{coker}[A^\times \rightarrow B^\times]$ has finite \mathbb{Q} -rank in one additional case:

(147.4) $\deg(k/\mathbb{Q}) < \infty$ and $A/\sqrt{0} \rightarrow B/\sqrt{0}$ is surjective. \square

Lemma 148. *Let k be a field and $A \rightarrow B$ a homomorphism of Artin k -algebras. Then $\ker[A^\times \rightarrow B^\times]$ is torsion iff one of the following holds.*

(148.1) $A \rightarrow B$ is injective.

(148.2) $\text{char } k > 0$ and $A/\sqrt{0} \rightarrow B/\sqrt{0}$ is injective.

(148.3) k is locally finite.

Moreover, $\ker[A^\times \rightarrow B^\times]$ has finite \mathbb{Q} -rank in one additional case:

(148.4) $\deg(k/\mathbb{Q}) < \infty$ and $A/\sqrt{0} \rightarrow B/\sqrt{0}$ is injective. \square

Lemma 149. *Let k be a field that is not locally finite. Let $A \supset k$ be a finite, reduced k -algebra. Let $k \subset L_1, L_2 \subset A$ be subfields. The following are equivalent.*

(149.1) $A^\times/(L_1^\times \cdot L_2^\times)$ is torsion.

(149.2) $A^\times/(L_1^\times \cdot L_2^\times)$ has finite \mathbb{Q} -rank.

(149.3) A is a field and A/L_i is purely inseparable for some $i = 1, 2$.

Proof. If A/L_i is purely inseparable then $A^q \subset L_i$ for some power q of $\text{char } k$, hence A^\times/L_i^\times is torsion. This proves (149.3) \Rightarrow (149.1) and (149.1) \Rightarrow (149.2) is clear.

Assume (149.2). We may replace A by its maximal separable subalgebra. Thus assume that A/k is separable. If $A = L_i$ for some i then we are done. Otherwise $\dim_k A = \dim_{L_i} A \cdot \dim_k L_i \geq 2 \dim_k L_i$.

If B is a reduced, separable k -algebra then B^\times is identified with the k -points of the k -torus $\mathcal{R}_k^B \mathbb{G}_m$. Thus $L_1^\times \cdot L_2^\times \rightarrow A^\times$ can be viewed as the k -points of a morphism of k -tori

$$\mu : \mathcal{R}_k^{L_1} \mathbb{G}_m \times \mathcal{R}_k^{L_2} \mathbb{G}_m \longrightarrow \mathcal{R}_k^A \mathbb{G}_m.$$

Both of the L_i contain k , thus

$$\dim_k \text{im}(\mu) \leq \dim_k L_1 + \dim_k L_2 - 1 < \dim A.$$

Thus $\text{coker}(\mu)$ is a positive dimensional k -torus, hence $\text{rank}_{\mathbb{Q}}(\text{coker}(\mu)(k)) = \infty$ by (141.4). Finally (144) shows that

$$\text{rank}_{\mathbb{Q}}(A^\times/(L_1^\times \cdot L_2^\times)) = \text{rank}_{\mathbb{Q}}(\text{coker}(\mu)(k)) = \infty. \quad \square$$

16. SPECIAL FIELDS

We discuss various classes of fields that were used earlier.

150 (Locally finite fields). A field k is called *locally finite*² if the following equivalent conditions hold.

(150.1) Every finitely generated subfield of k is finite.

(150.2) k is an algebraic extension of a finite field.

(150.3) k is isomorphic to a subfield of \mathbb{F}_p for some $p > 0$.

(150.4) $A(k)$ is a torsion group for every Abelian variety A over k .

²The terminology is not standard in English; it is an analog of the notion of locally finite group.

(150.5) There is a $C > 0$ such that $\text{rank}_{\mathbb{Q}} A(k) \leq C$ for every Abelian variety A over k .

The only non-obvious claims are (150.4–5). If k is not locally finite then it contains either \mathbb{Q} or $\mathbb{F}_p(t)$. In both cases, there is an Abelian variety A over k with arbitrarily large $\text{rank}_{\mathbb{Q}} A(k)$. For example, if E is an elliptic curve of rank ≥ 1 then E^m has rank $\geq m$.

151 (\mathbb{Q} -Mordell-Weil fields). A field k is *Mordell-Weil* (resp. \mathbb{Q} -*Mordell-Weil*) if for every Abelian variety A over k , the group of its k -points $A(k)$ is finitely generated (resp. has finite \mathbb{Q} -rank).

By [LN59], every finitely generated field is Mordell-Weil.

Weil restriction (cf. [BLR90, Sec.7.6]) shows that these properties are invariant under finite field extensions. Since every Abelian variety is a quotient of a Jacobian, it is equivalent to ask that $\text{Jac}(C)$ have these properties for every smooth projective curve C over k (144).

It is not clear how much the class of \mathbb{Q} -Mordell-Weil fields differs from the class of Mordell-Weil fields. For example, $\overline{\mathbb{F}}_p$ is \mathbb{Q} -Mordell-Weil but not Mordell-Weil.

152 (Anti-Mordell-Weil fields). Following [IL19] a field k is called *anti-Mordell-Weil* if

(152.1) the \mathbb{Q} -rank of $A(k)$ is infinite for every positive dimensional abelian variety.

In particular, k is not locally finite. If the latter holds then the \mathbb{Q} -rank of $T(k)$ is infinite for every k -torus T (141.4), hence (152.1) can be restated as:

(152.2) The \mathbb{Q} -rank of $A(k)$ is infinite for every positive dimensional semi-abelian variety A .

Finally, if $\text{char } k = 0$, then k is not a finite extension of \mathbb{Q} by the Mordell-Weil theorem, hence the \mathbb{Q} -rank of $U(k)$ is infinite for every unipotent group (141.3).

Thus (152.1–2) are further equivalent to:

(152.3) The \mathbb{Q} -rank of $A(k)$ is infinite for every positive dimensional commutative algebraic group A .

Warning. Note that if $\text{char } k = p > 0$ then $U(k)$ is p -power torsion for every unipotent group; this creates a crucial difference between 0 and positive characteristics for us.

Examples of anti-Mordell-Weil fields are the following.

(152.4) algebraically closed fields, save for $\overline{\mathbb{F}}_p$,

(152.5) \mathbb{R} and all real closed fields,

(152.6) \mathbb{Q}_p , more generally quotient fields of Henselian, local domains,

(152.7) large fields [Kob06, FP10, FP19] that are not locally finite, where a field k is *large* (also called ample, fertile or anti-mordell) if $C(k)$ is either empty or infinite for every smooth curve C .

The last case implies the earlier ones.

153 (Hilbertian fields). A field k is *Hilbertian* if for every irreducible polynomial $f(x, y) \in k[x, y]$ such that $\partial f / \partial y \neq 0$, there are infinitely many $c \in k$ such that $f(x, c) \in k[x]$ is irreducible. (We follow [FJ08, Chap.12] with adding the separability condition.)

Equivalently, for every smooth, irreducible curve C and every basepoint-free linear system $|M|$ that defines a separable map $C \rightarrow \mathbb{P}^1$, there are infinitely many

irreducible members $M_c \in |M|$. This also implies that, for every irreducible variety X over k , and every mobile linear system $|M|$ that defines a separable map $X \dashrightarrow \mathbb{P}^N$, there is a dense set $\Lambda \subset |M|(k)$ such that $M_\lambda \in |M|$ is irreducible for $\lambda \in \Lambda$.

Hilbert proved that number fields are Hilbertian. More generally, every finitely generated, infinite field is Hilbertian. A finite, separable extension of a Hilbertian field is Hilbertian, and so is any purely inseparable extension. See [Lan62, Chap.VIII] or [FJ08, Chaps.12–13] for these and many other facts.

In our proofs the Hilbertian condition is mostly used through the following consequence.

Lemma 154. *Let C be an irreducible, geometrically reduced, projective curve over a Hilbertian field k . Let $\Sigma \subset C$ be a finite subset and $Z \subset C$ a finite subscheme. Let L be a line bundle on C such that $\deg L \geq \deg Z + \deg \omega_C + 3$. Then every $s_Z \in H^0(Z, L|_Z)^\times$ can be lifted to $s_C \in H^0(C, L)$ such that $(s_C = 0)$ is irreducible, reduced and disjoint from Σ .*

Proof. The condition $s|_Z = c \cdot s_Z$ for some $c \in k$ determines a linear subsystem $|L, s_Z| \subset |L|$. The degree condition guarantees that it is basepoint-free and separable. Hence it has infinitely many irreducible members. Almost all of them are disjoint from Σ . \square

It turns out that versions of (154) hold for some non-Hilbertian fields and in our proofs a weakening of it is sufficient. In order to state it, we need a definition.

Definition 155. Let C be an irreducible, geometrically reduced, projective curve over a field k . Let $\Sigma \subset C$ be a finite subset, $Z \subset C$ a finite subscheme and L an ample line bundle on C . Let

$$\Gamma^{\text{irr}}(C, L, Z, \Sigma^c) \subset H^0(Z, \bigoplus_{m>0} L^m|_Z)^\times \quad (155.1)$$

be the subset consisting of the restrictions $s|_Z$ of those sections $s \in H^0(C, L^m)$, for which $(s = 0)$ is irreducible and disjoint from $\Sigma \cup Z$.

If k is Hilbertian then, by (154),

$$\Gamma^{\text{irr}}(C, L, Z, \Sigma^c) \supset H^0(Z, \bigoplus_{m \geq m_0} L^m|_Z)^\times \quad \text{for some } m_0. \quad (155.2)$$

Definition 156 (Weakly Hilbertian fields). Let C be a projective curve over a field k and L an ample line bundle on C . We say that (C, L) is *weakly Hilbertian* if the following holds.

(156.1) For every $s_Z \in H^0(Z, L^m|_Z)^\times$, some power of s_Z is in $\Gamma^{\text{irr}}(C, L, Z, \Sigma^c)$.

We call a field k *weakly Hilbertian* if (C, L) is weakly Hilbertian for every irreducible, geometrically reduced, projective curve over k and every ample line bundle L on C . We see in (157) that it is enough to check this for smooth curves.

By (154) every Hilbertian field is also weakly Hilbertian. Below we show that all Galois extensions $\mathbb{Q} \subset K \subsetneq \bar{\mathbb{Q}}$ are weakly Hilbertian. More generally, Galois extensions of Hilbertian fields, save those that are separably closed, are weakly Hilbertian; see (159). The fields \mathbb{Q}_p and $\mathbb{F}_p((t))$ are not weakly Hilbertian, but all smooth curves over them satisfy property (156.2) below; see (165). See (167) for finite fields.

Comment. We do not claim that the above is the ‘right’ notion of weakly Hilbertian fields. It works well for our purposes. However, from the field-theory point of view, the following even weaker version may seem more natural.

(156.2) Let C be an irreducible, geometrically reduced, projective curve over k and L an ample line bundle on C . Then there is an $m > 0$ and infinitely many $s_i \in H^0(C, L^{m_i})$ whose zero sets ($s_i = 0$) are irreducible and disjoint.

It is clear that (156.1) \Rightarrow (156.2); we do not know whether the converse holds.

Subfields of weakly Hilbertian fields

Let K/k be a finite, separable field extension. If k is Hilbertian, then so is K , but the converse fails. We prove in (158) that if K is weakly Hilbertian, then so is k . We do not know whether the converse holds.

Lemma 157. *Let C be an irreducible, geometrically reduced, projective curve and L an ample line bundle on C . Let $\pi : \bar{C} \rightarrow C$ denote the normalization. If (\bar{C}, \bar{L}) is weakly Hilbertian, then so is (C, L) .*

Proof. We are free to enlarge Z , hence we may assume that it contains the conductor subscheme of π and its ideal sheaf is also an ideal sheaf on \bar{C} , defining \bar{Z} . We have a natural inclusion $\tau : \mathcal{O}_Z \hookrightarrow \pi_* \mathcal{O}_{\bar{Z}}$. Thus if \bar{s} is a section of \bar{L}^m whose restriction to \bar{Z} agrees with $\tau(s_Z^m)$, then it descends to a section s of L^m . \square

Lemma 158. *Let K/k be a separable, algebraic field extension. If K is weakly Hilbertian then so is k .*

Proof. Start with C_k, L_k, Z_k, Σ_k and $s_{Z,k} \in H^0(Z_k, L_k|_{Z_k})^\times$. By base change we get C_K, L_K, Z_K and $s_{Z,K} \in H^0(Z_K, L_K|_{Z_K})^\times$. Let $C'_K \subset C_K$ be one of its irreducible components.

Assume that we have $s'_K \in H^0(C'_K, L'_K)$ such that $s'_K|_{Z'_K} = s_{Z,K}^m$. This s'_K is defined over a finite degree subextension $k \subset K_1 \subset K$.

Next assume first that C_k is smooth. Then C'_K is smooth and $C'_K \rightarrow C_k$ is flat. Thus $\text{norm}_{K_1/k}$ sends sections of $(L'_K)^m$ to sections of L_k^{md} where $d = \deg(K_1/k)$. Set $s_k := \text{norm}_{K_1/k}(s'_K)$. Then $s_k \in H^0(C_k, L_k^{md})$ and $s_k|_{Z_k} = s_{Z,k}^{md}$.

The singular case follows from (157). \square

Corollary 159. *Let k be a Hilbertian field and K/k a Galois extension that is not separably closed. Then K is weakly Hilbertian.*

Proof. By [Wei82], every nontrivial finite extension of such a field K is Hilbertian. Thus, if K is not separably closed, then K is weakly Hilbertian by (158). \square

As an example, let \mathbb{Q}^{solv} denote the composite of all Galois extensions of \mathbb{Q} with solvable Galois group. It is weakly Hilbertian by (159), but not Hilbertian, as shown by the polynomial $x^2 - y$.

Corollary 160. *Let $K \subset \bar{\mathbb{Q}}$ be a subfield whose Galois closure is different from $\bar{\mathbb{Q}}$. Then K is weakly Hilbertian.* \square

Not weakly Hilbertian fields

Next we show that $\bar{\mathbb{Q}}$ is not weakly Hilbertian, it does not even satisfy (156.2) for smooth curves. Note that a much stronger variant of (161) could be true; see (16) and (164). The same methods work for $\mathbb{R} \cap \bar{\mathbb{Q}}$. We do not know any other subfield of $\bar{\mathbb{Q}}$ that does not satisfy (156.2) for smooth curves, though presumably there are many.

Proposition 161. *There is a smooth projective curve C and an ample line bundle L , defined over $\bar{\mathbb{Q}}$, such that every section of L^m has at least 2 distinct zeros for every $m > 0$.*

Proof. Let $\pi : C \rightarrow B$ be a nonconstant morphism between smooth, projective curves such that $g(C) \geq g(B) + 2$ and $g(B) \geq 1$. Let $\Gamma \subset \text{Pic}(B)$ be as in (162).

The \mathbb{Q} -rank of $\text{Pic}(B)$ is infinite by (152.4), so there is an ample $L \in \text{Pic}(B)$ no power of which is in Γ . Then π^*L has the required property. \square

Lemma 162. *Let $\pi : C \rightarrow B$ be a nonconstant morphism between smooth, projective curves such that $g(C) \geq g(B) + 2$. Then*

$$\Gamma := \langle L \in \text{Pic}(B) : \pi^*L^m \sim \mathcal{O}_C(n[c]) \text{ for some } c \in C(k), n, m > 0 \rangle \subset \text{Pic}(B)$$

has finite \mathbb{Q} -rank.

Proof. This is clear if $C(k)$ is empty. Otherwise fix a point $c_0 \in C(k)$ and embed $C \hookrightarrow \mathbf{Jac}(C)$ sending c_0 to the origin. Set $A := \mathbf{Jac}(C)/\pi^*\mathbf{Jac}(B)$ with quotient map $\sigma : C \rightarrow A$. If $\pi^*L^m \sim \mathcal{O}_C(n[c])$ then $\sigma(c) \in A$ is a torsion point. Thus there are only finitely many such $c \in C(k)$ by [Zha98], and the \mathbb{Q} -rank of Γ is at most the number of such torsion points. \square

For nodal rational curves, there is an elementary proof.

Proposition 163. *Nodal rational curves over $\bar{\mathbb{Q}}$ are not weakly Hilbertian.*

Proof. A rational curve with r nodes is obtained from \mathbb{P}^1 by identifying r point pairs. Thus we start with $2r$ distinct points $a_1, \dots, a_{2r} \in \mathbb{A}^1$ and identify a_i with a_{r+i} to get a nodal rational curve C .

A line bundle on C is obtained by starting with some $L = \mathcal{O}_{\mathbb{P}^1}(m)$ and specifying isomorphisms $L|_{a_i} \cong L|_{a_{r+i}}$. Thus sections of the resulting line bundle are given by polynomials $p(x)$ of degree $\leq m$ such that $p(a_i) = u_i p(a_{r+i})$ for every i where $u_i \in k^\times$ specify the line bundle.

A polynomial with zeros $\{z_j : j \in J\}$ is $s(x) = \gamma \prod_j (x - z_j)^{m_j}$. Thus for nonzero u_1, \dots, u_r we aim to solve the r equations

$$\prod_{j \in J} \left(\frac{a_i - z_j}{a_{r+i} - z_j} \right)^{m_j} = u_i^n, \quad (163.1)$$

where $m_j, n \in \mathbb{Z}$ and $z_j \in \bar{\mathbb{Q}}$ are unknowns with $n \neq 0$.

For every p choose an extension v_p of the p -adic valuation to $\bar{\mathbb{Q}}$. The v_p -valuation of any $d \in \bar{\mathbb{Q}}^\times$ is 0 for all but finitely many p . We can thus choose p such that

$$v_p(a_i) = v_p(a_i - a_j) = 0 \quad \text{for every } i \neq j. \quad (163.2)$$

Thus taking the valuation of (163.1) we get the equations

$$\sum_j m_j v_p \left(\frac{a_i - z_j}{a_{r+i} - z_j} \right) = n v_p(u_i). \quad (163.3)$$

Choose the u_i such that $v_p(u_i) \neq 0$ for every i . By (163.3), for every i we get a $\sigma(i) \in J$ such that

$$v(a_i - z_{\sigma(i)}) > 0 \quad \text{or} \quad v(a_{r+i} - z_{\sigma(i)}) > 0. \quad (163.4)$$

If $r > |J|$ then the same z_j appears twice. Thus we have $v(a_{i_1} - z_j) > 0$ and $v(a_{i_2} - z_j) > 0$ for some $i_1 \neq i_2$ and j . Then

$$v(a_{i_1} - a_{i_2}) \geq \min\{v(a_{i_1} - z_j), v(a_{i_2} - z_j)\} > 0$$

gives a contradiction. \square

Claim 163.5. Let (R, v) be a valuation ring and $a, c \in R^\times$ such that $v(a) = v(c) = v(a - c) = 0$. Then

$$v\left(\frac{a-z}{c-z}\right) > 0 \Leftrightarrow v(a-z) > 0 \quad \text{and} \quad v\left(\frac{a-z}{c-z}\right) < 0 \Leftrightarrow v(c-z) > 0. \quad \square$$

The above proof shows the following stronger claim.

Corollary 164. *Let C be a rational curve with g nodes over $\bar{\mathbb{Q}}$. Then for ‘most’ ample line bundles L over C , every section of L^m has at least g distinct zeros. \square*

Quotient fields of discrete valuation rings

\mathbb{Q}_p , $\mathbb{F}_p((t))$, and, more generally, quotient fields of excellent DVRs, have very interesting behavior. Smooth curves have weak Hilbertian properties but singular curves do not.

Note that a DVR of characteristic 0 is excellent [Sta15, 07QW]. In positive characteristic, local rings of smooth curves are excellent and so are power series rings $K[[t]]$. However, there are many non-excellent DVRs; see [DS18] for especially simple examples. (The proof below uses excellence, but the result might hold for any DVR.)

Proposition 165. *Let (R, m) be an excellent DVR with quotient field K and locally finite residue field k . Let C_K be a smooth projective, irreducible curve over K and L_K an ample line bundle on C_K . Then $|L_K^n|$ has infinitely many irreducible members for n sufficiently divisible.*

Proof. We extend C_K to a flat morphism $C_R \rightarrow \text{Spec } R$. We may assume that C_R is regular by [Sha66]. Then L_K extends to a line bundle L_R on C_R .

Let E_1, \dots, E_r be the irreducible components of the central fiber $C_k = \sum m_i E_i$. The intersection matrix $(E_i \cdot E_j)$ is negative semidefinite with $[C_k]$ as the only null-vector. Thus the intersection matrix of E_2, \dots, E_r is negative definite. We can thus find a divisor F supported on E_2, \dots, E_r such that $L_R^{n_1}(F)$ has degree 0 on E_2, \dots, E_r .

By [Art62] the curves E_2, \dots, E_r can be contracted $C_R \rightarrow C_R^*$ and a suitable power of $L_R^{n_1}(F)$ descends to a line bundle L^* on C_R^* .

Now we have a normal scheme with a flat morphism $\pi : C_R^* \rightarrow R$ whose generic fiber is C_K and whose central fiber C_k^* is an irreducible curve. Furthermore there is a line bundle L^* whose restriction to C_K is a power of L_K .

Set $E^* = \text{red } C_k^*$ and pick any point $p \in E^*$ that is regular both on E^* and on C_R^* . Since $\text{rank}_{\mathbb{Q}} \text{Pic}(E^*) = 1$, after passing to a power of L^* , we may assume that

$$(165.1) \quad L^*|_{E^*} \text{ has a section } \bar{s}_0 \text{ that vanishes only at } p,$$

$$(165.2) \quad L^*(-E^*)/L^*(-2E^*) \text{ has a section } \bar{s}_1 \text{ that does not vanish at } p, \text{ and}$$

$$(165.3) \quad H^1(C_R^*, L^*(-E^*)) = H^1(C_R^*, L^*(-2E^*)) = 0.$$

By (165.3) we can lift \bar{s}_0 and \bar{s}_1 to $s_0 \in H^0(C_R^*, L^*)$ and $s_1 \in H^0(C_R^*, L^*(-E^*))$. For all but 1 residue value of $\lambda \in R$, $D_R(\lambda) := (s_0 + \lambda s_1 = 0)$ is regular at p . Since p is its sole point over k , $D_R(\lambda)$ is irreducible and reduced. \square

Proposition 166. *Let (R, m) be a Henselian DVR with quotient field K and residue field k . For $2g \leq |k|$ there are rational curves C with g nodes over K such that, for ‘most’ ample line bundles L over C , every section of L^n has at least g distinct zeros. \square*

Proof. We choose $a_i \in R \subset \mathbb{A}^1(K) \subset \mathbb{P}_K^1$ such that $\bar{a}_i \in k$ (their reduction mod m) are all distinct. As in (163), identifying a_i with a_{g+i} to get a nodal rational curve C and a line bundle L on C is obtained by starting with some $\mathcal{O}_{\mathbb{P}^1}(r)$ and specifying isomorphisms $\mathcal{O}_{\mathbb{P}^1}(r)|_{a_i} \cong \mathcal{O}_{\mathbb{P}^1}(r)|_{a_{g+i}}$. Thus sections of L^n are given by polynomials $f(x)$ of degree $\leq nr$ such that

$$f(a_i) = u_i^n f(a_{g+i}) \quad \text{for } i = 1, \dots, g, \quad (163.1)$$

where the $u_i \in K^\times$ determine L . We may assume that $f(x) \in R[x] \setminus m[x]$ and denote by \bar{f} its image in $k[x]$. If the m -adic valuation of u_i is not 0 for every i , this implies that

$$\bar{f}(\bar{a}_i) = 0 \quad \text{or} \quad \bar{f}(\bar{a}_{g+i}) = 0 \quad \text{for } i = 1, \dots, g. \quad (163.1)$$

Thus \bar{f} has at least g distinct zeros, so, by Hensel’s lemma, f has at least g distinct prime factors. \square

Note that, in the above case, the precise meaning of ‘most’ line bundles is the following. Choosing the point at infinity fixes an isomorphism $\text{Pic}(C) \cong \text{Pic}^\circ(C) \times \mathbb{Z}$.

For a node p_i the map $\rho_i : \text{Pic}^\circ(C) \rightarrow \mathbb{G}_m$ depends on the ordering of the preimages of the node. So we get 2 maps, that differ only by composing with the inverse map of \mathbb{G}_m . Composing ρ_i with the absolute value of the m -adic valuation gives a well-defined map $v_i : \text{Pic}^\circ(C) \rightarrow \mathbb{N}$. The g nodes together give $v : \text{Pic}(C) \rightarrow \mathbb{N}^g$. If all coordinates of $v(L)$ are nonzero, then every section of L^n has at least g distinct zeros for every n .

167. B. Poonen explained to us that, using geometric class field theory and the function field Chebotarev density theorem, one can prove that if C is a smooth projective curve over a finite field k and L is an ample line bundle, then L^m has a section with irreducible and reduced zero set for all $m \gg 1$. However, the probability that a random section has this property tends to 0 as $m \rightarrow \infty$.

REFERENCES

- [Amb18] Emiliano Ambrosi, *Specialization of Néron-Severi groups in positive characteristic*, arXiv e-prints (2018), arXiv:1810.06481.
- [And96] Yves André, *Pour une théorie inconditionnelle des motifs*, Inst. Hautes Etudes Sci. Publ. Math. **83** (1996), 5–49.
- [Art62] Michael Artin, *Some numerical criteria for contractability of curves on algebraic surfaces*, Amer. J. Math. **84** (1962), 485–496. MR MR0146182 (26 #3704)
- [BGS11] S. Boissière, O. Gabber, and O. Serman, *Sur le produit de variétés localement factorielles ou Q -factorielles*, ArXiv e-prints (2011).
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 21, Springer-Verlag, Berlin, 1990. MR 1045822 (91i:14034)
- [Bor91] Armand Borel, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer-Verlag, New York, 1991. MR 1102012 (92d:20001)
- [BPS16] Fedor A. Bogomolov, Alena Pirutka, and Aaron Michael Silberstein, *Families of disjoint divisors on varieties*, Eur. J. Math. **2** (2016), no. 4, 917–928. MR 3572551
- [Bri17a] Michel Brion, *Commutative algebraic groups up to isogeny*, Doc. Math. **22** (2017), 679–725. MR 3650225

- [Bri17b] ———, *Some structure theorems for algebraic groups*, Algebraic groups: structure and actions, Proc. Sympos. Pure Math., vol. 94, Amer. Math. Soc., Providence, RI, 2017, pp. 53–126. MR 3645068
- [BVRs09] Luca Barbieri-Viale, Andreas Rosenschon, and V. Srinivas, *The Néron-Severi group of a proper seminormal complex variety*, Math. Z. **261** (2009), no. 2, 261–276. MR 2457299
- [BVS93] L. Barbieri Viale and V. Srinivas, *On the Néron-Severi group of a singular variety*, J. Reine Angew. Math. **435** (1993), 65–82. MR 1203911
- [Chr18] Atticus Christensen, *Specialization of Néron-Severi groups in characteristic p* , arXiv e-prints (2018), arXiv:1810.06550.
- [CP16] François Charles and Bjorn Poonen, *Bertini irreducibility theorems over finite fields*, J. Amer. Math. Soc. **29** (2016), no. 1, 81–94. MR 3402695
- [DS18] Rankeya Datta and Karen E. Smith, *Excellence in prime characteristic*, Local and global methods in algebraic geometry, Contemp. Math., vol. 712, Amer. Math. Soc., Providence, RI, 2018, pp. 105–116. MR 3832401
- [FJ08] Michael D. Fried and Moshe Jarden, *Field arithmetic*, third ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 11, Springer-Verlag, Berlin, 2008, Revised by Jarden. MR 2445111
- [FP10] Arno Fehm and Sebastian Petersen, *On the rank of abelian varieties over ample fields*, Int. J. Number Theory **6** (2010), no. 3, 579–586. MR 2652895
- [FP19] ———, *Ranks of abelian varieties and the full Mordell-Lang conjecture in dimension one*, arXiv e-prints (2019), arXiv:1912.10710.
- [Ful98] William Fulton, *Intersection theory*, second ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 2, Springer-Verlag, Berlin, 1998. MR MR1644323 (99d:14003)
- [Gro60] Alexander Grothendieck, *Éléments de géométrie algébrique. I–IV.*, Inst. Hautes Études Sci. Publ. Math. (1960), no. 4,8,11,17,20,24,28,32.
- [Gro62] ———, *Fondements de la géométrie algébrique. [Extraits du Séminaire Bourbaki, 1957–1962.]*, Secrétariat mathématique, Paris, 1962. MR MR0146040 (26 #3566)
- [Gro68] ———, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, North-Holland Publishing Co., Amsterdam, 1968, Augmenté d’un exposé par Michèle Raynaud, Séminaire de Géométrie Algébrique du Bois-Marie, 1962, Advanced Studies in Pure Mathematics, Vol. 2. MR 0476737
- [Har62] Robin Hartshorne, *Complete intersections and connectedness*, Amer. J. Math. **84** (1962), 497–508. MR 0142547 (26 #116)
- [Har77] ———, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52. MR 0463157 (57 #3116)
- [IL19] Bo-Hae Im and Michael Larsen, *Abelian Varieties and Finitely Generated Galois Groups*, arXiv e-prints (2019), arXiv:1902.04011.
- [Ji20] Lena Ji, (*in preparation*), 2020.
- [KLOS20] János Kollár, Max Lieblich, Martin Olsson, and Will Sawin, *Revised and expanded version of [LO19]*, 2020.
- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KM09] János Kollár and Frédéric Mangolte, *Cremona transformations and diffeomorphisms of surfaces*, Adv. Math. **222** (2009), no. 1, 44–61. MR 2531367 (2010g:14016)
- [Kob06] Emi Kobayashi, *A remark on the Mordell-Weil rank of elliptic curves over the maximal abelian extension of the rational number field*, Tokyo J. Math. **29** (2006), no. 2, 295–300. MR 2284973
- [Koc18] Fabian Koch, *Jacobians of curves on surfaces*, <http://www.math.uni-bonn.de/people/huybrech/KochMaster.pdf>, 2018.
- [Kol96] János Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 32, Springer-Verlag, Berlin, 1996.
- [Kol18] ———, *Mumford divisors*, ArXiv e-prints (2018).

- [Lan62] Serge Lang, *Diophantine geometry*, Interscience Tracts in Pure and Applied Mathematics, No. 11, Interscience Publishers (a division of John Wiley & Sons), New York-London, 1962. MR 0142550
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I-II*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 48–49, Springer-Verlag, Berlin, 2004. MR 2095471 (2005k:14001a)
- [LN59] S. Lang and A. Néron, *Rational points of abelian varieties over function fields*, Amer. J. Math. **81** (1959), 95–118. MR 0102520
- [LO19] Max Lieblich and Martin Olsson, *A reconstruction theorem for varieties*, 2019.
- [Mat52] Teruhisa Matsusaka, *On the algebraic construction of the Picard variety. II*, Jap. J. Math. **22** (1952), 51–62. MR 0062471
- [Mil17] J. S. Milne, *Algebraic groups*, Cambridge Studies in Advanced Mathematics, vol. 170, Cambridge University Press, Cambridge, 2017, The theory of group schemes of finite type over a field. MR 3729270
- [MP12] Daves Maulik and Bjorn Poonen, *Néron-Severi groups under specialization*, Duke Math. J. **161** (2012), 2167–2206.
- [Mum66] David Mumford, *Lectures on curves on an algebraic surface*, With a section by G. M. Bergman. Annals of Mathematics Studies, No. 59, Princeton University Press, Princeton, N.J., 1966. MR 0209285 (35 #187)
- [Mum70] ———, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, No. 5, Published for the Tata Institute of Fundamental Research, Bombay, 1970. MR 0282985 (44 #219)
- [Nér52a] André Néron, *La théorie de la base pour les diviseurs sur les variétés algébriques*, Deuxième Colloque de Géométrie Algébrique, Liège, 1952, Georges Thone, Liège; Masson & Cie, Paris, 1952, pp. 119–126. MR 0052154
- [Nér52b] ———, *Problèmes arithmétiques et géométriques rattachés à la notion de rang d'une courbe algébrique dans un corps*, Bull. Soc. Math. France **80** (1952), 101–166. MR 56951
- [Per06] Jorge Vitório Pereira, *Fibrations, divisors and transcendental leaves*, J. Algebraic Geom. **15** (2006), no. 1, 87–110, With an appendix by Laurent Meersseman. MR 2177196
- [Pic95] Émile Picard, *Sur la théorie des groupes et des surfaces algébriques*, Rend. Circ. Math. Palermo **9** (1895).
- [Poo01] Bjorn Poonen, *Points having the same residue field as their image under a morphism*, J. Algebra **243** (2001), no. 1, 224–227. MR 1851662
- [Poo08] ———, *Smooth hypersurface sections containing a given subscheme over a finite field*, Math. Res. Lett. **15** (2008), no. 2, 265–271. MR 2385639
- [RS06] G. V. Ravindra and V. Srinivas, *The Grothendieck-Lefschetz theorem for normal projective varieties*, J. Algebraic Geom. **15** (2006), no. 3, 563–590. MR 2219849
- [RS09] ———, *The Noether-Lefschetz theorem for the divisor class group*, J. Algebra **322** (2009), no. 9, 3373–3391. MR 2567426
- [Ser59] Jean-Pierre Serre, *Morphismes universels et variété d'albanese*, Séminaire Claude Chevalley **4** (1958-1959) (fr), talk:10.
- [Ser89] ———, *Lectures on the Mordell-Weil theorem*, Aspects of Mathematics, E15, Friedr. Vieweg & Sohn, Braunschweig, 1989, Translated from the French and edited by Martin Brown from notes by Michel Waldschmidt. MR 1002324
- [Sev06] Francesco Severi, *Sulla totalità delle curve algebriche tracciate sopra una superficie algebrica*, Math. Ann. **62** (1906), no. 2, 194–225. MR 1511373
- [Sha66] Igor R. Shafarevich, *Lectures on minimal models and birational transformations of two dimensional schemes*, Notes by C. P. Ramanujam. Tata Institute of Fundamental Research Lectures on Mathematics and Physics, No. 37, Tata Institute of Fundamental Research, Bombay, 1966. MR 0217068 (36 #163)
- [Sil83] Joseph H. Silverman, *Heights and the specialization map for families of abelian varieties*, J. Reine Angew. Math. **342** (1983), 197–211. MR 703488
- [Sta15] The Stacks Project Authors, *Stacks Project*, <http://stacks.math.columbia.edu>, 2015.
- [Ter85] Tomohide Terasoma, *Complete intersections with middle Picard number 1 defined over \mathbb{Q}* , Math. Z. **189** (1985), 289–296.
- [Tot00] Burt Totaro, *The topology of smooth divisors and the arithmetic of abelian varieties*, Michigan Math. J. **48** (2000), 611–624, Dedicated to William Fulton on the occasion of his 60th birthday. MR 1786508

- [vdW37] B. L. van der Waerden, *Zur algebraischen Geometrie. X; Über lineare Scharen von reduziblen Mannigfaltigkeiten*, Math. Ann. **113** (1937), no. 1, 705–712. MR 1513118
- [VY08] Oswald Veblen and John Wesley Young, *A Set of Assumptions for Projective Geometry*, Amer. J. Math. **30** (1908), no. 4, 347–380. MR 1506049
- [Wei82] Rainer Weissauer, *Der Hilbertsche Irreduzibilitätssatz*, J. Reine Angew. Math. **334** (1982), 203–220. MR 667458
- [WK81] Roger Wiegand and William Krauter, *Projective surfaces over a finite field*, Proc. Amer. Math. Soc. **83** (1981), no. 2, 233–237. MR 624904
- [Zar41] Oscar Zariski, *Pencils on an algebraic variety and a new proof of a theorem of Bertini*, Trans. Amer. Math. Soc. **50** (1941), 48–70. MR 4241
- [Zha98] Shou-Wu Zhang, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), no. 1, 159–165. MR 1609518

Princeton University, Princeton NJ 08544-1000
kollar@math.princeton.edu