sec1

LIE BIDERIVATIONS ON TRIANGULAR ALGEBRAS

XINFENG LIANG, DANDAN REN AND FENG WEI

ABSTRACT. Let \mathcal{T} be a triangular algebra over a commutative ring \mathcal{R} and $\varphi : \mathcal{T} \times \mathcal{T} \longrightarrow \mathcal{T}$ be an arbitrary Lie biderivation of \mathcal{T} . We will address the question of describing the form of φ in the current work. It is shown that under certain mild assumptions, φ is the sum of an inner biderivation and an extremal biderivation and a some central bilinear mapping. Our results is immediately applied to block upper triangular algebras and Hilbert space nest algebras .

1. INTRODUCTION

Let \mathcal{R} be a commutative ring with identity and \mathcal{A} be an associative \mathcal{R} -algebra with center $\mathcal{Z}(\mathcal{A})$. An \mathcal{R} -linear mapping $d : \mathcal{A} \longrightarrow \mathcal{A}$ is called a *derivation* if d(xy) = d(x)y + xd(y) for all $x, y \in \mathcal{A}$, and is called a *Lie derivation* if

d([x, y]) = [d(x), y] + [x, d(x)]

for all $x \in A$. An \mathcal{R} -linear mapping $d : \mathcal{A} \longrightarrow \mathcal{A}$ of the form $a \mapsto am - ma$ for some $m \in \mathcal{A}$, is said to be an *inner derivation*. A \mathcal{R} -bilinear mapping $\varphi : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ is a *biderivation* if it is a derivation with respect to both components, that is

$$\varphi(xz,y) = \varphi(x,y)z + x\varphi(z,y)$$
 and $\varphi(x,yz) = \varphi(x,y)z + y\varphi(x,z)$

for all $x, y \in \mathcal{A}$. If the algebra \mathcal{A} is noncommutative, then the mapping $\varphi(x, y) = \lambda[x, y]$ for all $x, y \in \mathcal{A}$ and some $\lambda \in \mathcal{Z}(\mathcal{A})$ is called an *inner biderivation*. An \mathcal{R} -bilibear mapping $\varphi : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ is said to be an *extremal biderivation* if it is of the form $\varphi(x, y) = [x, [y, a]]$ for all $x, y \in \mathcal{A}$ and some $a \in \mathcal{A}, a \notin \mathcal{Z}(\mathcal{A})$. An \mathcal{R} -bilinear mapping $\varphi : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ is a *Lie biderivation* if it is a Lie derivation with respect to both components, implying that

$$\varphi([x, z], y) = [\varphi(x, y), z] + [x, \varphi(z, y)] \text{ and}$$
$$\varphi(x, [y, z]) = [\varphi(x, y), z] + [y, \varphi(x, z)]$$

for all $x, y \in \mathcal{A}$.

Suppose that A and B are two unital algebras over \mathcal{R} and M is a nonzero faithful bimodule as a left A-module and also right B-module. Then one can define

$$\left[\begin{array}{cc} A & M \\ 0 & B \end{array}\right] = \left\{ \left[\begin{array}{cc} a & m \\ 0 & b \end{array}\right] \middle| a \in A, b \in B, m \in M \right\}$$

to be an associative algebra under matrix-like addition and matrix-like multiplication. An algebra \mathcal{T} is called a *triangular algebra* if there exist algebras A, B and nonzero faithful (A, B)-bimodule M such that \mathcal{T} is (algebraically) isomorphic to

$$\left[\begin{array}{cc}A&M\\O&B\end{array}\right]$$

under matrix-like addition and matrix-like multiplication. Usually, we denote a triangular algebra by $\mathcal{T} = \begin{bmatrix} A & M \\ O & B \end{bmatrix}$. This kind of algebra was first introduced by Chase in [10]. He applied triangular algebras to show

Date: March 2, 2020.

²⁰⁰⁰ Mathematics Subject Classification. 16W25, 15A78, 47L35.

Key words and phrases. Lie biderivation, derivation, triangular algebra.

The work of the first author is partially supported by Key Projects of Natural Science Research in Anhui Province (Grant No. KJ2019A0107,KJ2018A0082); the second author is partially supported by the National Natural Science Foundation of China (11801008), Key Program of Scientific Research Fund for Young Teachers of AUST (Grant No.QN2017209), talent Introduction Project of Anhui University of Science & Technology(Grant No. 11690).

us the asymmetric behavior of semi-hereditary rings and constructed an classical example of a left semihereditary ring which is not right semi-hereditary. Harada referred to the triangular algebras as generalized triangular matrix rings in the literature [22] which he used triangular algebras to study the structure of hereditary semi-primary rings. The definition of triangular algebra is somewhat formal and hence they are said to be formal triangular matrix algebras in the situation of noncommutative algebras [21].

The concept of biderivation originates from difference or functional equations and their inequalities instead of associative algebras. It was Maksa [25] who initially introduced the concept of biderivations. Vukman [32, 33] investigated biderivations on prime and semiprime rings. Brešar et al. [8] have shown that each biderivation φ on a noncommutative prime ring \mathcal{R} is of the form $\varphi(x, y) = \lambda[x, y]$, for some element λ in the extended centroid of \mathcal{R} . It has turned out that this result can be applied to the problem of describing the form of commuting mappings. We encourage the reader to refer to the survey paper [7] where applications of biderivations to some other areas are provided. The study of commuting mappings and biderivations has its deep roots in the structure theory of associative algebras, where it has proved to be influential and far-reaching, see [7] and references therein. They have been becoming an active research topic in the theory of additive mappings of associative algebras since Brešar's elegant work [6, 8]. On the other hand, an interest in studying these mappings on Lie algebras has been increasing more recently, see [9, 31, 41].

The objective of this paper is to investigate Lie biderivations on triangular algebras. Many authors have made important and essantial contributions to the related topics, see [1, 2, 4–6, 9, 14, 16–20, 23, 26, 31, 34, 36, 37, 41]. Cheung in [11] initiated the study of linear mappings of abstract triangular algebras and obtained a number of elegant results. He gave detailed descriptions concerning automorphisms, derivations, commuting mappings and Lie derivations of triangular algebras in [11, 12]. Benkovič [3] considered Jordan derivations of triangular matrices over a commutative ring with identity and proved that any Jordan derivation from the algebra of all upper triangular matrices into its arbitrary bimodule is the sum of a derivation and an antiderivation. Zhang and Yu [35] observed that each Jordan derivation on a 2-torsion free triangular algebra is a derivation. Generalized Biderivations on nest algebras were also studied by Zhang et al. [36]. They provided a sufficient and necessary condition which enable each generalized biderivation on a complex separable Hilbert space nest algebra to be inner. Zhao et al. [37] investigated biderivations on upper triangular matrix algebras over a commutative ring \mathcal{R} . They proved that each biderivation on the algebra $\mathcal{T}_n(\mathcal{R})$ of all upper triangular $n \times n$ matrices over \mathcal{R} is the sum of an inner biderivation and an extremal biderivation.

Benkovič [4] paid special attention to biderivations on a certain class of triangular algebras. He obtained that a bilinear biderivation φ of a triangular algebra \mathcal{T} satisfying certain conditions (see the conditions (1)-(4) in Theorem 3.2) is of the form $\varphi(x,y) = \lambda[x,y] + [x,[y,r]]$ for some element $\lambda \in \mathcal{Z}(\mathcal{T})$ and some element $r \in \mathcal{T}$. On the other hand, Ghosseiri [19] considered biderivations of an arbitrary triangular ring \mathcal{T} (not assuming M is a faithful (A, B)-bimodule). He proved that each biderivation $\varphi: \mathcal{T} \times \mathcal{T} \longrightarrow \mathcal{T}$ can be decomposed into $\varphi = \tau + \psi + \delta$, where τ is a biderivation satisfying certain conditions, ψ is an extremal biderivation, δ is a special kind of biderivation. Basing on previous Benkovič's and Ghosseiri's works, our [23] use the same as condition (see the conditions (1)-(4) in Theorem 3.8) to obtain the form of Jordan biderivation of a triangular algebras \mathcal{T} ; Eremita [17] used the notion of the maximal left ring of quotients to describe the form of biderivations of a triangular ring. His distinguished approach permit him to achieve double purpose: generalizing Benkovič's result on biderivations [4, Theorem 4.11] and refining Ghosseiri's result on biderivations [19, Theorem 2.4]. More recently, Ghosseiri and his collaborators [14, 20] further characterized the structure of biderivations and superderivations on trivial extensions, which are natural generalizations of the corresponding results on triangular algebras. Wang et al^[38] investigated biderivations on Parabolic Subalgebras of Simple Lie Algebras \mathfrak{g} of rank l over an algebraically closed field of characteristic zero. Let \mathfrak{p} an arbitrary parabolic subalgebra of \mathfrak{g} , they proved that a bilinear map $\varphi: \mathfrak{p} \times \mathfrak{p} \to \mathfrak{p}$ is a biderivation if and only if it is a sum of an inner and an extremal biderivation. Wang and Yu[39] observed that each biderivation of Schr0dinger-Virasoro Lie algebra Ω over the complex field \mathfrak{C} is inner. As an application of biderivations, they show that every linear commuting map φ on Ω has the form $\varphi(x) = \lambda x + \mathfrak{f}(x)M_0$, where $\lambda \in \mathfrak{C}, M_0$ is a basis of the one-dimensional center of Ω , and \mathfrak{f} is a linear function from Ω to \mathfrak{C} . Cheng et al. [40] prove that each skew-symmetric biderivation of the Lie algebra \mathfrak{gca} over the complex field \mathfrak{C} is inner. As an application of biderivations, we will show that every linear commuting map ϕ on the Lie algebra gca has the form $\phi(x) = \lambda x$, where $\lambda \in \mathfrak{C}$. Liu et al.[41] determine the biderivations of the block Lie algebras $\mathcal{B}(q)$ for all $q \in \mathfrak{C}$. More precisely, they prove that the space of biderivations of $\mathcal{B}(q)$ is spanned by inner biderivations and one outer biderivation. Applying this result, they also research all commuting maps on $\mathcal{B}(q)$.

This paper is devoted to the treatment of Lie biderivations of triangular algebras, and its framework is as follows. After Introduction, the second section states some fundamental facts about triangular algebras and demonstrates three classical examples. The kernel question of the current work is to describe the decomposition forms of Lie biderivations on triangular algebras, which takes places in the mainbody–Section 3.

2. Preliminaries

Let \mathcal{R} be a commutative ring with identity. Let A and B be unital algebras over \mathcal{R} . Recall that an (A, B)-bimodule M is *faithful* if for any $a \in A$ and $b \in B$, aM = 0 (resp. Mb = 0) implies that a = 0 (resp. b = 0).

Let A, B be unital associative algebras over \mathcal{R} and M be a unital (A, B)-bimodule, which is faithful as a left A-module and also as a right B-module. We denote the *triangular algebra* consisting of A, B and M by

$$\mathcal{T} = \left[\begin{array}{cc} A & M \\ O & B \end{array} \right].$$

Then \mathcal{T} is an associative and noncommutative \mathcal{R} -algebra. The center $\mathcal{Z}(\mathcal{T})$ of \mathcal{T} is (see [13, Proposition 3])

$$\mathcal{Z}(\mathcal{T}) = \left\{ \left[\begin{array}{cc} a & 0 \\ 0 & b \end{array} \right] \right| am = mb, \ \forall \ m \in M \right\}.$$

Let us define two natural \mathcal{R} -linear projections $\pi_A : \mathcal{T} \to A$ and $\pi_B : \mathcal{T} \to B$ by

$$\pi_A : \begin{bmatrix} a & m \\ 0 & b \end{bmatrix} \longmapsto a \text{ and } \pi_B : \begin{bmatrix} a & m \\ 0 & b \end{bmatrix} \longmapsto b.$$

It is easy to see that $\pi_A(\mathcal{Z}(\mathcal{T}))$ is a subalgebra of $\mathcal{Z}(A)$ and that $\pi_B(\mathcal{Z}(\mathcal{T}))$ is a subalgebra of $\mathcal{Z}(B)$. Furthermore, there exists a unique algebraic isomorphism $\tau \colon \pi_A(\mathcal{Z}(\mathcal{T})) \longrightarrow \pi_B(\mathcal{Z}(\mathcal{T}))$ such that $am = m\tau(a)$ for all $a \in \pi_A(\mathcal{Z}(\mathcal{T}))$ and for all $m \in M$.

Let 1 (resp. 1') be the identity of the algebra A (resp. B), and let I be the identity of the triangular algebra \mathcal{T} . We will use the following notations:

$$e = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad f = I - e = \begin{bmatrix} 0 & 0 \\ 0 & 1' \end{bmatrix}$$

and

xxsec2.1

xxsec2

$$\mathcal{T}_{11} = e\mathcal{T}e, \ \mathcal{T}_{12} = e\mathcal{T}f, \ \mathcal{T}_{22} = f\mathcal{T}f_{22}$$

Thus the triangular algebra \mathcal{T} can be written as

$$\mathcal{T} = e\mathcal{T}e + e\mathcal{T}f + f\mathcal{T}f = \mathcal{T}_{11} + \mathcal{T}_{12} + \mathcal{T}_{22}.$$

Here, \mathcal{T}_{11} and \mathcal{T}_{22} are subalgebras of \mathcal{T} which are isomorphic to A and B, respectively. \mathcal{T}_{12} is a $(\mathcal{T}_{11}, \mathcal{T}_{22})$ bimodule which is isomorphic to the (A, B)-bimodule M. It should be remarked that $\pi_A(\mathcal{Z}(\mathcal{T}))$ and $\pi_B(\mathcal{Z}(\mathcal{T}))$ are isomorphic to $e\mathcal{Z}(\mathcal{T})e$ and $f\mathcal{Z}(\mathcal{T})f$, respectively. Then there is an algebra isomorphism $\tau: e\mathcal{Z}(\mathcal{T})e \longrightarrow f\mathcal{Z}(\mathcal{T})f$ such that $am = m\tau(a)$ for all $m \in e\mathcal{T}f$.

Let us see several classical triangular algebras which will be frequently invoked in the sequel discussion.

2.1. Upper triangular matrix algebras. Let \mathcal{R} be a commutative ring with identity. We denote the set of all $p \times q$ matrices over \mathcal{R} by $M_{p \times q}(\mathcal{R})$ and denote the set of all $n \times n$ upper triangular matrices over \mathcal{R} by $T_n(\mathcal{R})$. For $n \geq 2$ and each $1 \leq k \leq n-1$, the upper triangular matrix algebra $T_n(\mathcal{R})$ can be written as

$$T_n(\mathcal{R}) = \left[\begin{array}{cc} T_k(\mathcal{R}) & M_{k \times (n-k)}(\mathcal{R}) \\ O & T_{n-k}(\mathcal{R}) \end{array} \right].$$

xxsec2.2

xxsec2.3

2.2. Block upper triangular matrix algebras. Let \mathcal{R} be a commutative ring with identity. For each positive integer n and each positive integer m with $m \leq n$, we denote by $\bar{d} = (d_1, \dots, d_i, \dots, d_m) \in \mathbb{N}^m$ an ordered *m*-vector of positive integers such that $n = d_1 + \cdots + d_i + \cdots + d_m$. The block upper triangular matrix algebra $B_n^d(\mathcal{R})$ is a subalgebra of $M_n(\mathcal{R})$ of the form

$$B_{n}^{\bar{d}}(\mathcal{R}) = \begin{bmatrix} M_{d_{1}}(\mathcal{R}) & \cdots & M_{d_{1} \times d_{i}}(\mathcal{R}) & \cdots & M_{d_{1} \times d_{m}}(\mathcal{R}) \\ & \ddots & \vdots & & \vdots \\ & & M_{d_{i}}(\mathcal{R}) & \cdots & M_{d_{i} \times d_{m}}(\mathcal{R}) \\ & & O & \ddots & \vdots \\ & & & M_{d_{m}}(\mathcal{R}) \end{bmatrix} = \\ \vdots & \vdots & \vdots \\ T_{1,1} \cdots T_{1,d_{1}} & \cdots & T_{1,x+d_{i}} \\ \vdots & \ddots & \vdots \\ T_{d_{1},1} \cdots T_{d_{1},d_{1}} & \cdots & T_{1,x+d_{i}} \\ & \ddots & \vdots \\ T_{d_{1},x+1} \cdots T_{d_{1},x+d_{i}} & \cdots & T_{1,x+d_{m}} \\ & \vdots & \ddots & \vdots \\ T_{x+1,x+1} \cdots T_{x+1,x+d_{i}} & \cdots & T_{x+1,y+d_{m}} \\ & \vdots & \ddots & \vdots \\ T_{x+d_{i},x+1} \cdots T_{x+d_{i},x+d_{i}} & \cdots & T_{x+1,y+d_{m}} \\ & \vdots & \ddots & \vdots \\ T_{x+d_{i},y+1} \cdots T_{x+d_{i},y+d_{m}} \\ & \vdots & \ddots & \vdots \\ T_{y+1,y+1} \cdots T_{y+1,y+d_{m}} \\ & \vdots & \ddots & \vdots \\ T_{y+d_{m},y+1} \cdots T_{y+d_{m},y+d_{m}} \end{bmatrix}$$

Note that the full matrix algebra $M_n(\mathcal{R})$ of all $n \times n$ matrices over \mathcal{R} and the upper triangular matrix algebra $T_n(\mathcal{R})$ of all $n \times n$ upper triangular matrices over \mathcal{R} are two special cases of block upper triangular matrix algebras. If $n \ge 2$ and $B_n^{\bar{d}}(\mathcal{R}) \ne M_n(\mathcal{R})$, then $B_n^{\bar{d}}(\mathcal{R})$ is a triangular algebra and can be represented as

$$B_n^{\bar{d}}(\mathcal{R}) = \begin{bmatrix} B_j^{d_1}(\mathcal{R}) & M_{j \times (n-j)}(\mathcal{R}) \\ O_{(n-j) \times j} & B_{n-j}^{\bar{d}_2}(\mathcal{R}) \end{bmatrix},$$

where $1 \leq j < m$ and $\bar{d}_1 \in \mathbb{N}^j, \bar{d}_2 \in \mathbb{N}^{m-j}$.

2.3. Nest algebras. Let **H** be a complex Hilbert space and $\mathcal{B}(\mathbf{H})$ be the algebra of all bounded linear operators on **H**. Let I be a index set. A nest is a set \mathcal{N} of closed subspaces of **H** satisfying the following conditions:

- (1) $0, \mathbf{H} \in \mathcal{N};$
- (2) If $N_1, N_2 \in \mathcal{N}$, then either $N_1 \subseteq N_2$ or $N_2 \subseteq N_1$;

(3) If $\{N_i\}_{i\in I} \subseteq \mathcal{N}$, then $\bigcap_{i\in I} N_i \in \mathcal{N}$; (4) If $\{N_i\}_{i\in I} \subseteq \mathcal{N}$, then the norm closure of the linear span of $\bigcup_{i\in I} N_i$ also lies in \mathcal{N} .

If $\mathcal{N} = \{0, \mathbf{H}\}$, then \mathcal{N} is called a trivial nest, otherwise it is called a non-trivial nest.

The nest algebra associated with \mathcal{N} is the set

$$\mathcal{T}(\mathcal{N}) = \{ T \in \mathcal{B}(\mathbf{H}) \mid T(N) \subseteq N \text{ for all } N \in \mathcal{N} \}.$$

A nontrivial nest algebra is a triangular algebra. Indeed, if $N \in \mathcal{N} \setminus \{0, \mathbf{H}\}$ and E is the orthogonal projection onto N, then $\mathcal{N}_1 = E(\mathcal{N})$ and $\mathcal{N}_2 = (1-E)(\mathcal{N})$ are nests of N and \mathcal{N}^{\perp} , respectively. Moreover, $\mathcal{T}(\mathcal{N}_1) =$ $E\mathcal{T}(\mathcal{N})E, \mathcal{T}(\mathcal{N}_2) = (1-E)\mathcal{T}(\mathcal{N})(1-E)$ are nest algebras and

$$\mathcal{T}(\mathcal{N}) = \begin{bmatrix} \mathcal{T}(\mathcal{N}_1) & E\mathcal{T}(\mathcal{N})(1-E) \\ O & \mathcal{T}(\mathcal{N}_2) \end{bmatrix}$$

Note that any finite dimensional nest algebra is isomorphic to a complex block upper triangular matrix algebra. We refer the reader to [15] for the theory of nest algebras.

xxsec2.4

2.4. Matrix Incidence algebras. Let \mathbb{K} be a field and A be a unital algebra over \mathbb{K} . Let X be a partially ordered set with the partial order \leq . We define the *incidence algebra* of X over A as

$$I(X,A) = \{ f : X \times X \longrightarrow A \mid f(x,y) = 0 \text{ if } x \not\leq y \}$$

with algebraic operation given by

$$(f+g)(x,y) = f(x,y) + g(x,y),$$

$$(f*g)(x,y) = \sum_{x \le z \le y} f(x,z)g(z,y)$$

$$(k \cdot f)(x,y) = k \cdot f(x,y)$$

for all $f, g \in I(X, A), k \in \mathbb{K}$ and $x, y, z \in X$. Obviously, f is a A-valued function on $\{(x, y) \in X \times X | x \leq y\}$. The product * is usually called *convolution* in function theory. In particular, if X be finite partially ordered set with n elements, then I(X, A) is isomorphic to a subalgebra of the algebra $M_n(A)$ of square matrices over \mathbb{K} with elements $[a_{ij}]_{n \times n} \in M_n(A)$ satisfying $a_{ij} = 0$ if $i \leq j$, for some partial order \leq defined in the partial order set (poset) $\{1, \dots, n\}$ [27, Proposition 1.2.4]. More precisely, I(X, A) is isomorphic to an upper triangular matrix algebra with entries A or 0. We will call such incidence algebras matrix incidence algebras. In fact, any incidence algebra arising from a finite partially ordered set is isomorphic to some matrix incidence algebra I(X, A), where \leq is consistent with the natural order. Nevertheless, we can not say that each matrix incidence algebra is a triangular algebra in which M is a faithful (A, B)-bimodule in usual. Not all incidence algebras I(X, A) can be considered as a triangular algebra.

To illustrate this conclusion, let us see an intuitional example. Let $X = \{1, 2, 3, 4, 5, 6, 7\}$ be a partially ordered set and its relations generated by

$$\{1 \le 3, 2 \le 3, 3 \le 4, 4 \le 5, 5 \le 6, 5 \le 7\}$$

We represent this partially ordered set X by the following diagram

 $\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & A & 0 \\ 0 & 0 & 0 & 0 & 0 & A \end{bmatrix}$ The incidence algebra of a partially ordered set (poset) X is the algebra of functions from the segments of X into an K-algebra A, which extends the various convolutions in algebras of arithmetic functions. Incidence algebras, in fact, were first considered by Ward [30] as generalized algebras of arithmetic functions. Rota and Stanley developed incidence algebras as the fundamental structures of enumerative combinatorial theory and allied areas of arithmetic function theory. The theory of Möbius functions, including the classical Möbius function of number theory and the combinatorial inclusion-exclusion formula, is established in the context of incidence algebras are quite striking as well, including the fact that the lattice of ideals (in the finite-dimensional case) is distributive, and that the partial order can be recovered from the algebra. The latter has led to a complete description of the automorphisms and derivations of the algebra [28].

Then we have

lered set A D	y une	1011	Own	ig u	agra	2111		
$X = \begin{bmatrix} 1 \\ & \\ & \\ & \\ & \\ & 2 \end{bmatrix}$	3		→ 4		→ 5		6 7 7].
	$\int A$	0	A	A	A	A	A	1
	0	A	A	A	A	A	A	
	0	0	A	A	A	A	A	
$I(X, A) \cong$	0	0	0	A	A	A	A	.
	0	0	0	0	A	A	A	
		0	0	0	0	A	0	

In the theory of operator algebras, incidence algebras are referred to as "bigraph algebras" or "finite dimensional CSL algebras". For a finite dimensional Hilbert space **H** and the algebra $\mathcal{B}(\mathbf{H})$ of all bounded linear operators on **H**. A digraph algebra is a subalgebra A of $\mathcal{B}(\mathbf{H})$ which contains a maximal abelian self-adjoint subalgebra \mathcal{D} of $\mathcal{B}(\mathbf{H})$. Since \mathcal{D} is maximal abelian, the invariant projections for A, LatA, are elements of \mathcal{D} and so are mutually commuting. Thus A is a CSL-algebra (The abbreviation CSL denotes 'commutative subspace lattice'). Obviously, A is finite dimensional; on the other hand, every finite dimensional CSL-algebra acts on a finite dimensional Hilbert space and contains a mass. The term digraph algebra refers to the fact that associated with A there is a directed graph on the set of vertices $\{1, 2, \dots, n\}$. This graph contains all the self loops. Then A contains the matrix unit e_{ij} if and only if there is a (directed) edge from j to i in the digraph.

Let I(X, A) be an incidence algebra of X over A. The identity element ϵ of I(X, A) is given by $\epsilon(x, y) = \delta_{xy}$ for all $x \leq y$, where $\delta_{xy} \in \{0, 1\}$ is the Kronecker sign. For each pair $x, y \in X$ with $x \leq y$ we define $\epsilon_{xy}(u, v) = \delta_{xu}\delta_{yv}$ for all $u \leq v$. Then $\epsilon_{xy} * \epsilon_{zu} = \delta_{yz}\epsilon_{xu}$ and $\epsilon_{xy}a = a\epsilon_{xy}$ for all $a \in A$. Let $1 \leq n \leq \infty$, let $X = \{1, 2, \dots, n\}$ if $n < \infty$, or $X = \{1, 2, \dots, \}$ if $n = \infty$, and endow X with the usual linear ordering. Then I(X, A) can be identified with the upper triangular matrix algebra $T_n(A)$ by identifying ϵ_{xy} with the matrix $[\delta_{xi}\delta_{yj}]_{i,j=1}^n$. Note that the case $T_{\infty}(A)$ is of infinite matrices. As another extreme case, let $X = \{1, 2, \dots, n\}$ with $1 \leq n < \infty$. If X has the pre-order \leq' , where $i \leq' j$ for each pair $(i, j) \in X \times X$, then $I(X, A) \cong M_n(A)$, the full matrix algebras of $n \times n$ matrices over A. We now deduce some results for the upper triangular matrix algebras $T_n(A)$ with $1 \leq n \leq \infty$ and for the full matrix algebras $M_n(A)$ with $1 \leq n < \infty$ from the results for incidence algebras.

3. Lie Biderivations

This part is the main part of our work, this part is mainly research Lie biderivation of triangular algebras. In order to better explain our work, we prove the following important formulas.

Example 3.1. Let \mathcal{A} be a associative algebra over a commutative ring \mathcal{R} and $\phi : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ be a Lie biderivation on \mathcal{A} , then ϕ has the forwing properties:

$$[\phi(x,a), [b,y]] + [\phi(x,b), [y,a]] = [\phi(y,a), [x,b]] + [\phi(y,b), [x,a]]$$

for all $a, b, x, y \in \mathcal{A}$.

Proof. Let $\phi : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$ be a Lie biderivation of \mathcal{A} . For arbitrary $x, y, a, b \in \mathcal{A}$, let us compute $\phi([x, y], [a, b])$. Since ϕ is a Lie derivation with respective to the first component, we have

$$\begin{split} \phi([x,y],[a,b]) &= [\phi(x,[a,b]),y] + [x,\phi(y,[a,b])] \\ &= [[a,\phi(x,b)] + [\phi(x,a),b],y] + [x,[\phi(y,a),b] + [a,\phi(y,b)]] \\ &= [a\phi(x,b) - \phi(x,b)a + \phi(x,a)b - b\phi(x,a),y] \\ &+ [x,\phi(y,a)b - b\phi(y,a) + a\phi(y,b) - \phi(y,b)a] \\ &= (a\phi(x,b) - \phi(x,b)a + \phi(x,a)b - b\phi(x,a))y \\ &- y(a\phi(x,b) - \phi(x,b)a + \phi(x,a)b - b\phi(x,a)) \\ &+ x(\phi(y,a)b - b\phi(y,a) + a\phi(y,b) - \phi(y,b)a) \\ &- (\phi(y,a)b - b\phi(y,a) + a\phi(y,b) - \phi(y,b)a)x; \end{split}$$
(3.1)

Likewise, the mapping ϕ is a Lie biderivation with respect to the second component as well, we have

$$\begin{split} \phi([x,y],[a,b]) &= [\phi([x,y],a),b] + [a,\phi([x,y],b)] \\ &= [[\phi(x,a),y] + [x,\phi(y,a)],b] + [a,[\phi(x,b),y] + [x,\phi(y,b)]] \\ &= [\phi(x,a)y - y\phi(x,a) + x\phi(y,a) - \phi(y,a)x,b] \\ &+ [a,\phi(x,b)y - y\phi(x,b) + x\phi(y,b) - \phi(y,b)x]] \\ &= (\phi(x,a)y - y\phi(x,a) + x\phi(y,a) - \phi(y,a)x)b \\ &- b(\phi(x,a)y - y\phi(x,a) + x\phi(y,a) - \phi(y,a)x) \\ &+ a(\phi(x,b)y - y\phi(x,b) + x\phi(y,b) - \phi(y,b)x) \\ &- (\phi(x,b)y - y\phi(x,b) + x\phi(y,b) - \phi(y,b)x)a. \end{split}$$
(3.2)

Comparing (3.1) and (3.2), one can obtain

$$[\phi(x,a), [b,y]] + [\phi(y,b), [a,x]] = [\phi(x,b), [a,y]] + [\phi(y,a), [x,b]]$$

For arbitrary $x, y, a, b \in \mathcal{A}$.

Let M be a unital (A, B)-bimodule, the mapping $f : M \to M$ satisfying f(am) = af(m) and f(mb) = f(m)b for all $a \in A, m \in M, b \in B$ is called bimodule homomorphism. A bimodule homomorphism $f : M \to M$ is of the standard form if there exist $a_0 \in \mathcal{Z}(A), b_0 \in \mathcal{Z}(B)$ such that

$$f(m) = a_0 m + m b_0 \tag{3.3}$$

for all $m \in M$.

Below we give the main theorem of this paper.

EXAMPLE 1 Theorem 3.2. Let $\mathcal{T} = \begin{bmatrix} A & M \\ O & B \end{bmatrix}$ be a triangular algebra over a commutative ring \mathcal{R} and let $\phi : \mathcal{T} \times \mathcal{T} \longrightarrow \mathcal{T}$ be a Lie biderivation. If the following conditions holds:

- (i) $\pi_A(\mathcal{Z}(\mathcal{T})) = \mathcal{Z}(A)$ and $\pi_B(\mathcal{Z}(\mathcal{T})) = \mathcal{Z}(B)$;
- (ii) at least one of the algebras A and B is noncommutative;
- (iii) each bimodule homomorphism $f: M \to M$ is of the standard form;
- (iv) if $\alpha a = 0$, $\alpha \in \mathcal{Z}(A)$, $0 \neq a \in A$, then $\alpha = 0$.

Then every Lie biderivation $\phi : \mathcal{T} \times \mathcal{T} \to \mathcal{T}$ is of the form

$$\phi(x, y) = \lambda_0[x, y] + [x, [y, \phi(e, e)]] + \mu(x, y)$$

where for some $\lambda_0 \in \mathcal{Z}(\mathcal{T})$ and $\mu : \mathcal{T} \times \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ is a central mapping for arbitrary $x, y \in \mathcal{T}$.

In order to better prove the main theorem, we need to obtain the following series of lemmas.

Exerc3.3 Lemma 3.3. Let $\mathcal{T} = \begin{bmatrix} A & M \\ O & B \end{bmatrix}$ be a triangular algebra over a commutative ring \mathcal{R} and let $\phi : \mathcal{T} \times \mathcal{T} \longrightarrow \mathcal{T}$ be a Lie biderivation.

- (1) $\phi(0, x) = \phi(x, 0) = 0;$
- (2) $\phi(1,x) = e\phi(1,x)e \oplus f\phi(1,x)f \in \mathcal{Z}(\mathcal{T})$ and $\phi(x,1) = e\phi(x,1)e \oplus f\phi(x,1)f \in \mathcal{Z}(\mathcal{T})$;
- (3) $e\phi(e,e)f = -e\phi(f,e)f = -e\phi(e,f)f = e\phi(f,f)f.$

for all $x \in \mathcal{T}$

Proof.

(1) Since ϕ is a Lie derivation with respect to the first component, we have $\phi(0, x) = \phi([0, 0], x)$

$$(0, x) = \phi([0, 0], x)$$

= $[0, \phi(0, x)] + [\phi(0, x), 0] = 0$ (3.4)

for all $x \in \mathcal{T}$.

(2) Since ϕ is a Lie derivation with respect to the first component and also use the relation (3.4), we have

$$0 = \phi(0, x) = \phi([1, y], x)$$

= $[\phi(1, x), y] + [1, \phi(y, x)] = [\phi(1, x), y],$
7

for arbitrary $x, y \in \mathcal{T}$. Because of the arbitrariness of element $y \in \mathcal{T}$, one can obtain $\phi(1, x) \in \mathcal{Z}(\mathcal{T})$. Furthermore, we have By an analogous manner of $\phi(1, x)$, we have $\phi(x, 1) \in \mathcal{Z}(\mathcal{T})$ for all $x \in \mathcal{T}$. (3) In view of (2) one can obtain $e\phi(1, x)f = 0$ for all $x \in \mathcal{T}$. Further using the relation e + f = 1 and taking

(3) In view of (2), one can obtain $e\phi(1, x)f = 0$ for all $x \in \mathcal{T}$. Further using the relation e + f = 1 and taking x = e and x = f respectively, we get

$$e\phi(e,e)f = -e\phi(f,e)f$$
 and $e\phi(e,f)f = -e\phi(f,f)f.$ (3.5)

By an analogous manner, according to the relation (2), we obtain $e\phi(x,1)f = 0$ for all $x \in \mathcal{T}$. Furthermore, we have

$$e\phi(e,e)f = -e\phi(e,f)f$$
 and $e\phi(f,e)f = -e\phi(f,f)f.$ (3.6)

Comparing (3.5) with (3.6), we have

$$e\phi(e,e)f = -e\phi(f,e)f = -e\phi(e,f)f = e\phi(f,f)f.$$

xxsec3.4 Lemma 3.4. With notations as above, we have

(1) $\phi(a,m) = \alpha_0 am = -\phi(m,a);$ (2) $\phi(b,m) = \alpha_0 mb = -\phi(m,b)$ for all $a \in A, b \in B, m \in M.$

Proof. Since ϕ is a Lie derivation with respect to the second component, we have

$$\begin{split} \phi(a,m) &= \phi(a,[e,m]) = [\phi(a,e),m] + [e,\phi(a,m)] \\ &= \phi(a,e)m - m\phi(a,e) + e\phi(a,m) - \phi(a,m)e \end{split}$$

for all $a \in A, m \in M$. Multiplying the above equation by e on the left side and by e on the right side, we have $e\phi(a, m)e = 0$. Similarly, one can obtain $f\phi(a, m)f = 0$ and

$$e\phi(a,e)m = m\phi(a,e)f,\tag{3.7}$$

for all $a \in A, m \in M$. Based on the above three formulas, we can get

0

$$\phi(a,m) = e\phi(a,m)f \tag{3.8}$$

for all $a \in A, \in M$.

Since ϕ is a Lie derivation with respect to the second component, we have

$$= \phi(a, [b, e]) = [\phi(a, b), e] + [b, \phi(a, e)]$$

= $\phi(a, b)e - e\phi(a, b) + b\phi(a, e) - \phi(a, e)b$

for all $a \in A, b \in B$. Multiplying the above equation by f on the left side and f on the right side, we have $b\phi(a, e)f = f\phi(a, e)b$, and then

$$f\phi(a,e)f \in \mathcal{Z}(B) \tag{3.9}$$

for all $a \in A, b \in B$. Combining (3.7) and (3.9) together with the faithfulness of A-left module M, one can obtain

$$e\phi(a,e)e \oplus f\phi(a,e)f \in \mathcal{Z}(\mathcal{T}) \tag{3.10}$$

for all $a \in A, m \in M$. Similarly, we have

$$e\phi(a,f)e \oplus f\phi(a,f)f \in \mathcal{Z}(\mathcal{T})$$
(3.11)

for all $a \in A, m \in M$.

Since ϕ is a Lie derivation with respect to the first component, we have

$$0 = \phi(0,m) = \phi([a,e],m) = [\phi(a,m),e] + [a,\phi(e,m)]$$

= $\phi(a,m)e - e\phi(a,m) + a\phi(e,m) - \phi(e,m)a$ (3.12)

for all $a \in A, m \in M$. Multiplying in (3.12) by e on the left side and by f on the right side, we have

$$e\phi(a,m)f = a\phi(e,m)f$$
⁸
(3.13)

for all $a \in A, m \in M$. Considering (3.8) and (3.13), we conclude that

$$\phi(a,m) = a\phi(e,m)f \in M$$

for all $a \in A, m \in M$. In analogous manner, one can check that

$$\phi(m, a) = a\phi(m, e)f \text{ and } e\phi(e, a)e \oplus f\phi(e, a)f \in \mathcal{Z}(\mathcal{T}) \text{ and } e\phi(f, a)e \oplus f\phi(f, a)f \in \mathcal{Z}(\mathcal{T});$$

$$\phi(b, m) = e\phi(f, m)b \text{ and } e\phi(b, e)e \oplus f\phi(b, e)f \in \mathcal{Z}(\mathcal{T}) \text{ and } e\phi(b, f)e \oplus f\phi(b, f)f \in \mathcal{Z}(\mathcal{T});$$

$$\phi(m, b) = e\phi(m, f)b \text{ and } e\phi(e, b)e \oplus f\phi(e, b)f \in \mathcal{Z}(\mathcal{T}) \text{ and } e\phi(f, b)e \oplus f\phi(f, b)f \in \mathcal{Z}(\mathcal{T})$$
(3.14)

for all $a \in A, b \in B, m \in M$.

Let us see a mapping $\mathfrak{h}: M \to M$ be defined by $\mathfrak{h}(m) = e\phi(e, m)f$ for all $m \in M$, then \mathfrak{h} is a bimodule homomorphism as a left A-module and also right B-module for all $m \in M$.

Namely, for arbitrary $a \in A, b \in B, m \in M$, since ϕ is a Lie derivation with respect to the first argument, in light of the relation (3.10) and (3.14) we can have

$$\begin{split} \mathfrak{h}(am) &= e\phi(e, am)f \\ &= e\phi(e, [a, m])f \\ &= e([\phi(e, a), m] + [a, \phi(e, m)])f \\ &= e(\phi(e, a)m - m\phi(e, a)f + a\phi(m, e) - \phi(m, e)a)f \\ &= m(\eta(e\phi(e, a)e) - \phi(e, a)f) + a\phi(m, e)f \\ &= a\phi(m, e)f \\ &= a\mathfrak{h}(m) \end{split}$$

and

$$\begin{split} \mathfrak{h}(mb) &= e\phi(e,mb)f \\ &= e\phi(e,[m,b])f \\ &= e([\phi(e,m),b] + [m,\phi(e,b)]) \\ &= e(\phi(e,m)b - b\phi(e,m) + m\phi(e,b) - \phi(e,b)m)f \\ &= e\phi(e,m)b + m\phi(e,b)f - e\phi(e,b)m \\ &= e(\phi(e,m)b + m(f\phi(e,b)f - \eta(e\phi(e,b)e))f \\ &= e\phi(e,m)b \\ &= \mathfrak{h}(m)b. \end{split}$$

The assumption (iii) implies that the bimodule homomorphism \mathfrak{h} is of the **standard form**

 $\mathfrak{h}(m) = a_0 m + m b_0 = e \phi(e, m) f$

for some $a_0 \in \mathcal{Z}(A)$ and $b_0 \in \mathcal{Z}(B)$ and all $m \in M$. Now we use the assumption (i) to see that $a_0 \in \pi_A(\mathcal{Z}(\mathcal{T}))$ and $b_0 \in \pi_B(\mathcal{Z}(\mathcal{T}))$. We may write

$$\mathfrak{h}(m) = \phi(e, m) = e\phi(e, m)f = (a_0 + \eta^{-1}(b_0))m = \alpha_0 m$$

for all $m \in M$, where $\alpha_0 = a_0 + \eta^{-1}(b_0) \in \pi_A(\mathcal{Z}(\mathcal{T}))$.

Likely, one can define a mapping $\mathfrak{g}: M \to M$ defined by $\mathfrak{g}(m) = e\phi(m, e)f$ for all $m \in M$, which is a bimodule homomorphism as a left A-module and also right B-module. So there exists $\beta_0 \in \pi_A(\mathcal{Z}(\mathcal{T}))$ so that $\phi(m, e) = \beta_0 m$.

Let us next show that

$$\mathfrak{h}(m) = \alpha_0 m = -\mathfrak{g}(m), \text{ i.e., } e\phi(e,m)f == \alpha_0 m = -e\phi(m,e)f$$

for all $m \in M$. We need to prove that $\alpha_0 + \beta_0 = 0$.

According to the assumption (ii), we may assume that A is a noncommutative algebra. Choose $a, a' \in A$ such that $[a, a'] \neq 0$. Since $\phi(e, m) = \alpha_0 m$ and $\phi(m, e) = \beta_0 m$, we by Lemma 3.1 get we have

$$[\varphi(a, a'), [e, m]] + [\varphi(a, e), [a', m]] = [\varphi(m, e), [a, a']] + [\varphi(m, a'), [a, e]]$$

for all $a, a' \in A, m \in M$. In view of the relation (3.10), we have

$$[\varphi(a,a'),m] = -[a,a']\varphi(m,e)f = -[a,a']\alpha_0 m, \qquad (3.15)$$

for all $a, a' \in A, m \in M$.

Adopting similar methods and using Lemma 3.1, we have

$$[\varphi(a, a'), [m, e]] + [\varphi(a, m), [a', e]] = [\varphi(e, m), [a, a']] + [\varphi(e, a'), [a, m]]$$

for all $a, a' \in A, m \in M$. In view of the relation (3.10), we obtain

$$[\varphi(a, a'), m] = [a, a']\varphi(e, m)f = [a, a']\beta_0 m$$
(3.16)

for all $a, a' \in A, m \in M$.

It follows from the equalities (3.15) and (3.16) yields $(\alpha_0 + \beta_0)[a, a']m = 0$ for all $m \in M$. The faithfulness of the left A-module M now implies $(\alpha_0 + \beta_0)[a, a'] = 0$. Since $[a, a'] \neq 0$ we conclude using the condition (iv) that $\alpha_0 + \beta_0 = 0$. Considering $\alpha_0 + \beta_0 = 0$ and $\varphi(f, m) + \varphi(e, m) = 0$ together with $\varphi(m, e) + \varphi(m, f) = 0$, we see that

$$\varphi(m, f) = \alpha_0 m = -\varphi(f, m)$$

for all $m \in M$.

Let $a \in A$, and $m \in M$ be arbitrary elements, we can achieve

$$\varphi(a,m) = a\varphi(e,m)f = \alpha_0 am.$$

This proves the first equality. The other three equations can be proven in an analogous manner. \Box

xxsec3.5 Le

Lemma 3.5. With notations as above, we have

- (1) $\phi(a,b) = e\phi(a,b)e a\phi(e,e)b + f\phi(a,b)f$, where $e\phi(a,b)e \oplus f\phi(a,b)f \in \mathcal{Z}(\mathcal{T})$;
- (2) $\phi(b,a) = e\phi(b,a)e a\phi(f,f)b + f\phi(b,a)f$, where $e\phi(b,a)e \oplus f\phi(b,a)f \in \mathcal{Z}(\mathcal{T})$

for all $a \in A, b \in B$.

Proof.

(1) Since ϕ is a Lie derivation with respect to the first component, we have

$$0 = \phi([e, a], b)$$

= [e, $\phi(a, b)$] + [$\phi(e, b), a$]
= $e\phi(a, b) - \phi(a, b)e + \phi(e, b)a - a\phi(e, b)$

for all $a \in A, b \in B$. Multiplying the above equation by e on the left side and by f on the right side, we can obtain

$$e\phi(a,b)f = a\phi(e,b)f \tag{3.17}$$

for all $a \in A, b \in B$.

Since ϕ is a Lie derivation with respect to the second component, one can have

$$0 = \phi([b_1, a], b_2)$$

= $[b_1, \phi(a, b_2)] + [\phi(b_1, b_2), a]$
= $b_1\phi(a, b_2) - \phi(a, b_2)b_1 + \phi(b_1, b_2)a - a\phi(b_1, b_2)$

for all $b_1, b_2 \in B, a \in A$. Multiplying the above equation by e on the left side and by e on the right side, we can obtain $e\phi(b_1, b_2)a = a\phi(b_1, b_2)e$ for all $b_1, b_2 \in B$. And then

$$e\phi(b_1, b_2)e \in \mathcal{Z}(A) \tag{3.18}$$

for all $b_1, b_2 \in B$. Multiplying the above equation by e on the left side and by f on the right side, we can obtain

$$e\phi(a, b_2)b_1 = -a\phi(b_1, b_2)f; \tag{3.19}$$

for all $b_1, b_2 \in B, a \in A$. Multiplying the above equation by f on the left side and by f on the right side, we can obtain the relation $b_1\phi(a, b_2)f = f\phi(a, b_2)b_1$ for all $a \in A, b_1, b_2 \in B$. And then

$$f\phi(a,b_2)f \in \mathcal{Z}(B) \tag{3.20}$$

for all $a \in A, b_1, b_2 \in B$.

Similarly, Since ϕ is a Lie derivation with respect to the second component, we have

$$0 = \phi(a_1, [b, a_2])$$

= $[\phi(a_1, b), a_2] + [b, \phi(a_1, a_2)]$
= $\phi(a_1, b)a_2 - a_2\phi(a_1, b) + b\phi(a_1, a_2) - \phi(a_1, a_2)b$

for all $a_1, a_2 \in A, b \in B$. Multiplying the above equation by e on the left side and by e on the right side, one can check $e\phi(a_1, b)a_2 = a_2\phi(a_1, b)e$, ie.,

$$e\phi(a_1, b)e \in \mathcal{Z}(A) \tag{3.21}$$

for all $a_1, a_2 \in A, b \in B$. Multiplying the above equation by f on the left side and by f on the right side, we can obtain $b\phi(a_1, a_2)f = f\phi(a_1, a_2)b$, and then

$$f\phi(a_1, a_2)f \in \mathcal{Z}(B) \tag{3.22}$$

for all $a_1, a_2 \in A, b \in B$. Multiplying the above equation by e on the left side and f on the right side, we can obtain

$$a_2\phi(a_1,b)f = -e\phi(a_1,a_2)b \tag{3.23}$$

for all $a_1, a_2 \in A, b \in B$. Combining (3.19) and (3.23) with the conclusion (2) coming from Lemma 3.3, we have

$$e\phi(a,b)f = -a\phi(e,e)b$$

for all $a \in A, b \in B$. We therefore have

$$\phi(a,b) = e\phi(a,b)e - a\phi(e,e)b + f\phi(a,b)f$$

Now, we prove the following relation

$$e\phi(a,b)e \oplus f\phi(a,b)f \in \mathcal{Z}(\mathcal{T})$$

for all $a \in A, b \in B$.

Namely, according to Lemma 3.1, we receive the relation

$$[\phi(a,b),[m,e]] + [\phi(a,m),[e,b]] = [\phi(e,b),[m,a]] + [\phi(e,m),[a,b]]$$

for all $a \in A, b \in B, m \in M$. In view of the relation $e\phi(e, b)e \oplus f\phi(e, b)f \in \mathcal{Z}(\mathcal{T})$, we can obtain

 $e\phi(a,b)m - m\phi(a,b)f = -e\phi(e,b)am + am\phi(e,b)f$

$$= (\eta^{-1}(f\phi(e,b)f) - e\phi(e,b)e)am = 0$$

for all $a \in A, b \in B, m \in M$. Using the relations (3.20), (3.21) and the faithfulness of left A-module M, we have

$$e\phi(a,b)e \oplus f\phi(a,b)f \in \mathcal{Z}(\mathcal{T})$$

for all $a \in A, b \in B$.

(2) By an analogous manner of (1), we have

$$\phi(b,a) = e\phi(b,a)e - a\phi(f,f)b + f\phi(b,a)f,$$

where $e\phi(b, a)e \oplus f\phi(b, a)f \in \mathcal{Z}(\mathcal{T})$ for all $a \in A, b \in B$.

xxsec3.6 Lemma 3.6. With notations as above, we have

 $\phi(m,n) = 0$

for all $m, n \in M$.

Proof. For all $m, n \in M$, we have

$$\varphi(m,n) = \varphi([e,m],n)$$

= $[\varphi(e,n),m] + [e,\varphi(m,n)]$
= $\varphi(e,n)m - m\varphi(e,n) + e\varphi(m,n) + \varphi(m,n)e.$
11

Multiplying the above equation by e on the left and by e on the right side, we have $e\varphi(m, n)e = 0$ for all $m \in M, n \in N$. Similarly, one can obtain $f\varphi(m, n)f = 0$ for all $m \in M, n \in N$. By invoking to above relations, we immediately see that

$$\varphi(m,n) = e\varphi(m,n)f \in M \tag{3.24}$$

for all $m \in M, n \in N$.

Fix a element $m \in M$, then the mapping $\mathfrak{k} : M \to M$ defined by $\mathfrak{k}(m) = \varphi(m, n) = e\varphi(m, n)f$ for all $n \in M$ is a bimodule homomorphism as a left A-module and also right B-module.

In fact, using Lemma 3.8 and (3.24), we have

$$\begin{split} \mathfrak{k}(an) &= e\varphi(m, [a, n])f \\ &= e([\varphi(m, a), n] + [a, \varphi(m, n)])f \\ &= e(\varphi(m, a)n - n\varphi(m, a) + a\varphi(m, n) - \varphi(m, n)a)f \\ &= a\varphi(m, n)f \\ &= a\mathfrak{k}(n) \end{split}$$

and

xxsec3.7

$$\begin{split} \mathfrak{k}(nb) &= e\varphi(m,nb)f\\ &= \varphi(m,[n,b])\\ &= e([\varphi(m,n),b] + [n,\varphi(m,b]))\\ &= e(\varphi(m,n)b - b\varphi(m,n) + n\varphi(m,b) - \varphi(m,b)n)f\\ &= e\varphi(m,n)b\\ &= \mathfrak{k}(n)b \end{split}$$

for all $b \in B, m, n \in M$. For fixing $m \in M$, it follows from the assumption (iii) that there exists $\alpha_m \in \mathcal{Z}(A)$ such that

$$\varphi(m,n) = k(n) = \alpha_m n \text{ for all } n \in M.$$
(3.25)

Without loss of generality, we might assume that that A is a noncommutative algebra, and let $a, a' \in A$ be fixed elements such that $[a, a'] \neq 0$. Using Lemma 3.1, we have

$$[\varphi(a, a'), [n, m]] + [\varphi(a, n), [a', m]] = [\varphi(m, n), [a, a']] + [\varphi(m, a'), [a, m]],$$

for all $m, n \in M$. Using (3.24) and (3.25), we may write

$$0 = [\varphi(m, n), [a, a']] = [a, a']\varphi(m, n) = [a, a']\alpha_m n$$

for all $m, n \in M$. The faithfulness of the left A-module implies $[a, a']\alpha_m = 0$ for every $m \in M$. In view of the assumption (iv), we obtain that $\alpha_m = 0$ for all $m \in M$. We therefore say that $\varphi(m, n) = 0$ for all $m, n \in M$.

Lemma 3.7. With notations as above, we have

(1) For arbitrary $a_1, a_2 \in A$, we have

$$\begin{aligned} \phi(a_1, a_2) &= e\phi(a_1, a_2)e + a_1 a_2 \phi(e, e)f + f\phi(a_1, a_2)f \\ &= e\phi(a_1, a_2)e + a_2 a_1 \phi(e, e)f + f\phi(a_1, a_2)f, \end{aligned}$$

where $f\phi(a_1, a_2)f \in \mathcal{Z}(A)$ and $e\phi(a_1, a_2)e = \eta^{-1}(f\phi(a_1, a_2)f) - \alpha_0[a_1, a_2];$ (2) For arbitrary $a_1, a_2 \in A$, we have

$$\begin{split} \phi(b_1,b_2) &= e\phi(b_1,b_2)e + e\phi(e,e)b_1b_2 + f\phi(b_1,b_2)f \\ &= e\phi(b_1,b_2)e + e\phi(e,e)b_1b_2 + f\phi(b_1,b_2)f, \end{split}$$

where $e\phi(b_1, b_2)e \in \mathcal{Z}(A)$ and $f\phi(b_1, b_2)f = \eta(e\phi(b_1, b_2)e) - \eta(\alpha_0)[b_1, b_2]$ for all $b_1, b_2 \in B$.

Proof. The conclusion (1) and conclusion (2) can be obtained by the similar ways. For the sake of conciseness, we now only prove conclusion (1).

(1). Combining (3.19) and (3.23) together with the relation $e\phi(1, x)f = 0$ in Lemma 3.3, we obtain

$$e\phi(a_1, a_2)f = -a_2a_1\phi(e, f)f = a_2a_1\phi(e, e)f.$$
(3.26)

for all $a_1, a_2 \in A, x \in T$. In similar methods, we have

$$e\phi(a_1, a_2)f = a_1 a_2 \phi(e, e)f \tag{3.27}$$

for all $a_1, a_2 \in A$. In view of above relations ((3.26)) and (3.27), we arrive at

$$e\phi(a_1, a_2)f = a_2a_1\phi(e, e)f = a_1a_2\phi(e, e)f$$
(3.28)

for all $a_1, a_2 \in A$.

According to Lemma 3.1, we have

$$[\phi(a_1, a_2), [e, m]] + [\phi(a, e), [m, a_2]] = [\phi(m, a_2), [e, a]] + [\phi(m, e), [a, a_2]]$$

for all $a_1, a_2, m \in M$. In view of the relation , we receive at

$$[\phi(a_1, a_2), m] - [\phi(a, e), a_2m] = -[a, a_2]\phi(m, e)$$
(3.29)

for all $a_1, a_2 \in A, m \in M$ Combining (3.29) and (3.10) together with Lemma 3.8, we can achieve

$$(e\phi(a_1, a_2)e - \eta^{-1}(f\phi(a_1, a_2)f) - \alpha_0[a_1, a_2])m = 0$$

for all $a_1, a_2, m \in M$. The faithfulness of left A-module M now implies

$$e\phi(a_1, a_2)e = \eta^{-1}(f\phi(a_1, a_2)f) + \alpha_0[a, a_2]$$

for all $a_1, a_2 \in A$.

Based on the above process, one can check that

$$\phi(a_1, a_2) = \eta^{-1}(f\phi(a_1, a_2)f) + f\phi(a_1, a_2)f + a_1a_2\phi(e, e)f + \alpha_0[a, a_2]$$

for all $a_1, a_2 \in A$.

(2). In analogous manner of (1), we have

$$\phi(b_1, b_2) = e\phi(b_1, b_2)e + e\phi(e, e)b_1b_2 + f\phi(b_1, b_2)f,$$

where

$$f\phi(b_1, b_2)f = \eta^{-1}(e\phi(b_1, b_2)e) + \eta(\alpha_0)[b_1, b_2]$$

for all $b_1, b_2 \in B$.

Let us now give the proof of our main theorem.

Proof of Theorem 3.2.

At the end, in order to get the final main theorem, we need to obtain the following equation:

$$aa'\phi(e,e)f - a\phi(e,e)b' - a'\phi(e,e)b + e\phi(e,e)b' = [x, [y, \phi(e,e)]]$$

for all x = a + m + b and y = a' + m' + b' and all $a, a' \in A, b, b' \in B, m, m' \in M$. In fact, let $x = \begin{bmatrix} a & m \\ b \end{bmatrix} \in \mathcal{T}$ and $y = \begin{bmatrix} a' & m' \\ b' \end{bmatrix} \in \mathcal{T}$, Taking into account the relation $e\phi(e, e)e \oplus f\phi(e, e)f \in \mathcal{T}$.

 $\mathcal{Z}(\mathcal{T})$, we arrive at

$$\begin{bmatrix} x, [y, \phi(e, e)] \end{bmatrix}$$

$$= \begin{bmatrix} \begin{bmatrix} a & m \\ b \end{bmatrix}, \begin{bmatrix} \begin{bmatrix} a' & m' \\ b' \end{bmatrix}, \begin{bmatrix} e\phi(e, e)e & e\phi(e, e)f \\ f\phi(e, e)f \end{bmatrix} \end{bmatrix}$$

$$= \begin{bmatrix} \begin{bmatrix} a & m \\ b \end{bmatrix}, \begin{bmatrix} 0 & a'\phi(e, e)f - e\phi(e, e)b' \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & a(a'\phi(e, e)f - e\phi(e, e)b') - (a'\phi(e, e)f - e\phi(e, e)b')b \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & aa'\phi(e, e)f - a\phi(e, e)b' - a'\phi(e, e)b + e\phi(e, e)b' \\ 0 \end{bmatrix},$$

for all $a, a' \in A, b, b' \in B, m, m' \in M$

Let x = a + m + b and y = a' + m' + b', according to the bilinearity of the mapping ϕ , we get the following decomposition form

$$\begin{split} \phi(x,y) =& \phi(a,a') + \phi(a,m') + \phi(a,b') \\ &+ \phi(m,a') + \phi(m,m') + \phi(m,b') \\ &+ \phi(b,a') + \phi(b,m') + \phi(b,b') \\ &= \alpha_0[a,a'] - \alpha_0 am' + \alpha_0 a'm + \alpha_0 mb' - \alpha_0 m'b + \eta(\alpha_0)[b,b'] \\ &+ \eta^{-1}(f\phi(a_1,a_2)f) + f\phi(a_1,a_2)f + e\phi(b_1,b_2)e + \eta(e\phi(b_1,b_2)e) \\ &+ e\phi(a,b)e + f\phi(a,b)f + e\phi(b,a)e + f\phi(b,a)f \\ &+ aa'\phi(e,e)f - a\phi(e,e)b' - a'\phi(e,e)b + e\phi(e,e)b' \\ &= \lambda_0[x,y] + [x, [y, \phi(e,e)]] + \mu(x,y) \end{split}$$

where $\mu: \mathcal{T} \times \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ is a central mapping such that

$$\mu(x,y) = \eta^{-1}(f\phi(a_1,a_2)f) + f\phi(a_1,a_2)f + e\phi(b_1,b_2)e + \eta(e\phi(b_1,b_2)e) + e\phi(a,b)e + f\phi(a,b)f + e\phi(b,a)e + f\phi(b,a)f = \left[\begin{bmatrix} \eta^{-1}(f\phi(a_1,a_2)f) + e\phi(b_1,b_2)e + e\phi(a,b)e + e\phi(b,a)e & 0 \\ f\phi(a_1,a_2)f + \eta(e\phi(b_1,b_2)e) + f\phi(a,b)f + f\phi(b,a)f \end{bmatrix} \in \mathcal{Z}(\mathcal{T})$$

for all $a, a' \in A, b, b' \in B, m, m' \in M$.

As a direct corollary of Theorem 3.2, we get Lie biderivations of the (block) upper triangular matrix algebra and the nest algebra following from the main theorem.

- **Corollary 3.8.** Let C be a commutative domain with identity. If $n \geq 3$, then each Lie biderivation of (block) upper triangular matrix algebra $B_n^k(C)$ is the sum of an extremal biderivation and an inner biderivation. In particular, every biderivation of upper triangular matrix algebra $T_n(C)$ is the sum of an extremal biderivation and an inner biderivation and central mapping.
 - **Corollary 3.9.** Let \mathcal{N} be a nest of a Hilbert space H, where dimH > 3. Then each Lie biderivation φ of nest algebra $\mathcal{T}(\mathcal{N})$ is the sum of an extremal biderivation and an inner biderivation and and central mapping.

References

- [1] C. Abdioğlu and T.-K. Lee, A basic functional identity with applications to Jordan σ -biderivations, Comm. Algebra, 45 (2017), 1741-1756. 2
- [2] D. Aiat Hadj Ahmed, On Jordan biderivationa of trianular matrix rings, J. Math Res Appl., 36 (2016), 162-170. 2
- [3] D. Benkovič, Jordan derivations and antiderivations on triangular matrices, Linear Algebra Appl., 397 (2005), 235-244. 2
- [4] D. Benkovič, Biderivations of triangular algebras, Linear Algebra Appl., 431 (2009), 1587-1602. 2
- [5] D. Benkovič and D. Eremita, Commuting traces and commutativity preserving maps on triangular algebras, J. Algebra, 280 (2004), 797-824. 2
 - [6] M. Brešar, On generalized biderivation and related maps, J. Algebra., 172 (1995), 764-786. 2
- [7] M. Brešar, Commuting maps: A Survey, Taiwanese J. Math., 8 (2004), 361-397. 2
- [8] M. Brešar, W.S. Martindale 3rd, C.R. Miers Centalizing maps in prime rings with involution, J. Algebra., **161** (1993), 342-357. **2**
- [9] M. Brešar and K.-M. Zhao, Biderivations and commuting linear maps on Lie algebras, J. Lie Theory, **28** (2018), 885-900. 2
- [10] S. U. Chase, A generalization of the ring of triangular matrices, Nagoya Math. J., 18 (1961), 13-25. 1
- [11] W.-S. Cheung, Maps on triangular algebras, Ph.D. Dissertation, University of Victoria, 2000. 172pp. 2
- [12] W.-S. Cheung, Commuting maps of triangular algebras, J. London Math. Soc., 63 (2001), 117-127. 2

xxsec3.4

xxsec3.5

i	og	luI	Lee	e20)1	7

Ahmed2016

Benkovic2005

Benkovic2009 cEremita2004



esarZhao2018



- Cheung2003 hosseiri2019 Davidson1988 DuWang2013 Eremita2017 Fosner2015 hosseiri2013 hosseiri2017 HaghanyVara Harada ngRenWei2019 uGuoZhao2018 Maksa1980 MGRS02017
- lDonnell1997 Stanley1970
- Stanley1997
- Ward1937 Tang2018
- Vukman1989
- Vukman1990
- Wang2016
- ZhangYu2006
- FengLiWu2006
- loWangYao2009
- ngYuChen2011
- WangYu2013
- uGuoZhao2018 [41] X.-W.Liu, X.-Q. Guo and K.-M. Zhao Biderivations of the block Lie algebras, Linear Algebra Appl., **538** (2018), 43-55. 2, 3

- [13] W.-S. Cheung, Lie derivations of triangular algebras, Linear Multilinear Algebras., 51 (2003), 299-310. 3
- [14] H. Cheraghpour and N. M. Ghosseiri, On superderivations and superbiderivation of trivial extensions and triangular matrix rings, Comm. Algebra, 47 (2019), 1662-1670. 2
- [15]K. R. Davidson, Nest algebras, Pitman Research Notes in Mathematics Series, 191, Longman, London/New York, 1988. 4
- [16] Y.-Q. Du and Y. Wang, Biderivation of generalized matrix algebras, Linear Algebra Appl., 438 (2013), 4483-4499. 2
- [17] D. Eremita, Biderivations of triangular rings revisited, Bull. Malays. Math. Soc., 40 (2017), 505-527. 2
- [18] A. Fošner, On generalized α -biderivations, Mediterr. J. Math., 12 (2017), 1-7. 2
- [19] N. M. Ghosseiri, On biderivation of upper triangular matrix rings, Linear Algebra Appl., 438 (2013), 250-260. 2
- [20] N. M. Ghosseiri, On derivations and biderivation of trivial extensions and triangular matrix rings, Bull. Iranian Math. Soc., 43 (2017), 1629-1644. 2
- [21] A. Haghany and K. Varadarajan, Study of formal triangular matrix rings, Comm. Algebra, 27 (1999), 5507-5525. 2
- [22] M. Harada, Hereditary semi-primary rings and triangular matrix rings, Nagoya Math. J., 27 (1966), 463-484. 2
- [23] X-F. Liang, D-D. Ren and F. Wei, Jordan biderivations of triangular algebras, Submitted. 2
- [41] X.-W. Liu, X.-Q. Guo and K.-M. Zhao, *Biderivations of the block Lie algebras*, Linear Algebra Appl., **538** (2018), 43-55. 2, 3
- [25] G. Maksa, A remark on symmetric biadditive functions having nonnegative diagonalization, Glasnik Mat. Ser. III, 15(35) (1980), 279-282. 2
- [26] C. Martín González, J. Repka and J. Sánchez-Ortega, Automorphisms, σ -biderivations and σ -commuting maps of triangular algebras, Mediterr. J. Math., 14 (2017), 14:68. 2
- [27] E. Spiegel and C. J. O'Donnell, Incidence Algebras, Monographs and Textbooks in Pure and Applied Mathematics, vol. 206, Marcel Dekker, New York, 1997. 5
 - [28] R. P. Stanley, Structure of incidence algebras and their automorphism groups, Bull. Amer. Math. Soc., **76** (1970), 1236-1239. **5**
- [29] R. P. Stanley, Enumerative Combinatorics, Vol. 1. With a foreword by Gian-Carlo Rota. Corrected reprint of the 1986 original. Cambridge Studies in Advanced Mathematics, 49. Cambridge University Press, Cambridge, 1997. 5
- [30] M. Ward, Arithmetic functions on rings, Ann. of Math., 38 (1937), 725-732. 5
- [31] X.-M. Tang, Biderivations of finite-dimensional complex simple Lie algebras, Linear Multilinear Algebra, **66** (2018), 250-259. **2**
- [32] J. Vukman, Symmetric bi-derivations on prime and semi-prime rings, Aequationes Math., 38 (1989), 245-254. 2
- [33] J. Vukman, Two results concerning symmetric bi-derivations on prime rings, Aequationes Math., 40 (1990), 181-189. 2
- [34] Y. Wang, Biderivations of triangular rings, Linear Multilinear Algebra, 64 (2016), 1952-1959. 2
- [35] J.-H. Zhang and W.-Y. Yu, Jordan derivation of triangular algebras, Linear Algebra Appl., 419 (2006), 251-255. 2
- [36] J.-H. Zhang, S. Feng, H.-X. Li, R.-H. Wu, Generalized biderivations of nest algebras, Linear Algebra Appl., 418 (2006), 225-233. 2
- [37] Y.-X. Zhao, D.-Y. Wang and R.-P. Yao, Biderivations of upper triangular matrix algebras over commu*tative ring*, Int. J. Math. Game Theory Algebra, **18** (2009), 473-478. **2**
- [38] D.-Y. Wang X.-X Yu and Zh.-X. Chen Biderivations of the parabolic subalgebras of simple Lie algebras. Communications in Algebra., **39** (2011), 4097-4104. 2
- [39] D.-Y. Wang X.-X Yu Biderivations And Linear Commuting Maps on the Schrödinger-Virasoro Lie Algebra, Communications in Algebra., **39** (2013), 2166-2173.

SunZhang2017 [40] X.Cheng, M.-J. Wang, J.-C.Sun and H.-L. Zhang *Biderivations and linear commuting maps on the Lie algebra* gca, Communications in Algebra., 65 (2017), 2483-2493. 2

LIANG: SCHOOL OF MATHEMATICS AND STATISTICS, ANHUI UNIVERSITY OF SCIENCE & TECHNOLOGY, 232001, HUAINAN, P.R. CHINA

E-mail address: xfliang@aust.edu.cn

Ren: School of Mathematics and Statistics, AnHui university of science & technology, 232001, Huainan, P.R. China

 $E\text{-}mail\ address:\ \texttt{dandanren0225@163.com}$

WEI: SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, 100081, BEIJING, P. R. CHINA *E-mail address*: daoshuo@hotmail.com, daoshuowei@gmail.com