

Dp-finite fields IV: the rank 2 picture

Will Johnson

March 23, 2020

Abstract

We investigate fields of characteristic 0 and dp-rank 2. While we do not obtain a classification, we prove that any unstable field of characteristic 0 and dp-rank 2 admits a unique definable V-topology. If this statement could be generalized to higher ranks, we would obtain the expected classification of fields of finite dp-rank.

We obtain the unique definable V-topology by investigating the “canonical topology” defined in [8]. Contrary to the expectations of [9], the canonical topology need not be a V-topology. However, we are able to characterize the canonical topology (on fields of dp-rank 2 and characteristic 0) in terms of differential valued fields.

This differential valued structure is obtained through a partial classification of *2-inflat*ors, a sort of “generalized valuation” that arises naturally in fields of finite rank.

Additionally, we give an example of a dp-rank 2 expansion of ACVF with a definable set of full rank and empty interior. This example interferes with certain strategies for proving the henselianity conjecture.

1 Introduction

NIP structures play a central role in modern model theory, and so it would be desirable to classify the NIP theories of fields. NIP can be characterized via dp-rank: a structure M is NIP iff $\text{dp-rk}(M) < \infty$. For an overview of NIP and dp-rank, see ([12], Chapters 2 and 4).

From the point of view of dp-rank, the natural first step is fields of dp-rank 1 (dp-minimal fields). Dp-minimal fields were successfully classified in ([7], Chapter 9). The hope is to generalize this proof to the next simplest case—fields of finite dp-rank (dp-finite fields).

1.1 The story so far

The present paper continues [8], [9], [10], which made some partial progress on the classification of dp-finite fields. The overall strategy is to prove the dp-finite case of the Shelah conjecture:

Conjecture 1.1 (Shelah conjecture, dp-finite case). *Let K be a dp-finite field. Then one of the following holds:*

- K is finite
- K is algebraically closed
- K is real closed
- K admits a non-trivial henselian valuation.

Modulo this conjecture, the classification of dp-finite fields is known ([3], Theorem 3.11¹).

Let $(K, +, \cdot, \dots)$ be a dp-finite field, possibly with extra structure. If K is stable, then K must be algebraically closed or finite ([5], Proposition 7.2). Assume K is unstable. In [8] and [9], we constructed a field topology on K characterized by the fact that the following family is a neighborhood basis of 0:

$$\{X - X : X \subseteq K, X \text{ is definable, } \text{dp-rk}(X) = \text{dp-rk}(K)\}.$$

Here $X - Y$ denotes the set of differences

$$\{x - y : x \in X, y \in Y\},$$

rather than the set difference $X \setminus Y$.

We call this topology the *canonical topology*. In a monster model $\mathbb{K} \succeq K$, define the *K -infinitesimals* to be the intersection of all K -definable basic neighborhoods:

$$J_K = \bigcap \{X - X : X \subseteq \mathbb{K}, X \text{ is } K\text{-definable, } \text{dp-rk}(X) = \text{dp-rk}(K)\}.$$

Using the infinitesimals, we proved the Shelah conjecture for dp-finite fields of positive characteristic in ([8], Corollary 11.4).

In [9], we sketched a strategy for attacking fields of characteristic 0. Say that K is *valuation type* if the canonical topology is a V-topology. (See [11] for a reference on topological fields and V-topologies.) We conjectured

Conjecture 1.2 (Valuation conjecture). *If K is an unstable dp-finite field, then K is valuation type.*

Modulo this conjecture, we proved the Shelah conjecture. We also gave a seemingly weaker criterion which implies the valuation conjecture:

Fact 1.3 (Theorem 8.11 in [9]). *If the K -infinitesimals J_K contain a non-zero ideal of a multi-valuation ring on \mathbb{K} , then K is valuation type.*

Here, a *multi-valuation ring on \mathbb{K}* means a finite intersection of valuation rings on \mathbb{K} .

¹The classification in [3] is for strongly dependent fields, but the proof specializes to the case of dp-finite fields. The fields appearing in the conjectured classification of strongly dependent fields are all dp-finite ([3], Proposition 3.9).

1.2 Main results for dp-finite fields

In the present paper, we investigate unstable fields of dp-rank 2 and characteristic 0. We find a counterexample to the valuation conjecture:

Theorem 1.4. *There is a valued field (K, \mathcal{O}) and a subset $R \subseteq K$ such that*

- *The structure $(K, +, \cdot, \mathcal{O}, R)$ has dp-rank 2.*
- *The set R has full rank $\text{dp-rk}(R) = 2$, but has empty interior with respect to the valuation topology.*
- *The canonical topology is not a V-topology.*

The counterexample does not contradict the Shelah conjecture, or the expected classification of dp-finite fields and valued fields. In fact, $(K, \mathcal{O}) \models \text{ACVF}_{0,0}$.

In spite of the counterexample, we are able to prove the following, statement, which would imply the Shelah conjecture if generalized to higher ranks:

Theorem 1.5. *Let K be an unstable field of characteristic 0 and dp-rank 2.*

- *The canonical topology on K is definable, i.e., there is a uniformly definable basis of opens.*
- *There is a unique definable non-trivial V-topology on K .*

Additionally, we can give a rather explicit description of the canonical topology, in the cases where it is not a V-topology.

Theorem 1.6. *Let \mathbb{K} be a sufficiently resplendent unstable field of characteristic 0 and dp-rank 2. Suppose the canonical topology on \mathbb{K} is not a V-topology. Then there exists a valuation $\text{val} : \mathbb{K} \rightarrow \Gamma$ and a derivation $\delta : \mathbb{K} \rightarrow \mathbb{K}$ such that the following sets form a basis for the canonical topology:*

$$B_{a,b,\gamma} := \{x \in \mathbb{K} : \text{val}(x - a) > \gamma \text{ and } \text{val}(\delta x - b) > \gamma\}.$$

Moreover, every $B_{a,b,\gamma}$ is non-empty.

Non-emptiness of the $B_{a,b,\gamma}$ expresses some degree of independence between the derivation and the valuation. We call this sort of field topology a *DV-topology*. We investigate DV-topologies in §8.

Although we are primarily interested in the case $\text{char}(K) = 0$, we only use $\text{char}(K) \neq 2$. In fact, we prove the valuation conjecture in odd characteristic:

Theorem 1.7. *Let K be an unstable field of dp-rank 2, with $\text{char}(K) > 2$. Then the canonical topology on K is a V-topology.*

1.3 Inflators

Theorem 1.5 is obtained through an algebraic analysis of 2-inflators. An *inflator* is some sort of “generalized valuation” that occurs naturally when attempting to prove the valuation conjecture. The basic properties of inflators were investigated in [10].

The main point of [10] was that the infinitesimals J_K are “governed” by an r -inflator, for some $r \leq \text{dp-rk}(\mathbb{K})$.

Fact 1.8. *Let \mathbb{K} be a sufficiently saturated unstable dp-finite field. Then there are small models $k_0 \preceq K \preceq \mathbb{K}$ and a malleable k_0 -linear r -inflator ς on \mathbb{K} such that*

- $r \leq \text{dp-rk}(\mathbb{K})$.
- The group J_K of K -infinitesimals is an ideal in the fundamental ring R_ς of ς .
- If ς' is any mutation of ς , then there is a small model $K' \succeq K$ such that $J_{K'}$ is an ideal in $R_{\varsigma'}$.

For definitions of inflators, the fundamental ring, malleability, and mutation, see Definitions 4.1, 5.8, 5.31, and 10.2 (respectively) in [10]. We verify Fact 1.8 in §6.1 below.

An inflator ς is *weakly multi-valuation type* ([10], Definition 5.27) if its fundamental ring R_ς contains a non-zero ideal of a multi-valuation ring. Because of Fact 1.8, we can focus our attention on 2-inflators ς with the following properties:

1. ς is malleable.
2. No mutation of ς is weakly multi-valuation type. Otherwise, some $K' \equiv K$ would have valuation type, implying the same for K .
3. The underlying field K has characteristic 0. The classification of dp-finite fields is already known in positive characteristic.

Sections 2–5 carry out an algebraic analysis of inflators satisfying these assumptions. The original hope was to rule out these “wicked” 2-inflators. Instead, we get a rather explicit algebraic description.

Theorem 1.9. *Let $(K, \mathcal{O}, \mathfrak{m})$ be a valued field of characteristic 0, and k_0 be a small subfield on which the valuation is trivial. Let $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$ be a k_0 -linear derivation. Suppose that for every $x \in K/\mathfrak{m}$, the set*

$$\{y \in \mathcal{O} : \partial y = x\}$$

is dense in \mathcal{O} . Let

$$\begin{aligned} R &= \{x \in \mathcal{O} : \partial x \in \mathcal{O}/\mathfrak{m}\} \\ I &= \{x \in \mathfrak{m} : \partial x \in \mathfrak{m}/\mathfrak{m} = 0\}. \end{aligned}$$

Then R is a subring of K and I is an ideal in R , and the quotient R/I is isomorphic to $k[\varepsilon] = k[\varepsilon]/(\varepsilon^2)$, where k is the residue field of \mathcal{O} . There is a 2-inflator

$$\begin{aligned} \varsigma_n : \text{Sub}_K(K^n) &\rightarrow \text{Sub}_k((R/I)^n) \\ V &\mapsto (V \cap R^n + I^n)/I^n. \end{aligned}$$

We call these *diffevaluation inflators* and study them in §8. Actually, the definition is a little more general, allowing K/\mathfrak{m} to be replaced with a “mock K/\mathfrak{m} .”

Up to mutation, diffevaluation inflators account for all the “wicked” 2-inflators:

Theorem 1.10. *Let K be a field of characteristic 0. Let ς be a malleable 2-inflator on K . Suppose that no mutation of ς is weakly multi-valuation type. Then some mutation of ς is a diffevaluation inflator.*

Using this explicit characterization, we prove Theorem 1.5. The characterization strongly hints at how to build the example in Theorem 1.4; we check the details in §10.

1.4 Notation and conventions

A “field” may contain additional structure beyond the pure field structure. We will use bold \mathbb{K} for sufficiently saturated and resplendent fields.

Unlike [8], [9], and [10], we will write the K -infinitesimals as J_K , not I_K , to avoid conflict with the fundamental ideal I of a 2-inflator.

We will use the following definitions and facts from [10]:

- Directories (Definition 2.1) and the characterization of semisimple directories (Theorem 2.7).
- Inflators and equivalence of inflators (Definitions 4.1, 4.2).
- The basic inflator calculus of §5.1.
- The fundamental ring and ideal, and the generalized residue map (Proposition 5.7, Definition 5.8).
- Tame and wild elements (Lemma 5.22 and Definition 5.23).
- The notions of (weakly) multi-valuation type (Definitions 5.26, 5.27), and the characterization in terms of tame and wild elements (Proposition 5.25).
- Mutation (Theorem 10.1, Definition 10.2), transitivity of mutation (Proposition 10.5), and commutativity of mutation (Remark 10.7).
- Malleability (Definition 5.31), and its preservation under mutation (Proposition 10.13)

Unlike [10], we will use k_0 for the small ground field, rather than K_0 . Our inflators will be k_0 -linear. All rings will be k_0 -algebras, and all fields will extend k_0 .

Over the course of §2–5, we will analyze a 2-inflator $\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_S(M)$, satisfying the following assumptions:

1. The characteristic of K (or its subfield k_0) is not 2.
2. ς is malleable

3. No mutation of ς is weakly multi-valuation type.

For §3–5, we will add the additional assumption

4. ς is isotypic: if we write M as $N \oplus N'$ with N, N' simple, then $N \cong N'$.

We call (1-3) the “Weak Assumptions” and (1-4) the “Strong Assumptions.”

Remark 1.11. Until Lemma 5.20, we will use only the following weaker form of (3): no mutation of ς is multi-valuation type.

During our analysis, we will define a number of sets, rings, and functions. We include the following list as a reference:

- The 2-inflator will be

$$\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_S(M),$$

where M is a semisimple S -module of length 2. Beginning in §3, S will be a skew field k , and following §3.1, k will be commutative.

- $k[\varepsilon]$ will denote the ring of dual numbers $k[\varepsilon]/(\varepsilon^2)$.
- R and I will denote the fundamental ring and ideal of ς . The generalized residue map will be $\widehat{\text{res}} : R \rightarrow \text{End}_S(M)$, or later $\widehat{\text{res}} : R \rightarrow \text{End}_k(M)$. In §3, we will arrange for $M = k[\varepsilon]$, and show that $\widehat{\text{res}}$ factors through

$$k[\varepsilon] \cong \text{End}_{k[\varepsilon]}(k[\varepsilon]) \subseteq \text{End}_k(k[\varepsilon]) = \text{End}_k(M).$$

Beginning in §4, we will therefore view $\widehat{\text{res}}$ as a map

$$\widehat{\text{res}} : R \rightarrow k[\varepsilon].$$

- In §3, \mathcal{A}_0 will denote the image of $\widehat{\text{res}}$ in $\text{End}_k(M)$, and \mathcal{A} will denote the k -algebra generated by \mathcal{A}_0 . In Lemma 3.7 and Proposition 3.11, we will see

$$\mathcal{A}_0 = \mathcal{A} \cong k[\varepsilon].$$

- \mathfrak{p} and Q will denote the sets

$$\begin{aligned} \mathfrak{p} &= \{x \in R : \widehat{\text{res}}(x) \in k\varepsilon\} \\ Q &= \{x \in R : \widehat{\text{res}}(x) \in k\}, \end{aligned}$$

where $k[\varepsilon] = k \oplus k\varepsilon$. Then \mathfrak{p} will be an ideal in R , and Q will be a subring of R .

- \mathcal{O} will denote the integral closure of R . In Corollary 4.5, we will see that \mathcal{O} is a valuation ring. In Proposition 4.9, we will see that the residue field of \mathcal{O} is k . We will let

$$\begin{aligned} \text{val} : K^\times &\rightarrow \Gamma \\ \text{res} : \mathcal{O} &\rightarrow k \end{aligned}$$

denote the valuation and residue map. In Proposition 4.9, we will see that

$$\widehat{\text{res}}(x) = s + t\varepsilon \implies \text{res}(x) = s,$$

for $x \in R$ and $s, t \in k$.

- In §5, we will construct an \mathcal{O} -module D and a Q -linear derivation $\partial : \mathcal{O} \rightarrow D$, as well as a valuation

$$\text{val} : D \rightarrow \Gamma_{\leq 0} \cup \{+\infty\}.$$

(D is essentially K/\mathfrak{m}). Before constructing D and ∂ , we will define a map

$$\text{val}_{\partial} : \mathcal{O} \rightarrow \Gamma_{\leq 0} \cup \{+\infty\}$$

in Definition 5.12. Later, $\text{val}_{\partial}(x)$ will turn out to be $\text{val}(\partial x)$.

2 Reduction to the isotypic case

For the duration of §2-5,

$$\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_S(M)$$

will be a k_0 -linear 2-inflator, and the following Weak Assumptions will be in place:

- The characteristic of K (or its subfield k_0) is not 2.
- ς is malleable ([10], Definition 5.31).
- No mutation of ς is weakly of multi-valuation type ([10], Definition 5.27)

Remark 2.1. If ς satisfies the Weak Assumptions, then so does any mutation ς' , by ([10], Propositions 10.5 and 10.13).

Since M is semisimple of length 2, we can write it as an internal direct sum

$$M = A \oplus B,$$

with A, B simple. We say that ς is *isotypic* if $A \cong B$. This depends only on the isomorphism class of the directory $\text{Dir}_S(M)$.

By Theorem 2.7 in [10], we can assume that we are in one of two cases:

- $S = k$ and $M = k^2$, for some division algebra k over k_0 .
- $S = k_1 \times k_2$ and $M = k_1 \oplus k_2$, for two division algebras k_1, k_2 over k_0 .

The first case is isotypic, and the second case is non-isotypic.

2.1 The degeneracy subspace

For any $\alpha \in K$, let Θ_α denote the line

$$\Theta_\alpha := K \cdot (1, \alpha) = \{(x, \alpha x) : x \in K\}.$$

For any $\varphi \in \text{End}_S(M)$, let Θ_φ denote the graph of φ , i.e.,

$$\Theta_\varphi = \{(x, \varphi(X)) : x \in M\}.$$

Recall from ([10], Definition 5.8) that the *fundamental ring* of ς is the set

$$R = \{\alpha \in K \mid \exists \varphi \in \text{End}_S(M) : \varsigma_2(\Theta_\alpha) = \Theta_\varphi\}.$$

This is a subring of K .

For every $\alpha \in K$, one of two things occurs, by Lemma 5.22 in [10].

- The fundamental ring R contains all but at most two of the numbers

$$\{\alpha\} \cup \left\{ \frac{1}{\alpha - c} : c \in k_0 \right\}. \quad (1)$$

- The fundamental ring R contains none of the numbers in (1).

In the first case, α is called *tame*, and in the second case α is called *wild* ([10], Definition 5.23). The fact that ς does not have multi-valuation type implies that there is at least one wild $\alpha \in K$ ([10], Proposition 5.25).

Lemma 2.2. *There is an S -submodule $A \subseteq M$ of length 1 such that for every wild α ,*

$$\varsigma_2(\Theta_\alpha) = A \oplus A \subseteq M \oplus M.$$

Proof. Suppose α is wild. Then $\alpha \notin R$, so $\varsigma_2(\Theta_\alpha)$ is not the graph of an endomorphism. Counting lengths, this implies $0 \oplus A \subseteq \varsigma_2(\Theta_\alpha)$ for some length-1 submodule $A \subseteq M$. Similarly, $\alpha^{-1} \notin R$ implies that

$$0 \oplus A' \subseteq \varsigma_2(\Theta_{\alpha^{-1}})$$

for some length-1 submodule $A' \subseteq M$. Or equivalently, by permutation invariance,

$$A' \oplus 0 \subseteq \varsigma_2(\Theta_\alpha).$$

As $\varsigma_2(\Theta_\alpha)$ has length 2, we must have

$$\varsigma_2(\Theta_\alpha) = A' \oplus A.$$

We claim that $A = A'$. Suppose otherwise. Then M is an internal direct sum of A and A' . By $GL_2(k_0)$ -equivariance,

$$\begin{aligned} \varsigma_2(\Theta_{1/(\alpha+1)}) &= \varsigma_2(\{(\alpha x + x, x) : x \in K\}) \\ &= \{(y + x, x) : x \in A', y \in A\}. \end{aligned}$$

The latter expression is the graph of an endomorphism, so $1/(\alpha + 1) \in R$ and α is tame, a contradiction.

Thus, for any wild α there is some length-1 submodule $A_\alpha \subseteq M$ such that $\varsigma_2(\Theta_\alpha) = A_\alpha \oplus A_\alpha$. It remains to show that A_α doesn't depend on α . Suppose for the sake of contradiction that α, β are wild and $A_\alpha \neq A_\beta$. Then

$$\begin{aligned}\varsigma_3(\{(x, \alpha x, y) : x, y \in K\}) &= A_\alpha \oplus A_\alpha \oplus M \\ \varsigma_3(\{(w, y, \beta y) : w, y \in K\}) &= M \oplus A_\beta \oplus A_\beta \\ \varsigma_3(\{(x, \alpha x, \alpha \beta x) : x \in K\}) &= A_\alpha \oplus 0 \oplus A_\beta\end{aligned}$$

using Lemma 5.2.1 in [10] and the fact that $A_\alpha \cap A_\beta = 0$. Then

$$\begin{aligned}\varsigma_3(\{(x, \alpha x, \alpha \beta x) : x \in K\}) &= A_\alpha \oplus 0 \oplus A_\beta \\ \varsigma_3(\{(0, y, 0) : y \in K\}) &= 0 \oplus M \oplus 0 \\ \varsigma_3(\{(x, y, \alpha \beta x) : x, y \in K\}) &= A_\alpha \oplus M \oplus A_\beta,\end{aligned}$$

using Lemma 5.2.2 in [10]. But a symmetric argument shows

$$\varsigma_3(\{(x, y, \beta \alpha x) : x, y \in K\}) = A_\beta \oplus M \oplus A_\alpha$$

As $\alpha\beta = \beta\alpha$, it follows that

$$A_\alpha \oplus M \oplus A_\beta = A_\beta \oplus M \oplus A_\alpha,$$

and $A_\alpha = A_\beta$, a contradiction. □

We call $A = A_\alpha$ the *degeneracy subspace* of M .

2.2 Reduction to the isotypic case

Corollary 2.3. *Under the Weak Assumptions, ς has a mutation which is isotypic.*

Proof. Take wild α , and let $A \subseteq M$ be the degeneracy subspace, so that

$$\varsigma_2(K \cdot (1, \alpha)) = A \oplus A.$$

By definition of mutation, the mutation along $K \cdot (1, \alpha)$ is of the form

$$\varsigma' : \text{Dir}_K(K) \rightarrow \text{Dir}_S(M'),$$

where $M' = \varsigma_2(K \cdot (1, \alpha)) = A \oplus A$. The S -module M' is isotypic. □

3 The explicit formula for ς

For the duration of §3-5,

$$\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_S(M)$$

will be a k_0 -linear 2-inflator, and the following Strong Assumptions will be in place:

- $\text{char}(k_0) = \text{char}(K) \neq 2$
- ς is malleable
- No mutation of ς is weakly of multi-valuation type.
- ς is isotypic, i.e., if we write M as a direct sum of two simple S -modules A and B , then $A \cong B$.

Isotypy is the new assumption, not present in the Weak Assumptions of §2. Isotypy implies that

$$\text{Dir}_S(M) \cong \text{Dir}_k(k^2)$$

for some division k_0 -algebra k . Therefore, we may assume $S = k$ and M is a two-dimensional k -vector space.

Remark 3.1. As in Remark 2.1, the Strong Assumptions are preserved under mutations. For isotypy, note that any mutation of ς will have the form

$$\varsigma' : \text{Dir}_K(K) \rightarrow \text{Dir}_k(M')$$

for some k -module M' . The fact that k is a division ring ensures that M' is isotypic.

3.1 k is commutative

Proposition 3.2. *The division ring k is commutative (a field).*

Proof. Changing coordinates on k^2 , we may assume that the degeneracy subspace is $0 \oplus k$. If $\alpha \in K$ is wild, then

$$\varsigma_2(\{(x, \alpha x) : x \in K\}) = (0 \oplus k) \oplus (0 \oplus k). \quad (2)$$

We write elements of $M^n = (k^2)^n$ as tuples $(a_1, b_1; a_2, b_2; \dots; a_n, b_n)$.

Let a, b be two non-commuting elements of k . By malleability, we can find $\alpha, \beta \in K^\times$ such that

$$\begin{aligned} \varsigma_2(\{(x, \alpha x) : x \in K\}) &\supseteq \{(t, 0; ta, 0) : t \in k\} \\ \varsigma_2(\{(x, \beta x) : x \in K\}) &\supseteq \{(t, 0; tb, 0) : t \in k\} \end{aligned}$$

Neither α nor β can be wild, by equation (2). By $GL_2(k_0)$ -equivariance,

$$\begin{aligned}\varsigma_2(\{(x, \alpha x) : x \in K\}) &\supseteq \{(t, 0; ta, 0) : t \in k\} \\ \varsigma_2(\{(x, (\alpha - 1)x) : x \in K\}) &\supseteq \{(t, 0; t(a - 1), 0) : t \in k\} \\ \varsigma_2(\{(x, \alpha^{-1}x) : x \in K\}) &\supseteq \{(t, 0; ta^{-1}, 0) : t \in k\} \\ \varsigma_2(\{(x, (\alpha - 1)^{-1}x) : x \in K\}) &\supseteq \{(t, 0; t(a - 1)^{-1}, 0) : t \in k\}\end{aligned}$$

And one of $\alpha, 1/\alpha, 1/(\alpha - 1)$ is in R . So, replacing a with one of $\{a, a^{-1}, (a - 1)^{-1}\}$, we may assume that $\alpha \in R$. Similarly, we may assume $\beta \in R$. Then

$$\begin{aligned}\varsigma_2(\{(x, \alpha x) : x \in K\}) \\ \varsigma_2(\{(x, \beta x) : x \in K\})\end{aligned}$$

are graphs of endomorphisms $\varphi_A, \varphi_B : k^2 \rightarrow k^2$. Because

$$\begin{aligned}(t, 0; ta, 0) &\in \Gamma_A \\ (t, 0; tb, 0) &\in \Gamma_B\end{aligned}$$

it follows that $\varphi_A(t, 0) = (ta, 0)$ and $\varphi_B(t, 0) = (tb, 0)$. Thus φ_A and φ_B do not commute. But this is impossible, as φ_A, φ_B both lie in the image of the homomorphism $R \rightarrow \text{End}_k(k^2)$, and R is commutative. \square

Thus k is a field extending k_0 .

3.2 The algebra \mathcal{A}

Let \mathcal{A}_0 be the image of R in $\text{End}_k(k^2) = M_2(k)$. This is a commutative k_0 -algebra.

Lemma 3.3. *The algebra \mathcal{A}_0 is not contained in the center of $M_2(k)$, i.e., \mathcal{A}_0 contains a matrix not of the form $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$.*

Proof. Changing coordinates on $M \cong k^2$, we may assume that $0 \oplus k$ is the degeneracy subspace, and so

$$\varsigma_2(\{(x, \alpha x) : x \in K\}) = (0 \oplus k) \oplus (0 \oplus k). \quad (3)$$

for any wild α . By malleability, choose $\alpha \in K$ such that

$$\varsigma_2(\{(x, \alpha x) : x \in K\}) \supseteq \{(t, 0; 0, t) : t \in k\} \ni (1, 0; 0, 1).$$

Then α cannot be wild. By $GL_2(k_0)$ -equivariance, we have

$$\begin{aligned}(1, 0; 0, 1) &\in \varsigma_2(\{(x, \alpha x) : x \in K\}) \\ (0, 1; 1, 0) &\in \varsigma_2(\{(x, \alpha^{-1}x) : x \in K\}) \\ (1, 1; 1, 0) &\in \varsigma_2(\{(x, (\alpha + 1)^{-1}x) : x \in K\}).\end{aligned}$$

By tameness of α , one of the right-hand-sides is the graph of some endomorphism $\varphi \in \text{End}_k(k^2)$. By definition, $\varphi \in \mathcal{A}_0$. But no central matrix $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$ can map $(1, 0)$ to $(0, 1)$ or map $(0, 1)$ to $(1, 0)$ or map $(1, 1)$ to $(1, 0)$. \square

Let \mathcal{A} be the k -subalgebra of $M_2(k)$ generated by \mathcal{A}_0 . It is a commutative k -algebra. Note that $M = k^2$ is naturally an \mathcal{A} -module.

Proposition 3.4. *For any $V \subseteq K^n$, the specialization $\varsigma_n(V)$ is an \mathcal{A} -submodule of M^n .*

Proof. We already know that $\varsigma_n(V)$ is a k -submodule, so it remains to show that $\varsigma_n(V)$ is closed under multiplication by \mathcal{A}_0 . Let a be an element of R , specializing to $\varphi \in \mathcal{A}_0$. Then

$$\begin{aligned} \varsigma_{2n}(\{(\vec{x}, a\vec{x}) : \vec{x} \in K^n\}) &= \{(x_1, \dots, x_n, \varphi x_1, \dots, \varphi x_n) : \vec{x} \in M^n\} \\ \varsigma_{2n}(\{(\vec{x}, \vec{y}) : \vec{x} \in V, \vec{y} \in K^n\}) &= \{(x_1, \dots, x_n, y_1, \dots, y_n) : \vec{x} \in \varsigma_n(V), \vec{y} \in M^n\} \\ \varsigma_{2n}(\{(\vec{x}, \vec{y}) : \vec{x} \in K^n, \vec{y} \in V\}) &= \{(x_1, \dots, x_n, y_1, \dots, y_n) : \vec{x} \in M^n, \vec{y} \in \varsigma_n(V)\} \\ \varsigma_{2n}(\{(\vec{x}, a\vec{x}) : \vec{x} \in V\}) &= \{(x_1, x_2, \dots, x_n, \varphi x_1, \dots, \varphi x_n) : \vec{x} \in \varsigma_n(V)\}. \end{aligned}$$

The first line holds because a specializes to φ , using compatibility with \oplus and permutations. The second and third lines hold by compatibility with \oplus . The fourth line holds by intersecting the first and second lines (using Lemma 5.2.1 in [10]). Now V is a K -submodule of K^n , and hence an R -submodule. Therefore

$$\{(\vec{x}, a\vec{x}) : \vec{x} \in V\} \subseteq \{(\vec{x}, \vec{y}) : \vec{x} \in K^n, \vec{y} \in V\}.$$

As ς_{2n} is order-preserving,

$$\begin{aligned} &\{(x_1, x_2, \dots, x_n, \varphi x_1, \dots, \varphi x_n) : \vec{x} \in \varsigma_n(V)\} \\ &\subseteq \{(x_1, \dots, x_n, y_1, \dots, y_n) : \vec{x} \in M^n, \vec{y} \in \varsigma_n(V)\}, \end{aligned}$$

which implies that $\varsigma_n(V)$ is closed under multiplication by φ . \square

Proposition 3.4 says that the inflator $\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_k(M)$ factors through $\text{Dir}_{\mathcal{A}}(M) \subseteq \text{Dir}_k(M)$.

Corollary 3.5. *The degeneracy subspace $X \subseteq M$ is an \mathcal{A} -submodule of M .*

Proof. If $\alpha \in K$ is wild, then

$$\varsigma_2(K \cdot (1, \alpha)) = X \oplus X,$$

and so $X \oplus X$ is an \mathcal{A} -submodule of $M \oplus M$. This implies X is an \mathcal{A} -submodule of M . \square

Lemma 3.6. *The k -algebra \mathcal{A} is isomorphic to one of the following*

- $k \times k$
- $k[\varepsilon] = k[\varepsilon]/(\varepsilon^2)$.

Moreover, M is a free \mathcal{A} -module of rank 1.

Proof. View \mathcal{A} as a commutative k -subalgebra of $M_2(k)$. Take $\mu \in \mathcal{A}$ non-central. Changing the identification $M \cong k^2$, we may assume we are in one of three cases:

- $\mu = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ for some $a \neq b$ in k .
- $\mu = \begin{pmatrix} a & 0 \\ b & a \end{pmatrix}$ for some $a, b \in k$ with $b \neq 0$.
- $\mu = \begin{pmatrix} 0 & 1 \\ a & b \end{pmatrix}$ for some monic irreducible quadratic polynomial $x^2 - bx - a \in k[x]$.

The degeneracy subspace is a one-dimensional subspace of k^2 , preserved by μ , and so μ has an eigenvector. This rules out the third case.

In the first case, the k -subalgebra generated by μ is

$$\mathcal{A}' = \left\{ \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix} : x, y \in k \right\} \cong k \times k.$$

Then $\mathcal{A} \supseteq \mathcal{A}'$. In particular, \mathcal{A} contains the matrix $\mu' = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, and \mathcal{A} lies in the centralizer of μ' . By inspection, this centralizer is \mathcal{A}' . Thus $\mathcal{A} = \mathcal{A}' \cong k \times k$. Also, the vector $(1, 1)$ freely generates k^2 as an \mathcal{A}' -module.

In the second case, the k -subalgebra generated by μ is

$$\mathcal{A}'' = \left\{ \begin{pmatrix} x & 0 \\ y & x \end{pmatrix} : x, y \in k \right\} \cong k[\varepsilon].$$

Then $\mathcal{A} \supseteq \mathcal{A}''$. In particular, \mathcal{A} contains the matrix $\mu'' = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, and \mathcal{A} lies in the centralizer of μ'' . By inspection, this centralizer is \mathcal{A}'' . Thus $\mathcal{A} = \mathcal{A}'' \cong k[\varepsilon]$. Also, the vector $(1, 0)$ freely generates k^2 as an \mathcal{A}'' -module. \square

Lemma 3.7. *The two algebras \mathcal{A}_0 and \mathcal{A} are equal.*

Proof. Changing M up to isomorphism, we may assume $M = \mathcal{A}$. Let μ be any element of \mathcal{A} ; we will show $\mu \in \mathcal{A}_0$. By malleability, there is a line $L \subseteq K^2$ such that

$$\varsigma_2(L) \supseteq k \cdot (1, \mu).$$

But $\varsigma_2(L)$ is an \mathcal{A} -submodule of M^2 (Proposition 3.4), and the \mathcal{A} -submodule generated by $(1, \mu)$ is

$$\{(x, \mu x) : x \in \mathcal{A}\}.$$

Thus

$$\varsigma_2(L) \supseteq \{(x, \mu x) : x \in \mathcal{A}\}.$$

Both sides have length two as k -modules, and so equality holds. Let a be the slope of L . Evidently, $a \neq \infty$, or else $\varsigma_2(L)$ would be $0 \oplus \mathcal{A}$. So $a \in R$ and a specializes to μ . Therefore $\mu \in \mathcal{A}_0$. \square

In particular, the natural homomorphism

$$R \twoheadrightarrow \mathcal{A}_0 \hookrightarrow \mathcal{A}$$

is surjective. If I is the fundamental ideal (the kernel of $R \rightarrow \text{End}_k(M)$), then R/I is isomorphic to $k \times k$ or to $k[\varepsilon]$.

Proposition 3.8. *Fix any \mathcal{A} -module isomorphism $M \cong \mathcal{A}$. Then the specialization maps $\varsigma_n : \text{Sub}_K(K^n) \rightarrow \text{Sub}_k(\mathcal{A}^n)$ are given by the formula*

$$\varsigma_n(V) = \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : (x_1, \dots, x_n) \in V \cap R^n\}$$

where $\widehat{\text{res}}$ is the generalized residue map $R \twoheadrightarrow R/I \cong \mathcal{A}$.

Proof. First suppose that $(a_1, \dots, a_n) \in R^n \cap V$, and a_i specializes to $b_i \in \mathcal{A}$ for each i . Then the line

$$L = \{(x, a_1x, \dots, a_nx) : x \in K\}$$

is contained in $K \oplus V$, and so

$$\varsigma_{n+1}(L) = \{(x, b_1x, \dots, b_nx) : x \in \mathcal{A}\} \subseteq \varsigma_{n+1}(K \oplus V) = \mathcal{A} \oplus \varsigma_n(V).$$

In particular, $(1, b_1, \dots, b_n) \in \mathcal{A} \oplus \varsigma_n(V)$, and so

$$(\widehat{\text{res}}(a_1), \dots, \widehat{\text{res}}(a_n)) = (b_1, \dots, b_n) \in \varsigma_n(V).$$

We have seen

$$\varsigma_n(V) \supseteq \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : (x_1, \dots, x_n) \in V \cap R^n\}$$

Conversely, suppose that $(b_1, \dots, b_n) \in \varsigma_n(V)$. Then

$$(1, b_1, \dots, b_n) \in \mathcal{A} \oplus \varsigma_n(V) = \varsigma_{n+1}(K \oplus V).$$

By malleability, there is a line $L \subseteq K \oplus V$ such that

$$\varsigma_{n+1}(L) \supseteq k \cdot (1, b_1, \dots, b_n) \ni (1, b_1, \dots, b_n).$$

The left hand side is an \mathcal{A} -module, so

$$\varsigma_{n+1}(L) \supseteq \mathcal{A} \cdot (1, b_1, \dots, b_n).$$

Both sides have length two over k , so equality holds. Now L must be the graph of a K -linear function $K \rightarrow K^n$; otherwise $L \subseteq 0 \oplus K^n$ and $\varsigma_{n+1}(L)$ would be contained in $0 \oplus \mathcal{A}^n$, which is visibly false. Thus

$$L = K \cdot (1, a_1, \dots, a_n)$$

for some $a_i \in K$. So

$$\varsigma_{n+1}(\{(x, a_1x, \dots, a_nx) : x \in K\}) = \{(x, b_1x, \dots, b_nx) : x \in \mathcal{A}\}.$$

Joining this with

$$\varsigma_{n+1}(0 \oplus K^{i-1} \oplus 0 \oplus K^{n-i}) = 0 \oplus \mathcal{A}^{i-1} \oplus 0 \oplus \mathcal{A}^{n-i},$$

we obtain

$$\begin{aligned} \varsigma_{n+1}(\{(x, y_1, \dots, y_{i-1}, a_ix, y_{i+1}, \dots, y_n) : x, y_1, \dots, y_n \in K\}) \\ = \{(x, y_1, \dots, y_{i-1}, b_ix, y_{i+1}, \dots, y_n) : x, y_1, \dots, y_n \in \mathcal{A}\}. \end{aligned}$$

By permutation invariance and \oplus -compatibility, we see

$$\varsigma_2(\{(x, a_ix) : x \in K\}) = \{(x, b_ix) : x \in \mathcal{A}\}.$$

So each a_i is in R , and $\widehat{\text{res}}(a_i) = b_i$. The fact that $L \subseteq K \oplus V$ implies that $\vec{a} \in V$. So we have shown that

$$\varsigma_n(V) \subseteq \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : (x_1, \dots, x_n) \in V \cap R^n\}. \quad \square$$

Corollary 3.9. $K = \text{Frac}(R)$.

Proof. Let α be any element of K^\times . By Proposition 3.8,

$$\varsigma_2(\{(x, \alpha x) : x \in K\}) = \{(\widehat{\text{res}}(x), \widehat{\text{res}}(y)) : x, y \in R, y/x = \alpha\}.$$

The fact that $\varsigma_2(\{(x, \alpha x) : x \in K\}) \neq 0$ implies that there exist non-trivial $x, y \in R$ such that $y/x = \alpha$. \square

3.3 Ruling out the split case

Let I denote the *fundamental ideal* ([10], Definition 5.8), i.e., the kernel of the generalized residue map

$$\widehat{\text{res}} : R \rightarrow \mathcal{A}.$$

Remark 3.10. Every maximal ideal of R comes from a maximal ideal of the artinian ring $R/I \cong \mathcal{A}$. Indeed, this follows from the fact that $I \subseteq \text{Jac}(R)$ ([10], Proposition 5.7.4).

For example, if $\mathcal{A} \cong k \times k$, then R has two maximal ideals.

Proposition 3.11. *The algebra \mathcal{A} is isomorphic to $k[\varepsilon]$.*

Proof. Otherwise, $\mathcal{A} \cong k \times k$. Let $p_1, p_2 : R \rightarrow k$ be the two maps such that

$$\widehat{\text{res}}(x) = (p_1(x), p_2(x)) \in k \times k \cong \mathcal{A}.$$

The two maximal ideals of R are the kernels of p_1 and p_2 .

The degeneracy subspace of ς is some rank 1 submodule of \mathcal{A} , necessarily $k \times 0$ or $0 \times k$. Without loss of generality, it is $0 \times k$.

Take some wild element $a \in K$. By Proposition 3.8 and the definition of the degeneracy subspace,

$$\begin{aligned} \varsigma_2(K \cdot (1, a)) &= \{(\widehat{\text{res}}(x), \widehat{\text{res}}(y)) : x, y \in R, y/x = a\} \\ &= \{(p_1(x), p_2(x); p_1(y), p_2(y)) : x, y \in R, y/x = a\} \\ &= \{(0, t; 0, s) : t, s \in k\}. \end{aligned}$$

Therefore, we can find $x, y, x', y' \in R$ such that

$$\begin{aligned} (p_1(x), p_2(x)) &= (0, 0) \\ (p_1(y), p_2(y)) &= (0, 1) \\ y &= ax \\ (p_1(x'), p_2(x')) &= (0, 1) \\ (p_1(y'), p_2(y')) &= (0, 0) \\ y' &= ax'. \end{aligned}$$

Then $xy' = yx'$, and we obtain a contradiction:

$$1 = p_2(y)p_2(x') = p_2(yx') = p_2(xy') = p_2(x)p_2(y') = 0. \quad \square$$

Corollary 3.12. *The fundamental ring R is a local ring.*

Proof. Its maximal ideals come from $k[\varepsilon]$, which is a local ring. \square

The following corollary will be useful later:

Corollary 3.13. *If $\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_k(k^2)$ satisfies the Strong Assumptions, and $a \in K$ specializes to a matrix $\mu = \begin{pmatrix} b & c \\ d & e \end{pmatrix}$, then μ must have a repeated eigenvalue, and so*

$$(b + e)^2 = (\text{Tr}(\mu))^2 = 4 \det(\mu) = 4(be - cd).$$

Proof. The element a lies in R and its image in \mathcal{A} is μ . If μ is central, the identity is clear. Otherwise, the proof of Lemma 3.6 shows that μ must be conjugate to a matrix of the form $\begin{pmatrix} x & y \\ 0 & x \end{pmatrix}$ or $\begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix}$. In the second case, $\mathcal{A} \cong k \times k$, contradicting Proposition 3.11. So we may assume $\mu = \begin{pmatrix} x & y \\ 0 & x \end{pmatrix}$, and then the desired identity is clear. \square

Now that we have identified \mathcal{A} , we can specify the degeneracy locus:

Lemma 3.14. *Under any isomorphism of \mathcal{A} -modules $M \cong \mathcal{A}$, the degeneracy subspace is the principal ideal $k\varepsilon = (\varepsilon) \triangleleft k[\varepsilon] \cong \mathcal{A}$.*

Proof. There are only three $k[\varepsilon]$ -submodules of $k[\varepsilon]$, and $k\varepsilon$ is the only one having dimension 1 over k . \square

We summarize the picture in the following Theorem.

Theorem 3.15. *Let ς be an isotypic malleable k_0 -linear 2-inflator on a field K with $\text{char}(K) \neq 2$. Suppose that no mutation of ς is weakly of multi-valuation type. Then there are*

- A field k extending k_0 .
- A subring $R \subseteq K$.
- An ideal $I \triangleleft R$
- An isomorphism of k_0 -algebras $R/I \cong k[\varepsilon] := k[\varepsilon]/(\varepsilon^2)$

such that ς is isomorphic to

$$\begin{aligned} \varsigma &: \text{Dir}_K(K) \rightarrow \text{Dir}_k(k[\varepsilon]) \\ \varsigma_n(V) &= \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : \vec{x} \in V \cap R^n\} \end{aligned}$$

where $\widehat{\text{res}}$ is the quotient map

$$R \twoheadrightarrow R/I \cong k[\varepsilon].$$

Moreover,

- R is the fundamental ring of ς , I is the fundamental ideal, and $\widehat{\text{res}}$ is the generalized residue map (in the sense of Definition 5.8 in [10]).
- R is a local ring, whose unique maximal ideal \mathfrak{m} is the pullback of $k \cdot \varepsilon$ along $\widehat{\text{res}}$.
- $\text{Frac}(R) = K$.

4 The associated valuation

Continue the Strong Assumptions of §3. In light of Theorem 3.15, we assume that ς has the form

$$\begin{aligned} \varsigma &: \text{Dir}_K(K) \rightarrow \text{Dir}_k(k[\varepsilon]) \\ \varsigma_n &: \text{Sub}_K(K^n) \rightarrow \text{Sub}_k(k[\varepsilon]^n) \\ V &\mapsto \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : \vec{x} \in V \cap R^n\} \end{aligned}$$

where $\widehat{\text{res}} : R \twoheadrightarrow k[\varepsilon]$ is the generalized residue map. The fundamental ideal $I \triangleleft R$ is the kernel of $\widehat{\text{res}}$.

4.1 Finitely generated ideals

Let Q be the set of $a \in R$ such that $\widehat{\text{res}}(a)$ lies in k , i.e.,

$$\widehat{\text{res}}(a) = x + 0\varepsilon$$

for some $x \in k$. The set Q is a k_0 -subalgebra of R . Note that I is an ideal in Q . Moreover, if $a \in Q \setminus I$, then

$$\widehat{\text{res}}(a) = x + 0\varepsilon$$

for some non-zero x , and so $a \in R^\times$. Then

$$\widehat{\text{res}}(a^{-1}) = (x + 0\varepsilon)^{-1} = x^{-1} + 0\varepsilon,$$

so that $a^{-1} \in Q$. Thus Q is a local ring and I is its maximal ideal.

Lemma 4.1. *Let a, b, c be three elements of K . Then there exist $x, y, z \in Q$ such that $ax + by + cz = 0$, and at least one of x, y, z is 1. The same holds for R instead of Q .*

Proof. Consider the vector space $V = \{(x, y, z) \in K^3 : ax + by + cz = 0\}$. Then

$$\varsigma_3(V) = \{(\widehat{\text{res}}(x), \widehat{\text{res}}(y), \widehat{\text{res}}(z)) : (x, y, z) \in R^3 \text{ and } ax + by + cz = 0\}.$$

Also,

$$\dim_k(\varsigma_3(V)) = 2 \cdot \dim_K(V) = 4.$$

Counting dimensions, $\varsigma_3(V)$ must have non-trivial intersection with the subspace

$$k^3 = \{(s + 0\varepsilon, t + 0\varepsilon, u + 0\varepsilon) : s, t, u \in k^3\} \subseteq k[\varepsilon]^3.$$

Therefore, there exist $x, y, z \in R$ such that

$$\widehat{\text{res}}(x) = s + 0\varepsilon$$

$$\widehat{\text{res}}(y) = t + 0\varepsilon$$

$$\widehat{\text{res}}(z) = u + 0\varepsilon$$

for some $s, t, u \in k$, not all zero. Then $x, y, z \in Q$ and at least one of the three is in Q^\times . If $x \in Q^\times$, we may replace x, y, z with $x/x, y/x, z/x$, and arrange for $x = 1$. This handles the case of Q , and the case of R follows as $R \supseteq Q$. \square

We shall return to the ring Q in §5.2.

Corollary 4.2. *Any finitely-generated R -submodule of K is generated by at most two elements. In particular, any ideal of R is generated by at most two elements.*

Proof. It suffices to consider the case of three generators: $R \cdot a + R \cdot b + R \cdot c \leq K$. Then the lemma implies that one of a, b, c is in the R -submodule generated by the other two. \square

4.2 The integral closure of R

We let \mathfrak{p} denote the unique maximal ideal of R , i.e., the set of $x \in R$ such that $\widehat{\text{res}}(x)$ has the form $0 + t\varepsilon$ for some $t \in k$. Because R is local, $R^\times = R \setminus \mathfrak{p}$. Note that \mathfrak{p} is the pullback of the principal ideal $(\varepsilon) = k\varepsilon \triangleleft k[\varepsilon]$ along $\widehat{\text{res}}(-)$.

Lemma 4.3. *Suppose $\alpha \in K$ satisfies a monic quadratic equation over R :*

$$\alpha^2 + b\alpha + c = 0$$

for some $b, c \in R$. If α is wild, then $b^2 - 4c \in \mathfrak{p}$.

Proof. Let $b_0, b_1, c_0, c_1 \in k$ be such that

$$\begin{aligned}\widehat{\text{res}}(b) &= b_0 + b_1\varepsilon \\ \widehat{\text{res}}(c) &= c_0 + c_1\varepsilon.\end{aligned}$$

We must show $b_0^2 = 4c_0$. Let L be the line $K \cdot (1, \alpha)$. By Lemma 3.14,

$$\varsigma_2(L) = \varsigma_2(\{(x, \alpha x) : x \in K\}) = k\varepsilon \oplus k\varepsilon = \{(s\varepsilon, t\varepsilon) : s, t \in k\}.$$

It follows that

$$\begin{aligned}\varsigma_4(\{(x, \alpha x, -cx, -b\alpha x) : x \in K\}) &= \{(s\varepsilon, t\varepsilon, -c_0s\varepsilon, -b_0t\varepsilon) : s, t \in k\} \\ \varsigma_3(\{(x, \alpha x, -b\alpha x - cx) : x \in K\}) &= \{(s\varepsilon, t\varepsilon, (-b_0t - c_0s)\varepsilon) : s, t \in k\} \\ \varsigma_4(\{(x, \alpha x, \alpha x, \alpha^2 x) : x \in K\}) &= \{(s\varepsilon, t\varepsilon, t\varepsilon, (-b_0t - c_0s)\varepsilon) : s, t \in k\}\end{aligned}$$

Here, we are using the identities

$$\begin{aligned}\widehat{\text{res}}(b) \cdot (s\varepsilon) &= (b_0s)\varepsilon \\ \widehat{\text{res}}(c) \cdot (s\varepsilon) &= (c_0s)\varepsilon \\ \alpha^2 &= -b\alpha - c.\end{aligned}$$

Now let $\zeta' : \text{Dir}_K(K) \rightarrow \text{Dir}_k(M')$ be the mutation along L . Then

$$M' = \varsigma_2(L) = \{(s\varepsilon, t\varepsilon) : s, t \in k\} \cong k^2,$$

and

$$\zeta'_2(L) = \varsigma_4(\{(x, \alpha x; \alpha x, \alpha^2 x) : x \in K\}) = \{(s, t; t, (-b_0t - c_0s)) : s, t \in k\}.$$

Note that $\zeta'(L)$ is the graph of the k -linear map

$$k^2 \rightarrow k^2 \tag{4}$$

$$(s, t) \mapsto (t, -b_0t - c_0s) \tag{5}$$

Therefore α lies in the fundamental ring R' of ζ' . By Remark 3.1, ζ' continues to satisfy the Strong Assumptions, and then by Corollary 3.13, the linear map (5) must have a repeated eigenvalue. Therefore

$$b_0^2 = \left(\text{Tr} \begin{pmatrix} 0 & 1 \\ -c_0 & -b_0 \end{pmatrix} \right)^2 = 4 \det \begin{pmatrix} 0 & 1 \\ -c_0 & -b_0 \end{pmatrix} = 4c_0. \quad \square$$

Lemma 4.4. *Let α be an element of K^\times . Then α or α^{-1} is integral over R , and in fact one of α or α^{-1} satisfies a monic polynomial equation of degree d , where*

$$d = \begin{cases} 1 & \alpha \text{ tame} \\ 2 & \alpha \text{ wild.} \end{cases}$$

Proof. Let α be an element of K . First suppose α is tame. Then there is some $b \in k_0$ such that the number $\alpha' := 1/(\alpha - b)$ lies in R . Because R is a k_0 -algebra, it contains $b\alpha' + 1$. Because R is a local k_0 -algebra, at least one of α' and $b\alpha' + 1$ is invertible. Therefore at least one of the following lies in R :

$$\alpha = \frac{b\alpha' + 1}{\alpha'}$$

$$1/\alpha = \frac{\alpha'}{b\alpha' + 1}.$$

Next suppose α is wild. By Lemma 4.1, there are $x, y, z \in R$ such that

$$x + y\alpha + z\alpha^2 = 0,$$

and at least one of x, y, z is in R^\times . If z is invertible, then

$$\alpha^2 + (y/z)\alpha + (x/z) = 0,$$

so α is integral over R . Similarly, if x is invertible, then $1/\alpha$ is integral over R . So we may assume y is invertible and x, z are not. Then

$$0 \equiv x \equiv z \not\equiv y \pmod{\mathfrak{p}}.$$

Let $\beta = \frac{1+\alpha}{1-\alpha}$, so that $\alpha = \frac{\beta-1}{\beta+1}$. Then

$$\begin{aligned} x + y\alpha + z\alpha^2 &= 0 \\ (\beta + 1)^2x + (\beta - 1)(\beta + 1)y + (\beta - 1)^2z &= 0 \\ (\beta^2 + 2\beta + 1)x + (\beta^2 - 1)y + (\beta^2 - 2\beta + 1)z &= 0 \\ (x + y + z)\beta^2 + (2x - 2z)\beta + (x + z - y) &= 0 \\ \beta^2 + \frac{2x - 2z}{x + z + y}\beta + \frac{x + z - y}{x + z + y} &= 0 \end{aligned}$$

where in the final line we have used the fact that (R, \mathfrak{p}) is a local ring and

$$x + z + y \equiv y \not\equiv 0 \pmod{\mathfrak{p}}.$$

Note also that

$$\begin{aligned} \frac{2x - 2z}{x + z + y} &\equiv \frac{0}{y} \pmod{\mathfrak{p}} \\ \frac{x + z - y}{x + z + y} &\equiv \frac{-y}{y} \pmod{\mathfrak{p}}. \end{aligned}$$

Thus $\beta^2 + b'\beta + c' = 0$ for some $b', c' \in R$ with $b' \equiv 0$ and $c' \equiv -1$. Because β and α are related by a fractional linear transformation over k_0 , we know that β is wild. As we are not in characteristic 2,

$$(b')^2 = 0 \not\equiv -4 \equiv 4c' \pmod{\mathfrak{p}},$$

contradicting Lemma 4.3. □

Corollary 4.5. *Let \mathcal{O} denote the integral closure of R (in K). Then \mathcal{O} is a valuation ring on K .*

4.3 Tameness and \mathcal{O}

Let \mathfrak{m} denote the maximal ideal of the valuation ring \mathcal{O} .

Lemma 4.6. *The intersection $\mathfrak{m} \cap R$ is exactly the prime ideal \mathfrak{p} .*

Proof. Let α be an element of R . First suppose $\alpha \notin \mathfrak{p}$. Then $\alpha^{-1} \in R \subseteq \mathcal{O}$, so $\alpha \notin \mathfrak{m}$. Conversely, suppose $\alpha \in \mathfrak{p}$ but $\alpha \notin \mathfrak{m}$. Then $\alpha^{-1} \in \mathcal{O}$, so there exist $c_0, c_1, \dots, c_{n-1} \in R$ such that

$$\alpha^{-n} + c_{n-1}\alpha^{1-n} + \dots + c_1\alpha^{-1} + c_0 = 0,$$

or equivalently

$$-1 = c_{n-1}\alpha + c_{n-2}\alpha^2 + \dots + c_1\alpha^{n-1} + c_0\alpha^n.$$

But the right-hand side is in \mathfrak{p} and the left hand side is not, a contradiction. □

Corollary 4.7. *The valuation ring \mathcal{O} is non-trivial.*

Proof. By surjectivity of $\widehat{\text{res}} : R \rightarrow k[\varepsilon]$, we can find $x \in R$ with $\widehat{\text{res}}(x) = \varepsilon$. Then $x \neq 0$, but $x \in \mathfrak{p}$. Therefore $\mathfrak{m} \neq 0$ and $\mathcal{O} \neq K$. □

Lemma 4.8. *If $\alpha \in \mathcal{O}$, then α is tame if and only if $\alpha \in R$.*

Proof. If $\alpha \in R$ then α is tame by definition. Conversely, suppose α is tame. By Lemma 4.4, one of α or $1/\alpha$ is in R . If $\alpha \in R$, we are done. Otherwise, $1/\alpha \in R$ and $\alpha \notin R$, so $1/\alpha$ is a non-invertible element of R . Then $1/\alpha \in \mathfrak{p} \subseteq \mathfrak{m}$, so α has negative valuation, contradicting the assumption that $\alpha \in \mathcal{O}$. □

4.4 The residue map

Proposition 4.9. *The induced map*

$$k \cong R/\mathfrak{p} = R/(R \cap \mathfrak{m}) \hookrightarrow \mathcal{O}/\mathfrak{m}$$

is onto, hence an isomorphism. In particular, \mathcal{O} has residue field isomorphic to k .

Proof. We must show that for every $x \in \mathcal{O}$ there exists $y \in R$ such that

$$x \equiv y \pmod{\mathfrak{m}}.$$

If $x \in R$ we can take $y = x$, so we may assume $x \notin R$. Then x is wild by Lemma 4.8. If $x \in \mathfrak{m}$ we can take $y = 0$. So we may assume that $x^{-1} \in \mathcal{O}$. By Lemma 4.4, at least one of x or x^{-1} satisfies a monic quadratic polynomial equation over R .

First suppose it is x . Then

$$x^2 + c_1x + c_0 = 0 \tag{6}$$

for some $c_0, c_1 \in R$. By Lemma 4.3,

$$c_1^2 - 4c_0 \in \mathfrak{p}.$$

As we are not in characteristic 2, we may rewrite (6) as

$$\left(x + \frac{c_1}{2}\right)^2 = \left(\frac{c_1^2}{4} - c_0\right).$$

The right hand side is in $\mathfrak{p} \subseteq \mathfrak{m}$, so it has positive valuation. Therefore $x + c_1/2$ also has positive valuation:

$$x + \frac{c_1}{2} \in \mathfrak{m}.$$

Thus we can take $y = -c_1/2$.

Next suppose x^{-1} satisfies a monic quadratic polynomial equation over R :

$$x^{-2} + c_1x^{-1} + c_0 = 0.$$

The same argument shows that $x^{-1} \equiv b \pmod{\mathfrak{m}}$ for some $b \in R$. Then $x \equiv b^{-1} \pmod{\mathfrak{m}}$, and

$$x \notin \mathfrak{m} \implies b \notin \mathfrak{m} \implies b \notin \mathfrak{p} \implies b \in R^\times \implies b^{-1} \in R,$$

so we can take $y = b^{-1}$. □

We let $\text{res} : \mathcal{O} \rightarrow k$ denote the natural residue map. Note that if $a \in R$ and

$$\widehat{\text{res}}(a) = x + y\varepsilon,$$

then $\text{res}(a) = x$.

4.5 The limiting ring

Lemma 4.10. *Let α be an element of K . Suppose $\alpha \notin \mathcal{O}$, and $\alpha^{-1} \notin \mathfrak{p}$. Let ζ' denote the mutation along $K \cdot (1, \alpha^{-1})$, let R' denote the fundamental ring of ζ' , and let \mathfrak{p}' denote the maximal ideal of R' . Then $\alpha^{-1} \in \mathfrak{p}'$.*

Proof. First note that $\alpha^{-1} \in \mathfrak{m} \subseteq \mathcal{O}$. If α^{-1} is tame, then $\alpha^{-1} \in R$ by Lemma 4.8, and then $\alpha^{-1} \in \mathfrak{p}$ by Lemma 4.6. This contradicts the assumptions, and so α^{-1} and α are wild. By Lemma 4.4, one of α and α^{-1} satisfies a monic quadratic polynomial equation over R . Since α does not lie in the integral closure \mathcal{O} of R , it must be α^{-1} that satisfies the equation:

$$\alpha^{-2} = b\alpha^{-1} + c.$$

Then $b^2 + 4c \in \mathfrak{p}$ by Lemma 4.3. We claim that $c \in \mathfrak{p}$. Otherwise, $c \in R^\times$, and

$$c^{-1} = c^{-1}b\alpha + \alpha^2,$$

contradicting the fact that α is not integral over R . So

$$\begin{aligned} b^2 + 4c &\equiv 0 \pmod{\mathfrak{p}} \\ c &\equiv 0 \pmod{\mathfrak{p}} \end{aligned}$$

and therefore $b \in \mathfrak{p}$ as well. Let β, γ be $\widehat{\text{res}}(b)$ and $\widehat{\text{res}}(c)$, respectively. The fact that $b, c \in \mathfrak{p}$ implies that $\beta, \gamma \in k\varepsilon$, and therefore β, γ annihilate $k\varepsilon$.

As α^{-1} is wild and $k\varepsilon$ is the degeneracy subspace,

$$\varsigma_2(\{(x, \alpha^{-1}x) : x \in K\}) = \{(s\varepsilon, t\varepsilon) : s, t \in k\}.$$

By the usual inflator calculus, one sees that

$$\begin{aligned} \varsigma_3(\{(x, \alpha^{-1}x, \alpha^{-2}x) : x \in K\}) &= \varsigma_3(\{(x, \alpha^{-1}x, b\alpha^{-1}x + cx) : x \in K\}) \\ &= \{(s\varepsilon, t\varepsilon, \beta t\varepsilon + \gamma s\varepsilon) : s, t \in k\} \\ &= \{(s\varepsilon, t\varepsilon, 0) : s, t \in k\}. \end{aligned}$$

Now let ζ' be the mutation of σ along $K \cdot (1, \alpha^{-1})$. Then

$$\begin{aligned} \zeta'_2(\{(x, \alpha^{-1}x) : x \in K\}) &= \zeta_4(\{(x, \alpha^{-1}x; \alpha^{-1}x, \alpha^{-2}x) : x \in K\}) \\ &= \{(s, t; t, 0) : s, t \in k\}. \end{aligned}$$

Thus α^{-1} specializes to the endomorphism

$$(s, t) \mapsto (t, 0),$$

and so $\alpha^{-1} \in R'$. This endomorphism fails to be invertible, so $\alpha^{-1} \in \mathfrak{p}'$. □

Lemma 4.11. *Let ζ' be a mutation of ς , with fundamental ring R' . Let \mathfrak{p}' be the maximal ideal of R' . Then $\mathfrak{p} \subseteq \mathfrak{p}'$.*

Proof. Suppose ζ' is the mutation along $L = K \cdot (a_1, \dots, a_n)$. Then ζ' is a map

$$\text{Dir}_K(K) \rightarrow \text{Dir}_k(M'),$$

where $M' = \varsigma_n(L) \subseteq (k[\varepsilon])^n$. By Proposition 3.4, M' is a $k[\varepsilon]$ -submodule of $(k[\varepsilon])^n$.

Take $b \in \mathfrak{p}$. Then $\widehat{\text{res}}_\varsigma(b) = s\varepsilon$ for some $s \in k$. This means that $b \in R$ specializes (with respect to ς) to the endomorphism

$$\begin{aligned} k[\varepsilon] &\rightarrow k[\varepsilon] \\ z &\mapsto s\varepsilon z. \end{aligned}$$

With respect to the mutation ς' , the element b specializes to the endomorphism

$$\begin{aligned} M' &\rightarrow M' \\ \vec{v} &\mapsto s\varepsilon\vec{v}, \end{aligned}$$

by Lemma 10.3 in [10]. This map is not onto, by Nakayama's lemma (over the Noetherian ring $k[\varepsilon]$). Therefore $b \in R'$, but $b^{-1} \notin R'$, implying that $b \in \mathfrak{p}'$. \square

Proposition 4.12. *If ς' is a mutation of ς , with fundamental ring R' , then $R' \subseteq \mathcal{O}$. Consequently, R' has the same integral closure as R .*

Proof. Let b be an element of R' that is not in \mathcal{O} . First suppose that $b^{-1} \in \mathfrak{p}$. Then Lemma 4.11 implies $b^{-1} \in \mathfrak{p}'$. Therefore $b \notin R'$, a contradiction.

Next suppose that $b^{-1} \notin \mathfrak{p}$. Let τ and τ' be the mutations of ς and ς' along $K \cdot (1, b^{-1})$. By Lemma 4.10, $b^{-1} \in \mathfrak{p}_\tau$. By commutativity of mutation (Remark 10.7 in [10]), τ' is a mutation of τ . By Lemma 4.11,

$$b^{-1} \in \mathfrak{p}_\tau \subseteq \mathfrak{p}_{\tau'}.$$

This implies $b \notin R_{\tau'}$. But $b \in R' = R_{\varsigma'} \subseteq R_{\tau'}$, a contradiction. \square

Corollary 4.13. *The integral closure \mathcal{O} is the limiting ring R_∞ of [10], Definition 10.9.*

Proof. The limiting ring R_∞ is integrally closed, so $R_\infty \supseteq \mathcal{O}$. On the other hand R_∞ is a union of rings R' obtained by mutation. Proposition 4.12 says $R' \subseteq \mathcal{O}$. Thus $R_\infty \subseteq \mathcal{O}$. \square

5 Differential structure

Continue the Strong Assumptions of §3-4.

Remark 5.1. Over the next few sections, we will carry out a number of convoluted calculations. The motivated reader may wish to keep two running examples in mind:

- The diffevaluation inflators of §8.6 below.
- The “endless mutation” example of §12.3 in [10].

The second example doesn't actually satisfy the Strong Assumptions, but this won't matter until Lemma 5.20. (See Remark 1.11.)

5.1 Double mutation lemma

The idea of the next few lemmas is that we can calculate the residue $\text{res}(r)$ of an element r by passing to a mutation where r becomes tame; Proposition 4.12 ensures that the valuation does not change in the mutation.

Lemma 5.2. *Let r be an element of K , let q be an element of k , and let L be a line (a one-dimensional subspace) in K^n . Suppose that every element of*

$$\varsigma_{2n}(\{(\vec{x}, r\vec{x}) : \vec{x} \in L\})$$

is of the form $(\vec{x}, q\vec{x})$. Then $\text{val}(r) \geq 0$ and $\text{res}(r) = q$.

Proof. Let $M' = \varsigma_n(L)$. Then

$$\begin{aligned} \varsigma_{2n}(\{(\vec{x}, r\vec{x}) : \vec{x} \in L\}) &\subseteq \{(\vec{x}, q\vec{x}) : \vec{x} \in k[\varepsilon]^n\} \\ \varsigma_{2n}(\{(\vec{x}, r\vec{x}) : \vec{x} \in L\}) &\subseteq \varsigma_{2n}(L \oplus L) = M' \oplus M' \\ \varsigma_{2n}(\{(\vec{x}, r\vec{x}) : \vec{x} \in L\}) &\subseteq \{(\vec{x}, q\vec{x}) : \vec{x} \in M'\}. \end{aligned}$$

The first line is by assumption, the second line is by order-preservation and \oplus -compatibility, and the third line follows by intersecting the first two lines. Counting lengths, equality must hold in the second line. Let

$$\zeta' : \text{Dir}_K(K) \rightarrow \text{Dir}_k(M')$$

be the mutation of ζ along L . Let $R', \mathfrak{p}', I', \mathcal{O}', \mathfrak{m}'$ denote the analogues of $R, \mathfrak{p}, I, \mathcal{O}, \mathfrak{m}$ for the mutation ζ' . Then

$$\zeta'_2(K \cdot (1, r)) = \varsigma_{2n}(\{(\vec{x}, r\vec{x}) : \vec{x} \in L\}) = \{(\vec{x}, q\vec{x}) : \vec{x} \in M'\}.$$

It follows that r specializes with respect to ζ' to the endomorphism

$$\begin{aligned} M' &\rightarrow M' \\ x &\mapsto qx. \end{aligned}$$

Thus $r \in R' \subseteq \mathcal{O}'$. Choose some $p \in R$ such that $\widehat{\text{res}}(p) = q = q + 0\varepsilon$. By Lemma 10.3 in [10], the element p is also in R' , and also specializes to this endomorphism. Therefore

$$r - p \in I' \subseteq \mathfrak{p}' \subseteq \mathfrak{m}'.$$

By Proposition 4.12, $\mathcal{O} = \mathcal{O}'$ and $\mathfrak{m} = \mathfrak{m}'$, implying that $r \in \mathcal{O}$ and $r - p \in \mathfrak{m}$. Therefore $\text{val}(r) \geq 0$ and $\text{res}(r) = \text{res}(p) = q$. \square

Lemma 5.3. *Suppose $a \in K$ has $\text{val}(a) > 0$, and suppose*

$$(u, v) \in \varsigma_2(K \cdot (1, a))$$

for some $u, v \in k[\varepsilon]$ with $v \neq 0$. Then there is $a' \in K$ such that

$$(k \cdot v) \oplus (k \cdot v) = \varsigma_2(K \cdot (1, a')).$$

Proof. If a is wild, then

$$\varsigma_2(K \cdot (1, a)) = k\varepsilon \oplus k\varepsilon$$

by Lemmas 2.2 and 3.14. Then $v \in k\varepsilon$, and

$$(k \cdot v) \oplus (k \cdot v) = k\varepsilon \oplus k\varepsilon.$$

So we may take $a' = a$.

Otherwise, a is tame, and so $a \in R \cap \mathfrak{m} = \mathfrak{p}$ by Lemmas 4.6 and 4.8. Then $\widehat{\text{res}}(a) = b\varepsilon$ for some $b \in k$, and

$$(u, v) \in \varsigma_2(K \cdot (1, a)) = \{(x + y\varepsilon, (b\varepsilon)(x + y\varepsilon)) : x, y \in k\} = \{(x + y\varepsilon, 0 + (xb)\varepsilon) : x, y \in k\}.$$

Thus $v \in k\varepsilon$. Then we can take a' to be any wild element, and

$$(k \cdot v) \oplus (k \cdot v) = k\varepsilon \oplus k\varepsilon = \varsigma_2(K \cdot (1, a')). \quad \square$$

Lemma 5.4 (Double mutation lemma). *Let a, r be elements of K , with $\text{val}(a) > 0$. Suppose that*

$$(s, t, u, qu) \in \varsigma_4(K \cdot (1, r, a, ar))$$

for some $s, t, u \in k[\varepsilon]$ and $q \in k$ with u nonzero. Then $\text{val}(r) \geq 0$ and $\text{res}(r) = q$.

Proof. Let $\zeta' : \text{Dir}_K(K) \rightarrow \text{Dir}_k(M')$ be the mutation of ς along $K \cdot (1, r)$, where $M' = \varsigma_2(M') \subseteq (k[\varepsilon])^2$. Then

$$\zeta'_2(K \cdot (1, a)) = \varsigma_4(K \cdot (1, r, a, ar)) \ni (s, t, u, qu).$$

By Lemma 5.3 applied to ζ' , there is some $a' \in K$ such that

$$\zeta'_2(K \cdot (1, a')) = \{(xu, xqu; yu, yqu) : x, y \in k\}.$$

Equivalently, then

$$\begin{aligned} \varsigma_4(K \cdot (1, r, a', a'r)) &= \{(xu, xqu, yu, yqu) : x, y \in k\} \\ \varsigma_4(K \cdot (1, a', r, a'r)) &= \{(xu, yu, xqu, yqu) : x, y \in k\}. \end{aligned}$$

By Lemma 5.2 applied to the line $L = K \cdot (1, a')$, it follows that $\text{val}(r) \geq 0$ and $\text{res}(r) = q$. \square

5.2 Neutralizers

Recall from §4.1 that Q is the subring

$$Q = \{x \in R : \widehat{\text{res}}(x) \in k\},$$

where we view k as a subset of $k[\varepsilon]$ in the natural way.

Definition 5.5. If $a \in \mathcal{O}$, a *neutralizer* is an $a^\dagger \in Q$ such that $aa^\dagger \in R \setminus Q$.

Neutralizers need not be unique.

Lemma 5.6. *If $a \in \mathcal{O} \setminus Q$, then a has a neutralizer.*

Proof. First suppose $a \in R \setminus Q$. Then 1 is a neutralizer.

Next suppose $a \in \mathcal{O} \setminus R$. By Lemma 4.8, the element a is wild. Then

$$\varsigma_2(K \cdot (1, a)) = k\varepsilon \oplus k\varepsilon.$$

On the other hand, by Proposition 3.8,

$$\varsigma_2(K \cdot (1, a)) = \{(\widehat{\text{res}}(x), \widehat{\text{res}}(ax)) : x \in R, ax \in R\}.$$

Therefore, there is some $a^\dagger \in R$ such that $aa^\dagger \in R$, and

$$\begin{aligned}\widehat{\text{res}}(a^\dagger) &= 0 \\ \widehat{\text{res}}(aa^\dagger) &= \varepsilon.\end{aligned}$$

Then $a^\dagger \in Q$ and $aa^\dagger \in R \setminus Q$. □

Lemma 5.7. *Let a be a wild element.*

- *If $b \in R$ and $ab \in R$, then*

$$\begin{aligned}\widehat{\text{res}}(b) &= p\varepsilon \\ \widehat{\text{res}}(ab) &= q\varepsilon\end{aligned}$$

for some $p, q \in k$.

- *If a^\dagger is a neutralizer of a , then*

$$\begin{aligned}\widehat{\text{res}}(a^\dagger) &= 0 \\ \widehat{\text{res}}(aa^\dagger) &= q\varepsilon.\end{aligned}$$

for some nonzero $q \in k$.

Proof. Because a is wild and $k\varepsilon$ is the degeneracy subspace,

$$\{(\widehat{\text{res}}(x), \widehat{\text{res}}(ax)) : x \in R, ax \in R\} = \varsigma_2(K \cdot (1, a)) = k\varepsilon \oplus k\varepsilon.$$

The fact that $b, ab \in R$ thus implies that $\widehat{\text{res}}(b)$ and $\widehat{\text{res}}(ab)$ lie in $k\varepsilon$, and so

$$\begin{aligned}\widehat{\text{res}}(b) &= 0 + p\varepsilon \\ \widehat{\text{res}}(ab) &= 0 + q\varepsilon,\end{aligned}$$

for some $p, q \in k$. When b is a neutralizer a^\dagger , we must have $p = 0$ and $q \neq 0$, because $a^\dagger \in Q$ and $aa^\dagger \notin Q$. □

Lemma 5.8. *Let a be a wild element with $\text{val}(a) > 0$, and let a^\dagger be a neutralizer of a . Suppose $b \in R$ and $ab \in R$. Then $\text{val}(b) \geq \text{val}(a^\dagger)$. Moreover,*

$$ab \in Q \iff \text{val}(b) > \text{val}(a^\dagger).$$

Proof. By Lemma 5.7,

$$\begin{aligned}\widehat{\text{res}}(a^\dagger) &= 0 \\ \widehat{\text{res}}(b) &= p\varepsilon \\ \widehat{\text{res}}(aa^\dagger) &= s\varepsilon \\ \widehat{\text{res}}(ab) &= q\varepsilon,\end{aligned}$$

for some $p, q, s \in k$ with $s \neq 0$. Note $ab \in Q \iff q = 0$. It suffices to show that $\text{val}(b/a^\dagger) \geq 0$ and $\text{res}(b/a^\dagger) = q/s$.

By the inflator calculus,

$$\begin{aligned}\varsigma_5(K \cdot (1, a^\dagger, b, aa^\dagger, ab)) &= \varsigma_5(\{(x, a^\dagger x, bx, aa^\dagger x, abx) : x \in K\}) \\ &= \{(x, 0, (p\varepsilon)x, (s\varepsilon)x, (q\varepsilon)y) : x \in k[\varepsilon]\} \\ &= (k[\varepsilon]) \cdot (1, 0, p\varepsilon, s\varepsilon, q\varepsilon),\end{aligned}$$

and therefore

$$(0, p\varepsilon, s\varepsilon, q\varepsilon) \in \varsigma_4(K \cdot (a^\dagger, b, aa^\dagger, ab)) = \varsigma_4(K \cdot (1, b/a^\dagger, a, ab/a^\dagger)).$$

By the Double Mutation Lemma 5.4 with $r = b/a^\dagger$, it follows that $\text{val}(b/a^\dagger) \geq 0$ and $\text{res}(b/a^\dagger) = q/s$. \square

Lemma 5.9. *Let a be a wild element with $\text{val}(a) > 0$, and let a^\dagger be a neutralizer. Suppose $b \in R$ and $ab \notin R$. Then $\text{val}(b) < \text{val}(a^\dagger)$.*

Proof. Note that $b \in R \subseteq \mathcal{O}$ and $a \in \mathfrak{m} \subseteq \mathcal{O}$, so $ab \in \mathcal{O}$. The fact that $ab \notin R$ then implies that ab is wild, by Lemma 4.8. Also, $ab \notin Q$, as $Q \subseteq R$. By Lemma 5.6, ab has a neutralizer $(ab)^\dagger$. By Lemma 5.7,

$$\begin{aligned}\widehat{\text{res}}((ab)^\dagger) &= 0 \\ \widehat{\text{res}}(ab(ab)^\dagger) &= s\varepsilon,\end{aligned}$$

for some non-zero $s \in k$. Let $c = b(ab)^\dagger$. Then $c \in R$ (because $b \in R$ and $(ab)^\dagger \in Q \subseteq R$). Also

$$ac = (ab)(ab)^\dagger \in R \setminus Q.$$

By Lemma 5.8,

$$ac \notin Q \implies \text{val}(c) = \text{val}(a^\dagger).$$

Then

$$\text{val}(a^\dagger) = \text{val}(c) = \text{val}(b) + \text{val}((ab)^\dagger) > \text{val}(b),$$

because $\text{res}((ab)^\dagger) = 0$. \square

Lemmas 5.8 and 5.9 combine to yield the following:

Lemma 5.10. *Let a be a wild element with $\text{val}(a) > 0$, and let a^\dagger be a neutralizer of a . Suppose $b \in R$.*

$$\begin{aligned} ab \in R &\iff \text{val}(b) \geq \text{val}(a^\dagger) \\ ab \in Q &\iff \text{val}(b) > \text{val}(a^\dagger). \end{aligned}$$

Next, we weaken the assumption on a , allowing $\text{val}(a) = 0$:

Lemma 5.11. *Let a be a wild element with $\text{val}(a) \geq 0$, and let a^\dagger be a neutralizer of a . Suppose $b \in R$. Then*

$$ab \in R \iff \text{val}(b) \geq \text{val}(a^\dagger).$$

If $b \in Q \subseteq R$, then

$$ab \in Q \iff \text{val}(b) > \text{val}(a^\dagger).$$

Proof. Let $\text{res}(a) = \gamma$, and choose $c \in R$ with $\widehat{\text{res}}(c) = \gamma + 0\varepsilon$. Then $c \in Q$ and $\text{res}(c) = \gamma$. Let $a' = a - c$. Then $\text{res}(a') = \text{res}(a) - \text{res}(c) = \gamma - \gamma = 0$. So $\text{val}(a') > 0$. Because $a \notin R$ and $c \in R$, we have $a' = a - c \notin R$, and so a' is wild by Lemma 4.8. Moreover,

$$a'a^\dagger = aa^\dagger - ca^\dagger \in (R \setminus Q) - Q = R \setminus Q.$$

Therefore a^\dagger is a neutralizer of a' , and we can apply Lemma 5.10 to a', a^\dagger, b . Then

$$ab \in R \iff a'b \in R \iff \text{val}(b) \geq \text{val}(a^\dagger),$$

since $ab - a'b = cb \in R$. If $b \in Q$, then $cb \in Q$ and

$$ab \in Q \iff a'b \in Q \iff \text{val}(b) > \text{val}(a^\dagger). \quad \square$$

5.3 The secondary valuation

Definition 5.12. For $a \in \mathcal{O}$, let $\text{val}_\partial(a)$ denote

$$\text{val}_\partial(a) = \begin{cases} +\infty & \text{if } a \in Q \\ -\text{val}(a^\dagger) & \text{if } a \text{ has a neutralizer } a^\dagger \end{cases}$$

Lemma 5.13. $\text{val}_\partial(a)$ is well-defined: for any $a \in \mathcal{O}$, exactly one of the following holds

1. $a \in Q$
2. a has a neutralizer a^\dagger ,

and in case (2) the valuation $\text{val}(a^\dagger)$ is independent of the choice of a neutralizer a^\dagger .

Proof. If $a \notin Q$, then a neutralizer a^\dagger exists by Lemma 5.6. Conversely, if a neutralizer a^\dagger exists, then $a^\dagger \in Q$ and $aa^\dagger \in R \setminus Q$. As Q is a subring, $a \notin Q$.

Now suppose a^\dagger and b are two neutralizers of a . If a is wild, then Lemma 5.11 applies, so

$$ab \in R \setminus Q \implies \text{val}(b) = \text{val}(a^\dagger).$$

If a is tame, we can write

$$\begin{aligned}\widehat{\text{res}}(a) &= x + y\varepsilon \\ \widehat{\text{res}}(a^\dagger) &= z + 0\varepsilon \\ \widehat{\text{res}}(b) &= w + 0\varepsilon\end{aligned}$$

for some $x, y, z, w \in k$, using the fact that $a^\dagger, b \in Q$. Then

$$\begin{aligned}\widehat{\text{res}}(aa^\dagger) &= xz + yz\varepsilon \\ \widehat{\text{res}}(ab) &= xw + yw\varepsilon,\end{aligned}$$

The fact that $aa^\dagger, ab \in R \setminus Q$ implies that yz and yw are non-zero. Therefore z, w are non-zero and $\text{res}(a^\dagger)$ and $\text{res}(b)$ are non-zero, implying $\text{val}(a^\dagger) = 0 = \text{val}(b)$. \square

Lemma 5.14. *For $a \in \mathcal{O}$,*

- $\text{val}_\partial(a) \geq 0 \iff a \in R$
- $\text{val}_\partial(a) > 0 \iff a \in Q$.

Proof. First suppose $a \in Q$. Then $\text{val}_\partial(a) = +\infty$ by definition. Next suppose that $a \in R \setminus Q$. Then 1 is a neutralizer of a , so $\text{val}_\partial(a) = -\text{val}(1) = 0$.

Lastly, suppose $a \in \mathcal{O} \setminus R$. Let a^\dagger be a neutralizer. By Lemma 5.7, $\widehat{\text{res}}(a^\dagger) = 0$, implying $\text{res}(a^\dagger) = 0$, $\text{val}(a^\dagger) > 0$, and $\text{val}_\partial(a) < 0$. \square

Lemma 5.15. *If $a \in \mathcal{O}$ and $b \in Q$, then*

$$\text{val}_\partial(ab) = \text{val}_\partial(a) + \text{val}(b)$$

unless the right hand side is positive, in which case

$$\text{val}_\partial(ab) = +\infty.$$

Proof. First suppose $a \in Q$. Then $\text{val}_\partial(a) = +\infty$, and the conclusion says that $ab \in Q$, which is true (Q is a ring).

Next, suppose that $a \in R \setminus Q$. Then $\text{val}_\partial(a) = 0$. As, $a, ab \in R$ and $b \in Q$, we may write

$$\begin{aligned}\widehat{\text{res}}(a) &= x + y\varepsilon \\ \widehat{\text{res}}(b) &= z + 0\varepsilon \\ \widehat{\text{res}}(ab) &= xz + yz\varepsilon.\end{aligned}$$

for some $x, y, z \in k$, with $y \neq 0$. Then

$$\begin{aligned} \text{val}(b) > 0 &\implies z = \text{res}(b) = 0 \implies yz = 0 \implies ab \in Q \implies \text{val}_\partial(ab) = +\infty \\ \text{val}(b) = 0 &\implies z = \text{res}(b) \neq 0 \implies yz \neq 0 \implies ab \in R \setminus Q \implies \text{val}_\partial(ab) = 0. \end{aligned}$$

Lastly, suppose $a \in \mathcal{O} \setminus R$, so a is wild. Take a neutralizer a^\dagger . We break into cases according to the sign of $\text{val}_\partial(a) + \text{val}(b)$.

- If $\text{val}_\partial(a) + \text{val}(b) > 0$, then $\text{val}(b) > \text{val}(a^\dagger)$, so $ab \in Q$ by Lemma 5.11. Thus $\text{val}_\partial(ab) = +\infty$.
- If $\text{val}_\partial(a) + \text{val}(b) = 0$, then $\text{val}(b) = \text{val}(a^\dagger)$, so $ab \in R \setminus Q$, by Lemma 5.11. Thus $\text{val}_\partial(ab) = 0$.
- If $\text{val}_\partial(a) + \text{val}(b) < 0$, then $\text{val}(b) < \text{val}(a^\dagger)$, and $ab \notin R$ by Lemma 5.11. On the other hand, $a, b \in \mathcal{O}$, so $ab \in \mathcal{O} \setminus R$, and ab is wild (Lemma 4.8). Take a neutralizer $(ab)^\dagger$, and let $c = b(ab)^\dagger$. Then $b, (ab)^\dagger \in Q$, so $c \in Q$. Also $ac = (ab)(ab)^\dagger \in R \setminus Q$, and so c is a neutralizer of a . Then

$$\text{val}_\partial(ab) = -\text{val}((ab)^\dagger) = -\text{val}(c) + \text{val}(b) = \text{val}_\partial(a) + \text{val}(b). \quad \square$$

The next lemma says that for any γ , the set

$$\{x \in \mathcal{O} : \text{val}_\partial(x) \geq \gamma\}$$

is a subring of \mathcal{O} .

Lemma 5.16. *For any $a, b \in \mathcal{O}$, let $\gamma = \min(\text{val}_\partial(a), \text{val}_\partial(b))$. Then*

$$\begin{aligned} \text{val}_\partial(a + b) &\geq \gamma \\ \text{val}_\partial(ab) &\geq \gamma. \end{aligned}$$

Proof. If $\gamma > 0$ (i.e., $\gamma = +\infty$), this holds because Q is a ring. If $\gamma = 0$, this holds because R is a ring. So we may assume

$$0 > \gamma = \text{val}_\partial(a) \leq \text{val}_\partial(b),$$

swapping a and b if necessary. The fact that $\text{val}_\partial(a) < 0$ implies a is wild. Take a neutralizer a^\dagger of a . Then $a^\dagger \in Q$, so by Lemma 5.15,

$$\text{val}_\partial(a^\dagger b) \geq \text{val}(a^\dagger) + \text{val}_\partial(b) = \text{val}_\partial(b) - \text{val}_\partial(a) \geq 0,$$

so $a^\dagger b \in R$ by Lemma 5.14. Then

$$a^\dagger(a + b) = aa^\dagger + a^\dagger b \in R + R = R,$$

so

$$\text{val}_\partial(a^\dagger(a+b)) \geq 0.$$

By Lemma 5.15, it follows that

$$\text{val}_\partial(a+b) + \text{val}(a^\dagger) \geq 0,$$

or equivalently, that

$$\text{val}_\partial(a+b) \geq -\text{val}(a^\dagger) = \text{val}_\partial(a) = \gamma.$$

Also, Lemma 5.11 shows that

$$(a^\dagger b \in R \text{ and } \text{val}(a^\dagger b) \geq \text{val}(a^\dagger)) \implies aa^\dagger b \in R,$$

as a is wild. Thus $a^\dagger(ab) \in R$, and $\text{val}_\partial(a^\dagger(ab)) \geq 0$. As in the case of $a+b$, this implies that

$$\text{val}_\partial(ab) + \text{val}(a^\dagger) \geq 0,$$

or equivalently, that $\text{val}_\partial(ab) \geq \gamma$. □

Later (Corollary 5.26), we will get an improved rule for $\text{val}_\partial(ab)$, but for now we content ourselves with the following cases:

Lemma 5.17. *If $a \in \mathcal{O}$ and $\text{val}(a) + \text{val}_\partial(a) > 0$, then $a^2 \in R$, i.e., $\text{val}_\partial(a^2) \geq 0$.*

Proof. We may assume $a \notin R$, so a is wild. Take a neutralizer a^\dagger . Then

$$0 < \text{val}(a) + \text{val}_\partial(a) = \text{val}(a) - \text{val}(a^\dagger),$$

and so $\text{val}(a) > \text{val}(a^\dagger)$. Therefore a^\dagger/a is not integral over R . By Lemma 4.4, the inverse a/a^\dagger satisfies a monic quadratic polynomial equation over R . Therefore

$$a^2 = baa^\dagger + c(a^\dagger)^2$$

for some $b, c \in R$. As $aa^\dagger, b, c, a^\dagger$ are all in R , this implies $a^2 \in R$. □

Lemma 5.18. *Let γ be a positive element of the valuation group, and a, b be elements of \mathcal{O} . Suppose*

$$\begin{aligned} \text{val}(a) &> \gamma \\ \text{val}(b) &> \gamma \\ \text{val}_\partial(a) &> -\gamma \\ \text{val}_\partial(b) &> -\gamma. \end{aligned}$$

Then $ab \in R$, i.e., $\text{val}_\partial(ab) \geq 0$.

Proof. Note that $\text{val}(a) + \text{val}_\partial(a) > 0$ and $\text{val}(b) + \text{val}_\partial(b) > 0$, so $a^2, b^2 \in R$ by Lemma 5.17. By Lemma 5.16, $\text{val}_\partial(a+b) > -\gamma$, and so similarly $(a+b)^2 \in R$. Since R is an algebra over a field k_0 of characteristic $\neq 2$,

$$ab = \frac{(a+b)^2 - a^2 - b^2}{2} \in R. \quad \square$$

5.4 Density

Lemma 5.19. *For every non-zero $a \in K$, there is non-zero $b \in Q$ such that $\text{val}(b) \geq \text{val}(a)$.*

Proof. We may assume $a \in \mathcal{O}$ (otherwise take $b = 1$). By Corollary 3.9, there are non-zero $x, y \in R$ such that $x = ay$. By surjectivity of $\widehat{\text{res}} : R \rightarrow k[\varepsilon]$, there is $z \in R$, necessarily non-zero, such that $\widehat{\text{res}}(z) = \varepsilon$. Set $b = xz^2 = ayz^2$. Then

$$\text{val}(b) \geq \text{val}(a)$$

because $y, z \in R \subseteq \mathcal{O}$. And $b \neq 0$, because $y, z \neq 0$. Lastly,

$$\widehat{\text{res}}(b) = \widehat{\text{res}}(x)\widehat{\text{res}}(z^2) = \widehat{\text{res}}(x) \cdot \varepsilon^2 = 0,$$

and so $b \in Q$. □

Lemma 5.20. *For every γ in the value group, there exists a with $\text{val}(a) > \gamma$ and $\text{val}_\partial(a) < -\gamma$.*

Proof. Increasing γ , we may assume $0 < \gamma = \text{val}(b)$ for some $b \in Q$, by Lemma 5.19. Since ς is not weakly multi-valuation type (Definition 5.27 in [10]), the ball of valuative radius 2γ cannot be contained in R . Therefore there is $c \in K$ with $\text{val}(c) > 2\gamma$ and $c \notin R$. Let $a = c/b$. Then

$$\text{val}(a) = \text{val}(c) - \text{val}(b) > 2\gamma - \gamma = \gamma > 0.$$

Therefore $a \in \mathcal{O}$ and $\text{val}_\partial(a)$ is meaningful. If $\text{val}_\partial(a) \geq -\gamma$, then

$$0 \leq \text{val}_\partial(a) + \gamma = \text{val}_\partial(a) + \text{val}(b) \leq \text{val}_\partial(ab) = \text{val}_\partial(c),$$

by Lemma 5.15. Then $c \in R$ by Lemma 5.14, a contradiction. □

Lemma 5.21. *If $a, b \in \mathfrak{m}$ and $\text{val}_\partial(a) < \text{val}_\partial(b)$, then*

$$b = p + qa$$

for some $p, q \in Q$.

Proof. Applying Lemma 4.1 to the set $\{1, a, b\}$ we obtain one of three cases:

- $1 = pa + qb$ for some $p, q \in Q$. This cannot happen, as

$$pa + qb \in \mathcal{O}\mathfrak{m} + \mathcal{O}\mathfrak{m} = \mathfrak{m} \not\cong 1.$$

- $a = p + qb$ for some $p, q \in Q$. By Lemma 5.16,

$$\text{val}_\partial(a) = \text{val}_\partial(p + qb) \geq \min(\text{val}_\partial(p), \text{val}_\partial(q), \text{val}_\partial(b)) = \text{val}_\partial(b),$$

as $\text{val}_\partial(p) = \text{val}_\partial(q) = +\infty$. This contradicts the assumption.

- $b = p + qa$ for some $p, q \in Q$. □

Proposition 5.22. *The set Q is dense in \mathcal{O} , with respect to the valuation topology.*

Proof. We first show that the closure of Q contains \mathfrak{m} . Let b be some element of \mathfrak{m} . Let γ be a given positive element of the valuation group. We will find $c \in Q$ such that $\text{val}(c - b) > \gamma$. If $b \in Q$, we can take $c = b$. Otherwise, $\text{val}_\partial(b) \leq 0$. By Lemma 5.20, there is a such that $\text{val}(a) > \gamma > 0$ and $\text{val}_\partial(a) < \text{val}_\partial(b) \leq 0$. In particular, $a \in \mathfrak{m}$. By Lemma 5.21, there are $c, q \in Q$ such that

$$b = c + qa.$$

Then $\text{val}(b - c) = \text{val}(q) + \text{val}(a) \geq \text{val}(a) > \gamma$, because $q \in Q \subseteq \mathcal{O}$.

Next let b be any element of \mathcal{O} . Take $d \in R$ such that

$$\widehat{\text{res}}(d) = \text{res}(b) + 0\varepsilon.$$

Then $d \in Q$ and $b - d \in \mathfrak{m}$. So we can approximate $b - d$ arbitrarily closely by elements of Q . Equivalently, b is in the closure of $d + Q = Q$. □

5.5 The derivation ∂

Let D be the Q -module

$$D := \mathcal{O}/Q$$

and let $\partial : \mathcal{O} \rightarrow D$ be the natural Q -linear map.

Proposition 5.23. *The Q -module structure on D extends to an \mathcal{O} -module structure as follows: for $a, b \in \mathcal{O}$,*

$$a \cdot \partial b := \partial(a' \cdot b),$$

where $a' \in Q$ and $\text{val}(a' - a) + \text{val}_\partial(b) > 0$. In particular, the choice of a' doesn't matter.

Proof. We first check that $a \cdot \partial b$ is well defined. We can find an $a' \in Q$ such that $\text{val}(a' - a) + \text{val}_\partial(b) > 0$ by Proposition 5.22. If a'' is another such choice, then $a'' - a' \in Q$ and $\text{val}(a'' - a') + \text{val}_\partial(b) > 0$. By Lemma 5.15, it follows that

$$\text{val}_\partial((a'' - a')b) = +\infty,$$

and $(a'' - a')b \in Q$ by Lemma 5.14. Thus $a''b - a'b \in Q$ and $\partial(a''b) = \partial(a'b)$. So the action of \mathcal{O} on D is well-defined. Furthermore, the action of \mathcal{O} on D extends the action of Q . (If $a \in Q$, we can take $a' = a$.)

Next we check the module axioms. For the associative law

$$(a_1 \cdot a_2)\partial b \stackrel{?}{=} a_1(a_2\partial b),$$

take $a'_1, a'_2 \in Q$ such that

$$\begin{aligned} \text{val}(a'_1 a'_2 - a_1 a_2) &> -\text{val}_\partial(b) \\ \text{val}(a'_2 - a_2) &> -\text{val}_\partial(b) \\ \text{val}(a'_1 - a_1) &> -\text{val}_\partial(b). \end{aligned}$$

This is possible by density of Q and the fact that multiplication is continuous. As $a'_2 \in Q$,

$$\text{val}_\partial(a'_2 b) \geq \text{val}(a'_2) + \text{val}_\partial(b) \geq \text{val}_\partial(b),$$

by Lemma 5.15, and so

$$\text{val}(a'_1 - a_1) > -\text{val}_\partial(b) \geq -\text{val}_\partial(a'_2 b).$$

Thus

$$(a_1 a_2) \partial b = \partial(a'_1 a'_2 b) = a_1 \partial(a'_2 b) = a_1 \cdot (a_2 \partial b).$$

The other three module axioms

$$\begin{aligned} (a_1 + a_2) \partial b &= a_1 \partial b + a_2 \partial b \\ a(\partial b_1 + \partial b_2) &= (a \partial b_1) + (a \partial b_2) \\ 1 \partial b &= b \end{aligned}$$

are proven similarly: one replaces the a 's with very close elements of Q .² □

Proposition 5.24. *The map $\partial : \mathcal{O} \rightarrow D$ is a Q -linear derivation:*

- $\partial q = 0$ for $q \in Q$.
- $\partial(ab) = a \partial b + b \partial a$ for $a, b \in \mathcal{O}$.

Proof. The map is Q -linear with kernel Q by construction. Note that $I \neq 0$, as R is a domain and $R/I \cong k[\varepsilon]$ is not. Take non-zero $u \in I$. Then $u \in Q$ and $\text{val}(u) > 0$. Take γ a positive element of the value group such that

$$\begin{aligned} \min(\text{val}_\partial(a), \text{val}_\partial(b)) &> -\gamma \\ \text{val}(u) &< \gamma. \end{aligned}$$

By Proposition 5.22, we can find $a_Q, b_Q \in Q$ and $a', b' \in \mathcal{O}$ such that

$$\begin{aligned} a &= a_Q + a' \\ b &= b_Q + b' \\ \text{val}(a') &> 3\gamma \\ \text{val}(b') &> 3\gamma. \end{aligned}$$

By Lemmas 5.16 and 5.14,

$$\text{val}_\partial(a') \geq \min(\text{val}_\partial(a), \text{val}_\partial(-a_Q)) = \min(\text{val}_\partial(a), +\infty) = \text{val}_\partial(a) > -\gamma.$$

²For the second distributive law, one must choose $a' \in Q$ such that

$$\text{val}(a' - a) + \min\{\text{val}_\partial(b_1), \text{val}_\partial(b_2), \text{val}_\partial(b_1 + b_2)\} > 0.$$

If $\text{val}_\partial(a'/u) \leq -2\gamma$, then

$$\text{val}_\partial(a') = \text{val}_\partial(a'/u) + \text{val}(u) < -2\gamma + \gamma = -\gamma < 0$$

by Lemma 5.15, a contradiction. Thus

$$\begin{aligned}\text{val}_\partial(a'/u) &> -2\gamma \\ \text{val}_\partial(b'/u) &> -2\gamma,\end{aligned}$$

where the second line follows by a similar argument. Also,

$$\begin{aligned}\text{val}(a'/u) &= \text{val}(a') - \text{val}(u) > 3\gamma - \gamma = 2\gamma \\ \text{val}(b'/u) &= \text{val}(b') - \text{val}(u) > 3\gamma - \gamma = 2\gamma.\end{aligned}$$

Thus, by Lemma 5.18, $(a'/u)(b'/u) \in R$. Then

$$a'b' = (a'/u)(b'/u)(u^2) \in R \cdot u^2 \subseteq I \subseteq Q,$$

so we see that

$$\partial(a'b') = 0.$$

Also,

$$\text{val}(0 - a') + \text{val}_\partial(b') = \text{val}(a') + \text{val}_\partial(b) > 3\gamma - \gamma > 0,$$

so $a'\partial b' = \partial(0 \cdot b') = 0$. Similarly, $b'\partial a' = 0$. So

$$\partial(a'b') = 0 = a'\partial b' + b'\partial a'.$$

The other three equations

$$\begin{aligned}\partial(a_Q b') &= a_Q \partial b' + b' \partial a_Q \\ \partial(a' b_Q) &= a' \partial b_Q + b_Q \partial a' \\ \partial(a_Q b_Q) &= a_Q \partial b_Q + b_Q \partial a_Q\end{aligned}$$

hold by Q -linearity and the fact that ∂ vanishes on Q . Adding these four equations, we obtain the desired Leibniz rule

$$\partial(ab) = a\partial b + b\partial a. \quad \square$$

Note that

$$Q = \{a \in \mathcal{O} : \partial a = 0\}.$$

Let Γ be the value group of \mathcal{O} . We define

$$\text{val} : D \rightarrow \Gamma_{\leq 0} \cup \{+\infty\}$$

by the equation

$$\text{val}(\partial a) = \text{val}_\partial(a).$$

By Lemmas 5.14 and 5.16, this is well-defined, and satisfies the identities

$$\begin{aligned}\text{val}(a + b) &\geq \min(\text{val}(a), \text{val}(b)) \\ \text{val}(a) = +\infty &\iff a = 0\end{aligned}$$

for $a, b \in D$.

Lemma 5.25. For $a \in \mathcal{O}$ and $b \in D$,

$$\text{val}(ab) = \text{val}(a) + \text{val}(b),$$

unless the right hand side is positive, in which case

$$\text{val}(ab) = +\infty.$$

Proof. Write b as ∂c for some $c \in \mathcal{O}$. Take $a' \in Q$ such that $\text{val}(a - a') + \text{val}(\partial c) > 0$ and $\text{val}(a - a') > \text{val}(a)$. Then by definition, $a\partial c = \partial(a'c)$. Also $\text{val}(a) = \text{val}(a')$. By Lemma 5.15,

$$\text{val}(a\partial c) = \text{val}(\partial(a'c)) = \text{val}_{\partial}(a'c) = \text{val}(a') + \text{val}_{\partial}(c) = \text{val}(a) + \text{val}(\partial c),$$

unless the right hand side is positive, in which case $\text{val}(ab) = +\infty$. \square

Using this and the fact that ∂ is a derivation, we get an improved version of the multiplication statement in Lemma 5.16.

Corollary 5.26. If $a, b \in \mathcal{O}$, then

$$\text{val}_{\partial}(ab) \geq \min(\text{val}(a) + \text{val}_{\partial}(b), \text{val}_{\partial}(a) + \text{val}(b)).$$

5.6 The module of differentials

The \mathcal{O} -module D of differentials shares many properties with K/\mathfrak{m} .

Lemma 5.27. For any $\gamma \leq 0$ in the value group, there is $b \in D$ such that $\text{val}(b) = \gamma$.

Proof. By Lemma 5.25, it suffices to show that

$$\{\text{val}(b) : b \in D\}$$

has no lower bound, which follows by Lemma 5.20. \square

Lemma 5.28. If $a, b \in D$ and $\text{val}(a) < \text{val}(b)$, then $b \in \mathcal{O} \cdot a$.

Proof. We may assume $b \neq 0$, in which case $\text{val}(a) < \text{val}(b) \leq 0$. By Proposition 5.22, $\mathcal{O} = \mathfrak{m} + Q$. Therefore, we may write $a = \partial a'$ and $b = \partial b'$ for some $a', b' \in \mathfrak{m}$. By Lemma 5.21, we can write

$$b' = p + qa'$$

for some $p, q \in Q$. Then $\partial p = \partial q = 0$, so

$$b = \partial b' = q\partial a' = qa. \quad \square$$

Of course, we can replace $\text{val}(a) < \text{val}(b)$ with a non-strict inequality:

Proposition 5.29. If $a, b \in D$ and $\text{val}(a) \leq \text{val}(b)$, then $b \in \mathcal{O} \cdot a$.

Proof. We may assume $b, a \neq 0$. By Lemma 5.27, there is $z \in D$ with $\text{val}(z) < \text{val}(a)$. Then $a = \alpha z$ and $b = \beta z$ for some $\alpha, \beta \in \mathcal{O}$. By Lemma 5.25

$$\begin{aligned}\text{val}(\alpha) &= \text{val}(a) - \text{val}(z) \\ \text{val}(\beta) &= \text{val}(b) - \text{val}(z).\end{aligned}$$

Then $\text{val}(\alpha) \leq \text{val}(\beta)$, and so $\beta \in \mathcal{O}\alpha$ as \mathcal{O} is a valuation ring. Therefore

$$b = \beta z = \gamma \alpha z = \gamma a \in \mathcal{O}a$$

for some $\gamma \in \mathcal{O}$. □

Proposition 5.30. *D is divisible as an \mathcal{O} -module: for any $b \in D$ and non-zero $a \in \mathcal{O}$, there is $x \in D$ such that $ax = b$.*

Proof. We may assume $b \neq 0$. By Lemma 5.27, there is $c \in D$ such that $\text{val}(c) \leq \text{val}(b) - \text{val}(a)$. Then $\text{val}(ac) \leq \text{val}(b)$, and so

$$b \in \mathcal{O} \cdot ac \subseteq a \cdot D. \quad \square$$

Proposition 5.31. *Let D_0 be the \mathcal{O} -submodule of $x \in D$ such that $\text{val}(x) \geq 0$.*

- D_0 is the image of R under ∂ .
- Viewing k as the \mathcal{O} -module \mathcal{O}/\mathfrak{m} , there is a unique \mathcal{O} -module isomorphism $\text{res}_2 : D_0 \rightarrow k$ such that

$$\widehat{\text{res}}(x) = \text{res}(x) + \text{res}_2(\partial x)\varepsilon$$

for any $x \in R$.

Proof. The first point is clear from Lemma 5.14:

$$\text{val}(\partial x) \geq 0 \iff \text{val}_\partial(x) \geq 0 \iff x \in R.$$

Take some $w \in R$ such that $\widehat{\text{res}}(w) = 0 + \varepsilon$. Then $w \notin Q$, so

$$\text{val}(\partial w) = \text{val}_\partial(w) = 0$$

by Lemma 5.14. By Proposition 5.29, ∂w generates D_0 as an \mathcal{O} -module. Also, $\text{Ann}_\mathcal{O}(\partial w)$ is \mathfrak{m} by Lemma 5.25. Thus there is an isomorphism

$$\begin{aligned}\text{res}_2 : D_0 &\rightarrow \mathcal{O}/\mathfrak{m} \\ y\partial w &\mapsto \text{res}(y).\end{aligned}$$

Now let $x \in R$ be given. Then

$$\widehat{\text{res}}(x) = s + t\varepsilon$$

for some $s, t \in k$. We already know that $s = \text{res}(k)$, and we must show that

$$\text{res}_2(\partial x) = t.$$

Take $y \in R$ with $\widehat{\text{res}}(y) = t + 0\varepsilon$. Then $y \in Q$. Also,

$$\widehat{\text{res}}(x - wy) = \widehat{\text{res}}(x) - \widehat{\text{res}}(w)\widehat{\text{res}}(y) = s + t\varepsilon - \varepsilon t = s,$$

so $x - wy \in Q$. Then

$$\partial x = \partial(wy) = y\partial w,$$

and so

$$\text{res}_2(\partial x) = \text{res}_2(y\partial w) = \text{res}(y) = t.$$

This proves the formula

$$\widehat{\text{res}}(x) = \text{res}(x) + \text{res}_2(\partial x)\varepsilon.$$

Finally, this formula uniquely determines res_2 , because $\partial : R \rightarrow D_0$ is onto. \square

5.7 Odd positive characteristic

Proposition 5.32. *If K is perfect, then $\text{char}(K) = 0$.*

Proof. Suppose for the sake of contradiction that $\text{char}(K) = p > 2$. By construction, the derivation $\partial : \mathcal{O} \rightarrow D$ is onto. Also, D cannot vanish, since we have constructed a submodule D_0 isomorphic to k . Therefore $\partial a \neq 0$ for some $a \in \mathcal{O}$. By perfection of K , we can write $a = b^p$. Then

$$\partial a = \partial(b^p) = pb^{p-1}\partial b = 0,$$

a contradiction. \square

6 Application to fields of dp-rank 2

Recall that a topology on a structure is *definable* if it admits a uniformly definable basis of open sets.

Theorem 6.1. *Let $(K, +, \cdot, 0, 1, \dots)$ be a field of characteristic 0, possibly with extra structure. Suppose K has dp-rank 2 and is unstable.*

1. K does not admit two independent definable valuation rings.
2. K admits a definable non-trivial V -topology.
3. The canonical topology on K is definable. (See §1.1).

We prove these statements in §6.2–6.5, but for now, we give some motivation.

Proposition 6.2. *Under the assumptions of Theorem 6.1, K admits a unique definable non-trivial V -topology.*

Proof. Existence follows from Theorem 6.1.2. For uniqueness, suppose K admits two independent definable V -topologies. We may replace K with an \aleph_0 -saturated elementary extension. Then the two V -topologies are induced by externally definable valuation rings $\mathcal{O}_1, \mathcal{O}_2$, by Proposition 3.5 in [4]. Replacing K with its Shelah expansion K^{Sh} , we obtain two independent valuation rings. The Shelah expansion continues to have dp-rank 2—this is a simple exercise using quantifier elimination in the Shelah expansion ([12], Proposition 3.23). \square

Proposition 6.3. *Let $(K, \mathcal{O}_1, \mathcal{O}_2, \dots)$ be a field with two definable valuation rings $\mathcal{O}_1, \mathcal{O}_2$, and possibly additional structure. If $\text{dp-rk}(K) \leq 2$, then \mathcal{O}_1 and \mathcal{O}_2 are comparable.*

Proof. Suppose $\mathcal{O}_1, \mathcal{O}_2$ are incomparable. The join $\mathcal{O}_1 \cdot \mathcal{O}_2$ is itself a valuation ring. Let K' be the residue field of $\mathcal{O}_1 \cdot \mathcal{O}_2$. Then \mathcal{O}_1 and \mathcal{O}_2 induce two independent valuation rings \mathcal{O}'_1 and \mathcal{O}'_2 on K' . Indeed, there is an isomorphism between

- The poset of valuation rings on K' .
- The poset of valuation rings on K that are contained in $\mathcal{O}_1 \cdot \mathcal{O}_2$.

Thus \mathcal{O}'_1 and \mathcal{O}'_2 are incomparable, and $\mathcal{O}'_1 \cdot \mathcal{O}'_2$ must be the maximal valuation ring on K' , which is K' itself. Thus \mathcal{O}'_1 and \mathcal{O}'_2 are incomparable and independent.

Replacing $(K, \mathcal{O}_1, \mathcal{O}_2)$ with $(K', \mathcal{O}'_1, \mathcal{O}'_2)$, we may assume that \mathcal{O}_1 and \mathcal{O}_2 are independent. Then we get a contradiction:

- If $\text{dp-rk}(K) \leq 1$, use Lemma 9.4.14 in [7].
- If $\text{char}(K) > 0$, use Lemma 2.6 in [8].
- If $\text{dp-rk}(K) = 2$ and $\text{char}(K) = 0$, then Theorem 6.1 applies. \square

Proposition 6.4. *Suppose parts 1 and 2 of Theorem 6.1 hold for all ranks. In other words, suppose the following hold:*

- *No dp-finite field of characteristic 0 admits two independent definable valuation rings.*
- *Every unstable dp-finite field of characteristic 0 admits a (non-trivial) definable V -topology.*

Then the Shelah conjecture holds for dp-finite fields: every dp-finite field is either finite, algebraically closed, real closed, or henselian. Therefore, the conjectured classification of ([3], Theorem 3.11) holds.

Proof. As in Proposition 6.3, we conclude that any two definable valuation rings $\mathcal{O}_1, \mathcal{O}_2$ on a dp-finite field are comparable. As in the proof of Theorem 2.8 in [8], this implies the henselianity conjecture for dp-finite fields: every definable valuation ring on a dp-finite field is henselian.

Because the Shelah expansion of a dp-finite structure is dp-finite, it follows that any externally definable valuation ring must also be henselian.

Claim 6.5. If K is dp-finite of characteristic 0, then one of the following holds:

- K is algebraically closed.
- K is real closed.
- K is finite
- K admits a definable non-trivial valuation ring.

Proof. We may replace K with a sufficiently saturated elementary extension. If K is stable, then K is finite or algebraically closed, by Proposition 7.2 in [5]. Otherwise, K admits a non-trivial definable V-topology, by assumption. By Proposition 3.5 in [4], K admits a non-trivial externally definable valuation ring \mathcal{O} . This valuation ring must be henselian. By Theorem 5.2 in [6], either K admits a definable valuation ring or K is real closed or algebraically closed.³ □_{Claim}

This in turn implies the Shelah conjecture for dp-finite fields of characteristic 0. The case of positive characteristic is Corollary 11.4 [8].

The classification in ([3], Theorem 3.11) is proven conditional on the Shelah conjecture. (The proof is for strongly dependent fields, but can be restricted to the smaller class of dp-finite fields.) □

6.1 The pedestal machine

We review the setup from ([10], Part II). Let \mathbb{K} be an unstable monster field, possibly with extra structure, with $\text{dp-rk}(\mathbb{K}) \leq 2$.

Fix a *magic subfield* $k_0 \preceq \mathbb{K}$, i.e., a small model with the following property (Definition 8.3 in [9]):

For every k_0 -linear subspace $G \leq (\mathbb{K}, +)$, if G is type-definable (over any small set), then $G = G^{00}$.

Magic subfields exist by ([8], Corollary 8.7).

Let Λ denote the lattice of type-definable k_0 -linear subspaces of \mathbb{K} . Recall from ([8], Definition 9.13) that a *strict n -cube* in Λ is an injection

$$\begin{aligned} \text{Pow}(n) &\hookrightarrow \Lambda \\ S &\mapsto G_S \end{aligned}$$

that preserves the unbounded lattice operations:

$$\begin{aligned} G_{S_1 \cup S_2} &= G_{S_1} + G_{S_2} \\ G_{S_1 \cap S_2} &= G_{S_1} \cap G_{S_2}. \end{aligned}$$

³If K is separably closed, then K is algebraically closed, because dp-finite fields are perfect.

We call G_\emptyset the *base* of the cube; the base need not be 0.

The *reduced rank* of Λ is the maximum r such that a strict r -cube exists (Definition 9.17 in [8]). By Proposition 10.1.7 in [8], the reduced rank is 1 or 2.

If r is the reduced rank, a *pedestal* in Λ is a group $G \in \Lambda$ that is the base of a strict r -cube (Definition 8.4 in [9]⁴). Since r is small, we can describe what this means explicitly:

- If $r = 1$, then an r -cube is a chain of length two, and a pedestal is any $G \in \Lambda$ other than \mathbb{K} itself.
- If $r = 2$, then an r -cube is

$$\{G \cap H, G, H, G + H\}$$

for two incomparable $G, H \in \Lambda$. Therefore, a pedestal is a group of the form $G \cap H$ where G, H are incomparable elements of Λ .

Fact 6.6 (Proposition 10.4.1 in [8]). *Non-zero pedestals exist.*

In Theorem 9.3 of [10], we associated an r -inflator to any non-zero pedestal H .

Fact 6.7. *Let H be a non-zero pedestal with associated r -inflator ς .*

1. ς is malleable.
2. The fundamental ring R_H of ς is given as

$$R_H = \{x \in \mathbb{K} : xH \subseteq H\}.$$

3. If H is type-definable over a small model K containing k_0 , then the infinitesimals J_K are contained in the fundamental ideal I_H .
4. If H is type-definable over a small model K containing k_0 , then $R_H \cdot J_K \subseteq J_K$, and so J_K is a sub-ideal of the fundamental ideal I_H .
5. If ς' is obtained by mutating ς along a line $\mathbb{K} \cdot (a_1, \dots, a_n)$, then ς' is the r -inflator associated to the group

$$H' = (a_1^{-1}H) \cap \dots \cap (a_n^{-1}H)$$

In particular, H' is itself a non-zero pedestal.

This follows from ([10], Theorem 9.3, Remark 9.5, Proposition 10.15), ([8], Proposition 10.15.5, Lemma 10.20), and Lemma 6.9 below.

Remark 6.8. By construction ([8], Theorem 4.20.4, Definition 6.3), the family of basic neighborhoods is uniformly ind-definable across all models. In other words, there is a set of formulas $\{\psi_i(x; \vec{z}_i)\}_{i \in I}$ such that for any model K , the collection of basic neighborhoods on K is exactly

$$\{\psi_i(K; \vec{c}) : i \in I, \vec{c} \in K^{|\vec{z}_i|}\}.$$

⁴Pedestals were called “special groups” in §10 of [8]

Lemma 6.9. *Let G be a non-zero pedestal, type-definable over a small model $K \preceq \mathbb{K}$, with K extending k_0 . Let R be the stabilizer ring of G :*

$$R = \{x \in \mathbb{K} : xG \subseteq G\}.$$

Let J_K be the group of K -infinitesimals. Then J_K is an ideal in R .

The following proof was sketched in Remark 10.19 of [8].

Proof. First of all, $J_K \subseteq R$ by Proposition 10.15.(2,5) in [8]. It remains to show that $R \cdot J_K \subseteq J_K$. Take a non-zero element $j_0 \in G$. Take a small model $K' \preceq \mathbb{K}$ with $K' \supseteq K \cup \{j_0\}$. As G is type-definable over the larger model K' , we see that

$$J_{K'} \cdot G \subseteq J_{K'}$$

by ([8], Proposition 10.4.3). Now for any $\varepsilon \in J_{K'}$ and $a \in R$, we have

$$\varepsilon \cdot a \cdot j_0 \in J_{K'} \cdot R \cdot G \subseteq J_{K'} \cdot G \subseteq J_{K'},$$

implying that $\varepsilon \cdot a \in j_0^{-1} J_{K'}$. As $J_{K'}$ is invariant under scaling by elements of $(K')^\times$ ([8], Remark 6.9.3), we see that $j_0^{-1} J_{K'} = J_{K'}$, and

$$\varepsilon \cdot a \in J_{K'}.$$

As $a \in R$ and $\varepsilon \in J_{K'}$ were arbitrary,

$$R \cdot J_{K'} \subseteq J_{K'}. \tag{7}$$

Claim 6.10. If $S \subseteq R$ is type-definable over K , and U is a K -definable basic neighborhood, then there is a K -definable basic neighborhood V such that

$$S \cdot V \subseteq U.$$

Proof. Since $K' \supseteq K$, the neighborhood U is K' -definable and contains $J_{K'}$. Therefore,

$$S \cdot J_{K'} \subseteq R \cdot J_{K'} \subseteq J_{K'} \subseteq U,$$

by (7). By compactness, there is a K -definable set $S' \supseteq S$, and a K' -definable basic neighborhood $V' \supseteq J_{K'}$ such that

$$S' \cdot V' \subseteq U.$$

We can write V' as $\psi_i(\mathbb{K}; \vec{b})$ for one of the formulas ψ_i in Remark 6.8. Since S' and U are K -definable, we can find \vec{c} from K such that

$$S' \cdot \psi_i(\mathbb{K}; \vec{c}) \subseteq U.$$

Take $V = \psi_i(\mathbb{K}; \vec{c})$. Then

$$S \cdot V \subseteq S' \cdot V \subseteq U. \tag{Claim}$$

Now by compactness, it follows that for any subset $S \subseteq R$ that is type-definable over K , we have

$$S \cdot J_K \subseteq J_K.$$

As the ring R is K -invariant, it is a union of such subsets S , and therefore

$$R \cdot J_K \subseteq J_K,$$

as desired. □

6.2 The valuation-type case

In [9], we considered the case where the canonical topology is a V-topology. We say that \mathbb{K} is *valuation type* if this holds. We showed in this case that

- The canonical topology is a definable V-topology ([9], Lemma 7.1).
- Any two definable valuation rings are dependent ([9], Lemma 9.5).

Thus, the three parts of Theorem 6.1 are automatic in this case.

Fact 6.11 (Theorem 8.11 in [9]). *If K is a small submodel and if J_K contains a non-zero ideal of some multi-valuation ring on \mathbb{K} , then the canonical topology on \mathbb{K} is a V-topology.*

This has several consequences:

Corollary 6.12. *Let G be a non-zero pedestal with stabilizer R and associated r -inflator ς .*

1. *If R contains a non-zero ideal of a multi-valuation ring, then \mathbb{K} is valuation type.*
2. *If ς is weakly multi-valuation type, then \mathbb{K} is valuation type.*
3. *If $r = 1$, then \mathbb{K} is valuation type.*
4. *If some mutation of ς is weakly multi-valuation type, then \mathbb{K} is valuation type.*

Proof. 1. Lemma 6.9—the point is that if R' is a multi-valuation ring, and

$$a_1 R' \subseteq R,$$

then $a_2 a_1 R' \subseteq a_2 R \subseteq J_K$ for any non-zero $a_2 \in J_K$.

2. R is the fundamental ring of ς , and “weakly multi-valuation type” means that the fundamental ring contains a non-zero multi-valuation ideal ([10], Definition 5.27).
3. 1-inflators are multi-valuation type (Proposition 5.19 in [10]).
4. If ς' is obtained from ς by mutation, then ς' is the r -inflator associated to some other non-zero pedestal G' (Fact 6.7.5). In particular, if ς' is weakly of multi-valuation type, then G' shows that \mathbb{K} is valuation type. \square

Theorem 6.13. *If \mathbb{K} is not valuation type (and characteristic 0 and unstable), then there is a small model K and a 2-inflator ς satisfying the Strong Assumptions of §3–5, such that the infinitesimals J_K are an ideal in the fundamental ring R of ς .*

Proof. By Fact 6.6, non-zero pedestals exist. Let G be some non-zero pedestal and ς be the associated inflator. Then G satisfies the Weak Assumptions:

- K has characteristic 0 by assumption.

- ς is malleable by Fact 6.7.1.
- $r = 2$ by Corollary 6.12.3.
- No mutation of ς is weakly multi-valuation type, by Corollary 6.12.4.

By Corollary 2.3, there is a mutation ς' of ς such that ς' is isotypic. Then ς' inherits the other properties from ς (see Remark 2.1), and therefore ς' satisfies the Strong Assumptions. By Fact 6.7.5, ς' is the 2-inflator coming from some other pedestal G' . Let K be a small model containing k_0 , and type-defining G' . By Fact 6.7.4, J_K is an ideal in the fundamental ring of ς' . \square

In the remainder of §6, we therefore assume

1. \mathbb{K} is a monster model of an unstable field of dp-rank 2 and characteristic 0.
2. k_0 is a magic subfield.
3. ς is a k_0 -linear 2-inflator on \mathbb{K} satisfying the Strong Assumptions of §3-5, including isotopy.
4. R and I are the fundamental ring and ideal of ς , and D and ∂ are as in §5.5.
5. J is the group of K -infinitesimals over some small model $K \preceq \mathbb{K}$ containing k_0 . In particular,
 - J is type-definable
 - J is contained in every K -definable basic neighborhood.
 - J is non-zero ([8], Remark 6.9.1).
6. J is an ideal in R , contained in the fundamental ideal I .

6.3 Independent valuation rings

Recall that the valuation ring \mathcal{O} is the integral closure of R .

Proposition 6.14. *Let \mathcal{O}' be a valuation ring on \mathbb{K} , independent from \mathcal{O} . Then $\mathcal{O}' \not\supseteq J$.*

Proof. Assume for the sake of contradiction that $\mathcal{O}' \supseteq J$. Take nonzero $e \in J$. Then $R \cdot e \subseteq J$, so

$$R \subseteq e^{-1} \cdot J \subseteq e^{-1} \mathcal{O}'.$$

Let val' be the valuation from \mathcal{O}' and let $\gamma = \text{val}'(e^{-1})$. Then

$$x \in R \implies \text{val}'(x) \geq \text{val}'(e^{-1}) = \gamma.$$

We claim that for all $a \in K$,

$$x \in \mathfrak{m} \implies \text{val}'(x) \geq \min(\gamma, \gamma/2)$$

Indeed, suppose $x \in \mathfrak{m}$. Then x^{-1} isn't integral over R , so by Lemma 4.4,

$$x^2 + bx + c = 0$$

for some $b, c \in R$. By Newton polygons,

$$\text{val}'(x) \geq \min\left(\text{val}'(b), \frac{\text{val}'(c)}{2}\right) \geq \min(\gamma, \gamma/2).$$

On the other hand, \mathcal{O} is independent from \mathcal{O}' , so by the approximation theorem, there is $x \in K$ with $\text{val}(x) > 0$ and $\text{val}'(x) < \min(\gamma, \gamma/2)$, a contradiction. \square

Corollary 6.15. *If $\mathcal{O}_1, \mathcal{O}_2$ are two 0-definable valuation rings on \mathbb{K} , then \mathcal{O}_1 and \mathcal{O}_2 are not independent.*

Proof. The definable set \mathcal{O}_i has full dp-rank for $i = 1, 2$. It follows that $\mathcal{O}_i - \mathcal{O}_i$ is a 0-definable basic neighborhood, and so

$$J \subseteq \mathcal{O}_i - \mathcal{O}_i = \mathcal{O}_i$$

for $i = 1, 2$. By Proposition 6.14, both \mathcal{O}_1 and \mathcal{O}_2 induce the same topology as the valuation ring \mathcal{O} —the integral closure of R . \square

6.4 The definable V-topology

Let $\text{val} : \mathbb{K} \rightarrow \Gamma$ be the valuation associated to \mathcal{O} . We will show that the associated valuation topology is definable.

Lemma 6.16. *There is a type-definable set $B \subseteq \mathbb{K}$ and some $\gamma \in \Gamma$ such that*

$$\text{val}(x) > \gamma \implies x \in B \implies \text{val}(x) \geq 0.$$

for $x \in \mathbb{K}$.

Proof. Let B be the set

$$B = \{x \in \mathbb{K} \mid \exists y, z \in J : x^2 = yx + z\}.$$

Then B is type-definable. If $x \in B$, then

$$x^2 = yx + z$$

for some $y, z \in J \subseteq R$, and so x lies in the integral closure \mathcal{O} of R .

Now take non-zero $c \in J$, and let $\gamma = \text{val}(c)$. Note $c \in R \subseteq \mathcal{O}$, so $\gamma \geq 0$. Suppose $\text{val}(x) > \gamma$. Then $\text{val}(x/c) > 0$, so $x/c \in \mathfrak{m}$ and $c/x \notin \mathcal{O}$. By Lemma 4.4, either x/c or c/x satisfies a monic polynomial equation of degree 2 over R . As c/x is not in the integral closure \mathcal{O} of R , we see that x/c satisfies the equation:

$$(x/c)^2 = (x/c)y_0 + z_0$$

for some $y_0, z_0 \in R$. Then

$$x^2 = (cy_0)x + c^2z_0,$$

and $cy_0, c^2z_0 \in J$. Thus $x \in B$. \square

Say that two subsets $X, Y \subseteq \mathbb{K}$ are “co-embeddable” if there exist $a, b \in \mathbb{K}^\times$ such that

$$\begin{aligned} a \cdot X &\subseteq Y \\ b \cdot Y &\subseteq X. \end{aligned}$$

This is an equivalence relation.

Remark 6.17. Suppose X and Y are co-embeddable, X is type-definable, and Y is \vee -definable. Then there is a definable set Z co-embeddable with X and Y . Indeed, after rescaling, we may assume

$$X \subseteq Y.$$

Then we may find a definable set Z interpolating X and Y , by compactness:

$$X \subseteq Z \subseteq Y.$$

Lemma 6.18. *There is a definable set B that is co-embeddable with \mathcal{O} .*

Proof. By Lemma 6.16, there is a type-definable set B_0 and $\gamma \in \Gamma$ such that

$$\text{val}(x) > \gamma \implies x \in B_0 \implies \text{val}(x) \geq 0.$$

Therefore B_0 is co-embeddable with \mathcal{O} . Let B_1 be the \vee -definable set $B_1 = \{0\} \cup \{y \in \mathbb{K}^\times : y^{-1} \notin B_0\}$. Note that

$$\text{val}(y) > 0 \implies y \in B_1 \implies \text{val}(y) \geq -\gamma.$$

Thus B_1 is co-embeddable with \mathcal{O} . By Remark 6.17, there is a definable set in the co-embeddability class of B_0, B_1 , and \mathcal{O} . \square

Recall from Lemma 2.1(d) in [11], that a set S in a topological field K is *bounded* if and only if for every open neighborhood $U \ni 0$, there is non-zero $a \in K^\times$ such that

$$a \cdot S \subseteq U.$$

Theorem 6.19. *The V -topology induced by \mathcal{O} is definable.*

Proof. Take a definable set B that is co-embeddable with \mathcal{O} . Then B is a bounded neighborhood of 0, with respect to the V -topology induced by \mathcal{O} . Therefore, the following definable family is a neighborhood basis of 0, by Lemma 2.1(e) in [11]:

$$\{aB : a \in \mathbb{K}^\times\}.$$

This proves definability, by Lemma 6.20 below. \square

Lemma 6.20. *Let $(K, +, \cdot, \dots)$ be a field, possibly with extra structure. Let τ be a field topology on K . Then τ is definable if and only if there is a definable neighborhood basis of 0.*

Proof. If $\{U_a\}_{a \in Y}$ is a definable basis of opens, then

$$\{U_a : a \in Y \text{ and } 0 \in U_a\}$$

is a definable neighborhood basis of 0. Conversely, suppose $\{N_a\}_{a \in Y}$ is a definable neighborhood basis of 0. Let

$$N_a^{\text{int}} = \{x \in K \mid \exists b \in Y : x + N_b \subseteq N_a\}.$$

Then $\{N_a^{\text{int}}\}_{a \in Y}$ is a definable basis of open neighborhoods around 0, and

$$\{b + N_a^{\text{int}} : b \in K, a \in Y\}.$$

is a definable basis of open sets. □

6.5 Definability of the canonical topology

Lemma 6.21. *There is nonnegative $\gamma \in \Gamma$, and a type-definable set S , such that for $x \in \mathbb{K}$ with $\text{val}(x) > \gamma$, we have*

$$\text{val}(\partial x) < -\gamma \implies x \in S \implies \text{val}(\partial x) \leq 0.$$

Proof. Take some non-zero $c \in J$. Take $\gamma = \text{val}(c)$. Then $\gamma \geq 0$, as

$$J \subseteq R \subseteq \mathcal{O}.$$

Take $e_0 \in R$ such that $\widehat{\text{res}}(e_0) = s + t\varepsilon$, with $t \neq 0$. Then $e_0 \in R \setminus Q$. Let B be the open ball of valuative radius γ . By Proposition 5.22, $\mathcal{O} = B + Q$. Therefore there is $e \in B$ with $e - e_0 \in Q$. Then $e \in R \setminus Q$, and $\text{val}(e) > \gamma = \text{val}(c)$.

Let S be the type-definable set of $x \in \mathbb{K}$ such that

$$\exists y, z \in J : e = xy + z.$$

Suppose $\text{val}(x) > \gamma$ and $\text{val}(\partial x) < -\gamma$. Apply Lemma 4.1 to the set $\{xc, e, c\}$. There are three cases:

- c is generated by xc and e . This cannot happen, since $R \subseteq \mathcal{O}$, since $\text{val}(c) < \text{val}(e)$ (by choice of e), and since $\text{val}(c) < \text{val}(xc)$ as $\text{val}(x) > \gamma \geq 0$.
- xc is generated by e and c . As $e, c \in R$, this would imply $xc \in R$. But $c \in J \subseteq I \subseteq Q$, so by Lemma 5.15,

$$\text{val}(\partial(xc)) = \text{val}(c) + \text{val}(\partial x) < \text{val}(c) - \gamma = 0.$$

By Lemma 5.14, $xc \notin R$, a contradiction.

- e is generated by c and xc . Then

$$e = xcy_0 + cz_0,$$

for some $y_0, z_0 \in R$. If $y = cy_0$ and $z = cz_0$, then $y, z \in J$ (as $J \triangleleft R$), and $e = xy + z$. So $x \in S$.

Conversely, suppose $x \in S$ and $\text{val}(x) > \gamma$. Then there are $y, z \in J \subseteq I \subseteq Q$ such that

$$e = xy + z.$$

Now Q is a subring, and $y, z \in Q$, $e \notin Q$. Therefore $x \notin Q$. On the other hand, $\text{val}(x) > \gamma \geq 0$, so $x \in \mathcal{O}$. Therefore $x \in \mathcal{O} \setminus Q$, which implies $\text{val}(\partial x) \leq 0$ by Lemma 5.14. \square

Lemma 6.22. *Some \vee -definable set is co-embeddable with R .*

Proof. Take γ and S as in Lemma 6.21. By Theorem 6.19, we can find γ' and a definable set B such that

$$\text{val}(x) > \gamma' \implies x \in B \implies \text{val}(x) > \gamma.$$

Claim 6.23. If $\text{val}(x) > \gamma'$ and $\text{val}(\partial x) > 0$, then $x \in B \setminus S$.

Proof. Because $\text{val}(x) > \gamma'$, we have $x \in B$ and $\text{val}(x) > \gamma$, and so Lemma 6.21 applies. Then $\text{val}(\partial x) > 0$ implies $x \notin S$, by the contrapositive to Lemma 6.21. \square_{Claim}

Claim 6.24. If $x \in B \setminus S$, then $\text{val}(x) > \gamma$ and $\text{val}(\partial x) \geq -\gamma$.

Proof. The fact that x is in B implies that $\text{val}(x) > \gamma$, and thus that Lemma 6.21 applies. By the contrapositive to Lemma 6.21, $x \notin S$ implies $\text{val}(\partial x) \geq -\gamma$. \square_{Claim}

By Lemma 5.19, there is $b \in Q$ such that $\text{val}(b) > \gamma'$. Then

$$bR \subseteq B \setminus S. \tag{8}$$

Indeed, if $x \in R$, then

$$\begin{aligned} \text{val}(bx) &= \text{val}(b) + \text{val}(x) > \gamma' + 0 \\ \text{val}(\partial(bx)) &\geq \text{val}(b) + \text{val}(\partial x) > \gamma' + 0 \geq 0, \end{aligned}$$

and Claim 6.23 applies. Also,

$$b \cdot (B \setminus S) \subseteq R. \tag{9}$$

Indeed, if $x \in B \setminus S$, then

$$\begin{aligned} \text{val}(x) &> \gamma \\ \text{val}(\partial x) &\geq -\gamma \end{aligned}$$

by Claim 6.24. But then $bx \in R$:

$$\begin{aligned} \text{val}(bx) &= \text{val}(b) + \text{val}(x) > \gamma' + \gamma \geq 0 \\ \text{val}(\partial(bx)) &\geq \text{val}(b) + \text{val}(\partial x) > \gamma' - \gamma \geq 0. \end{aligned}$$

By (8)-(9), the \vee -definable set $B \setminus S$ is co-embeddable with R . \square

Lemma 6.25. *The set J is bounded with respect to the canonical topology on \mathbb{K} : for any basic neighborhood U , there is $a \in \mathbb{K}^\times$ such that $aJ \subseteq U$.*

Proof. Recall that J is the set J_K of K -infinitesimals. Let K' be a small model containing K and defining U . Then U contains the group $J_{K'}$ of K' -infinitesimals. By Corollary 8.9 in [9], there is non-zero a such that

$$a \cdot J_K \subseteq J_{K'} \subseteq U. \quad \square$$

Recall from [11], §2, that a ring topology is *locally bounded* if there is a bounded neighborhood of 0. If R is a proper subring of a field $K = \text{Frac}(R)$, then R induces a locally bounded ring topology on K , as in [11], Example 1.2 and Theorem 2.2(a).

Theorem 6.26. *The canonical topology on \mathbb{K} is locally bounded, definable, and induced by R .*

Proof. Note that J and R are co-embeddable, as

$$cR \subseteq J \subseteq R$$

for any non-zero $c \in J$. By Lemma 6.22, some \vee -definable set U is co-embeddable with R and J . As J itself is type-definable, we can take U to be *definable* by Remark 6.17. Rescaling U , we may assume $J \subseteq U$. By compactness, there is a K -definable basic neighborhood V such that

$$J \subseteq V \subseteq U,$$

as J is the directed intersection of such neighborhoods. Therefore U is a neighborhood of 0. Also, U is bounded, because it is co-embeddable with the bounded set J . Therefore the canonical topology is locally bounded. By Lemma 2.1(e) in [11], the family

$$\{aU : a \in \mathbb{K}^\times\}$$

is a neighborhood basis of 0. Then the canonical topology is definable by Lemma 6.20. The family

$$\{aR : a \in \mathbb{K}^\times\}$$

is also a neighborhood basis of 0, because U and R are co-embeddable. □

Once the canonical topology is definable on the monster, it is uniformly definable on all models:

Theorem 6.27.

1. *There is a formula $\varphi(x; \vec{y})$ such that for every small model $K \preceq \mathbb{K}$, the family of sets*

$$\{\varphi(K; \vec{b}) : \vec{b} \in K^{|\vec{y}|}\}$$

is a neighborhood basis of 0 for the canonical topology on K .

2. *If K, K' are two small submodels, then K and K' with their canonical topologies are “locally equivalent” in the sense of [11].*

Proof. Let $\{\psi_i(x; \vec{z}_i)\}_{i \in I}$ be as in Remark 6.8, so that

$$\{\psi_i(K; \vec{c}) : i \in I, \vec{c} \in K^{|\vec{z}_i|}\}$$

is the set of basic neighborhoods on any $K \equiv \mathbb{K}$.

On \mathbb{K} , Theorem 6.26 gives a (\mathbb{K} -)definable neighborhood basis \mathcal{N} . By saturation, there must be a finite subset $I_0 \subseteq I$ such that every set in \mathcal{N} has the form $\psi_i(K; \vec{c})$ for some $i \in I_0$. The fact that \mathcal{N} is a neighborhood basis implies that

$$\forall j \in I \forall \vec{c} \exists i \in I_0 \exists \vec{c}' : \psi_i(\mathbb{K}; \vec{c}') \subseteq \psi_j(\mathbb{K}; \vec{c}).$$

This is a small conjunction of first-order sentences, so it holds in submodels $K \preceq \mathbb{K}$. Then for any small model K , the family

$$\{\psi_i(K; \vec{c}) : i \in I_0, \vec{c} \in K^{|\vec{z}_i|}\}$$

is a neighborhood basis of 0. Because I_0 is finite, this can be written as

$$\{\varphi(K; \vec{b}) : \vec{b} \in K^{|\vec{y}|}\}$$

for some formula $\varphi(x; \vec{y})$.

This proves the first point. The second point is immediate, because local sentences can be evaluated on a neighborhood basis ([11], Theorem 1.1(a)). \square

6.6 Odd positive characteristic

In Theorem 6.13, we can weaken the assumption $\text{char}(K) = 0$ to $\text{char}(K) \neq 2$; the same proof works. But the positive characteristic case then leads to a contradiction:

Proposition 6.28. *Let \mathbb{K} be a monster model of an unstable field with $\text{dp-rk}(\mathbb{K}) \leq 2$. If \mathbb{K} is not of valuation type, then $\text{char}(\mathbb{K})$ is 0 or 2.*

Proof. Suppose $\text{char}(\mathbb{K}) > 2$. As in the proof of Theorem 6.13, there would be a 2-inflator satisfying the Strong Assumptions of §3-5. But strongly dependent fields are perfect, and so $\text{char}(\mathbb{K}) = 0$ by Proposition 5.32. \square

Therefore

Theorem 6.29. *If K is a field with $\text{dp-rk}(K) \leq 2$ and $\text{char}(K) > 2$, then either K is stable, or K is valuation type.*

Perhaps this can be proven in characteristic 2 as well.

7 Reduced rank and generators

Let R be a noncommutative ring, and M be an R -module. Say that M has *property W_n* if the following holds: for any $a_0, a_1, \dots, a_n \in M$, there is some $0 \leq i \leq n$ such that

$$a_i \in R \cdot a_1 + \dots + R \cdot a_{i-1} + R \cdot a_{i+1} + \dots + R \cdot a_n.$$

In other words, any submodule of M generated by a set S of size $n + 1$ is generated by an n -element subset of S .

Remark 7.1. This property appeared in Lemma 4.1, which said that K has property W_2 as a Q -module or R -module.

Lemma 7.2.

1. If M_0, \dots, M_n are non-zero, then $M_0 \oplus \dots \oplus M_n$ does not have property W_n .
2. If M has property W_n and $N \leq M$, then N has property W_n .
3. If M has property W_n and $N \leq M$, then M/N has property W_n .

Proof.

1. Take a_i a non-zero element of M_i , viewed as an element of the direct sum. Then $\{a_0, a_1, \dots, a_n\}$ violates property W_n .
2. Clear.
3. Given $a_i \in M/N$, lift them to $\tilde{a}_i \in M$, apply property W_n in M to obtain i and $r_0, \dots, r_n \in R$ such that

$$\tilde{a}_i = r_0 \tilde{a}_0 + \dots + r_{i-1} \tilde{a}_{i-1} + r_{i+1} \tilde{a}_{i+1} + \dots + r_n \tilde{a}_n,$$

and then project back to M/N . □

Proposition 7.3. M has property W_n if and only if the reduced rank of $\text{Sub}_R(M)$ is at most n .

Proof. If the reduced rank of $\text{Sub}_R(M)$ is greater than n , then there is a strict $(n + 1)$ -cube in M . This corresponds to submodules $M^- \leq M^+ \leq M$ and an isomorphism

$$M^+/M^- \cong N_0 \oplus \dots \oplus N_n$$

where the N_i are non-zero R -modules. By Lemma 7.2, the right hand side does not satisfy W_n , and therefore neither do M^+ or M .

Conversely, suppose W_n fails, witnessed by $a_0, \dots, a_n \in M$. Let $N_i = M \cdot a_i$. Then

$$N_0 + \dots + N_n > N_0 + \dots + N_{i-1} + N_{i+1} + \dots + N_n,$$

for any $0 \leq i \leq n$. By Proposition 6.3.2 in [10], $\text{Sub}_R(M)$ has reduced rank greater than n . □

8 Diffeoevaluation data

In §8, all fields will have characteristic 0, and all rings will be \mathbb{Q} -algebras.

8.1 Mock K/\mathfrak{m} 's

Let K be a valued field with valuation ring \mathcal{O} , maximal ideal \mathfrak{m} , and residue field $k = \mathcal{O}/\mathfrak{m}$.

Definition 8.1. A *mock* K/\mathfrak{m} is a divisible \mathcal{O} -module D extending k satisfying the following property: for any $x, y \in D$,

$$x \in \mathcal{O} \cdot y \text{ or } y \in \mathcal{O} \cdot x.$$

Note that K/\mathfrak{m} is naturally a mock K/\mathfrak{m} .

There is a theory T whose models are pairs (K, D) , where K is a valued field and $k \hookrightarrow D$ is a mock K/\mathfrak{m} .

Proposition 8.2. *Let (K, D) be a model of T . If (K, D) is countable or \aleph_1 -resplendent, then D is isomorphic (as an extension of k) to K/\mathfrak{m} .*

Proof. The resplendent case follows from the countable case. Assume K, D are countable. Then the value group Γ has countable cofinality. Take a sequence

$$a_0, a_1, \dots$$

in K such that $a_0 = 1$ and the sequence

$$\text{val}(a_0), \text{val}(a_1), \dots$$

is descending with no lower bound. Then

$$\mathcal{O} = \mathcal{O} \cdot a_0 \subseteq \mathcal{O} \cdot a_1 \subseteq \dots$$

and the union of this chain is K .

By divisibility, we can find a sequence

$$b_0, b_1, \dots$$

in D such that

- b_0 is the image of 1 under the embedding $k \hookrightarrow D$.
- $b_{i-1} = (a_{i-1}/a_i)b_i$, for all $i \geq 1$.

By induction on i , the b_i are all non-zero. Define $f_i : \mathcal{O} \cdot a_i \rightarrow D$ by $f_i(x) = (x/a_i)b_i$. If $x \in \mathcal{O} \cdot a_i$, then

$$f_i(x) = (x/a_i)b_i = (x/a_i)(a_i/a_{i+1})b_{i+1} = (x/a_{i+1})b_{i+1} = f_{i+1}(x).$$

Therefore the f_i glue together to yield a morphism

$$f : K \rightarrow D.$$

Moreover, $f(a_i) = f_i(a_i) = (a_i/a_i)b_i = b_i$ for all i .

Claim 8.3. For any $x \in K$,

$$f(x) = 0 \iff \text{val}(x) > 0.$$

Proof. First suppose $\text{val}(x) \geq 0$. Then $x \in \mathcal{O} = \mathcal{O} \cdot a_0 = \text{dom}(f_0)$, and so

$$f(x) = f_0(x) = (x/a_0)b_0 = xb_0.$$

By choice of b_0 , the annihilator $\text{Ann}_{\mathcal{O}}(b_0)$ is exactly \mathfrak{m} , and so

$$f(x) = 0 \iff xb_0 = 0 \iff x \in \mathfrak{m} \iff \text{val}(x) > 0.$$

Next suppose $\text{val}(x) \leq 0$. Then $1/x \in \mathcal{O}$, and so

$$f(1) = f((1/x)x) = (1/x)f(x),$$

because f is \mathcal{O} -linear. By the first case,

$$\text{val}(1) = 0 \implies f(1) \neq 0 \implies f(x) \neq 0.$$

□_{Claim}

Therefore $\ker(f) = \mathfrak{m}$, and f induces an embedding

$$K/\mathfrak{m} \hookrightarrow D.$$

Restricted to $k = \mathcal{O}/\mathfrak{m}$, this embedding is

$$(x + \mathfrak{m}) \mapsto f(x) = f_0(x) = (x/a_0)b_0 = xb_0.$$

By choice of b_0 , this is the given embedding of k into D .

It remains to show that f is onto. Suppose not. Take an element y of D that is not in the image of f . Then

$$y \notin \mathcal{O} \cdot b_i,$$

for any i , since $\mathcal{O} \cdot b_i$ is the image of f_i . By definition of mock K/\mathfrak{m} , it follows that

$$b_i \in \mathcal{O} \cdot y$$

for all i . In particular, there are $c_i \in \mathcal{O}$ such that $b_i = c_i y$. Take i large enough that $\text{val}(a_i c_0) < 0$. Then

$$(1/a_i)y = \frac{1}{a_i c_0} c_0 y = \frac{1}{a_i c_0} b_0 = 0,$$

because $\text{Ann}(b_0) = \mathfrak{m}$. But then

$$b_0 = f_i(a_0) = f_i(1) = (1/a_i)b_i = (1/a_i)c_i y = c_i(1/a_i)y = c_i 0 = 0,$$

contradicting the choice of b_0 . □

Corollary 8.4. *Let D be a mock K/\mathfrak{m} . Then there is a map $\text{val} : D \rightarrow \Gamma \cup \{+\infty\}$ with the following properties:*

1. $\text{val}(x) \leq 0$ or $\text{val}(x) = +\infty$ for all $x \in D$.
2. $\text{val}(x) = +\infty$ if and only if $x = 0$.
3. $\text{val}(x) \geq 0$ if and only if x is in the image of $k \hookrightarrow D$.
4. For any $a \in \mathcal{O}$ and $x \in D$,

$$\text{val}(ax) = \begin{cases} \text{val}(a) + \text{val}(x) & \text{if } \text{val}(a) + \text{val}(x) \leq 0 \\ +\infty & \text{if } \text{val}(a) + \text{val}(x) > 0. \end{cases}$$

5. For any $x, y \in D$,

$$\text{val}(x + y) \geq \min(\text{val}(x), \text{val}(y)).$$

6. If $\text{val}(x) \leq \text{val}(y)$, then $y \in \mathcal{O} \cdot x$.

7. For any $\gamma \in \Gamma$, there is $x \in D$ such that $\text{val}(x) \leq \gamma$.

Proof. Let D^* be the image of $k \setminus \{0\}$ in D . We claim that

- For every non-zero x in D , there is an $a \in \mathcal{O}$ such that $ax \in D^*$.
- If $ax \in D^*$ and $bx \in D^*$, then $\text{val}(a) = \text{val}(b)$.
- If we define $\text{val} : D \rightarrow \Gamma \cup \{+\infty\}$ as

$$\text{val}(x) = \begin{cases} +\infty & \text{if } x = 0 \\ -\text{val}(a) & \text{if } ax \in D^*, \end{cases}$$

then val satisfies the listed conditions.

These three claims can be expressed by a first-order sentence, so we may pass to a resplendent elementary extension. Then we may assume D is K/\mathfrak{m} , in which case the three claims are straightforward. \square

8.2 Diffeovalued fields

Definition 8.5. A *diffeovalued field* is a structure $(K, \mathcal{O}, D, \partial)$ where

- (K, \mathcal{O}) is a valued field (of equicharacteristic 0).
- $k \hookrightarrow D$ is a mock K/\mathfrak{m} .
- $\partial : \mathcal{O} \rightarrow D$ is a derivation.

The theory of diffeovalued fields is first-order.

Definition 8.6. A *normalization* of a diffeovalued field is a choice of an isomorphism $D \cong K/\mathfrak{m}$ (respecting the embedding $k \hookrightarrow D$).

Every sufficiently resplendent diffeovalued field admits a normalization, by Proposition 8.2.

Definition 8.7. A *normalized diffeovalued field* is a diffeovalued field with a choice of a normalization.

Equivalently, a normalized diffeovalued field is a valued field (K, \mathcal{O}) with a derivation $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$.

Definition 8.8. Let K be a normalized diffeovalued field. A *lifting* is a derivation $\delta : K \rightarrow K$ such that

$$\partial x = (\delta x) + \mathfrak{m}$$

for $x \in \mathcal{O}$.

Proposition 8.9. *Let K be a sufficiently resplendent diffeovalued field. Suppose the value group Γ is p -divisible for at least one prime. Then K admits a normalization and a lifting.*

Proof. The normalization comes from Proposition 8.2, and the lifting comes from Corollary A.21 in the appendix. \square

Some assumption on the value group is necessary: Proposition A.22 in the appendix gives an example of a normalized diffeovalued field which cannot be lifted, even after passing to an elementary extension.

Definition 8.10. A *lifted diffeovalued field* is a normalized diffeovalued field with a choice of a lifting.

Equivalently, a lifted diffeovalued field is a field with a derivation and a valuation.

Lemma 8.11. *Let K be a diffeovalued field. If $x \in \mathcal{O}^\times$, then $\partial(x^{-1}) = -x^{-2}\partial x$.*

Proof. This follows as usual from

$$0 = \partial(xx^{-1}) = x^{-1}\partial x + x\partial(x^{-1}). \quad \square$$

8.3 Dense diffeovalued fields

Definition 8.12. A diffeovalued field is *dense* if for every $x \in D$, the fiber

$$\{y \in \mathcal{O} : \partial y = x\}$$

is dense in \mathcal{O} , with respect to the valuation topology.

The theory of dense diffeovalued fields is first-order.

Remark 8.13. Denseness implies that the value group Γ is non-trivial.

Proof. If the valuation is trivial, then $\mathcal{O} = K$ and the valuation topology is discrete. Then every fiber

$$\{y \in K : \partial y = x\}$$

is dense in \mathcal{O} , hence equal to \mathcal{O} . This is absurd unless D is a singleton. But D contains a submodule isomorphic to $k \cong K$. \square

8.4 The diffevaluation topology

Fix a *dense* diffeovalued field K . Define

$$\begin{aligned} R &= \{x \in K : \text{val}(x) \geq 0 \text{ and } \text{val}(\partial x) \geq 0\} \\ Q &= \{x \in K : \text{val}(x) \geq 0 \text{ and } \text{val}(\partial x) > 0\} \\ I &= \{x \in K : \text{val}(x) > 0 \text{ and } \text{val}(\partial x) > 0\}. \end{aligned}$$

Note that in the definition of Q, I ,

$$\text{val}(\partial x) > 0 \iff \partial x = 0,$$

because D is a mock K/\mathfrak{m} . In fact, Q is merely the kernel of $\partial : \mathcal{O} \rightarrow D$.

Lemma 8.14.

1. R, Q are proper subrings of K .
2. I is a proper ideal in R and in Q .
3. $\text{Frac}(Q) = \text{Frac}(R) = K$.
4. Q is a local ring with maximal ideal I .
5. $I \neq 0$.

Proof.

1. Easy. Properness holds because $R, Q \subseteq \mathcal{O}$, and \mathcal{O} is a proper subring by Remark 8.13.
2. Easy. Properness holds because $1 \notin I$.
3. As $Q \subseteq R$, it suffices to show $\text{Frac}(Q) = K$. Given $a \in K$, we must show $a \in \text{Frac}(Q)$. Replacing a with a^{-1} , we may assume $a \in \mathcal{O}$. If $a \in Q$, we are done. Otherwise, $\text{val}(\partial a) \leq 0$. By density, there is b such that

$$\begin{aligned} \text{val}(b) &> -\text{val}(\partial a) \\ b &\neq 0 \\ \partial b &= 0. \end{aligned}$$

Then $b \in Q$, and so $ab \in \mathcal{O}$. Also,

$$\partial(ab) = a\partial b + b\partial a = b\partial a = 0.$$

because $\partial b = 0$, and $\text{val}(b)$ is high enough for b to annihilate ∂a . Thus $ab \in \mathcal{O}$ and $a = (ab)/b \in \text{Frac}(Q)$.

4. Since I is a proper ideal in Q , it suffices to show

$$x \in Q \setminus I \implies x^{-1} \in Q.$$

Suppose $x \in Q \setminus I$, so that $\text{val}(x) = 0$ and $\partial x = 0$. Then $x^{-1} \in \mathcal{O}$, and

$$\partial(x^{-1}) = -x^{-2}\partial x = 0,$$

by Lemma 8.11. So $x^{-1} \in Q$.

5. By (4), Q/I is a field. If $I = 0$, then Q is a field, and $Q = K$ by (3). This contradicts (1). \square

Proposition 8.15. *Let K be a dense diffeovalued field. There is a locally bounded field topology on K characterized by the fact that either of the following are a neighborhood basis of 0:*

$$\begin{aligned} &\{aR : a \in K^\times\} \\ &\{aQ : a \in K^\times\}. \end{aligned}$$

The topology is locally bounded, non-discrete, and Hausdorff.

Proof. The ring Q induces a locally bounded, non-discrete, Hausdorff field topology because Q is a proper local subring of $K = \text{Frac}(Q)$. For example, see Theorem 2.2(b) in [11]. If a is a non-zero element of I , then

$$R \cdot a \subseteq R \cdot I = I \subseteq Q,$$

so R and Q induce the same topology. \square

Definition 8.16. The *diffeoevaluation topology* is the topology induced by Q or R as in Proposition 8.15.

Proposition 8.17. *The diffeoevaluation topology is not a V -topology.*

Proof. In a V -topology, the following local sentence holds, where U, V range over neighborhoods of 0:

$$\forall U \exists V \forall x, y : ((xy \in V) \rightarrow (x \in U \text{ or } y \in U)).$$

However, this fails for $U = R$. Indeed, suppose $V = aR$ is such that

$$xy \in aR \implies (x \in R \vee y \in R).$$

Shrinking the set aR , we may assume $a \in R$. By density, there is x such that

$$\begin{aligned}\text{val}(x) &> \text{val}(a) \\ \text{val}(\partial x) &< 0.\end{aligned}$$

Let $y = a/x$. Then

$$xy = a \in aR.$$

On the other hand, $x \notin R$ by choice of $\text{val}(\partial x)$, and $y \notin \mathcal{O} \supseteq R$, by choice of $\text{val}(x)$. \square

Definition 8.18. A *DV-topology* is a field topology that is locally equivalent to a dense diffevaluation topology.

8.5 Lifted diffeovalued fields

For lifted diffeovalued fields, we can characterize density and the diffevaluation topology more naturally.

Proposition 8.19. *Let (K, δ, val) be a lifted diffeovalued field. Then K is dense if and only if for every $a, b \in K$ and every $\gamma \in \Gamma$, there is $x \in K$ such that*

$$\begin{aligned}\text{val}(x - a) &> \gamma \\ \text{val}(\delta x - b) &> \gamma\end{aligned}$$

Proof. Unwinding the definition, density says that we can solve equations of the form

$$\begin{aligned}\text{val}(x - a) &> \gamma \\ \text{val}(\delta x - b) &> 0\end{aligned}$$

when $a \in \mathcal{O}$. So the listed conditions certainly imply density. Conversely, suppose density holds.

Claim 8.20. For any $\gamma \in \Gamma$ and $b \in K$, there is x such that

$$\begin{aligned}\text{val}(x) &> \gamma \\ \text{val}(\delta x - b) &> \gamma.\end{aligned}$$

Proof. Take some non-zero a such that $\text{val}(a) > \gamma$.

By density, there is y such that

$$\begin{aligned}\text{val}(y) &> \max(\gamma - \text{val}(\delta a), 0) \\ \text{val}(\delta y - b/a) &> 0.\end{aligned}$$

Let $x = ya$. Then

$$\begin{aligned}\text{val}(x) &= \text{val}(ya) = \text{val}(y) + \text{val}(a) > 0 + \gamma \\ \text{val}(a\delta y - b) &> \text{val}(a) > \gamma.\end{aligned}$$

Also,

$$\text{val}(y\delta a) = \text{val}(y) + \text{val}(\delta a) > \gamma.$$

So we see that

$$\text{val}(\delta x - b) = \text{val}(y\delta a + (a\delta y - b)) > \gamma.$$

□Claim

Now given any a, b, γ , we can find ε such that

$$\begin{aligned} \text{val}(\varepsilon) &> \gamma \\ \text{val}(\delta\varepsilon + \delta a - b) &> \gamma. \end{aligned}$$

Set $x = a + \varepsilon$. Then

$$\begin{aligned} \text{val}(x - a) &= \text{val}(\varepsilon) > \gamma \\ \text{val}(\delta x - b) &= \text{val}(\delta\varepsilon + \delta a - b) > \gamma. \end{aligned}$$

□

Proposition 8.21. *If (K, δ, val) is a dense, lifted diffeovalued field, then sets of the form*

$$\{x \in K : \text{val}(x - a_1) > \gamma_1 \text{ and } \text{val}(\delta x - a_2) > \gamma_2\} \text{ for } \gamma_1, \gamma_2 \in \Gamma, a_1, a_2 \in K$$

form a basis of opens in the diffeoevaluation topology.

Proof. Let $B_\gamma = \{x \in K : \text{val}(x) > \gamma \text{ and } \text{val}(\delta x) > \gamma\}$. It suffices to show that the B_γ form a neighborhood basis of 0. First of all, one sees by a straightforward calculation that B_γ is an R -submodule of K . The strong form of density in Proposition 8.19 can be used to show that B_γ is strictly bigger than $\{0\}$; for example take a, b very small relative to γ , and then take x with $x - a$ and $\delta x - b$ very small relative to a, b, γ . Thus B_γ is a neighborhood of 0 in the diffeoevaluation topology. Conversely, given any $a \in K^\times$, we claim that

$$B_\gamma \subseteq a^{-1} \cdot R$$

for sufficiently large γ . Indeed, if γ is sufficiently large relative to a , and $x \in B_\gamma$, then

$$\begin{aligned} \text{val}(ax) &= \text{val}(a) + \text{val}(x) \geq \text{val}(a) + \gamma \geq 0 \\ \text{val}(\delta(ax)) &= \text{val}(a\delta x + x\delta a) \geq \gamma + \min(\text{val}(a), \text{val}(\delta a)) \geq 0. \end{aligned}$$

□

8.6 Diffeoevaluation inflators

Fix a **dense** diffeovalued field K . Let D_0 be the image of the embedding $k \hookrightarrow D$, and let $\text{res}' : D_0 \rightarrow k$ be the inverse of this embedding. Let Q, R, I be as in §8.4. Note that if $a \in R$, then $\partial a \in D_0$, and so $\text{res}'(\partial a)$ makes sense.

Lemma 8.22.

1. *If $a \in \mathcal{O}$ and $b \in R$, then*

$$\text{res}'(a\partial b) = \text{res}(a) \text{res}'(\partial b).$$

2. The quotient Q/I is isomorphic (as a ring) to k via the map $\text{res}(-)$. Therefore, we can regard k -modules as Q -modules.
3. R and I are Q -submodules of K .
4. For $x \in R$, let $\widehat{\text{res}}(x) = (\text{res}(x), \text{res}'(\partial x)) \in k^2$. Then $\widehat{\text{res}}$ induces an isomorphism of Q -modules from R/I to k^2 . In particular, R/I is a semisimple Q -module of length 2.

Proof. 1. This holds because res' is an \mathcal{O} -linear map from D_0 to k , and the \mathcal{O} -module structure on k comes from $\text{res} : \mathcal{O} \rightarrow k$.

2. The ring homomorphism $\text{res} : Q \rightarrow k$ is onto, by density. Indeed, given any $x \in k$, we can find $y \in \mathcal{O}$ such that

$$\begin{aligned}\text{res}(x) &= y \\ \partial x &= 0.\end{aligned}$$

Then $x \in Q$ and $\text{res}(x) = y$. The kernel of $\text{res} : Q \rightarrow k$ is I , by definition of Q and I .

3. R is a Q -module because R is a superring of Q . I is a Q -module because I is an ideal in R .
4. The map $\widehat{\text{res}} : R \rightarrow k^2$ is obviously \mathbb{Z} -linear. It is surjective by density. The kernel is I , by definition of R and I . For Q -linearity, suppose $x \in Q$ and $y \in R$. Then

$$\text{res}(xy) = \text{res}(x) \text{res}(y).$$

Also, $x \in Q$ implies $\partial x = 0$, and so

$$\text{res}'(\partial(xy)) = \text{res}'(x\partial y + y\partial x) = \text{res}'(x\partial y) = \text{res}(x) \text{res}'(\partial y)$$

by part (1). Thus

$$\widehat{\text{res}}(xy) = \text{res}(x) \cdot (\text{res}(y), \text{res}'(\partial y)) = \text{res}(x) \widehat{\text{res}}(y). \quad \square$$

Lemma 8.23. *For any $a, b, c \in K$, the Q -submodule generated by $\{a, b, c\}$ is generated by a two-element subset of $\{a, b, c\}$.*

Proof. Without loss of generality, $\text{val}(c) \leq \text{val}(a)$ and $\text{val}(c) \leq \text{val}(b)$. Rescaling, we may assume $c = 1$. Then $a, b \in \mathcal{O}$.

Without loss of generality, $\text{val}(\partial a) \leq \text{val}(\partial b)$. Now, if $\text{val}(\partial b) > 0$, then $b \in Q \cdot 1$, and we are done. So we may assume

$$\text{val}(\partial a) \leq \text{val}(\partial b) \leq 0.$$

Take $x_0 \in \mathcal{O}$ such that $\partial b = x_0 \partial a$. By density, there is some x such that $\partial x = 0$ and

$$\text{val}(x - x_0) > -\text{val}(\partial a) \geq 0.$$

Note that $\text{val}(x - x_0) \geq 0$, so $x - x_0 \in \mathcal{O}$. As $x_0 \in \mathcal{O}$, we see $x \in \mathcal{O}$. Then $x \in Q$.

Now $\text{val}(x - x_0)$ is large enough that $x - x_0$ annihilates ∂a , so

$$x\partial a = x_0\partial a = \partial b.$$

Then

$$\partial(xa) = x\partial a + a\partial x = \partial b + 0.$$

Let $y = b - xa$. Then

$$\partial y = \partial b - \partial(xa) = 0.$$

Also, $b, x, a \in \mathcal{O}$, and so $y \in \mathcal{O}$. Thus $y \in Q$. Then

$$b = xa + y \in Q \cdot a + Q \cdot 1. \quad \square$$

Theorem 8.24. *Let K be a dense diffeovalued field. There is a malleable \mathbb{Q} -linear 2-inflator*

$$\begin{aligned} \text{Dir}_K(K) &\rightarrow \text{Dir}_k(k^2) \\ \text{Sub}_K(K^n) &\rightarrow \text{Sub}_k(k^{2n}) \\ V &\mapsto \{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : \vec{x} \in V \cap R^n\}, \end{aligned}$$

where $\widehat{\text{res}}(x) = (\text{res}(x), \text{res}'(\partial x))$.

Proof. By Proposition 7.3 and Lemma 8.23, the reduced rank of $\text{Sub}_Q(K)$ is at most 2. As R/I is a semisimple Q -module of length 2, we see that $\text{Sub}_Q(K)$ has reduced rank exactly two. Let

- \mathcal{C} the category of Q -modules.
- $F : K \text{ Vect} \rightarrow Q \text{ Mod}$ the forgetful functor.
- $G : Q \text{ Mod} \rightarrow \mathbb{Q} \text{ Vect}$ the forgetful functor.

Then Assumptions 8.1 and 8.11 of [10] hold. Applying Propositions 8.9 and 8.12 in [10], we obtain a malleable 2-inflator

$$\begin{aligned} \text{Dir}_K(K) &\rightarrow \text{Dir}_Q(R/I) \\ \text{Sub}_K(K^n) &\rightarrow \text{Sub}_Q((R/I)^n) \\ V &\mapsto (V \cap R^n + I^n)/I^n. \end{aligned}$$

Now $(V \cap R^n + I^n)/I^n$ can be described as the image of $V \cap R^n$ under the projection $R^n \twoheadrightarrow (R/I)^n$. Under the isomorphism $\text{Dir}_Q(R/I) \cong \text{Dir}_Q(k^2) \cong \text{Dir}_k(k^2)$, this is exactly

$$\{(\widehat{\text{res}}(x_1), \dots, \widehat{\text{res}}(x_n)) : \vec{x} \in V \cap R^n\}. \quad \square$$

Definition 8.25. A *diffeoevaluation inflator* on K is a 2-inflator on K arising from a dense diffeoevaluation on K via Theorem 8.24.

8.7 Characterization of diffevaluation inflators

We can summarize §3–5 as follows:

Theorem 8.26. *Let ς be an isotypic, malleable 2-inflator on a field K of characteristic 0. If no mutation of ς is weakly multi-valuation type, then ς is a diffevaluation inflator.*

Proof. By Corollary 4.5, we have a valuation ring \mathcal{O} . By Propositions 5.23 and 5.24, we have an \mathcal{O} -module D and a derivation $\partial : \mathcal{O} \rightarrow D$. By Propositions 5.29, 5.30, and 5.31 (with Proposition 4.9), D is a mock K/\mathfrak{m} , and the sets R, I, Q of §3–5 are exactly

$$\begin{aligned} R &= \{x \in \mathcal{O} : \text{val}(x) \geq 0\} \\ I &= \{x \in \mathfrak{m} : \partial x = 0\} \\ Q &= \{x \in \mathcal{O} : \partial x = 0\}. \end{aligned}$$

Thus we have a diffevaluation, and the sets R, I, Q agree with the ones defined in §8.4.

The map $\partial : \mathcal{O} \rightarrow D$ is surjective, by its construction in §5.5. Proposition 5.22 says that Q is dense in \mathcal{O} . Every other fiber of ∂ is a translate of Q , by surjectivity. Therefore every fiber is dense. So the diffevaluation data is dense.

Finally, by Propositions 3.8 and 5.31, ς has the same form as the diffevaluation inflator constructed in Theorem 8.24. \square

Under the Weak Assumptions of §2–5, we can say the following:

Corollary 8.27. *Let ς be a malleable 2-inflator on a field K of characteristic 0. Then some mutation ς' of ς is either weakly multi-valuation type, or a diffevaluation inflator.*

Proof. If no mutation of ς is weakly multi-valuation type, then ς satisfies the Weak Assumptions of §2. By Corollary 2.3, there is some mutation ς' which is isotypic. By Remark 2.1, ς' satisfies the Strong Assumptions of §3–5, and so ς' is a diffevaluation inflator by Theorem 8.26. \square

We can use Theorem 8.26 to characterize diffevaluation inflators. We first need some lemmas.

Lemma 8.28. *Let K be a field of characteristic 0 and $\mathcal{O}_1, \dots, \mathcal{O}_n$ be some valuation rings on K . Let b be an element of K . Then there is non-zero $q \in \mathbb{Q}$ such that $1/(b-q) \in \mathcal{O}_1 \cap \dots \cap \mathcal{O}_n$.*

Proof. Let \mathfrak{m}_i be the maximal ideal of \mathcal{O}_i . We need non-zero q such that $b - q \notin \mathfrak{m}_i$ for $1 \leq i \leq n$. For each i , let

$$B_i = \{q \in \mathbb{Q} : b - q \in \mathfrak{m}_i\}.$$

We must show that $B_1 \cup \dots \cup B_n \cup \{0\}$ fails to cover all of \mathbb{Q} . There are three possibilities for each B_i :

- If $b \notin \mathbb{Q} + \mathfrak{m}_i$, then B_i is empty.
- Otherwise, if \mathcal{O}_i has residue characteristic 0, then B_i is a singleton.

- Otherwise, if \mathcal{O}_i has residue characteristic $p > 0$, then B_i is a p -adic ball in \mathbb{Q} (of radius $1/p$).

We can take $q = 1/n!$, for $n \gg 0$. □

Lemma 8.29. *Let K be a dense diffeovalued field. As usual, let*

$$R = \{x \in \mathcal{O} : \text{val}(\partial x) \geq 0\}.$$

Let S be a multi-valuation ring on K . If $aS \subseteq R$, then $a = 0$.

Proof. Note that $a = a \cdot 1 \in a \cdot S \subseteq R$, so $a \in R \subseteq \mathcal{O}$.

Suppose $a \neq 0$. By density, we can find $b \in K$ such that

$$\begin{aligned} \text{val}(b) &> 0 \\ \text{val}(\partial b) &< \min(\text{val}(\partial a), 0) - \text{val}(a) \end{aligned}$$

Then $b \in \mathfrak{m}$. By Lemma 8.28, there is non-zero $q \in \mathbb{Q}$ such that $1/(b - q) \in S$. Then

$$\frac{a}{b - q} \in aS \subseteq R.$$

Now $b - q \in \mathcal{O}^\times$ because \mathcal{O} is equicharacteristic 0 and $q \neq 0$. By Lemma 8.11,

$$\partial \left(\frac{1}{b - q} \right) = \frac{-\partial b}{(b - q)^2},$$

and

$$\text{val} \left(\frac{-a\partial b}{(b - q)^2} \right) = \text{val}(a) + \text{val}(\partial b),$$

because the right hand side is less than 0. Then

$$\partial \left(\frac{a}{b - q} \right) = \frac{\partial a}{b - q} + \frac{-a\partial b}{(b - q)^2}.$$

But

$$\begin{aligned} \text{val} \left(\frac{\partial a}{b - q} \right) &= \text{val}(\partial a) \\ \text{val} \left(\frac{-a\partial b}{(b - q)^2} \right) &= \text{val}(a) + \text{val}(\partial b) < \text{val}(\partial a). \end{aligned}$$

So

$$\text{val} \left(\partial \left(\frac{a}{b - q} \right) \right) = \text{val}(a) + \text{val}(\partial b) < 0.$$

So $a/(b - q) \notin R$, a contradiction. □

Proposition 8.30. *Let K be a dense diffeovalued field, and let $\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_k(k^2)$ be the induced 2-inflator. Then no mutation of ς is weakly multi-valuation type.*

Proof. Recall that ς is the 2-inflator induced by the pedestal I in $\text{Sub}_Q(K)$. Suppose ς' is the mutation of ς along the line $K \cdot (a_1, a_2, \dots, a_m)$. By Proposition 10.15 in [10], ς' is the 2-inflator induced by the pedestal

$$I' = a_1^{-1}I \cap \dots \cap a_m^{-1}I.$$

Note that I' is non-zero, because it is open in the diffeoevaluation topology on K .

By Proposition 8.10 in [10], the fundamental ring R' of ς' is the “stabilizer”

$$R' = \{x \in K : xI' \subseteq I'\}.$$

Suppose for the sake of contradiction that ς' is weakly multi-valuation type. Then there is a multivaluation ring S on K such that R' contains a non-zero S -module. So there is some non-zero $a \in K$ such that

$$a \cdot S \subseteq R'.$$

Let b be a non-zero element in I' . Choose i so that $a_i \neq 0$. Then

$$a_i \cdot b \cdot a \cdot S \subseteq a_i \cdot b \cdot R' \subseteq a_i \cdot I' \subseteq I \subseteq R.$$

By Lemma 8.29, $a_i b a = 0$, which is absurd. □

Theorem 8.31. *Let ς be a 2-inflator on a field K of characteristic 0. Then ς is a diffeoevaluation inflator if and only if the following conditions hold:*

- ς is malleable.
- ς is isotypic.
- No mutation of ς is weakly multi-valuation type.

Proof. If the listed properties hold, then ς is a diffeoevaluation inflator by Theorem 8.26. Conversely, suppose $\varsigma : \text{Dir}_K(K) \rightarrow \text{Dir}_k(k^2)$ is a diffeoevaluation inflator. Then ς is plainly isotypic, and malleable by Theorem 8.24. The final property holds by Proposition 8.30. □

9 The canonical topology in characteristic 0

Let $(\mathbb{K}, +, \cdot, \dots)$ be a field, possibly with extra structure. Assume

- \mathbb{K} is sufficiently resplendent
- $\text{dp-rk}(\mathbb{K}) \leq 2$ and $\text{char}(\mathbb{K}) = 0$.
- \mathbb{K} is unstable, and the canonical topology is not a V-topology.

Theorem 9.1. *There is a valuation $\text{val} : \mathbb{K} \rightarrow \Gamma$ and a derivation $\delta : \mathbb{K} \rightarrow \mathbb{K}$ such that for every $a, b \in \mathbb{K}$ and $\gamma \in \Gamma$, the set*

$$\{x \in \mathbb{K} : \text{val}(x - a) > \gamma \text{ and } \text{val}(\delta x - b) > \gamma\}$$

is non-empty, and these sets form a basis for the canonical topology on \mathbb{K} .

Proof. By respndence and the uniform definability of the canonical topology (Theorem 6.27), we may replace \mathbb{K} with an elementarily equivalent field K . By Propositions 8.19 and 8.21, it suffices to produce a dense, lifted diffevaluation structure on K such that the diffevaluation topology agrees with the canonical topology.

Take a magic subfield $k_0 \preceq \mathbb{K}$. By Corollary 4.6 in [1], there is a prime p such that the embedding

$$k_0^\times / (k_0^\times)^p \hookrightarrow \mathbb{K}^\times / (\mathbb{K}^\times)^p$$

is an isomorphism, and so $\mathbb{K}^\times = k_0^\times \cdot (\mathbb{K}^\times)^p$. Thus $\mathbb{K}^\times / k_0^\times$ is p -divisible.

By Theorem 6.13, there is a k_0 -linear 2-inflator ς on \mathbb{K} satisfying the Strong Assumptions of §3–5. By Theorem 6.26, its fundamental ring R induces the canonical topology on \mathbb{K} . By Theorem 8.26 (and its proof), ς is the 2-inflator induced by some diffevaluation data $(\mathcal{O}, D, \partial)$, and

$$R = \{x \in \mathcal{O} : \text{val}(\partial x) \geq 0\}.$$

Thus, the canonical topology agrees with the diffevaluation topology.

Note that R is a k_0 -algebra, and therefore $k_0 \subseteq R \subseteq \mathcal{O}$. So the value group $\mathbb{K}^\times / \mathcal{O}^\times$ is a quotient of $\mathbb{K}^\times / k_0^\times$, and is p -divisible.

Let \mathbb{K}^+ be the expansion of \mathbb{K} by the diffevaluation data. By Theorem 6.27, there is a sentence σ holding in \mathbb{K}^+ , expressing that

- the diffevaluation is dense
- the value group is p -divisible
- the diffevaluation topology agrees with the canonical topology (of the reduct).

Let K be a sufficiently resplendent elementary extension of \mathbb{K}^+ . Then σ holds in K , and K admits a lifting, by Proposition 8.9. □

Recall from Definition 8.18 that a *DV-topology* is a field topology that is “locally equivalent” in the sense of [11] to a diffevaluation topology on a dense diffevalued field.

Corollary 9.2. *If K is a field of dp -rank 2 and characteristic 0, then one of the following holds:*

- K is stable.
- The canonical topology on K is a V -topology.
- The canonical topology on K is a DV -topology.

Proof. Theorem 9.1 and Theorem 6.27.2. □

10 Counterexample to the valuation conjecture

As outlined in §10 of [9], it would be very helpful if the Valuation Conjecture 1.2 were true. Unfortunately, algebraically closed dense diffeovalued fields turn out to be a counterexample, as hinted by Theorem 9.1 and Corollary 9.2.

Theorem 10.1. *Let ADVF be the theory of algebraically closed, dense diffeovalued fields of residue characteristic 0. Then ADVF is consistent, complete, unstable, has dp-rank 2, and is not valuation type.*

This will take some work to prove. In order to get a cleaner quantifier elimination result, it helps to work in a slightly different theory expanding ADVF.

Definition 10.2. Let $(K, \mathcal{O}, \mathfrak{m})$ be a valued field of residue characteristic 0.

- Let M be an \mathcal{O} -module. An M -valued log derivation on K is a group homomorphism

$$\partial \log : K^\times \rightarrow (M, +)$$

such that

$$(x + y)\partial \log(x + y) = x\partial \log x + y\partial \log y$$

for $x, y \in \mathcal{O}$.

- A *truncated log derivation* on K is a log derivation taking values in K/\mathfrak{m} .

If $\partial \log : K^\times \rightarrow (M, +)$ is a log derivation, and we define $\partial x = x \cdot \partial \log x$ for $x \in \mathcal{O}$, then $\partial : \mathcal{O} \rightarrow M$ is a derivation.

Definition 10.3. *LDVF* is the theory of $(K, \text{val}, \partial \log)$, where

- $(K, \text{val}) \models \text{ACVF}_{0,0}$
- $\partial \log$ is a truncated log derivation on K .
- Every fiber of $\partial \log$ is dense in K , with respect to the valuation topology.

Remark 10.4. The notion of “log derivation” used here is probably related to a standard construction of *logarithmic differentials* in log geometry. Specifically, an M -valued log derivation on \mathcal{O} is probably the same thing as an \mathcal{O} -linear morphism $\Omega_{\mathcal{O}/\mathbb{Q}}(\log \Gamma_{>0}) \rightarrow M$, where the module of log differentials $\Omega_{\mathcal{O}/\mathbb{Q}}(\log \Gamma_{>0})$ is as defined in §6.4.14 of [2]. I have not traced through the definitions to verify this.

10.1 Consistency

Recall that if L/K is an extension of fields of characteristic 0, if V is an L -vector space, and if $\partial : K \rightarrow V$ is a derivation, then we can extend ∂ to a derivation $\partial' : L \rightarrow V$. In the case where $L = K(t)$ (a pure transcendental extension), we can arrange for $\partial't$ to equal any value we want in V .

Lemma 10.5. *There is an algebraically closed dense, lifted diffeovalued field. In other words, there is an algebraically closed field K of characteristic 0, a non-trivial valuation $\text{val} : K \rightarrow \Gamma$ with residue characteristic 0, and a derivation $\partial : K \rightarrow K$ such that for any $a, b \in K$ and $\gamma \in \Gamma$, there is $x \in K$ such that*

$$\begin{aligned}\text{val}(x - a) &\geq \gamma \\ \text{val}(\partial x - b) &\geq \gamma.\end{aligned}$$

(Compare with Proposition 8.19.)

Proof. Choose some extension of the t -adic valuation on $\mathbb{Q}(t)$ to $\mathbb{Q}(t)^{alg}$. Let K be the completion of $\mathbb{Q}(t)^{alg}$. Then K is an algebraically closed field with a complete rank 1 valuation of residue characteristic 0. Moreover, the valuation topology on K is metrizable, separable, and complete. As $(K, +)$ is a non-discrete topological group, it has no isolated points. As K is a perfect Polish space, it is uncountable.

Let $\{U_i \times V_i\}_{i \in \mathbb{N}}$ be a countable basis of opens in $K \times K$. Let $K_0 \preceq K$ be a countable elementary substructure which defines the U_i and V_i , and is dense in K . Recursively choose $t_i, s_i \in K$ such that

- $t_i \in U_i$ and $s_i \in V_i$
- t_i is transcendental over $K_0(t_0, t_1, \dots, t_{i-1})$.

This is possible because $K_0(t_0, t_1, \dots, t_{i-1})$ is countable, so at least one transcendental $t' \in K$ exists. Replacing t' with its inverse, we may arrange for $t' \in \mathcal{O}$. Then, K_0 -definability of U_i ensures that there are $a, b \in K_0^\times$ such that $a \cdot \mathcal{O} + b \subseteq U_i$. Take $t_i = at' + b$.

Take the trivial derivation $K_0 \rightarrow K$ and extend it successively to $K_0(t_0)$, $K_0(t_0, t_1)$, \dots , arranging for $\partial t_i = s_i$. This determines a derivation $K(t_0, t_1, \dots) \rightarrow K$, which we can then extend to a derivation $K \rightarrow K$. The collection of t_i witnesses the required density statement. \square

Lemma 10.6. *Let (K, val, δ) be an algebraically closed, dense, lifted diffeovalued field. Let $\partial \log : K^\times \rightarrow K/\mathfrak{m}$ be the composition*

$$K^\times \rightarrow K \twoheadrightarrow K/\mathfrak{m},$$

where the first map is the usual log derivation $x \mapsto (\delta x)/x$, and the second map is the quotient map $x \mapsto x + \mathfrak{m}$. Then $\partial \log$ is a truncated log derivation, and $(K, \text{val}, \partial \log)$ is a model of LDVF.

Proof. If we set $\partial x = x \cdot \partial \log x$ for $x \in \mathcal{O}$, then $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$ is exactly the composition

$$K \xrightarrow{\delta} K \twoheadrightarrow K/\mathfrak{m}.$$

Thus ∂ is a derivation, and $\partial \log$ is a truncated log derivation.

By choice of (K, val) , it is a model of $\text{ACVF}_{0,0}$. Finally, we verify the density axiom. Given $b \in K$, we must show that the fiber

$$\left\{ x \in K : \frac{\delta x}{x} \in b + \mathfrak{m} \right\}$$

is dense in K , or equivalently, dense in K^\times . Fix $a \in K^\times$ and γ in the value group. By continuity of division, there is γ' such that for any $x, y \in K$,

$$(\text{val}(x - a) > \gamma' \text{ and } \text{val}(y - ab) > \gamma') \implies \text{val}\left(\frac{y}{x} - b\right) > 0.$$

By choice of (K, val, δ) , there is x such that

$$\begin{aligned} \text{val}(x - a) &> \max(\gamma', \gamma) \\ \text{val}(\delta x - ab) &> \gamma'. \end{aligned}$$

Then

$$\begin{aligned} \text{val}(x - a) &> \gamma \\ \text{val}\left(\frac{\delta x}{x} - b\right) &> 0. \end{aligned}$$

Thus x is within γ of a , and $\partial \log x = b + \mathfrak{m}$. So $(K, \text{val}, \partial \log)$ is a model of LDVF. \square

As an immediate corollary,

Proposition 10.7. *The theory LDVF is consistent.*

10.2 Calculations in ACVF

Lemma 10.8. *Let K be a model of $\text{ACVF}_{0,0}$. Let S be a subset of K and S' be a subset of K/\mathfrak{m} . Let $D \subseteq K$ be definable over $S \cup S'$. Then one of the following holds:*

- D has interior.
- D is finite, and every element is field-theoretically algebraic over S .

Proof. Replacing K with an elementary extension, and we may assume that K is a monster model. We may assume S, S' are finite. For $A \subseteq K$, let A^{alg} denote the field-theoretic algebraic closure, i.e., the algebraic closure of the subfield generated by A .

The swiss cheese decomposition ensures that D has interior unless D is finite. So we may assume D is finite. Then $D \subseteq \text{acl}(S \cup S')$. Let $\{x_1, \dots, x_n\}$ enumerate the elements of S' . Enlarging S' , we may assume $x_1 = 0$. Let $\pi : K \rightarrow K/\mathfrak{m}$ be the quotient map. The fibers of π are infinite, and thus uncountable, by saturation of the monster. Therefore we can find $y_i, y'_i \in \pi^{-1}(x_i)$ such that

$$\begin{aligned} y_i &\notin (Sy_1y_2 \cdots y_{i-1})^{\text{alg}} \\ y'_i &\notin (Sy_1y_2 \cdots y_n y'_1 y'_2 \cdots y'_{i-1})^{\text{alg}}. \end{aligned}$$

Then the sequence $y_1, y_2, \dots, y'_1, y'_2, \dots, y'_n$ is a sequence of independent transcendentals over S^{alg} .

We arranged for $y_1, y'_1 \in \pi^{-1}(x_1) = \pi^{-1}(0) = \mathfrak{m}$. Therefore, y_1 and y'_1 have non-trivial valuation. Then

$$\begin{aligned} M &:= (Sy_1y_2 \cdots y_n)^{\text{alg}} \preceq K \\ M' &:= (Sy'_1y'_2 \cdots y'_n)^{\text{alg}} \preceq K, \end{aligned}$$

by model completeness of ACVF. Note that $x_i \in \text{dcl}^{\text{eq}}(y_i)$, and so $S' \subseteq \text{dcl}^{\text{eq}}(M)$. Then D is M -definable, and so $D \subseteq M$ because D is finite. Similarly, $D \subseteq M'$.

On the other hand, we arranged for the following to hold in the ACF reduct:

$$y_1y_2 \cdots y_n \underset{S}{\perp} y'_1y'_2 \cdots y'_n.$$

Therefore $M \cap M' = S^{\text{alg}}$, and so $D \subseteq S^{\text{alg}}$. □

Lemma 10.9. *Let K be a model of ACVF. Let $P(x) = a_nx^n + \cdots + a_1x + a_0$ be a polynomial such that $\min_{i \leq n}(\text{val}(a_i)) = 0$. Then the number of roots of P in \mathcal{O} , counted with multiplicities, is equal to the largest i such that $\text{val}(a_i) = 0$.*

Proof. This is a basic statement about Newton polygons. Let r_1, \dots, r_n be the roots of $P(x)$, counted with multiplicity. Reordering, we may assume $r_1, \dots, r_m \in \mathcal{O}$, and $r_{m+1}, \dots, r_n \notin \mathcal{O}$. Then

$$\begin{aligned} P(x) &= cQ(x) \\ Q(x) &= \prod_{i=1}^m (x - r_i) \cdot \prod_{i=m+1}^n (1 - x/r_i). \end{aligned}$$

for some $c \in K^\times$. Then $Q(x) \in \mathcal{O}[x]$, and its reduction modulo \mathfrak{m} is

$$\prod_{i=1}^m (x - \text{res}(r_i)),$$

a nonzero polynomial in $k[x]$, of degree m . If we write $Q(x) = b_nx^n + \cdots + b_1x + b_0$, then $\min_{i \leq n}(\text{val}(b_i)) = 0$, and so $\text{val}(c) = 0$. Then

$$\max\{i \leq n : \text{val}(a_i) = 0\} = \max\{i \leq n : \text{val}(b_i) = 0\} = m. \quad \square$$

Some form of Rolle's theorem holds in models of $\text{ACVF}_{0,0}$:

Lemma 10.10. *Let K be a model of $\text{ACVF}_{0,0}$. Let B be a ball. Let $P(x)$ be a polynomial in $K[x]$. If P has two distinct zeros in B , then $P'(x)$ has a zero in B .*

Proof. We may assume P is non-zero; otherwise the result is trivial. Let r_1, r_2 be two zeros in B . Shrinking B to the smallest ball containing r_1, r_2 , we may assume B is a closed ball. Shifting everything by an affine transformation, we may assume $B = \mathcal{O}$. Let $P(x) = a_n x^n + \cdots + a_1 x + a_0$. Multiplying P by a constant from K^\times , we may assume $\min_{i \leq n} \text{val}(a_i) = 0$. The polynomial $P(x)$ has at least two roots in \mathcal{O} , so by Lemma 10.9, there is some $m \geq 2$ such that $\text{val}(a_m) = 0$. Note

$$P'(x) = na_n x^{n-1} + \cdots + 2a_2 x + a_1.$$

Also, $\text{val}(ia_i) = \text{val}(a_i)$ for $i \geq 1$, because of residue characteristic 0. Therefore, $\text{val}(ia_i) \geq 0$, and $\text{val}(ma_m) = 0$. So the coefficient of x^{m-1} has valuation 0 for some $m \geq 2$. By Lemma 10.9, $P'(x)$ has at least $2 - 1$ roots in \mathcal{O} . \square

10.3 Calculations with log derivations

Lemma 10.11. *Let $(K, \mathcal{O}, \mathfrak{m})$ be a valued field, let M be an \mathcal{O} -module, and let $\partial \log : K^\times \rightarrow M$ be a log derivation. If $x, y \in K$ satisfy*

$$\text{val}(x - y) \leq \max(\text{val}(x), \text{val}(y)), \tag{10}$$

then $x/(x - y), y/(x - y) \in \mathcal{O}$, and

$$\partial \log(x - y) = \frac{x}{x - y} \cdot \partial \log(x) - \frac{y}{x - y} \cdot \partial \log(y).$$

Proof. First note that

$$\text{val}(x - y) \leq \min(\text{val}(x), \text{val}(y)).$$

This is automatic if $\text{val}(x) \neq \text{val}(y)$, and equivalent to (10) otherwise.

Recall the derivation $\partial : \mathcal{O} \rightarrow M$ given by $\partial x = x \cdot \partial \log x$. Then

$$\begin{aligned} 0 &= \partial(1) = \partial \left(\frac{x - y}{x - y} \right) = \partial \left(\frac{x}{x - y} \right) - \partial \left(\frac{y}{x - y} \right) \\ &= \frac{x}{x - y} \partial \log \left(\frac{x}{x - y} \right) - \frac{y}{x - y} \partial \log \left(\frac{y}{x - y} \right) \\ &= \frac{x}{x - y} [\partial \log(x) - \partial \log(x - y)] - \frac{y}{x - y} [\partial \log(y) - \partial \log(x - y)] \\ &= \left(\frac{x}{x - y} \cdot \partial \log(x) - \frac{y}{x - y} \cdot \partial \log(y) \right) - \left(\frac{x}{x - y} - \frac{y}{x - y} \right) \partial \log(x - y) \\ &= \left(\frac{x}{x - y} \cdot \partial \log(x) - \frac{y}{x - y} \cdot \partial \log(y) \right) - \partial \log(x - y). \end{aligned} \quad \square$$

Proposition 10.12. *Let (K, \mathcal{O}) be an algebraically closed field with a log derivation $\partial \log : K \rightarrow M$, for some \mathcal{O} -module M . Suppose that $\partial \log$ vanishes on some subfield $F \subseteq K$, and K is algebraic over F (so that $K = F^{alg}$). Then $\partial \log$ vanishes on K .*

Proof. Increasing F , we may assume F is maximal among subfields on which $\partial \log$ vanishes. Suppose for the sake of contradiction that $F \subsetneq K$. Take minimal $n > 1$ such that F has a finite extension of degree n . If $P(x) \in F[x]$ has degree $\leq n$, then one of the following happens:

- $P(x)$ factors into linear polynomials
- $P(x)$ is irreducible of degree n .

Claim 10.13. If $a \in K$ and $[F(a) : F] = n$, then

- $F(a)^\times$ is generated by F^\times and the elements $a - b$ with $b \in F$.
- There is $b \in F$ such that $\partial \log(a - b) \neq 0$.

Proof. Every element of $F(a)$ is of the form $P(a)$ for some polynomial $P(x) \in F[x]$ of degree less than n . Then P splits into linear factors, so

$$P(a) = c(a - b_1)(a - b_2) \cdots (a - b_n)$$

for some $c, b_1, b_2, \dots, b_n \in F$. This proves the first point. If $\partial \log(a - b) = 0$ for all $b \in F$, then $\partial \log$ must vanish on $F(a)^\times$, contradicting the choice of F . □_{Claim}

Take an arbitrary extension L/F of degree n , and break into cases:

- If $\text{val}(L)$ is strictly larger than $\text{val}(F)$, take $\gamma \in \text{val}(L) \setminus \text{val}(F)$. The inequality

$$|\text{val}(L)/\text{val}(F)| \leq [L : F]$$

implies that $m\gamma \in \text{val}(F)$ for some $m \leq n$. Take $c \in F$ with $\text{val}(c) = m\gamma$. The polynomial $x^m - c$ has no roots in F , so $m = n$ and $x^n - c$ is irreducible. Take $a \in K$ such that $a^n = c$. Note that

$$\partial \log(a) = (1/n)\partial \log(c) = 0,$$

because $c \in F$ and the residue characteristic is 0.

By Claim 10.13, there is $b \in F(a)$ such that $\partial \log(a - b) \neq 0$. Now $\text{val}(b) \neq \gamma = \text{val}(a)$, by choice of γ , and so

$$\text{val}(a - b) = \min(\text{val}(a), \text{val}(b)).$$

Also $\partial \log(a) = \partial \log(b) = 0$. By Lemma 10.11, $\partial \log(a - b) = 0$, a contradiction.

- If $\text{res}(L)$ is strictly larger than $\text{res}(F)$, take $\alpha \in \text{res}(L) \setminus \text{res}(F)$. The inequality

$$[\text{res}(L) : \text{res}(F)] \leq [L : F]$$

implies that $[\text{res}(F)(\alpha) : \text{res}(F)] \leq n$. Let

$$x^m + \beta_{m-1}x^{m-1} + \cdots + \beta_1x + \beta_0$$

be the monic irreducible polynomial of α over $\text{res}(F)$. Because of residue characteristic 0, this polynomial is separable, and so

$$m\alpha^{m-1} + (m-1)\beta_{m-1}\alpha^{m-2} + \cdots + 2\beta_2\alpha + \beta_1 \neq 0. \quad (11)$$

Take $b_i \in F$ with $\text{res } b_i = \beta_i$, and let $P(x)$ be the polynomial

$$x^m + b_{m-1}x^{m-1} + \cdots + b_1x + b_0 \in F[x].$$

Then $P(x)$ is irreducible, and so $m = [\text{res}(F)(\alpha) : \text{res}(F)] = n$. Let $a \in K$ be the root of $P(x)$ with $\text{res}(a) = \alpha$. Then

$$a^n + b_{n-1}a^{n-1} + \cdots + b_1a + b_0 = 0.$$

Applying the derivation $\partial : \mathcal{O} \rightarrow M$, which vanishes on the b_i , we obtain

$$(na^{n-1} + (n-1)b_{n-1}a^{n-2} + \cdots + 2b_2a + b_1)\partial a = 0.$$

The expression inside the parentheses has nonzero residue, by (11), and so it is an element of \mathcal{O}^\times . Therefore $\partial a = 0$. Now $\text{res}(a) = \alpha \notin \text{res}(F)$, so $\text{res}(a) \neq 0$ and a is invertible as well. Therefore $\partial \log a = (\partial a)/a = 0$.

By Claim 10.13, there is some $b \in F$ such that $\partial \log(a - b) \neq 0$. Then

$$\text{val}(a - b) \leq \max(\text{val}(a), \text{val}(b)).$$

(Otherwise, $\text{res}(a) = \text{res}(b) \in F$, contradicting the choice of a and α .) By Lemma 10.11, $\partial \log(a - b) = 0$, a contradiction.

- Lastly, suppose that L/F is an immediate extension. By maximality of F , there is $a \in L$ with $\partial \log(a) \neq 0$. Let \mathcal{C} be the collection of balls containing a , with center and radius from F . Let I be the intersection $\bigcap \mathcal{C}$. As usual, $I \cap F = \emptyset$. (Suppose $b \in I \cap F$. Then $rv(a - b) = rv(b' - b)$ for some $b' \in F$, because the extension is immediate. The ball centered around b' of radius $\text{val}(a - b')$ does not contain b , contradicting the choice of b .)

Let $P(x)$ be the minimal polynomial of a over F . Then $P(x)$ has degree n . Let a_1, \dots, a_n be the roots of $P(x)$, with $a_1 = a$. Note that $P'(x)$ has degree $n - 1$, and therefore splits over F . So no root of $P'(x)$ is in I . By Rolle's Theorem (Lemma 10.10), a is the unique root of $P(x)$ in I .

Therefore I has empty intersection with the finite set $\{0, a_2, \dots, a_n\}$. We can find $b \in F$ such that

$$\text{val}(a - b) > \max(\text{val}(0 - b), \text{val}(a_2 - b), \text{val}(a_3 - b), \dots, \text{val}(a_n - b)).$$

Take $c \in F$ with $\text{val}(a - b) = \text{val}(c)$, and let $e_i = (a_i - b)/c$. Then $\text{val}(e_1) = 0$, and $\text{val}(e_i) < 0$ for $i > 1$. The e_i are the roots of the irreducible polynomial

$$Q(x) = P(cx + b) = s_n x^n + s_{n-1} x^{n-1} + \dots + s_1 x + s_0 \in F[x].$$

By Newton polygons, $\text{val}(s_0) = \text{val}(s_1) < \text{val}(s_i)$ for $i > 1$. Then we can apply ∂ to the equation

$$(s_n/s_1)e_1^n + \dots + (s_2/s_1)e_1^2 + e_1 + (s_0/s_1) = 0,$$

and obtain

$$(n(s_n/s_1)e_1^{n-1} + \dots + 2(s_2/s_1)e_1 + 1)\partial e_1 = 0,$$

because the coefficients s_n/s_1 lie in F , where ∂ vanishes. But the expression in parentheses has valuation 0, because $e_1 \in \mathcal{O}$ and $s_i/s_1 \in \mathfrak{m}$ for $2 \leq i \leq n$. Therefore $\partial e_1 = 0$. As $\text{val}(e_1) = 0$, we have $e_1 \in \mathcal{O}^\times$ as well, and then $\partial \log e_1 = (\partial e_1)/e_1 = 0$. Then $\partial \log(a - b) = \partial \log e_1 + \partial \log c = 0$, as $c \in F$. Finally,

$$\text{val}(a - b) > \text{val}(b) = \text{val}(a),$$

and so $\partial \log(a - b)$ and $\partial \log(b)$ determine $\partial \log(a)$, by Lemma 10.11. Thus $\partial \log(a) = 0$, contradicting the choice of a . \square

Proposition 10.12 is probably a consequence of Lemma 6.5.12 and Claim 6.5.14 in [2], but I am not entirely certain.

10.4 Quantifier elimination and completeness

Let \mathcal{L}_0 be the language for $\text{ACVF}_{0,0}$ with two sorts, K and K/\mathfrak{m} , and the following functions and relations:

- The field operations on K , including the constants 0, 1, and *division*.
- The \mathcal{O} -module structure on K/\mathfrak{m} , i.e., the group structure (including 0 and negation) and the multiplication map

$$\mathcal{O} \times K/\mathfrak{m} \rightarrow K/\mathfrak{m},$$

understood as a partial function on $K \times K/\mathfrak{m}$.

- All \emptyset -definable relations on K and K/\mathfrak{m} .

Then $\text{ACVF}_{0,0}$ has quantifier elimination in \mathcal{L}_0 , because we Morleyized. If $K \models \text{ACVF}_{0,0}$, an \mathcal{L}_0 -substructure of K consists of a pair (F, D) , where

- F is a subfield of K .
- D is an \mathcal{O}_F -submodule of K/\mathfrak{m} .

Note that D need not contain the image of F under $K \rightarrow K/\mathfrak{m}$, as we did not include this map as one of the functions in the signature.

Let \mathcal{L} be the language for LDVF obtained by expanding \mathcal{L}_0 with a function symbol for the map $\partial \log : K^\times \rightarrow K/\mathfrak{m}$. If K is a model of LDVF, then an \mathcal{L} -substructure is a pair (F, D) , where

- F is a subfield of K
- D is an \mathcal{O}_F -submodule of K/\mathfrak{m}
- D contains $\partial \log x$ for $x \in F$.

Lemma 10.14. *Let K be a model of LDVF. Let K' be a $|K|^+$ -saturated model of LDVF. Let (F, D) be a proper \mathcal{L} -substructure of K . Let $f : (F, D) \hookrightarrow K'$ be an \mathcal{L} -embedding (an isomorphism onto a substructure of K'). Then f can be extended to an \mathcal{L} -embedding $f' : (F', D') \hookrightarrow K'$ for some strictly larger \mathcal{L} -substructure (F', D') .*

Proof. First suppose $D < K/\mathfrak{m}$. Note that $(F, K/\mathfrak{m})$ is an \mathcal{L} -substructure of K . By quantifier elimination of $\text{ACVF}_{0,0}$ in the language \mathcal{L}_0 , we can extend f to an \mathcal{L}_0 -embedding $f' : (F, K/\mathfrak{m}) \hookrightarrow K'$. Then f' is already an \mathcal{L} -embedding, because

$$f'(\partial \log(x)) = f(\partial \log(x)) = \partial \log(f(x)) = \partial \log(f'(x))$$

for any $x \in F$. (The first equation holds because $x \in F \implies \partial \log(x) \in D$, and f' extends f on D .)

So we may assume $D = K/\mathfrak{m}$, and $F < K$.

Claim 10.15. If F' is a subfield of K containing F , and $f' : (F', K/\mathfrak{m}) \hookrightarrow K'$ is an \mathcal{L}_0 -embedding extending f , then

- f' induces a map from the valuation ring of F' to the valuation ring of K' , and so we can regard K'/\mathfrak{m}' as a module over the valuation ring of F' .
- If $\Delta : F' \rightarrow K'/\mathfrak{m}'$ is defined by

$$\Delta(x) = f'(\partial \log(x)) - \partial \log(f'(x)),$$

then Δ is a log derivation $F' \rightarrow K'/\mathfrak{m}'$.

- Δ vanishes on F .
- If Δ vanishes on F' , then f' is an \mathcal{L} -embedding.

Proof. The first point is clear, since f' is a partial elementary map in the ACVF reduct. The second point is a direct calculation:

$$\begin{aligned}
\Delta(xy) &= f'(\partial \log(xy)) - \partial \log(f'(xy)) \\
&= f'(\partial \log(x) + \partial \log(y)) - \partial \log(f'(x)f'(y)) \\
&= f'(\partial \log(x)) + f'(\partial \log(y)) - \partial \log(f'(x)) - \partial \log(f'(y)) \\
&= \Delta(x) + \Delta(y) \\
(x+y)\Delta(x+y) &= f'(x+y)\Delta(x+y) \\
&= f'(x+y)f'(\partial \log(x+y)) - f'(x+y)\partial \log(f'(x+y)) \\
&= f'((x+y)\partial \log(x+y)) - (f'(x) + f'(y))\partial \log(f'(x) + f'(y)) \\
&= f'(x\partial \log x + y\partial \log y) - f'(x)\partial \log(f'(x)) - f'(y)\partial \log(f'(y)) \\
&= f'(x)f'(\partial \log x) + f'(y)f'(\partial \log y) - f'(x)\partial \log(f'(x)) - f'(y)\partial \log(f'(y)) \\
&= f'(x)\Delta(x) + f'(y)\Delta(y).
\end{aligned}$$

The third point expresses that f is an \mathcal{L} -embedding. The fourth point is clear. □_{Claim}

Next suppose $F \neq F^{alg}$. By quantifier elimination of $\text{ACVF}_{0,0}$, we can extend f to an \mathcal{L}_0 -embedding $f' : (F^{alg}, K/\mathfrak{m}) \hookrightarrow K'$. Let $\Delta : F^{alg} \rightarrow K'/\mathfrak{m}'$ be as in Claim 10.15. Then Δ vanishes on F , and therefore on F^{alg} , by Proposition 10.12. Therefore f' is an \mathcal{L} -embedding.

Finally, suppose that $F = F^{alg}$. Take a transcendental $a \in K \setminus F$. Let \vec{s} be an infinite tuple enumerating F , and \vec{t} be an infinite tuple enumerating K/\mathfrak{m} . Let $\Sigma(x; \vec{y}; \vec{z})$ be the complete \mathcal{L}_0 -type of $(a; \vec{s}; \vec{t})$. Note that $a' \in K'$ satisfies $\Sigma(x; f(\vec{s}); f(\vec{t}))$, if and only if there is an \mathcal{L}_0 -embedding $f' : (F(a), K/\mathfrak{m}) \hookrightarrow K'$ extending f and sending $a \mapsto a'$.

For any $b \in F$, let $\psi_b(x)$ be the type in K' asserting that

$$\partial \log(x - f(b)) = f(\partial \log(a - b)).$$

(The right hand side makes sense, because we arranged $D = K/\mathfrak{m}$.)

Claim 10.16. For any $b \in F$, the type $\Sigma(x; f(\vec{s}); f(\vec{t})) \cup \{\psi_b(x)\}$ is realized in K' .

Proof. By saturation, it suffices to show finite satisfiability. Suppose $\varphi(x; \vec{y}; \vec{z})$ is an \mathcal{L}_0 -formula satisfied by $(a; \vec{s}; \vec{t})$. We must find $a' \in K'$ satisfying

$$\varphi(a'; f(\vec{s}); f(\vec{t})) \wedge \psi_b(a')$$

The definable set $\varphi(K; \vec{s}; \vec{t})$ has interior, by Lemma 10.8 and transcendence of a over \vec{s} . As f is a partial elementary map in the ACVF reduct, the definable set $\varphi(K'; f(\vec{s}); f(\vec{t}))$ has interior as well. By the density axiom of LDVF, there is $x \in K'$ such that

$$\begin{aligned}
x + f(b) &\in \varphi(K'; f(\vec{s}); f(\vec{t})) \\
\partial \log(x) &= f(\partial \log(a - b)).
\end{aligned}$$

Take $a' = x + f(b)$. □_{Claim}

Claim 10.17. The type $\Sigma(x; f(\vec{s}); f(\vec{t})) \cup \{\psi_b(x) : b \in F\}$ is realized in K' .

Proof. By saturation, it suffices to show finite satisfiability. Let b_1, \dots, b_n be elements of F . We claim that the type

$$\Sigma(x; f(\vec{s}); f(\vec{t})) \cup \{\psi_{b_1}(x), \dots, \psi_{b_n}(x)\}$$

is realized in K' . Without loss of generality,

$$\text{val}(a - b_1) \geq \text{val}(a - b_2) \geq \dots \geq \text{val}(a - b_n).$$

By Claim 10.16 there is a' realizing $\Sigma(x; f(\vec{s}); f(\vec{t})) \cup \{\psi_{b_1}(x)\}$. Let $f' : (F(a), K/\mathfrak{m}) \rightarrow K'$ be the \mathcal{L}_0 -embedding extending f and sending a to a' . Note that

$$\partial \log(f'(a - b_1)) = \partial \log(f'(a) - f'(b_1)) = \partial \log(a' - f(b_1)) \stackrel{*}{=} f(\partial \log(a - b_1)) = f'(\partial \log(a - b_1)).$$

The starred equation holds because of $\psi_{b_1}(a')$. By Claim 10.15, there is a log derivation $F(a) \rightarrow K'/\mathfrak{m}'$ given by

$$\Delta(x) := f'(\partial \log(x)) - \partial \log(f'(x)).$$

Then $\Delta(a - b_1) = 0$. Also, $\Delta(b_i - b_1) = 0$ for any i , because $b_i - b_1 \in F$. Moreover,

$$\text{val}((a - b_1) - (b_i - b_1)) = \text{val}(a - b_i) \leq \text{val}(a - b_1),$$

and so $\Delta(a - b_i) = 0$, by Lemma 10.11. Then for each i ,

$$\partial \log(a' - f(b_i)) = \partial \log(f'(a) - f'(b_i)) = \partial \log(f'(a - b_i)) = f'(\partial \log(a - b_i)) = f(\partial \log(a - b_i)).$$

Therefore $\psi_{b_i}(a')$ holds. \square

Using Claim 10.17, take $a' \in K'$ realizing

$$\Sigma(x; f(\vec{s}); f(\vec{t})) \cup \{\psi_b(x) : b \in F\}.$$

Let $f' : (F(a), K/\mathfrak{m}) \hookrightarrow K'$ be the \mathcal{L}_0 -embedding extending f and mapping a to a' . Let $\Delta(x) = f'(\partial \log(x)) - \partial \log(f'(x))$ as in Claim 10.15. The statement $\psi_b(a')$ implies that

$$\partial \log(a' - f(b)) = f(\partial \log(a - b)).$$

Therefore

$$\partial \log(f'(a - b)) = \partial \log(f'(a) - f'(b)) = \partial \log(a' - f(b)) = f(\partial \log(a - b)) = f'(\partial \log(a - b)).$$

So $\Delta(a - b) = 0$ for any $b \in F$. As F is algebraically closed and a is transcendental, the multiplicative group $F(a)^\times$ is generated by

$$F^\times \cup \{a - b : b \in F\}.$$

Then Δ must vanish on $F(a)$, because it vanishes on the generators. By Claim 10.15, f' is an \mathcal{L} -embedding. \square

Theorem 10.18. LDVF has quantifier elimination in the language \mathcal{L} .

Proof. This follows from Lemma 10.14, by well-known model-theoretic techniques. \square

Corollary 10.19. LDVF is complete.

Proof. LDVF is consistent by Proposition 10.7. Take two models M_1 and M_2 , viewed as \mathcal{L} -structures. Note that $(\mathbb{Q}, 0)$ is an \mathcal{L} -substructure of M_i , for each i . The identity map $(\mathbb{Q}, 0) \rightarrow (\mathbb{Q}, 0)$ is an isomorphism of \mathcal{L} -structures:

- For the \mathcal{L}_0 -structure, this holds because $\text{ACVF}_{0,0}$ is complete.
- For the map $\partial \log$, this holds because $\partial \log$ is trivial on both copies of \mathbb{Q} .

By quantifier elimination, $M_1 \equiv M_2$. \square

Corollary 10.20. Let ADVF be the theory of algebraically closed dense diffeovalued fields (K, ∂, D) .

1. If $(K, \partial \log)$ is a model of LDVF, then $(K, \partial, K/\mathfrak{m})$ is a model of ADVF.
2. Up to elementary equivalence, every model of ADVF arises in this way.
3. ADVF is complete.

Proof. It suffices to prove Point 2. Let $(K, \partial, \text{val})$ be a model of ADVF. Passing to a resplendent elementary extension, we may assume that a lifting exists, by Proposition 8.9. So we obtain a lifted diffeovalued field (K, δ, val) . Define

$$\begin{aligned} \partial \log : K^\times &\rightarrow K/\mathfrak{m} \\ \partial \log(x) &= \frac{\delta x}{x} + \mathfrak{m}. \end{aligned}$$

Then $(K, \partial \log, \text{val}) \models \text{LDVF}$, by Lemma 10.6. The derivation $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$ determined by $\partial \log$ is the original derivation ∂ . \square

10.5 Upper bound on dp-rank

Let T be the theory of dense, lifted diffeovalued fields. By Proposition 8.19, a model of T is a field K with a non-trivial valuation (of residue characteristic 0) and a derivation $\delta : K \rightarrow K$ such that for any $a, b \in K$ and γ in the value group, there is $x \in K$ such that

$$\begin{aligned} \text{val}(x - a) &\geq \gamma \\ \text{val}(\delta x - b) &\geq \gamma. \end{aligned}$$

The theory T probably has no nice properties, other than being consistent.

Remark 10.21. Lemma 10.6 says that if $(K, \delta, \text{val}) \models T$, and we define

$$\partial \log(x) := \frac{\delta x}{x} + \mathfrak{m},$$

then $(K, \partial \log, \text{val}) \models \text{LDVF}$. As LDVF is complete, it follows that every sufficiently re-splendent model of LDVF can be expanded to a model of T .

Fix some one-sorted language for T . Say that $\varphi(\vec{x})$ is a *ACVF-formula* if $\varphi(\vec{x})$ is defined in the ACVF-reduct, and similarly for *LDVF-formulas*. Let $\text{tp}_{\text{ACVF}}(\vec{a}/B)$ be the set of all ACVF-formulas with parameters in B , satisfied by \vec{a} . Define $\text{tp}_{\text{LDVF}}(\vec{a}/B)$ similarly.

If $M \models T$ and \vec{a} is a tuple in M , define $\delta(\vec{a})$ coordinatewise.

Lemma 10.22. *Let M_1, M_2 be two models of T . Let \vec{a} be a tuple in M_1 and \vec{b} be a tuple of the same length in M_2 . Suppose that $\text{tp}_{\text{ACVF}}(\vec{a}\delta(\vec{a})/\emptyset) = \text{tp}_{\text{ACVF}}(\vec{b}\delta(\vec{b})/\emptyset)$. Then $\text{tp}_{\text{LDVF}}(\vec{a}/\emptyset) = \text{tp}_{\text{LDVF}}(\vec{b}/\emptyset)$.*

Proof. After replacing M_1 and M_2 with elementary extensions, there is an isomorphism of the ACVF-reducts

$$f : (M_1, \text{val}) \rightarrow (M_2, \text{val})$$

such that $f(\vec{a}) = \vec{b}$ and $f(\delta(\vec{a})) = \delta(\vec{b})$. With \mathcal{L}_0 as in the previous section, this induces an isomorphism of \mathcal{L}_0 -structures

$$f' : (M_1, M_1/\mathfrak{m}^{M_1}) \rightarrow (M_2, M_2/\mathfrak{m}^{M_2}).$$

Restricting the first sort, we obtain an \mathcal{L}_0 -isomorphism

$$f'' : (\mathbb{Q}(\vec{a}), M_1/\mathfrak{m}^{M_1}) \rightarrow (\mathbb{Q}(\vec{b}), M_2/\mathfrak{m}^{M_2}),$$

because $f(\vec{a}) = \vec{b}$. We claim that f'' preserves $\partial \log$. If c is in $\mathbb{Q}[\vec{a}]$, then

$$c = P(a_1, \dots, a_n)$$

for some $P(x_1, \dots, x_n) \in \mathbb{Q}[x_1, \dots, x_n]$. Then

$$\delta(c) = \sum_{i=1}^n \frac{\partial P}{\partial x_i}(\vec{a}) \cdot \delta(a_i).$$

Since f sends $\delta(a_i)$ to $\delta(b_i)$,

$$f(\delta(c)) = \sum_{i=1}^n \frac{\partial P}{\partial x_i}(\vec{b}) \cdot \delta(b_i) = \delta(P(\vec{b})) = \delta(f(c)).$$

Then $f(\delta(c)) = \delta(f(c))$, implying that $f'(\partial \log(c)) = \partial \log(f'(c))$. More generally, if $c \in \mathbb{Q}(\vec{a})$, then $c = c_1/c_2$ for $c_i \in \mathbb{Q}[\vec{a}]$, and

$$\begin{aligned} f'(\partial \log(c)) &= f'(\partial \log(c_1) - \partial \log(c_2)) = f'(\partial \log(c_1)) - f'(\partial \log(c_2)) \\ &= \partial \log(f'(c_1)) - \partial \log(f'(c_2)) = \partial \log(f'(c_1)/f'(c_2)) \\ &= \partial \log(f'(c_1/c_2)) = \partial \log(f'(c)). \end{aligned}$$

Thus $f'(\partial \log(c)) = \partial \log(f'(c))$ for $c \in \mathbb{Q}(\vec{a})$, and f'' is an \mathcal{L} -isomorphism. By quantifier elimination of LDVF in the language \mathcal{L} , it follows that $\text{tp}_{\text{LDVF}}(\vec{a}/\emptyset) = \text{tp}_{\text{LDVF}}(\vec{b}/\emptyset)$. \square

Lemma 10.23. *For every LDVF-formula $\varphi(\vec{x})$, there is an ACVF-formula $\psi(\vec{x}; \vec{y})$ such that*

$$T \vdash \varphi(\vec{x}) \iff \psi(\vec{x}; \delta(\vec{x})).$$

Proof. A standard compactness argument. □

Theorem 10.24. *If $(K, \partial \log, \text{val}) \models \text{LDVF}$, then $\text{dp-rk}(K) \leq 2$.*

Proof. Suppose there is an ict-pattern of depth 3:

$$\begin{aligned} &\varphi_0(x; \vec{b}_{0,0}), \varphi_0(x; \vec{b}_{0,1}), \dots \\ &\varphi_1(x; \vec{b}_{1,0}), \varphi_0(x; \vec{b}_{1,1}), \dots \\ &\varphi_2(x; \vec{b}_{2,0}), \varphi_0(x; \vec{b}_{2,1}), \dots \end{aligned}$$

Replacing K with an elementary extension, we may assume that K can be expanded to a model of T , by Remark 10.21. Applying Lemma 10.23, we obtain an ict-pattern of depth 3,

$$\begin{aligned} &\varphi'_0(x, y; \vec{c}_{0,0}), \varphi'_0(x, y; \vec{c}_{0,1}), \dots \\ &\varphi'_1(x, y; \vec{c}_{1,0}), \varphi'_0(x, y; \vec{c}_{1,1}), \dots \\ &\varphi'_2(x, y; \vec{c}_{2,0}), \varphi'_0(x, y; \vec{c}_{2,1}), \dots \end{aligned}$$

made of ACVF-formulas. But in ACVF, the set K^2 has dp-rank less than 3. □

10.6 Non-valuation type

Lemma 10.25. *If R is an interpretable integral domain in some structure, and $K = \text{Frac}(R)$, then $\text{dp-rk}(K) = \text{dp-rk}(R)$.*

Proof. Work in a monster model. The inequality $\text{dp-rk}(R) \leq \text{dp-rk}(K)$ is clear. Conversely, suppose there is an ict-pattern of depth κ in K . Then there are formulas $\{\varphi_\alpha(x; y_\alpha)\}_{\alpha < \kappa}$, coefficients $\{b_{\alpha,i}\}_{\alpha < \kappa, i < \omega}$, and witnesses $\{a_\eta\}_{\eta: \kappa \rightarrow \omega}$ in K , such that

$$\varphi_\alpha(a_\eta, b_{\alpha,i}) \iff \eta(\alpha) = i.$$

For any $x_1, \dots, x_n \in K$, we can find a non-zero common denominator $s \in R$ such that

$$\{x_1 s, \dots, x_n s\} \subseteq R.$$

By saturation, we can find some non-zero $s \in R$ such that sa_η lies in R for every η . Let $a'_\eta = sa_\eta$, and let $\psi_\alpha(x, y, z)$ be the formula

$$\psi_\alpha(x, y, z) \equiv \varphi_\alpha(x/y, z).$$

Then there is an ict-pattern of depth κ in R :

$$\psi_\alpha(a'_\eta, s, b_{\alpha,i}) \iff \eta(\alpha) = i. \quad \square$$

Theorem 10.26. *If $(K, \partial \log, \text{val}) \models \text{LDVF}$, then K is unstable, not of valuation type, and has dp-rank exactly 2.*

Proof. Consider the reduct $(K, \partial, \text{val})$, a dense diffeovalued field by Corollary 10.20. Let R be the ring

$$R = \{x \in \mathcal{O} : \partial x \in \mathcal{O}/\mathfrak{m}\},$$

as in §8.4. By Proposition 8.15, the sets $\{aR : a \in K^\times\}$ form a neighborhood basis for a definable non-trivial Hausdorff topology. So certainly K is unstable. Also $\text{Frac}(R) = K$, so $\text{dp-rk}(R) = \text{dp-rk}(K)$ by Lemma 10.25. Therefore $R - R = R$ is a basic neighborhood in the canonical topology. So the canonical topology is finer than the diffeoevaluation topology. If the canonical topology is a V-topology, so is every coarsening, by Theorem 3.2 in [11]. But the diffeoevaluation topology is not a V-topology, by Proposition 8.17. So $(K, \partial \log, \text{val})$ does not have valuation type. Then $\text{dp-rk}(K) \neq 1$, because dp-minimal fields have valuation type (Theorem 9.3.28 in [7]). Therefore, K has dp-rank 2. \square

Remark 10.27. The same argument applies to intermediate reducts between the full model of LDVF, and the reduct $(K, +, \cdot, R)$.

The structure $(K, +, \cdot, \mathcal{O}, R)$ is similar to Example 7.1 in [4]: both are NIP valued fields in which some infinite definable set has empty interior. (In our case, the set is R .) Unlike [4], our example is not a *pure* valued field, but an expansion.

11 Concluding remarks

The example of §10 derails some promising strategies to attack the Shelah conjecture and henselianity conjecture. For example, it disproves our “valuation conjecture” (Conjecture 1.2), which would have implied the Shelah conjecture ([9], Theorem 9.9).

Consider the even simpler conjecture:

Conjecture 11.1. *If $(K, +, \cdot, \mathcal{O}, \dots)$ is a dp-finite valued field, and $S \subseteq K$ is a definable set of full dp-rank ($\text{dp-rk}(S) = \text{dp-rk}(K)$), then S has non-empty interior.*

Conjecture 11.1 would imply the Henselianity conjecture for dp-finite fields, by Theorem 7.5 in [4]. To the best of my knowledge, all the known results on the henselianity conjecture use this strategy.

However, the theory LDVF of §10 is a counterexample to Conjecture 11.1. Indeed, the definable set R has full rank, but empty interior with respect to \mathcal{O} .

Conjecture 11.1 probably holds for *pure* valued fields $(K, +, \cdot, \mathcal{O})$, but this is probably impossible to prove without first classifying dp-finite valued fields using some other strategy. (The purity assumption is hard to use in proofs.)

It seems we need a new strategy to attack the dp-finite Shelah and henselianity conjectures. Perhaps the analysis of §2–5 can be extended to higher ranks.

Here is a conjectural sketch. For any $n \geq 1$, there should be a class of “field topologies of type W_n ,” cut out by a local sentence (in the sense of [11]). The canonical topology on

an unstable dp-finite field K should be a definable W_n -topology for some $n \leq \text{dp-rk}(K)$. For $n = 1$, a W_1 -topology should be the same thing as a V-topology. For $n = 2$, a W_2 -topology should either be a DV-topology in the sense of §8.4, or a topology generated by two independent V-topologies.

For $n = 3$, there should be four types:

- A topology generated by three independent V-topologies.
- A topology generated by a V-topology and an independent DV-topology.
- A topology that is like a DV-topology, but with basic opens

$$\{x \in K : \text{val}(x - a) \geq \gamma, \text{val}(\delta_1 x - b) \geq \gamma, \text{val}(\delta_2 x - c) \geq \gamma\}.$$

for *two* derivations $\delta_1, \delta_2 : K \rightarrow K$.

- A topology that is like a DV-topology, but involving second derivatives, with basic opens

$$\{x \in K : \text{val}(x - a) \geq \gamma, \text{val}(\delta x - b) \geq \gamma, \text{val}(\delta^2 x - c) \geq \gamma\}.$$

Now suppose K is a field of dp-rank 3. Using results from [9], it should be possible to prove that the squaring map $f(x) = x^2$ is an open map from K^\times to K^\times . This should exclude the first two cases. In the latter two cases, it should be possible to show that the W_3 -topology has a unique V-topology coarsening. This would imply that K admits a unique definable V-topology. Generalizations of these arguments should work for $n > 3$.

From this point of view, the rank 2 case is too easy: the rank is so small that there is no room for two independent topologies, unless both are V-topologies.

A Appendix: Resplendent lifting

In the appendix, we assume that all rings are \mathbb{Q} -algebras, all fields extend \mathbb{Q} , and all valued fields have residue characteristic 0.

A.1 Extending derivations

Say that an ordered abelian group Γ is \mathbb{Z} -less if it satisfies the following equivalent conditions:

- For every $a > 0$ in Γ , if Δ^+ is the minimal convex subgroup containing a and Δ^- is the maximal convex subgroup avoiding a , then $\Delta^+/\Delta^- \not\cong \mathbb{Z}$.
- For every $a > 0$ in Γ , there is $b \in \Gamma$ such that

$$(1/3)a < b < (2/3)a,$$

i.e., $a < 3b < 2a$.

- For every $a > 0$ in Γ and every $p < q$ in \mathbb{Q} , there is $b \in \Gamma$ such that

$$pa < b < qa.$$

For example, if Γ is p -divisible for some prime p , then Γ is \mathbb{Z} -less.

Remark A.1. Let $\Gamma' > \Gamma$ be an extension of ordered abelian groups. Suppose Γ'/Γ is torsion (i.e., $\Gamma' \leq \Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$). If Γ is \mathbb{Z} -less, then Γ' is \mathbb{Z} -less.

Lemma A.2. *Let L/K be an algebraic extension of valued fields. Suppose that the value group of K is \mathbb{Z} -less. Let a be a nonzero element of \mathcal{O}_L with positive valuation. Suppose that $a^n \in K$. Then there are $b, c \in \mathcal{O}_L$ such that*

- $a = bc^n$
- bc^{n-1} is in K .
- c^n is in K .

Proof. By \mathbb{Z} -lessness, there is $\gamma \in \Gamma_K$ such that

$$\frac{n-1}{n} \text{val}(a) < \gamma < \text{val}(a).$$

Let $e \in K$ have $\text{val}(e) = \gamma$. Let $b = e^n a^{1-n}$ and $c = ae^{-1}$. Then

$$\begin{aligned} \text{val}(b) &= n \cdot \gamma - (n-1) \text{val}(a) > 0 \\ \text{val}(c) &= \text{val}(ae^{-1}) = \text{val}(a) - \gamma > 0 \\ bc^n &= e^n a^{1-n} a^n e^{-n} = a \\ bc^{n-1} &= e^n a^{1-n} a^{n-1} e^{1-n} = e \in K. \\ c^n &= a^n e^{-n} \in K. \end{aligned}$$

□

Proposition A.3. *Let (K, \mathcal{O}) be an algebraically closed field with a derivation $\partial : \mathcal{O} \rightarrow M$ for some \mathcal{O} -module M . Suppose that ∂ vanishes on some subfield $F \subseteq K$, and K is algebraic over F (so that $K = F^{\text{alg}}$). Suppose $\text{val}(F)$ is \mathbb{Z} -less. Then ∂ vanishes on K .*

The proof is nearly identical to the proof of Proposition 10.12.

Proof. Increasing F , we may assume F is maximal among subfields on which ∂ vanishes. Suppose for the sake of contradiction that $F \subsetneq K$. Take minimal $n > 1$ such that F has a finite extension of degree n . If $P(x) \in F[x]$ has degree $\leq n$, then one of the following happens:

- $P(x)$ factors into linear polynomials
- $P(x)$ is irreducible of degree n .

Take an arbitrary extension L/F of degree n , and break into cases:

- If $\text{val}(L)$ is strictly larger than $\text{val}(F)$, take $\gamma \in \text{val}(L) \setminus \text{val}(F)$. The inequality

$$|\text{val}(L)/\text{val}(F)| \leq [L : F]$$

implies that $m\gamma \in \text{val}(F)$ for some $m \leq n$. Take $c \in F$ with $\text{val}(c) = m\gamma$. The polynomial $x^m - c$ has no roots in F , so $m = n$ and $x^n - c$ is irreducible. Take $a \in K$ such that $a^n = c$.

By maximality of F , there is some $x \in \mathcal{O}_{F(a)}$ such that $\partial x \neq 0$. We can write

$$x = y_0 + y_1 a + \cdots + y_{n-1} a^{n-1}$$

for some $y_i \in F$. Note that $i\gamma \notin \text{val}(F)$ for $i < n$, and so the non-zero terms $y_i a^i$ have pairwise distinct valuations. Therefore

$$0 \leq \text{val}(x) = \min_i (\text{val}(y_i a^i)).$$

So every $y_i a^i$ is in \mathcal{O} , and

$$\partial x = \sum_{i=0}^{n-1} \partial(y_i a^i) = \sum_{i=1}^{n-1} \partial(y_i a^i).$$

For each $i > 0$, we have $\text{val}(y_i a^i) \neq 0$. Then Lemma A.2 gives $b, c \in \mathcal{O}_{F(a)}$ and $e \in \mathcal{O}_F$ such that

$$\begin{aligned} e &= b c^{n-1} \\ y_i a^i &= b c^n = e c \\ c^n &\in F. \end{aligned}$$

Therefore

$$\partial(y_i a^i) = \partial(ec) = e\partial(c) = b c^{n-1} \partial(c) = \frac{b}{n} \partial(c^n) = 0,$$

as $e, c^n \in \mathcal{O}_F$, and $1/n \in \mathcal{O}_K$ (by the assumption of residue characteristic 0). So $\partial(x) = 0$, a contradiction.

- If $\text{res}(L)$ is strictly larger than $\text{res}(F)$, take $\alpha \in \text{res}(L) \setminus \text{res}(F)$. The inequality

$$[\text{res}(L) : \text{res}(F)] \leq [L : F]$$

implies that $[\text{res}(F)(\alpha) : \text{res}(F)] \leq n$. Let

$$x^m + \beta_{m-1} x^{m-1} + \cdots + \beta_1 x + \beta_0$$

be the monic irreducible polynomial of α over $\text{res}(F)$. Because of residue characteristic 0, this polynomial is separable, and so

$$m\alpha^{m-1} + (m-1)\beta_{m-1}\alpha^{m-2} + \cdots + 2\beta_2\alpha + \beta_1 \neq 0. \quad (12)$$

Take $b_i \in F$ with $\text{res } b_i = \beta_i$, and let $P(x)$ be the polynomial

$$x^m + b_{m-1}x^{m-1} + \cdots + b_1x + b_0 \in F[x].$$

Then $P(x)$ is irreducible, and so $m = [\text{res}(F)(\alpha) : \text{res}(F)] = n$. Let $a \in K$ be the root of $P(x)$ with $\text{res}(a) = \alpha$. Then

$$a^n + b_{n-1}a^{n-1} + \cdots + b_1a + b_0 = 0.$$

Applying the derivation $\partial : \mathcal{O} \rightarrow M$, which vanishes on the b_i , we obtain

$$(na^{n-1} + (n-1)b_{n-1}a^{n-2} + \cdots + 2b_2a + b_1)\partial a = 0.$$

The expression inside the parentheses has nonzero residue, by (12), and so it is an element of \mathcal{O}^\times . Therefore $\partial a = 0$.

Now if

$$x = y_0 + y_1a + \cdots + y_{n-1}a^{n-1}$$

is any element of $F(a)$, then $\text{val}(x) = \min_i \text{val}(y_i)$. To see this, one reduces to the case where $\min_i(\text{val}(y_i)) = 0$; then

$$\text{res}(y_0) + \text{res}(y_1)\alpha + \cdots + \text{res}(y_{n-1})\alpha^{n-1} \neq 0,$$

by linear independence of $\{1, \alpha, \dots, \alpha^{n-1}\}$ over $\text{res}(K)$.

Consequently, $\mathcal{O}_{F(a)} = \mathcal{O}_F[a]$. As ∂ vanishes on \mathcal{O}_F and a , it vanishes on $\mathcal{O}_{F(a)}$, contradicting the maximality of F .

- Lastly, suppose that L/F is an immediate extension. By maximality of F , there is $a \in \mathcal{O}_L$ with $\partial(a) \neq 0$. Let \mathcal{C} be the collection of balls containing a , with center and radius from F . Let I be the intersection $\bigcap \mathcal{C}$. As usual, $I \cap F = \emptyset$.

Let $P(x)$ be the minimal polynomial of a over F . Then $P(x)$ has degree n . Let a_1, \dots, a_n be the roots of $P(x)$, with $a_1 = a$. Note that $P'(x)$ has degree $n-1$, and therefore splits over F . So no root of $P'(x)$ is in I . By Rolle's Theorem (Lemma 10.10), a is the unique root of $P(x)$ in I .

Therefore I has empty intersection with the finite set $\{0, a_2, \dots, a_n\}$. We can find $b \in F$ such that

$$\text{val}(a - b) > \max(\text{val}(0 - b), \text{val}(a_2 - b), \text{val}(a_3 - b), \dots, \text{val}(a_n - b)).$$

Take $c \in F$ with $\text{val}(a - b) = \text{val}(c)$, and let $e_i = (a_i - b)/c$. Then $\text{val}(e_1) = 0$, and $\text{val}(e_i) < 0$ for $i > 1$. The e_i are the roots of the irreducible polynomial

$$Q(x) = P(cx + b) = s_nx^n + s_{n-1}x^{n-1} + \cdots + s_1x + s_0 \in F[x].$$

By Newton polygons, $\text{val}(s_0) = \text{val}(s_1) < \text{val}(s_i)$ for $i > 1$. Then we can apply ∂ to the equation

$$(s_n/s_1)e_1^n + \cdots + (s_2/s_1)e_1^2 + e_1 + (s_0/s_1) = 0,$$

and obtain

$$(n(s_n/s_1)e_1^{n-1} + \cdots + 2(s_2/s_1)e_1 + 1)\partial e_1 = 0,$$

because the coefficients s_n/s_1 lie in F , where ∂ vanishes. But the expression in parentheses has valuation 0, because $e_1 \in \mathcal{O}$ and $s_i/s_1 \in \mathfrak{m}$ for $2 \leq i \leq n$. Therefore $\partial e_1 = 0$.

Meanwhile, $\text{val}(a - b) > \text{val}(b)$ implies that

$$0 \leq \text{val}(a) = \min(\text{val}(a - b), \text{val}(b)) = \min(\text{val}(c), \text{val}(b)),$$

and so $b, c \in \mathcal{O}_F$. But $a = b + e_1c$, and so $\partial(a) = c\partial e_1 = 0$, contradicting the choice of a . \square

A.2 Review of Kähler differentials

If $A \rightarrow B$ is a morphism of (commutative unital) rings, then $\Omega_{B/A}$ denotes the module of Kähler differentials. This is the B -module generated by terms db for $b \in B$, subject to the relations

$$\begin{aligned} d(b_1 + b_2) &= db_1 + db_2 \\ d(b_1b_2) &= b_1db_2 + b_2db_1 \\ da &= 0 \quad \text{if } a \in A. \end{aligned}$$

If M is a B -module, there is an isomorphism

$$\text{Hom}_B(\Omega_{B/A}, M) \cong \text{Der}_A(B, M)$$

natural in M , where $\text{Der}_A(B, M)$ denotes the set of A -linear derivations $B \rightarrow M$.

The following facts about Kähler differentials are well-known:

Fact A.4. *If $A \rightarrow B \rightarrow C$ is a morphism of rings, then*

$$\Omega_{B/A} \otimes_B C \rightarrow \Omega_{C/A} \rightarrow \Omega_{C/B} \rightarrow 0$$

is exact.

Fact A.5. *If $A \rightarrow B$ is a morphism of rings and $S \subseteq B$ is a multiplicative subset, then*

$$S^{-1}\Omega_{B/A} \cong \Omega_{S^{-1}B/A}.$$

Fact A.6. *If L/K is an extension of (characteristic 0) fields and $\{t_i\}_{i \in I}$ is a transcendence basis (possibly infinite), then $\{dt_i\}_{i \in I}$ is an L -basis of $\Omega_{L/K}$.*

Remark A.7. If \mathcal{O} is a valuation ring and M is an \mathcal{O} -module, the following are equivalent:

1. M is flat.
2. M is torsionless.
3. Every finitely-generated submodule of M is free.
4. M is a direct limit of free modules.
5. The natural map $M \rightarrow M \otimes_{\mathcal{O}} K$ is an injection.

We will use two flatness results from ([2], Corollary 6.5.21 and Theorem 6.5.15).

Fact A.8. *If \mathcal{O} is a valuation ring with residue characteristic 0, then $\Omega_{\mathcal{O}/\mathbb{Q}}$ is flat as an \mathcal{O} -module.*

Fact A.9. *Let \mathcal{O}'/\mathcal{O} be an extension of valuation rings. Suppose $\text{Frac}(\mathcal{O}) \models \text{ACF}$. Then $\Omega_{\mathcal{O}'/\mathcal{O}}$ is flat as an \mathcal{O} -module.*

A.3 Flatness and extensions

Lemma A.10. *Let \mathcal{O} be a valuation ring and $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a short exact sequence of \mathcal{O} -modules.*

- *If B is flat, then A is flat.*
- *If A and C are flat, then B is flat.*

Proof. By Remark A.7, an \mathcal{O} -module is flat if and only if it is torsionless.

For the first point: submodules of torsionless modules are torsionless.

For the second point, suppose that A and C are torsionless. For any nonzero $r \in \mathcal{O}$, there is a diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \end{array}$$

where the rows are exact and the vertical maps are multiplication by r . Because A and C are torsionless, the outer vertical maps are injective. By the snake lemma, the inner vertical map is injective. As r is arbitrary, C is torsionless. \square

Lemma A.11. *Let \mathcal{O}'/\mathcal{O} be an extension of valuation rings. Let M be an \mathcal{O} -module. Then M is flat (as an \mathcal{O} -module) if and only if $M \otimes_{\mathcal{O}} \mathcal{O}'$ is flat (as an \mathcal{O}' -module).*

Proof. If M is flat, then M is a direct limit of free \mathcal{O} -modules, and so $M \otimes_{\mathcal{O}} \mathcal{O}'$ is a direct limit of free \mathcal{O}' -modules. Conversely, suppose that M is not flat. Then there is an injection

$$\mathcal{O}/I \hookrightarrow M$$

for some non-zero proper ideal I in \mathcal{O} . As \mathcal{O}' is torsionless over \mathcal{O} , it is flat as an \mathcal{O} -module. Therefore the functor $- \otimes_{\mathcal{O}} \mathcal{O}'$ is exact, and the map

$$(\mathcal{O}/I) \otimes_{\mathcal{O}} \mathcal{O}' \hookrightarrow M \otimes_{\mathcal{O}} \mathcal{O}'$$

is injective. But $(\mathcal{O}/I) \otimes_{\mathcal{O}} \mathcal{O}' \cong \mathcal{O}'/I\mathcal{O}'$. The ideal $I\mathcal{O}'$ is non-trivial, because it contains the non-trivial elements of I . And $I\mathcal{O}'$ is a proper ideal, because it is generated by elements of positive valuation. Therefore $\mathcal{O}'/I\mathcal{O}'$ is not torsionless, and neither is the larger module $M \otimes_{\mathcal{O}} \mathcal{O}'$. \square

Lemma A.12. *Let $K_1 \subseteq K_2 \subseteq K_3$ be a chain of three fields (of characteristic 0). Then the map*

$$\Omega_{K_2/K_1} \otimes_{K_2} K_3 \rightarrow \Omega_{K_3/K_1}$$

is injective.

Proof. Let B be a transcendence basis of K_2/K_1 , and B' be a transcendence basis of K_3/K_1 extending B . By Fact A.6, the set $\{dt : t \in B\}$ is a K_3 -linear basis of $\Omega_{K_2/K_1} \otimes_{K_2} K_3$, and the set $\{dt : t \in B'\}$ is a K_3 -linear basis of Ω_{K_3/K_1} . The map in question is induced by the inclusion $B \hookrightarrow B'$, and is therefore injective. \square

Lemma A.13. *Let \mathcal{O}'/\mathcal{O} be an extension of valuation rings (with residue characteristic 0). Then the map $\Omega_{\mathcal{O}/\mathbb{Q}} \otimes_{\mathcal{O}} \mathcal{O}' \rightarrow \Omega_{\mathcal{O}'/\mathbb{Q}}$ is injective.*

Proof. This follows from the commuting diagram

$$\begin{array}{ccccccc} \Omega_{\mathcal{O}/\mathbb{Q}} \otimes_{\mathcal{O}} \mathcal{O}' & \hookrightarrow & (\Omega_{\mathcal{O}/\mathbb{Q}} \otimes_{\mathcal{O}} \mathcal{O}') \otimes_{\mathcal{O}'} K' & \xlongequal{\quad} & (\Omega_{\mathcal{O}/\mathbb{Q}} \otimes_{\mathcal{O}} K) \otimes_K K' & \xlongequal{\quad} & \Omega_{K/\mathbb{Q}} \otimes_K K' , \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{\mathcal{O}'/\mathbb{Q}} & \hookrightarrow & \Omega_{\mathcal{O}'/\mathbb{Q}} \otimes_{\mathcal{O}'} K' & \xlongequal{\quad} & \Omega_{\mathcal{O}'/\mathbb{Q}} \otimes_{\mathcal{O}'} K' & \xlongequal{\quad} & \Omega_{K'/\mathbb{Q}} \end{array}$$

where the left horizontal arrows are injective by flatness (Fact A.8), the right horizontal arrows are isomorphisms by Fact A.5, and the rightmost vertical map is injective by Lemma A.12. \square

Lemma A.14. *Let $\mathcal{O}_1 \subseteq \mathcal{O}_2 \subseteq \mathcal{O}_3$ be a chain of two valuation ring extensions⁵. Then the map $\Omega_{\mathcal{O}_2/\mathcal{O}_1} \otimes_{\mathcal{O}_2} \mathcal{O}_3 \rightarrow \Omega_{\mathcal{O}_3/\mathcal{O}_1}$ is injective.*

⁵Meaning that the inclusions are local homomorphisms.

Proof. There is a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \Omega_{\mathcal{O}_1/\mathbb{Q}} \otimes_{\mathcal{O}_1} \mathcal{O}_3 & \longrightarrow & \Omega_{\mathcal{O}_2/\mathbb{Q}} \otimes_{\mathcal{O}_2} \mathcal{O}_3 & \longrightarrow & \Omega_{\mathcal{O}_2/\mathcal{O}_1} \otimes_{\mathcal{O}_2} \mathcal{O}_3 \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \longrightarrow & \Omega_{\mathcal{O}_1/\mathbb{Q}} \otimes_{\mathcal{O}_1} \mathcal{O}_3 & \longrightarrow & \Omega_{\mathcal{O}_3/\mathbb{Q}} & \longrightarrow & \Omega_{\mathcal{O}_3/\mathcal{O}_1} \longrightarrow 0
\end{array} \tag{13}$$

The bottom row is right exact by Fact A.4, and left exact by Lemma A.13. The same argument shows that

$$0 \rightarrow \Omega_{\mathcal{O}_1/\mathbb{Q}} \otimes_{\mathcal{O}_1} \mathcal{O}_2 \rightarrow \Omega_{\mathcal{O}_2/\mathbb{Q}} \rightarrow \Omega_{\mathcal{O}_2/\mathcal{O}_1} \rightarrow 0$$

is an exact sequence. Applying the exact functor $-\otimes_{\mathcal{O}_2} \mathcal{O}_3$ yields the exactness of the top row of (13). The middle vertical map of (13) is injective by Lemma A.13. The snake lemma then implies that the right vertical map is injective. \square

Definition A.15. Let \mathcal{O}'/\mathcal{O} be an extension of valuation rings. Then \mathcal{O}'/\mathcal{O} is *pseudosmooth* if $\Omega_{\mathcal{O}'/\mathcal{O}}$ is flat (as an \mathcal{O}' -module).

Proposition A.16. Let $\mathcal{O}_1 \subseteq \mathcal{O}_2 \subseteq \mathcal{O}_3$ be a chain of two valuation ring extensions.

1. If $\mathcal{O}_2/\mathcal{O}_1$ and $\mathcal{O}_3/\mathcal{O}_2$ are pseudosmooth, then $\mathcal{O}_3/\mathcal{O}_1$ is pseudosmooth.
2. If $\mathcal{O}_3/\mathcal{O}_1$ is pseudosmooth, then $\mathcal{O}_2/\mathcal{O}_1$ is pseudosmooth.

Proof. The sequence

$$0 \rightarrow \Omega_{\mathcal{O}_2/\mathcal{O}_1} \otimes_{\mathcal{O}_2} \mathcal{O}_3 \rightarrow \Omega_{\mathcal{O}_3/\mathcal{O}_1} \rightarrow \Omega_{\mathcal{O}_3/\mathcal{O}_2} \rightarrow 0$$

is right exact by Fact A.4, and left exact by Lemma A.14. Then

$$\begin{aligned}
\mathcal{O}_2/\mathcal{O}_1 \text{ is pseudosmooth} &\iff \Omega_{\mathcal{O}_2/\mathcal{O}_1} \text{ is flat} \iff \Omega_{\mathcal{O}_2/\mathcal{O}_1} \otimes_{\mathcal{O}_2} \mathcal{O}_3 \text{ is flat} \\
\mathcal{O}_3/\mathcal{O}_1 \text{ is pseudosmooth} &\iff \Omega_{\mathcal{O}_3/\mathcal{O}_1} \text{ is flat} \\
\mathcal{O}_3/\mathcal{O}_2 \text{ is pseudosmooth} &\iff \Omega_{\mathcal{O}_3/\mathcal{O}_2} \text{ is flat,}
\end{aligned}$$

using Lemma A.11 in the first line. The desired statements follow from Lemma A.10. \square

Proposition A.17. Let \mathcal{O}'/\mathcal{O} be an extension of valued fields of residue characteristic 0. Suppose the value group of \mathcal{O} is \mathbb{Z} -less. Then $\Omega_{\mathcal{O}'/\mathcal{O}}$ is flat as an \mathcal{O}' -module.

Proof. We must show that \mathcal{O}'/\mathcal{O} is pseudosmooth. By Proposition A.16.2, we may replace \mathcal{O}' with a larger valued field. Since valuations can be extended along any field extension, we may assume that $\text{Frac}(\mathcal{O}')$ contains the algebraic closure of $\text{Frac}(\mathcal{O})$. Let \mathcal{O}'' be the induced valuation ring on $\text{Frac}(\mathcal{O})^{alg}$. Then $\mathcal{O} \subseteq \mathcal{O}'' \subseteq \mathcal{O}'$. Now $\mathcal{O}'/\mathcal{O}''$ is pseudosmooth by Fact A.9, as $\text{Frac}(\mathcal{O}'')$ is algebraically closed. By Proposition A.16.1, it remains to show that $\mathcal{O}''/\mathcal{O}$ is pseudosmooth. In fact, $\Omega_{\mathcal{O}''/\mathcal{O}}$ vanishes, by Proposition A.3. \square

A.4 Resplendent lifting in the \mathbb{Z} -less case

Lemma A.18. *Let R be a ring, and*

$$\begin{array}{ccc} A & \xrightarrow{g'} & M \\ f' \downarrow & & \downarrow f \\ A \oplus B & \xrightarrow{g} & N \end{array}$$

be a diagram of R -modules, with f surjective, and f' the inclusion of the first factor. If B is free, then there is a diagonal map $h : A \oplus B \rightarrow M$ making the diagram commute:

$$\begin{array}{ccc} A & \xrightarrow{g'} & M \\ f' \downarrow & \nearrow h & \downarrow f \\ A \oplus B & \xrightarrow{g} & N. \end{array}$$

The proof is well-known but included for completeness.

Proof. Consider the morphism $B \rightarrow N$ given by $x \mapsto g(0, x)$. Because B is free, there is $h_1 : B \rightarrow M$ lifting this, so that

$$g(0, x) = f(h_1(x))$$

for $x \in B$. Define $h : A \oplus B \rightarrow M$ by the formula $h(x, y) = g'(x) + h_1(y)$. Then

$$h(f'(x)) = h(x, 0) = g'(x)$$

$$f(h(x, y)) = f(g'(x)) + f(h_1(y)) = g(f'(x)) + g(0, y) = g(x, 0) + g(0, y) = g(x, y). \quad \square$$

Lemma A.19. *Let R be a valuation ring. Let*

$$\begin{array}{ccc} A & \xrightarrow{g'} & M \\ f' \downarrow & & \downarrow f \\ B & \xrightarrow{g} & N \end{array}$$

be a diagram of R -modules, with $f' : A \hookrightarrow B$ injective and $f : M \rightarrow N$ surjective. Suppose the following hold:

- $\text{coker}(f')$ is flat.
- Let (M, N) be the two-sorted structure with the R -module structure on M and N , and the surjection $f : M \rightarrow N$. Then (M, N) is $(|A| + |B| + |R|)^+$ -saturated.

Then there is a morphism $h : B \rightarrow M$ making the diagram commute

$$\begin{array}{ccc} A & \xrightarrow{g'} & M \\ f' \downarrow & \nearrow h & \downarrow f \\ B & \xrightarrow{g} & N. \end{array}$$

Proof. Without loss of generality, $f' : A \hookrightarrow B$ is an inclusion. Then B/A is flat.

Let $\vec{x} = \langle x_b \rangle_{b \in B}$ be a tuple of variables indexed by B . For every submodule $C \subseteq B$ containing A , let $\Sigma_C(\vec{x})$ be the $*$ -type in (M, N) asserting the following:

- If $b \in C$, then $x_b \in M$ and $f(x_b) = g(b)$.
- If $b \in A$, then $x_b = g'(x_b)$.
- If $r \in R$ and $b \in C$, then $rx_b = x_{rb}$.
- If $b, b' \in C$, then $x_{b+b'} = x_b + x_{b'}$.

Then $\Sigma_C(\vec{x})$ is realized in (M, N) if and only if there is a morphism $h_C : C \rightarrow M$ such that the diagram commutes

$$\begin{array}{ccc}
 A & \xrightarrow{g'} & M \\
 \subseteq \downarrow & \nearrow h_C & \downarrow f \\
 C & \xrightarrow{g_C} & N
 \end{array} \tag{14}$$

It suffices to realize $\Sigma_B(\vec{x})$. This type is a directed union of the types

$$\{\Sigma_C(\vec{x}) : C/A \text{ is finitely generated}\}.$$

By saturation, it suffices to realize the types in this family. Suppose C/A is finitely generated. Then C/A injects into B/A , so C/A is free by Remark A.7. The sequence

$$0 \rightarrow A \rightarrow C \rightarrow C/A \rightarrow 0$$

therefore splits. By Lemma A.18, there is a dashed arrow making (14) commute. \square

Note that \mathbb{Z} -lessness is a conjunction of first-order axioms, so it is preserved in elementary equivalence of ordered abelian groups.

Theorem A.20. *Let (K, \mathcal{O}) be a valued field with residue characteristic 0 and \mathbb{Z} -less value group. Let $f : M \twoheadrightarrow N$ be a surjective morphism of \mathcal{O} -modules. Let $\partial : \mathcal{O} \rightarrow N$ be a derivation. Consider the three-sorted structure (\mathcal{O}, M, N) with the ring structure on \mathcal{O} , the module structures on M, N , the epimorphism f , and the derivation ∂ .*

- *If the structure (\mathcal{O}, M, N) is sufficiently saturated and resplendent, then there is a derivation $\delta : \mathcal{O} \rightarrow M$ making the diagram commute:*

$$\begin{array}{ccc}
 & & M \\
 & \nearrow \delta & \downarrow f \\
 \mathcal{O} & \xrightarrow{\partial} & N
 \end{array}$$

- *In general, such a lifting exists after passing to an elementary extension.*

Proof. The two statements are clearly equivalent, by definition of resplendence and existence of resplendent elementary extensions. We prove the second statement. Consider an elementary chain

$$(\mathcal{O}, M, N) = (\mathcal{O}_0, M_0, N_0) \preceq (\mathcal{O}_1, M_1, N_1) \preceq (\mathcal{O}_2, M_2, N_2) \preceq \cdots$$

where each structure is saturated over the previous structure. Let f_i, ∂_i be the structure maps in $(\mathcal{O}_i, M_i, N_i)$. We will recursively build a sequence of derivations $\delta_i : \mathcal{O}_i \rightarrow M_{i+1}$ such that

$$\begin{aligned} f_{i+1}(\delta_i(x)) &= \partial_i(x) \\ \delta_{i+1}(x) &= \delta_i(x) \end{aligned}$$

for $i \geq 0$ and $x \in \mathcal{O}_i$. If this can be done successfully, then the union of the δ_i 's is the desired lifting of ∂ on the structure $\bigcup_i (\mathcal{O}_i, M_i, N_i)$, an elementary extension of (\mathcal{O}, M, N) , and we are done.

At step $i = 0$, we must find an \mathcal{O}_0 -linear map $\Omega_{\mathcal{O}_0/\mathbb{Q}} \rightarrow M_1$ making the diagram commute

$$\begin{array}{ccccc} & & & & M_1 \\ & & & & \downarrow f_1 \\ \Omega_{\mathcal{O}_0/\mathbb{Q}} & \xrightarrow{\partial_0} & N_0 & \xrightarrow{\subseteq} & N_1 \\ & \nearrow & & & \downarrow f_1 \end{array} \quad (15)$$

At step $i > 0$, we must find an \mathcal{O}_i -linear map $\Omega_{\mathcal{O}_i/\mathbb{Q}} \rightarrow M_{i+1}$ making the diagram commute

$$\begin{array}{ccccc} (\mathcal{O}_i \otimes_{\mathcal{O}_{i-1}} \Omega_{\mathcal{O}_{i-1}/\mathbb{Q}})^{\delta_i^{-1}} & \xrightarrow{\subseteq} & M_i & \xrightarrow{\subseteq} & M_{i+1} \\ \downarrow & \nearrow \text{---} & & & \downarrow f_{i+1} \\ \Omega_{\mathcal{O}_i/\mathbb{Q}} & \xrightarrow{\partial_i} & N_i & \xrightarrow{\subseteq} & N_{i+1} \end{array} \quad (16)$$

The dashed map exists in both cases by Lemma A.19. For (15), Fact A.8 shows that $\Omega_{\mathcal{O}_0/\mathbb{Q}}$ is flat. For (16), the map

$$\mathcal{O}_i \otimes_{\mathcal{O}_{i-1}} \Omega_{\mathcal{O}_{i-1}/\mathbb{Q}} \rightarrow \Omega_{\mathcal{O}_i/\mathbb{Q}}$$

is injective by Lemma A.13, the cokernel is $\Omega_{\mathcal{O}_i/\mathcal{O}_{i-1}}$ by Fact A.4, and $\Omega_{\mathcal{O}_i/\mathcal{O}_{i-1}}$ is flat by Proposition A.17. \square

Corollary A.21. *Let (K, ∂) be a sufficiently resplendent normalized diffeovalued field. If the value group of K is \mathbb{Z} -less, or p -divisible for some p , then (K, ∂) admits a lifting.*

Proof. If the value group is p -divisible, then it is \mathbb{Z} -less. Theorem A.20 allows us to lift the given derivation $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$ to a derivation $\delta_0 : \mathcal{O} \rightarrow K$. This corresponds to an \mathcal{O} -linear map $\Omega_{\mathcal{O}/\mathbb{Q}} \rightarrow K$, which in turn yields a K -linear map

$$\Omega_{K/\mathbb{Q}} \cong \Omega_{\mathcal{O}/\mathbb{Q}} \otimes_{\mathcal{O}} K \rightarrow K$$

by Fact A.5. Thus $\delta_0 : \mathcal{O} \rightarrow K$ extends to a derivation $\delta : K \rightarrow K$. \square

A.5 An unliftable example

The assumption that Γ is \mathbb{Z} -less is necessary in Corollary A.21.

Proposition A.22. *There is a valued field $(K, \mathcal{O}, \mathfrak{m})$ of residue characteristic 0, and a derivation $\partial : \mathcal{O} \rightarrow K/\mathfrak{m}$, such that in any elementary extension $(K^*, \mathcal{O}^*, \mathfrak{m}^*, \partial) \succeq (K, \mathcal{O}, \mathfrak{m}, \partial)$, there is no derivation $\delta : \mathcal{O}^* \rightarrow K^*$ making the diagram commute:*

$$\begin{array}{ccc} & & K^* \\ & \nearrow \delta & \downarrow \\ \mathcal{O}^* & \xrightarrow{\partial} & K^*/\mathfrak{m}^* \end{array}$$

One can even take (K, \mathcal{O}) to be dp-minimal as a pure valued field.

Proof. Let $\mathbb{Z} + \mathbb{Z}\omega$ be the free abelian group on two generators $1, \omega$, ordered so that $\omega > n \cdot 1$ for all $n \in \mathbb{Z}$. In other words, $\mathbb{Z} + \mathbb{Z}\omega$ is the lexicographic product $\mathbb{Z} \times \mathbb{Z}$, with generators $\omega := (1, 0)$ and $1 := (0, 1)$.

Let L be the Hahn field $\mathbb{Q}^{alg}((t^{\mathbb{Z} + \mathbb{Z}\omega}))$, and let K be the relative algebraic closure of $\mathbb{Q}(t, t^\omega)$ in L . Let $\mathcal{O}_L, \mathcal{O}_K$ denote the valuation rings on L and K , and $\mathfrak{m}_L, \mathfrak{m}_K$ denote their maximal ideals. The valued field (K, \mathcal{O}_K) is henselian with residue characteristic 0, algebraically closed residue field, and dp-minimal value group, so (K, \mathcal{O}_K) is a dp-minimal valued field.

Let val' be the coarsening of val by the convex subgroup $\mathbb{Z} \leq \mathbb{Z} + \mathbb{Z}\omega$, and let $\mathfrak{p} \triangleleft \mathcal{O}_L$ be the associated maximal ideal. If $\text{val}(x) = i + j\omega$, then $\text{val}'(x) = j$. Moreover, for any x ,

$$x \in \mathfrak{p} \iff \text{val}'(x) > 0 \iff \text{val}(x) > \mathbb{Z}.$$

Note that for $x \in L$,

$$\text{val}'(x) \geq 0 \implies x \in \mathfrak{m}_L + \mathbb{Q}^{alg}[t^{-1}], \quad (17)$$

because one can split $x = \sum_{i,j} a_{i,j} t^{i+j\omega}$ as

$$x = \sum_{i+j\omega \leq 0} a_{i,j} t^{i+j\omega} + \sum_{i+j\omega > 0} a_{i,j} t^{i+j\omega}.$$

The assumption $\text{val}'(x) \geq 0$ ensures that the first sum only involves $i + j\omega$ with $j = 0$. The support is well-ordered, so the first sum is finite, and belongs to $\mathbb{Q}^{alg}[t^{-1}]$. The other sum is in \mathfrak{m}_L , proving (17).

Choose

$$u = 1 + a_1 t + a_2 t^2 + \cdots \in 1 + t\mathbb{Q}^{alg}[[t]] \subseteq \mathbb{Q}^{alg}((t)) = \mathbb{Q}^{alg}((t^{\mathbb{Z}})) \subseteq \mathbb{Q}^{alg}((t^{\mathbb{Z} + \mathbb{Z}\omega})) = L$$

such that $u \not\equiv v \pmod{\mathfrak{p}}$ for all $v \in K$. Such a u exists because $1 + t\mathbb{Q}^{alg}[[t]]$ is uncountable, K is countable, and the elements of $\mathbb{Q}^{alg}((t))$ are pairwise distinct modulo \mathfrak{p} . (The valuation val' restricts to the trivial valuation on $\mathbb{Q}^{alg}((t))$.)

Consider the derivation

$$\begin{aligned} \partial_0 : L &\rightarrow L \\ \sum_{i,j} a_{i,j} t^{i+j\omega} &\mapsto \sum_{i,j} a_{i,j} j t^{i+(j-1)\omega}. \end{aligned}$$

Note that for $x \in L$,

$$x \in \mathcal{O}_L \implies \text{val}'(\partial_0 x) \geq 0 \tag{18}$$

Let $\partial_1 : L \rightarrow L$ be the derivation $\partial_1 x := u\partial_0 x$. Let ∂ be the composition

$$\mathcal{O}_K \hookrightarrow L \xrightarrow{\partial_1} L \rightarrow L/\mathfrak{m}_L.$$

We claim that ∂ factors through the inclusion $K/\mathfrak{m}_K \hookrightarrow L/\mathfrak{m}_L$. Indeed,

$$x \in \mathcal{O}_K \implies x \in \mathcal{O}_L \implies \text{val}'(\partial_0 x) \geq 0 \iff \text{val}'(u\partial_0 x) \geq 0,$$

because $\text{val}(u) = 0$. By (17),

$$\begin{aligned} \text{val}'(u\partial_0 x) \geq 0 &\implies u\partial_0 x \in \mathfrak{m}_L + \mathbb{Q}^{alg}[t^{-1}] \\ &\implies \partial x \in (\mathbb{Q}^{alg}[t^{-1}] + \mathfrak{m}_L)/\mathfrak{m}_L \subseteq (K + \mathfrak{m}_L)/\mathfrak{m}_L \cong K/(K \cap \mathfrak{m}_K) = K/\mathfrak{m}_K. \end{aligned}$$

So ∂ is a well-defined derivation from \mathcal{O}_K to K/\mathfrak{m}_K .

We claim that the following first-order statement σ holds in the structure $(K, \mathcal{O}_K, \partial)$:

There is an $a \in \mathcal{O}_K$ such that for every $a' \in K$, there are $b, c \in \mathcal{O}_K$ such that $a = bc$ and for every $b', c' \in K$, the following identities do *not* all hold:

$$\begin{aligned} b' &\equiv \partial b \pmod{\mathfrak{m}_K} \\ c' &\equiv \partial c \pmod{\mathfrak{m}_K} \\ a' &= bc' + cb'. \end{aligned}$$

Before proving this, note that this would complete the proof:

- The statement σ is first-order, so it remains true in any elementary extension of $(K, \mathcal{O}, \partial)$.
- If the lifting $\delta : \mathcal{O} \rightarrow K$ exists, the statement σ is false, because an adversary can choose

$$\begin{aligned} a' &= \delta a \\ b' &= \delta b \\ c' &= \delta c. \end{aligned}$$

and the three equations would hold.

We now prove σ . For our opening move, we choose $a = t^\omega \in \mathcal{O}_K$. The opponent chooses $a' \in K$. Note

$$\begin{aligned}\partial_0 a &= \partial_0 t^\omega = 1 \\ \partial_1 a &= u \partial_0 a = u.\end{aligned}$$

By choice of u , we know that $a' - u \notin \mathfrak{p}$, so $\text{val}(a' - u) < n$ for some $n \in \mathbb{Z}$.

For our next move, we take $b = t^n$ and $c = t^{\omega-n}$. The condition $a = bc$ holds, so we haven't lost the game yet. The opponent chooses $b', c' \in K$. Suppose that all three identities hold:

$$\begin{aligned}b' &\equiv \partial b \pmod{\mathfrak{m}_K} \\ c' &\equiv \partial c \pmod{\mathfrak{m}_K} \\ a' &= bc' + cb'.\end{aligned}$$

Then

$$\begin{aligned}\partial_1 b &\equiv b' \pmod{\mathfrak{m}_L} \\ \partial_1 c &\equiv c' \pmod{\mathfrak{m}_L}.\end{aligned}$$

Now b, c are divisible by t^n , so

$$\begin{aligned}c \partial_1 b &\equiv cb' \pmod{t^n \mathfrak{m}_L} \\ b \partial_1 c &\equiv bc' \pmod{t^n \mathfrak{m}_L}\end{aligned}$$

Adding the two equations, and using the identities

$$\begin{aligned}a' &= bc' + cb' \\ \partial_1(bc) &= b \partial_1 c + c \partial_1 b,\end{aligned}$$

we obtain

$$\partial_1(bc) \equiv a' \pmod{t^n \mathfrak{m}_L}$$

On the other hand,

$$\partial_1(bc) = \partial_1(a) = u,$$

so $u \equiv a' \pmod{t^n \mathfrak{m}_L}$. Then $\text{val}(u - a') > n$, contradicting the choice of n . So it is impossible for all three identities to hold, and we have won the game. This proves the sentence σ and completes the proof. \square

Acknowledgments. The author would like to thank Meng Chen, Hagen Knaf, and Franz-Viktor Kuhlmann for some helpful information on Kähler differentials. This material is based upon work supported by the National Science Foundation under Award No. DMS-1803120. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author and do not necessarily reflect the views of the National Science Foundation.

References

- [1] Artem Chernikov, Itay Kaplan, and Pierre Simon. Groups and fields with NTP_2 . *Proc. Amer. Math. Soc.*, 143:395–406, 2015.
- [2] Ofer Gabber and Lorenzo Ramero. *Almost Ring Theory*. Number 1800 in Lectures Notes in Mathematics. Springer, 2003.
- [3] Yatir Halevi, Assaf Hasson, and Franziska Jahnke. A conjectural classification of strongly dependent fields. *Bulletin of Symbolic Logic*, 25(2):182–195, June 2019.
- [4] Yatir Halevi, Assaf Hasson, and Franziska Jahnke. Definable V-topologies, henselianity and NIP. arXiv:1901.05920v2 [math.LO], 2019.
- [5] Yatir Halevi and Daniel Palacín. The dp-rank of abelian groups. *Journal of Symbolic Logic*, 84:957–986, September 2019.
- [6] Franziska Jahnke and Jochen Koenigsmann. Uniformly defining p-henselian valuations. *Annals of Pure and Applied Logic*, 166(7-8):741–754, July-August 2015.
- [7] Will Johnson. *Fun with Fields*. PhD thesis, University of California, Berkeley, 2016. Available at <https://math.berkeley.edu/~willij/drafts/will-thesis.pdf>.
- [8] Will Johnson. Dp-finite fields I: infinitesimals and positive characteristic. arXiv:1903.11322v2 [math.LO], March 2019.
- [9] Will Johnson. Dp-finite fields II: the canonical topology and its relation to henselianity. arXiv:1910.05932v2 [math.LO], October 2019.
- [10] Will Johnson. Dp-finite fields III: inflators and directories. arXiv:1911.04727v1 [math.LO], November 2019.
- [11] Alexander Prestel and Martin Ziegler. Model theoretic methods in the theory of topological fields. *Journal für die reine und angewandte Mathematik*, 299-300:318–341, 1978.
- [12] Pierre Simon. *A guide to NIP theories*. Lecture Notes in Logic. Cambridge University Press, July 2015.