Characterizing uninorms on bounded lattices[☆]

Xinxing Wu^{a,b,c,*}, Guanrong Chen^d

^aSchool of Sciences, Southwest Petroleum University, Chengdu, Sichuan 610500, China ^bInstitute for Artificial Intelligence, Southwest Petroleum University, Chengdu, Sichuan 610500, China ^cZhuhai College of Jilin University, Zhuhai, Guangdong 519041, China ^dDepartment of Electrical Engineering, City University of Hong Kong, Hong Kong SAR, China

Abstract

This paper establishes some equivalent conditions of a uninorm, extending an arbitrary triangular norm on [0, e] or an arbitrary triangular conorm on [e, 1] to the whole lattice.

Keywords: Lattice; Uninorm; Incomparability.

2010 MSC: 03B52, 06B20, 03E72.

1. Introduction and preliminaries

As an extension of the logic connective conjunction and disjunction in classical two-valued logic, triangular norms (t-norms) with the neutral 1 and triangular conorms (t-conorms) with the neutral 0 on the unit interval [0,1] were introduced by Menger [10] and by Schweizer and Sklar [12], respectively. Then, the concepts were extended to uniorms by Yager and Rybalov [14], replacing the neural element 1 of the t-norm or neural element 0 of the t-conorm by a value e lying anywhere in [0,1].

On the other hand, a lattice [2] is a partially ordered set (L, \leq) satisfying that every pair of two elements $x, y \in L$ have a greatest lower bound, called *infimum* and denoted as $x \wedge y$, as well as a smallest upper bound, called *supremum* and denoted as $x \vee y$. A lattice is bounded if it has a top element and a bottom element, written as 0 and 1, respectively. Let $(L, \leq, 0, 1)$ denote a bounded lattice with top element 1 and bottom element 0 throughout this paper.

For $x, y \in L$, the expression x < y means that $x \le y$ and $x \ne y$. The elements x and y in L are comparable if $x \le y$ or $y \le x$. Otherwise, x and y are called incomparable if $x \nleq y$ and $y \nleq x$, and in this case, it is denoted as x || y. Let $I_e = \{x \in L : x || e\}$ for $e \in L$.

[★]This work was supported by the National Natural Science Foundation of China (No. 11601449), the Science and Technology Innovation Team of Education Department of Sichuan for Dynamical System and its Applications (No. 18TD0013), the Youth Science and Technology Innovation Team of Southwest Petroleum University for Nonlinear Systems (No. 2017CXTD02), and the National Natural Science Foundation of China (Key Program) (No. 51534006).

^{*}Corresponding author

Email addresses: wuxinxing5201314@163.com (Xinxing Wu), gchen@ee.cityu.edu.hk (Guanrong Chen)

Definition 1. [2] Let $(L, \leq, 0, 1)$ be a bounded lattice and $a, b \in L$ with $a \leq b$. Define a subinterval [a, b] as

$$[a, b] = \{x \in L : a \le x \le b\}.$$

Other subintervals such as [a, b) and (a, b) are defined similarly. Let $A(e) = ([0, e] \times [e, 1]) \cup ([e, 1] \times [0, e])$.

Definition 2. [9, 14] Let $(L, \leq, 0, 1)$ be a bounded lattice. A binary operation $U: L^2 \longrightarrow L$ is called a *uninorm on* L if it is commutative, associative, increasing with respect to both variables and there exist some element $e \in L$ called the *neutral element* such that U(e, x) = x for all $x \in L$. In particular, an operation $T: L^2 \longrightarrow L$ is called a t-norm (t-conorm) if it is commutative, associative, increasing with respect to both variables and has a neutral element e = 1 (e = 0).

Uninorms on bounded lattices were originated from the work of Karaçal and Mesiar [9], whose results guarantee the existence of uninorms with any neural element e on any bounded lattice L. Meanwhile, they also constructed the smallest uninorm and the greatest uninorm with the neural element $e \in L$. Recently, constructing new uninorms on bounded lattices was investigated by many authors (see [3, 4, 5, 6, 7, 8, 11, 13]).

Caylı and Karaçal [5] obtained the following results:

Theorem 1. [5, Theorem 3.9] Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. Suppose that $x \vee y \in I_e$ for all $x, y \in I_e$. If T_e is a t-norm on [0, e], then the function $U_T : L \times L \longrightarrow L$ defined by

$$U_{T}(x,y) = \begin{cases} T_{e}(x,y), & (x,y) \in [0,e]^{2}, \\ 1, & (x,y) \in (e,1]^{2}, \\ y, & (x,y) \in [0,e] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [0,e], \\ x \vee y, & otherwise, \end{cases}$$

$$(1.1)$$

is a uninorm on L with the neutral element e.

Theorem 2. [5, Theorem 3.12] Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. Suppose that $x \land y \in I_e$ for all $x, y \in I_e$. If S_e is a t-conorm on [e, 1], then the function $U_S: L \times L \longrightarrow L$ defined by

$$U_{S}(x,y) = \begin{cases} 0, & (x,y) \in [0,e)^{2}, \\ S_{e}(x,y), & (x,y) \in [e,1]^{2}, \\ y, & (x,y) \in [0,e] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [0,e], \\ x \wedge y, & otherwise, \end{cases}$$
(1.2)

is a uninorm on L with the neutral element e.

Theorem 3. [5, Theorem 3.1] Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. Suppose that either $x \vee y > e$ for all $x, y \in I_e$ or $x \vee y \in I_e$ for all $x, y \in I_e$. If T_e is a t-norm on [0, e], then the function $U_t : L \times L \longrightarrow L$ defined by

$$U_{t}(x,y) = \begin{cases} T_{e}(x,y), & (x,y) \in [0,e]^{2}, \\ x \vee y, & (x,y) \in A(e) \cup (I_{e} \times I_{e}), \\ y, & (x,y) \in [0,e] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [0,e], \\ 1, & otherwise, \end{cases}$$
(1.3)

is a uninorm on L with the neutral element e.

Theorem 4. [5, Theorem 3.5] Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. Suppose that either $x \wedge y < e$ for all $x, y \in I_e$ or $x \wedge y \in I_e$ for all $x, y \in I_e$. If S_e is a t-conorm on [e, 1], then the function $U_s : L \times L \longrightarrow L$ defined by

$$U_{s}(x,y) = \begin{cases} S_{e}(x,y), & (x,y) \in [e,1]^{2}, \\ x \wedge y, & (x,y) \in A(e) \cup (I_{e} \times I_{e}), \\ y, & (x,y) \in [e,1] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [e,1], \\ 0, & otherwise, \end{cases}$$
(1.4)

is a uninorm on L with the neutral element e.

For $e \in L \setminus \{0,1\}$, it is clear that $(0,e) \in [0,e]^2 \cap A(e)$ and $(e,1) \in [e,1]^2 \cap A(e)$. According to the method of Çaylı and Karaçal [5] in proving the above Theorems 3 and 4, it follows that

$$0 = T_e(0, e) = U_t(0, e) = 0 \lor e = e,$$

and

$$1 = S_e(e, 1) = U_s(e, 1) = e \land 1 = e,$$

are both impossible. Meanwhile, consider the complete lattice L_1 with the Hasse diagram shown in Fig. 1 and take $S_e: \{e, a, 1\}^2 \longrightarrow \{e, a, 1\}$ by $S_e(x, y) = x \lor y$ for $(x, y) \in \{e, a, 1\}^2$. Clearly, $I_e = \{b\}$. This implies that $x \land y = b \in I_e$ for all $x, y \in I_e$, i.e., e satisfies the hypothesis in Theorem 2. Applying Theorem 2 (i.e., [5, Theorem 3.12]) yields that U_S defined by (1.2) is a uninorm on L_1 . However, $U_S(0, a) = 0 < b = U_S(0, b)$, which is impossible, because a > b. This shows that [5, Theorem 3.12] does not hold. Thus, we modify the above Theorems 2, 3, and 4 in the following (see Theorems 5, 6, and 7) and obtain some equivalent characterizations of the binary operation defined by 1.1 to be uninorms. Similarly, we also give some equivalent characterizations of the binary operations constructed by Aşıcı and Mesiar [1] to be uninorms. These extend the main results in [1, 5].




Figure 1: The Hasse diagram of the lattice L_1

2. Modification of Theorem 2

This section is devoted to correcting Theorem 2 above and characterizing uninorm property for the operation induced by a conorm on [e,1] (see Theorem 5 below). In particular, our result shows that this operation is a uninorm with the neural element e if and only if $I_e \wedge I_e \subset I_e \cup \{0\}$, implying that the uninorm property of this operation is only related to the property of e in the lattice L.

Theorem 5. Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. If S_e is a t-conorm on [e, 1] and the function $U_S : L \times L \longrightarrow L$ is defined by

$$U_{S}(x,y) = \begin{cases} 0, & (x,y) \in [0,e)^{2}, \\ S_{e}(x,y), & (x,y) \in [e,1]^{2}, \\ y, & (x,y) \in [e,1] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [e,1], \\ x \wedge y, & otherwise, \end{cases}$$

$$(2.1)$$

then the following statements are equivalent:

- (i) U_S is a uninorm on L with the neutral element e;
- (ii) $I_e = \emptyset$; otherwise, for any $y, z \in I_e$, it holds that $y \land z \in I_e$ or $y \land z = 0$;
- (iii) $I_e \wedge I_e \subset I_e \cup \{0\}$;
- (iv) \wedge is a t-norm on $I_e \cup \{0,1\}$.

Proof. Clearly, (ii) \iff (iv). It remains to show that (i) \iff (ii).

(i) \Longrightarrow (ii). Suppose, on the contrary, that there exist $y_0, z_0 \in I_e$ such that $y_0 \wedge z_0 \notin I_e$ and $y_0 \wedge z_0 > 0$. This implies that $y_0 \wedge z_0 \in (0, e)$. Then,

$$U_S(y_0 \wedge z_0, U_S(y_0, z_0)) = U_S(y_0 \wedge z_0, y_0 \wedge z_0) = 0,$$

and

$$U_S(U_S(y_0 \wedge z_0, y_0), z_0) = U_S(y_0 \wedge z_0, z_0) = y_0 \wedge z_0 > 0,$$




Figure 2: Structure of U_S in Theorem 5.

which contradict with the associativity of U_S .

- (ii) \Longrightarrow (i). Clearly, U_S is a uninorm when $I_e = \emptyset$. Without loss of generality, assume that $I_e \neq \emptyset$. Firstly, it is not difficult to verify that U_S is a commutative binary operation with the neutral element e.
- i.1) Monotonicity. For any $x, y \in L$ with $x \leq y$, it holds that $U_S(x, z) \leq U_S(y, z)$ for all $z \in L$. Consider the following cases:
 - 1. $x \in [0, e)$.
 - 1.1. If $y \in [0, e)$, then

$$U_S(x,z) = \begin{cases} 0, & z \in [0,e), \\ x, & z = e, \\ x, & z \in (e,1], \\ x \wedge z, & z \in I_e, \end{cases}$$
 (2.2)

and

$$U_S(y,z) = \begin{cases} 0, & z \in [0,e), \\ y, & z = e, \\ y, & z \in (e,1], \\ y \land z, & z \in I_e, \end{cases}$$

implying that $U_S(x,z) \leq U_S(y,z)$.

1.2. If $y \in (e, 1]$, then,

$$U_S(y,z) = \begin{cases} z, & z \in [0,e), \\ y, & z = e, \\ S_e(y,z), & z \in (e,1], \\ z, & z \in I_e. \end{cases}$$

This, together with (2.2) and $S_e(y,z) \geq y$, implies that $U_S(x,z) \leq U_S(y,z)$.

1.3. If $y \in I_e$, then

$$U_{S}(y,z) = \begin{cases} y \wedge z, & z \in [0,e), \\ y, & z = e, \\ y, & z \in (e,1], \\ y \wedge z, & z \in I_{e}. \end{cases}$$
 (2.3)

This, together with (2.2), implies that $U_S(x,z) \leq U_S(y,z)$.

- 1.4. If y = e, then $U_S(y, z) = z$. This, together with (2.2), implies that $U_S(x, z) \le U_S(y, z)$.
- 2. $x \in (e, 1]$. From $x \leq y$, it follows that $y \in (e, 1]$. Then

$$U_S(x,z) = \begin{cases} z, & z \in [0,e), \\ x, & z = e, \\ S_e(x,z), & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$
 (2.4)

and

$$U_{S}(y,z) = \begin{cases} z, & z \in [0,e), \\ y, & z = e, \\ S_{e}(y,z), & z \in (e,1], \\ z, & z \in I_{e}, \end{cases}$$
 (2.5)

implying that $U_S(x,z) \leq U_S(y,z)$, because T_e is increasing.

- 3. x = e. This implies that $y \in [e, 1]$. Clearly, $U_S(x, z) = z = U_S(y, z)$ if y = e. If $y \in (e, 1]$, applying (2.5) and $U_S(x, z) = z$, it follows that $U_S(x, z) \leq U_S(y, z)$.
- 4. $x \in I_e$. From $x \leq y$, it follows that $y \in (e, 1] \cup I_e$. Applying (2.1) yields that

$$U_{S}(x,z) = \begin{cases} x \wedge z, & z \in [0,e), \\ x, & z = e, \\ x, & z \in (e,1], \\ x \wedge z, & z \in I_{e}. \end{cases}$$
 (2.6)

- 4.1. If $y \in (e, 1]$, from (2.5) and (2.6), it follows that $U_S(x, z) \leq U_S(y, z)$.
- 4.2. If $y \in I_e$, from (2.3) and (2.6), it is clear that $U_S(x,z) \leq U_S(y,z)$.
- i.2) Associativity. For any $x, y, z \in L$, it holds that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$. If one of the elements x, y and z is equal to e, then the equality is always satisfied. Otherwise, consider the following cases:

- 1. $x \in [0, e)$.
 - 1.1. $y \in [0, e)$.
 - 1.1.1. If $z \in [0, e)$, then $U_S(x, U_S(y, z)) = 0 = U_S(U_S(x, y), z)$.
 - 1.1.2. If $z \in (e, 1]$, then $U_S(x, U_S(y, z)) = U_S(x, y) = 0$ and $U_S(U_S(x, y), z) = U_S(0, z) = 0$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.1.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, y \wedge z) = 0$ and $U_S(U_S(x, y), z) = U_S(0, z) = 0$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.2. $y \in (e, 1]$.
 - 1.2.1. If $z \in [0, e)$, then $U_S(x, U_S(y, z)) = U_S(x, z) = 0$ and $U_S(U_S(x, y), z) = U_S(x, z) = 0$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.2.2. If $z \in (e, 1]$, then $U_S(x, U_S(y, z)) = U_S(x, S_e(y, z)) = x$ and $U_S(U_S(x, y), z) = U_S(x, z) = x$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.2.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, z) = x \wedge z$ and $U_S(U_S(x, y), z) = U_S(x, z) = x \wedge z$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.3. $y \in I_e$.
 - 1.3.1. If $z \in [0, e)$, then $U_S(x, U_S(y, z)) = U_S(x, y \wedge z) = 0$ and $U_S(U_S(x, y), z) = U_S(x \wedge y, z) = 0$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.3.2. If $z \in (e, 1]$, then $U_S(x, U_S(y, z)) = U_S(x, y) = x \wedge y$ and $U_S(U_S(x, y), z) = U_S(x \wedge y, z) = x \wedge y$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
 - 1.3.3. If $z \in I_e$, then

$$U_S(x, U_S(y, z)) = U_S(x, y \wedge z)$$

$$= \begin{cases} x \wedge y \wedge z, & y \vee z \in I_e, \\ 0, & y \wedge z = 0, \end{cases}$$

$$= x \wedge y \wedge z.$$

and $U_S(U_S(x,y),z) = U_S(x \wedge y,z) = x \wedge y \wedge z$, implying that $U_S(x,U_S(y,z)) = U_S(U_S(x,y),z)$.

- 2. $x \in (e, 1]$.
 - $2.1. y \in [0, e).$
 - 2.1.1. If $z \in [0, e)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.1.2)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

2.1.2. If $z \in (e, 1]$, then $U_S(x, U_S(y, z)) = U_S(x, z)$ and $U_S(U_S(x, y), z) = U_S(x, z)$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.

- 2.1.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, z)$ and $U_S(U_S(x, y), z) = U_S(x, z)$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
- 2.2. $y \in (e, 1]$.
 - 2.2.1. If $z \in [0, a_1)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.2.2)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

- 2.2.2. If $z \in (e,1]$, then $U_S(x, U_S(y,z)) = U_S(x, S_e(y,z)) = S_e(x, S_e(y,z)) = S_e(S_e(x,y),z) = S_e(U_S(x,y),z) = U_S(U_S(x,y),z)$.
- 2.2.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, z) = z$ and $U_S(U_S(x, y), z) = U_S(S_e(x, y), z) = z$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
- 2.3. $y \in I_e$.
 - 2.3.1. If $z \in [0, e)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.3.2)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

- 2.3.2. If $z \in (e, 1]$, then $U_S(x, U_S(y, z)) = U_S(x, y) = y$ and $U_S(U_S(x, y), z) = U_S(y, z) = y$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
- 2.3.3. If $z \in I_e$, then

$$U_S(x, U_S(y, z)) = U_S(x, y \lor z)$$

$$= \begin{cases} y \land z, & y \land z \in I_e, \\ 0, & y \land z = 0, \end{cases}$$

$$= y \land z,$$

and $U_S(U_S(x,y),z) = U_S(y,z) = y \wedge z$, implying that $U_S(x,U_S(y,z)) = U_S(U_S(x,y),z)$.

- 3. $x \in I_e$.
 - 3.1. $y \in [0, e)$.
 - 3.1.1. If $z \in [0, e)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.1.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

3.1.2. If $z \in (e, 1]$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 2.1.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

- 3.1.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, y \wedge z) = x \wedge y \wedge z$ and $U_S(U_S(x, y), z) = U_S(x \wedge y, z) = x \wedge y \wedge z$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
- 3.2. $y \in (e, 1]$.
 - 3.2.1. If $z \in [0, e)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.2.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

3.2.2. If $z \in (e, 1]$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 2.2.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

- 3.2.3. If $z \in I_e$, then $U_S(x, U_S(y, z)) = U_S(x, z) = x \wedge z$ and $U_S(U_S(x, y), z) = U_S(x, z) = x \wedge z$, implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.
- 3.3. $y \in I_e$.
 - 3.3.1. If $z \in [0, e)$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 1.3.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

3.3.2. If $z \in (e, 1]$, then

$$U_S(x, U_S(y, z)) = U_S(U_S(z, y), x)$$
 (commutativity of U_S)
= $U_S(z, U_S(y, x))$ (by 2.3.3)
= $U_S(U_S(x, y), z)$ (commutativity of U_S).

3.3.3. If $z \in I_e$, then

$$U_S(x, U_S(y, z)) = U_S(x, y \wedge z)$$

$$= \begin{cases} x \wedge y \wedge z, & y \wedge z \in I_e, \\ 0, & y \wedge z = 0, \end{cases}$$

$$= x \wedge y \wedge z,$$

and

$$U_S(U_S(x,y),z) = U_S(x \wedge y, z)$$

$$= \begin{cases} x \wedge y \wedge z, & x \wedge y \in I_e, \\ 0, & x \wedge y = 0, \end{cases}$$

$$= x \wedge y \wedge z,$$

implying that $U_S(x, U_S(y, z)) = U_S(U_S(x, y), z)$.

3. Modifications of Theorems 3 and 4

Similarly to Section 2, this section is devoted to correcting Theorems 3 and 4 and characterizing the uninorm property (see Theorems 5 and 7) for some new binary operations on the lattices.

Theorem 6. Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. If T_e is a t-norm on [0, e] and the function $U_t : L \times L \longrightarrow L$ is defined by

$$U_{t}(x,y) = \begin{cases} T_{e}(x,y), & (x,y) \in [0,e]^{2}, \\ x \vee y, & (x,y) \in ([0,e] \times (e,1]) \cup ((e,1] \times [0,e]) \cup (I_{e} \times I_{e}), \\ y, & (x,y) \in [0,e] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [0,e], \\ 1, & otherwise, \end{cases}$$
(3.1)

then the following statements are equivalent:

- (i) U_t is a uninorm on L with the neutral element e;
- (ii) $I_e = \emptyset$; otherwise, for any $y, z \in I_e$, it holds that $y \lor z \in I_e$ or $y \lor z = 1$;
- (iii) $I_e \vee I_e \subset I_e \cup \{1\};$
- (iv) \vee is a t-conorm on $I_e \cup \{0,1\}$.

Proof. Clearly, (ii) \iff (iii) \iff (iv). It remains to show that (i) \iff (ii).

(i) \Longrightarrow (ii). Suppose, on the contrary, that there exist $y_0, z_0 \in I_e$ such that $y_0 \lor z_0 \notin I_e$ and $y_0 \lor z_0 < 1$. This implies that $y_0 \lor z_0 \in (e, 1)$. Consequently,

$$U_t(y_0, U_t(y_0, z_0)) = U_t(y_0, y_0 \lor z_0) = 1,$$

and

$$U_t(U_t(y_0, y_0), z_0) = U_t(y_0, z_0) = y_0 \lor z_0 < 1,$$




Figure 3: Structure of U_t in Theorem 6.

which contradict with the associativity of U_t .

(ii) \Longrightarrow (i). Clearly, U_t is a uninorm when $I_e = \emptyset$. Without loss of generality, assume that $I_e \neq \emptyset$. Firstly, it is not difficult to verify that U_t is a commutative binary operation with the neutral element e.

i.1) Monotonicity. For any $x, y \in L$ with $x \leq y$, it holds that $U_t(x, z) \leq U_t(y, z)$ for all $z \in L$. Consider the following cases:

1. $x \in [0, e)$.

1.1. If $y \in [0, e)$, then

$$U_{t}(x,z) = \begin{cases} T_{e}(x,z), & z \in [0,e), \\ x, & z = e, \\ z, & z \in (e,1], \\ z, & z \in I_{e}, \end{cases}$$
(3.2)

and

$$U_t(y,z) = \begin{cases} T_e(y,z), & z \in [0,e), \\ y, & z = e, \\ z, & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$

implying that $U_t(x, z) \leq U_t(y, z)$, because T_e is increasing.

1.2. If $y \in (e, 1]$, then,

$$U_t(y, z) = \begin{cases} y, & z \in [0, e), \\ y, & z = e, \\ 1, & z \in (e, 1], \\ 1, & z \in I_e. \end{cases}$$

This, together with (3.2) and $T_e(x,z) \leq x$, implies that $U_t(x,z) \leq U_t(y,z)$.

1.3. If $y \in I_e$, then

$$U_{t}(y,z) = \begin{cases} y, & z \in [0,e), \\ y, & z = e, \\ 1, & z \in (e,1], \\ y \lor z, & z \in I_{e}. \end{cases}$$
 (3.3)

This, together with (3.2) and $T_e(x,z) \leq x$, implies that $U_t(x,z) \leq U_t(y,z)$.

- 1.4. If y = e, then $U_t(y, z) = z$. This, together with (3.2), implies that $U_t(x, z) \le U_t(y, z)$.
- 2. $x \in (e, 1]$. From $x \leq y$, it follows that $y \in (e, 1]$. Then,

$$U_t(x,z) = \begin{cases} x, & z \in [0,e), \\ x, & z = e, \\ 1, & z \in (e,1], \\ 1, & z \in I_e, \end{cases}$$
 (3.4)

and

$$U_t(y,z) = \begin{cases} y, & z \in [0,e), \\ y, & z = e, \\ 1, & z \in (e,1], \\ 1, & z \in I_e, \end{cases}$$
 (3.5)

implying that $U_t(x,z) \leq U_t(y,z)$.

- 3. x = e. This implies that $y \in [e, 1]$. Clearly, $U_t(x, z) = z = U_t(y, z)$ if y = e. If $y \in (e, 1]$, applying (3.5) and $U_t(x, z) = z$, it follows that $U_t(x, z) \leq U_t(y, z)$.
- 4. $x \in I_e$. From $x \leq y$, it follows that $y \in (e, 1] \cup I_e$. Applying (3.1) yields that

$$U_t(x,z) = \begin{cases} x, & z \in [0,e), \\ x, & z = e, \\ 1, & z \in (e,1], \\ x \lor z, & z \in I_e. \end{cases}$$
 (3.6)

4.1. If $y \in (e, 1]$, from (3.5) and (3.6), it follows that $U_t(x, z) \leq U_t(y, z)$.

- 4.2. If $y \in I_e$, from (3.3) and (3.6), it is clear that $U_t(x,z) \leq U_t(y,z)$.
- i.2) Associativity. For any $x, y, z \in L$, it holds that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$. If one of the elements x, y and z is equal to e, then the equality is always satisfied. Otherwise, consider the following cases:
 - 1. $x \in [0, e)$.
 - 1.1. $y \in [0, e)$.
 - 1.1.1. If $z \in [0, e)$, then $U_t(x, U_t(y, z)) = U_t(x, T_e(y, z)) = T_e(x, T_e(y, z)) = T_e(T_e(x, y), z) = T_e(U_t(x, y), z) = U_t(U_t(x, y), z)$.
 - 1.1.2. If $z \in (e, 1]$, then $U_t(x, U_t(y, z)) = U_t(x, z) = z$ and $U_t(U_t(x, y), z) = U_t(T_e(x, y), z) = z$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.1.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, z) = z$ and $U_t(U_t(x, y), z) = U_t(T_e(x, y), z) = z$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.2. $y \in (e, 1]$.
 - 1.2.1. If $z \in [0, e)$, then $U_t(x, U_t(y, z)) = U_t(x, y) = y$ and $U_t(U_t(x, y), z) = U_t(y, z) = y$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.2.2. If $z \in (e, 1]$, then $U_t(x, U_t(y, z)) = U_t(x, 1) = 1$ and $U_t(U_t(x, y), z) = U_t(y, z) = 1$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.2.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, 1) = 1$ and $U_t(U_t(x, y), z) = U_t(y, z) = 1$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.3. $y \in I_e$.
 - 1.3.1. If $z \in [0, e)$, then $U_t(x, U_t(y, z)) = U_t(x, y) = y$ and $U_t(U_t(x, y), z) = U_t(y, z) = y$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
 - 1.3.2. If $z \in (e,1]$, then $U_t(x, U_t(y,z)) = U_T(x,1) = 1$ and $U_t(U_t(x,y),z) = U_t(y,z) = 1$, implying that $U_t(x, U_t(y,z)) = U_t(U_t(x,y),z)$.
 - 1.3.3. If $z \in I_e$, then

$$U_t(x, U_t(y, z)) = U_t(x, y \lor z)$$

$$= \begin{cases} y \lor z, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= y \lor z,$$

and $U_t(U_t(x,y),z) = U_t(y,z) = y \vee z$, implying that $U_t(x,U_t(y,z)) = U_t(U_t(x,y),z)$.

- 2. $x \in (e, 1]$.
 - 2.1. $y \in [0, e)$.

2.1.1. If $z \in [0, e)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x)$$
 (commutativity of U_t)
= $U_t(z, U_t(y, x))$ (by 1.1.2)
= $U_t(U_t(x, y), z)$ (commutativity of U_t).

- 2.1.2. If $z \in (e, 1]$, then $U_t(x, U_t(y, z)) = U_t(x, z)$ and $U_t(U_t(x, y), z) = U_t(x, z)$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
- 2.1.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, z)$ and $U_t(U_t(x, y), z) = U_t(x, z)$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
- 2.2. $y \in (e, 1]$.
 - 2.2.1. If $z \in [0, a_1)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x) \quad \text{(commutativity of } U_t)$$
$$= U_t(z, U_t(y, x)) \quad \text{(by 1.2.2)}$$
$$= U_t(U_t(x, y), z) \quad \text{(commutativity of } U_t).$$

- 2.2.2. If $z \in (e,1]$, then $U_t(x, U_t(y,z)) = U_t(x,1) = 1$ and $U_t(U_t(x,y),z) = U_t(1,z) = 1$, implying that $U_t(x, U_t(y,z)) = U_t(U_t(x,y),z)$.
- 2.2.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, 1) = 1$ and $U_t(U_t(x, y), z) = U_t(1, z) = 1$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
- 2.3. $y \in I_e$.
 - 2.3.1. If $z \in [0, e)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x) \quad \text{(commutativity of } U_t)$$

$$= U_t(z, U_t(y, x)) \quad \text{(by 1.3.2)}$$

$$= U_t(U_t(x, y), z) \quad \text{(commutativity of } U_t).$$

- 2.3.2. If $z \in (e,1]$, then $U_t(x, U_t(y,z)) = U_t(x,1) = 1$ and $U_t(U_t(x,y),z) = U_t(1,z) = 1$, implying that $U_t(x, U_t(y,z)) = U_t(U_t(x,y),z)$.
- 2.3.3. If $z \in I_e$, then

$$U_t(x, U_t(y, z)) = U_t(x, y \lor z)$$

$$= \begin{cases} 1, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= 1,$$

and $U_t(U_t(x,y),z) = U_t(1,z) = 1$, implying that $U_t(x,U_t(y,z)) = U_t(U_t(x,y),z)$.

- 3. $x \in I_e$.
 - 3.1. $y \in [0, e)$.

3.1.1. If $z \in [0, e)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x)$$
 (commutativity of U_t)
= $U_t(z, U_t(y, x))$ (by 1.1.3)
= $U_t(U_t(x, y), z)$ (commutativity of U_t).

3.1.2. If $z \in (e, 1]$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x) \quad \text{(commutativity of } U_t)$$
$$= U_t(z, U_t(y, x)) \quad \text{(by 2.1.3)}$$
$$= U_t(U_t(x, y), z) \quad \text{(commutativity of } U_t).$$

- 3.1.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, z)$ and $U_t(U_t(x, y), z) = U_t(x, z)$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
- 3.2. $y \in (e, 1]$.
 - 3.2.1. If $z \in [0, e)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x)$$
 (commutativity of U_t)
= $U_t(z, U_t(y, x))$ (by 1.2.3)
= $U_t(U_t(x, y), z)$ (commutativity of U_t).

3.2.2. If $z \in (e, 1]$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x)$$
 (commutativity of U_t)
= $U_t(z, U_t(y, x))$ (by 2.2.3)
= $U_t(U_t(x, y), z)$ (commutativity of U_t).

- 3.2.3. If $z \in I_e$, then $U_t(x, U_t(y, z)) = U_t(x, 1) = 1$ and $U_t(U_t(x, y), z) = U_t(1, z) = 1$, implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.
- 3.3. $y \in I_e$.
 - 3.3.1. If $z \in [0, e)$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x)$$
 (commutativity of U_t)
= $U_t(z, U_t(y, x))$ (by 1.3.3)
= $U_t(U_t(x, y), z)$ (commutativity of U_t).

3.3.2. If $z \in (e, 1]$, then

$$U_t(x, U_t(y, z)) = U_t(U_t(z, y), x) \quad \text{(commutativity of } U_t)$$
$$= U_t(z, U_t(y, x)) \quad \text{(by 2.3.3)}$$
$$= U_t(U_t(x, y), z) \quad \text{(commutativity of } U_t).$$

3.3.3. If $z \in I_e$, then

$$U_t(x, U_t(y, z)) = U_t(x, y \lor z)$$

$$= \begin{cases} x \lor y \lor z, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= x \lor y \lor z,$$

and

$$U_t(U_t(x,y),z) = U_t(x \lor y,z)$$

$$= \begin{cases} x \lor y \lor z, & x \lor y \in I_e, \\ 1, & x \lor y = 1, \end{cases}$$

$$= x \lor y \lor z,$$

implying that $U_t(x, U_t(y, z)) = U_t(U_t(x, y), z)$.

Theorem 7. Let $(X, \leq, 0, 1)$ be a bounded lattice and $e \in L \setminus \{0, 1\}$. If S_e is a t-conorm on [e, 1] and the function $U_s : L \times L \longrightarrow L$ is defined by

$$U_{s}(x,y) = \begin{cases} S_{e}(x,y), & (x,y) \in [e,1]^{2}, \\ x \wedge y, & (x,y) \in ([0,e) \times [e,1]) \cup ([e,1] \times [0,e)) \cup (I_{e} \times I_{e}), \\ y, & (x,y) \in [e,1] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [e,1], \\ 0, & otherwise, \end{cases}$$
(3.7)

then the following statements are equivalent:

- (i) U_s is a uninorm on L with the neutral element e;
- (ii) $I_e = \emptyset$; otherwise, for any $y, z \in I_e$, it holds that $y \land z \in I_e$ or $y \land z = 0$;
- (iii) $I_e \wedge I_e \subset I_e \cup \{0\};$
- (iv) \wedge is a t-norm on $I_e \cup \{0,1\}$.

Proof. It can be proved similarly to the proof of Theorem 6.

4. Characterizations of binary operations defined by 1.1

Theorem 8. Let $(X, \leq, 0, 1)$ be a bounded lattice, $e \in L \setminus \{0, 1\}$, and T_e be a t-norm on [0, e]. Then, the following statements are equivalent:

- (i) U_T defined by (1.1) is a uninorm on L with the neutral element e;
- (ii) $I_e = \emptyset$; otherwise, for any $y, z \in I_e$, it holds that $y \lor z \in I_e$ or $y \lor z = 1$;
- (iii) $I_e \vee I_e \subset I_e \cup \{1\};$
- (iv) \vee is a t-conorm on $I_e \cup \{0,1\}$.

Proof. Clearly, (ii) \iff (iv). It remains to show that (i) \iff (ii).

(i) \Longrightarrow (ii). Suppose, on the contrary, that there exist $y_0, z_0 \in I_e$ such that $y_0 \lor z_0 \notin I_e$ and $y_0 \lor z_0 < 1$. This implies that $y_0 \lor z_0 \in (e, 1)$. Then,

$$U_T(y_0 \lor z_0, U_T(y_0, z_0)) = U_T(y_0 \lor z_0, y_0 \lor z_0) = 1,$$

and

$$U_T(U_T(y_0 \lor z_0, y_0), z_0) = U_T(y_0 \lor z_0, z_0) = y_0 \lor z_0 < 1,$$

which contradict with the associativity of U_T .

- (ii) \Longrightarrow (i). Clearly, U_T is a uninorm when $I_e = \emptyset$. Without loss of generality, assume that $I_e \neq \emptyset$. Firstly, it is not difficult to verify that U_T is a commutative binary operation with the neutral element e.
- i.1) Monotonicity. For any $x, y \in L$ with $x \leq y$, it holds that $U_T(x, z) \leq U_T(y, z)$ for all $z \in L$. Consider the following cases:
 - 1. $x \in [0, e)$.
 - 1.1. If $y \in [0, e)$, then

$$U_T(x,z) = \begin{cases} T_e(x,z), & z \in [0,e), \\ x, & z = e, \\ z, & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$
(4.1)

and

$$U_T(y,z) = \begin{cases} T_e(y,z), & z \in [0,e), \\ y, & z = e, \\ z, & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$

implying that $U_T(x,z) \leq U_T(y,z)$, because T_e is increasing.

1.2. If $y \in (e, 1]$, then,

$$U_T(y,z) = \begin{cases} y, & z \in [0,e), \\ y, & z = e, \\ 1, & z \in (e,1], \\ y \lor z, & z \in I_e. \end{cases}$$

This, together with (4.1) and $T_e(x,z) \leq x$, implies that $U_T(x,z) \leq U_T(y,z)$.

1.3. If $y \in I_e$, then

$$U_{T}(y,z) = \begin{cases} y, & z \in [0,e), \\ y, & z = e, \\ y \lor z, & z \in (e,1], \\ y \lor z, & z \in I_{e}. \end{cases}$$
(4.2)

This, together with (4.1) and $T_e(x,z) \leq x$, implies that $U_T(x,z) \leq U_T(y,z)$.

- 1.4. If y = e, then $U_T(y, z) = z$. This, together with (4.1), implies that $U_T(x, z) \le U_T(y, z)$.
- 2. $x \in (e, 1]$. From $x \leq y$, it follows that $y \in (e, 1]$. Then,

$$U_T(x,z) = \begin{cases} x, & z \in [0,e), \\ x, & z = e, \\ 1, & z \in (e,1], \\ x \lor z, & z \in I_e, \end{cases}$$
(4.3)

and

$$U_T(y,z) = \begin{cases} y, & z \in [0,e), \\ y, & z = e, \\ 1, & z \in (e,1], \\ y \lor z, & z \in I_e, \end{cases}$$
(4.4)

implying that $U_T(x,z) \leq U_T(y,z)$.

- 3. x = e. This implies that $y \in [e, 1]$. Clearly, $U_T(x, z) = z = U_T(y, z)$ if y = e. If $y \in (e, 1]$, applying (4.4) and $U_T(x, z) = z$, it follows that $U_T(x, z) \leq U_T(y, z)$.
- 4. $x \in I_e$. From $x \leq y$, it follows that $y \in (e, 1] \cup I_e$. Applying (1.1) yields that

$$U_{T}(x,z) = \begin{cases} x, & z \in [0,e), \\ x, & z = e, \\ x \lor z, & z \in (e,1], \\ x \lor z, & z \in I_{e}. \end{cases}$$
(4.5)

- 4.1. If $y \in (e, 1]$, from (4.4) and (4.5), it follows that $U_T(x, z) \leq U_T(y, z)$.
- 4.2. If $y \in I_e$, from (4.2) and (4.5), it is clear that $U_T(x,z) \leq U_T(y,z)$.
- i.2) Associativity. For any $x, y, z \in L$, it holds that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$. If one of the elements x, y and z is equal to e, then the equality is always satisfied. Otherwise, consider the following cases:
 - 1. $x \in [0, e)$.

- 1.1. $y \in [0, e)$.
 - 1.1.1. If $z \in [0, e)$, then $U_T(x, U_T(y, z)) = U_T(x, T_e(y, z)) = T_e(x, T_e(y, z)) = T_e(T_e(x, y), z) = T_e(U_T(x, y), z) = U_T(U_T(x, y), z)$.
 - 1.1.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, z) = z$ and $U_T(U_T(x, y), z) = U_T(T_e(x, y), z) = z$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
 - 1.1.3. If $z \in I_e$, then $U_T(x, U_T(y, z)) = U_T(x, z) = z$ and $U_T(U_T(x, y), z) = U_T(T_e(x, y), z) = z$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 1.2. $y \in (e, 1]$.
 - 1.2.1. If $z \in [0, e)$, then $U_T(x, U_T(y, z)) = U_T(x, y) = y$ and $U_T(U_T(x, y), z) = U_T(y, z) = y$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
 - 1.2.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, 1) = 1$ and $U_T(U_T(x, y), z) = U_T(y, z) = 1$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
 - 1.2.3. If $z \in I_e$, then $U_T(x, U_T(y, z)) = U_T(x, y \vee z) = y \vee z$ and $U_T(U_T(x, y), z) = U_T(y, z) = y \vee z$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 1.3. $y \in I_e$.
 - 1.3.1. If $z \in [0, e)$, then $U_T(x, U_T(y, z)) = U_T(x, y) = y$ and $U_T(U_T(x, y), z) = U_T(y, z) = y$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
 - 1.3.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, y \lor z) = y \lor z$ and $U_T(U_T(x, y), z) = U_T(y, z) = y \lor z$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
 - 1.3.3. If $z \in I_e$, then

$$U_T(x, U_T(y, z)) = U_T(x, y \lor z)$$

$$= \begin{cases} y \lor z, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= y \lor z,$$

and $U_T(U_T(x,y),z) = U_T(y,z) = y \vee z$, implying that $U_T(x,U_T(y,z)) = U_T(U_T(x,y),z)$.

- 2. $x \in (e, 1]$.
 - $2.1. y \in [0, e).$
 - 2.1.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.1.2)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

2.1.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, z)$ and $U_T(U_T(x, y), z) = U_T(x, z)$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.

2.1.3. If
$$z \in I_e$$
, then $U_T(x, U_T(y, z)) = U_T(x, z)$ and $U_T(U_T(x, y), z) = U_T(x, z)$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.

- 2.2. $y \in (e, 1]$.
 - 2.2.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.2.2)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

- 2.2.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, 1) = 1$ and $U_T(U_T(x, y), z) = U_T(1, z) = 1$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 2.2.3. If $z \in I_e$, then $U_T(x, U_T(y, z)) = U_T(x, y \vee z) = 1$ and $U_T(U_T(x, y), z) = U_T(1, z) = 1$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 2.3. $y \in I_e$.
 - 2.3.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.3.2)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

- 2.3.2. If $z \in (e, 1]$, then $U_T(x, U_T(y, z)) = U_T(x, y \vee z) = 1$ and $U_T(U_T(x, y), z) = U_T(x \vee y, z) = 1$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 2.3.3. If $z \in I_e$, then

$$U_T(x, U_T(y, z)) = U_T(x, y \lor z)$$

$$= \begin{cases} x \lor y \lor z, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= x \lor y \lor z,$$

and $U_T(U_T(x,y),z) = U_T(x \lor y,z) = x \lor y \lor z$, implying that $U_T(x,U_T(y,z)) = U_T(U_T(x,y),z)$.

- 3. $x \in I_e$.
 - 3.1. $y \in [0, e)$.
 - 3.1.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.1.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

3.1.2. If $z \in (e, 1]$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 2.1.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

- 3.1.3. If $z \in I_e$, then $U_T(x, U_T(y, z)) = U_T(x, z)$ and $U_T(U_T(x, y), z) = U_T(x, z)$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 3.2. $y \in (e, 1]$.
 - 3.2.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.2.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

3.2.2. If $z \in (e, 1]$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 2.2.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

- 3.2.3. If $z \in I_e$, then $U_T(x, U_T(y, z)) = U_T(x, y \lor z) = x \lor y \lor z$ and $U_T(U_T(x, y), z) = U_T(x \lor y, z) = x \lor y \lor z$, implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$.
- 3.3. $y \in I_e$.
 - 3.3.1. If $z \in [0, e)$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 1.3.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

3.3.2. If $z \in (e, 1]$, then

$$U_T(x, U_T(y, z)) = U_T(U_T(z, y), x)$$
 (commutativity of U_T)
= $U_T(z, U_T(y, x))$ (by 2.3.3)
= $U_T(U_T(x, y), z)$ (commutativity of U_T).

3.3.3. If $z \in I_e$, then

$$U_T(x, U_T(y, z)) = U_T(x, y \lor z)$$

$$= \begin{cases} x \lor y \lor z, & y \lor z \in I_e, \\ 1, & y \lor z = 1, \end{cases}$$

$$= x \lor y \lor z,$$

and

$$U_T(U_T(x,y),z) = U_T(x \lor y,z)$$

$$= \begin{cases} x \lor y \lor z, & x \lor y \in I_e, \\ 1, & x \lor y = 1, \end{cases}$$

$$= x \lor y \lor z,$$

implying that $U_T(x, U_T(y, z)) = U_T(U_T(x, y), z)$

5. Characterizations of uninorms constructed by Aşıcı and Mesiar [1]

This section is devoted to characterizing the uninorm property for the operations introduced by Aşıcı and Mesiar [1] (see Theorems 11 and 12). First, the incomparability is defined for two subsets of a lattice.

Definition 3. Let $(L, \leq, 0, 1)$ be a bounded lattice and $A, B \subset L$. Then, A and B are comparable if there exist $x \in A$ and $y \in B$ such that x and y are comparable. Otherwise, A and B are incomparable, i.e., x||y for all $x \in A$ and $y \in B$, and in this case, it is denoted as A||B.

Recently, Asici and Mesiar [1] obtained the following results:

Theorem 9. [1, Theorem 6] Let $(L, \leq, 0, 1)$ be a bounded lattice with $e \in L \setminus \{0, 1\}$ such that for all $x \in I_e$ and $y \in (0, e]$ it holds $x \| y$, i.e., $I_e \| (0, e]$. If T_e is a t-norm on [0, e], then the function $U_{T_e}: L^2 \longrightarrow L$ defined by

$$U_{T_{e}}(x,y) = \begin{cases} T_{e}(x,y), & (x,y) \in [0,e]^{2}, \\ y, & (x,y) \in [e,1] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [e,1], \\ 0, & (x,y) \in ([0,e) \times I_{e}) \cup (I_{e} \times [0,e)), \\ x \wedge y, & (x,y) \in ([0,e) \times [e,1]) \cup ([e,1] \times [0,e)) \cup (I_{e} \times I_{e}), \\ x \vee y, & otherwise, \end{cases}$$

$$(5.1)$$

is a uninorm on L with the neutral element e.

Theorem 10. [1, Theorem 7] Let $(L, \leq, 0, 1)$ be a bounded lattice with $e \in L \setminus \{0, 1\}$ such that for all $x \in I_e$ and $y \in (0, e]$ it holds x || y, i.e., $I_e || (0, e]$. If S_e is a t-conorm on [e, 1], then the function $U_{S_e}: L^2 \longrightarrow L$ defined by

$$U_{S_{e}}(x,y) = \begin{cases} S_{e}(x,y), & (x,y) \in [e,1]^{2}, \\ y, & (x,y) \in [e,1] \times I_{e}, \\ x, & (x,y) \in I_{e} \times [e,1], \\ 0, & (x,y) \in ([0,e) \times I_{e}) \cup (I_{e} \times [0,e)), \\ x \wedge y, & (x,y) \in ([0,e) \times [e,1]) \cup ([e,1] \times [0,e)) \cup (I_{e} \times I_{e}), \\ x \vee y, & otherwise, \end{cases}$$

$$(5.2)$$

is a uninorm on L with the neutral element e.

Theorem 11. Let $(X, \leq, 0, 1)$ be a bounded lattice, $e \in L \setminus \{0, 1\}$, and T_e be a t-norm on [0, e]. Then, the following statements are equivalent:

- (i) U_{T_e} defined by (5.1) is a uninorm on L with the neutral element e;
- (ii) The following hold:
 - ii-1) $\mathscr{P} = \emptyset$ or $T_e|_{\mathscr{P} \times [0,e) \cup [0,e) \times \mathscr{P}} \equiv 0$, where $\mathscr{P} = \{x \in (0,e) : \exists y \in I_e, \ x \leq y\};$
 - ii-2) $I_e \wedge I_e \subset I_e \cup \{0\}$.

Proof. $(i) \Longrightarrow (ii)$.

(i) \Longrightarrow ii.1). Suppose, on the contrary, that there exist $x_0 \in [0, e)$ and $y_0 \in \mathscr{P}$ such that $T_e(x_0, y_0) > 0$. From $y_0 \in \mathscr{P}$, it follows that there exists $z \in I_e$ such that $y_0 \leq z_0$. Since U_{T_e} is increasing, applying (5.1) yields that

$$0 < T_e(x_0, y_0) = U_{T_e}(x_0, y_0) \le U_{T_e}(x_0, z_0) = 0,$$

which is a contradiction.

(i) \Longrightarrow ii.2). Suppose that $I_e \wedge I_e \nsubseteq I_e \cup \{0\}$, i.e., there exist $y_1, z_1 \in I_e$ such that $0 < y_1 \wedge z_1$ and $y_1 \wedge z_1 \notin I_e$. This implies that $y_1 \wedge z_1 \in (0, e)$. From (5.1) and the associativity of U_{T_e} , it follows that

$$0 = U_{T_e}(y_1, U_{T_e}(y_1, z_1)) = U_{T_e}(U_{T_e}(y_1, y_1), z_1) = y_1 \land z_1 > 0,$$

which is a contradiction.

- (ii) \Longrightarrow (i). When $\mathscr{P} = \emptyset$, this has been proved in [1, Theorem 6]. Now assume that $T_e|_{\mathscr{P}\times[0,e)\cup[0,e)\times\mathscr{P}}\equiv 0$ and $I_e\wedge I_e\subset I_e\cup\{0\}$. Firstly, it is not difficult to verify that U_{T_e} is a commutative binary operation with the neutral element e.
- i.1) Monotonicity. For any $x, y \in L$ with $x \leq y$, it holds that $U_{T_e}(x, z) \leq U_{T_e}(y, z)$ for all $z \in L$. Consider the following cases:
 - 1. $x \in [0, e)$.
 - 1.1. If $y \in [0, e)$, then

$$U_{T_e}(x,z) = \begin{cases} T_e(x,z), & z \in [0,e), \\ x, & z = e, \\ x, & z \in (e,1], \\ 0, & z \in I_e, \end{cases}$$
 (5.3)

and

$$U_{T_e}(y, z) = \begin{cases} T_e(y, z), & z \in [0, e), \\ y, & z = e, \\ y, & z \in (e, 1], \\ 0, & z \in I_e, \end{cases}$$

implying that $U_{T_e}(x,z) \leq U_{T_e}(y,z)$, because T_e is increasing.

1.2. If $y \in (e, 1]$, then,

$$U_{T_e}(y, z) = \begin{cases} z, & z \in [0, e), \\ y, & z = e, \\ y \lor z, & z \in (e, 1], \\ z, & z \in I_e. \end{cases}$$

This, together with (5.3) and $T_e(x,z) \leq z$, implies that $U_{T_e}(x,z) \leq U_{T_e}(y,z)$.

1.3. If $y \in I_e$, then $x \in \mathscr{P}$. Applying (5.3) and (5.1) follows that

$$U_{T_e}(x,z) = \begin{cases} 0, & z \in [0,e), \\ x, & z = e, \\ x, & z \in (e,1], \\ 0, & z \in I_e, \end{cases}$$

and

$$U_{T_e}(y,z) = \begin{cases} 0, & z \in [0,e), \\ y, & z = e, \\ y, & z \in (e,1], \\ y \wedge z, & z \in I_e, \end{cases}$$
 (5.4)

implying that $U_{T_e}(x, z) \leq U_{T_e}(y, z)$.

- 1.4. If y = e, then $U_{T_e}(y, z) = z$. This, together with (5.3), implies that $U_{T_e}(x, z) \leq U_{T_e}(y, z)$.
- 2. $x \in (e, 1]$. From $x \leq y$, it follows that $y \in (e, 1]$. Then,

$$U_{T_e}(x,z) = \begin{cases} z, & z \in [0,e), \\ x, & z = e, \\ x \lor z, & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$
 (5.5)

and

$$U_{T_e}(y,z) = \begin{cases} z, & z \in [0,e), \\ y, & z = e, \\ y \lor z, & z \in (e,1], \\ z, & z \in I_e, \end{cases}$$
 (5.6)

implying that $U_{T_e}(x,z) \leq U_{T_e}(y,z)$.

3. x = e. This implies that $y \in [e, 1]$. Clearly, $U_{T_e}(x, z) = z = U_{T_e}(y, z)$ if y = e. If $y \in (e, 1]$, applying (5.6) and $U_{T_e}(x, z) = z$, it follows that $U_{T_e}(x, z) \leq U_{T_e}(y, z)$.

4. $x \in I_e$. From $x \leq y$, it follows that $y \in (e, 1] \cup I_e$. Applying (5.1) yields that

$$U_{T_e}(x,z) = \begin{cases} 0, & z \in [0,e), \\ x, & z = e, \\ x, & z \in (e,1], \\ x \wedge z, & z \in I_e. \end{cases}$$
 (5.7)

- 4.1. If $y \in (e, 1]$, from (5.6) and (5.7), it follows that $U_{T_e}(x, z) \leq U_{T_e}(y, z)$.
- 4.2. If $y \in I_e$, from (5.4) and (5.7), it is clear that $U_{T_e}(x,z) \leq U_{T_e}(y,z)$.
- i.2) Associativity. For any $x, y, z \in L$, it holds that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$. If one of the elements x, y and z is equal to e, then the equality is always satisfied. Otherwise, consider the following cases:
 - 1. $x \in [0, e)$.
 - 1.1. $y \in [0, e)$.
 - 1.1.1. If $z \in [0, e)$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, T_e(y, z)) = T_e(x, T_e(y, z)) = T_e(T_e(x, y), z) = T_e(U_{T_e}(x, y), z) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.1.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y) = T_e(x, y)$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(T_e(x, y), z) = T_e(x, y)$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.1.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, 0) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(T_e(x, y), z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.2. $y \in (e, 1]$.
 - 1.2.1. If $z \in [0, e)$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, z) = T_e(x, z)$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x, z) = T_e(x, z)$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.2.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y \vee z) = x$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x, z) = x$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.2.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, z) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x, z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.3. $y \in I_e$.
 - 1.3.1. If $z \in [0, e)$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, 0) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(0, z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.3.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(0, z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
 - 1.3.3. If $z \in I_e$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y \wedge z)$$

$$= \begin{cases} 0, & y \wedge z \in I_e, \\ 0, & y \wedge z = 0, \end{cases}$$

$$= 0,$$

and
$$U_{T_e}(U_{T_e}(x,y),z) = U_{T_e}(0,z) = 0$$
, implying that $U_{T_e}(x,U_{T_e}(y,z)) = U_{T_e}(U_{T_e}(x,y),z)$.

- 2. $x \in (e, 1]$.
 - 2.1. $y \in [0, e)$.
 - 2.1.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x) \quad \text{(commutativity of } U_{T_e})$$

$$= U_{T_e}(z, U_{T_e}(y, x)) \quad \text{(by 1.1.2)}$$

$$= U_{T_e}(U_{T_e}(x, y), z) \quad \text{(commutativity of } U_{T_e}).$$

- 2.1.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y) = y$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(y, z) = y$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 2.1.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, 0) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(y, z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 2.2. $y \in (e, 1]$.
 - 2.2.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x)$$
 (commutativity of U_{T_e})
= $U_{T_e}(z, U_{T_e}(y, x))$ (by 1.2.2)
= $U_{T_e}(U_{T_e}(x, y), z)$ (commutativity of U_{T_e}).

- 2.2.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y \lor z) = x \lor y \lor z$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x \lor y, z) = x \lor y \lor z$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 2.2.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, z) = z$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x \vee y, z) = z$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 2.3. $y \in I_e$.
 - 2.3.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x) \quad \text{(commutativity of } U_{T_e})$$

$$= U_{T_e}(z, U_{T_e}(y, x)) \quad \text{(by 1.3.2)}$$

$$= U_{T_e}(U_{T_e}(x, y), z) \quad \text{(commutativity of } U_{T_e}).$$

- 2.3.2. If $z \in (e, 1]$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y) = y$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(y, z) = y$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 2.3.3. If $z \in I_e$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y \wedge z)$$

$$= \begin{cases} y \wedge z, & y \wedge z \in I_e, \\ 0, & y \wedge z = 0, \end{cases}$$

$$= y \wedge z,$$

and
$$U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(y, z) = y \wedge z$$
, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.

- 3. $x \in I_e$.
 - 3.1. $y \in [0, e)$.
 - 3.1.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x) \quad \text{(commutativity of } U_{T_e})$$

$$= U_{T_e}(z, U_{T_e}(y, x)) \quad \text{(by 1.1.3)}$$

$$= U_{T_e}(U_{T_e}(x, y), z) \quad \text{(commutativity of } U_{T_e}).$$

3.1.2. If $z \in (e, 1]$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x) \quad \text{(commutativity of } U_{T_e})$$

$$= U_{T_e}(z, U_{T_e}(y, x)) \quad \text{(by 2.1.3)}$$

$$= U_{T_e}(U_{T_e}(x, y), z) \quad \text{(commutativity of } U_{T_e}).$$

- 3.1.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, 0) = 0$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(0, z) = 0$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 3.2. $y \in (e, 1]$.
 - 3.2.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x) \quad \text{(commutativity of } U_{T_e})$$

$$= U_{T_e}(z, U_{T_e}(y, x)) \quad \text{(by 1.2.3)}$$

$$= U_{T_e}(U_{T_e}(x, y), z) \quad \text{(commutativity of } U_{T_e}).$$

3.2.2. If $z \in (e, 1]$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x)$$
 (commutativity of U_{T_e})
= $U_{T_e}(z, U_{T_e}(y, x))$ (by 2.2.3)
= $U_{T_e}(U_{T_e}(x, y), z)$ (commutativity of U_{T_e}).

- 3.2.3. If $z \in I_e$, then $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, z) = x \wedge z$ and $U_{T_e}(U_{T_e}(x, y), z) = U_{T_e}(x, z) = x \wedge z$, implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.
- 3.3. $y \in I_e$.
 - 3.3.1. If $z \in [0, e)$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(z, y), x)$$
 (commutativity of U_{T_e})
= $U_{T_e}(z, U_{T_e}(y, x))$ (by 1.3.3)
= $U_{T_e}(U_{T_e}(x, y), z)$ (commutativity of U_{T_e}).

3.3.2. If $z \in (e, 1]$, then

$$\begin{split} U_{T_e}(x,U_{T_e}(y,z)) &= U_{T_e}(U_{T_e}(z,y),x) \quad \text{(commutativity of } U_{T_e}) \\ &= U_{T_e}(z,U_{T_e}(y,x)) \quad \text{(by 2.3.3)} \\ &= U_{T_e}(U_{T_e}(x,y),z) \quad \text{(commutativity of } U_{T_e}). \\ &\qquad \qquad 27 \end{split}$$

3.3.3. If $z \in I_e$, then

$$U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(x, y \lor z)$$

$$= \begin{cases} x \land y \land z, & y \land z \in I_e, \\ 0, & y \land z = 0, \end{cases}$$

$$= x \land y \land z,$$

and

$$U_{T_e}(U_{T_e}(x,y),z) = U_{T_e}(x \wedge y, z)$$

$$= \begin{cases} x \wedge y \wedge z, & x \wedge y \in I_e, \\ 0, & x \wedge y = 0, \end{cases}$$

$$= x \wedge y \wedge z,$$

implying that $U_{T_e}(x, U_{T_e}(y, z)) = U_{T_e}(U_{T_e}(x, y), z)$.

Theorem 12. Let $(X, \leq, 0, 1)$ be a bounded lattice, $e \in L \setminus \{0, 1\}$, and S_e be a t-conorm on [e, 1]. Then, the following statements are equivalent:

- (i) U_{S_e} defined by (5.2) is a uninorm on L with the neutral element e;
- (ii) $\mathscr{P} := \{x \in (0, e) : \exists y \in I_e, \ x \leq y\} = \emptyset;$
- (iii) $I_e || (0, e]$.

Proof. Applying Theorem 10, it is clear that $(iii) \iff (ii) \Longrightarrow (i)$.

(i) \Longrightarrow (iii). Suppose that I_e and (0, e] are comparable, i.e., there exist $x_0 \in (0, e]$ and $y_0 \in I_e$ such that $x_0 \leq y_0$. Clearly, $x_0 \neq e$. Since U_{S_e} is increasing, applying (5.2) yields that

$$0 < x_0 = U_{S_e}(x_0, x_0) \le U_{S_e}(x_0, y_0) = 0,$$

which is a contradiction.

Remark 1. (1) Theorems 1 and 2 are respectively direct corollaries of Theorems 8 and 5.

(2) Theorem 9 is a direct corollary of Theorem 11, when $\mathscr{P} = \emptyset$.

Example 1. Consider the complete lattice L with the Hasse diagram shown in Fig. 4 and take $T_e: \{0, a, e\}^2 \longrightarrow \{0, a, e\}$ by $T_e(x, y) = x \wedge y$ for $(x, y) \in \{0, a, e\}^2$. Clearly, $I_e = \{b, c\}$. The binary operation U_T defined by (1.1) is given in Table 1. Applying Theorem 8 shows that U_T is a uninorm on L, since $b \vee c = 1$. However, applying Theorem 1 can not conclude that U_T is a uninorm on L.




Figure 4: The Hasse diagram of the lattice L in Example 1

U_T	0	a	b	c	e	1
0	0	0	b	c	0	1
a	0	a	b	c	a	1
b	b	b	b	1	b	1
c	c	c	1	c	c	1
e	0	a	b	c	e	1
1	1	1	1	1	1	1

Table 1: The binary operation U_T in Example 1

References

- [1] E. Aşıcı and R. Mesiar. On the construction of uninorms on bounded lattices. Fuzzy Sets Syst., 2020. https://doi.org/10.1016/j.fss.2020.02.007.
- [2] G. Birkhoff. *Lattice Theory*. American Mathematical Society Colloquium Publications, Rhode Island, 1973.
- [3] S. Bodjanova and M. Kalina. Construction of uninorms on bounded lattices. In 2014 IEEE 12th International Symposium on Intelligent Systems and Informatics (SISY), pages 61–66. IEEE, 2014.
- [4] G. D. Çaylı. On the structure of uninorms on bounded lattices. Fuzzy Sets Syst., 357:2–26, 2019.
- [5] G. D. Çaylı and F. Karaçal. Construction of uninorms on bounded lattices. Kybernetika, 53(3):394–417, 2017.
- [6] G. D. Çaylı, F. Karaçal, and R. Mesiar. On a new class of uninorms on bounded lattices. *Information Sciences*, 367:221–231, 2016.
- [7] Y. Dan, B. Q. Hu, and J. Qiao. New constructions of uninorms on bounded lattices. *Int. J. Approx. Reason.*, 110:185–209, 2019.
- [8] F. Karacal, Ü. Ertuğrul, and R. Mesiar. Characterization of uninorms on bounded lattices. Fuzzy Set. Syst., 308:54–71, 2017.
- [9] F. Karaçal and R. Mesiar. Uninorms on bounded lattices. Fuzzy Sets Syst., 261:33–43, 2015.
- [10] K. Menger. Statistical metrics. Proc. Nat. Acad. Sci. USA, 37:535–537, 1942.
- [11] Y. Ouyang and H.-P. Zhang. Constructing uninorms via closure operations on a bounded lattice. Fuzzy Sets Syst., 2020. https://doi.org/10.1016/j.fss.2019.05.006.
- [12] B. Schweizer and A. Sklar. Associative functions and statistical triangle inequalities. *Publ. Math.*, 8:169–186, 1961.
- [13] A. Xie and S. Li. On constructing the largest and smallest uninorms on bounded lattices. *Fuzzy Sets Syst.*, 2020. https://doi.org/10.1016/j.fss.2019.04.020.
- [14] R. R. Yager and A. Rybalov. Uninorm aggregation operators. Fuzzy Sets Syst., 80(1):111–120, 1996.