

A p -ADIC MONODROMY THEOREM FOR DE RHAM LOCAL SYSTEMS

KOJI SHIMIZU

ABSTRACT. We study horizontal semistable and horizontal de Rham representations of the absolute Galois group of a certain smooth affinoid over a p -adic field. In particular, we prove that a horizontal de Rham representation becomes horizontal semistable after a finite extension of the base field. As an application, we show that every de Rham local system on a smooth rigid analytic variety becomes horizontal semistable étale locally around every classical point. We also discuss potentially crystalline loci of de Rham local systems and cohomologically potentially good reduction loci of smooth proper morphisms.

1. INTRODUCTION

Let p be a prime and let K be a complete discrete valuation field of characteristic zero with perfect residue field of characteristic p .

Fontaine introduced the notions of crystalline, semistable, and de Rham representations of the absolute Galois group of K to study p -adic étale cohomology of algebraic varieties over K . It is now known that Galois representations arising from geometry in this way are de Rham. He also conjectured that every de Rham representation is potentially semistable. This reflects the semistable reduction conjecture stating that every smooth proper algebraic variety over K acquires semistable reduction after a finite field extension. Fontaine's conjecture was proved by Berger [Ber02] and is often referred to as p -adic monodromy theorem.

This article discusses a similar question for families of p -adic Galois representations parametrized by a variety. More precisely, let X be a smooth rigid analytic variety over K (regarded as an adic space) and let \mathbb{L} be an étale \mathbb{Z}_p -local system on X . For every classical point $x \in X$ (i.e., a point whose residue class field $k(x)$ is finite over K) and a geometric point \bar{x} above x , the stalk $\mathbb{L}_{\bar{x}}$ is a Galois representation of $k(x)$. In this way, we regard \mathbb{L} as a family of Galois representations. In particular, we can ask whether \mathbb{L} is de Rham, semistable, etc., at every classical point. Note that building on earlier work [Fal88, Hyo89, Fal02, Bri08, AI13], Scholze [Sch13] defines the notion of de Rham local systems, and Liu and Zhu [LZ17] prove that \mathbb{L} is a de Rham local system if and only if \mathbb{L} is de Rham at every classical point of X (or even at a single classical point on each connected component). In this context, we prove the following p -adic monodromy theorem:

Theorem 1.1 (cf. Theorem 9.2). *Let \mathbb{L} be a de Rham \mathbb{Z}_p -local system on a smooth rigid analytic variety X over K . For every K -rational point $x \in X$, there exist an analytic open neighborhood $U \subset X$ of x and a finite extension L of K such that $\mathbb{L}|_{U \otimes_K L}$ is semistable at every classical point. Moreover, if $\mathbb{L}_{\bar{x}}$ is potentially*

Date: March 25, 2020.

crystalline as a Galois representation of $k(x)$, then we can choose U and L so that $\mathbb{L}|_{U \otimes_K L}$ is crystalline at every classical point.

We obtain a similar result for every classical point. Note that X can have a proper open subset containing all the K -rational points or even all the classical points in general. Hence the union of open sets U in the theorem may not be the entire X .

Remark 1.2. Currently, there seems no good way to define crystalline or semistable local systems on X without fixing a good formal model of X . This is the reason why our p -adic monodromy theorem concerns pointwise semistability. See [AI13, AI12, Fal89, Fal90, Fal02, TT19, Tsu] for definitions of crystalline or semistable local systems when starting with a nice integral or formal model.

Remark 1.3. In [Kis99], Kisin studies torsion and ℓ -adic étale local systems on X for a prime $\ell \neq p$, and shows that around each K -rational point, such a local system is locally constant in the analytic topology of X . In contrast, p -adic étale local systems on X are not necessarily locally constant in the analytic topology. Our result can be regarded as a p -adic analogue of Kisin’s result.

Remark 1.4. Note that Theorem 1.1 is of local nature. We may also ask a global version of p -adic monodromy conjecture (cf. [LZ17, Remark 1.4]): Let X be a smooth rigid analytic variety over K , and let \mathbb{L} be a de Rham \mathbb{Z}_p -local system on X . Does there exist a rigid analytic variety X' finite étale over X such that $\mathbb{L}|_{X'}$ is semistable at every classical point? We know an affirmative answer in the following two cases:

- (i) the case where \mathbb{L} is the p -adic Tate module of an abelian variety over X (use Raynaud’s criterion [SGA7, Exposé IX, Proposition 4.7]).
- (ii) the case where \mathbb{L} has a single Hodge–Tate weight [Shi18, Theorem 1.6].

On the other hand, Lawrence and Li [LL20] show that there exist an algebraic variety X over K and a de Rham \mathbb{Z}_p -local system \mathbb{L} on X such that for every finite extension L of K , $\mathbb{L}|_{X \otimes_K L}$ is not semistable at some closed point.

Remark 1.5. It is an interesting attempt to formulate a semistable reduction conjecture in families over a field of mixed characteristic; Theorem 1.1 will be p -adic Hodge-theoretic evidence for such a conjecture. In the case of fields of characteristic zero, Abramovich and Karu [AK00] make a semistable reduction conjecture in families, and Adiprasito, Liu, and Temkin [ALT18] prove the conjecture of Abramovich and Karu in a larger generality. See these articles for the details.

Let us turn to applications of Theorem 1.1. Note that Liu and Zhu’s result implies that if an étale \mathbb{Q}_p -local system \mathbb{L} on X is de Rham at a classical point on each connected component, then \mathbb{L} is de Rham at every classical point. Such a strong result does not hold for potentially crystalline representations. However, we prove that if \mathbb{L} is potentially crystalline at a classical point x , then it is so at every classical point in an analytic open neighborhood of x .

Theorem 1.6 (Theorem 10.2). *If \mathbb{L} is a de Rham \mathbb{Q}_p -local system on a smooth rigid analytic variety X over K , then there exists an analytic open subset $U \subset X$ such that the following holds for every classical point $x \in X$:*

- (i) *if $x \in U$, then \mathbb{L} is potentially crystalline at x .*

(ii) if $x \notin U$, then \mathbb{L} is not potentially crystalline at x .

We call such U a *potentially crystalline locus* of \mathbb{L} .

Note that our definition takes only classical points into consideration and thus a potentially crystalline locus of \mathbb{L} is not unique. On the other hand, the union of potentially crystalline loci is also a potentially crystalline locus. Hence we obtain the largest potentially crystalline locus of \mathbb{L} .

The second application concerns the special case where \mathbb{L} comes from geometry. For example, suppose that we have a family of elliptic curves $f: E \rightarrow X$. For a point $x \in X(\overline{K})$, the elliptic curve E_x has good reduction if and only if the j -invariant of E_x is a p -adic integer. The set of such points may well be called the (potentially) good reduction locus of E . In general, if we have a smooth proper family of algebraic varieties, the étale cohomology of the fibers yields both ℓ -adic and p -adic étale local systems on the base. By combining known cases of ℓ -adic/ p -adic monodromy-weight conjecture and the ℓ -independence with our result or Kisin's result, we obtain the following theorem:

Theorem 1.7 (Theorem 10.15). *Assume that the residue field of K is finite. Let $f: Y \rightarrow X$ be a smooth proper morphism between smooth algebraic varieties over K such that the relative dimension of f is at most two. For each $m \in \mathbb{N}$, there exists an analytic open subset U of the adic space X^{ad} associated to X such that the following holds for every classical point x of X^{ad} :*

- (i) if $x \in U$, then $R^m f_* \mathbb{Q}_\ell$ is potentially unramified at x for $\ell \neq p$ and $R^m f_* \mathbb{Q}_p$ is potentially crystalline at x .
- (ii) if $x \notin U$, then $R^m f_* \mathbb{Q}_\ell$ is not potentially unramified at x for $\ell \neq p$ and $R^m f_* \mathbb{Q}_p$ is not potentially crystalline at x .

We call such U a *cohomologically potentially good reduction locus* of f of degree m .

As in the previous theorem, such a locus is not unique but we obtain the largest cohomologically potentially good reduction locus.

Remark 1.8. Imai and Mieda [IM20] introduce a notion of the potentially good reduction locus of a Shimura variety using automorphic local systems. They prove the existence of the potentially good reduction loci of Shimura varieties of preabelian type, and use these loci to study the étale cohomology of Shimura varieties.

We now explain ideas of the proof of Theorem 1.1. We start with an example that motivates our proof: Let $\mathfrak{f}: \mathfrak{Y} \rightarrow \mathfrak{X}$ be a smooth proper morphism of smooth p -adic formal schemes over $\text{Spf } \mathcal{O}_K$, and let $f: Y \rightarrow X$ denote its adic generic fiber. Then $\mathbb{L} := R^m f_* \mathbb{Q}_p$ is a de Rham \mathbb{Q}_p -local system. In fact, we even know that it is crystalline at every classical point (hence there is nothing to prove in this case). Let z be a closed point in the special fiber of \mathfrak{X} and let $U \subset X$ be the residue open disk of z . For K -rational points $x_1, x_2 \in U$, there is a canonical identification of φ -modules¹

$$D_{\text{cris}}(\mathbb{L}_{\overline{x_1}}) \cong H_{\text{cris}}^m(\mathfrak{Y}_z/W(k))[p^{-1}] \cong D_{\text{cris}}(\mathbb{L}_{\overline{x_2}}).$$

One key property of U is that the Gauss–Manin connection on the vector bundle $H_{\text{dR}}^m(Y/X)$ on X has a full set of solutions when restricted to U (cf. [Kat73] and [BO83, Remark 2.9]). Heuristically, this means that we can compare φ -modules

¹They need not be isomorphic as *filtered* φ -modules, and thus $\mathbb{L}_{\overline{x_1}}$ and $\mathbb{L}_{\overline{x_2}}$ may not be isomorphic as Galois representations of K .

associated to stalks of \mathbb{L} at different points on U by parallel transport. Our proof articulates this idea despite the absence of integral models of X nor motives for \mathbb{L} .

Let us return to the general setting of Theorem 1.1. Liu and Zhu [LZ17] introduce a natural functor D_{dR} from the category of étale \mathbb{Q}_p -local systems on X to the category of filtered vector bundles with integrable connections on X . When $X = \text{Spa}(K, \mathcal{O}_K)$, this agrees with Fontaine's D_{dR} -functor for Galois representations of K . The functor commutes with arbitrary pullbacks, and an étale \mathbb{Q}_p -local system \mathbb{L} on X is de Rham if and only if $\text{rank } D_{\text{dR}}(\mathbb{L}) = \text{rank } \mathbb{L}$. Note that in the above geometric situation, $D_{\text{dR}}(R^m f_* \mathbb{Q}_p)$ coincides with $H_{\text{dR}}^m(Y/X)$ equipped with the Gauss–Manin connection and the Hodge filtration.

Suppose that \mathbb{L} is a de Rham \mathbb{Z}_p -local system and fix a K -rational point x of X . Since X is smooth, the theory of p -adic differential equations over a poly-disk (Theorem 9.7) tells that there is an analytic open neighborhood $U \subset X$ of x such that $D_{\text{dR}}(\mathbb{L}|_U)$ has a full set of solutions. By shrinking U if necessary, we may further assume that U is isomorphic to a rigid torus $\text{Spa}(A, A^\circ)$, where $A^\circ := \mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle$, the p -adic completion of the Laurent polynomial ring $\mathcal{O}_K[T_1^{\pm 1}, \dots, T_n^{\pm 1}]$, and $A := A^\circ[p^{-1}]$.

Let \mathcal{G}_A denote the absolute Galois group of A . Following work of Colmez [Col08] and Brinon [Bri08], we define horizontal crystalline, semistable, and de Rham period rings $B_{\text{max}}^\nabla(A^\circ), B_{\text{st}}^\nabla(A^\circ), B_{\text{dR}}^\nabla(A^\circ)$. Using these period rings, we define the notions of horizontal crystalline, semistable, and de Rham representations of \mathcal{G}_A (Definition 4.16). In our setting, $\mathbb{L}|_U$ corresponds to a p -adic representation V of \mathcal{G}_A . Since $\mathbb{L}|_U$ is de Rham and $D_{\text{dR}}(\mathbb{L}|_U)$ has a full set of solutions, V is horizontal de Rham. Now the first part of Theorem 1.1 follows from a p -adic monodromy theorem for horizontal de Rham representations:

Theorem 1.9 (Theorem 7.4). *If a p -adic representation V of \mathcal{G}_A is horizontal de Rham, then there exists a finite extension L of K such that $V|_{\mathcal{G}_{A_L}}$ is horizontal semistable, where $A_L := A \otimes_K L$.*

The second part of Theorem 1.1 follows from the study of the monodromy operator (cf. Theorem 7.5).

To explain the proof of Theorem 1.9, we need some more notation. Let \mathcal{O}_K denote the p -adic completion of the localization $A_{(\pi)}^\circ$, where π is a uniformizer of K . Then $\mathcal{K} := \mathcal{O}_K[p^{-1}]$ is a complete discrete valuation field with imperfect residue field. We similarly have the notions of horizontal semistable and de Rham representations of the absolute Galois group of \mathcal{K} . In [Ohk13], Ohkubo shows that every horizontal de Rham representation becomes horizontal semistable after restricting to the absolute Galois group of the composite LK for some finite extension L of K . Hence Theorem 1.9 follows from Ohkubo's result and the following purity theorem for horizontal semistable representations of \mathcal{G}_A :

Theorem 1.10 (Theorem 6.1). *If a horizontal de Rham representation of \mathcal{G}_A is horizontal semistable when restricted to the absolute Galois group of \mathcal{K} , then it is also horizontal semistable as a representation of \mathcal{G}_A .*

This is the key technical result of our article and the proof requires detailed analysis of period rings (e.g. Lemma 2.40). Our purity theorem and its proof are inspired by similar results in different contexts: a p -adic monodromy theorem for de Rham representations of the absolute Galois group of K with values in a reduced

affinoid algebra by Berger and Colmez [BC08], and a purity theorem for crystalline local systems in the case of good reduction by Tsuji [Tsu, Theorem 5.4.8].

At this point, we emphasize that the main goal of our work is to develop p -adic Hodge theory of horizontal crystalline, semistable, and de Rham representations of \mathcal{G}_A . These classes of representations are much more restrictive than those of crystalline and de Rham representations studied by Brinon [Bri08]. However, it is still worth studying them. In fact, as we have explained, de Rham local systems come from horizontal de Rham representations (étale) locally around each classical point. For another instance, Faltings [Fal10] studies the admissible loci of certain p -adic period domains using horizontal crystalline representations (see also [Har13, Ked10b, Far18] for relevant work).

This article establishes two results in p -adic Hodge theory of \mathcal{G}_A . First we prove the p -adic monodromy theorem for horizontal de Rham representations as we have discussed. Second we define Fontaine's functors D_{pst} and V_{pst} in our setting (Section 5). We prove that these functors induce an equivalence of categories between the category of horizontal de Rham representations of \mathcal{G}_A and that of admissible discrete filtered $(\varphi, N, \text{Gal}(\overline{K}/K), A)$ -modules (Theorem 5.10). What is missing in our work is to characterize the admissibility condition in the latter category in terms of more intrinsic properties like semistability (weak admissibility). In the case of p -adic Hodge theory for K , this is first established by Colmez and Fontaine [CF00]. Unfortunately, results on p -adic period domains already suggest that the admissibility in our setting cannot be characterized by (classical) pointwise weak admissibility (cf. [Har13, Example 6.7]).

Finally, we explain the organization of this paper. The first part (Sections 2-7) develops p -adic Hodge theory for rings (i.e., representations of \mathcal{G}_A) and the second part (Sections 8-10) studies p -adic Hodge theory for rigid analytic varieties.

In the first part, we need to define horizontal period rings such as $B_{\text{dR}}^{\nabla}(A^{\circ})$. We define them in two steps. First we define horizontal period rings such as $\mathbb{B}_{\text{dR}}(\Lambda, \Lambda^+)$ for every perfectoid Banach pair (Λ, Λ^+) over the completed algebraic closure of K . For a base ring R (e.g. A°), the p -adic completion \widehat{R} of the integral closure of R in the algebraic closure of $R[p^{-1}]$ gives such a perfectoid Banach pair, and we set $B_{\text{dR}}^{\nabla}(R) := \mathbb{B}_{\text{dR}}(\widehat{R}[p^{-1}], \widehat{R})$.

In Section 2, we define horizontal period rings for perfectoid Banach pairs, following [Col08] and [Bri08]. Most of the arguments in this section are standard in p -adic Hodge theory. One exception might be Lemma 2.40, which will be used crucially in the proof of the purity theorem in Section 6. Section 3 studies the class of base rings R over which Brinon [Bri08] develops p -adic Hodge theory. In Section 4, we define horizontal crystalline, semistable, and de Rham representations. Section 5 introduces filtered $(\varphi, N, \text{Gal}(L/K), A)$ -modules, and the functors D_{pst} and V_{pst} . This section is not used in later sections. After these preparations, we state and prove the purity theorem in Section 6. We prove Theorem 1.9 in Section 7.

The second part starts with Section 8, where we review étale local systems on smooth rigid analytic varieties and period sheaves on the pro-étale site. Our p -adic monodromy theorem is proved in Section 9. Finally, we discuss potentially crystalline loci and cohomologically potentially good reduction loci in Section 10.

Acknowledgments. The author thanks Hélène Esnault, Teruhisa Koshikawa and Alex Youcis for valuable discussions and comments about this work. The author

was supported by the National Science Foundation under Grant No. DMS-1638352 through membership at the Institute for Advanced Study.

Notation.

For a ring S of characteristic p , the Frobenius map $x \mapsto x^p$ is denoted by φ . We say that S is *perfect* if φ is an isomorphism. In this case, $W(S)$ denotes the ring of p -typical Witt vectors of S . By functoriality, $W(S)$ admits the Witt vector Frobenius $W(\varphi)$, which we simply denote by φ .

A rigid analytic variety over K refers to a quasi-separated adic space locally of finite type over $\mathrm{Spa}(K, \mathcal{O}_K)$ (cf. [Hub94, Proposition 4.5(iv)]). For a rigid analytic variety X over K , an (analytic) open subset of X refers to an open subset of X as an adic space. Definitions of étale \mathbb{Z}_p -local systems and étale \mathbb{Q}_p -local systems on X are explained in Subsection 8.1.

2. PERIOD RINGS

In this section, we introduce horizontal period rings $\mathbb{B}_{\mathrm{dR}}, \mathbb{B}_{\mathrm{max}}, \mathbb{B}_{\mathrm{st}}$, etc., and study their basic properties. We will use these period rings to define horizontal de Rham, crystalline, and semistable representations in Section 4. Colmez define these period rings for sympathetic algebras in [Col08]. For our later technical purpose, we define them in a more general setting. Most proofs in this section have essentially appeared in [Fon94a, Col08, Bri08, KL15]. Lemma 2.40 is a key technical result used in the proof of the purity theorem in Section 6.

2.1. Review of Banach algebras and perfectoid pairs. In this subsection, let L be a field that is complete with respect to a non-archimedean p -adic norm $|\cdot|$ satisfying $|p| = p^{-1}$.

Definition 2.1 (cf. [Col08, §2.2]).

- (i) Let Λ be an L -algebra. An L -algebra seminorm $|\cdot|_\Lambda$ on Λ is a function $|\cdot|_\Lambda : \Lambda \rightarrow \mathbb{R}_{\geq 0}$ such that
 - $|0|_\Lambda = 0$ and $|1|_\Lambda = 1$;
 - $|x + y|_\Lambda \leq \max\{|x|_\Lambda, |y|_\Lambda\}$ and $|xy|_\Lambda \leq |x|_\Lambda \cdot |y|_\Lambda$ for $x, y \in \Lambda$;
 - $|ax|_\Lambda = |a| \cdot |x|_\Lambda$ for $a \in L$ and $x \in \Lambda$.

We say that an L -algebra seminorm is an L -algebra norm if $|x|_\Lambda = 0$ holds only when $x = 0$. In this case, we call the pair $(\Lambda, |\cdot|_\Lambda)$ an L -normed algebra. If there is no confusion, we simply say that Λ is an L -normed algebra and denote its fixed L -algebra norm by $|\cdot|_\Lambda$.

- (ii) An L -Banach algebra is an L -normed algebra $(\Lambda, |\cdot|_\Lambda)$ such that Λ is complete with respect to the topology defined by $|\cdot|_\Lambda$. A morphism of L -Banach algebras will refer to a continuous L -algebra homomorphism.
- (iii) Let Λ be an L -Banach algebra. An element $x \in \Lambda$ is called *power-bounded* if the subset $S = \{x^n \mid n \in \mathbb{N}\} \subset \Lambda$ is bounded, i.e., for every open neighborhood U of 0, there exists an open neighborhood V of 0 such that $VS \subset U$. The set of power-bounded elements forms a subring of Λ , which we denote by Λ° .
- (iv) ([KL15, Definition 2.8.1]) An L -Banach algebra Λ is called *uniform* if the subring Λ° of Λ is bounded. Note that Λ is uniform if and only if the norm on Λ is equivalent to its spectral (semi)norm².

²If Λ is uniform, then Λ with the spectral norm is uniform in the sense of [Ber90, p. 16].

- (v) ([KL15, Definition 2.4.1]) An L -Banach pair³ is a pair (Λ, Λ^+) in which Λ is an L -Banach algebra and Λ^+ is a subring of Λ° which is open and integrally closed in Λ . Such a Λ^+ is also called a *ring of integral elements*. A morphism of L -Banach pairs $(\Lambda, \Lambda^+) \rightarrow (\Lambda', \Lambda'^+)$ is a morphism $f: \Lambda \rightarrow \Lambda'$ of L -Banach algebras such that $f(\Lambda^+) \subset \Lambda'^+$.

Remark 2.2. Note that in [KL15] a morphism of L -Banach algebras is required to be bounded.

Remark 2.3. If (Λ, Λ^+) is an L -Banach pair and if Λ is uniform, then Λ^+ is reduced, p -torsion free, and p -adically complete and separated.

Definition 2.4. Assume further that L is algebraically closed. An L -Banach pair (Λ, Λ^+) is called *perfectoid* if Λ is uniform and if the Frobenius morphism $x \mapsto x^p$ on $\Lambda^+/p\Lambda^+$ is surjective.

Remark 2.5. Under the assumption that L is algebraically closed, an L -Banach pair (Λ, Λ^+) is perfectoid in the above sense if and only if it is perfectoid in the sense of [KL15, Definition 3.6.1] by [KL15, Proposition 3.6.2 (a)]. In the case where $\Lambda^+ = \Lambda^\circ$, this definition is equivalent to the condition that Λ is a perfectoid L -algebra in the sense of [Sch12, Definition 5.1 (i)], in which case we will simply say that Λ is perfectoid (over L). By [KL15, Proposition 3.6.2 (d)], (Λ, Λ^+) is perfectoid if and only if Λ is perfectoid, i.e., (Λ, Λ°) is perfectoid.

2.2. Period rings for perfectoid pairs. Let k be a perfect field of characteristic p and set $K_0 := W(k)[p^{-1}]$. Let K be a totally ramified finite extension of K_0 . We fix a uniformizer π of K and an algebraic closure \bar{K} of K . Let C denote the p -adic completion of \bar{K} . Then C is algebraically closed by Krasner's lemma. We denote by \mathcal{O}_C the ring of integers of C .

Set-up 2.6. Let (Λ, Λ^+) be a perfectoid C -Banach pair such that Λ^+ is an \mathcal{O}_C -algebra. Note that $\Lambda = \Lambda^+[p^{-1}]$.

Remark 2.7. When we discuss a morphism $(\Lambda, \Lambda^+) \rightarrow (\Lambda', \Lambda'^+)$ of such objects, we consider a morphism of K -Banach pairs as opposed to that of C -Banach pairs. For example, we will consider $\sigma: (C, \mathcal{O}_C) \rightarrow (C, \mathcal{O}_C)$ for $\sigma \in \text{Gal}(\bar{K}/K)$.

We set

$$\Lambda^{+b} := \varprojlim_{\varphi} \Lambda^+/p\Lambda^+.$$

By [KL15, Remark 3.4.10], this is a ring of characteristic p (called the *tilt* of Λ^+) and there is a natural multiplicative monoid isomorphism

$$\varprojlim_{\lambda \mapsto \lambda^p} \Lambda^+ \xrightarrow{\cong} \Lambda^{+b}.$$

We denote by $\Lambda^{+b} \rightarrow \Lambda^+; x \mapsto x^\sharp$ the inverse of the above isomorphism followed by the first projection. Note that Λ^{+b} has a norm $|\cdot|$ defined by $|x| := |x^\sharp|_\Lambda$ and it is complete with respect to this norm ([KL15, Lemma 3.4.5]).

We set

$$\begin{aligned} \mathbb{A}_{\text{inf}}(\Lambda^+) &= \mathbb{A}_{\text{inf}}(\Lambda, \Lambda^+) := W(\Lambda^{+b}), \quad \text{and} \\ \mathbb{A}_{\text{inf}, K}(\Lambda^+) &= \mathbb{A}_{\text{inf}, K}(\Lambda, \Lambda^+) := \mathcal{O}_K \otimes_{W(k)} \mathbb{A}_{\text{inf}}(\Lambda^+). \end{aligned}$$

³It is called an adic L -Banach ring in [KL15].

By functoriality of p -typical Witt vectors, $\mathbb{A}_{\text{inf}}(\Lambda^+)$ carries the Frobenius φ lifting $x \mapsto x^p$. There exists a unique surjective $W(k)$ -algebra homomorphism

$$\theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)} : \mathbb{A}_{\text{inf}}(\Lambda^+) \rightarrow \Lambda^+$$

characterized by

$$\theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)} \left(\sum_{i=0}^{\infty} [x_i] p^i \right) = \sum_{i=0}^{\infty} x_i^{\sharp} p^i,$$

where $[x_i] \in \mathbb{A}_{\text{inf}}(\Lambda^+)$ is the Teichmüller lift of x_i (cf. [KL15, Lemma 3.2.2, Definition 3.4.3]). This map also extends to a surjective \mathcal{O}_K -algebra homomorphism

$$\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)} : \mathbb{A}_{\text{inf},K}(\Lambda^+) \rightarrow \Lambda^+.$$

Choose a compatible system $(p_m)_{m \in \mathbb{N}}$ of p -power roots of p in \overline{K} , i.e., $p_m \in \overline{K}$ with $p_0 = p$ and $p_{m+1}^p = p_m$. Set $p^{\flat} := (p_0 \bmod p, p_1 \bmod p, \dots) \in \mathcal{O}_C^{\flat} \subset \Lambda^{+\flat}$. Similarly, choose a compatible system $(\pi_m)_{m \in \mathbb{N}}$ of p -power roots of π in \overline{K} and set $\pi^{\flat} := (\pi_0 \bmod p, \pi_1 \bmod p, \dots) \in \mathcal{O}_C^{\flat} \subset \Lambda^{+\flat}$.

Lemma 2.8. *The ideal $\text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$ is principal. In fact, it is generated by $[p^{\flat}] - p$. Similarly, $\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$ is principal with a generator $[\pi^{\flat}] - \pi$.*

Proof. The first assertion is [KL15, Lemma 3.6.3]⁴. We prove the second. Note $[\pi^{\flat}] - \pi \in \text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$. Since $\mathbb{A}_{\text{inf},K}(\Lambda^+)$ is π -adically complete and Λ^+ is π -torsion free, it suffices to prove that the ideal $(\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}, \pi)$ is equal to $([\pi^{\flat}] - \pi, \pi)$. Take $x \in \text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$. Note that x is uniquely written as $x = \sum_{i=0}^{\infty} [x_i] \pi^i$ for some $x_i \in \Lambda^{+\flat}$. Since $0 = \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}(x) = \sum_{i=0}^{\infty} x_i^{\sharp} \pi^i$, we see that π divides x_0^{\sharp} in Λ^+ . Using the multiplicative monoid isomorphism $\varprojlim_{\lambda \mapsto \lambda^p} \Lambda^+ \xrightarrow{\cong} \Lambda^{+\flat}$ and the fact that Λ^+ is integrally closed, we see that π^{\flat} divides x_0 in $\Lambda^{+\flat}$ and thus $[\pi^{\flat}]$ divides $[x_0]$ in $\mathbb{A}_{\text{inf},K}(\Lambda^+)$. Write $[x_0] = y[\pi^{\flat}]$ for some $y \in \mathbb{A}_{\text{inf},K}(\Lambda^+)$. Then we have

$$x = y([\pi^{\flat}] - \pi) + \pi \left(y + \sum_{i=1}^{\infty} [x_i] \pi^{i-1} \right).$$

This implies $(\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}, \pi) = ([\pi^{\flat}] - \pi, \pi)$. \square

Let us define the horizontal de Rham period rings:

Definition 2.9. We set

$$\begin{aligned} \mathbb{B}_{\text{dR}}^+(\Lambda^+) &= \mathbb{B}_{\text{dR}}^+(\Lambda, \Lambda^+) := \varprojlim_m \mathbb{A}_{\text{inf}}(\Lambda^+) [p^{-1}] / (\text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)} [p^{-1}])^m, \quad \text{and} \\ \mathbb{B}_{\text{dR},K}^+(\Lambda^+) &= \mathbb{B}_{\text{dR},K}^+(\Lambda, \Lambda^+) := \varprojlim_m \mathbb{A}_{\text{inf},K}(\Lambda^+) [p^{-1}] / (\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)} [p^{-1}])^m. \end{aligned}$$

We will prove that these two rings are naturally isomorphic in Lemma 2.12, and write simply $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ for them afterward.

Lemma 2.10. *The map $\theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$ extends to*

$$\theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)} : \mathbb{B}_{\text{dR}}^+(\Lambda^+) \rightarrow \Lambda = \Lambda^+ [p^{-1}],$$

⁴It is also a special case of the second assertion by considering the case where $K = K_0$.

and $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is complete with respect to the ideal $\text{Ker } \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}$. Similarly, the map $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$ extends to

$$\theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)} : \mathbb{B}_{\text{dR},K}^+(\Lambda^+) \rightarrow \Lambda,$$

and $\mathbb{B}_{\text{dR},K}^+(\Lambda^+)$ is complete with respect to the ideal $\text{Ker } \theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)}$.

Proof. It suffices to prove the second assertion as the first is a special case of the second where $K = K_0$. Obviously, $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$ extends to $\theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)} : \mathbb{B}_{\text{dR},K}^+(\Lambda^+) \rightarrow \Lambda$. Since Λ is p -torsion free, the ideal $\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}]$ of $\mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]$ is also generated by $[\pi^b] - \pi$, and thus finitely generated. It follows that $\mathbb{B}_{\text{dR},K}^+(\Lambda^+)$ is complete with respect to $\text{Ker } \theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)}$. \square

Note that the natural homomorphisms

$$\mathbb{A}_{\text{inf}}(\Lambda^+)[p^{-1}]/(\text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}[p^{-1}])^m \rightarrow \mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]/(\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}])^m$$

give rise to a ring homomorphism

$$\mathbb{B}_{\text{dR}}^+(\Lambda^+) \rightarrow \mathbb{B}_{\text{dR},K}^+(\Lambda^+).$$

Choose a compatible system $(\varepsilon_m)_{m \in \mathbb{N}}$ of nontrivial p -power roots of unity in \overline{K} , i.e., $\varepsilon_m \in \overline{K}$ with $\varepsilon_0 = 1$, $\varepsilon_1 \neq 1$, and $\varepsilon_{m+1}^p = \varepsilon_m$. Set

$$\varepsilon := (\varepsilon_0 \bmod p, \varepsilon_1 \bmod p, \dots) \in \mathcal{O}_C^b \subset \Lambda^{+b}.$$

Note that $[\varepsilon] - 1 \in \mathbb{A}_{\text{inf}}(\Lambda^+)$ satisfies $\theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}([\varepsilon] - 1) = 0$. In particular, we define

$$t := \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} ([\varepsilon] - 1)^m$$

as an element of $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$.

Lemma 2.11. *The ideal $\text{Ker } \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}$ is generated by t . Similarly, $\text{Ker } \theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)}$ is generated by t .*

Proof. Since the first assertion is a special case of the second, we only show the second. We know that $\text{Ker } \theta_{\mathbb{B}_{\text{dR},K}^+(\Lambda^+)}$ is generated by $([\pi^b] - \pi)$. Hence it suffices to prove that t divides $[\pi^b] - \pi$ in $\mathbb{B}_{\text{dR},K}^+(\Lambda^+)$. By functoriality of our construction, it is enough to prove this for $(\Lambda, \Lambda^+) = (C, \mathcal{O}_C)$. This is [Fon94a, 1.5.4]. \square

Lemma 2.12. *The natural homomorphism*

$$\mathbb{B}_{\text{dR}}^+(\Lambda^+) \rightarrow \mathbb{B}_{\text{dR},K}^+(\Lambda^+)$$

is an isomorphism.

Proof. This follows from Lemmas 2.10 and 2.11. \square

As mentioned before, we identify these rings and denote them by $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ in what follows. In particular, $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is a K -algebra.

Lemma 2.13. *The natural map $\mathbb{A}_{\text{inf},K}(\Lambda^+) \rightarrow \mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is injective.*

Proof. We claim

$$\mathbb{A}_{\text{inf},K}(\Lambda^+) \cap (\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}])^m = (\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)})^m, \quad \text{and} \\ \bigcap_m (\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}])^m = 0.$$

Note that the assertion follows from the second claim. The first claim follows from the fact that $\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}]$ is generated by $[\pi^b] - \pi$ and Λ^+ is p -torsion free. By the first claim, the second claim is reduced to $\bigcap_m (\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)})^m = 0$, which follows from the fact that $\mathbb{A}_{\text{inf},K}(\Lambda^+)/(\pi) = \Lambda^{+b}$ has a norm $||$ satisfying $|\pi^b| = |\pi|_\Lambda > 0$. \square

Lemma 2.14. *The ring $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is t -torsion free.*

Proof. It is enough to show that $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is $([p^b] - p)$ -torsion free. Since $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is the completion of $\mathbb{A}_{\text{inf}}(\Lambda^+)[p^{-1}]$ with respect to the ideal $([p^b] - p)$, it remains to prove that the latter ring is $([p^b] - p)$ -torsion free. Since Λ^+ is p -torsion free, Λ^{+b} is p^b -torsion free. It follows that $\mathbb{A}_{\text{inf}}(\Lambda^+)$ is $([p^b] - p)$ -torsion free and so is $\mathbb{A}_{\text{inf}}(\Lambda^+)[p^{-1}]$. \square

Definition 2.15. Set

$$\mathbb{B}_{\text{dR}}(\Lambda^+) = \mathbb{B}_{\text{dR}}(\Lambda, \Lambda^+) := \mathbb{B}_{\text{dR}}^+(\Lambda^+)[t^{-1}].$$

We equip $\mathbb{B}_{\text{dR}}(\Lambda^+)$ with the decreasing filtration defined by

$$\text{Fil}^m \mathbb{B}_{\text{dR}}(\Lambda^+) := t^m \mathbb{B}_{\text{dR}}^+(\Lambda^+) \quad (m \in \mathbb{Z}).$$

This is separated and exhaustive.

Definition 2.16. Let $\mathbb{A}_{\text{inf}}(\Lambda^+)[\frac{\text{Ker } \theta}{p}]$ be the $\mathbb{A}_{\text{inf}}(\Lambda^+)$ -subalgebra of $\mathbb{A}_{\text{inf}}(\Lambda^+)[p^{-1}]$ generated by $p^{-1} \text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$. We define $\mathbb{A}_{\text{max}}(\Lambda^+) = \mathbb{A}_{\text{max}}(\Lambda, \Lambda^+)$ to be the p -adic completion of $\mathbb{A}_{\text{inf}}(\Lambda^+)[\frac{\text{Ker } \theta}{p}]$.

Similarly, let $\mathbb{A}_{\text{inf},K}(\Lambda^+)[\frac{\text{Ker } \theta}{\pi}]$ be the $\mathbb{A}_{\text{inf},K}(\Lambda^+)$ -subalgebra of $\mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]$ generated by $\pi^{-1} \text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$. We define $\mathbb{A}_{\text{max},K}(\Lambda^+) = \mathbb{A}_{\text{max},K}(\Lambda, \Lambda^+)$ to be the p -adic completion of $\mathbb{A}_{\text{inf},K}(\Lambda^+)[\frac{\text{Ker } \theta}{\pi}]$.

Lemma 2.17. *The Frobenius φ on $\mathbb{A}_{\text{inf}}(\Lambda^+)$ extends to an endomorphism on $\mathbb{A}_{\text{inf}}(\Lambda^+)[\frac{\text{Ker } \theta}{p}]$ and thus on $\mathbb{A}_{\text{max}}(\Lambda^+)$.*

We call this endomorphism on $\mathbb{A}_{\text{max}}(\Lambda^+)$ the Frobenius and denote it by φ .

Proof. Recall that $\text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$ is generated by $[p^b] - p$. Hence the assertion follows from

$$\varphi\left(\frac{[p^b] - p}{p}\right) = \frac{[p^b]^p - p}{p} = p^{p-1} \left(\frac{[p^b] - p}{p} + 1\right)^p - 1 \in \mathbb{A}_{\text{inf}}(\Lambda^+) \left[\frac{\text{Ker } \theta}{p}\right].$$

\square

Lemma 2.18. *The rings $\mathbb{A}_{\text{max}}(\Lambda^+)$ and $\mathbb{A}_{\text{max},K}(\Lambda^+)$ are p -torsion free.*

Proof. It is enough to show that $\mathbb{A}_{\text{max},K}(\Lambda^+)$ is p -torsion free. Note that $\mathbb{A}_{\text{inf},K}(\Lambda^+)$ is a finite free module over $\mathbb{A}_{\text{inf}}(\Lambda^+)$ and thus p -torsion free. Hence $\mathbb{A}_{\text{inf},K}(\Lambda^+)[\frac{\text{Ker } \theta}{\pi}]$ is also p -torsion free. The assertion for $\mathbb{A}_{\text{max},K}(\Lambda^+)$ follows from this. \square

Definition 2.19. Set

$$\begin{aligned}\mathbb{B}_{\max}^+(\Lambda^+) &= \mathbb{B}_{\max}^+(\Lambda, \Lambda^+) := \mathbb{A}_{\max}(\Lambda^+)[p^{-1}], \quad \text{and} \\ \mathbb{B}_{\max, K}^+(\Lambda^+) &= \mathbb{B}_{\max, K}^+(\Lambda, \Lambda^+) := \mathbb{A}_{\max, K}(\Lambda^+)[p^{-1}].\end{aligned}$$

Lemma 2.20. *The element*

$$t = \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} ([\varepsilon] - 1)^m$$

converges in $\frac{1}{p^r} \mathbb{A}_{\max, K}(\Lambda^+)$ for a sufficiently large $r \in \mathbb{N}$. It also converges in $\mathbb{A}_{\max}(\Lambda^+)$.

Proof. Since the proofs are similar (and the second assertion is easier to prove), we show the first. Write

$$\frac{(-1)^{m-1}}{m} ([\varepsilon] - 1)^m = \frac{(-1)^{m-1} \pi^m}{m} \left(\frac{[\varepsilon] - 1}{\pi} \right)^m.$$

Observe that $\frac{(-1)^{m-1} \pi^m}{m}$ approaches zero p -adically when $m \rightarrow \infty$ and thus there exists $r \in \mathbb{N}$ such that $\frac{(-1)^{m-1} \pi^m}{m} \in \frac{1}{p^r} \mathbb{A}_{\max, K}(\Lambda^+)$ for every m . Hence the first assertion follows from the above observation and $[\varepsilon] - 1 \in \text{Ker } \theta_{\mathbb{A}_{\max, K}(\Lambda^+)}$. \square

Proposition 2.21. *We have*

$$\mathcal{O}_K \otimes_{W(k)} \mathbb{A}_{\inf}(\Lambda^+) \left[\frac{\text{Ker } \theta}{p} \right] \subset \mathbb{A}_{\inf, K}(\Lambda^+) \left[\frac{\text{Ker } \theta}{\pi} \right] \subset \frac{1}{p} \left(\mathcal{O}_K \otimes_{W(k)} \mathbb{A}_{\inf}(\Lambda^+) \left[\frac{\text{Ker } \theta}{p} \right] \right).$$

In particular,

$$K \otimes_{K_0} \mathbb{B}_{\max}^+(\Lambda^+) \cong \mathbb{B}_{\max, K}^+(\Lambda^+).$$

Proof. Set $e = [K : K_0]$ and write $p = a\pi^e$ and $p^b = a^b(\pi^b)^e$ for some $a \in \mathcal{O}_K^\times$ and $a^b \in (\mathcal{O}_K^\times)^\times$. For every $m \in \mathbb{N}$, we have

$$\left(\frac{[p^b]}{p} \right)^m = [a^b]^m a^{-m} \left(\frac{[\pi^b]}{\pi} \right)^{em}.$$

On the other hand, if we write $m = qe + r$ with $0 \leq r < e$, we have

$$\left(\frac{[\pi^b]}{\pi} \right)^m = \frac{1}{p} \left(a[a^b]^{-1} \frac{[p^b]}{p} \right)^q \frac{p}{\pi^r} [\pi^b]^r.$$

The first assertion follows from these equalities and Lemma 2.8. Taking the p -adic completion of the first part yields

$$\mathcal{O}_K \otimes_{W(k)} \mathbb{A}_{\max}(\Lambda^+) \subset \mathbb{A}_{\max, K}(\Lambda^+) \subset \frac{1}{p} \left(\mathcal{O}_K \otimes_{W(k)} \mathbb{A}_{\max}(\Lambda^+) \right).$$

Hence the second assertion follows by inverting p . \square

Let us now turn to Colmez's description of the horizontal period rings $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ and $\mathbb{B}_{\max, K}^+(\Lambda^+)$ as K -vector spaces [Col08, §8.4].

Construction 2.22. Fix a family of elements $(e_i)_{i \in I}$ of Λ^+ such that the images of e_i 's in $\Lambda^+/\pi\Lambda^+$ form a k -basis of $\Lambda^+/\pi\Lambda^+$. Then every element of Λ is written uniquely as $\sum_{i \in I} a_i e_i$ with $a_i \in K$ such that for every $r > 0$ there are only

finitely many i 's with $|a_i|_\Lambda \geq r$. For each $i \in I$, choose $\tilde{e}_i \in \mathbb{A}_{\text{inf},K}(\Lambda^+)$ such that $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}(\tilde{e}_i) = e_i$. Consider the map

$$s: \Lambda \rightarrow \mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]; \quad \sum_{i \in I} a_i e_i \mapsto \sum_{i \in I} a_i \tilde{e}_i.$$

This is a K -linear map and satisfies $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)} \circ s = \text{id}$ and $s(\Lambda^+) \subset \mathbb{A}_{\text{inf},K}(\Lambda^+)$. In particular, s is continuous with respect to p -adic topology.

Set

$$v = \pi^{-1}([\pi^b] - \pi).$$

Recall $\mathbb{B}_{\text{dR}}^+(\Lambda^+) = \varprojlim_m \mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]/(\text{Ker } \theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}[p^{-1}])^m$. Note that v generates $\text{Ker } \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}$ by Lemmas 2.8 and 2.12. For $x \in \mathbb{B}_{\text{dR}}^+(\Lambda^+)$, we define two sequences $(a_m(x))_{m \in \mathbb{N}}$ and $(b_m(x))_{m \in \mathbb{N}}$ with $a_m(x) \in \mathbb{B}_{\text{dR}}^+(\Lambda^+)$ and $b_m(x) \in \Lambda$ by

$$a_0(x) = x,$$

$$b_m(x) = \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}(a_m(x)), \quad a_{m+1}(x) = \frac{1}{v}(a_m(x) - s(b_m(x))).$$

Define

$$\tilde{\theta}_{s,v}: \mathbb{B}_{\text{dR}}^+(\Lambda^+) \rightarrow \Lambda[[X]] \quad \text{by} \quad \tilde{\theta}_{s,v}(x) := \sum_{m=0}^{\infty} b_m(x) X^m.$$

Lemma 2.23.

- (i) *The map $\tilde{\theta}_{s,v}$ is a K -linear isomorphism, and the inverse map is given by $\sum_{m=0}^{\infty} c_m X^m \mapsto \sum_{m=0}^{\infty} s(c_m) v^m$.*
- (ii) *For $F(X) \in K[[X]]$ and $x \in \mathbb{B}_{\text{dR}}^+(\Lambda^+)$, we have*

$$\tilde{\theta}_{s,v}(xF(v)) = \tilde{\theta}_{s,v}(x)F(X).$$

- (iii) *We have the following inclusions of \mathcal{O}_K -modules:*

$$\Lambda^+[[\pi^2 X]] \subset \tilde{\theta}_{s,v}(\mathbb{A}_{\text{inf},K}(\Lambda^+)) \subset \Lambda^+[[\pi X]].$$

- (iv) *The map $\tilde{\theta}_{s,v}$ induces natural identifications (as K -vector spaces)*

$$\mathbb{A}_{\text{max},K}(\Lambda^+) \cong \Lambda^+\langle X \rangle, \quad \text{and} \quad \mathbb{B}_{\text{max},K}^+(\Lambda^+) \cong \Lambda\langle X \rangle.$$

Here $\Lambda^+\langle X \rangle$ denotes the p -adic completion of the polynomial algebra $\Lambda^+[X]$ and $\Lambda\langle X \rangle := \Lambda^+\langle X \rangle[p^{-1}]$.

Proof. Part (i) follows from the fact that $\text{Ker } \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}$ is generated by $v = \pi^{-1}([\pi^b] - \pi)$, and Part (ii) follows from the definition of $\tilde{\theta}_{s,v}$ and the fact that $\theta_{\mathbb{B}_{\text{dR}}^+(\Lambda^+)}$ and s are both K -linear.

We show Part (iii). For $x \in \mathbb{A}_{\text{inf}}(\Lambda^+)$, we see $b_m(x) \in \pi^m \Lambda^+$ by induction on m . Hence the second inclusion follows. For the first inclusion, choose any $f \in \Lambda^+[[\pi^2 X]]$ and write $f = \sum_{m=0}^{\infty} c_m (\pi^2 X)^m$ with $c_m \in \Lambda^+$. Then $s(c_m) \in \mathbb{A}_{\text{inf},K}(\Lambda^+)$ and $y := \sum_{m=0}^{\infty} s(c_m) \pi^m ([\pi^b] - \pi)^m$ converges in $\mathbb{A}_{\text{inf},K}(\Lambda^+)$. It is now easy to see $\tilde{\theta}_{s,v}(y) = f$.

For Part (iv), first note that Parts (ii) and (iii) imply

$$\Lambda^+[[\pi^2 X]][X] \subset \tilde{\theta}_{s,v} \left(\mathbb{A}_{\text{inf},K}(\Lambda^+) \left[\frac{\text{Ker } \theta}{\pi} \right] \right) \subset \Lambda^+[[\pi X]][X].$$

Hence the identification $\mathbb{A}_{\max,K}(\Lambda^+) \cong \Lambda^+ \langle X \rangle$ follows from

$$\varprojlim_m \Lambda^+ [[\pi^2 X]][X] / p^m \Lambda^+ [[\pi^2 X]][X] \cong \varprojlim_m \Lambda^+ [[\pi X]][X] / p^m \Lambda^+ [[\pi X]][X] \cong \Lambda^+ \langle X \rangle.$$

By inverting p , we obtain $\mathbb{B}_{\max,K}^+(\Lambda^+) \cong \Lambda \langle X \rangle$. \square

We will show that there is an injective K -algebra homomorphism

$$\mathbb{B}_{\max,K}^+(\Lambda^+) \rightarrow \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+).$$

Since $\mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ is $\mathrm{Ker} \theta_{\mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)}^+$ -adically complete, the natural K -algebra homomorphism $\mathbb{A}_{\mathrm{inf},K}(\Lambda^+) \left[\frac{\mathrm{Ker} \theta}{\pi} \right] \rightarrow \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ extends to $\mathbb{A}_{\max,K}(\Lambda^+) \rightarrow \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ and thus to $\mathbb{B}_{\max,K}^+(\Lambda^+) \rightarrow \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$. We deduce the injectivity from the description of these period rings in terms of $\tilde{\theta}_{s,v}$ as follows:

Proposition 2.24. *The map $\tilde{\theta}_{s,v}$ induces identifications (as K -vector spaces)*

$$\begin{array}{ccccc} \mathbb{A}_{\max,K}(\Lambda^+) & \longrightarrow & \mathbb{B}_{\max,K}^+(\Lambda^+) & \longrightarrow & \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \tilde{\theta}_{v,s} \cong \\ \Lambda^+ \langle X \rangle & \hookrightarrow & \Lambda \langle X \rangle & \hookrightarrow & \Lambda[[X]], \end{array}$$

making the diagram commutative. In particular, the top horizontal maps are injective, and $\mathbb{B}_{\max,K}^+(\Lambda)$ is t -torsion free.

Proof. We have already proved that the vertical maps are all isomorphisms. Since the bottom horizontal maps are injective, so are the top horizontal maps. Since $\mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ is t -torsion free, the last assertion follows. \square

Definition 2.25. We set

$$\begin{aligned} \mathbb{B}_{\max}(\Lambda^+) &= \mathbb{B}_{\max}(\Lambda, \Lambda^+) := \mathbb{B}_{\max}^+(\Lambda^+)[t^{-1}], \quad \text{and} \\ \mathbb{B}_{\max,K}(\Lambda^+) &= \mathbb{B}_{\max,K}(\Lambda, \Lambda^+) := \mathbb{B}_{\max,K}^+(\Lambda^+)[t^{-1}]. \end{aligned}$$

Since $\varphi(t) = pt$, the Frobenius φ on $\mathbb{A}_{\max}(\Lambda^+)$ extends to an endomorphism on $\mathbb{B}_{\max}(\Lambda^+)$, which we still call the Frobenius and denote by φ .

Let us define the horizontal semistable period rings. Consider the elements

$$\log \frac{[p^b]}{p} := \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} \left(\frac{[p^b]}{p} - 1 \right)^m, \quad \log \frac{[\pi^b]}{\pi} := \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} \left(\frac{[\pi^b]}{\pi} - 1 \right)^m \in \mathbb{B}_{\mathrm{dR}}^+(\Lambda^+).$$

Definition 2.26. Define $\mathbb{B}_{\mathrm{st}}^+(\Lambda^+) = \mathbb{B}_{\mathrm{st}}^+(\Lambda, \Lambda^+)$ to be the $\mathbb{B}_{\max}^+(\Lambda^+)$ -subalgebra of $\mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ generated by $\log \frac{[p^b]}{p}$. Similarly, define $\mathbb{B}_{\mathrm{st},K}^+(\Lambda^+) = \mathbb{B}_{\mathrm{st},K}^+(\Lambda, \Lambda^+)$ to be the $\mathbb{B}_{\max,K}^+(\Lambda^+)$ -subalgebra of $\mathbb{B}_{\mathrm{dR}}^+(\Lambda^+)$ generated by $\log \frac{[p^b]}{p}$.⁵

Lemma 2.27. *The definitions of $\mathbb{B}_{\mathrm{st}}^+(\Lambda^+)$ and $\mathbb{B}_{\mathrm{st},K}^+(\Lambda^+)$ are independent of the choice of a compatible system of p -power roots (p_m) of p .*

⁵Strictly speaking, our definitions slightly differ from the ones in [Col08, §8.6]. However, our horizontal semistable period rings are isomorphic to Colmez's period rings by Corollary 2.31.

Proof. If (p'_m) is a different choice, there exists $\alpha \in \mathbb{Z}_p$ such that $p'_m = \varepsilon_m^\alpha p_m$ for every $m \in \mathbb{N}$. It follows

$$\log \frac{[p'^b]}{p} = \log \frac{[p^b]}{p} + \alpha t \quad \text{in } \mathbb{B}_{\text{dR}}^+(\Lambda^+) \text{ with } \alpha t \in \mathbb{B}_{\text{max}}^+(\Lambda^+).$$

□

Lemma 2.28. *The ring $\mathbb{B}_{\text{st},K}^+(\Lambda^+)$ coincides with the $\mathbb{B}_{\text{max},K}^+(\Lambda^+)$ -subalgebra of $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ generated by $\log \frac{[\pi^b]}{\pi}$.*

Proof. Set $e = [K : K_0]$. Write $p = a\pi^e$ for some $a \in \mathcal{O}_K^\times$ and take $a^b \in (\mathcal{O}_C^b)^\times$ such that $p^b = a^b(\pi^b)^e$. Then $[a^b]a^{-1} \in \mathbb{A}_{\text{inf},K}(\Lambda^+)$ satisfies $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}([a^b]a^{-1} - 1) = 0$. It follows that the series

$$\sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} \left(\frac{[a^b]}{a} - 1 \right)^m$$

converges in $\mathbb{A}_{\text{max},K}(\Lambda^+)$. We denote the limit by $\log \frac{[a^b]}{a}$. We can also check that the image of $\log \frac{[a^b]}{a}$ in $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$ is given by the same series and satisfies

$$\log \frac{[p^b]}{p} = \log \frac{[a^b]}{a} + e \log \frac{[\pi^b]}{\pi}.$$

Hence the assertion follows. □

We will describe $\mathbb{B}_{\text{st},K}^+(\Lambda^+)$ via $\tilde{\theta}_{s,v}$. For this we need the following fact:

Lemma 2.29. *Let A be a uniform \mathbb{Q}_p -Banach algebra. Then the power series $\log(1+X) := \sum_{m=0}^{\infty} \frac{(-1)^{m-1}}{m} X^m \in A[[X]]$ is transcendental over $A\langle X \rangle$. Hence the $A\langle X \rangle$ -subalgebra $A\langle X \rangle[\log(1+X)]$ of $A[[X]]$ generated by $\log(1+X)$ is isomorphic to a polynomial algebra over $A\langle X \rangle$.*

Proof. Since A is uniform, equip A with its spectral norm. Then the Gel'fand transform $A \rightarrow \prod_{x \in \mathcal{M}(A)} \mathcal{H}(x)$ is isometric with each $\mathcal{H}(x)$ a field ([Ber90, Corollary 1.3.2 (ii)]). We can thus reduce the general case to the case where A is a field. We may further assume that A is a complete algebraically closed field. In this case, the assertion is well-known. In fact, suppose that there exist $f_0(X), \dots, f_m(X) \in A\langle X \rangle$ such that

$$f_0(X) + f_1(X) \log(1+X) + \dots + f_m(X) (\log(1+X))^m = 0.$$

If ζ is a p -power root of unity, then $X = \zeta - 1$ is a root of $\log(1+X)$, and thus it is also a root of $f_0(X)$. Since any nonzero element of $A\langle X \rangle$ has at most finitely many roots in \mathcal{O}_A , we have $f_0(X) = 0$. By repeating the same argument, we conclude that $f_0(X), \dots, f_m(X)$ are all zero. □

Proposition 2.30. *The map $\tilde{\theta}_{s,v}$ induces identifications (as K -vector spaces)*

$$\begin{array}{ccccc} \mathbb{B}_{\text{max},K}^+(\Lambda^+) & \longrightarrow & \mathbb{B}_{\text{st},K}^+(\Lambda^+) & \longrightarrow & \mathbb{B}_{\text{dR}}^+(\Lambda^+) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ \Lambda\langle X \rangle & \longrightarrow & \Lambda\langle X \rangle[\log(1+X)] & \longrightarrow & \Lambda[[X]], \end{array}$$

sending $\log \frac{[\pi^b]}{\pi}$ to $\log(1+X)$.

Proof. Recall $v = \frac{[\pi^b]}{\pi} - 1$. Hence we have $\tilde{\theta}_{s,v}((\log \frac{[\pi^b]}{\pi})^m) = (\log(1+X))^m$ by Lemma 2.23(ii). By Lemma 2.28,

$$\mathbb{B}_{\text{st},K}^+(\Lambda^+) = \sum_{m=0}^{\infty} \mathbb{B}_{\text{max},K}^+(\Lambda^+) \left(\log \frac{[\pi^b]}{\pi} \right)^m$$

as K -vector subspaces of $\mathbb{B}_{\text{dR}}^+(\Lambda^+)$. It follows from Lemma 2.23(ii) and (iv) that

$$\tilde{\theta}_{s,v}(\mathbb{B}_{\text{st},K}^+(\Lambda^+)) = \sum_{m=0}^{\infty} \Lambda\langle X \rangle (\log(1+X))^m \subset \Lambda[[X]].$$

□

Corollary 2.31. *The $\mathbb{B}_{\text{max}}^+(\Lambda^+)$ -algebra homomorphism from the polynomial ring $\mathbb{B}_{\text{max}}^+(\Lambda^+)[u]$ to $\mathbb{B}_{\text{st}}^+(\Lambda^+)$ sending u to $\log \frac{[p^b]}{p}$ is an isomorphism. Similarly, the $\mathbb{B}_{\text{max},K}^+(\Lambda^+)$ -algebra homomorphism $\mathbb{B}_{\text{max},K}^+(\Lambda^+)[u] \rightarrow \mathbb{B}_{\text{st},K}^+(\Lambda^+); u \mapsto \log \frac{[p^b]}{p}$ is an isomorphism. In particular, $K \otimes_{K_0} \mathbb{B}_{\text{st}}^+(\Lambda^+) \cong \mathbb{B}_{\text{st},K}^+(\Lambda^+)$.*

Proof. For the second assertion, we only need to show that the homomorphism is bijective. Hence it follows from Lemmas 2.23(ii), 2.28, 2.29, and Proposition 2.30. The first assertion is reduced to the second assertion by considering the case $K = K_0$. The last assertion follows from the first two since $K \otimes_{K_0} \mathbb{B}_{\text{max}}^+(\Lambda^+) \cong \mathbb{B}_{\text{max},K}^+(\Lambda^+)$. □

Definition 2.32. We extend the Frobenius on $\mathbb{B}_{\text{max}}(\Lambda^+)$ to the polynomial ring $\mathbb{B}_{\text{max}}^+(\Lambda^+)[u]$ by setting $\varphi(u) = pu$. Then the $\mathbb{B}_{\text{max}}^+(\Lambda^+)$ -linear derivation $N = -\frac{d}{du}$ on $\mathbb{B}_{\text{max}}^+(\Lambda^+)[u]$ satisfies $N\varphi = p\varphi N$. Via the $\mathbb{B}_{\text{max}}^+(\Lambda^+)$ -algebra isomorphism

$$\mathbb{B}_{\text{max}}^+(\Lambda^+)[u] \xrightarrow{\cong} \mathbb{B}_{\text{st}}^+(\Lambda^+); \quad u \mapsto \log \frac{[p^b]}{p},$$

we define the Frobenius φ and the monodromy operator N on $\mathbb{B}_{\text{st}}^+(\Lambda^+)$. Note $N\varphi = p\varphi N$ and

$$\varphi\left(\log \frac{[p^b]}{p}\right) = p \log \frac{[p^b]}{p}, \quad \text{and} \quad N\left(\left(\log \frac{[p^b]}{p}\right)^m\right) = -m \left(\log \frac{[p^b]}{p}\right)^{m-1}.$$

As in Lemma 2.27, we can check that the definitions of φ and N on $\mathbb{B}_{\text{st}}^+(\Lambda^+)$ are independent of the choice of a compatible system of p -power roots of p .

Definition 2.33. Set

$$\begin{aligned} \mathbb{B}_{\text{st}}(\Lambda^+) &= \mathbb{B}_{\text{st}}(\Lambda, \Lambda^+) := \mathbb{B}_{\text{st}}^+(\Lambda^+)[t^{-1}], \quad \text{and} \\ \mathbb{B}_{\text{st},K}(\Lambda^+) &= \mathbb{B}_{\text{st},K}(\Lambda, \Lambda^+) := \mathbb{B}_{\text{st},K}^+(\Lambda^+)[t^{-1}]. \end{aligned}$$

Note that there are isomorphisms

$$\mathbb{B}_{\text{max}}(\Lambda^+)[u] \xrightarrow{\cong} \mathbb{B}_{\text{st}}(\Lambda^+), \quad \text{and} \quad \mathbb{B}_{\text{max},K}(\Lambda^+)[u] \xrightarrow{\cong} \mathbb{B}_{\text{st},K}(\Lambda^+)$$

sending u to $\log \frac{[p^b]}{p}$. Moreover, the Frobenius φ (resp. the monodromy operator N) on $\mathbb{B}_{\text{st}}^+(\Lambda^+)$ extends to a ring (resp. $\mathbb{B}_{\text{max}}^+(\Lambda^+)$ -linear) endomorphism on $\mathbb{B}_{\text{st}}(\Lambda^+)$, which we still denote by φ (resp. N).

Lemma 2.34. *The sequence*

$$0 \longrightarrow \mathbb{B}_{\max}(\Lambda^+) \longrightarrow \mathbb{B}_{\text{st}}(\Lambda^+) \xrightarrow{N} \mathbb{B}_{\text{st}}(\Lambda^+) \longrightarrow 0$$

is exact.

Proof. This follows easily from the isomorphism $\mathbb{B}_{\max}(\Lambda^+)[u] \xrightarrow{\cong} \mathbb{B}_{\text{st}}(\Lambda^+)$ commuting with N . \square

We also introduce the horizontal crystalline period ring for completeness:

Definition 2.35. Let $\mathbb{A}'_{\text{cris}}(\Lambda^+)$ denote the PD-envelope of $\mathbb{A}_{\text{inf}}(\Lambda^+)$ relative to the ideal $\text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$ compatible with the canonical PD-structure on $p\mathbb{A}_{\text{inf}}(\Lambda^+)$. Define $\mathbb{A}_{\text{cris}}(\Lambda^+) = \mathbb{A}_{\text{cris}}(\Lambda, \Lambda^+)$ to be the p -adic completion of $\mathbb{A}'_{\text{cris}}(\Lambda^+)$.

Note that the natural inclusion $\mathbb{A}'_{\text{cris}}(\Lambda^+) \hookrightarrow \mathbb{A}_{\text{inf}}(\Lambda^+)[\frac{\text{ker } \theta}{p}]$ yields an inclusion

$$\mathbb{A}_{\text{cris}}(\Lambda^+) \hookrightarrow \mathbb{A}_{\max}(\Lambda^+).$$

Lemma 2.36. *The Frobenius φ on $\mathbb{A}_{\text{inf}}(\Lambda^+)$ extends to a ring endomorphism φ on $\mathbb{A}_{\text{cris}}(\Lambda^+)$.*

Proof. Since $\varphi(x) \equiv x^p \pmod{p\mathbb{A}_{\text{inf}}(\Lambda^+)}$ for $x \in \mathbb{A}_{\text{inf}}(\Lambda^+)$, the Frobenius φ on $\mathbb{A}_{\text{inf}}(\Lambda^+)$ extends to a ring endomorphism on $\mathbb{A}'_{\text{cris}}(\Lambda^+)$. Passing to the p -adic completion, it further extends to a ring endomorphism on $\mathbb{A}_{\text{cris}}(\Lambda^+)$. \square

By construction, the inclusion $\mathbb{A}_{\text{cris}}(\Lambda^+) \hookrightarrow \mathbb{A}_{\max}(\Lambda^+)$ is φ -equivariant.

Lemma 2.37. *We have*

$$\varphi(\mathbb{A}_{\max}(\Lambda^+)) \subset \mathbb{A}_{\text{cris}}(\Lambda^+) \subset \mathbb{A}_{\max}(\Lambda^+).$$

Proof. We need to prove the first inclusion. For this, it is enough to show

$$\varphi\left(\mathbb{A}_{\text{inf}}(\Lambda^+)\left[\frac{\text{ker } \theta}{p}\right]\right) \subset \mathbb{A}'_{\text{cris}}(\Lambda^+).$$

Choose $x \in \text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\Lambda^+)}$ and write $\varphi(x) = x^p + py$ for some $y \in \mathbb{A}_{\text{inf}}(\Lambda^+)$. Then we have

$$\varphi\left(\frac{x}{p}\right) = (p-1)!x^{[p]} + y \in \mathbb{A}'_{\text{cris}}(\Lambda^+).$$

\square

Definition 2.38. Set

$$\begin{aligned} \mathbb{B}_{\text{cris}}^+(\Lambda^+) &= \mathbb{B}_{\text{cris}}^+(\Lambda, \Lambda^+) := \mathbb{A}_{\text{cris}}(\Lambda)[p^{-1}], \quad \text{and} \\ \mathbb{B}_{\text{cris}}(\Lambda^+) &= \mathbb{B}_{\text{cris}}(\Lambda, \Lambda^+) := \mathbb{B}_{\text{cris}}(\Lambda)[t^{-1}]. \end{aligned}$$

The Frobenius φ on $\mathbb{A}_{\text{cris}}(\Lambda^+)$ naturally extends over these rings, and the previous lemma implies

$$\varphi(\mathbb{B}_{\max}(\Lambda^+)) \subset \mathbb{B}_{\text{cris}}(\Lambda^+) \subset \mathbb{B}_{\max}(\Lambda^+).$$

Remark 2.39. The construction of all the period rings is functorial on perfectoid C -Banach pairs (Λ, Λ^+) with morphisms of K -Banach pairs.

The following lemma will be heavily used in the proof of the purity theorem for horizontal semistable representations (Theorem 6.1).

Lemma 2.40. *Let (Λ', Λ'^+) be another perfectoid C -Banach pair such that Λ'^+ is an \mathcal{O}_C -algebra, and let $f^+ : \Lambda^+ \hookrightarrow \Lambda'^+$ be an injective \mathcal{O}_K -algebra homomorphism with p -torsion free cokernel. Then the natural commutative diagram*

$$\begin{array}{ccc} \mathbb{B}_{\text{st},K}^+(\Lambda^+) & \longrightarrow & \mathbb{B}_{\text{dR},K}^+(\Lambda^+) \\ \downarrow & & \downarrow \\ \mathbb{B}_{\text{st},K}^+(\Lambda'^+) & \longrightarrow & \mathbb{B}_{\text{dR},K}^+(\Lambda'^+) \end{array}$$

is Cartesian with injective vertical maps.

Proof. By construction of the period rings, we have a natural commutative diagram as in the statement. We will prove that it is Cartesian.

Let $f : \Lambda \rightarrow \Lambda'$ denote the induced K -algebra homomorphism from f^+ by inverting p . We also change the norm on Λ (resp. Λ') so that Λ^+ (resp. Λ'^+) becomes the unit ball. Then f is a closed isometric embedding of K -Banach algebras; by p -torsion freeness of $\text{Coker } f^+$, the induced map $\Lambda^+/p^n\Lambda^+ \rightarrow \Lambda'^+/p^n\Lambda'^+$ is injective for all $n \in \mathbb{N}$, and thus f is closed and isometric.

The commutative diagram in question only depends on f^+ as an \mathcal{O}_K -algebra homomorphism and not on the entire C -Banach structures on Λ and Λ' . Hence by replacing the C -Banach structure $C \rightarrow \Lambda'$ by precomposing an isometric K -algebra automorphism $C \rightarrow C$, we may further assume $f(\pi_n) = \pi_n$ for every $n \in \mathbb{N}$.

We will explain that Construction 2.22 is made compatibly with $f^+ : \Lambda^+ \hookrightarrow \Lambda'^+$. Since the map $\Lambda^+/\pi\Lambda^+ \rightarrow \Lambda'^+/\pi\Lambda'^+$ is injective, choose a family of elements $(e_i)_{i \in I}$ of Λ^+ and a family of elements $(e'_j)_{j \in J}$ of Λ'^+ such that the images of e_i 's in $\Lambda^+/\pi\Lambda^+$ form a k -basis of $\Lambda^+/\pi\Lambda^+$ and such that the images of $f^+(e_i)$'s and e'_j 's in $\Lambda'^+/\pi\Lambda'^+$ form a k -basis of $\Lambda'^+/\pi\Lambda'^+$; this is possible by lifting k -bases of $\Lambda^+/\pi\Lambda^+ \subset \Lambda'^+/\pi\Lambda'^+$. We also choose a lift $\tilde{e}_i \in \mathbb{A}_{\text{inf},K}(\Lambda^+)$ of e_i with respect to $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda^+)}$ and a lift $\tilde{e}'_j \in \mathbb{A}_{\text{inf},K}(\Lambda'^+)$ of e'_j with respect to $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda'^+)}$.

Consider the K -linear maps

$$\begin{aligned} s : \Lambda &\rightarrow \mathbb{A}_{\text{inf},K}(\Lambda^+)[p^{-1}]; \quad \sum_{i \in I} a_i e_i \mapsto \sum_{i \in I} a_i \tilde{e}_i, \quad \text{and} \\ s' : \Lambda' &\rightarrow \mathbb{A}_{\text{inf},K}(\Lambda'^+)[p^{-1}]; \quad \sum_{i \in I} a_i f(e_i) + \sum_{j \in J} b_j e'_j \mapsto \sum_{i \in I} a_i \mathbb{A}_{\text{inf},K}(f)(\tilde{e}_i) + \sum_{j \in J} b_j \tilde{e}'_j. \end{aligned}$$

Then we have $\mathbb{A}_{\text{inf},K}(f) \circ s = s' \circ f$.

Since $f(\pi_n) = \pi_n$ for every $n \in \mathbb{N}$, we have $\mathbb{A}_{\text{inf},K}(f)([\pi^b]) = [\pi^b]$. Hence we apply Construction 2.22 to the maps s and s' with $v = \pi^{-1}([\pi^b] - \pi)$ and obtain the commutative diagram

$$\begin{array}{ccc} \mathbb{B}_{\text{dR},K}^+(\Lambda^+) & \xrightarrow[\cong]{\tilde{\theta}_{s,v}} & \Lambda[[X]] \\ \downarrow & & \downarrow \\ \mathbb{B}_{\text{dR},K}^+(\Lambda'^+) & \xrightarrow[\cong]{\tilde{\theta}_{s',v}} & \Lambda'[[X]]. \end{array}$$

By these maps and Proposition 2.30, it suffices to show that the commutative diagram

$$\begin{array}{ccc} \Lambda\langle X \rangle[\log(1+X)] & \longrightarrow & \Lambda[[X]] \\ \downarrow & & \downarrow \\ \Lambda'\langle X \rangle[\log(1+X)] & \longrightarrow & \Lambda'[[X]] \end{array}$$

is Cartesian. This follows from Lemma 2.41 below. \square

Lemma 2.41. *Let A be a \mathbb{Q}_p -Banach algebra and B a closed subalgebra of A . If*

$$g(x) \in \bigoplus_{i=0}^h A\langle X \rangle \log(1+X)^i$$

satisfies $g(x) \in B[[X]]$, then

$$g(x) \in \bigoplus_{i=0}^h B\langle X \rangle \log(1+X)^i.$$

Proof. This is stated and proved in the proof of [BC08, Lemme 6.3.1]. \square

Remark 2.42. Assume that Λ is sympathetic (see [Col08, p.356] for definition). Under this assumption, Colmez shows that the natural sequence

$$0 \rightarrow \mathbb{Q}_p \rightarrow \mathbb{B}_{\max}(\Lambda^\circ)^{\varphi=1} \rightarrow \mathbb{B}_{\mathrm{dR}}(\Lambda^\circ)/\mathbb{B}_{\mathrm{dR}}^+(\Lambda^\circ) \rightarrow 0$$

is exact [Col08, Proposition 9.25]. The sequence is called the fundamental exact sequence. The exactness does not hold for perfectoid C -Banach pairs (Λ, Λ^+) in general. See also Proposition 4.20.

3. PRELIMINARIES ON BASE RINGS

In this section, we review the class of base rings which is considered in [Bri08]. As in the previous section, let k be a perfect field of characteristic p and set $K_0 := W(k)[p^{-1}]$. Let K be a totally ramified finite extension of K_0 and fix a uniformizer π of K . We denote by C the p -adic completion of an algebraic closure \overline{K} of K .

Let $\mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle$ denote the p -adic completion of $\mathcal{O}_K[T_1^{\pm 1}, \dots, T_n^{\pm 1}]$.

Set-up 3.1 ([Bri08, p. 7]). Let \tilde{R} be a ring obtained from $\mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle$ by a finite number of iterations of the following operations:

- p -adic completion of an étale extension;
- p -adic completion of a localization;
- completion with respect to an ideal containing p .

We further assume that either $\mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle \rightarrow \tilde{R}$ has geometrically regular fibers or \tilde{R} has Krull dimension less than two, and that $k \rightarrow \tilde{R}/\pi\tilde{R}$ is geometrically integral. Let R be an \tilde{R} -algebra such that

- (i) R is an integrally closed domain;
- (ii) R is finite and flat over \tilde{R} ;
- (iii) $R[p^{-1}]$ is étale over $\tilde{R}[p^{-1}]$;
- (iv) K is algebraically closed in $R[p^{-1}]$.

Definition 3.2 ([Bri08, p. 9]). We say that R satisfies (BR) if $R = \widetilde{R}$. This is a condition on good reduction (bonne réduction), and [Bri08] often assumes it in the study of (horizontal) crystalline representations.

Set $R_K := R[p^{-1}]$. Fix an algebraic closure $\overline{\text{Frac } R}$ of $\text{Frac } R$ containing \overline{K} . Let \overline{R} be the union of finite R -subalgebras R' of $\overline{\text{Frac } R}$ such that $R'[p^{-1}]$ is étale over R_K . Note that \overline{R} is a normal domain as it can be written as an increasing union of such (by replacing each R' with its normalization). Let $\widehat{\overline{R}}$ denote the p -adic completion of \overline{R} . Then $\widehat{\overline{R}}$ is an \mathcal{O}_C -algebra. Note that $\widehat{\overline{R}}[p^{-1}]$ is the union of finite Galois extensions of R_K , and thus we set⁶

$$\mathcal{G}_{R_K} := \text{Gal}(\widehat{\overline{R}}[p^{-1}]/R_K).$$

It is a profinite group acting on \overline{R} , and this action extends to $\widehat{\overline{R}}$ continuously.

Let us summarize the properties of $\widehat{\overline{R}}$.

Lemma 3.3 ([Bri08, Proposition 2.0.3]). *The ring $\widehat{\overline{R}}$ is p -torsion free. Moreover, $\overline{R} \rightarrow \widehat{\overline{R}}$ is injective, and $\overline{R} \cap p\widehat{\overline{R}} = p\overline{R}$.*

We will explain that the ring $\widehat{\overline{R}}[p^{-1}]$ admits a C -Banach algebra structure and that $(\widehat{\overline{R}}[p^{-1}], \widehat{\overline{R}})$ is a perfectoid C -Banach pair (hence we can apply the results in the previous section). For $x \in \widehat{\overline{R}}[p^{-1}]$, set

$$|x| = \inf\{|a|_C \mid a \in C, x \in a\widehat{\overline{R}}\}.$$

Note that we normalize the norm $|\cdot|_C$ on C by $|p|_C = p^{-1}$.

Lemma 3.4. *The function $|\cdot|$ is a C -algebra norm on $\widehat{\overline{R}}[p^{-1}]$, and $\widehat{\overline{R}}[p^{-1}]$ is complete with respect to this norm. Namely, $\widehat{\overline{R}}[p^{-1}]$ becomes a C -Banach algebra with this norm. Moreover, $\widehat{\overline{R}}$ is an open subring of $\widehat{\overline{R}}[p^{-1}]$, and the induced topology on $\widehat{\overline{R}}$ agrees with the p -adic topology (i.e., the topology defined by the ideal $p\widehat{\overline{R}}$).*

Proof. By construction, we see $\widehat{\overline{R}} \cap C = \mathcal{O}_C$ inside $\widehat{\overline{R}}[p^{-1}]$. Since $\widehat{\overline{R}}$ is p -torsion free and p -adically separated, the function $|\cdot|$ is a C -algebra norm. The p -adic completeness of $\widehat{\overline{R}}$ implies that $\widehat{\overline{R}}[p^{-1}]$ is complete with respect to $|\cdot|$. Finally, the second assertion follows from the inclusions

$$p^{m+1}\widehat{\overline{R}} \subset \{x \in \widehat{\overline{R}}[p^{-1}] \mid |x| \leq p^{-(m+1)}\} \subset p^m\widehat{\overline{R}}.$$

□

Lemma 3.5. *The C -Banach algebra $\widehat{\overline{R}}[p^{-1}]$ is uniform.*

Proof. Since $p^{-1}\widehat{\overline{R}}$ is a bounded subset, it suffices to show $(\widehat{\overline{R}}[p^{-1}])^\circ \subset p^{-1}\widehat{\overline{R}}$.

First we prove $(\widehat{\overline{R}}[p^{-1}])^\circ \cap \overline{R}[p^{-1}] \subset p^{-1}\overline{R}$. Take $x \in (\widehat{\overline{R}}[p^{-1}])^\circ \cap \overline{R}[p^{-1}]$. Then the sequence $(p^n x^n)_n$ converges to zero as $n \rightarrow \infty$. In particular, $(px)^n \in \overline{R}$ for some n . Since \overline{R} is a normal domain, we have $px \in \overline{R}$ and thus $x \in p^{-1}\overline{R}$.

⁶Our \mathcal{G}_{R_K} is denoted by \mathcal{G}_R in [Bri08].

For $x \in (\widehat{R}[p^{-1}])^\circ$, write $x = u + v$ with $u \in \overline{R}[p^{-1}]$ and $v \in \widehat{R}$. Since $(\widehat{R}[p^{-1}])^\circ$ is a ring containing \widehat{R} , we have $u \in (\widehat{R}[p^{-1}])^\circ$. Hence $u \in p^{-1}\overline{R}$ by what we have proved, and we conclude $x \in p^{-1}\widehat{R}$. \square

Lemma 3.6. *The subring \widehat{R} of $\widehat{R}[p^{-1}]$ is a ring of integral elements.*

Proof. We need to show that \widehat{R} is integrally closed in $\widehat{R}[p^{-1}]$.

Let $x \in \widehat{R}[p^{-1}]$ be an element that is integral over \widehat{R} and suppose that x satisfies

$$x^m + a_{m-1}x^{m-1} + \cdots + a_0 = 0$$

for some $a_0, \dots, a_{m-1} \in \widehat{R}$. For each $l \in \mathbb{N}$, there exist $y_l \in \overline{R}[p^{-1}]$ and $z_l \in p^l \widehat{R}$ such that $x = y_l + z_l$. Similarly, for each $i = 0, \dots, m-1$, there exists $a_{i,l} \in \overline{R}$ such that $a_i - a_{i,l} \in p^l \widehat{R}$. We compute

$$\begin{aligned} & |y_l^m + a_{m-1,l}y_l^{m-1} + \cdots + a_{0,l}| \\ &= |(y_l^m + a_{m-1,l}y_l^{m-1} + \cdots + a_{0,l}) - (x^m + a_{m-1}x^{m-1} + \cdots + a_0)| \\ &\leq \max\{|(y_l^m + a_{m-1,l}y_l^{m-1} + \cdots + a_{0,l}) - (x^m + a_{m-1,l}x^{m-1} + \cdots + a_{0,l})|, \\ &\quad |(x^m + a_{m-1,l}x^{m-1} + \cdots + a_{0,l}) - (x^m + a_{m-1}x^{m-1} + \cdots + a_0)|\} \\ &\leq |p^l| \max\{1, |x|^{m-1}\}. \end{aligned}$$

Since $\{y \in \widehat{R}[p^{-1}] \mid |y| < 1\} \subset \widehat{R}$ and $\overline{R}[p^{-1}] \cap \widehat{R} = \overline{R}$, we see that $y_l^m + a_{m-1,l}y_l^{m-1} + \cdots + a_{0,l} \in \overline{R}$ for $l \gg 0$. For such l , we have $y_l \in \overline{R}$ by normality of \overline{R} and thus $x = y_l + z_l \in \widehat{R}$. \square

Lemma 3.7. *The C -Banach pair $(\widehat{R}[p^{-1}], \widehat{R})$ is perfectoid.*

Proof. It remains to prove that the Frobenius on $\widehat{R}/p\widehat{R}$ is surjective. Note that $\widehat{R}/p\widehat{R} = \overline{R}/p\overline{R}$. Let $\bar{x} \in \overline{R}/p\overline{R}$ and choose a lift $x \in \overline{R}$ of \bar{x} . Set $R' := R[x][T]/(T^{p^2} + pT - x)$. Then R' is finite over R and $R'[p^{-1}]$ is étale over R_K . Choose an embedding of R' into $\overline{\text{Frac}} \overline{R}$ and let y denote the image of T . Then $y \in \overline{R}$ and it satisfies $(y^p)^p \equiv x \pmod{p\overline{R}}$. Hence the Frobenius on $\overline{R}/p\overline{R}$ is surjective. \square

4. HORIZONTAL CRYSTALLINE, SEMISTABLE, AND DE RHAM REPRESENTATIONS

In this section, we define horizontal crystalline, semistable, and de Rham representations of \mathcal{G}_{R_K} , and study their basic properties. We mainly follow [Bri08, §8], where horizontal crystalline and de Rham representations are studied.

4.1. Admissible representations. We review the notion of admissible representations. Let E be a finite extension of \mathbb{Q}_p and G a topological group.

Definition 4.1. An E -representation of G is a finite-dimensional E -vector space equipped with a continuous action of G . A morphism of E -representations of G is an E -linear map commuting with actions of G . We denote the category of E -representations of G by $\text{Rep}_E(G)$. It is naturally a Tannakian category.

Remark 4.2. Note that [Fon94b, §1] deals with an abstract field E and an abstract group G but that properties discussed there continue to hold in our setting.

Definition 4.3 (cf. [Fon94b, 1.3]).

- (i) An (E, G) -ring is an E -algebra B equipped with an E -linear action of G .
- (ii) Let B be an (E, G) -ring and let $V \in \text{Rep}_E(G)$. We set

$$D_B(V) := (B \otimes_E V)^G,$$

and we let

$$\alpha_B(V) : B \otimes_{B^G} D_B(V) \rightarrow B \otimes_E V$$

denote the B -linear map induced from the inclusion $D_B(V) \subset B \otimes_E V$.

We say that V is B -admissible if $\alpha_B(V)$ is an isomorphism of B -modules.

Lemma 4.4. *Let G' be another topological group and let $\alpha : G' \rightarrow G$ be a continuous group homomorphism. Let B be an (E, G) -ring and B' an (E, G') -ring equipped with an E -algebra homomorphism $\beta : B \rightarrow B'$ satisfying*

$$\beta(\alpha(g') \cdot b) = g' \cdot \beta(b) \quad \text{for all } g' \in G' \text{ and } b \in B.$$

Let $V \in \text{Rep}_E(G)$ and let $V|_{G'}$ denote the corresponding E -representation of G' . If V is B -admissible and $\alpha_{B'}(V|_{G'})$ is injective, then $V|_{G'}$ is B' -admissible, and the natural map

$$(B')^{G'} \otimes_{B^G} D_B(V) \rightarrow D_{B'}(V|_{G'})$$

is an isomorphism.

Proof. Consider the commutative diagram

$$\begin{array}{ccc} B \otimes_{B^G} D_B(V) & \xrightarrow[\cong]{\alpha_B(V)} & B \otimes_E V \\ \downarrow & & \downarrow \\ B' \otimes_{(B')^{G'}} D_{B'}(V|_{G'}) & \xrightarrow{\alpha_{B'}(V|_{G'})} & B' \otimes_E V. \end{array}$$

It follows that $V \subset B' \otimes_E V$ is contained in the image of $\alpha_{B'}(V|_{G'})$. Since $\alpha_{B'}(V|_{G'})$ is injective and B' -linear, it is an isomorphism. We then have a G' -equivariant isomorphism

$$B' \otimes_B (B \otimes_{B^G} D_B(V)) \xrightarrow{\cong} B' \otimes_{(B')^{G'}} D_{B'}(V|_{G'}).$$

Taking G' -invariants yields an isomorphism $(B')^{G'} \otimes_{B^G} D_B(V) \xrightarrow{\cong} D_{B'}(V|_{G'})$. \square

Proposition 4.5 ([Bri06, Proposition 3.5]). *Let B be an (E, G) -ring and assume that B is faithfully flat over $F := B^G$ and that $\alpha_B(V)$ is injective for every $V \in \text{Rep}_E(G)$. Let $V_1, V_2 \in \text{Rep}_E(G)$.*

- (i) *If V_1 and V_2 are B -admissible, then so is $V_1 \otimes_E V_2$ and the natural map $D_B(V_1) \otimes_F D_B(V_2) \rightarrow D_B(V_1 \otimes_E V_2)$ is an isomorphism.*
- (ii) *Let $V \in \text{Rep}_E(G)$ be an extension of V_2 by V_1 . If V is B -admissible, then so are V_1 and V_2 . Moreover, the sequence $0 \rightarrow D_B(V_1) \rightarrow D_B(V) \rightarrow D_B(V_2) \rightarrow 0$ is exact.*
- (iii) *Let $h \in \mathbb{N}$ and let $V \in \text{Rep}_E(G)$ be B -admissible. Then $\bigwedge_E^h V$ is also B -admissible and the natural map $\bigwedge_F^h D_B(V) \rightarrow D_B(\bigwedge_E^h V)$ is an isomorphism.*

Proposition 4.6 ([Bri06, Proposition 3.7]). *Let B be an (E, G) -ring satisfying the following conditions:*

- (i) $F := B^G$ is Noetherian and B is faithfully flat over F ;
- (ii) $\alpha_B(V)$ is injective for every $V \in \text{Rep}_E(G)$;
- (iii) if $V \in \text{Rep}_E(G)$ is B -admissible and $\dim_E V = 1$, then the dual representation V^\vee is also B -admissible.

Then for every B -admissible representation $V \in \text{Rep}_E(G)$, the dual V^\vee is also B -admissible and the natural map

$$D_B(V^\vee) \rightarrow \text{Hom}_F(D_B(V), F)$$

is an isomorphism.

4.2. Horizontal crystalline, semistable, and de Rham representations. Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1. Recall that we set $\mathcal{G}_{R_K} := \text{Gal}(\overline{R_K}/R_K)$ and that $(\widehat{R}[p^{-1}], \widehat{R})$ is a perfectoid C -Banach pair.

Definition 4.7. Set

$$\begin{aligned} B_{\max}^\nabla(R) &:= \mathbb{B}_{\max}(\widehat{R}), & B_{\max, K}^\nabla(R) &:= \mathbb{B}_{\max, K}(\widehat{R}), \\ B_{\text{cris}}^\nabla(R) &:= \mathbb{B}_{\text{cris}}(\widehat{R}), \\ B_{\text{st}}^\nabla(R) &:= \mathbb{B}_{\text{st}}(\widehat{R}), & B_{\text{st}, K}^\nabla(R) &:= \mathbb{B}_{\text{st}, K}(\widehat{R}), \\ B_{\text{dR}}^\nabla(R) &:= \mathbb{B}_{\text{dR}}(\widehat{R}) = \mathbb{B}_{\text{dR}, K}(\widehat{R}), & \text{and } B_{\text{dR}}^{\nabla+}(R) &:= \mathbb{B}_{\text{dR}}^+(\widehat{R}). \end{aligned}$$

By functoriality of the construction, all these period rings have an action of \mathcal{G}_{R_K} . In particular, they are $(\mathbb{Q}_p, \mathcal{G}_{R_K})$ -rings. To simplify the notation, for $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, we write $D_{\max}^\nabla(V)$ for $D_{B_{\max}^\nabla(R)}(V)$ and $\alpha_{\max}^\nabla(V)$ for $\alpha_{B_{\max}^\nabla(R)}(V)$. We use similar conventions for other period rings.

Remark 4.8.

- (i) By definition, our period rings $B_{\max}^\nabla(R)$, $B_{\text{cris}}^\nabla(R)$, and $B_{\text{dR}}^\nabla(R)$ coincide with the horizontal period rings defined in [Bri08, Définitions 5.1.2, 6.1.2] and denoted by the same symbols.
- (ii) The period rings $B_{\max}^\nabla(R)$ and $B_{\text{cris}}^\nabla(R)$ admit the Frobenius φ . Similarly, $B_{\text{st}}^\nabla(R)$ admits the Frobenius φ and the monodromy operator N satisfying $N\varphi = p\varphi N$.
- (iii) Recall the following relations among period rings:

$$\begin{aligned} K \otimes_{K_0} B_{\max}^\nabla(R) &= B_{\max, K}^\nabla(R), & K \otimes_{K_0} B_{\text{st}}^\nabla(R) &= B_{\text{st}, K}^\nabla(R), \\ (B_{\text{st}}^\nabla(R))^{N=0} &= B_{\max}^\nabla(R), & \text{and } \varphi(B_{\max}^\nabla(R)) &\subset B_{\text{cris}}^\nabla(R) \subset B_{\max}^\nabla(R). \end{aligned}$$

- (iv) The action of \mathcal{G}_{R_K} on $B_{\max}^\nabla(R)$ and $B_{\text{cris}}^\nabla(R)$ commutes with the Frobenius. In particular, for $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, the modules $D_{\max}^\nabla(V)$ and $D_{\text{cris}}^\nabla(V)$ are equipped with the Frobenius φ by restricting $\varphi \otimes \text{id}$ on $B_{\max}^\nabla(R) \otimes_{\mathbb{Q}_p} V$ and $B_{\text{cris}}^\nabla(R) \otimes_{\mathbb{Q}_p} V$. The maps $\alpha_{\max}^\nabla(V)$ and $\alpha_{\text{cris}}^\nabla(V)$ are φ -equivariant. We also have

$$\varphi(D_{\max}^\nabla(V)) \subset D_{\text{cris}}^\nabla(V) \subset D_{\max}^\nabla(V).$$

- (v) The action of \mathcal{G}_{R_K} on $B_{\text{st}}^\nabla(R)$ commutes with the Frobenius and the monodromy operator. In particular, for $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, the module $D_{\text{st}}^\nabla(V)$ is equipped with the Frobenius φ and the monodromy operator N by restricting $\varphi \otimes \text{id}$ and $N \otimes \text{id}$ on $B_{\text{st}}^\nabla(R) \otimes_{\mathbb{Q}_p} V$. These endomorphisms satisfy

$$N\varphi = p\varphi N, \quad \text{and} \quad (D_{\text{st}}^\nabla(V))^{N=0} = D_{\max}^\nabla(V).$$

Proposition 4.9 ([Bri08, Corollaire 5.3.7]). *We have*

$$(B_{\text{dR}}^\nabla(R))^{\mathcal{G}_{R_K}} = K.$$

Corollary 4.10. *We have*

$$\begin{aligned} (B_{\text{max},K}^\nabla(R))^{\mathcal{G}_{R_K}} &= (B_{\text{st},K}^\nabla(R))^{\mathcal{G}_{R_K}} = K; \\ (B_{\text{cris}}^\nabla(R))^{\mathcal{G}_{R_K}} &= (B_{\text{max}}^\nabla(R))^{\mathcal{G}_{R_K}} = (B_{\text{st}}^\nabla(R))^{\mathcal{G}_{R_K}} = K_0. \end{aligned}$$

Proof. Since we have \mathcal{G}_{R_K} -equivariant K -algebra homomorphisms

$$K \otimes_{K_0} B_{\text{max}}^\nabla(R) \xrightarrow{\cong} B_{\text{max},K}^\nabla(R) \hookrightarrow B_{\text{dR}}^\nabla(R),$$

we obtain

$$(B_{\text{max},K}^\nabla(R))^{\mathcal{G}_{R_K}} = K, \quad \text{and} \quad (B_{\text{max}}^\nabla(R))^{\mathcal{G}_{R_K}} = K_0.$$

By the inclusions $\varphi(B_{\text{max}}^\nabla(R)) \subset B_{\text{cris}}^\nabla(R) \subset B_{\text{max}}^\nabla(R)$, we have $(B_{\text{cris}}^\nabla(R))^{\mathcal{G}_{R_K}} = K_0$. We can prove the remaining assertions in the same way. \square

The following is clear from what we have discussed:

Lemma 4.11. *Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$.*

- (i) $D_{\text{max}}^\nabla(V)$ and $D_{\text{st}}^\nabla(V)$ are K_0 -vector spaces, and $D_{\text{max},K}^\nabla(V)$, $D_{\text{st},K}^\nabla(V)$, and $D_{\text{dR}}^\nabla(V)$ are K -vector spaces.
- (ii) *We have*

$$K \otimes_{K_0} D_{\text{max}}^\nabla(V) \cong D_{\text{max},K}^\nabla(V), \quad \text{and} \quad K \otimes_{K_0} D_{\text{st}}^\nabla(V) \cong D_{\text{st},K}^\nabla(V).$$

Moreover, $\alpha_{\text{max}}^\nabla(V)$ is an isomorphism if and only if $\alpha_{\text{max},K}^\nabla(V)$ is an isomorphism. Similarly, $\alpha_{\text{st}}^\nabla(V)$ is an isomorphism if and only if $\alpha_{\text{st},K}^\nabla(V)$ is an isomorphism.

- (iii) *There are natural injective maps of K -vector spaces:*

$$D_{\text{max},K}^\nabla(V) \hookrightarrow D_{\text{st},K}^\nabla(V) \hookrightarrow D_{\text{dR}}^\nabla(V).$$

Lemma 4.12. *For every $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, the maps*

$$\alpha_{\text{cris}}^\nabla(V), \quad \alpha_{\text{max}}^\nabla(V), \quad \alpha_{\text{max},K}^\nabla(V), \quad \alpha_{\text{st}}^\nabla(V), \quad \alpha_{\text{st},K}^\nabla(V), \quad \alpha_{\text{dR}}^\nabla(V)$$

are all injective. Moreover, we have

$$\begin{aligned} \dim_{K_0} D_{\text{max}}^\nabla(V) &= \dim_K D_{\text{max},K}^\nabla(V) \leq \dim_{K_0} D_{\text{st}}^\nabla(V) = \dim_K D_{\text{st},K}^\nabla(V) \\ &\leq \dim_K D_{\text{dR}}^\nabla(V) \\ &\leq \dim_{\mathbb{Q}_p} V. \end{aligned}$$

Proof. The map $\alpha_{\text{dR}}^\nabla(V)$ is injective by [Bri08, Proposition 8.2.9]. Consider the commutative diagram

$$\begin{array}{ccc} K \otimes_{K_0} (B_{\text{max}}^\nabla(R) \otimes_{K_0} D_{\text{max}}^\nabla(V)) & \xrightarrow{K \otimes_{K_0} \alpha_{\text{max}}^\nabla(V)} & K \otimes_{K_0} (B_{\text{max}}^\nabla(R) \otimes_{\mathbb{Q}_p} V) \\ \parallel & & \parallel \\ B_{\text{max},K}^\nabla(R) \otimes_K D_{\text{max},K}^\nabla(V) & \xrightarrow{\alpha_{\text{max},K}^\nabla(V)} & B_{\text{max},K}^\nabla(R) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ B_{\text{dR}}^\nabla(R) \otimes_K D_{\text{dR}}^\nabla(V) & \xrightarrow{\alpha_{\text{dR}}^\nabla(V)} & B_{\text{dR}}^\nabla(R) \otimes_{\mathbb{Q}_p} V. \end{array}$$

The vertical equalities follow from Lemma 4.11(ii). Hence the injectivity of $\alpha_{\text{dR}}^\nabla(V)$ implies that of $\alpha_{\text{max}}^\nabla(V)$ and $\alpha_{\text{max},K}^\nabla(V)$. The same argument works for the injectivity of $\alpha_{\text{st}}^\nabla(V)$ and $\alpha_{\text{st},K}^\nabla(V)$. The injectivity of $\alpha_{\text{cris}}^\nabla(V)$ follows from that of $\alpha_{\text{max}}^\nabla(V)$ and the inclusion $B_{\text{cris}}^\nabla(R) \subset B_{\text{max}}^\nabla(R)$ by a similar argument. The second assertion follows from Lemma 4.11(iii). \square

Remark 4.13. The injectivity of $\alpha_{\text{cris}}^\nabla(V)$ is also proved in [Bri08, Proposition 8.2.9] under the additional assumption (BR) on R .

Lemma 4.14. *For $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, $\alpha_{\text{max}}^\nabla(V)$ is an isomorphism if and only if $\alpha_{\text{cris}}^\nabla(V)$ is an isomorphism. Similarly, $\alpha_{\text{st}}^\nabla(V)$ is an isomorphism if and only if $\alpha_{B_{\text{cris}}^\nabla(R)[\log \frac{[p^b]}{p}]}^\nabla(V)$ is an isomorphism.*

Proof. The sufficiency follows from Lemma 4.4. Since the proof of the necessity of the two assertions is similar, we deal with the necessity for the first. Assume that $\alpha_{\text{max}}^\nabla(V)$ is an isomorphism. Then the Frobenius $\varphi: B_{\text{max}}^\nabla(R) \rightarrow B_{\text{max}}^\nabla(V)$ induces the commutative diagram

$$\begin{array}{ccc} B_{\text{max}}^\nabla(R) \otimes_{K_0} D_{\text{max}}^\nabla(V) & \xrightarrow[\cong]{\alpha_{\text{max}}^\nabla(V)} & B_{\text{max}}^\nabla(R) \otimes_{\mathbb{Q}_p} V \\ \varphi \otimes \varphi \downarrow & & \downarrow \varphi \otimes \text{id} \\ B_{\text{max}}^\nabla(R) \otimes_{K_0} D_{\text{max}}^\nabla(V) & \xrightarrow[\cong]{\alpha_{\text{max}}^\nabla(V)} & B_{\text{max}}^\nabla(R) \otimes_{\mathbb{Q}_p} V. \end{array}$$

Since $\varphi: B_{\text{max}}^\nabla(R) \rightarrow B_{\text{max}}^\nabla(V)$ factors through $B_{\text{cris}}^\nabla(R) \subset B_{\text{max}}^\nabla(R)$, the above diagram also factors as

$$\begin{array}{ccc} B_{\text{max}}^\nabla(R) \otimes_{K_0} D_{\text{max}}^\nabla(V) & \xrightarrow[\cong]{\alpha_{\text{max}}^\nabla(V)} & B_{\text{max}}^\nabla(R) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ B_{\text{cris}}^\nabla(R) \otimes_{K_0} D_{\text{cris}}^\nabla(V) & \xrightarrow[\subset]{\alpha_{\text{cris}}^\nabla(V)} & B_{\text{cris}}^\nabla(R) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ B_{\text{max}}^\nabla(R) \otimes_{K_0} D_{\text{max}}^\nabla(V) & \xrightarrow[\cong]{\alpha_{\text{max}}^\nabla(V)} & B_{\text{max}}^\nabla(R) \otimes_{\mathbb{Q}_p} V. \end{array}$$

In particular, the image of $\alpha_{\text{cris}}^\nabla(V)$ contains V . Since $\alpha_{\text{cris}}^\nabla(V)$ is injective and $B_{\text{cris}}^\nabla(R)$ -linear, it is an isomorphism. \square

Lemma 4.15. *For $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, $\alpha_{\text{max}}^\nabla(V)$ is an isomorphism if and only if $\alpha_{\text{st}}^\nabla(V)$ is an isomorphism and the monodromy operator N on $D_{\text{st}}^\nabla(V)$ is zero.*

Proof. The necessity follows from Lemma 4.4 and $(D_{\text{st}}^\nabla(V))^{N=0} = D_{\text{max}}^\nabla(V)$. We prove the sufficiency. Note that the multiplication map

$$B_{\text{st}}^\nabla(R) \otimes_{K_0} B_{\text{st}}^\nabla(R) \rightarrow B_{\text{st}}^\nabla(R)$$

is compatible with $N \otimes \text{id} + \text{id} \otimes N$ on the source and N on the target. Since $\alpha_{\text{st}}^\nabla(V)$ is an isomorphism, this yields the following diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & B_{\max}^\nabla \otimes_{\mathbb{Q}_p} V & \longrightarrow & B_{\text{st}}^\nabla \otimes_{\mathbb{Q}_p} V & \xrightarrow{N \otimes \text{id}} & B_{\text{st}}^\nabla \otimes_{\mathbb{Q}_p} V \longrightarrow 0 \\ & & \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\ 0 & \longrightarrow & \text{Ker}(f) & \longrightarrow & B_{\text{st}}^\nabla \otimes_{K_0} D_{\text{st}}^\nabla(V) & \xrightarrow{f} & B_{\text{st}}^\nabla \otimes_{K_0} D_{\text{st}}^\nabla(V) \longrightarrow 0, \end{array}$$

where $B_{\max}^\nabla = B_{\max}^\nabla(R)$, $B_{\text{st}}^\nabla = B_{\text{st}}^\nabla(R)$, and f denotes $N \otimes \text{id} + \text{id} \otimes N$. Since the monodromy operator N on $D_{\text{st}}^\nabla(V)$ is assumed to be zero, we conclude $\text{Ker } f = B_{\max}^\nabla(R) \otimes_{K_0} D_{\max}^\nabla(V)$ and the left vertical isomorphism is $\alpha_{\max}^\nabla(V)$. \square

Definition 4.16. Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$.

- (i) We say that V is *horizontal crystalline* if $\alpha_{\max}^\nabla(V)$ is an isomorphism.
- (ii) We say that V is *horizontal semistable* if $\alpha_{\text{st}}^\nabla(V)$ is an isomorphism.
- (iii) We say that V is *horizontal de Rham* if $\alpha_{\text{dR}}^\nabla(V)$ is an isomorphism.

See Lemma 4.11(ii), 4.14, and 4.15 for equivalent definitions. By Lemma 4.4, a horizontal crystalline representation is horizontal semistable, and a horizontal semistable representation is horizontal de Rham.

Remark 4.17. Our definition of horizontal de Rham representations coincides with that of [Bri08, p. 117]. By Lemma 4.14, our definition of horizontal crystalline representations coincides with the definition of horizontal R_0 -crystalline representations in [Bri08, p. 117].

Remark 4.18. Observe that the period rings $B_{\max}^\nabla(R)$, $B_{\text{st}}^\nabla(R)$, and $B_{\text{dR}}^\nabla(R)$ all satisfy the assumptions of Proposition 4.5. Hence the full subcategory of horizontal crystalline (resp. semistable, resp. de Rham) representations of \mathcal{G}_{R_K} is stable under taking tensor products, subquotients, and exterior products.

Lemma 4.19. *The full subcategory of horizontal de Rham representations of \mathcal{G}_{R_K} is stable under taking duals. If R satisfies Condition (BR), then the full subcategory of horizontal crystalline (resp. semistable) representations of \mathcal{G}_{R_K} is stable under taking duals.*

Proof. By Proposition 4.6, it suffices to prove the assertion for one-dimensional representations. For horizontal de Rham representations (resp. horizontal crystalline representations under Condition (BR)), it is [Bri08, Théorème 8.4.2]. In general, if $D_{\text{st}}^\nabla(V)$ is one-dimensional over K_0 for $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, then $N = 0$ by the relation $N\varphi = p\varphi N$. In particular, every one-dimensional horizontal semistable representation is horizontal crystalline, and thus so is its dual. \square

Proposition 4.20. *The sequence*

$$0 \rightarrow \mathbb{Q}_p \rightarrow B_{\max}^\nabla(R)^{\varphi=1} \rightarrow B_{\text{dR}}^\nabla(R)/B_{\text{dR}}^{\nabla,+}(R) \rightarrow 0$$

is exact.

Proof. Since $\varphi(B_{\max}^\nabla(R)) \subset B_{\text{cris}}^\nabla(R) \subset B_{\max}^\nabla(R)$, we get

$$B_{\text{cris}}^\nabla(R)^{\varphi=1} = B_{\max}^\nabla(R)^{\varphi=1}.$$

Hence the assertion is [Bri08, Proposition 6.2.24]. \square

The exact sequence in the proposition is called *the fundamental exact sequence*.

4.3. Functoriality. Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1 and fix an algebraic closure $\overline{\text{Frac } R}$ of $\text{Frac } R$ as before. Let us discuss functorial aspects of our formalism in two special cases.

Let R' be another \mathcal{O}_R -algebra satisfying the conditions in Set-up 3.1 and let $f: R \rightarrow R'$ be an \mathcal{O}_K -algebra homomorphism. Recall

$$\mathcal{G}_{R_K} = \text{Gal}(\overline{R}[p^{-1}]/R_K) = \pi_1^{\text{ét}}(\text{Spec } R_K, \overline{\text{Frac } R}).$$

Note that $R_K \rightarrow R'_K \rightarrow \overline{\text{Frac } R'}$ defines another geometric point of $\text{Spec } R_K$. Choose a path on $\text{Spec } R_K$ from $\overline{\text{Frac } R'}$ to $\overline{\text{Frac } R}$ (cf. [SGA1, Exposé V, 7]). Concretely, this is equivalent to choosing an \mathcal{O}_K -algebra homomorphism $\overline{f}: \overline{R} \rightarrow \overline{R'}$ extending f . The path defines a group homomorphism

$$\begin{aligned} \mathcal{G}_{R'_K} &= \pi_1^{\text{ét}}(\text{Spec } R'_K, \overline{\text{Frac } R'}) \xrightarrow{\text{Spec } f^*} \pi_1^{\text{ét}}(\text{Spec } R_K, \overline{\text{Frac } R'}) \\ &\longrightarrow \pi_1^{\text{ét}}(\text{Spec } R_K, \overline{\text{Frac } R}) = \mathcal{G}_{R_K}. \end{aligned}$$

In what follows (e.g. Proposition 4.23), we always choose a path so that \overline{f} is an $\mathcal{O}_{\overline{K}}$ -algebra homomorphism. Then \overline{f} gives rise to a homomorphism $(\widehat{R}[p^{-1}], \widehat{R}) \rightarrow (\widehat{R'}[p^{-1}], \widehat{R'})$ of C -Banach pairs and induces homomorphisms

$$\text{B}_{\max}^{\nabla}(R) \rightarrow \text{B}_{\max}^{\nabla}(R'), \quad \text{B}_{\text{st}}^{\nabla}(R) \rightarrow \text{B}_{\text{st}}^{\nabla}(R'), \quad \text{and} \quad \text{B}_{\text{dR}}^{\nabla}(R) \rightarrow \text{B}_{\text{dR}}^{\nabla}(R').$$

Moreover they are compatible with $\mathcal{G}_{R'_K} \rightarrow \mathcal{G}_{R_K}$.

Lemma 4.21. *If a \mathbb{Q}_p -representation V of \mathcal{G}_{R_K} is horizontal crystalline (resp. semistable, resp. de Rham), then so is the representation $V|_{\mathcal{G}_{R'_K}}$ of $\mathcal{G}_{R'_K}$.*

Proof. This follows from Lemma 4.4. \square

We also consider the case of changing the base field. Let L be a finite extension of K , and set

$$R_{\mathcal{O}_L} := R \otimes_{\mathcal{O}_K} \mathcal{O}_L \quad \text{and} \quad R_L := R_{\mathcal{O}_L}[p^{-1}] = R_K \otimes_K L.$$

Observe that $R_{\mathcal{O}_L}$ satisfies the conditions in Set-up 3.1 with respect to L . Since $\overline{\text{Frac } R}$ is an algebraic closure of $\text{Frac } R_{\mathcal{O}_L}$, we have $\overline{R_{\mathcal{O}_L}} = \overline{R}$. It follows \mathcal{G}_{R_L} is an open subgroup of \mathcal{G}_{R_K} of finite index.

Lemma 4.22. *Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$.*

- (i) *If V is horizontal crystalline, then the representation $V|_{\mathcal{G}_{R_L}}$ of \mathcal{G}_{R_L} is also horizontal crystalline. Similarly, if V is horizontal semistable, then so is $V|_{\mathcal{G}_{R_L}}$.*
- (ii) *V is horizontal de Rham if and only if $V|_{\mathcal{G}_{R_L}}$ is horizontal de Rham.*

Proof. Part (i) and the necessity of Part (ii) follow from Lemma 4.4. Assume that $V|_{\mathcal{G}_{R_L}}$ is horizontal de Rham. Using the necessity, we may reduce to the case where L is Galois over K . Then \mathcal{G}_{R_L} is a normal subgroup of \mathcal{G}_{R_K} with quotient $\text{Gal}(L/K)$. Since $\text{B}_{\text{dR}}^{\nabla}(R) = \text{B}_{\text{dR}}^{\nabla}(R_{\mathcal{O}_L})$, the finite-dimensional L -vector space $D_{\text{dR}}^{\nabla}(V|_{\mathcal{G}_{R_{\mathcal{O}_L}}})$ has a semilinear $\text{Gal}(L/K)$ -action and

$$D_{\text{dR}}^{\nabla}(V) = D_{\text{dR}}^{\nabla}(V|_{\mathcal{G}_{R_L}})^{\text{Gal}(L/K)}.$$

By Galois descent, we have

$$L \otimes_K D_{\text{dR}}^{\nabla}(V) \xrightarrow{\cong} D_{\text{dR}}^{\nabla}(V|_{\mathcal{G}_{R_L}}).$$

It follows that $\alpha_{\text{dR}}^\nabla(V) = \alpha_{\text{dR}}^\nabla(V|_{\mathcal{G}_{R_L}})$ and it is an isomorphism by assumption. \square

By combining these two cases, we can consider the following setting:

Proposition 4.23. *Let L be a finite extension of K and R' an \mathcal{O}_L -algebra satisfying the conditions in Set-up 3.1 with respect to L . Let $f: R \rightarrow R'$ be an \mathcal{O}_K -algebra homomorphism. Fix a path on $\text{Spec } R_K$ from $\overline{\text{Frac } R'}$ to $\overline{\text{Frac } R}$ and consider the associated homomorphism $\mathcal{G}_{R'_L} \rightarrow \mathcal{G}_{R_K}$. Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$ and set $f^*V := V|_{\mathcal{G}_{R'_L}}$. If V is horizontal semistable and f^*V is horizontal crystalline, then V is horizontal crystalline.*

Proof. Denote the residue field of L by k_L and set $L_0 = W(k_L)[p^{-1}]$. By Lemma 4.4, we have $D_{\text{st}}^\nabla(f^*V) \cong L_0 \otimes_{K_0} D_{\text{st}}^\nabla(V)$. Moreover, it follows from the construction that this isomorphism is compatible with the monodromy operator N on the left hand side and $\text{id} \otimes N$ on the right hand side. Since f^*V is horizontal crystalline, we have $N = 0$ on $D_{\text{st}}^\nabla(f^*V)$. By faithful flatness of $K_0 \rightarrow L_0$, we also have $N = 0$ on $D_{\text{st}}^\nabla(V)$. Hence V is horizontal crystalline by Lemma 4.15. \square

Remark 4.24. We will use the proposition when $R' = \mathcal{O}_L$. In this case, it is informally phrased as follows: if a horizontal semistable representation is crystalline at one classical point, then it is horizontal crystalline.

4.4. Horizontal de Rham representations and de Rham representations.

In this subsection, we review the de Rham period ring studied in [Bri08]. This ring will be used in the next section to define the V_{st} -functor. Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1. Set

$$\widehat{\Omega}_R := \varprojlim_m \Omega_{R/\mathbb{Z}}^1 / p^m \Omega_{R/\mathbb{Z}}^1.$$

Definition 4.25. Let $B_{\text{dR}}(R)$ denote the de Rham period ring of R defined in [Bri08, Définition 5.1.5]. We briefly recall the definition below.

The map $\theta_{\mathbb{A}_{\text{inf}}(\widehat{R})} : \mathbb{A}_{\text{inf}}(\widehat{R}) \rightarrow \widehat{R}$ extends R -linearly to the map

$$\theta_R : R \otimes_{\mathbb{Z}} \mathbb{A}_{\text{inf}}(\widehat{R}) \rightarrow \widehat{R}.$$

Let $\mathbb{A}_{\text{inf}}(\widehat{R}/R)$ denote the completion of the ring $R \otimes_{\mathbb{Z}} \mathbb{A}_{\text{inf}}(\widehat{R})$ with respect to the topology defined by the ideal $\theta_R^{-1}(p\widehat{R})$ ([Bri08, Définition 5.1.3]). The map θ_R naturally extends to a map $\mathbb{A}_{\text{inf}}(\widehat{R}/R)[p^{-1}] \rightarrow \widehat{R}[p^{-1}]$, which we still denote by θ_R . We set

$$B_{\text{dR}}^+(R) := \varprojlim_m \mathbb{A}_{\text{inf}}(\widehat{R}/R)[p^{-1}] / (\text{Ker } \theta_R)^m,$$

and denote the natural map $B_{\text{dR}}^+(R) \rightarrow \widehat{R}[p^{-1}]$ by θ_R . For $m \in \mathbb{N}$, we set $\text{Fil}^m B_{\text{dR}}^+(R) := (\text{Ker } \theta_R)^m$. This is a decreasing separated and exhaustive filtration on $B_{\text{dR}}^+(R)$. We define

$$B_{\text{dR}}(R) := B_{\text{dR}}^+(R)[t^{-1}].$$

The ring has an action of \mathcal{G}_{R_K} , a decreasing separated and exhaustive filtration Fil^\bullet given by

$$\text{Fil}^0 B_{\text{dR}}(R) = \sum_{i=0}^{\infty} t^{-i} \text{Fil}^i B_{\text{dR}}^+(R), \text{ and } \text{Fil}^m B_{\text{dR}}(R) = t^m \text{Fil}^0 B_{\text{dR}}(R) \quad (m \in \mathbb{Z}),$$

and a $B_{\text{dR}}^{\nabla}(R)$ -linear integrable connection

$$\nabla: B_{\text{dR}}(R) \rightarrow B_{\text{dR}}(R) \otimes_R \widehat{\Omega}_R$$

satisfying the Griffiths transversality

$$\nabla(\text{Fil}^m B_{\text{dR}}(R)) \subset \text{Fil}^{m-1} B_{\text{dR}}(R) \otimes_R \widehat{\Omega}_R.$$

We also have $(B_{\text{dR}}(R))^{\mathcal{G}_{R_K}} = R_K$ and $B_{\text{dR}}^{\nabla}(R) = B_{\text{dR}}(R)^{\nabla=0}$ ([Bri08, Propositions 5.2.12, 5.3.3]). The ring $B_{\text{dR}}(R)$ is a $(\mathbb{Q}_p, \mathcal{G}_{R_K})$ -ring.

Definition 4.26. For $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, we write $D_{\text{dR}}(V)$ for $D_{B_{\text{dR}}(R)}(V)$ and $\alpha_{\text{dR}}(V)$ for $\alpha_{B_{\text{dR}}(R)}(V)$. We say that V is *de Rham* if it is $B_{\text{dR}}(R)$ -admissible, i.e., $\alpha_{\text{dR}}(V)$ is an isomorphism of $B_{\text{dR}}(R)$ -modules. Note that $\alpha_{\text{dR}}(V)$ is always injective by [Bri08, Proposition 8.2.4].

The R_K -module $D_{\text{dR}}(V)$ is projective of rank at most $\dim_{\mathbb{Q}_p} V$ by [Bri08, Proposition 8.3.1], and it is equipped with induced connection

$$\nabla: D_{\text{dR}}(V) \rightarrow D_{\text{dR}}(V) \otimes_R \widehat{\Omega}_R,$$

and filtration $\text{Fil}^{\bullet} D_{\text{dR}}(V)$. We also have a natural R_K -linear map

$$\beta_{\text{dR}}(V): R_K \otimes_K D_{\text{dR}}^{\nabla}(V) \rightarrow D_{\text{dR}}(V).$$

Lemma 4.27 ([Bri08, Proposition 8.2.1]). *For $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, it is horizontal de Rham if and only if it is de Rham and $\beta_{\text{dR}}(V)$ is an isomorphism.*

Lemma 4.28 ([Bri08, Proposition 8.4.3]). *Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$. If V is de Rham, then the isomorphism*

$$\alpha_{\text{dR}}(V): B_{\text{dR}}(R) \otimes_{R_K} D_{\text{dR}}(V) \xrightarrow{\cong} B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V$$

is compatible with \mathcal{G}_{R_K} -actions and connections, and strictly compatible with filtrations.

Lemma 4.29 ([Bri08, Corollaire 5.2.11]). *For $m \in \mathbb{Z}$, we have*

$$B_{\text{dR}}^{\nabla}(R) \cap \text{Fil}^m B_{\text{dR}}(R) = t^m B_{\text{dR}}^{\nabla+}(R).$$

Remark 4.30. Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$ be a horizontal de Rham representation. The filtration $\text{Fil}^m B_{\text{dR}}^{\nabla}(R) := t^m B_{\text{dR}}^{\nabla+}(R)$ on $B_{\text{dR}}^{\nabla}(R)$ induces a filtration on $D_{\text{dR}}^{\nabla}(V)$. The maps $\alpha_{\text{dR}}^{\nabla}(V)$ and $\beta_{\text{dR}}(V)$ preserve filtrations, i.e.,

$$\begin{aligned} \alpha_{\text{dR}}^{\nabla}(V)(\text{Fil}^m(B_{\text{dR}}^{\nabla}(R) \otimes_K D_{\text{dR}}^{\nabla}(V))) &\subset \text{Fil}^m B_{\text{dR}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V, \\ \beta_{\text{dR}}(V)(\text{Fil}^m(R_K \otimes_K D_{\text{dR}}^{\nabla}(V))) &\subset \text{Fil}^m D_{\text{dR}}(V). \end{aligned}$$

However, neither $\alpha_{\text{dR}}^{\nabla}(V)$ nor $\beta_{\text{dR}}(V)$ is strictly compatible with filtrations in general⁷. See the example below.

Example 4.31. We will give an example of a horizontal crystalline representation (hence horizontal de Rham) such that $\alpha_{\text{dR}}^{\nabla}(V)$ is not strictly compatible with filtrations.

⁷[Bri08, Proposition 8.4.3] claims that $\alpha_{\text{dR}}^{\nabla}(V)$ is strictly compatible with filtrations, but a proof is not given.

Assume $K = K_0$ for simplicity. Let $R = \mathcal{O}_K \langle T^\pm \rangle$ and let $h(T) = 1 + pT \in R$. Fix a nontrivial compatible system $(1, \varepsilon_1, \dots)$ of p -power roots of 1 ($\varepsilon_1 \neq 1$) and a compatible system $(h, h^{1/p}, \dots)$ of p -power roots of h . They define elements of $\widehat{\widehat{R}}^b$:

$$\varepsilon := (1 \bmod p, \varepsilon_1 \bmod p, \dots), \quad h^b := (h \bmod p, h^{1/p} \bmod p, \dots).$$

Consider the two-dimensional \mathbb{Q}_p -representation V of \mathcal{G}_{R_K} corresponding to (ε, h^b) . Namely, set $V = \mathbb{Q}_p e_1 + \mathbb{Q}_p e_2$ and define the action of $g \in \mathcal{G}_{R_K}$ on V by

$$ge_1 = \chi(g)e_1, \quad ge_2 = \eta(g)e_1 + e_2,$$

where $\chi: \mathcal{G}_{R_K} \rightarrow \text{Gal}(\overline{K}/K) \rightarrow \mathbb{Z}_p^\times$ is the p -adic cyclotomic character defined by $g(\varepsilon_m) = \varepsilon_m^{\chi(g)}$, and $\eta: \mathcal{G}_{R_K} \rightarrow \mathbb{Z}_p$ is a map defined by $g(h^{1/p^m}) = \varepsilon_m^{\eta(g)} h^{1/p^m}$.

We will show that V is a horizontal crystalline representation.

Claim. *The series*

$$\log[h^b] := \sum_{m=1}^{\infty} (-1)^{m-1} \frac{([h^b] - 1)^m}{m}$$

converges in $A_{\max}^\nabla(R) := A_{\max}(\widehat{\widehat{R}})$ and satisfies

$$g(\log[h^b]) = \eta(g)t + \log[h^b], \quad \varphi(\log[h^b]) = p \log[h^b].$$

Moreover, we have $\log[h^b] \in \text{Fil}^0 B_{\text{dR}}^\nabla(R) \setminus \text{Fil}^1 B_{\text{dR}}^\nabla(R)$.

Proof. Observe $\theta_{\mathbb{A}_{\text{inf}}(\widehat{\widehat{R}})}([h^b] - 1) = h - 1 = pT \in p\widehat{\widehat{R}}$. Hence we can write $[h^b] - 1 = x + py$ for some $x \in \text{Ker } \theta_{\mathbb{A}_{\text{inf}}(\widehat{\widehat{R}})}$ and $y \in \mathbb{A}_{\text{inf}}(\widehat{\widehat{R}})$. It follows

$$(-1)^{m-1} \frac{([h^b] - 1)^m}{m} = (-1)^{m-1} \frac{p^m}{m} \left(\frac{[h^b] - 1}{p} \right)^m = (-1)^{m-1} \frac{p^m}{m} \left(\frac{x}{p} + y \right)^m.$$

Since $\frac{p^m}{m} \in \mathbb{Z}_p$ and it converges to zero as $m \rightarrow \infty$, we conclude that the series $\sum_{m=1}^{\infty} (-1)^{m-1} \frac{([h^b] - 1)^m}{m}$ converges in $A_{\max}^\nabla(R)$.

We can easily check the remaining statements (cf. [Fon94a, 3.1.3] and [BC09, Lemma 9.2.2]). For example, since $\theta_{A_{\max}^\nabla(R)}: A_{\max}^\nabla(R) \rightarrow \widehat{\widehat{R}}$ is continuous with respect to the p -adic topology, we have

$$\theta(\log[h^b]) = \sum_{m=1}^{\infty} (-1)^{m-1} \frac{(h - 1)^m}{m} = \sum_{m=1}^{\infty} (-1)^{m-1} \frac{(pT)^m}{m} \neq 0.$$

Hence $\log[h^b] \in \text{Fil}^0 B_{\text{dR}}^\nabla(R) \setminus \text{Fil}^1 B_{\text{dR}}^\nabla(R)$. \square

Set

$$f_1 = \frac{1}{t}e_1, \quad f_2 = e_2 - \frac{\log[h^b]}{t}e_1 \in B_{\max}^\nabla(R) \otimes_{\mathbb{Q}_p} V.$$

Then we can check that f_1 and f_2 are fixed by \mathcal{G}_{R_K} and generate $B_{\max}^\nabla(R) \otimes_{\mathbb{Q}_p} V$ as a $B_{\max}^\nabla(R)$ -module. This implies that V is horizontal crystalline with $D_{\max}^\nabla(V) = Kf_1 + Kf_2$. We also see that $\varphi(f_1) = \frac{1}{p}f_1$ and $\varphi(f_2) = f_2$. Note that $D_{\text{dR}}^\nabla(V) = D_{\max}^\nabla(V)$ in this case and if we equip $D_{\text{dR}}^\nabla(V)$ with a decreasing filtration induced from $\text{Fil}^m(B_{\text{dR}}^\nabla(R) \otimes_{\mathbb{Q}_p} V) := t^m B_{\text{dR}}^{\nabla+}(R) \otimes_{\mathbb{Q}_p} V$, we have

$$\text{Fil}^{-1} D_{\text{dR}}^\nabla(V) = D_{\text{dR}}^\nabla(V), \quad \text{Fil}^0 D_{\text{dR}}^\nabla(V) = 0.$$

In particular, the $B_{\text{dR}}^{\nabla}(R)$ -linear isomorphism

$$\alpha_{\text{dR}}^{\nabla}(V): B_{\text{dR}}^{\nabla}(R) \otimes_K D_{\text{dR}}^{\nabla}(V) \xrightarrow{\cong} B_{\text{dR}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V$$

is *not* strictly compatible with filtrations. Indeed, we have $e_2 = 1 \otimes f_2 + \log[h^b] \otimes f_1 \notin \text{Fil}^0(B_{\text{dR}}^{\nabla}(R) \otimes_K D_{\text{dR}}^{\nabla}(V))$ while $e_2 \in \text{Fil}^0(B_{\text{dR}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V)$.

Let now turn to $D_{\text{dR}}(V) = R_K f_1 + R_K f_2$. We will give another R_K -basis. Note that h is invertible in R and $[h^b]/h - 1 \in \text{Ker } \theta_R$. It follows that the series

$$\log \frac{[h^b]}{h} := \sum_{m=1}^{\infty} (-1)^{m-1} \frac{([h^b]/h - 1)^m}{m}$$

converges in $B_{\text{dR}}^+(R)$. Let

$$\log h := \sum_{m=1}^{\infty} (-1)^{m-1} \frac{(h-1)^m}{m} = \sum_{m=1}^{\infty} (-1)^{m-1} \frac{p^m}{m} T^m \in R.$$

We can check

$$\log[h^b] = \log \frac{[h^b]}{h} + \log h \in B_{\text{dR}}^+(R).$$

Set

$$f_3 = e_2 - \frac{\log[h^b]/h}{t} e_1 = f_2 + \frac{\log h}{t} e_1 \in B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V.$$

Then f_3 is fixed by \mathcal{G}_{R_K} , and $D_{\text{dR}}(V) = R_K f_1 + R_K f_3$. Consider the filtration on $D_{\text{dR}}(V)$ induced from $\text{Fil}^m(B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V) := (\text{Fil}^m B_{\text{dR}}(R)) \otimes_{\mathbb{Q}_p} V$. Since $\log \frac{[h^b]}{h} \in \text{Fil}^1 B_{\text{dR}}(R)$, we see

$$\text{Fil}^{-1} D_{\text{dR}}(V) = D_{\text{dR}}(V), \quad \text{Fil}^0 D_{\text{dR}}(V) = R_K f_3, \quad \text{Fil}^1 D_{\text{dR}}(V) = 0.$$

It follows that the natural R_K -linear isomorphism

$$\beta_{\text{dR}}(V): R_K \otimes_K D_{\text{dR}}^{\nabla}(V) \xrightarrow{\cong} D_{\text{dR}}(V)$$

is *not* strictly compatible with filtrations. Note that the $B_{\text{dR}}(R)$ -linear isomorphism

$$\alpha_{\text{dR}}(V): B_{\text{dR}}(R) \otimes_{R_K} D_{\text{dR}}(V) \xrightarrow{\cong} B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V$$

is strictly compatible with filtrations.

5. FILTERED $(\varphi, N, \text{Gal}(L/K), R_K)$ -MODULES

In this section, we follow [Fon94b] and define filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -modules and the functors D_{pst} and V_{pst} .

As before, let k be a perfect field of characteristic p and set $K_0 := W(k)[p^{-1}]$. Let K be a totally ramified finite extension of K_0 . We denote the maximal unramified extension of K_0 inside \overline{K} by K_0^{ur} . Let L be a Galois extension of K inside \overline{K} . Let L_0 be the maximal unramified extension of K_0 in L and σ the absolute Frobenius on L_0 . Note $(\overline{K})_0 = K_0^{\text{ur}}$.

Let us first recall the definition of $(\varphi, N, \text{Gal}(L/K))$ -modules:

Definition 5.1 ([Fon94b, 4.2.1]). A $(\varphi, N, \text{Gal}(L/K))$ -module is a finite-dimensional L_0 -vector space D equipped with

- (i) an injective σ -semilinear map $\varphi: D \rightarrow D$ (*Frobenius*),
- (ii) an L_0 -linear endomorphism $N: D \rightarrow D$ (*monodromy operator*), and
- (iii) a semilinear action of $\text{Gal}(L/K)$

such that they satisfy the following compatibilities:

- $N\varphi = p\varphi N$;
- for every $g \in \text{Gal}(L/K)$, $g\varphi = \varphi g$, and $gN = Ng$.

When $L = K$, we simply call it a (φ, N) -module.

A morphism of $(\varphi, N, \text{Gal}(L/K))$ -modules is an L_0 -linear map that commutes with φ, N , and $\text{Gal}(L/K)$ -action.

We say that a $(\varphi, N, \text{Gal}(L/K))$ -module D is *discrete* if the action of $\text{Gal}(L/K)$ on D is discrete, i.e., for every $d \in D$, the stabilizer subgroup of d is open in $\text{Gal}(L/K)$.

Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1. We set

$$R_{\mathcal{O}_L} := \mathcal{O}_L \otimes_{\mathcal{O}_K} R, \quad \text{and} \quad R_L := L \otimes_K R_K.$$

Note that $R_{\mathcal{O}_L}$ is p -adically complete if L is finite over K but that it is not the case in general. For an L_0 -vector space D , we write D_{R_L} for $R_L \otimes_{L_0} D$.

Definition 5.2 (cf. [Fon94b, 4.3.2]). A *filtered* $(\varphi, N, \text{Gal}(L/K), R_K)$ -module is a pair consisting of

- a $(\varphi, N, \text{Gal}(L/K))$ -module D , and
- a separated and exhaustive decreasing filtration $(\text{Fil}^i D_{R_L})_{i \in \mathbb{Z}}$ of D_{R_L}

such that each $\text{Fil}^i D_{R_L}$ is a $\text{Gal}(L/K)$ -stable projective R_L -submodule of D_{R_L} . When $L = K$, we simply call it a (φ, N, R_K) -module.

A morphism of filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -modules is defined to be a morphism of $(\varphi, N, \text{Gal}(L/K))$ -modules whose scalar extension from L_0 to R_L preserves filtrations. A filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -module is called *discrete* if it is discrete as a $(\varphi, N, \text{Gal}(L/K))$ -module.

We denote the full subcategory of discrete filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -modules by $\text{MF}_{L/K}(\varphi, N, R_K)$.

Proposition 5.3. *Let L be a finite Galois extension of K and let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$. If $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable, then $D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}})$ is a $(\varphi, N, \text{Gal}(L/K))$ -module. Moreover, if we equip $D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}})$ with the filtration induced from $D_{\text{dR}}(V|_{\mathcal{G}_{R_L}})$ via the isomorphism*

$$R_L \otimes_{L_0} D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}}) \xrightarrow{\cong} D_{\text{dR}}(V|_{\mathcal{G}_{R_L}}),$$

then $D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}})$ is a filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -module.

Proof. Recall that $B_{\text{st}}^{\nabla}(R_{\mathcal{O}_L}) \otimes_{\mathbb{Q}_p} V$ admits an injective σ -semilinear map $\varphi \otimes \text{id}$ and an L_0 -linear endomorphism $N \otimes \text{id}$ satisfying $(N \otimes \text{id})(\varphi \otimes \text{id}) = p(\varphi \otimes \text{id})(N \otimes \text{id})$. Moreover, by the identification $B_{\text{st}}^{\nabla}(R_{\mathcal{O}_L}) \otimes_{\mathbb{Q}_p} V = B_{\text{st}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V$, it is equipped with diagonal semilinear \mathcal{G}_{R_K} -action commuting with $\varphi \otimes \text{id}$ and $N \otimes \text{id}$. The first assertion follows from these remarks and $\mathcal{G}_{R_K}/\mathcal{G}_{R_L} \cong \text{Gal}(L/K)$. The second assertion follows from the first and [Bri08, Proposition 8.3.4(a)]. \square

Definition 5.4. Assume that L is finite over K . Recall the natural identifications $B_{\text{st}}^{\nabla}(R_{\mathcal{O}_L}) = B_{\text{st}}^{\nabla}(R)$ and $B_{\text{dR}}^{\nabla}(R_{\mathcal{O}_L}) = B_{\text{dR}}^{\nabla}(R)$. For a filtered $(\varphi, N, \text{Gal}(L/K), R_K)$ -module D , set

$$V_{\text{st},L}(D) := (B_{\text{st}}^{\nabla}(R) \otimes_{L_0} D)^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R) \otimes_{R_L} D_{R_L}),$$

where φ (resp. N) on $B_{\text{st}}^{\nabla}(R) \otimes_{L_0} D$ denotes $\varphi \otimes \varphi$ (resp. $N \otimes \text{id} + \text{id} \otimes N$) and Fil^{\bullet} on $B_{\text{dR}}(R) \otimes_{R_L} D_{R_L}$ denotes $\sum_{m \in \mathbb{Z}} \text{Fil}^m B_{\text{dR}}(R) \otimes_{R_L} \text{Fil}^{\bullet-m} D_{R_L}$. It is a \mathbb{Q}_p -vector space equipped with a \mathbb{Q}_p -linear action of \mathcal{G}_{R_K} .

Proposition 5.5. *Let L be a finite Galois extension of K and let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$. If $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable, then the natural \mathbb{Q}_p -linear map*

$$V \rightarrow V_{\text{st},L}(D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}}))$$

is a \mathcal{G}_{R_K} -equivariant isomorphism.

Proof. We use the identification $B_{\text{st}}^{\nabla}(R_{\mathcal{O}_L}) = B_{\text{st}}^{\nabla}(R)$. For simplicity, we write V_L for $V|_{\mathcal{G}_{R_L}}$. Since V_L is horizontal semistable, we have the commutative diagram

$$\begin{array}{ccc} B_{\text{st}}^{\nabla}(R) \otimes_{L_0} D_{\text{st}}^{\nabla}(V_L) & \xrightarrow[\cong]{\alpha_{\text{st}}^{\nabla}(V_L)} & B_{\text{st}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ B_{\text{dR}}(R) \otimes_{R_L} D_{\text{dR}}(V_L) & \xrightarrow[\cong]{\alpha_{\text{dR}}(V_L)} & B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V. \end{array}$$

Furthermore, the isomorphism $\alpha_{\text{st}}^{\nabla}(V_L)$ is compatible with \mathcal{G}_{R_K} -actions, Frobenii, and monodromy operators, and the isomorphism $\alpha_{\text{dR}}(V_L)$ is compatible with \mathcal{G}_{R_K} -actions and strictly compatible with filtrations. Note also $D_{\text{st}}^{\nabla}(V_L)_{R_L} \cong D_{\text{dR}}(V_L)$. Hence we have \mathcal{G}_{R_K} -equivariant isomorphisms

$$\begin{aligned} V_{\text{st}}(D_{\text{st}}^{\nabla}(V_L)) &= (B_{\text{st}}^{\nabla}(R) \otimes_{L_0} D_{\text{st}}^{\nabla}(V_L))^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R) \otimes_{R_L} D_{\text{st}}^{\nabla}(V_L)_{R_L}) \\ &\cong (B_{\text{st}}^{\nabla}(R) \otimes_{L_0} D_{\text{st}}^{\nabla}(V_L))^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R) \otimes_{R_L} D_{\text{dR}}(V_L)) \\ &\cong (B_{\text{st}}^{\nabla}(R) \otimes_{\mathbb{Q}_p} V)^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R) \otimes_{\mathbb{Q}_p} V) \\ &\cong ((B_{\text{st}}^{\nabla}(R))^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R))) \otimes_{\mathbb{Q}_p} V. \end{aligned}$$

By Lemma 4.29 and Proposition 4.20, we have

$$\begin{aligned} (B_{\text{st}}^{\nabla}(R))^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R)) &= (B_{\text{st}}^{\nabla}(R))^{\varphi=1, N=0} \cap (B_{\text{dR}}^{\nabla}(R) \cap \text{Fil}^0(B_{\text{dR}}(R))) \\ &= (B_{\text{max}}^{\nabla}(R))^{\varphi=1} \cap B_{\text{dR}}^{\nabla+}(R) = \mathbb{Q}_p. \end{aligned}$$

Hence $((B_{\text{st}}^{\nabla}(R))^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R))) \otimes_{\mathbb{Q}_p} V$ is \mathcal{G}_{R_K} -equivariantly isomorphic to V . \square

Definition 5.6. For $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$, set

$$D_{\text{pst}}^{\nabla}(V) := \varinjlim_L D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}}),$$

where L ranges over all the finite Galois extensions of K inside \overline{K} . It is a vector space over K_0^{ur} equipped with a Frobenius φ , a monodromy operator N , and a semi-linear action of $\text{Gal}(\overline{K}/K)$. By unwinding the definitions, we see $D_{\text{pst}}^{\nabla}(V)^{\text{Gal}(\overline{K}/L)} = D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}})$ for every finite Galois extension L of K inside \overline{K} .

Let $\text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$ be the full subcategory of $\text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$ consisting of objects V with the following property: there exists a finite Galois extension L of K such that $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable.

Remark 5.7. By Lemma 4.22(ii), every object of $\text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$ is horizontal de Rham. We will prove that $\text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$ is exactly the full subcategory of horizontal de Rham representations if R satisfies Condition (BR) (Theorem 7.4). In this case, $\text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$ is a Tannakian subcategory of $\text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$ by Remark 4.18, Lemma 4.19 and by Theorem 7.4 or the fact that $R_{\mathcal{O}_L}$ satisfies Condition (BR) with respect to L for every finite extension L of K .

Lemma 5.8. *Let $V \in \text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$. Let L be a finite Galois extension of K inside \overline{K} such that $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable. Then there is a natural identification $D_{\text{pst}}^{\nabla}(V)_{R_{\overline{K}}} \cong R_{\overline{K}} \otimes_L D_{\text{dR}}(V|_{\mathcal{G}_{R_L}})$. Moreover, if we set*

$$\text{Fil}^{\bullet} D_{\text{pst}}^{\nabla}(V)_{R_{\overline{K}}} := R_{\overline{K}} \otimes_{R_L} \text{Fil}^{\bullet} D_{\text{dR}}(V|_{\mathcal{G}_{R_L}}),$$

then $D_{\text{pst}}^{\nabla}(V)$ is a discrete filtered $(\varphi, N, \text{Gal}(\overline{K}/K), R_K)$ -module and the filtration is independent of the choice of L .

Proof. For every finite Galois extension L' of K containing L , $V|_{\mathcal{G}_{R_{L'}}}$ is horizontal semistable and we have

$$\begin{aligned} D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_{L'}}}) &\cong L'_0 \otimes_{L_0} D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}}), \quad \text{and} \\ D_{\text{dR}}(V|_{\mathcal{G}_{R_{L'}}}) &\cong R_{L'} \otimes_{R_L} D_{\text{dR}}(V|_{\mathcal{G}_{R_L}}) \cong R_{L'} \otimes_{L_0} D_{\text{st}}^{\nabla}(V|_{\mathcal{G}_{R_L}}). \end{aligned}$$

Moreover, the first isomorphism is compatible with Frobenii, monodromy operators and $\text{Gal}(L'/K)$ -actions, and the second isomorphism is compatible with $\text{Gal}(L'/K)$ -actions and strictly compatible with filtrations. The assertions follow from these remarks and Proposition 5.3. \square

Definition 5.9. Let D be a filtered $(\varphi, N, \text{Gal}(\overline{K}/K), R_K)$ -module. Set

$$V_{\text{pst}}(D) := (B_{\text{st}}^{\nabla}(R) \otimes_{K_0^{\text{ur}}} D)^{\varphi=1, N=0} \cap \text{Fil}^0(B_{\text{dR}}(R) \otimes_{R_{\overline{K}}} D_{R_{\overline{K}}}),$$

where φ (resp. N) on $B_{\text{st}}^{\nabla}(R) \otimes_{K_0^{\text{ur}}} D$ denotes $\varphi \otimes \varphi$ (resp. $N \otimes \text{id} + \text{id} \otimes N$) and Fil^{\bullet} on $B_{\text{dR}}(R) \otimes_{R_{\overline{K}}} D_{R_{\overline{K}}}$ denotes $\sum_{m \in \mathbb{Z}} \text{Fil}^m B_{\text{dR}}(R) \otimes_{R_{\overline{K}}} \text{Fil}^{\bullet-m} D_{R_{\overline{K}}}$. It is a \mathbb{Q}_p -vector space equipped with a \mathbb{Q}_p -linear action of \mathcal{G}_{R_K} .

We say that D is *admissible* if it is isomorphic to $D_{\text{pst}}^{\nabla}(V)$ for some $V \in \text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$. Let $\text{MF}_{\overline{K}/K}^{\text{ad}}(\varphi, N, R_K)$ be the full subcategory of $\text{MF}_{\overline{K}/K}(\varphi, N, R_K)$ consisting of admissible objects.

Theorem 5.10. *Assume that R satisfies Condition (BR). Then the functor*

$$D_{\text{pst}}^{\nabla} : \text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K}) \rightarrow \text{MF}_{\overline{K}/K}^{\text{ad}}(\varphi, N, R_K)$$

is an equivalence of categories with quasi-inverse given by V_{pst} .

Proof. By Proposition 5.5, we see that the \mathbb{Q}_p -linear map $V \rightarrow V_{\text{pst}}(D_{\text{pst}}^{\nabla}(V))$ is a \mathcal{G}_{R_K} -equivariant isomorphism. For $V_1, V_2 \in \text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$, this result for $V = \text{Hom}(V_1, V_2) \in \text{Rep}_{\text{pst}}^{\nabla}(\mathcal{G}_{R_K})$ implies that D_{pst} is fully faithful. Hence the assertion follows. \square

Remark 5.11. As mentioned in the introduction, it is desirable to characterize admissible filtered $(\varphi, N, \text{Gal}(\overline{K}/K), R_K)$ -module intrinsically. When $R = K$, admissibility is equivalent to weak admissibility by [CF00]. Results on p -adic period domains suggest that admissibility will not be characterized by weak admissibility after every specialization of R to a finite extension of K (see [Har13, Example 6.7]).

In [Bri08], Brinon defines crystalline representations of \mathcal{G}_{R_K} and associates to them filtered (φ, ∇) -modules over R_K under Condition (BR). He also defines the notion of pointwise weak admissibility for filtered (φ, ∇) -modules over R_K [Bri08, Définition 7.1.11] by considering the specialization of R to the p -adic completion of the localization of R at the prime ideal (π) (when R satisfies Condition (BR)). Brinon proves that filtered (φ, ∇) -modules over R_K coming from crystalline representations are pointwisely weakly admissible ([Bri08, Proposition 8.3.4]). However, Moon [Moo18] proves that there are pointwisely weakly admissible filtered (φ, ∇) -modules that do not arise from crystalline representations.

6. PURITY FOR HORIZONTAL SEMISTABLE REPRESENTATIONS

6.1. Statement. We first recall our set-up. Let k be a perfect field of characteristic p and set $K_0 := W(k)[p^{-1}]$. Let K be a totally ramified finite extension of K_0 and fix a uniformizer π of K . We denote by C the p -adic completion of an algebraic closure \overline{K} of K . Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1. We further assume that R satisfies Condition (BR). In particular, $(\pi) \subset R$ is a prime ideal.

Consider the localization $R_{(\pi)}$ of R at the prime ideal (π) . Let \mathcal{O}_K denote the p -adic completion of $R_{(\pi)}$. It is a complete discrete valuation ring with residue field admitting a finite p -basis. Set $\mathcal{K} := \mathcal{O}_K[p^{-1}]$ and fix an algebraic closure $\overline{\mathcal{K}}$ of \mathcal{K} . Let $\mathcal{O}_{\overline{\mathcal{K}}}$ denote the ring of integers of $\overline{\mathcal{K}}$. We denote the p -adic completion of $\mathcal{O}_{\overline{\mathcal{K}}}$ by $\widehat{\mathcal{O}_{\overline{\mathcal{K}}}}$. Representations of $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K})$ are studied in [Bri06, Mor14, Ohk13]. In particular, we can discuss whether such a representation is horizontal crystalline, semistable, or de Rham (cf. [Ohk13]). Note that these classes of representations are defined by admissibility of the period rings $\mathbb{B}_{\text{cris}}(\Lambda^+)$ (or equivalently $\mathbb{B}_{\text{max}}(\Lambda^+)$), $\mathbb{B}_{\text{st}}(\Lambda^+)$, and $\mathbb{B}_{\text{dR}}(\Lambda^+)$ for $\Lambda^+ = \widehat{\mathcal{O}_{\overline{\mathcal{K}}}}$, respectively (see Lemma 4.14 and [Ohk13, §4] for horizontal semistable representations).

Let T denote the set of prime ideals of \overline{R} above $(\pi) \subset R$. Note that \mathcal{G}_{R_K} acts transitively on T . For $\mathfrak{p} \in T$, consider the decomposition subgroup

$$\mathcal{G}_{R_K}(\mathfrak{p}) := \{g \in \mathcal{G}_{R_K} \mid g(\mathfrak{p}) = \mathfrak{p}\}.$$

We now fix an R -algebra embedding $\overline{R} \hookrightarrow \mathcal{O}_{\overline{\mathcal{K}}}$. This determines a prime ideal of \overline{R} above $(\pi) \subset R$, which we denote by \mathfrak{p}_0 , and it gives rise to a homomorphism

$$\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0) := \text{Gal}(\overline{\mathcal{K}}/\mathcal{K}) \rightarrow \mathcal{G}_{R_K}.$$

Note that the map factors as $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0) \twoheadrightarrow \mathcal{G}_{R_K}(\mathfrak{p}_0) \hookrightarrow \mathcal{G}_{R_K}$ by [Bri08, Lemme 3.3.1]. We also remark that a different choice of the embedding $\overline{R} \hookrightarrow \mathcal{O}_{\overline{\mathcal{K}}}$ defines another ideal $\mathfrak{p}'_0 \in T$ and it changes the homomorphism $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K}) \rightarrow \mathcal{G}_{R_K}$ with its conjugate by an element g of \mathcal{G}_{R_K} such that $g(\mathfrak{p}_0) = \mathfrak{p}'_0$.

Theorem 6.1. *Assume that R satisfies Condition (BR). Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$. If V is horizontal de Rham and $V|_{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)}$ is horizontal semistable, then V is horizontal semistable.*

Remark 6.2. The converse of Theorem 6.1 also holds by Lemma 4.4.

The rest of this section is devoted to the proof of Theorem 6.1. Our proof is modeled after Tsuji's work on a purity theorem on crystalline local systems [Tsu, Theorem 5.4.8].

6.2. Preliminaries on relevant period rings.

Lemma 6.3. *Consider $d: \mathcal{K} \rightarrow \mathcal{K} \otimes_{\mathcal{O}_{\mathcal{K}}} \widehat{\Omega}_{\mathcal{O}_{\mathcal{K}}}$, where $\widehat{\Omega}_{\mathcal{O}_{\mathcal{K}}} := \varprojlim_m \Omega^1_{\mathcal{O}_{\mathcal{K}}/\mathbb{Z}}/p^m \Omega^1_{\mathcal{O}_{\mathcal{K}}/\mathbb{Z}}$. We have*

$$\mathrm{Ker}(d: \mathcal{K} \rightarrow \mathcal{K} \otimes_{\mathcal{O}_{\mathcal{K}}} \widehat{\Omega}_{\mathcal{O}_{\mathcal{K}}}) = K.$$

Proof. Since K is algebraically closed in R , it is also algebraically closed in \mathcal{K} . Hence the lemma follows from [Bri06, Proposition 2.28]. \square

Remark 6.4. In [Ohk13], this kernel is denoted by $\mathcal{K}_{\mathrm{can}}$ (cf. [Ohk13, Definition 1.3 (i), Remark 1.4 (i)]).

Note that $\widehat{\mathcal{O}_{\mathcal{K}}}[p^{-1}]$ is a complete algebraically closed field containing C . Hence the pair $(\widehat{\mathcal{O}_{\mathcal{K}}}[p^{-1}], \widehat{\mathcal{O}_{\mathcal{K}}})$ is a perfectoid C -Banach pair.

For $\mathfrak{p} \in T$, let $\widehat{R}_{\mathfrak{p}}$ denote the p -adic completion of the localization $\overline{R}_{\mathfrak{p}}$ of \overline{R} at \mathfrak{p} . It is naturally an \mathcal{O}_C -algebra. Since $\overline{R}_{\mathfrak{p}}$ can be written as the union of one-dimensional Noetherian normal local domains with p non-unit, $\widehat{R}_{\mathfrak{p}}[p^{-1}]$ is a complete valuation field with respect to p -adic valuation; the associated norm is given by

$$|x| = \inf \{ |a|_C \mid a \in C, x \in a\widehat{R}_{\mathfrak{p}} \}.$$

In particular, $\widehat{R}_{\mathfrak{p}}[p^{-1}]$ is a uniform C -Banach algebra and $\widehat{R}_{\mathfrak{p}}$ is a ring of integral elements.

Lemma 6.5. *The C -Banach pair $(\widehat{R}_{\mathfrak{p}}[p^{-1}], \widehat{R}_{\mathfrak{p}})$ is perfectoid.*

Proof. Since $\widehat{R}_{\mathfrak{p}}/p\widehat{R}_{\mathfrak{p}} = \overline{R}_{\mathfrak{p}}/p\overline{R}_{\mathfrak{p}}$ is the localization of $\overline{R}/p\overline{R}$ at the ideal $\mathfrak{p} \bmod p\overline{R}$, the Frobenius is surjective on $\widehat{R}_{\mathfrak{p}}/p\widehat{R}_{\mathfrak{p}}$ as it is so on $\overline{R}/p\overline{R}$. \square

To simplify our notation, set

$$\Lambda_R^+ := \widehat{R}, \quad \Lambda_{\mathfrak{p}}^+ := \widehat{R}_{\mathfrak{p}} \quad (\mathfrak{p} \in T), \quad \text{and} \quad \Lambda_{\mathcal{K}}^+ := \widehat{\mathcal{O}_{\mathcal{K}}}.$$

We also set $\Lambda_* := \Lambda_*^+[p^{-1}]$ for $*$ in $\{R, \mathfrak{p}, \mathcal{K}\}$.

Consider the $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -equivariant inclusion $\overline{R}_{\mathfrak{p}_0} \hookrightarrow \mathcal{O}_{\mathcal{K}}$, where $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ acts on $\overline{R}_{\mathfrak{p}_0}$ via $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0) \twoheadrightarrow \mathcal{G}_{R_K}(\mathfrak{p}_0)$. This extends to $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -equivariant injective maps $\Lambda_{\mathfrak{p}_0}^+ \hookrightarrow \Lambda_{\mathcal{K}}^+$ and $\Lambda_{\mathfrak{p}_0} \hookrightarrow \Lambda_{\mathcal{K}}$. Note that the former is a map of complete valuation rings of rank one and thus it has p -torsion free cokernel.

Lemma 6.6. *The natural $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -equivariant commutative diagram*

$$\begin{array}{ccc} \mathbb{B}_{\mathrm{st}, K}^+(\Lambda_{\mathfrak{p}_0}^+) & \longrightarrow & \mathbb{B}_{\mathrm{dR}}^+(\Lambda_{\mathfrak{p}_0}^+) \\ \downarrow & & \downarrow \\ \mathbb{B}_{\mathrm{st}, K}^+(\Lambda_{\mathcal{K}}^+) & \longrightarrow & \mathbb{B}_{\mathrm{dR}}^+(\Lambda_{\mathcal{K}}^+) \end{array}$$

is Cartesian.

Proof. This follows from Lemma 2.40. \square

Let us recall the notion of G -regularity.

Definition 6.7 (cf. [Fon94b, 1.4.1]). Let E be a finite extension of \mathbb{Q}_p and G a topological group. An (E, G) -ring B is called *G -regular* if the following three conditions hold:

- ($G \cdot R_1$) The ring B is reduced.
- ($G \cdot R_2$) For all $V \in \text{Rep}_E(G)$, the map $\alpha_B(V)$ is injective.
- ($G \cdot R_3$) Every one-dimensional E -vector subspace of B that is stable under G -action is generated by an invertible element of B over E .

Proposition 6.8. *Let E be a finite extension of \mathbb{Q}_p and G a topological group. Let B be an (E, G) ring that is G -regular. Then $F := B^G$ is a field, and the following holds:*

- (i) *For every $V \in \text{Rep}_E(G)$,*

$$\dim_F D_B(V) \leq \dim_E V.$$

- (ii) *Moreover, if $\dim_F D_B(V) = \dim_E V$ holds, then the map $\alpha_B(V)$ is an isomorphism.*

Proof. The proof of [Fon94b, 1.4.2] works in this setting. \square

Lemma 6.9. *We have*

$$(\mathbb{B}_{\text{st}}(\Lambda_{\mathcal{K}}^+))^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)} = K_0, \quad \text{and} \quad (\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+))^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)} = K.$$

Moreover, the $(\mathbb{Q}_p, \widehat{\mathcal{G}}_R(\mathfrak{p}_0))$ -rings $\mathbb{B}_{\text{st}, K}(\Lambda_{\mathcal{K}}^+)$ and $\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+)$ are $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -regular.

Proof. The first assertion follows from [Ohk13, Corollary 4.3, third], Lemma 6.3, and the inclusion $K \otimes_{K_0} \mathbb{B}_{\text{st}}^+(\Lambda_{\mathcal{K}}^+) \subset \mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+)$. Since $\Lambda_{\mathcal{K}}$ is a field, so is $\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+)$. Hence $\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+)$ is $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -regular by [Fon94b, Proposition 1.6.3].

By [Fon94b, Proposition 1.6.5], it remains to verify that $\mathbb{B}_{\text{st}, K}(\Lambda_{\mathcal{K}}^+)$ satisfies Condition ($G \cdot R_3$). We follow [Ohk13, Lemma 4.5] and [Fon94b, Lemme 5.1.3 (ii)].

There exists a complete valuation field $\mathcal{K}^{\text{pf}} \subset \overline{\mathcal{K}}$ containing \mathcal{K} such that the relative ramification index of \mathcal{K}^{pf} over \mathcal{K} is one and such that the residue field of \mathcal{K}^{pf} is the perfect closure of \mathcal{K} inside the residue field of $\overline{\mathcal{K}}$ (cf. [Ohk13, 1G]). Let \mathcal{L} denote the p -adic completion of the maximal unramified extension of \mathcal{K}^{pf} inside $\Lambda_{\mathcal{K}} = \widehat{\mathcal{O}_{\overline{\mathcal{K}}}}[p^{-1}]$ and let P_0 denote the fraction field of the ring of Witt vectors with coefficients in the residue field of \mathcal{L} . Then \mathcal{L} is naturally a subfield of $\Lambda_{\mathcal{K}}$, and $\mathbb{B}_{\text{st}}(\Lambda_{\mathcal{K}}^+)$ is a P_0 -algebra. Note

$$\mathcal{L} \cong K \otimes_{K_0} P_0 \quad \text{and} \quad \mathbb{B}_{\text{st}, K}(\Lambda_{\mathcal{K}}^+) = \mathcal{L} \otimes_{P_0} \mathbb{B}_{\text{st}}(\Lambda_{\mathcal{K}}^+).$$

Let $\overline{\mathcal{L}}$ denote the algebraic closure of \mathcal{L} in $\Lambda_{\mathcal{K}}$. So $\Lambda_{\mathcal{K}}$ is the p -adic completion of $\overline{\mathcal{L}}$.

We will show that every one-dimensional \mathcal{L} -vector subspace Δ of $\mathbb{B}_{\text{st}, K}(\Lambda_{\mathcal{K}}^+)$ that is stable by $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L})$ is written as $\mathcal{L}t^i$ for some $i \in \mathbb{Z}$. Since t is invertible in $\mathbb{B}_{\text{st}, K}(\Lambda_{\mathcal{K}}^+)$ and $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L})$ can be identified with a subgroup of $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$, this will complete the proof.

Let $b \in \Delta$ be a generator. By replacing b by bt^{-i} for some $i \in \mathbb{Z}$, we may assume $b \in \mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+) \setminus t\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+)$. It is enough to show $b \in \mathcal{L}$. Consider $\theta = \theta_{\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+)} : \mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+) \rightarrow \Lambda_{\mathcal{K}}$. The image $\theta(\Delta)$ is a one-dimensional \mathcal{L} -vector subspace generated by $\theta(b)$ and stable by $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L})$. By a result of Sen (cf. [Fon94b, Remarque 3.3 (ii)]), we have $\theta(b) \in \overline{\mathcal{L}} \subset \Lambda_{\mathcal{K}}$. In particular, the action of $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L})$ on

$\theta(\Delta)$ factors through $\text{Gal}(\mathcal{L}'/\mathcal{L})$ for some finite Galois extension \mathcal{L}' of \mathcal{L} inside $\overline{\mathcal{L}}$ and so does its action on Δ .

Let $\mathcal{L}(b)$ denote the \mathcal{L} -subalgebra of $\mathbb{B}_{\text{st},K}(\Lambda_{\mathcal{K}}^+)$ generated by b . Since $b \in \mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+) \setminus t\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+)$, we see that $\mathcal{L}(b)$ is $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L})$ -equivariantly isomorphic to its image $\theta(\mathcal{L}(b))$. It follows that $\mathcal{L}(b)$ is isomorphic to a subextension of \mathcal{L} inside \mathcal{L}' . This implies that $\mathcal{L}' \otimes_{\mathcal{L}} \mathcal{L}(b)$ is not an integral domain unless $\mathcal{L}(b) = \mathcal{L}$. On the other hand, we know that $\mathcal{L}' \otimes_{\mathcal{L}} \mathbb{B}_{\text{st},K}(\Lambda_{\mathcal{K}}^+) = \mathcal{L}' \otimes_{P_0} \mathbb{B}_{\text{st}}(\Lambda_{\mathcal{K}}^+)$ embeds into $\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+)$ by Corollary 2.31 (with respect to \mathcal{L}' instead of K) and thus it is an integral domain. Hence we conclude $b \in \mathcal{L}$. \square

Lemma 6.10. *We have*

$$(\mathbb{B}_{\text{st}}(\Lambda_{\mathfrak{p}}^+))^{\mathcal{G}_{R_K}(\mathfrak{p})} = K_0, \quad \text{and} \quad (\mathbb{B}_{\text{dR}}(\Lambda_{\mathfrak{p}}^+))^{\mathcal{G}_{R_K}(\mathfrak{p})} = K.$$

Moreover, the $(\mathbb{Q}_p, \mathcal{G}_{R_K}(\mathfrak{p}))$ -rings $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)$ and $\mathbb{B}_{\text{dR}}(\Lambda_{\mathfrak{p}}^+)$ are $\mathcal{G}_{R_K}(\mathfrak{p})$ -regular.

Proof. Without loss of generality we may assume $\mathfrak{p} = \mathfrak{p}_0$. The first assertion follows from Lemmas 6.6 and 6.9. Since $\Lambda_{\mathfrak{p}_0}$ is a field, so is $\mathbb{B}_{\text{dR}}(\Lambda_{\mathfrak{p}_0}^+)$. Hence $\mathbb{B}_{\text{dR}}(\Lambda_{\mathfrak{p}_0}^+)$ is $\mathcal{G}_{R_K}(\mathfrak{p}_0)$ -regular. It remains to verify that $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)$ satisfies Condition $(G \cdot R_3)$ by [Fon94b, Proposition 1.6.5]. This follows from the $\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)$ -equivariant Cartesian diagram in Lemma 6.6 and the fact that $\mathbb{B}_{\text{dR}}(\Lambda_{\mathfrak{p}_0}^+)$, $\mathbb{B}_{\text{st},K}(\Lambda_{\mathcal{K}}^+)$, and $\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+)$ satisfy Condition $(G \cdot R_3)$. \square

6.3. Proof of Theorem 6.1. Let us go back to the setting of Theorem 6.1. By assumption, $V|_{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)}$ is $\mathbb{B}_{\text{st}}(\Lambda_{\mathcal{K}}^+)$ -admissible and thus $\mathbb{B}_{\text{st},K}(\Lambda_{\mathcal{K}}^+)$ -admissible. By replacing V by its Tate twist $V(-n)$ for $n \gg 0$, we may assume

$$(\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}} \cong (\mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}$$

and

$$(\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)} \xrightarrow{\cong} (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)} \xrightarrow{\cong} (\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)}.$$

In particular, the natural map $(\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}} \rightarrow (\mathbb{B}_{\text{dR}}(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)}$ is an isomorphism.

Lemma 6.11. *The representation $V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)}$ is $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)$ -admissible. Moreover, $(\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} = (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)}$.*

Proof. Consider the commutative diagram

$$\begin{array}{ccc} (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} & \hookrightarrow & (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} \\ \downarrow & & \downarrow \\ (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)} & \xrightarrow{\cong} & (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathcal{K}}^+) \otimes_{\mathbb{Q}_p} V)^{\widehat{\mathcal{G}}_{R_K}(\mathfrak{p}_0)}. \end{array}$$

It follows from the remark before the lemma that the right vertical map is an isomorphism of K -vector spaces of dimension $\dim_{\mathbb{Q}_p} V$. By Lemma 6.6, we conclude that all the four K -vector spaces in the diagram are equal, and we obtain the second assertion. We also have

$$\dim_{\mathbb{Q}_p} V = \dim_K(\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} \leq \dim_K(\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)}.$$

By Proposition 6.8 and Lemma 6.10, we have $\dim_K(\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} = \dim_{\mathbb{Q}_p} V$ and thus $V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)}$ is $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)$ -admissible. \square

Observe that each $\sigma \in \mathcal{G}_{R_K}$ induces an R -algebra isomorphism $\Lambda_{\mathfrak{p}}^+ \rightarrow \Lambda_{\sigma(\mathfrak{p})}^+$ for $\mathfrak{p} \in T$, and $\mathcal{G}_{R_K}(\sigma(\mathfrak{p})) = \sigma \mathcal{G}_{R_K}(\mathfrak{p}) \sigma^{-1}$ in \mathcal{G}_{R_K} .

Lemma 6.12. *For $\mathfrak{p} \in T$, the representation $V|_{\mathcal{G}_{R_K}(\mathfrak{p})}$ is $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)$ -admissible. Moreover, $(\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})} = (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}$.*

Proof. Choose $\sigma \in \mathcal{G}_{R_K}$ with $\sigma(\mathfrak{p}_0) = \mathfrak{p}$. Hence $\mathcal{G}_{R_K}(\mathfrak{p}) = \sigma \mathcal{G}_{R_K}(\mathfrak{p}_0) \sigma^{-1}$ in \mathcal{G}_{R_K} , and σ induces isomorphisms $\sigma: \Lambda_{\mathfrak{p}_0}^+ \xrightarrow{\cong} \Lambda_{\mathfrak{p}}^+$ and $\sigma: \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \xrightarrow{\cong} \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)$. These maps are compatible with the group homomorphism $\mathcal{G}_{R_K}(\mathfrak{p}_0) \rightarrow \mathcal{G}_{R_K}(\mathfrak{p})$ given by σ -conjugation.

Hence for every $\tau \in \mathcal{G}_{R_K}(\mathfrak{p}_0)$, we have a commutative diagram

$$\begin{array}{ccc} \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow[\cong]{\sigma \otimes \sigma} & \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V \\ \tau \otimes \tau \downarrow & & \downarrow \sigma \tau \sigma^{-1} \otimes \sigma \tau \sigma^{-1} \\ \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow[\cong]{\sigma \otimes \sigma} & \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V. \end{array}$$

In particular, if $x \in \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V$ is $\mathcal{G}_{R_K}(\mathfrak{p}_0)$ -invariant, then $(\sigma \otimes \sigma)(x) \in \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V$ is $\mathcal{G}_{R_K}(\mathfrak{p})$ -invariant. It follows that $\sigma \otimes \sigma$ restricts to a K -linear isomorphism $D_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0})}(V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)}) \xrightarrow{\cong} D_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}})}(V|_{\mathcal{G}_{R_K}(\mathfrak{p})})$. Hence we have

$$\dim_K D_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}})}(V|_{\mathcal{G}_{R_K}(\mathfrak{p})}) = \dim_{\mathbb{Q}_p} V.$$

By Proposition 6.8 and Lemma 6.10, we conclude that $V|_{\mathcal{G}_{R_K}(\mathfrak{p})}$ is $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)$ -admissible.

By a similar argument, we obtain the commutative diagram with isomorphic vertical maps

$$\begin{array}{ccc} (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} & \hookrightarrow & (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p}_0)} \\ \sigma \otimes \sigma \downarrow \cong & & \downarrow \sigma \otimes \sigma \cong \\ (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})} & \hookrightarrow & (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}. \end{array}$$

Hence the second assertion follows from the one for \mathfrak{p}_0 . \square

Set

$$\Lambda_T^+ := \prod_{\mathfrak{p} \in T} \Lambda_{\mathfrak{p}}^+ = \prod_{\mathfrak{p} \in T} \widehat{\overline{R}_{\mathfrak{p}}}, \quad \text{and} \quad \Lambda_T := \Lambda_T[p^{-1}].$$

Hence Λ_T is the product of $\Lambda_{\mathfrak{p}}$ over $\mathfrak{p} \in T$ as a C -Banach algebra.

Lemma 6.13. *The map*

$$\Lambda_R^+ \rightarrow \Lambda_T^+$$

is injective with p -torsion free cokernel.

Proof. Recall that \overline{R} can be written as the union of Noetherian normal domains finite over R . Since $R/\pi R$ is an integral domain, we deduce that the natural map $\overline{R}/\pi \overline{R} \rightarrow \prod_{\mathfrak{p} \in T} \overline{R}_{\mathfrak{p}}/\pi \overline{R}_{\mathfrak{p}}$ is injective. It follows that $\overline{R}/\pi^m \overline{R} \rightarrow \prod_{\mathfrak{p} \in T} \overline{R}_{\mathfrak{p}}/\pi^m \overline{R}_{\mathfrak{p}}$ is injective for every $m \in \mathbb{N}$. By taking the inverse limit over m , we see that $\Lambda_R^+ \rightarrow \prod_{\mathfrak{p} \in T} \Lambda_{\mathfrak{p}}^+ = \Lambda_T^+$ is injective with π -torsion free cokernel. Hence the cokernel is also p -torsion free. \square

Each $\sigma \in \mathcal{G}_{R_K}$ induces an R -algebra isomorphism $\Lambda_{\mathfrak{p}}^+ \rightarrow \Lambda_{\sigma(\mathfrak{p})}^+$ for $\mathfrak{p} \in T$. This gives rise to a natural \mathcal{G}_{R_K} -action on Λ_T^+ making the injection $\Lambda_R^+ \rightarrow \Lambda_T^+ \mathcal{G}_{R_K}$ -equivariant (cf. [Bri08, Remarque 3.3.2]).

Since products commute with inverse limits and the Witt vector functor, we have $\mathbb{A}_{\text{inf}}(\Lambda_T^+) = \prod_{\mathfrak{p} \in T} \mathbb{A}_{\text{inf}}(\Lambda_{\mathfrak{p}}^+)$. Since \mathcal{O}_K is finite free over $W(k)$, we conclude

$$\mathbb{A}_{\text{inf},K}(\Lambda_T^+) = \prod_{\mathfrak{p} \in T} \mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}}^+),$$

and $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda_T^+)}$ is the product of $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}}^+)}$ over $\mathfrak{p} \in T$.

Lemma 6.14. *The natural commutative diagram*

$$\begin{array}{ccc} \mathbb{B}_{\text{st},K}^+(\Lambda_R^+) & \longrightarrow & \mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \\ \downarrow & & \downarrow \\ \mathbb{B}_{\text{st},K}^+(\Lambda_T^+) & \longrightarrow & \mathbb{B}_{\text{dR}}^+(\Lambda_T^+) \end{array}$$

is Cartesian.

Proof. This follows from Lemmas 2.40 and 6.13. \square

For each $\mathfrak{p} \in T$, choose $\sigma_{\mathfrak{p}} \in \mathcal{G}_{R_K}$ such that $\sigma_{\mathfrak{p}}(\mathfrak{p}_0) = \mathfrak{p}$ and such that $\sigma_{\mathfrak{p}}|_{\overline{K}} = \text{id}$. This is possible since $R/\pi R$ is geometrically integral over k . We also assume $\sigma_{\mathfrak{p}_0} = \text{id}$. We have an isomorphism

$$\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}}: \mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V \xrightarrow{\cong} \mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V$$

compatible with the homomorphism $\mathcal{G}_{R_K}(\mathfrak{p}_0) \rightarrow \mathcal{G}_{R_K}(\mathfrak{p})$ given by $\sigma_{\mathfrak{p}}$ -conjugation.

Lemma 6.15.

(i) *The map*

$$\mathbb{B}_{\text{st},K}^+(\Lambda_T^+) \rightarrow \prod_{\mathfrak{p} \in T} \mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+)$$

is injective.

(ii) *If $x = (x_{\mathfrak{p}})_{\mathfrak{p}} \in \prod_{\mathfrak{p} \in T} (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)$ satisfies*

$$(\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x_{\mathfrak{p}_0}) = x_{\mathfrak{p}}$$

for every $\mathfrak{p} \in T$, then

$$x \in \mathbb{B}_{\text{st},K}^+(\Lambda_T^+) \otimes_{\mathbb{Q}_p} V.$$

Proof. Consider $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}_0}^+)}: \mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}_0}^+) \rightarrow \Lambda_{\mathfrak{p}_0}^+$ and choose an \mathcal{O}_K -linear section $s_{\mathfrak{p}_0}: \Lambda_{\mathfrak{p}_0}^+ \rightarrow \mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}_0}^+)$ of $\theta_{\mathbb{A}_{\text{inf},K}(\Lambda_{\mathfrak{p}_0}^+)}$ as in Construction 2.22. By the argument there, it defines a K -linear isomorphism

$$\tilde{\theta}_{v,\mathfrak{p}_0}: \mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \xrightarrow{\cong} \Lambda_{\mathfrak{p}_0} \langle X \rangle [\log(1+X)].$$

For each $\mathfrak{p} \in T$, we set

$$s_{\mathfrak{p}} := \sigma_{\mathfrak{p}} \circ s_{\mathfrak{p}_0} \circ \sigma_{\mathfrak{p}}^{-1} : \Lambda_{\mathfrak{p}}^+ \rightarrow \Lambda_{\mathfrak{p}_0}^+ \rightarrow \mathbb{A}_{\text{inf}, K}(\Lambda_{\mathfrak{p}_0}^+) \rightarrow \mathbb{A}_{\text{inf}, K}(\Lambda_{\mathfrak{p}}^+).$$

Then $s_{\mathfrak{p}}$ is an \mathcal{O}_K -linear section of $\theta_{\mathbb{A}_{\text{inf}, K}(\Lambda_{\mathfrak{p}}^+)}$ satisfying $s_{\mathfrak{p}} \circ \sigma_{\mathfrak{p}} = \sigma_{\mathfrak{p}} \circ s_{\mathfrak{p}_0}$. It also yields a K -linear isomorphism

$$\tilde{\theta}_{v, \mathfrak{p}} : \mathbb{B}_{\text{st}, K}^+(\Lambda_{\mathfrak{p}}^+) \xrightarrow{\cong} \Lambda_{\mathfrak{p}} \langle X \rangle [\log(1 + X)].$$

Let $\sigma_{\mathfrak{p}} : \Lambda_{\mathfrak{p}_0} \langle X \rangle [\log(1 + X)] \rightarrow \Lambda_{\mathfrak{p}} \langle X \rangle [\log(1 + X)]$ denote the continuous K -linear map extending $\sigma_{\mathfrak{p}} : \Lambda_{\mathfrak{p}_0} \rightarrow \Lambda_{\mathfrak{p}}$ by $\sigma_{\mathfrak{p}}(X^l (\log(1 + X))^m) = X^l (\log(1 + X))^m$. We see $\tilde{\theta}_{v, \mathfrak{p}} \circ \sigma_{\mathfrak{p}} = \sigma_{\mathfrak{p}} \circ \tilde{\theta}_{v, \mathfrak{p}_0}$ by Lemma 2.23(ii), Proposition 2.30, and the assumption $\sigma_{\mathfrak{p}}|_{\overline{K}} = \text{id}$.

Similarly,

$$\prod_{\mathfrak{p} \in T} s_{\mathfrak{p}} : \Lambda_T^+ = \prod_{\mathfrak{p} \in T} \Lambda_{\mathfrak{p}}^+ \rightarrow \prod_{\mathfrak{p} \in T} \mathbb{A}_{\text{inf}, K}(\Lambda_{\mathfrak{p}}^+) = \mathbb{A}_{\text{inf}, K}(\Lambda_T^+)$$

is an \mathcal{O}_K -linear section of $\theta_{\mathbb{A}_{\text{inf}, K}(\Lambda_T^+)}$, and it yields a K -linear isomorphism

$$\tilde{\theta}_{v, T} : \mathbb{B}_{\text{st}, K}^+(\Lambda_T^+) \xrightarrow{\cong} \Lambda_T \langle X \rangle [\log(1 + X)].$$

Then we have a commutative diagram

$$\begin{array}{ccc} \mathbb{B}_{\text{st}, K}^+(\Lambda_T^+) & \longrightarrow & \prod_{\mathfrak{p} \in T} \mathbb{B}_{\text{st}, K}^+(\Lambda_{\mathfrak{p}}^+) \\ \tilde{\theta}_{v, T} \downarrow \cong & & \prod \tilde{\theta}_{v, \mathfrak{p}} \downarrow \cong \\ \Lambda_T \langle X \rangle [\log(1 + X)] & \longrightarrow & \prod_{\mathfrak{p} \in T} \Lambda_{\mathfrak{p}} \langle X \rangle [\log(1 + X)]. \end{array}$$

For $h \in \mathbb{N}$ and $*$ $\in \{T, \mathfrak{p}\}$, let $\mathbb{A}_{\text{st}, K}^{\leq h}(\Lambda_*^+)$ denote the $\mathbb{A}_{\text{max}, K}(\Lambda_*^+)$ -submodule $\bigoplus_{i=0}^h \mathbb{A}_{\text{max}, K}(\Lambda_*^+) (\log[\pi^b]/\pi)^i$ of $\mathbb{B}_{\text{st}, K}^+(\Lambda_*^+)$. Note $\mathbb{B}_{\text{st}, K}^+(\Lambda_*^+) = \bigcup_h \mathbb{A}_{\text{st}, K}^{\leq h}(\Lambda_*^+) [p^{-1}]$. By Lemma 2.23(ii) and (iv), we have the functorial identification

$$\tilde{\theta}_{v, *} : \mathbb{A}_{\text{st}, K}^{\leq h}(\Lambda_*^+) \xrightarrow{\cong} \bigoplus_{i=0}^h \Lambda_*^+ \langle X \rangle (\log(1 + X))^i =: \Lambda_*^+ \langle X \rangle [\log(1 + X)]^{\leq h}$$

as \mathcal{O}_K -modules.

Recall

$$\Lambda_*^+ \langle X \rangle = \left\{ \sum_{i=0}^{\infty} \lambda_i X^i \in \Lambda_*^+ [[X]] \mid \forall l \in \mathbb{N}, \lambda_i \in p^l \Lambda_*^+ \text{ for all but finitely many } i \right\}.$$

In particular, $\Lambda_T^+ \langle X \rangle \rightarrow \prod_{\mathfrak{p} \in T} (\Lambda_{\mathfrak{p}}^+ \langle X \rangle)$ is injective.

We now prove Part (i). By the above remark and the transcendence of $\log(1 + X)$ (Lemma 2.29), we see that the map

$$\Lambda_T^+ \langle X \rangle [\log(1 + X)]^{\leq h} \rightarrow \prod_{\mathfrak{p} \in T} (\Lambda_{\mathfrak{p}}^+ \langle X \rangle [\log(1 + X)]^{\leq h})$$

is injective. Hence the map $\mathbb{A}_{\text{st}, K}^{\leq h}(\Lambda_T^+) \rightarrow \prod_{\mathfrak{p} \in T} \mathbb{A}_{\text{st}, K}^{\leq h}(\Lambda_{\mathfrak{p}}^+)$ is injective and Part (i) follows.

For Part (ii), consider the commutative diagram

$$\begin{array}{ccc}
\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow[\cong]{\tilde{\theta}_{v,\mathfrak{p}_0} \otimes \text{id}} & \Lambda_{\mathfrak{p}_0} \langle X \rangle [\log(1+X)] \otimes_{\mathbb{Q}_p} V \\
\downarrow \sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}} & & \downarrow \sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}} \\
\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow[\cong]{\tilde{\theta}_{v,\mathfrak{p}} \otimes \text{id}} & \Lambda_{\mathfrak{p}} \langle X \rangle [\log(1+X)] \otimes_{\mathbb{Q}_p} V.
\end{array}$$

Then the assertion is equivalent to the following:

If $x = (x_{\mathfrak{p}})_{\mathfrak{p}} \in \prod_{\mathfrak{p} \in T} (\Lambda_{\mathfrak{p}} \langle X \rangle [\log(1+X)] \otimes_{\mathbb{Q}_p} V)$ satisfies $(\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x_{\mathfrak{p}_0}) = x_{\mathfrak{p}}$ for every $\mathfrak{p} \in T$, then $x \in \Lambda_T \langle X \rangle [\log(1+X)] \otimes_{\mathbb{Q}_p} V$.

Fix a \mathcal{G}_{R_K} -stable \mathbb{Z}_p -lattice $V_{\mathbb{Z}_p}$ of V and choose such $x = (x_{\mathfrak{p}})_{\mathfrak{p}}$. There exist $m, h \in \mathbb{N}$ such that $p^m x_{\mathfrak{p}_0} \in \Lambda_{\mathfrak{p}_0}^+ \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}$. Then for every $\mathfrak{p} \in T$, we have

$$p^m x_{\mathfrak{p}} = (\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(p^m x_{\mathfrak{p}_0}) \in \Lambda_{\mathfrak{p}}^+ \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}.$$

Moreover, we see that if there exists $l \in \mathbb{N}$ and $f(X) \in \Lambda_{\mathfrak{p}_0}^+ [X] [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}$ (i.e., a polynomial in X and $\log(1+X)$) such that

$$p^m x_{\mathfrak{p}_0} - f(X) \in p^l (\Lambda_{\mathfrak{p}_0}^+ \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}),$$

then

$$p^m x_{\mathfrak{p}} - (\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(f(X)) \in p^l (\Lambda_{\mathfrak{p}}^+ \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}).$$

This implies $p^m x \in \Lambda_T^+ \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}$, and thus $x \in \Lambda_T \langle X \rangle [\log(1+X)]^{\leq h} \otimes_{\mathbb{Q}_p} V$. \square

Remark 6.16. With the notation as in the proof, the same argument as in the last part of the proof shows the following: Let $D_{\mathcal{O}_K}$ be a finite free \mathcal{O}_K -module and let $W \subset \mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$ be a finite \mathbb{Z}_p -submodule. Then the submodule $\prod_{\mathfrak{p} \in T} ((\sigma_{\mathfrak{p}} \otimes \text{id})(W))$ of $\prod_{\mathfrak{p} \in T} (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K})$ lands in $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_T^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$. In particular, if $x = (x_{\mathfrak{p}}) \in \prod_{\mathfrak{p} \in T} (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K})$ satisfies $x_{\mathfrak{p}} \in (\sigma_{\mathfrak{p}} \otimes \text{id})(W) \subset \mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$ for every $\mathfrak{p} \in T$, then $x \in \mathbb{A}_{\text{st},T}^{\leq h}(\Lambda_T^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$.

Lemma 6.17. *Let $x \in (\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}$ and let $x_{\mathfrak{p}}$ denote the image of x in $(\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}$. Then for each $\mathfrak{p} \in T$, we have*

$$(\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x_{\mathfrak{p}_0}) = x_{\mathfrak{p}}.$$

Proof. This follows from the commutative diagram

$$\begin{array}{ccc}
\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow{\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}}} & \mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V \\
\downarrow & & \downarrow \\
\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V & \xrightarrow{\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}}} & \mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V.
\end{array}$$

Namely, we have $(\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x_{\mathfrak{p}_0}) = ((\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x))_{\mathfrak{p}} = x_{\mathfrak{p}}$. \square

Lemma 6.18. *The inclusion*

$$(\mathbb{B}_{\text{st},K}^+(\Lambda_R) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}} \subset (\mathbb{B}_{\text{dR}}^+(\Lambda_R) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}.$$

is an equality.

Proof. Let $x \in (\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}$. For each $\mathfrak{p} \in T$, let $x_{\mathfrak{p}}$ denote the image of x in $(\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}$. Then we have $(\sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}})(x_{\mathfrak{p}_0}) = x_{\mathfrak{p}}$ by the previous lemma. Since $(\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})} = (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}$ by Lemma 6.12, we have $(x_{\mathfrak{p}})_{\mathfrak{p}} \in \prod_{\mathfrak{p} \in T} (\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)$. Hence we get $x \in \mathbb{B}_{\text{st},K}^+(\Lambda_T^+) \otimes_{\mathbb{Q}_p} V$ by Lemma 6.15 (ii). Since the commutative diagram

$$\begin{array}{ccc} \mathbb{B}_{\text{st},K}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V & \longrightarrow & \mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ \mathbb{B}_{\text{st},K}^+(\Lambda_T^+) \otimes_{\mathbb{Q}_p} V & \longrightarrow & \mathbb{B}_{\text{dR}}^+(\Lambda_T^+) \otimes_{\mathbb{Q}_p} V \end{array}$$

is Cartesian by Lemma 6.14, we conclude $x \in \mathbb{B}_{\text{st},K}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$. \square

Let us complete the proof of Theorem 6.1. Set

$$D := (\mathbb{B}_{\text{st},K}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}} = (\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}} = (\mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}.$$

This is a K -vector space and $\dim_K D = \dim_{\mathbb{Q}_p} V$. Consider the commutative diagram

$$\begin{array}{ccc} \mathbb{B}_{\text{st},K}(\Lambda_R^+) \otimes_K D & \xrightarrow{\alpha_{\text{st},K}^{\nabla}(V)} & \mathbb{B}_{\text{st},K}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V \\ \downarrow & & \downarrow \\ \mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_K D & \xrightarrow[\cong]{\alpha_{\text{dR}}^{\nabla}(V)} & \mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V. \end{array}$$

We need to show that $\alpha_{\text{st},K}^{\nabla}(V)$ is surjective. Choose $m \in \mathbb{N}$ such that the \mathbb{Q}_p -vector subspace $\mathbb{Q}_p t^m \otimes_{\mathbb{Q}_p} V$ of $\mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$ is contained in the image of $\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_K D$ under $\alpha_{\text{dR}}^{\nabla}(V)$. Fix a \mathcal{G}_{R_K} -stable \mathbb{Z}_p -lattice $V_{\mathbb{Z}_p}$ of V and set $V'_{\mathbb{Z}_p} := \mathbb{Z}_p t^m \otimes_{\mathbb{Z}_p} V_{\mathbb{Z}_p}$. It suffices to prove that $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{st},K}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$, is contained in the image of $\alpha_{\text{st},K}^{\nabla}(V)$.

For $h \in \mathbb{N}$ and $* \in \{R, T, \mathfrak{p}\}$, consider the $\mathbb{A}_{\text{max},K}(\Lambda_*^+)$ -submodule

$$\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_*^+) := \bigoplus_{i=0}^h \mathbb{A}_{\text{max},K}(\Lambda_*^+) \left(\log \frac{[\pi^b]}{\pi} \right)^i \subset \mathbb{B}_{\text{st},K}^+(\Lambda_*^+).$$

Note $\mathbb{B}_{\text{st},K}^+(\Lambda_*^+) = \bigcup_h \mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_*^+)[p^{-1}]$. Observe that each $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_R^+)[p^{-1}]$ is stable under the action of \mathcal{G}_{R_K} and

$$(\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_R^+)[p^{-1}])^{\mathcal{G}_{R_K}} = (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+)[p^{-1}])^{\mathcal{G}_{R_K}(\mathfrak{p})} = K.$$

Since D is finite-dimensional over K , there exists $h \in \mathbb{N}$ such that

$$D = (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_R^+)[p^{-1}] \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}}.$$

It follows that for each $\mathfrak{p} \in T$, we have the isomorphism

$$D \xrightarrow{\cong} (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+)[p^{-1}] \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})} = (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V)^{\mathcal{G}_{R_K}(\mathfrak{p})}.$$

By increasing h if necessary, we may further assume that $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V$, is contained in the image of $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}_0}^+)[p^{-1}] \otimes_K D$ under the

isomorphism

$$\alpha_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)}(V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)}): \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_K D \xrightarrow{\cong} \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V.$$

Choose an \mathcal{O}_K -lattice $D_{\mathcal{O}_K}$ of D such that $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V$, is contained in the image of $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$ under $\alpha_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)}(V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)})$.

For each $\mathfrak{p} \in T$, the functoriality of the construction gives a commutative diagram

$$\begin{array}{ccc} \mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K} & \xrightarrow{\alpha_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+)}(V|_{\mathcal{G}_{R_K}(\mathfrak{p}_0)})} & \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}_0}^+) \otimes_{\mathbb{Q}_p} V \\ \downarrow \sigma_{\mathfrak{p}} \otimes \text{id} & & \downarrow \sigma_{\mathfrak{p}} \otimes \sigma_{\mathfrak{p}} \\ \mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K} & \xrightarrow{\alpha_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)}(V|_{\mathcal{G}_{R_K}(\mathfrak{p})})} & \mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V. \end{array}$$

Since $V'_{\mathbb{Z}_p}$ is \mathcal{G}_{R_K} -stable, we conclude that $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{st},K}^+(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathbb{Q}_p} V$, is contained in the image of $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$ under $\alpha_{\mathbb{B}_{\text{st},K}(\Lambda_{\mathfrak{p}}^+)}(V|_{\mathcal{G}_{R_K}(\mathfrak{p})})$.

Consider the isomorphism

$$\alpha_{\text{dR}}^{\nabla}(V): \mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_K D \xrightarrow{\cong} \mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$$

and let $W \subset \mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_K D$ denote the inverse image of $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{dR}}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$, under $\alpha_{\text{dR}}^{\nabla}(V)$. Observe $W \subset \mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_K D$. By the above discussion, the image of W under the map

$$\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_K D \rightarrow \prod_{\mathfrak{p} \in T} (\mathbb{B}_{\text{dR}}^+(\Lambda_{\mathfrak{p}}^+) \otimes_K D)$$

lands in the subspace $\prod_{\mathfrak{p} \in T} (\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_{\mathfrak{p}}^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K})$. It follows from Remark 6.16 that the image of W under the map

$$\mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_K D \rightarrow \mathbb{B}_{\text{dR}}^+(\Lambda_T^+) \otimes_K D$$

lands in the subspace $\mathbb{A}_{\text{st},K}^{\leq h}(\Lambda_T^+) \otimes_{\mathcal{O}_K} D_{\mathcal{O}_K}$ and thus in $\mathbb{B}_{\text{st},K}^+(\Lambda_T^+) \otimes_K D$. By Lemma 6.14, we conclude that W lands in the subspace

$$\mathbb{B}_{\text{st},K}^+(\Lambda_R^+) \otimes_K D \subset \mathbb{B}_{\text{dR}}^+(\Lambda_R^+) \otimes_K D.$$

This implies that $V'_{\mathbb{Z}_p}$, as a submodule of $\mathbb{B}_{\text{st},K}(\Lambda_R^+) \otimes_{\mathbb{Q}_p} V$, is contained in the image of $\alpha_{\text{st},K}^{\nabla}(V)$. Hence V is a horizontal semistable representation of \mathcal{G}_{R_K} . This completes the proof of Theorem 6.1.

7. p -ADIC MONODROMY THEOREM FOR HORIZONTAL DE RHAM REPRESENTATIONS

7.1. Horizontal version of p -adic monodromy theorem for Galois representations. Let us recall a result of Ohkubo. We use a slightly different notation from [Ohk13]. Let \mathcal{K} be a complete discrete valuation field of mixed characteristic $(0, p)$ and let $k_{\mathcal{K}}$ denote the residue field of \mathcal{K} ⁸. Fix an algebraic closure $\overline{\mathcal{K}}$ of \mathcal{K} . Define the subfield \mathcal{K}_{can} of \mathcal{K} as the algebraic closure of $W(k_{\mathcal{K}}^p)[p^{-1}]$ in \mathcal{K} , where

⁸[Ohk13] uses K for our \mathcal{K} . In our paper, K is a finite extension of \mathbb{Q}_p .

$k_{\mathcal{K}}^{p^\infty} := \bigcap_{m \in \mathbb{N}} k_{\mathcal{K}}^{p^m}$. Note that \mathcal{K}_{can} coincides with the kernel of the canonical derivation $d: \mathcal{K} \rightarrow \mathcal{K} \otimes_{\mathcal{O}_{\mathcal{K}}} \widehat{\Omega}_{\mathcal{O}_{\mathcal{K}}}$ (cf. [Ohk13, Remark 1.4 (i), Proposition 1.13]). We use a similar notation for finite extensions of \mathcal{K} .

Ohkubo proves the p -adic monodromy theorem for horizontal de Rham representations of \mathcal{K} :

Theorem 7.1 ([Ohk13, Theorem 7.4]). *Let $V_{\mathcal{K}}$ be a horizontal de Rham representation of $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K})$. Then there exists a finite extension L of \mathcal{K}_{can} such that $V_{\mathcal{K}}|_{\text{Gal}(\overline{\mathcal{K}}/\mathcal{K}L)}$ is horizontal semistable, where $\mathcal{K}L$ is the composite field of \mathcal{K} and L in $\overline{\mathcal{K}}$.*

Remark 7.2. When $k_{\mathcal{K}}$ is perfect, the above theorem is the usual p -adic monodromy theorem, and it was first proved by Berger [Ber02, Théorème 0.7]. When $k_{\mathcal{K}}$ is imperfect, we can also consider the statement that a de Rham representation is potentially semistable. This statement is proved by Morita when $k_{\mathcal{K}}$ admits a finite p -basis [Mor14, Corollary 1.2] and by Ohkubo in general [Ohk13, Main Theorem]. Ohkubo deduces Theorem 7.1 from this version of p -adic monodromy theorem.

Remark 7.3. [Ohk13, Theorem 7.6] claims that every horizontal de Rham representation of $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K})$ comes from a de Rham representation of $\text{Gal}(\overline{\mathcal{K}_{\text{can}}}/\mathcal{K}_{\text{can}})$ via the pullback along $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K}) \rightarrow \text{Gal}(\overline{\mathcal{K}_{\text{can}}}/\mathcal{K}_{\text{can}})$, which is not correct. The error comes from an incorrect argument in the proof of [Ohk13, Proposition 7.3] (cf. Remark 4.30 and Example 4.31). However, these errors are irrelevant to the proofs of the p -adic monodromy theorem and the horizontal p -adic monodromy theorem [Ohk13, Main Theorem, Theorem 7.4], and the latter theorems are correct.

7.2. p -adic monodromy theorem for horizontal de Rham representations.

Let K be a complete discrete valuation field of characteristic zero with perfect residue field k of characteristic p . Fix a uniformizer π of K . Let R be an \mathcal{O}_K -algebra satisfying the conditions in Set-up 3.1. We further assume that R satisfies Condition (BR). For each finite extension L of K , we denote $\mathcal{O}_L \otimes_{\mathcal{O}_K} R$ by $R_{\mathcal{O}_L}$. Then $R_{\mathcal{O}_L}$ also satisfies the conditions in Set-up 3.1 and Condition (BR) with respect to L . As before, we write R_K (resp. R_L) for $R[p^{-1}]$ (resp. $R_{\mathcal{O}_L}[p^{-1}]$).

Theorem 1.9 in the introduction is a special case of the following p -adic monodromy theorem for horizontal de Rham representations:

Theorem 7.4. *Assume that R satisfies Condition (BR). Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_{R_K})$. If V is horizontal de Rham, then there exists a finite extension L of K such that $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable.*

Proof. We use the notation in Subsection 6.1. Consider the complete discrete valuation field $\mathcal{K} := \widehat{R_{(\pi)}}[p^{-1}]$. Fix an algebraic closure $\overline{\mathcal{K}}$ of \mathcal{K} and an R -algebra embedding $\overline{R} \hookrightarrow \overline{\mathcal{K}}$. Note $\mathcal{K}_{\text{can}} = K$ by Lemma 6.3.

Since V is a horizontal de Rham representation of \mathcal{G}_{R_K} , $V_{\mathcal{K}} := V|_{\text{Gal}(\overline{\mathcal{K}}/\mathcal{K})}$ is a horizontal de Rham representation of $\text{Gal}(\overline{\mathcal{K}}/\mathcal{K})$. Hence by Theorem 7.1, there exists a finite extension L of K such that $V_{\mathcal{K}}|_{\text{Gal}(\overline{\mathcal{K}}/\mathcal{K}L)}$ is horizontal semistable.

We prove that $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable. Let π_L denote a uniformizer of L . Since $R/\pi R$ is geometrically integral over k , $(\pi_L) \subset R_{\mathcal{O}_L}$ is a prime ideal and thus the ring of integers of $\mathcal{K}L$ coincides with the p -adic completion $(R_{\mathcal{O}_L})_{(\pi_L)}$ of the localization of $R_{\mathcal{O}_L}$ at (π_L) . Hence $\mathcal{K}L = (\widehat{R_{\mathcal{O}_L}})_{(\pi_L)}[p^{-1}]$.

We know that $V|_{\mathcal{G}_{R_L}}|_{\mathrm{Gal}(\overline{K}/LK)} = V_K|_{\mathrm{Gal}(\overline{K}/LK)}$ is a horizontal semistable representation of $\mathrm{Gal}(\overline{K}/LK)$. The embedding

$$\overline{R_{\mathcal{O}_L}} = \overline{R} \hookrightarrow \overline{K}$$

determines a prime ideal \mathfrak{p}_0 of $\overline{R_{\mathcal{O}_L}}$ above $(\pi_L) \subset R_{\mathcal{O}_L}$, and $\mathrm{Gal}(\overline{K}/LK)$ coincides with $\widehat{\mathcal{G}}_{R_L}(\mathfrak{p}_0)$. Therefore $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable by Theorem 6.1. \square

Theorem 7.5. *Assume R satisfies Condition (BR). Let V be a horizontal de Rham representation of \mathcal{G}_{R_K} . Assume that there exists a finite extension K' of K and a map $f: R \rightarrow K'$ such that f^*V is a potentially crystalline representation of $\mathrm{Gal}(\overline{K}/K')$. Then there exists a finite extension L of K such that $V|_{\mathcal{G}_{R_L}}$ is horizontal crystalline.*

Proof. By extending K' if necessary, we may assume that the representation f^*V of $\mathrm{Gal}(\overline{K}/K')$ is crystalline (equivalently, horizontal crystalline). By Theorem 7.4, there exists a finite extension L of K' such that $V|_{\mathcal{G}_{R_L}}$ is horizontal semistable. Then the assertion follows from Proposition 4.23. \square

8. p -ADIC HODGE THEORY FOR RIGID ANALYTIC VARIETIES

In this section, we review p -adic local systems and period sheaves. Let K be a complete discrete valuation field of characteristic zero with perfect residue field of characteristic p .

We denote the n -dimensional unit polydisk $\mathrm{Spa}(K\langle T_1, \dots, T_n \rangle, \mathcal{O}_K\langle T_1, \dots, T_n \rangle)$ by \mathbb{B}_K^n . We also write \mathbb{T}_K^n for $\mathrm{Spa}(K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle, \mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle)$, the n -dimensional rigid torus.

Let X be a smooth adic space over $\mathrm{Spa}(K, \mathcal{O}_K)$. We call a point x of X *classical* if the residue class field $k(x)$ of x is finite over K .

We work on the étale site $X_{\mathrm{\acute{e}t}}$ (cf. [Hub96, Definition 2.3.1]) and the pro-étale site $X_{\mathrm{pro\acute{e}t}}$. For the pro-étale site, we use the one defined in [Sch13, Sch16], and thus we have the projection

$$\nu: X_{\mathrm{pro\acute{e}t}} \rightarrow X_{\mathrm{\acute{e}t}}.$$

Recall that $U \in X_{\mathrm{pro\acute{e}t}}$ is called *affinoid perfectoid* if U has a pro-étale presentation $U = \varprojlim U_i \rightarrow X$ by affinoid $U_i = \mathrm{Spa}(A_i, A_i^+)$ such that, denoting by A^+ the p -adic completion of $\varinjlim A_i^+$, and $A = A^+[p^{-1}]$, the pair (A, A^+) is a perfectoid affinoid (L, L^+) -algebra for a perfectoid field L over K and an open and bounded valuation subring $L^+ \subset L$ (cf. [Sch13, Definition 4.3 (i)]). In this case, we write \widehat{U} for $\mathrm{Spa}(A, A^+)$, which is independent of the pro-étale presentation $U = \varprojlim U_i$.

8.1. p -adic local systems on $X_{\mathrm{\acute{e}t}}$. Let us clarify our terminology of local systems.

Definition 8.1 (cf. [Sch13, Definition 8.1], [KL15, §1.4, §8.4]). A *lisse \mathbb{Z}_p -sheaf* $\mathbb{L} = (\mathbb{L}_n)$ on $X_{\mathrm{\acute{e}t}}$ is an inverse system of sheaves of \mathbb{Z}/p^n -modules \mathbb{L}_n on $X_{\mathrm{\acute{e}t}}$ such that each \mathbb{L}_n is locally on $X_{\mathrm{\acute{e}t}}$ a constant sheaf associated with a finitely generated flat \mathbb{Z}/p^n -module and such that this inverse system is isomorphic in the procategory to an inverse system for which $\mathbb{L}_{n+1}/p^n \cong \mathbb{L}_n$. We also call a lisse \mathbb{Z}_p -sheaf on $X_{\mathrm{\acute{e}t}}$ an *étale \mathbb{Z}_p -local system* on X or a \mathbb{Z}_p -local system on $X_{\mathrm{\acute{e}t}}$.

Definition 8.2 (cf. [KL15, §1.4, §8.4]). An *isogeny \mathbb{Z}_p -local system* on $X_{\mathrm{\acute{e}t}}$ is an object of the isogeny category of \mathbb{Z}_p -local systems on $X_{\mathrm{\acute{e}t}}$. A *\mathbb{Q}_p -local system* on $X_{\mathrm{\acute{e}t}}$ is an object of the stack associated to the fibered category of isogeny \mathbb{Z}_p -local

systems on $X_{\text{ét}}$. We also call a \mathbb{Q}_p -local system on $X_{\text{ét}}$ an *étale \mathbb{Q}_p -local system* on X .

Remark 8.3. Assume that $X = \text{Spa}(A, A^+)$ is an affinoid. In this case, the category of adic spaces which are finite and étale over $\text{Spa}(A, A^+)$ is canonically equivalent to the category of schemes which are finite and étale over $\text{Spec } A$ by [Hub96, Example 1.6.6 ii)]. The natural functor from the category of lisse \mathbb{Z}_p -sheaves on the étale site $(\text{Spec } A)_{\text{ét}}$ to the category of \mathbb{Z}_p -local systems on $\text{Spa}(A, A^+)_{\text{ét}}$ is an equivalence of categories (cf. [KL15, Remark 8.4.5]).

Moreover, if $\text{Spec } A$ is connected and \bar{x} is a geometric point of $\text{Spec } A$, we have the étale fundamental group $\pi_1^{\text{ét}}(\text{Spec } A, \bar{x})$. Then the category of continuous representations of $\pi_1^{\text{ét}}(\text{Spec } A, \bar{x})$ in finite free \mathbb{Z}_p -modules is equivalent to the category of \mathbb{Z}_p -local systems on $\text{Spa}(A, A^+)_{\text{ét}}$. Similarly, the category of continuous representations of $\pi_1^{\text{ét}}(\text{Spec } A, \bar{x})$ in finite-dimensional \mathbb{Q}_p -vector spaces is equivalent to the category of isogeny \mathbb{Z}_p -local systems on $\text{Spa}(A, A^+)_{\text{ét}}$ (cf. [KL15, Remark 1.4.4]).

In particular, if A^+ satisfies conditions in Set-up 3.1 relative to K , then the category $\text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_A)$ is equivalent to the category of isogeny \mathbb{Z}_p -local systems on $\text{Spa}(A, A^+)_{\text{ét}}$.

Definition 8.4 ([Sch13, Definition 8.1]). Let $\widehat{\mathbb{Z}}_p = \varprojlim \mathbb{Z}/p^n$ as sheaves on $X_{\text{proét}}$. Then a lisse $\widehat{\mathbb{Z}}_p$ -sheaf on $X_{\text{proét}}$ is a sheaf of $\widehat{\mathbb{Z}}_p$ -modules on $X_{\text{proét}}$ such that locally in $X_{\text{proét}}$, it is isomorphic to $\widehat{\mathbb{Z}}_p \otimes_{\mathbb{Z}_p} M$, where M is a finitely generated \mathbb{Z}_p -module.

Similarly, set $\widehat{\mathbb{Q}}_p = \widehat{\mathbb{Z}}_p[p^{-1}]$. A lisse $\widehat{\mathbb{Q}}_p$ -sheaf on $X_{\text{proét}}$ is a sheaf of $\widehat{\mathbb{Q}}_p$ -modules on $X_{\text{proét}}$ such that locally in $X_{\text{proét}}$, it is isomorphic to $\widehat{\mathbb{Q}}_p \otimes_{\mathbb{Q}_p} M$, where M is a finite-dimensional \mathbb{Q}_p -vector space.

For a \mathbb{Z}_p -local system $\mathbb{L} = (\mathbb{L}_n)$ on $X_{\text{ét}}$, we set

$$\widehat{\mathbb{L}} = \varprojlim \nu^* \mathbb{L}_n.$$

Then $\widehat{\mathbb{L}}$ is a lisse $\widehat{\mathbb{Z}}_p$ -sheaf on $X_{\text{proét}}$. This functor is an equivalence of categories. Moreover, $R^j \varprojlim \nu^* \mathbb{L}_n = 0$ for $j > 0$ by [Sch13, Proposition 8.2]. Similarly, we can naturally associate to every \mathbb{Q}_p -local system \mathbb{L} on $X_{\text{ét}}$ a lisse $\widehat{\mathbb{Q}}_p$ -sheaf on $X_{\text{proét}}$, which we denote by $\widehat{\mathbb{L}}$.

8.2. Period sheaves and the Riemann–Hilbert functor D_{dR} . We will use the following sheaves on $X_{\text{proét}}$:

- The de Rham sheaf \mathbb{B}_{dR} (cf. [Sch13, Definition 6.1(iii)]). By [Sch13, Theorem 6.5(i)], for an affinoid perfectoid $U \in X_{\text{proét}}$ with $\widehat{U} = \text{Spa}(\Lambda, \Lambda^+)$, we have

$$\mathbb{B}_{\text{dR}}(U) = \mathbb{B}_{\text{dR}}(\Lambda, \Lambda^+),$$

where the latter is the period ring defined in Section 2, which agrees with the one defined in [Sch13, page 49].

- The structural de Rham sheaf $\mathcal{O}\mathbb{B}_{\text{dR}}$ (cf. [Sch13, Sch16]). This sheaf is a \mathbb{B}_{dR} -algebra equipped with a filtration and a \mathbb{B}_{dR} -linear connection

$$\nabla: \mathcal{O}\mathbb{B}_{\text{dR}} \rightarrow \mathcal{O}\mathbb{B}_{\text{dR}} \otimes_{\mathcal{O}_X} \Omega_X^1$$

satisfying the Griffiths transversality and $(\mathcal{O}\mathbb{B}_{\text{dR}})^{\nabla=0} = \mathbb{B}_{\text{dR}}$.

We have the following description of $\mathcal{O}\mathbb{B}_{\mathrm{dR}}$: Let $\mathrm{Spa}(A, A^+)$ be an affinoid admitting standard étale maps to both X and \mathbb{T}_L^n for some finite extension L over K . If we replace L with its algebraic closure in A , then $R = A^+$ satisfies conditions in Set-up 3.1 relative to L . Write $\overline{R} = \varinjlim_{i \in I} A_i^+$ where A_i^+ is a finite normal R -subalgebra of $\overline{\mathrm{Frac}} \overline{R}$ such that $A_i = A_i^+[p^{-1}]$ is étale over $A = R_K$. Set $U := \varprojlim_i \mathrm{Spa}(A_i, A_i^+) \in X_{\mathrm{proét}}$; this is affinoid perfectoid with $\widehat{U} = \mathrm{Spa}(\widehat{R}[p^{-1}], \widehat{R})$. In this setting, $\mathbb{B}_{\mathrm{dR}}(U) = \mathbb{B}_{\mathrm{dR}}(\widehat{R}) = \mathbb{B}_{\mathrm{dR}}^\nabla(R)$, and there is a natural $\mathbb{B}_{\mathrm{dR}}^\nabla(R)$ -linear isomorphism

$$\mathcal{O}\mathbb{B}_{\mathrm{dR}}(U) \cong \mathbb{B}_{\mathrm{dR}}(R),$$

compatible with filtrations and connections, where the right hand side is the de Rham period ring recalled in Subsection 4.4. To see this, note that $\mathcal{O}\mathbb{B}_{\mathrm{dR}}$ is defined to be the sheafification of the presheaf defined in [Sch16, (3)]. From the definitions, we have a $\mathbb{B}_{\mathrm{dR}}^\nabla(R)$ -algebra homomorphism $\mathbb{B}_{\mathrm{dR}}(R) \rightarrow \mathcal{O}\mathbb{B}_{\mathrm{dR}}(U)$. By comparing [Sch13, Proposition 6.10], [Sch16, (3)], and [Bri08, Proposition 5.2.2], we conclude that this map is an isomorphism compatible with filtrations and connections.

Lemma 8.5. *Let $U \in X_{\mathrm{proét}}$ be affinoid perfectoid. Then $H^q(U, \mathcal{O}\mathbb{B}_{\mathrm{dR}}) = 0$ for every $q > 0$.*

Proof. Consider the positive structural de Rham sheaf $\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ ([Sch13, Definition 6.8 (iii)]). It is equipped with a decreasing filtration $\mathrm{Fil}^\bullet \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ of $\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ defined by $\mathrm{Fil}^i \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ = (\mathrm{Ker} \theta)^i$. Then $\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ = \varprojlim_i \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ / \mathrm{Fil}^i \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ and $\mathrm{gr}^\bullet \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ is naturally a polynomial algebra over $\widehat{\mathcal{O}}_X$ by [Sch13, Proposition 6.10].

For every affinoid perfectoid U , we have $H^q(U, \widehat{\mathcal{O}}_X) = 0$ for $q > 0$ by [Sch13, Lemma 4.10 (v)]. Hence $H^q(U, \mathrm{Fil}^i \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ / \mathrm{Fil}^{i+1} \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+) = 0$ for $q > 0$ by coherence of U ([Sch13, Proposition 3.12 (iii)]). It follows that $H^0(U, \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ / \mathrm{Fil}^{i+1} \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+) \rightarrow H^0(U, \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+ / \mathrm{Fil}^i \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+)$ is surjective. By [Sch13, Lemma 3.18, Proposition 4.8], we have $H^q(U, \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+) = 0$ for every $q > 0$. Since $\mathcal{O}\mathbb{B}_{\mathrm{dR}} = \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+[t^{-1}]$ is written as $\varinjlim \mathcal{O}\mathbb{B}_{\mathrm{dR}}^+$ with multiplication-by- t transition maps, we conclude $H^q(U, \mathcal{O}\mathbb{B}_{\mathrm{dR}}) = 0$ for every $q > 0$ by coherence of U . \square

Definition 8.6 (cf. [Sch13, Definition 8.3]). Let \mathbb{L} be a \mathbb{Q}_p -local system on $X_{\mathrm{ét}}$. We say that \mathbb{L} is *de Rham* if there exists a filtered \mathcal{O}_X -module \mathcal{E} with an integrable connection such that there is an isomorphism of sheaves on $X_{\mathrm{proét}}$

$$\widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}} \cong \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\mathrm{dR}}$$

compatible with filtrations and connections. Similarly, we say that a(n isogeny) \mathbb{Z}_p -local system on $X_{\mathrm{ét}}$ is de Rham if the associated \mathbb{Q}_p -local system is de Rham.

Based on the work of Kedlaya–Liu and Scholze, Liu and Zhu proved the following theorem:

Theorem 8.7 ([LZ17, Theorem 3.9]). *For a \mathbb{Q}_p -local system \mathbb{L} on $X_{\mathrm{ét}}$, set*

$$D_{\mathrm{dR}}(\mathbb{L}) := \nu_*(\widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}}).$$

- (i) $D_{\mathrm{dR}}(\mathbb{L})$ is a vector bundle on $X_{\mathrm{ét}}$ equipped with an integrable connection

$$\nabla_{\mathbb{L}} := \nu_*(\mathrm{id} \otimes \nabla): D_{\mathrm{dR}}(\mathbb{L}) \rightarrow D_{\mathrm{dR}}(\mathbb{L}) \otimes_{\mathcal{O}_{X_{\mathrm{ét}}}} \Omega_{X_{\mathrm{ét}}}$$

and a decreasing filtration Fil^\bullet satisfying the Griffiths transversality.

- (ii) The functor $\mathbb{L} \mapsto (D_{\mathrm{dR}}(\mathbb{L}), \nabla_{\mathbb{L}}, \mathrm{Fil}^\bullet)$ commutes with pullbacks.
- (iii) The following conditions are equivalent for \mathbb{L} :
 - (a) \mathbb{L} is de Rham;
 - (b) $\mathrm{rank} D_{\mathrm{dR}}(\mathbb{L}) = \mathrm{rank} \mathbb{L}$;
 - (c) for each connected component, there exists a classical point x of X and a geometric point \bar{x} above x such that the stalk $\mathbb{L}_{\bar{x}}$ is a de Rham representation of the absolute Galois group of $k(x)$.

In this case, the natural map $\nu^* D_{\mathrm{dR}}(\mathbb{L}) \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\mathrm{dR}} \rightarrow \widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}}$ is an isomorphism of sheaves on $X_{\mathrm{pro\acute{e}t}}$ compatible with filtrations and connections.

Remark 8.8. The category of vector bundles on X (with the analytic topology) is equivalent to the category of vector bundles on $X_{\acute{e}t}$ by [Sch13, Lemma 7.3]. Hence we also regard $D_{\mathrm{dR}}(\mathbb{L})$ as a vector bundle on X .

Lemma 8.9. *Assume that $X = \mathrm{Spa}(A, A^+)$ is affinoid such that $R = A^+$ satisfies conditions in Set-up 3.1 relative to K and it is equipped with the p -adic topology. Let \mathbb{L} be an isogeny \mathbb{Z}_p -local system on $X_{\acute{e}t}$ and let $V \in \mathrm{Rep}_{\mathbb{Q}_p}(\mathcal{G}_A)$ be the corresponding representation. Then the vector bundle $D_{\mathrm{dR}}(\mathbb{L})$ is associated to the A -module $D_{\mathrm{dR}}(V)$ and this identification is compatible with connections and filtrations. Moreover, \mathbb{L} is de Rham if and only if V is a de Rham representation.*

Proof. Since $D_{\mathrm{dR}}(\mathbb{L})$ is a coherent sheaf on X , it suffices to show that $\Gamma(X, D_{\mathrm{dR}}(\mathbb{L}))$ is naturally identified with $D_{\mathrm{dR}}(V)$. By definition, we have

$$\Gamma(X, D_{\mathrm{dR}}(\mathbb{L})) = \Gamma(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}}).$$

Write $\overline{R} = \varinjlim_{i \in I} A_i^+$ where A_i^+ is a finite normal R -subalgebra of $\overline{\mathrm{Frac} R}$ such that $A_i = A_i^+[p^{-1}]$ is étale over $A = R_K$. Set $U := \varprojlim_i \mathrm{Spa}(A_i, A_i^+) \in X_{\mathrm{pro\acute{e}t}}$; this is affinoid perfectoid with $\widehat{U} = \mathrm{Spa}(\widehat{R}[p^{-1}], \widehat{R})$. Since every finite étale A -algebra splits over $\overline{R}[p^{-1}]$, the lisse $\widehat{\mathbb{Q}}_p$ -sheaf $\widehat{\mathbb{L}}$ is trivial over U and naturally identified with $\widehat{\mathbb{Q}}_p \otimes_{\mathbb{Q}_p} V$.

For each $i \in \mathbb{N}$, the i -fold fiber product $U^{i/X} := U \times_X U \times \cdots \times_X U$ of U over X exists and is affinoid perfectoid. By Lemma 8.5, we have $H^q(U^{i/X}, V \otimes_{\mathbb{Q}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}}) = 0$ for every $q > 0$ and $i \in \mathbb{N}$. Applying the Cartan–Leray spectral sequence to the cover $U \rightarrow X$ in $X_{\mathrm{pro\acute{e}t}}$, we compute

$$\begin{aligned} \Gamma(X, D_{\mathrm{dR}}(\mathbb{L})) &= H_{\mathrm{cont}}^0(\mathcal{G}_A, \Gamma(U, \widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}})) = H_{\mathrm{cont}}^0(\mathcal{G}_A, \Gamma(U, V \otimes_{\mathbb{Q}_p} \mathrm{B}_{\mathrm{dR}}(R))) \\ &= D_{\mathrm{dR}}(V). \end{aligned}$$

It is easy to see that this identification is compatible with connections and filtrations.

If V is a de Rham representation, then $D_{\mathrm{dR}}(V)$ is a finite projective A -module of rank equal to $\dim_{\mathbb{Q}_p} V = \mathrm{rank} \mathbb{L}$. Hence \mathbb{L} is de Rham. Conversely, if \mathbb{L} is de Rham, we have an isomorphism $\nu^* D_{\mathrm{dR}}(\mathbb{L}) \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\mathrm{dR}} \rightarrow \widehat{\mathbb{L}} \otimes_{\widehat{\mathbb{Q}}_p} \mathcal{O}\mathbb{B}_{\mathrm{dR}}$ of sheaves on $X_{\mathrm{pro\acute{e}t}}$.

Evaluating both sides at $U \in X_{\mathrm{pro\acute{e}t}}$ yields $D_{\mathrm{dR}}(V) \otimes_A \mathrm{B}_{\mathrm{dR}}(R) \xrightarrow{\cong} V \otimes_{\mathbb{Q}_p} \mathrm{B}_{\mathrm{dR}}(R)$ and this is identified with $\alpha_{\mathrm{dR}}(V)$. Hence V is a de Rham representation. \square

9. A p -ADIC MONODROMY THEOREM FOR DE RHAM LOCAL SYSTEMS

Let K be a complete discrete valuation field of characteristic zero with perfect residue field of characteristic p . Recall that a rigid analytic variety over K refers to a quasi-separated adic space locally of finite type over $\mathrm{Spa}(K, \mathcal{O}_K)$ (cf. [Hub94, Proposition 4.5(iv)]). For a rigid analytic variety X over K and a complete extension L of K , we write X_L for $X \times_{\mathrm{Spa}(K, \mathcal{O}_K)} \mathrm{Spa}(L, \mathcal{O}_L)$.

9.1. Statement of the theorem.

Definition 9.1. Let X be a smooth rigid analytic variety over K , and let \mathbb{L} be an isogeny \mathbb{Z}_p -local system on $X_{\text{ét}}$.

- (i) We say that \mathbb{L} is *horizontal de Rham* if \mathbb{L} is de Rham and $(D_{\text{dR}}(\mathbb{L}), \nabla)$ has a full set of solutions, i.e., $(D_{\text{dR}}(\mathbb{L}), \nabla) \cong (\mathcal{O}_X, d)^{\text{rank } \mathbb{L}}$.
- (ii) Let x be a classical point of X . We say that \mathbb{L} is *semistable* (resp. *potentially crystalline*) at x if there exists a geometric point \bar{x} above x such that the stalk $\mathbb{L}_{\bar{x}}$ is a semistable (resp. (potentially) crystalline) representation of the absolute Galois group of $k(x)$. This is equivalent to the condition that the stalk $\mathbb{L}_{\bar{x}}$ is a semistable (resp. (potentially) crystalline) representation for every geometric point \bar{x} above x .

Theorem 1.1 in the introduction is a special case of the following p -adic monodromy theorem for de Rham local systems:

Theorem 9.2. *Let X be a smooth rigid analytic variety over K , and let \mathbb{L} be a de Rham isogeny \mathbb{Z}_p -local system on $X_{\text{ét}}$. Then for every classical point $x \in X$, there exists an open neighborhood $U \subset X_{k(x)}$ of x and a finite extension L of $k(x)$ such that $\mathbb{L}|_{U_L}$ is horizontal de Rham and semistable at every classical point. Moreover, if $\mathbb{L}_{\bar{x}}$ is potentially crystalline, then U and L can be chosen in such a way that $\mathbb{L}|_{U_L}$ is horizontal de Rham and crystalline at every classical point.*

Let us first make a remark and discuss its consequence.

Remark 9.3. If x runs over all the classical points of X , the images in X of U 's in the theorem form a collection of open subsets of X . In general, the union of such open subsets may not be the entire X .

Corollary 9.4. *Let X be a quasi-compact smooth rigid analytic variety over K , and let \mathbb{L} be a de Rham \mathbb{Q}_p -local system on $X_{\text{ét}}$. Then there exists a finite extension L of K such that for every K -rational point $x \in X$ and every geometric point \bar{x} above x , the representation $\mathbb{L}_{\bar{x}}$ of the absolute Galois group of $k(x)$ becomes semistable over L .*

Proof of Corollary 9.4. Note that the set of K -valued points of X is compact with respect to the analytic topology; the general case is reduced to the case where $X = \mathbb{B}_K^n$, and the assertion is obvious in this case. Since \mathbb{L} is étale locally an isogeny \mathbb{Z}_p -local system and étale maps are open, the corollary follows from Theorem 9.2 and the compactness of the set of K -valued points of X . \square

We prove Theorem 9.2 in the rest of this section. For this, we show that $(D_{\text{dR}}(\mathbb{L}), \nabla)$ is locally horizontal de Rham. Then the assertion follows from Theorems 7.4 and 7.5.

9.2. Preliminaries on smooth rigid analytic varieties. The following lemma is easy to prove and we omit the proof.

Lemma 9.5. *Let $x \in \mathbb{B}_K^n$ be a point with residue class field K .*

- (i) *There exists a $\mathrm{Spa}(K, \mathcal{O}_K)$ -isomorphism*

$$\mathbb{B}_K^n \rightarrow \mathbb{B}_K^n$$

sending x to 0.

- (ii) *For every open neighborhood V of x in \mathbb{B}_K^n , there exists an open immersion of the form*

$$\mathbb{B}_K^n \hookrightarrow V$$

sending x to 0.

Proposition 9.6. *Let X be an n -dimensional smooth rigid analytic variety over K . Let L be a finite extension of K and $x \in X$ a classical point with residue class field L . Then there exists an open immersion*

$$\mathbb{B}_L^n \rightarrow X_L$$

sending $0 \in \mathbb{B}_L^n$ to $x \in X_L$.

Proof. By replacing X by X_L , we may assume $L = K$. By shrinking X if necessary, we may assume that the structure morphism $X \rightarrow \mathrm{Spa}(K, \mathcal{O}_K)$ factors as

$$X \xrightarrow{f} \mathbb{B}_K^n \rightarrow \mathrm{Spa}(K, \mathcal{O}_K),$$

where f is étale and the second is the structure morphism ([Hub96, Corollary 1.6.10]). We may further assume that $f(x) = 0$ by Lemma 9.5(i).

By [Hub96, Corollary 1.7.4, Lemma 1.6.4], the map of local rings

$$f^*: \mathcal{O}_{\mathbb{B}_K^n, 0} \rightarrow \mathcal{O}_{X, x}$$

is finite with the same residue class field K . Recall that $\mathcal{O}_{\mathbb{B}_K^n, 0}$ is Henselian (cf. [Hou61, p.9, Corollaire 2], [Ber93, Theorem 2.1.5]). Hence we conclude that $\mathcal{O}_{\mathbb{B}_K^n, 0} \cong \mathcal{O}_{X, x}$.

It follows that there exist an affinoid open neighborhood U of $x \in X$, an affinoid open neighborhood V of $0 \in \mathbb{B}_K^n$ and a morphism $g: V \rightarrow U$ such that $f \circ g$ is the inclusion $V \subset \mathbb{B}_K^n$. Note that g is injective and that $g^*: \mathcal{O}_{U, x} \rightarrow \mathcal{O}_{V, 0}$ is an isomorphism. After shrinking V and U if necessary, we may further assume that g is an isomorphism by [BGR84, 7.3.3 Proposition 5]. By Lemma 9.5(ii), we get an open immersion

$$\mathbb{B}_K^n \rightarrow V \xrightarrow{g} U \rightarrow X$$

sending 0 to x . □

9.3. Vector bundles with integrable connections on the polydisk \mathbb{B}_K^n . Let $A := K\langle T_1, \dots, T_n \rangle$ and let $d: A \rightarrow \Omega_A$ denote the universal K -derivation of A ([Hub96, Definition 1.6.1]). Concretely, Ω_A is a finite free A -module of rank n with generators dT_1, \dots, dT_n , and $d: A \rightarrow \Omega_A$ sends f to $df = \sum_{i=1}^n \frac{\partial f}{\partial T_i} dT_i$ for $f \in A$ ([Hub96, 1.6.2]). In other words, $\Omega_A = A \otimes_{A^\circ} \widehat{\Omega}_{A^\circ}$ (cf. Subsection 4.4).

The following result will be well known to experts. For example, see [DGS94, Appendix III], [Ked10a, Proposition 9.3.3] for the case $n = 1$.

Theorem 9.7. *Let (M, ∇) be a finite free A -module equipped with an integrable connection. Then there exists $l \in \mathbb{N}$ such that the induced connection on*

$$M \otimes_A K\langle T_1/p^l, \dots, T_n/p^l \rangle$$

has a full set of solutions, i.e., it is isomorphic to $(K\langle T_1/p^l, \dots, T_n/p^l \rangle, d)^{\text{rank}_A M}$.

Proof. Consider the natural inclusion

$$M \subset M \otimes_A K[[T_1, \dots, T_n]]$$

and extend the integrable connection to the latter. By a classical result (cf. [Kat70, Proposition 8.9]), we know that $M \otimes_A K[[T_1, \dots, T_n]]$ has a full set of solutions. In fact, we can construct an explicit K -linear surjection

$$P: M \rightarrow (M \otimes_A K[[T_1, \dots, T_n]])^{\nabla=0}$$

using the Taylor expansion, which we now explain.

For each $i = 1, \dots, n$, we set $D_i := \nabla_{\frac{\partial}{\partial T_i}}: M \rightarrow M$. Then the integrability of ∇ is equivalent to $D_i \circ D_{i'} = D_{i'} \circ D_i$ for every $i, i' = 1, \dots, n$. For each $\underline{j} = (j_1, \dots, j_n) \in \mathbb{N}^n$, we set

$$|\underline{j}| = j_1 + \dots + j_n, \quad \underline{j}! := (j_1!) \cdots (j_n!), \quad T^{\underline{j}} := T_1^{j_1} \cdots T_n^{j_n},$$

and

$$D_{\underline{j}} := \prod_{i=1}^n (D_i)^{j_i}: M \rightarrow M.$$

We define $P: M \rightarrow M \otimes_A K[[T_1, \dots, T_n]]$ by

$$P(f) := \sum_{\underline{j} \in \mathbb{N}^n} \frac{(-1)^{|\underline{j}|}}{\underline{j}!} T^{\underline{j}} D_{\underline{j}}(f).$$

Then we can check $P(f)(0, \dots, 0) = f(0, \dots, 0)$, $\text{Im } P \subset (M \otimes_A K[[T_1, \dots, T_n]])^{\nabla=0}$, $\text{Ker } P = (T_1, \dots, T_n)M$, and that P induces an isomorphism

$$P: M/(T_1, \dots, T_n) \xrightarrow{\cong} (M \otimes_A K[[T_1, \dots, T_n]])^{\nabla=0}.$$

Let $r = \text{rank}_A M$ and choose an A -basis of M so that we identify $M = A^r$. For each $f \in A^r$, we can uniquely write $P(f)$ as

$$P(f) = \sum_{\underline{j} \in \mathbb{N}^n} A_{\underline{j}}(f) T^{\underline{j}}, \quad A_{\underline{j}}(f) \in K^r.$$

More concretely, we see

$$A_{\underline{j}}(f) = \left(\left(\frac{\partial}{\partial T} \right)^{\underline{j}} (P(f)) \right) \Big|_{(0, \dots, 0)} = \left(\left(\frac{\partial}{\partial T} \right)^{\underline{j}} \left(\sum_{\underline{j}' \leq \underline{j}} \frac{(-1)^{|\underline{j}'|}}{\underline{j}'!} T^{\underline{j}'} D_{\underline{j}'}(f) \right) \right) \Big|_{(0, \dots, 0)},$$

where $\left(\frac{\partial}{\partial T} \right)^{\underline{j}} := \left(\frac{\partial}{\partial T_1} \right)^{j_1} \cdots \left(\frac{\partial}{\partial T_n} \right)^{j_n}$ for $\underline{j} = (j_1, \dots, j_n)$, and $\underline{j}' \leq \underline{j}$ refers to the inequality entrywise. In this way, we get a K -linear map $A_{\underline{j}}: A^r \rightarrow K^r$.

Consider the Gauss norm on A and equip $A^r (= M)$, $M_r(A)$ with the norm defined by the maximum of the Gauss norm of each entry. Similarly, equip K^r with the p -adic norm. We normalize these norms so that the norm of p is p^{-1} . We will estimate the operator norm $\|A_{\underline{j}}\|$. For each $i = 1, \dots, n$, write

$$D_i = \frac{\partial}{\partial T_i} + B_i, \quad B_i \in M_r(A).$$

Choose $C' > 0$ such that $\|B_i\| \leq C'$ for $i = 1, \dots, n$. Set $C = \max\{1, C'\}$. Note that both $\frac{\partial}{\partial T_i}: A^r \rightarrow A^r$ and the multiplication by T_i on A^r have operator norm 1 and that the p -adic norm of $\underline{j}!$ is greater than or equal to $(\frac{1}{p^{1/(p-1)}})^{|\underline{j}|}$. It follows that

$$\left\| \frac{(-1)^{|\underline{j}|}}{\underline{j}!} T^{\underline{j}} D_{\underline{j}} \right\| \leq C^{|\underline{j}|} p^{|\underline{j}|/(p-1)},$$

and thus

$$\|A_{\underline{j}}\| = \left\| \left(\left(\frac{\partial}{\partial T} \right)^{\underline{j}} \left(\sum_{\underline{j}' \leq \underline{j}} \frac{(-1)^{|\underline{j}'|}}{\underline{j}'!} T^{\underline{j}'} D_{\underline{j}'} \right) \right) \right\|_{(0, \dots, 0)} \leq C^{|\underline{j}|} p^{|\underline{j}|/(p-1)}.$$

Choose $l \in \mathbb{N}$ such that

$$C p^{1/(p-1)} p^{-l} < 1.$$

By the estimate on $\|A_{\underline{j}}\|$, we have $\text{Im } P \subset M \otimes_A K \langle T_1/p^l, \dots, T_n/p^l \rangle$. In other words, we have

$$(M \otimes_A K[[T_1, \dots, T_n]])^{\nabla=0} \subset M \otimes_A K \langle T_1/p^l, \dots, T_n/p^l \rangle.$$

We again identify $M = A^r$ and $M \otimes_A K \langle T_1/p^l, \dots, T_n/p^l \rangle = K \langle T_1/p^l, \dots, T_n/p^l \rangle^r$. Choose

$$s_1 = (s_{11}, \dots, s_{1r}), \dots, s_r = (s_{r1}, \dots, s_{rr}) \in M \otimes_A K \langle T_1/p^l, \dots, T_n/p^l \rangle$$

such that s_1, \dots, s_r form a K -basis of $(M \otimes_A K[[T_1, \dots, T_n]])^{\nabla=0}$ and such that the values at $(T_1, \dots, T_n) = (0, \dots, 0)$ satisfy

$$s_1(0, \dots, 0) = (1, 0, \dots, 0), \dots, s_r(0, \dots, 0) = (0, \dots, 0, 1).$$

Set

$$s := \det(s_{ij})_{i,j=1, \dots, r} \in K \langle T_1/p^l, \dots, T_n/p^l \rangle.$$

Then s satisfies $s(0, \dots, 0) = 1 \neq 0$. In particular, there exists $l' \geq l$ such that s is invertible in $K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle$.

Let us consider the induced connection ∇ on $M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle$. Since $s_1, \dots, s_r \in M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle$, we have

$$(M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle)^{\nabla=0} \subset M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle.$$

Moreover, since s is invertible in $K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle$, we conclude that the natural map

$$(M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle)^{\nabla=0} \otimes_K K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle \rightarrow M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle$$

is an isomorphism. This means that $(M \otimes_A K \langle T_1/p^{l'}, \dots, T_n/p^{l'} \rangle, \nabla)$ has a full set of solutions. \square

Remark 9.8. Note that A is a UFD by [BGR84, 5.2.6 Theorem 1]. Hence any finite projective A -module is finite free.

9.4. Proof of Theorem 9.2. We prove Theorem 9.2. By Proposition 9.6 and Theorem 9.7, there exists an open neighborhood $U \subset X_{k(x)}$ of x such that $U \cong \mathbb{B}_{k(x)}^n$ and $(D_{\text{dR}}(\mathbb{L})|_U, \nabla)$ has a full set of solutions. Hence $\mathbb{L}|_U$ is horizontal de Rham because $D_{\text{dR}}(\mathbb{L})|_U = D_{\text{dR}}(\mathbb{L}|_U)$. By shrinking U again, we may assume that U is isomorphic to $\mathbb{T}_{k(x)}^n$. Write $\mathbb{T}_{k(x)}^n = \text{Spa}(A, A^\circ)$. Then $R = A^\circ$ satisfies the conditions in Set-up 3.1 relative to $k(x)$. Let $V \in \text{Rep}_{\mathbb{Q}_p}(\mathcal{G}_A)$ be the representation corresponding to $\mathbb{L}|_U$. Then V is horizontal de Rham by Lemma 8.9. By Theorem 7.4, there exists a finite extension L of $k(x)$ such that $V|_{\mathcal{G}_{A_L}}$ is horizontal semistable. In particular, $\mathbb{L}|_{U_L}$ is horizontal de Rham and semistable at every classical point. The second assertion follows from Theorem 7.5.

10. APPLICATIONS: POTENTIALLY CRYSTALLINE LOCI AND POTENTIALLY GOOD REDUCTION LOCI

As an application of our p -adic monodromy theorem, we discuss potentially crystalline loci of de Rham local systems and cohomologically potentially good reduction loci of smooth proper families of relative dimension at most two.

Let K be a complete discrete valuation field of characteristic zero with perfect residue field of characteristic p . In Subsections 10.2 and 10.3, we will further assume that the residue field of K is finite.

10.1. Potentially crystalline loci.

Definition 10.1. Let X be a smooth rigid analytic variety over K , and let \mathbb{L} be a de Rham \mathbb{Q}_p -local system on $X_{\text{ét}}$. We say that an open subset U of X is a *potentially crystalline locus* of \mathbb{L} if U satisfies the following property: for every classical point x of X ,

- (i) if $x \in U$, then \mathbb{L} is potentially crystalline at x ;
- (ii) if $x \notin U$, then \mathbb{L} is not potentially crystalline at x .

Theorem 10.2. *For every smooth rigid analytic variety X over K and every de Rham \mathbb{Q}_p -local system \mathbb{L} on $X_{\text{ét}}$, there exists a potentially crystalline locus of \mathbb{L} .*

Proof. Observe that if there exists an étale covering $\{X_i\}_i$ of X such that each $\mathbb{L}|_{X_i}$ admits a potentially crystalline locus $U_i \subset X_i$, then the union of the images of U_i 's in X is a potentially crystalline locus of X . By passing to an étale cover of X , we may assume that \mathbb{L} is a de Rham isogeny \mathbb{Z}_p -local system on $X_{\text{ét}}$.

Let x be a classical point of X such that \mathbb{L} is potentially crystalline at x . By Theorem 9.2, there exists an open neighborhood $U'_x \subset X_{k(x)}$ of x such that $\mathbb{L}|_{U'_x}$ is potentially crystalline at every classical point. Let U_x denote the image of U'_x in X ; it is an open subset of X . Then the union of such U_x 's satisfies the desired properties. \square

Remark 10.3. Our definition takes only classical rank 1 points into consideration, and thus a potentially crystalline locus of \mathbb{L} is not unique. On the other hand, the union of potentially crystalline loci is also a potentially crystalline locus and thus we have the largest potentially crystalline locus.

10.2. The monodromy-weight conjecture and ℓ -independence. We will discuss cohomologically potentially good reduction loci of smooth proper maps in the next subsection. For this, let us briefly review basic results and conjectures on cohomologies of smooth proper varieties over p -adic fields.

Let K be a complete discrete valuation field of characteristic zero with *finite* residue field k of characteristic p . Set $\text{Gal}_K := \text{Gal}(\overline{K}/K)$. Let $q = p^h$ denote the cardinality of k . Set $P_0 := W(\overline{k})[p^{-1}]$. It is equipped with an action of Gal_K . Write $\sigma: P_0 \rightarrow P_0$ for the Witt vector Frobenius lifting $x \mapsto x^p$.

Let I_K denote the inertia subgroup of Gal_K and let W_K denote the Weil group of K ; they fit in the following exact sequence:

$$0 \longrightarrow I_K \longrightarrow W_K \xrightarrow{v} h\mathbb{Z} \longrightarrow 0,$$

where v is normalized so that $h \in h\mathbb{Z} \subset h\widehat{\mathbb{Z}} = \text{Gal}_k$ is the geometric Frobenius $a \mapsto a^{-q}$ of k . Set $W_K^+ := \{g \in W_K \mid v(g) \geq 0\}$.

Let ℓ be a prime. For $\ell \neq p$, let $t_\ell: I_K \rightarrow \mathbb{Z}_\ell(1)$ be the map sending $g \in I_K$ to $(g(\pi^{\ell-m})/\pi^{\ell-m})_m \in \mathbb{Z}_\ell(1)$, where π is a uniformizer of K ; the map t_ℓ is independent of the choice of a uniformizer π nor its ℓ -power roots.

Let X be a smooth proper scheme purely of dimension n over K . For each ℓ , the ℓ -adic étale cohomology $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$ is an ℓ -adic representation of Gal_K .

Let us first assume $\ell \neq p$. By Grothendieck's monodromy theorem, there exist a nilpotent endomorphism $N \in \text{End}(H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)(-1))$ and an open subgroup $I \subset I_K$ such that for $g \in I$, the action of g on $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$ is given by $\exp(t_\ell(g)N)$. The endomorphism N defines the monodromy filtration Fil_\bullet^N on $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$; it is the unique separated and exhaustive increasing filtration such that $N(\text{Fil}_s^N) \subset \text{Fil}_{s-2}^N(-1)$ for $s \in \mathbb{Z}$ and such that the induced map $N^s: \text{gr}_s^N H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell) \rightarrow \text{gr}_{-s}^N H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)(-s)$ is an isomorphism for $s \geq 0$.

Conjecture 10.4 (ℓ -adic monodromy-weight conjecture for $\ell \neq p$). *For $s \in \mathbb{Z}$ and $g \in W_K$, the eigenvalues of g on $\text{gr}_s^N H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$ are algebraic numbers and the complex absolute values of their conjugates are $p^{(m+s)v(g)/2}$.*

Remark 10.5. When X has good reduction, the conjecture follows from Deligne's purity theorem on weights in [Del80]. The conjecture is also known for $m \leq 2$; When X has a strictly semistable model and $\dim X = 2$, it is [RZ82, Satz 2.13]. Using [dJ96], we can reduce the general case to this case (see [Sai03, Lemma 3.9] or the proof of Theorem 10.8 below).

We next discuss the case $\ell = p$. For this we need to recall the general recipe of Fontaine [Fon94c].

Let V be a de Rham \mathbb{Q}_p -representation of Gal_K . By the p -adic monodromy theorem [Ber02, Théorème 0.7], V is potentially semistable. We set

$$\widehat{D}_{\text{pst}}(V) := \varprojlim_{H \subset I_K} (B_{\text{st}} \otimes_{\mathbb{Q}_p} V)^H,$$

where H runs over open subgroups of I_K . It is a P_0 -vector space of dimension $\dim_{\mathbb{Q}_p} V$ equipped with a semilinear action of Gal_K , a Frobenius $\varphi: \widehat{D}_{\text{pst}}(V) \rightarrow \widehat{D}_{\text{pst}}(V)$ (i.e., a σ -semilinear injective map commuting with Gal_K -action), and a monodromy operator $N: \widehat{D}_{\text{pst}}(V) \rightarrow \widehat{D}_{\text{pst}}(V)$ (i.e., a P_0 -linear map commuting with Gal_K -action) satisfying $N\varphi = p\varphi N$. Such an object is called a *p -module of Deligne* in [Fon94c, 1.1]. It follows that φ is bijective and that N is nilpotent.

We associate to $\widehat{D}_{\text{pst}}(V)$ a Weil–Deligne representation $W\widehat{D}_{\text{pst}}(V)$ over P_0 as follows: Set $W\widehat{D}_{\text{pst}}(V) := \widehat{D}_{\text{pst}}(V)$ as a P_0 -vector space equipped with the monodromy operator N . Define a P_0 -linear action of W_K on $W\widehat{D}_{\text{pst}}(V)$ by letting

$g \in W_K$ act as $g \cdot \varphi^{n(g)}$. Since N acts nilpotently, there exists a unique separated and exhaustive increasing filtration Fil_\bullet^N on $W\widehat{D}_{\text{pst}}(V)$ such that $N(\text{Fil}_s^N) \subset \text{Fil}_{s-2}^N$ for $s \in \mathbb{Z}$ and such that $N^s: \text{gr}_s^N \rightarrow \text{gr}_{-s}^N$ is an isomorphism for every $s \geq 0$.

Let us return to the p -adic representation $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p)$. It is de Rham by [Tsu99] and [dJ96] (cf. [Tsu02, Theorem A1]). We denote by $H_{\text{pst},p}^m(X)$ the associated Weil–Deligne representation $W\widehat{D}_{\text{pst}}(H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p))$ of K .

Conjecture 10.6 (p -adic monodromy-weight conjecture). *For $s \in \mathbb{Z}$ and $g \in W_K$, the eigenvalues of g on $\text{gr}_s^N H_{\text{pst},p}^m(X)$ are algebraic numbers and the complex absolute values of their conjugates are $p^{(m+s)v(g)/2}$.*

Remark 10.7. This is part of the conjecture stated in [Jan88, p.347] and [Jan10, Conjecture 5.1].

The following result will be well known to experts, but we give a proof for completeness.

Theorem 10.8. *Conjecture 10.6 holds for $m \leq 2$.*

Proof. We follow the proof of [Sai03, Lemma 3.9]. Note that for a finite extension L of K , $H_{\text{pst},p}^m(X)$ and $H_{\text{pst},p}^m(X_L)$ are isomorphic as Weil–Deligne representations of L (after rescaling the monodromy operator; see [Fon94c, 2.2.6]). It follows that the assertion for $H_{\text{pst},p}^m(X)$ is equivalent to the one for $H_{\text{pst},p}^m(X_L)$.

We may assume that X is geometrically connected over K . By [dJ96, Theorem 6.5], there exist a finite extension L of K inside \overline{K} , a smooth projective scheme X' over L , and a proper surjective and generically finite morphism $f: X' \rightarrow X$. The morphism f induces a Gal_L -equivariant injection $f^*: H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p) \rightarrow H_{\text{ét}}^m(X'_{\overline{K}}, \mathbb{Q}_p)$ and thus an injective map $f^*: H_{\text{pst},p}^m(X_L) \rightarrow H_{\text{pst},p}^m(X'_L)$ of Weil–Deligne representations of L . In this case, it suffice to prove the assertion for the latter, and thus we may further assume that X is projective by replacing X by X' and K by L . By replacing X by its hyperplane section, we may also assume $\dim X \leq 2$.

By applying [dJ96, Theorem 6.5] again, we may assume that X admits a projective and strictly semistable model \mathcal{X} over \mathcal{O}_K . Denote the special fiber $\mathcal{X} \otimes_{\mathcal{O}_K} k$ by Y . Let $M_{\mathcal{X}}$ be the fine log structure on \mathcal{X} associated to the simple normal crossing divisor $Y \subset \mathcal{X}$ and let M_Y be the inverse image of $M_{\mathcal{X}}$ on Y . Write $H_{\log \text{cris}}^m(Y/W)$ for the log-crystalline cohomology of $(Y, M_Y)/(k, \mathbb{N} \oplus k^\times)$ over $W := W(k)$ and set $H_{\log \text{cris}}^m(\mathcal{X}) := H_{\log \text{cris}}^m(Y/W) \otimes_W K_0$, where $K_0 := W[p^{-1}]$. Then $H_{\log \text{cris}}^m(\mathcal{X})$ has a natural structure of (φ, N) -module relative to K^9 . By [Tsu99, Theorem 0.2], we have an isomorphism $D_{\text{st}}(H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p)) \cong H_{\log \text{cris}}^m(\mathcal{X})$ of (φ, N) -modules relative to K^{10} . It follows that $\widehat{D}_{\text{pst}}(H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p)) \cong H_{\log \text{cris}}^m(\mathcal{X}) \otimes_{K_0} P_0$ as p -modules of Deligne.

Note that φ^h acts K_0 -linearly on $H_{\log \text{cris}}^m(\mathcal{X})$ and the monodromy operator defines the monodromy filtration Fil_\bullet^N on $H_{\log \text{cris}}^m(\mathcal{X})$. By the definition of the Weil–Deligne representation $H_{\text{pst},p}^m(X)$ and the above isomorphism of p -modules of

⁹A choice of a uniformizer of K equips $H_{\log \text{cris}}^m(\mathcal{X})$ with a filtration over K by the comparison isomorphism to the log-de Rham cohomology, but we do not need this structure.

¹⁰Depending on the convention of the monodromy operators, we may need to change the sign of one of the monodromy operators, but this does not affect our argument.

Deligne, it suffices to prove that the eigenvalues of φ^h on $\mathrm{gr}_s^N H_{\log \text{ cris}}^m(\mathcal{X})$ are algebraic numbers and the complex absolute values of their conjugates are $p^{(m+s)h/2} = q^{(m+s)/2}$.

For $r \geq 1$, let $Y^{(r)}$ denote the disjoint union of all r -fold intersections of the different irreducible components of Y . Then we have the p -adic weight spectral sequence

$$E_1^{-l, m+l} = \bigoplus_{j \geq \max\{-l, 0\}} H_{\text{cris}}^{m-2j-l}(Y^{(2j+l+1)}/W)(-j-l) \otimes_W K_0 \implies H_{\log \text{ cris}}^m(\mathcal{X}).$$

This spectral sequence is φ -equivariant (see [Nak05, (9.11.1)] and [Nak06, §2 (B)] for the details). By the purity of weights of the E_1 -term ([KM74]), the spectral sequence degenerates at E_2 , inducing the filtration Fil_{\bullet}^W on $H_{\log \text{ cris}}^m(\mathcal{X})$ such that the eigenvalues of φ^h on $\mathrm{gr}_s^W H_{\log \text{ cris}}^m(\mathcal{X})$ are algebraic numbers whose complex conjugates have absolute value $q^{(m+s)/2}$. By [Mok93, Théorème 5.3, Corollaire 6.2.3] and [Nak06, Theorem 8.3, Corollary 8.4], two filtrations Fil_{\bullet}^N and Fil_{\bullet}^W on $H_{\log \text{ cris}}^m(\mathcal{X})$ agree. This completes the proof. \square

We now want to compare the results for ℓ and p . For this, we need the following theorem on ℓ -independence.

Theorem 10.9 ([Och99, Theorem B, Theorem 3.1]). *Let X be a smooth proper scheme over K of dimension n . For every $g \in W_K^+$, the alternating sum*

$$\sum_{m=0}^{2n} \mathrm{tr}(g^*; H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_{\ell}))$$

is a rational integer that is independent of ℓ with $\ell \neq p$. Moreover, it is equal to

$$\sum_{m=0}^{2n} \mathrm{tr}(g^*; H_{\text{pst}, p}^m(X)).$$

Corollary 10.10. *Let X be a smooth proper scheme over K of dimension at most two. For every $m \in \mathbb{N}$ and $g \in W_K^+$, the trace $\mathrm{tr}(g^*; H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_{\ell}))$ is a rational integer that is independent of ℓ with $\ell \neq p$. Moreover, it is equal to $\mathrm{tr}(g^*; H_{\text{pst}, p}^m(X))$.*

Proof. The cases $m = 0, \dim X$ are obvious. The same assertion for an abelian variety with $m = 1$ is also known (cf. [Fon94c, Remarque 2.4.6 (iii)] and [Noo17, Corollary 2.2]). From these results, it remains to deal with the case where $\dim X = 2$ and $m = 2$; this follows from the above theorem on ℓ -independence of the alternating sum. \square

Remark 10.11. For $\ell \neq p$, we can also associate to the ℓ -adic representation $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_{\ell})$ a Weil–Deligne representation of K over \mathbb{Q}_{ℓ} , which we denote by $H_{\text{pst}, \ell}^m(X)$. It is expected that for every smooth proper scheme X over K , the Weil–Deligne representations $H_{\text{pst}, \ell}^m(X)$ with ℓ ranging all the primes including p are compatible in the sense of [Fon94c, 2.4.2]. See [Fon94c, 2.4] for the details.

Corollary 10.12. *Let X be a smooth proper scheme of dimension at most two over K and let $m \in \mathbb{N}$. For a rational prime $\ell \neq p$, the ℓ -adic representation $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_{\ell})$ is potentially unramified if and only if the p -adic representation $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_p)$ is potentially crystalline.*

Proof. The former condition in the assertion is equivalent to the condition that $N = 0$ on $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$. Since ℓ -adic monodromy-weight conjecture is known for X (Remark 10.5), it is also equivalent to the condition that, for every $g \in W_K^+$, the eigenvalues of g on $H_{\text{ét}}^m(X_{\overline{K}}, \mathbb{Q}_\ell)$, which are algebraic numbers, satisfy the property that the complex absolute values of their conjugates are $p^{mv(g)/2}$. Similarly, by Theorem 10.8, the latter condition in the assertion is equivalent to the condition that, for every $g \in W_K^+$, the eigenvalues of g on $H_{\text{pst}, p}^m(X)$, which are algebraic numbers, satisfy the property that the complex absolute values of their conjugates are $p^{mv(g)/2}$. Hence the assertion easily follows from Corollary 10.10. \square

10.3. Cohomologically potentially good reduction locus. Let K be a complete discrete valuation field of characteristic zero with *finite* residue field of characteristic p .

Definition 10.13. Let $f: Y \rightarrow X$ be a smooth proper morphism between smooth algebraic varieties over K and let X^{ad} denote the adic space associated to X . We say that an open subset U of X^{ad} is a *cohomologically potentially good reduction locus* of f of degree m if U satisfies the following property: for every classical point x of X^{ad} and every geometric point \overline{x} above x ,

- (i) if $x \in U$, then $(R^mf_*\mathbb{Q}_\ell)_{\overline{x}}$ is potentially unramified for $\ell \neq p$ and $(R^mf_*\mathbb{Q}_p)_{\overline{x}}$ is potentially crystalline as Galois representations of $k(x)$;
- (ii) if $x \notin U$, then $(R^mf_*\mathbb{Q}_\ell)_{\overline{x}}$ is not potentially unramified for $\ell \neq p$ and $(R^mf_*\mathbb{Q}_p)_{\overline{x}}$ is not potentially crystalline as Galois representations of $k(x)$.

Remark 10.14. Note that our conditions do not characterize cohomologically potentially good reduction loci. On the other hand, the union of two cohomologically potentially good reduction loci is also a cohomologically potentially good reduction locus and thus we can consider the largest cohomologically potentially good reduction locus if there is at least one.

Theorem 10.15. *Let $f: X \rightarrow S$ be a smooth proper morphism between smooth algebraic varieties over K . Assume that the relative dimension of f is at most two. Then there exists a cohomologically potentially good reduction locus of f of any degree.*

Proof. By Corollary 10.12, it follows from Theorem 10.2 or [Kis99, Proposition 5.2 (2)]. \square

Remark 10.16. As mentioned in the introduction, Imai and Mieda introduce a notion of the potentially good reduction locus of a Shimura variety (cf. [IM20, Definition 5.19]). It is a quasi-compact constructible open subset of the associated adic space where every ℓ -adic automorphic étale local system is potentially unramified for $\ell \neq p$ and every p -adic automorphic étale local system is potentially crystalline at every classical point. Due to the constructibility, such a locus is unique if it exists. They prove the existence of the potentially good reduction locus of Shimura varieties of preabelian type, and use them to study the étale cohomology of Shimura varieties (cf. [IM20, Theorem 6.1]).

REFERENCES

- [AK00] D. Abramovich and K. Karu, *Weak semistable reduction in characteristic 0*, Invent. Math. **139** (2000), no. 2, 241–273. MR 1738451

- [ALT18] Karim Adiprasito, Gaku Liu, and Michael Temkin, *Semistable reduction in characteristic 0*, arXiv e-prints (2018), arXiv:1810.03131v2.
- [AI12] Fabrizio Andreatta and Adrian Iovita, *Semistable sheaves and comparison isomorphisms in the semistable case*, Rend. Semin. Mat. Univ. Padova **128** (2012), 131–285 (2013). MR 3076834
- [AI13] ———, *Comparison isomorphisms for smooth formal schemes*, J. Inst. Math. Jussieu **12** (2013), no. 1, 77–151. MR 3001736
- [Ber02] Laurent Berger, *Représentations p -adiques et équations différentielles*, Invent. Math. **148** (2002), no. 2, 219–284. MR 1906150
- [BC08] Laurent Berger and Pierre Colmez, *Familles de représentations de de Rham et monodromie p -adique*, Astérisque (2008), no. 319, 303–337, Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules. MR 2493221
- [Ber90] Vladimir G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Mathematical Surveys and Monographs, vol. 33, American Mathematical Society, Providence, RI, 1990. MR 1070709
- [Ber93] ———, *Étale cohomology for non-Archimedean analytic spaces*, Inst. Hautes Études Sci. Publ. Math. (1993), no. 78, 5–161 (1994). MR 1259429
- [BO83] P. Berthelot and A. Ogus, *F -isocrystals and de Rham cohomology. I*, Invent. Math. **72** (1983), no. 2, 159–199. MR 700767
- [BGR84] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 261, Springer-Verlag, Berlin, 1984, A systematic approach to rigid analytic geometry. MR 746961
- [Bri06] Olivier Brinon, *Représentations cristallines dans le cas d'un corps résiduel imparfait*, Ann. Inst. Fourier (Grenoble) **56** (2006), no. 4, 919–999. MR 2266883
- [Bri08] ———, *Représentations p -adiques cristallines et de de Rham dans le cas relatif*, Mém. Soc. Math. Fr. (N.S.) (2008), no. 112, vi+159. MR 2484979
- [BC09] Olivier Brinon and Brian Conrad, *CMI summer school notes on p -adic Hodge theory (preliminary version)*, Available at <http://math.stanford.edu/~conrad/papers/notes.pdf>, 2009.
- [Col08] Pierre Colmez, *Espaces vectoriels de dimension finie et représentations de de Rham*, Astérisque (2008), no. 319, 117–186, Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules. MR 2493217
- [CF00] Pierre Colmez and Jean-Marc Fontaine, *Construction des représentations p -adiques semi-stables*, Invent. Math. **140** (2000), no. 1, 1–43. MR 1779803
- [Del80] Pierre Deligne, *La conjecture de Weil. II*, Inst. Hautes Études Sci. Publ. Math. (1980), no. 52, 137–252. MR 601520 (83c:14017)
- [dJ96] A. J. de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math. (1996), no. 83, 51–93. MR 1423020
- [DGS94] Bernard Dwork, Giovanni Gerotto, and Francis J. Sullivan, *An introduction to G -functions*, Annals of Mathematics Studies, vol. 133, Princeton University Press, Princeton, NJ, 1994. MR 1274045
- [Fal88] Gerd Faltings, *p -adic Hodge theory*, Journal of the American Mathematical Society **1** (1988), no. 1, 255–299.
- [Fal89] ———, *Crystalline cohomology and p -adic Galois-representations*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 25–80. MR 1463696
- [Fal90] ———, *F -isocrystals on open varieties: results and conjectures*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 219–248. MR 1106900
- [Fal02] ———, *Almost étale extensions*, Astérisque (2002), no. 279, 185–270, Cohomologies p -adiques et applications arithmétiques, II. MR 1922831
- [Fal10] ———, *Coverings of p -adic period domains*, J. Reine Angew. Math. **643** (2010), 111–139. MR 2658191
- [Far18] Laurent Fargues, *La courbe*, Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures, World Sci. Publ., Hackensack, NJ, 2018, pp. 291–319. MR 3966767

- [Fon94a] Jean-Marc Fontaine, *Le corps des périodes p -adiques*, Astérisque (1994), no. 223, 59–111, With an appendix by Pierre Colmez, Périodes p -adiques (Bures-sur-Yvette, 1988). MR 1293971
- [Fon94b] ———, *Représentations p -adiques semi-stables*, Astérisque (1994), no. 223, 113–184, With an appendix by Pierre Colmez, Périodes p -adiques (Bures-sur-Yvette, 1988). MR 1293972
- [Fon94c] ———, *Représentations l -adiques potentiellement semi-stables*, Astérisque (1994), no. 223, 321–347, Périodes p -adiques (Bures-sur-Yvette, 1988). MR 1293977
- [Har13] Urs Hartl, *On a conjecture of Rapoport and Zink*, Invent. Math. **193** (2013), no. 3, 627–696. MR 3091977
- [Hou61] Christian Houzel, *Géométrie analytique locale, II. Théorie des morphismes finis*, Séminaire Henri Cartan **13** (1960-1961), no. 2 (fr), talk:19.
- [Hub94] Roland Huber, *A generalization of formal schemes and rigid analytic varieties*, Math. Z. **217** (1994), no. 4, 513–551. MR 1306024
- [Hub96] ———, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, E30, Friedr. Vieweg & Sohn, Braunschweig, 1996. MR 1734903
- [Hyo89] Osamu Hyodo, *On variation of Hodge-Tate structures*, Math. Ann. **284** (1989), no. 1, 7–22. MR 995378
- [IM20] Naoki Imai and Yoichi Mieda, *Potentially good reduction loci of Shimura varieties*, Tunis. J. Math. **2** (2020), no. 2, 399–454. MR 3990825
- [Jan88] Uwe Jannsen, *On the Galois cohomology of l -adic representations attached to varieties over local or global fields*, Séminaire de Théorie des Nombres, Paris 1986–87, Progr. Math., vol. 75, Birkhäuser Boston, Boston, MA, 1988, pp. 165–182. MR 990511
- [Jan10] ———, *Weights in arithmetic geometry*, Jpn. J. Math. **5** (2010), no. 1, 73–102. MR 2609323
- [Kat70] Nicholas M. Katz, *Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin*, Inst. Hautes Études Sci. Publ. Math. (1970), no. 39, 175–232. MR 0291177
- [Kat73] ———, *Travaux de Dwork*, Séminaire Bourbaki, 24ème année (1971/1972), Exp. No. 409, 1973, pp. 167–200. Lecture Notes in Math., Vol. 317. MR 0498577
- [KM74] Nicholas M. Katz and William Messing, *Some consequences of the Riemann hypothesis for varieties over finite fields*, Invent. Math. **23** (1974), 73–77. MR 332791
- [Ked10a] Kiran S. Kedlaya, *p -adic differential equations*, Cambridge Studies in Advanced Mathematics, vol. 125, Cambridge University Press, Cambridge, 2010. MR 2663480
- [Ked10b] ———, *Relative p -adic Hodge theory and Rapoport-Zink period domains*, Proceedings of the International Congress of Mathematicians. Volume II, Hindustan Book Agency, New Delhi, 2010, pp. 258–279. MR 2827795
- [KL15] Kiran S. Kedlaya and Ruochuan Liu, *Relative p -adic Hodge theory: Foundations*, Astérisque (2015), no. 371, 239. MR 3379653
- [Kis99] Mark Kisin, *Local constancy in p -adic families of Galois representations*, Math. Z. **230** (1999), no. 3, 569–593. MR 1680032
- [LL20] Brian Lawrence and Shizhang Li, *A counterexample to an optimistic guess about étale local systems*, arXiv e-prints (2020), arXiv:2002.03799v1.
- [LZ17] Ruochuan Liu and Xinwen Zhu, *Rigidity and a Riemann-Hilbert correspondence for p -adic local systems*, Invent. Math. **207** (2017), no. 1, 291–343. MR 3592758
- [Mok93] Abdellah Mokrane, *La suite spectrale des poids en cohomologie de Hyodo-Kato*, Duke Math. J. **72** (1993), no. 2, 301–337. MR 1248675
- [Moo18] Yong Suk Moon, *Relative crystalline representations and weakly admissible modules*, arXiv e-prints (2018), arXiv:1806.00867v2.
- [Mor14] Kazuma Morita, *Crystalline and semi-stable representations in the imperfect residue field case*, Asian J. Math. **18** (2014), no. 1, 143–157. MR 3215344
- [Nak05] Yuki Yoshi Nakajima, *p -adic weight spectral sequences of log varieties*, J. Math. Sci. Univ. Tokyo **12** (2005), no. 4, 513–661. MR 2206357
- [Nak06] ———, *Signs in weight spectral sequences, monodromy-weight conjectures, log Hodge symmetry and degenerations of surfaces*, Rend. Sem. Mat. Univ. Padova **116** (2006), 71–185. MR 2287344
- [Noo17] Rutger Noot, *The p -adic representation of the Weil-Deligne group associated to an abelian variety*, J. Number Theory **172** (2017), 301–320. MR 3573156

- [Och99] Tadashi Ochiai, *l*-independence of the trace of monodromy, Math. Ann. **315** (1999), no. 2, 321–340. MR 1715253
- [Ohk13] Shun Ohkubo, *The p-adic monodromy theorem in the imperfect residue field case*, Algebra Number Theory **7** (2013), no. 8, 1977–2037. MR 3134041
- [RZ82] M. Rapoport and Th. Zink, *Über die lokale Zetafunktion von Shimuravarietäten. Monodromiefiltration und verschwindende Zyklen in ungleicher Charakteristik*, Invent. Math. **68** (1982), no. 1, 21–101. MR 666636
- [Sai03] Takeshi Saito, *Weight spectral sequences and independence of l*, J. Inst. Math. Jussieu **2** (2003), no. 4, 583–634. MR 2006800
- [Sch12] Peter Scholze, *Perfectoid spaces*, Publ. Math. Inst. Hautes Études Sci. **116** (2012), 245–313. MR 3090258
- [Sch13] ———, *p-adic Hodge theory for rigid-analytic varieties*, Forum Math. Pi **1** (2013), e1, 77. MR 3090230
- [Sch16] ———, *p-adic Hodge theory for rigid-analytic varieties—corrigendum*, Forum Math. Pi **4** (2016), e6, 4. MR 3535697
- [SGA1] *Revêtements étales et groupe fondamental*, Lecture Notes in Mathematics, Vol. 224, Springer-Verlag, Berlin-New York, 1971, Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de M. Raynaud. MR 0354651
- [SGA7] *Groupes de monodromie en géométrie algébrique. I*, Lecture Notes in Mathematics, Vol. 288, Springer-Verlag, Berlin-New York, 1972, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 I), Dirigé par A. Grothendieck. Avec la collaboration de M. Raynaud et D. S. Rim. MR 0354656
- [Shi18] Koji Shimizu, *Constancy of generalized Hodge-Tate weights of a local system*, Compos. Math. **154** (2018), no. 12, 2606–2642. MR 3873527
- [TT19] Fucheng Tan and Jilong Tong, *Crystalline comparison isomorphisms in p-adic Hodge theory: the absolutely unramified case*, Algebra Number Theory **13** (2019), no. 7, 1509–1581. MR 4009670
- [Tsu99] Takeshi Tsuji, *p-adic étale cohomology and crystalline cohomology in the semi-stable reduction case*, Invent. Math. **137** (1999), no. 2, 233–411. MR 1705837
- [Tsu02] ———, *Semi-stable conjecture of Fontaine-Jannsen: a survey*, Astérisque (2002), no. 279, 323–370, Cohomologies p-adiques et applications arithmétiques, II. MR 1922833
- [Tsu] ———, *Crystalline sheaves and filtered convergent F-isocrystals on log schemes*, preprint.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ, USA
E-mail address: kshimizu@ias.edu