

Positive elliptic-elliptic rotopulsators on Clifford tori of nonconstant size project onto regular polygons

Pieter Tibboel

Department of Mathematical Sciences
Xi'an Jiaotong-Liverpool University
Suzhou, China
Pieter.Tibboel@xjtlu.edu.cn

August 3, 2020

Abstract

Let $q_1, \dots, q_n$ be the position vectors of the point masses of the curved n -body problem. Consider any positive elliptic-elliptic rotopulsator solution $q_i^T = (r \cos(\theta + \alpha_i), r \sin(\theta + \alpha_i), \rho \cos(\phi + \beta_i), \rho \sin(\phi + \beta_i))$, $i \in \{1, \dots, n\}$, where $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n \in [0, 2\pi)$ are constants, ϕ , θ , r and ρ are twice-differentiable, continuous, nonconstant functions, $r^2 + \rho^2 = 1$, $r \geq 0$ and $\rho \geq 0$. We prove that if the configuration of the point masses is of nonconstant size, the configuration of the vectors $(r \cos(\theta + \alpha_i), r \sin(\theta + \alpha_i))^T$ is a regular polygon, as is the configuration of the vectors $(\rho \cos(\phi + \beta_i), \rho \sin(\phi + \beta_i))^T$, $i \in \{1, \dots, n\}$.

1 Introduction

Let $\sigma = \pm 1$. The n -body problem in spaces of constant Gaussian curvature, or curved n -body problem for short, is the problem of finding the dynamics of point masses

$$q_1, \dots, q_n \in \mathbb{M}_\sigma^3 = \{(x_1, x_2, x_3, x_4)^T \in \mathbb{R}^4 | x_1^2 + x_2^2 + x_3^2 + \sigma x_4^2 = \sigma\},$$

with respective masses $m_1 > 0, \dots, m_n > 0$, determined by the system of differential equations

$$\ddot{q}_i = \sum_{j=1, j \neq i}^n \frac{m_j(q_j - \sigma(q_i \odot q_j)q_i)}{(\sigma - \sigma(q_i \odot q_j)^2)^{\frac{3}{2}}} - \sigma(\dot{q}_i \odot \dot{q}_i)q_i, \quad i \in \{1, \dots, n\}, \quad (1.1)$$

where for $x, y \in \mathbb{M}_\sigma^3$ the product $\cdot \odot \cdot$ is defined as

$$x \odot y = x_1y_1 + x_2y_2 + x_3y_3 + \sigma x_4y_4.$$

The study of the curved n -body problem has applications to for example geometric mechanics, Lie groups and algebras, non-Euclidean and differential geometry and stability theory, the theory of polytopes and topology (see for example [10]) and for $n = 2$ goes back as far as the 1830s (see [1], [2], [3], [12], [13], [14], [15] and [20] for a historical overview and recent results). However, the first paper giving an explicit n -body problem in spaces of constant Gaussian curvature for general $n \geq 2$ was published in 2008 by Diacu, Pérez-Chavela and Santoprete (see [13], [14] and [15]). This breakthrough then gave rise to further results for the $n \geq 2$ case in [4], [5]–[12], [16], [19], [23]–[34] and the references therein.

Rotopulsators are solutions to (1.1) for which the configuration of the point masses may only rotate or change size, but retains its shape over time. They were first introduced by Diacu and Kordlou in [12] and can be divided into five classes, two for the positive curvature case ($\sigma = 1$) and three for the negative curvature case ($\sigma = -1$). Positive elliptic-elliptic rotopulsators are one of the two possible types for the positive curvature case and can be defined as follows: Let

$$R(x) = \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}.$$

If we write $q_i = (q_{i1}, q_{i2}, q_{i3}, q_{i4})^T$, $i \in \{1, \dots, n\}$, then we call $q_1, \dots, q_n$ a *positive elliptic-elliptic rotopulsator* if there exist nonnegative scalar functions r_i, ρ_i for which $r_i^2 + \rho_i^2 = 1$, scalar functions θ, ϕ and constants $\alpha_i, \beta_i \in \mathbb{R}$, such that

$$\begin{pmatrix} q_{i1} \\ q_{i2} \end{pmatrix} = r_i R(\theta + \alpha_i) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} q_{i3} \\ q_{i4} \end{pmatrix} = \rho_i R(\phi + \beta_i) \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (1.2)$$

Another definition we will need is the definition of a Clifford torus: Let $a > 0$ and $b > 0$. By a Clifford torus, we mean any set

$$\{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 | x_1^2 + x_2^2 = a^2, x_3^2 + x_4^2 = b^2\}$$

Finally, before we get to this paper's main result, a remark on terminology is needed: Throughout this paper, if $v_1, v_2, \dots, v_n \in \mathbb{R}^2$ are n not necessarily distinct vectors, then if among these vectors there are k distinct vectors $v_{j_1}, \dots, v_{j_k}$, such that for all other vectors v_i , $i \in \{1, \dots, n\} \setminus \{j_1, \dots, j_k\}$ we have

that there is an $l \in \{1, \dots, k\}$ such that $v_i = v_{j_l}$, then we say that the vectors $v_1, \dots, v_n$ represent the vertices of a polygon with k vertices. With that said, we will continue on the topic of positive elliptic-elliptic rotopulsators: In [26] it was proven that if the β_i are all equal and the r_i and ρ_i are independent of i and not constant, then the configuration of the point masses has to be a regular polygon and in [30] it was shown that in that case all masses have to be equal. For the case that the r_i and ρ_i are independent of i and not constant and the β_i are not necessarily equal, almost nothing is known, except that in [12] for $n = 3$ nonexistence was proven for Lagrangian configurations and in [19], for $n = 4$, nonexistence was proven for rectangular configurations and all masses equal. In this paper we will prove for general n that positive elliptic-elliptic rotopulsators on Clifford tori of nonconstant size project onto regular polygons. Specifically:

Theorem 1.1. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with r_i and ρ_i independent of i and not constant and the size of the configuration not fixed. Then both the $(q_{i1}, q_{i2})^T$, $i \in \{1, \dots, n\}$ and the $(q_{i3}, q_{i4})^T$, $i \in \{1, \dots, n\}$ represent vertices of a regular polygon.

We will now first formulate a criterion and lemmas needed to prove Theorem 1.1 in section 2, after which we will prove Theorem 1.1 in section 3.

2 Background theory

Throughout this paper we will use the notation introduced in the previous section for rotopulsators. Additionally, we will adopt the notation $\langle \cdot, \cdot \rangle$ for the Euclidean inner product. To prove Theorem 1.1, ideally we would like to repeat the main argument used to prove that for positive elliptic-elliptic rotopulsators for which $\beta_i = \beta_j$ for all $i, j \in \{1, \dots, n\}$ the configuration of the point masses has to be a regular polygon (see [25], [26]). The problem with positive elliptic-elliptic rotopulsators for which r_i and ρ_i are independent of i , $i \in \{1, \dots, n\}$, is, as we will see, that there is a possibility that there are $i, j \in \{1, \dots, n\}$, $i \neq j$ for which $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) = 0$. To illustrate that, we will start by formulating a lemma and a criterion for the existence of positive elliptic-elliptic rotopulsators we need. These were proven for general r_i and ρ_i , $i \in \{1, \dots, n\}$ in [12], but as the proofs are very short, we will add them here:

Lemma 2.1. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with $r_i = r$ and $\rho_i = \rho$ independent of i . Then $2\dot{r}\dot{\theta} + r\ddot{\theta} = 0$ and $2\dot{\rho}\dot{\phi} + \rho\ddot{\phi} = 0$.

Proof. It was proven in [7] that $\sum_{j=1}^n m_j q_j \wedge \dot{q}_j = \mathbf{0}$ where $\wedge$ represents the wedge product and $\mathbf{0}$ is the zero bivector. If e_1, e_2, e_3 and e_4 are the standard basis vectors of $\mathbb{R}^4$, then

$$0e_1 \wedge e_2 = \sum_{j=1}^n m_j (q_{j1}\ddot{q}_{j2} - q_{j2}\ddot{q}_{j1})e_1 \wedge e_2.$$

Using (1.2), $q_{j1}\ddot{q}_{j2} - q_{j2}\ddot{q}_{j1}$ can be rewritten as $2\dot{r}\dot{\theta} + r\ddot{\theta}$, so $0 = \sum_{j=1}^n m_j (2\dot{r}\dot{\theta} + r\ddot{\theta})$, which gives that $2\dot{r}\dot{\theta} + r\ddot{\theta} = 0$. Repeating this argument replacing e_1 and e_2 with e_3 and e_4 and q_{j1} and q_{j2} with q_{j3} and q_{j4} gives that $2\dot{\rho}\dot{\phi} + \rho\ddot{\phi} = 0$. $\square$

Criterion 2.1. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with $r_i = r$ and $\rho_i = \rho$ independent of i . Then

$$0 = \sum_{j=1, j \neq i}^n \frac{m_j \sin(\alpha_j - \alpha_i)}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}}, \quad (2.1)$$

$$0 = \sum_{j=1, j \neq i}^n \frac{m_j \sin(\beta_j - \beta_i)}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}} \text{ and} \quad (2.2)$$

$$\delta = r\rho^2 \sum_{j=1, j \neq i}^n \frac{m_j (\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i))}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}}, \quad (2.3)$$

where $\delta = \ddot{r} + r\rho^2((\dot{\phi})^2 - (\dot{\theta})^2) + r\left(\frac{\dot{r}}{\rho}\right)^2$.

Proof. Inserting (1.2) into (1.1), writing out the expressions for the first two coordinates of the vectors on both sides of (1.1) and multiplying both sides of the resulting equations from the left with $R(\theta + \alpha_i)^{-1}$ and using that

$r^2 + \rho^2 = 1$, we get that

$$\begin{aligned}
& (\ddot{r} - r\dot{\theta}^2) \begin{pmatrix} 1 \\ 0 \end{pmatrix} + (2\dot{r}\dot{\theta} + r\ddot{\theta}) \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\
&= r \sum_{j=1, j \neq i}^n \frac{m_j \left(\frac{\rho^2 (\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i))}{\sin(\alpha_j - \alpha_i)} \right)}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}} \\
&\quad - r(\dot{r}^2 + r^2\dot{\theta}^2 + \dot{\rho}^2 + \rho^2\dot{\phi}^2) \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \tag{2.4}
\end{aligned}$$

Collecting terms for the first coordinates of the vectors on both sides of (2.4) gives (2.3) and by Lemma 2.1 collecting terms for the second coordinates of the vectors on both sides of (2.4) gives (2.1). Repeating the argument used so far writing out the expressions for the third and fourth coordinates of the vectors on both sides of (1.1) instead then gives (2.2). The identity for the third coordinates can be rewritten as (2.3), which proves this criterion. $\square$

We can now explain the problem touched upon at the start of this section in more detail: The idea behind our proof of Theorem 1.1, which is essentially the idea behind proving that the configurations of the point masses have to be regular polygons in [25] and [26], is to use that the terms in the identities of Criterion 2.1 are linearly independent under certain conditions on the α_i and β_i , $i \in \{1, \dots, n\}$, if $r_i = r$ and $\rho_i = \rho$ are independent of i and not constant. However, we immediately see that a prerequisite to using such a linear independence argument is that $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0$ for those terms. We will therefore now prove three lemmas leading to the fact that we can exclude the possibility of $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) = 0$ and then give the exact conditions for the terms of the sums on the right-hand side of (2.1), (2.2) and (2.3) to be linearly independent:

Lemma 2.2. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with $r_i = r$ and $\rho_i = \rho$ independent of i and not constant and the size of the configuration of the point masses not fixed. Then for all but possibly one $i \in \{1, \dots, n\}$ we have that

$$\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0 \text{ for at least one } j \in \{1, \dots, n\} \setminus \{i\}.$$

Proof. If there are $i_1, i_2 \in \{1, \dots, n\}$ such that

$$\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) = 0 \text{ for all } j \in \{1, \dots, n\}, \quad i \in \{i_1, i_2\},$$

then

$$\begin{aligned}\langle q_i, q_j \rangle &= r^2 \cos(\alpha_j - \alpha_i) + \rho^2 \cos(\beta_j - \beta_i) = r^2 \cos(\alpha_j - \alpha_i) + \rho^2 \cos(\alpha_j - \alpha_i) \\ &= (r^2 + \rho^2) \cos(\alpha_j - \alpha_i) = \cos(\alpha_j - \alpha_i)\end{aligned}\quad (2.5)$$

for $i \in \{i_1, i_2\}$, which means that the angles between q_j , q_{i_1} and q_{i_2} are constant. As the length of each position vector is 1, we have that q_{i_1} and q_{i_2} are linearly dependent if and only if $q_{i_1} = -q_{i_2}$, which would mean that $1 - \langle q_{i_1}, q_{i_2} \rangle^2 = 0$, which would make (1.1) undefined, so q_{i_1} and q_{i_2} are linearly independent. As the lengths of q_{i_1} , q_{i_2} and q_j are all equal to 1 and therefore fixed and the angles between q_{i_1} and q_{i_2} , between q_{i_1} and q_j and between q_{i_2} and q_j are fixed by (2.5), the configuration of the point masses q_{i_1} , q_{i_2} and q_j is fixed for any j , which means that the configuration of the point masses can only rotate, but otherwise stays unchanged over time, which contradicts that the size of the configuration of the point masses is not constant. This means by extension that for all but perhaps one $i \in \{1, \dots, n\}$, (2.1), (2.2) and (2.3) have to have at least one term for which $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0$. $\square$

Next we will prove a lemma that serves as a check on restrictions on the q_i , $i \in \{1, \dots, n\}$ due to the q_i being the position vectors of a positive elliptic-elliptic rotopulsator for which the r_i and ρ_i are independent of i and not constant:

Lemma 2.3. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with $r_i = r$ and $\rho_i = \rho$ independent of i and not constant. Let $i_1, i_2, i_3 \in \{1, \dots, n\}$. Then

$$\begin{aligned}& \left((1 + B_{i_1 i_2, i_1 i_3}) + r^2 (A_{i_1 i_2, i_1 i_3} - B_{i_1 i_2, i_1 i_3}) \right)^2 \\ &= 4(\cos^2 \gamma_{i_1 i_2, i_1 i_3}) (1 - \cos(\beta_{i_2} - \beta_{i_1}) - r^2(\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1}))) \\ &\cdot (1 - \cos(\beta_{i_3} - \beta_{i_1}) - r^2(\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1})))\end{aligned}\quad (2.6)$$

and

$$\begin{aligned}& \left((1 + A_{i_1 i_2, i_1 i_3}) + \rho^2 (B_{i_1 i_2, i_1 i_3} - A_{i_1 i_2, i_1 i_3}) \right)^2 \\ &= 4(\cos^2 \gamma_{i_1 i_2, i_1 i_3}) (1 - \cos(\alpha_{i_2} - \alpha_{i_1}) - \rho^2(\cos(\beta_{i_2} - \beta_{i_1}) - \cos(\alpha_{i_2} - \alpha_{i_1}))) \\ &\cdot (1 - \cos(\alpha_{i_3} - \alpha_{i_1}) - \rho^2(\cos(\beta_{i_3} - \beta_{i_1}) - \cos(\alpha_{i_3} - \alpha_{i_1}))),\end{aligned}\quad (2.7)$$

where $\gamma_{i_1 i_2, i_1 i_3}$ is the constant angle between $q_{i_2} - q_{i_1}$ and $q_{i_3} - q_{i_1}$,

$$A_{i_1 i_2, i_1 i_3} = \cos(\alpha_{i_3} - \alpha_{i_2}) - \cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\alpha_{i_3} - \alpha_{i_1})$$

and

$$B_{i_1 i_2, i_1 i_3} = \cos(\beta_{i_3} - \beta_{i_2}) - \cos(\beta_{i_2} - \beta_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1}).$$

Proof. Because the q_i , $i \in \{1, \dots, n\}$ are the position vectors of the point masses of a rotopulsator, the shape of the configuration of the point masses has to remain unchanged over time. This means in particular that

$$\langle q_{i_2} - q_{i_1}, q_{i_3} - q_{i_1} \rangle = \|q_{i_2} - q_{i_1}\| \|q_{i_3} - q_{i_1}\| \cos \gamma_{i_1 i_2, i_1 i_3},$$

which means that

$$\langle q_{i_2} - q_{i_1}, q_{i_3} - q_{i_1} \rangle^2 = \|q_{i_2} - q_{i_1}\|^2 \|q_{i_3} - q_{i_1}\|^2 \cos^2 \gamma_{i_1 i_2, i_1 i_3}. \quad (2.8)$$

Writing out the left-hand side of (2.8) gives

$$\begin{aligned} \langle q_{i_2} - q_{i_1}, q_{i_3} - q_{i_1} \rangle^2 &= (1 - \langle q_{i_1}, q_{i_2} + q_{i_3} \rangle + \langle q_{i_2}, q_{i_3} \rangle)^2 \\ &= (1 - (r^2 (\cos(\alpha_{i_2} - \alpha_{i_1}) + \cos(\alpha_{i_3} - \alpha_{i_1})) \\ &\quad + \rho^2 (\cos(\beta_{i_2} - \beta_{i_1}) + \cos(\beta_{i_3} - \beta_{i_1}))) \\ &\quad + r^2 \cos(\alpha_{i_3} - \alpha_{i_2}) + \rho^2 \cos(\beta_{i_3} - \beta_{i_2})))^2, \end{aligned}$$

which, as $r^2 + \rho^2 = 1$, gives

$$\langle q_{i_2} - q_{i_1}, q_{i_3} - q_{i_1} \rangle^2 = ((1 + B_{i_1 i_2, i_1 i_3}) + r^2 (A_{i_1 i_2, i_1 i_3} - B_{i_1 i_2, i_1 i_3}))^2. \quad (2.9)$$

Writing out the right-hand side of (2.8) gives

$$\begin{aligned} &\|q_{i_2} - q_{i_1}\|^2 \|q_{i_3} - q_{i_1}\|^2 \cos^2 \gamma_{i_1 i_2, i_1 i_3} \\ &= 4(1 - \langle q_{i_1}, q_{i_2} \rangle)(1 - \langle q_{i_1}, q_{i_3} \rangle) \cos^2 \gamma_{i_1 i_2, i_1 i_3} \\ &= 4(1 - (r^2 \cos(\alpha_{i_2} - \alpha_{i_1}) + \rho^2 \cos(\beta_{i_2} - \beta_{i_1}))) \\ &\quad \cdot (1 - (r^2 \cos(\alpha_{i_3} - \alpha_{i_1}) + \rho^2 \cos(\beta_{i_3} - \beta_{i_1}))) \cos^2 \gamma_{i_1 i_2, i_1 i_3}, \end{aligned}$$

which, using again that $r^2 + \rho^2 = 1$, gives

$$\begin{aligned} &\|q_{i_2} - q_{i_1}\|^2 \|q_{i_3} - q_{i_1}\|^2 \cos^2 \gamma_{i_1 i_2, i_1 i_3} \\ &= 4(\cos^2 \gamma_{i_1 i_2, i_1 i_3}) (1 - \cos(\beta_{i_2} - \beta_{i_1}) - r^2 (\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1}))) \\ &\quad \cdot (1 - \cos(\beta_{i_3} - \beta_{i_1}) - r^2 (\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1}))). \quad (2.10) \end{aligned}$$

Combining (2.9) and (2.10) then gives (2.6) and interchanging the roles of the α s and the β s, of the A s and the B s and of r and ρ gives (2.7). $\square$

Lemma 2.4. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotopulsator solution of (1.1) with $r_i = r$ and $\rho_i = \rho$ independent of i and not constant and a configuration of nonconstant size. Then for all $i, j \in \{1, \dots, n\}$, $i \neq j$, we have that $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0$ and for all $k \in \{1, \dots, n\} \setminus \{i, j\}$, we have that

$$\frac{1 - \cos(\beta_i - \beta_j)}{\cos(\alpha_i - \alpha_j) - \cos(\beta_i - \beta_j)} = \frac{1 - \cos(\beta_k - \beta_j)}{\cos(\alpha_k - \alpha_j) - \cos(\beta_k - \beta_j)}. \quad (2.11)$$

and

$$\frac{1 - \cos(\alpha_i - \alpha_j)}{\cos(\alpha_i - \alpha_j) - \cos(\beta_i - \beta_j)} = \frac{1 - \cos(\alpha_k - \alpha_j)}{\cos(\alpha_k - \alpha_j) - \cos(\beta_k - \beta_j)}. \quad (2.12)$$

Proof. Note that in (2.6), if $4 \cos^2 \gamma_{i_1 i_2, i_1 i_3} \neq 0$, as r is not constant, we have that the left-hand side of (2.6) is a polynomial in terms of r^2 of degree two with a double root and the right-hand side of (2.6) is a polynomial of degree two in terms of r^2 that therefore also needs to have a double root, which means that

$$\frac{1 - \cos(\beta_{i_2} - \beta_{i_1})}{\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1})} = \frac{1 - \cos(\beta_{i_3} - \beta_{i_1})}{\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1})}$$

if $\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1}) \neq 0$ and $\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1}) \neq 0$. If $\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1}) = 0$, or $\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1}) = 0$, then again by (2.6) we have that both $\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1}) = 0$ and $\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1}) = 0$, as otherwise the degrees of the polynomials on the left-hand side and the right-hand side of (2.6) would not match. Suppose that $\gamma_{i_1 i_2, i_1 i_3} = \frac{\pi}{2}$. Then we repeat this argument for $\gamma_{i_3 i_2, i_3 i_1}$ and $\gamma_{i_2 i_1, i_2 i_3}$, which cannot be equal to $\frac{\pi}{2}$, as a triangle can have at most one right angle, instead, finding that either

$$\begin{aligned} \frac{1 - \cos(\beta_{i_2} - \beta_{i_1})}{\cos(\alpha_{i_2} - \alpha_{i_1}) - \cos(\beta_{i_2} - \beta_{i_1})} &= \frac{1 - \cos(\beta_{i_3} - \beta_{i_1})}{\cos(\alpha_{i_3} - \alpha_{i_1}) - \cos(\beta_{i_3} - \beta_{i_1})} \\ &= \frac{1 - \cos(\beta_{i_3} - \beta_{i_2})}{\cos(\alpha_{i_3} - \alpha_{i_2}) - \cos(\beta_{i_3} - \beta_{i_2})}, \end{aligned}$$

or that $\cos(\alpha_{i_k} - \alpha_{i_s}) - \cos(\beta_{i_k} - \beta_{i_s}) = 0$ for all $k, s \in \{1, 2, 3\}$. As this last result holds true for every triangle of point masses, we have that if there are $i, j \in \{1, \dots, n\}$, $i \neq j$ for which $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) = 0$, then we have that $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) = 0$ for all $i, j \in \{1, \dots, n\}$. As by Lemma 2.2 there are $i, j \in \{1, \dots, n\}$ for which

$$\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0,$$

this means that $\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i) \neq 0$ for all $i, j \in \{1, \dots, n\}, i \neq j$, so that means that (2.11) holds for all $i, j, k \in \{1, \dots, n\}, i \neq j, i \neq k, j \neq k$. Repeating the argument so far after interchanging the roles of the α s and the β s and r and ρ then proves that (2.12) holds for all $i, j, k \in \{1, \dots, n\}, i \neq j, i \neq k, j \neq k$ as well. This completes the proof. $\square$

We can now conclude:

Lemma 2.5. Let $q_1, \dots, q_n$ be a positive elliptic-elliptic rotapulsator solution of (1.1) with a configuration of nonconstant size, $r_i = r$ and $\rho_i = \rho$ independent of i and not constant. Let $A_{j_1 i_1}$ and $A_{j_2 i_2}$ be nonzero constants, $i_1, i_2, j_1, j_2 \in \{1, \dots, n\}, j_1 \neq i_1$ and $j_2 \neq i_2$. Then for terms

$$\frac{A_{j_1 i_1}}{(1 - (\cos(\beta_{j_1} - \beta_{i_1}) + r^2(\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})))^2)^{\frac{3}{2}}}$$

to cancel out against terms

$$\frac{A_{j_2 i_2}}{(1 - (\cos(\beta_{j_2} - \beta_{i_2}) + r^2(\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})))^2)^{\frac{3}{2}}}$$

in the sums in the right-hand sides of (2.1), (2.2) and (2.3), we need that

$$\cos(\alpha_{j_1} - \alpha_{i_1}) = \cos(\alpha_{j_2} - \alpha_{i_2}) \text{ and } \cos(\beta_{j_1} - \beta_{i_1}) = \cos(\beta_{j_2} - \beta_{i_2}).$$

Proof. Note that if $\cos(\alpha_{j_1} - \alpha_{i_1}) \neq \cos(\beta_{j_1} - \beta_{i_1})$, then terms

$$\frac{A_{j_1 i_1}}{(1 - (\cos(\beta_{j_1} - \beta_{i_1}) + r^2(\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})))^2)^{\frac{3}{2}}}$$

are linearly independent for different values of i_1, j_1 if and only if the roots of the polynomials $(1 - (\cos(\beta_{j_1} - \beta_i) + x(\cos(\alpha_{j_1} - \alpha_i) - \cos(\beta_{j_1} - \beta_i)))^2)$ differ for different values of j_1 . As the roots of

$$(1 - (\cos(\beta_{j_1} - \beta_{i_1}) + x(\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})))^2)$$

are

$$\frac{1 - \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})} \text{ and } -\frac{1 + \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})}$$

if $\cos(\alpha_{j_1} - \alpha_{i_1}) \neq \cos(\beta_{j_1} - \beta_{i_1})$, we have that terms

$$\frac{A_{j_1 i_1}}{(1 - (\cos(\beta_{j_1} - \beta_{i_1}) + r^2(\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})))^2)^{\frac{3}{2}}}$$

can cancel out against terms

$$\frac{A_{j_2 i_2}}{(1 - (\cos(\beta_{j_2} - \beta_{i_2}) + r^2(\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})))^2)^{\frac{3}{2}}}$$

if and only if

$$\frac{1 - \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})} = \frac{1 - \cos(\beta_{j_2} - \beta_{i_2})}{\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})}$$

and

$$-\frac{1 + \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})} = -\frac{1 + \cos(\beta_{j_2} - \beta_{i_2})}{\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})},$$

or

$$\frac{1 - \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})} = -\frac{1 - \cos(\beta_{j_2} - \beta_{i_2})}{\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})}$$

and

$$\frac{1 + \cos(\beta_{j_1} - \beta_{i_1})}{\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1})} = -\frac{1 + \cos(\beta_{j_2} - \beta_{i_2})}{\cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2})}.$$

The first of these possibilities is equivalent with $\cos(\beta_{j_1} - \beta_{i_1}) = \cos(\beta_{j_2} - \beta_{i_2})$
and

$$\cos(\alpha_{j_1} - \alpha_{i_1}) - \cos(\beta_{j_1} - \beta_{i_1}) = \cos(\alpha_{j_2} - \alpha_{i_2}) - \cos(\beta_{j_2} - \beta_{i_2}),$$

which, as

$$\begin{aligned} \frac{1 - \cos(\beta_j - \beta_i)}{\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)} &= \frac{1 - \cos(\alpha_j - \beta_i) + \cos(\alpha_j - \beta_i) - \cos(\alpha_j - \beta_i)}{\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)} \\ &= 1 + \frac{1 - \cos(\alpha_j - \beta_i)}{\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)}, \end{aligned}$$

also means that $\cos(\alpha_{j_1} - \alpha_{i_1}) = \cos(\alpha_{j_2} - \alpha_{i_2})$.

The second of these possibilities can be discarded by (2.11) and (2.12). This completes the proof. $\square$

3 Proof of Theorem 1.1

We will assume that there are $i, j \in \{1, \dots, n\}$ such that $\alpha_j - \alpha_i \neq 0$. If this is not the case, then we switch the roles of the α s and the β s. Additionally, let $S_{\alpha j} = \{s \in \{1, \dots, n\} | \alpha_s = \alpha_j\}$ and select $j_1, \dots, j_k \in \{1, \dots, n\}$ such that $S_{\alpha s_1} \cap S_{\alpha s_2} = \emptyset$ for $s_1, s_2 \in \{j_1, \dots, j_k\}$, $s_1 \neq s_2$ and $\bigcup_{u=1}^k S_{\alpha j_u} = \{1, \dots, n\}$, $0 \leq \alpha_{j_1} < \alpha_{j_2} < \dots < \alpha_{j_k} < 2\pi$ and $\alpha_{j_2} - \alpha_{j_1} \leq \alpha_{j_{u+1}} - \alpha_{j_u}$ for all $u \in \{1, \dots, k\}$, relabeling the point masses if necessary, where we define $\alpha_{j_{k+1}} = 2\pi + \alpha_{j_1}$. We will use a proof by contradiction. Assume that the vectors $(q_{i1}, q_{i2})^T$, $i \in \{1, \dots, n\}$ do not form a regular polygon. Then there has to be a $j_u \in \{j_1, \dots, j_k\}$ such that $\alpha_{j_2} - \alpha_{j_1} < \alpha_{j_{u+1}} - \alpha_{j_u}$, as otherwise we have that $\alpha_{j_2} - \alpha_{j_1} = \alpha_{j_{u+1}} - \alpha_{j_u}$ for all $j_u \in \{j_1, \dots, j_k\}$, which then means that the $(q_{j_1}, q_{j_2})^T$, $j \in \{1, \dots, n\}$ form a regular polygon with k vertices. Let

$$C_{ij}(r) = \frac{\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}}.$$

Then subtracting the identity obtained by taking $i = j_1$ in (2.3) from the identity obtained by taking $i = j_2$ in (2.3) we get that

$$0 = \left(\sum_{s \in S_{\alpha j_2}} m_s - \sum_{s \in S_{\alpha j_1}} m_s \right) C_{j_1 j_2}(r) + \sum_{v=3}^k \left(\sum_{s \in S_{\alpha j_v}} m_s \right) (C_{j_2 j_v} - C_{j_1 j_v}). \quad (3.1)$$

Note that (3.1) does not involve sums $\sum_{j \neq j_1, j \in S_{\alpha j_1}} m_j C_{j_1 j}$ and $\sum_{j \neq j_2, j \in S_{\alpha j_2}} m_j C_{j_2 j}$,

as for $\tilde{j} \neq j_1$, $\tilde{j} \in S_{\alpha j_1}$ and $\hat{j} \neq j_2$, $\hat{j} \in S_{\alpha j_2}$ we have that $\cos(\alpha_{j_1} - \alpha_{\tilde{j}}) = 1$ and $\cos(\alpha_{j_2} - \alpha_{\hat{j}}) = 1$, so by Lemma 2.5 we have that $C_{\tilde{j} j_1}$ and $C_{\hat{j} j_2}$ are linearly independent from $C_{j_1 j_2}$, $C_{j_1 j_v}$, $C_{j_2 j_v}$, $v \in \{3, \dots, k\}$ in (3.1).

The only way for any term involving $C_{j_1 j_u}$ in (3.1) to cancel out against a term $C_{j_2 j_{\tilde{v}}}$ is by Lemma 2.5 if $\cos(\alpha_{j_u} - \alpha_{j_1}) = \cos(\alpha_{j_{\tilde{v}}} - \alpha_{j_2})$, which means that $\alpha_{j_u} - \alpha_{j_1} = \alpha_{j_{\tilde{v}}} - \alpha_{j_2}$, or $\alpha_{j_u} - \alpha_{j_1} = 2\pi - (\alpha_{j_{\tilde{v}}} - \alpha_{j_2})$. If $\alpha_{j_u} - \alpha_{j_1} = \alpha_{j_{\tilde{v}}} - \alpha_{j_2}$, then $\alpha_{j_{\tilde{v}}} - \alpha_{j_u} = \alpha_{j_2} - \alpha_{j_1} < \alpha_{j_{u+1}} - \alpha_{j_u}$, giving $\alpha_{j_{\tilde{v}}} < \alpha_{j_{u+1}}$ and $\alpha_{j_{\tilde{v}}} - \alpha_{j_u} = \alpha_{j_2} - \alpha_{j_1} > 0$, giving $\alpha_{j_{\tilde{v}}} > \alpha_{j_u}$ and therefore $\alpha_{j_u} < \alpha_{j_{\tilde{v}}} < \alpha_{j_{u+1}}$, which is impossible. So that means that for any term involving $C_{j_1 j_u}$ in (3.1) to cancel out against a term $C_{j_2 j_{\tilde{v}}}$ we have that $\alpha_{j_u} - \alpha_{j_1} = 2\pi - (\alpha_{j_{\tilde{v}}} - \alpha_{j_2})$. Additionally there might be a $C_{j_1 j_w}$,

$j_w \in \{j_1, \dots, j_k\}$ such that $\cos(\alpha_{j_u} - \alpha_{j_1}) = \cos(\alpha_{j_w} - \alpha_{j_1})$. Let

$$\begin{aligned} V_{\alpha j_{\hat{v}}} &= \{s \in S_{\alpha j_{\hat{v}}} \mid \cos(\beta_s - \beta_{j_2}) = \cos(\beta_{j_{\hat{v}}} - \beta_{j_2})\}, \\ V_{\alpha j_u} &= \{s \in S_{\alpha u} \mid \cos(\beta_s - \beta_{j_1}) = \cos(\beta_{j_u} - \beta_{j_1})\} \end{aligned}$$

and $V_{\alpha j_w} = \{s \in S_{\alpha j_w} \mid \cos(\beta_w - \beta_{j_1}) = \cos(\beta_{j_w} - \beta_{j_1})\}$. We then have by (3.1) that

$$0 = \left(\sum_{s \in V_{\alpha j_{\hat{v}}}} m_s \right) - \left(\sum_{s \in V_{\alpha j_u}} m_s \right) - \left(\sum_{s \in V_{\alpha j_w}} m_s \right), \quad (3.2)$$

as $C_{j_1 j_u} = C_{j_2 j_{\hat{v}}} = C_{j_1 j_w} \neq 0$. Additionally, let

$$\widehat{S}_{ij}(r) = \frac{\sin(\alpha_j - \alpha_i)}{(1 - (\cos(\beta_j - \beta_i) + r^2(\cos(\alpha_j - \alpha_i) - \cos(\beta_j - \beta_i)))^2)^{\frac{3}{2}}}.$$

Then by (2.1) we have that

$$0 = \left(\sum_{s \in V_{\alpha j_{\hat{v}}}} m_s \right) \widehat{S}_{j_2 j_{\hat{v}}}. \quad (3.3)$$

If $\widehat{S}_{j_1 j_u} = \widehat{S}_{j_2 j_{\hat{v}}} = 0$, then $\sin(\alpha_{j_u} - \alpha_{j_1}) = 0$, which means that $\alpha_{j_u} - \alpha_{j_1} = 0$ or $\alpha_{j_u} - \alpha_{j_1} = \pi$ and $\alpha_{j_u} - \alpha_{j_1} = 2\pi - (\alpha_{j_{\hat{v}}} - \alpha_{j_2})$, so $\alpha_{j_u} - \alpha_{j_1} = \pi$ and $\alpha_{j_{\hat{v}}} - \alpha_{j_2} = \pi$, which contradicts the construction of j_u . We now therefore find the following: If there is a $\hat{v}$ such that $\cos(\alpha_{j_u} - \alpha_{j_1}) = \cos(\alpha_{j_{\hat{v}}} - \alpha_{j_2})$, then we have by (3.3) that $0 \neq 0$, which is a contradiction. If there is no such $\hat{v}$, then we have by (3.2) that $0 = 0 - \left(\sum_{s \in V_{\alpha j_u}} m_s \right) - \left(\sum_{s \in V_{\alpha j_w}} m_s \right) < 0$, which is again a contradiction. This means that our initial assumption about the existence of j_u is incorrect, which means that this proves that the configuration of the vectors $(q_{i1}, q_{i2})^T$, $i \in \{1, \dots, n\}$, is a regular polygon and by repeating the argument so far after interchanging the roles of the α s and β s, we find that the same is true for the configuration of the vectors $(q_{i3}, q_{i4})^T$. This completes the proof.

References

- [1] A.V. Borisov, I.S. Mamaev, The Restricted Two-Body Problem in Constant Curvature Spaces, *Celestial Mechanics and Dynamical Astronomy*, **96**(1): 117, (2006).

- [2] A.V. Borisov, I.S. Mamaev, A.A. Kilin, Two-Body Problem on a Sphere, Reduction, Stochasticity, Periodic Orbits, *Regular and Chaotic Dynamics*, **3** (9): 265–279, (2004).
- [3] J.F. Cariñena, M.F. Rañada and M. Santander, Central potentials on spaces of constant curvature: The Kepler problem on the two-dimensional sphere $\mathbb{S}^2$ and the hyperbolic plane $\mathbb{H}^2$, *J. Math. Phys.* **46** (2005), 052702.
- [4] Y. Deng, F. Diacu, S. Zhu, Variational property of Keplerian orbits by Maslov-type index, *J. Differential Equations* **267** (2019), no. 10, 5851–5869.
- [5] F. Diacu, On the singularities of the curved n -body problem, *Trans. Amer. Math. Soc.* **363** (2011), 2249–2264.
- [6] F. Diacu, Polygonal homographic orbits of the curved n -body problem, *Trans. Amer. Math. Soc.* **364** 5 (2012), 2783–2802.
- [7] F. Diacu, Relative equilibria in the 3-dimensional curved n -body problem, *Memoirs Amer. Math. Soc.* **228** 1071, (2013).
- [8] F. Diacu, Relative Equilibria of the Curved N -Body Problem, *Atlantis Studies in Dynamical Systems*, vol. 1, Atlantis Press, Amsterdam, 2012.
- [9] F. Diacu, The non-existence of centre-of-mass and linear-momentum integrals in the curved n -body problem, *arXiv:1202.4739*, 12 p.
- [10] F. Diacu, The curved N -body problem: risks and rewards, *Math. Intelligencer* **35** (2013), no. 3, 24–33.
- [11] F. Diacu, The classical N -body problem in the context of curved space, *Canad. J. Math.* (to appear).
- [12] F. Diacu, S. Kordlou, Rotopulsators of the curved N -body problem, *J. Differ. Equations* **255** (2013), 2709–2750.
- [13] F. Diacu, E. Pérez-Chavela and M. Santoprete, The n -body problem in spaces of constant curvature, *arXiv:0807.1747*, 54 p.
- [14] F. Diacu, E. Pérez-Chavela and M. Santoprete, The n -body problem in spaces of constant curvature. Part I: Relative equilibria, *J. Nonlinear Sci.* **22** (2012), no. 2, 247–266.

- [15] F. Diacu, E. Pérez-Chavela and M. Santoprete, The n-body problem in spaces of constant curvature. Part II: Singularities, *J. Nonlinear Sci.* **22** (2012), no. 2, 267–275.
- [16] F. Diacu, S. Popa, All the Lagrangian relative equilibria of the curved 3-body problem have equal masses, *J. Math. Phys.* **55** (2014), 112701.
- [17] F. Diacu, C. Stoica, S. Zhu, Central Configurations of the Curved N-Body Problem, *J. Nonlinear Sci.* (2018) 28:19992046 <https://doi.org/10.1007/s00332-018-9473-y>.
- [18] F. Diacu, J.M. Sánchez-Cerritos, S. Zhu, Stability of xed points and associated relative equilibria of the 3-body problem on $\mathbb{S}^1$ and $\mathbb{S}^2$, *J. Dynam. Differential Equations* 30 (2018), no. 1, 209-225. Modication after publication at arXiv:1603.03339.
- [19] F. Diacu, B. Thorn, Rectangular orbits of the curved 4-body problem, *Proc. Amer. Math. Soc.* **143** (2015), 1583–1593.
- [20] V. V. Kozlov and A. O. Harin, Keplers problem in constant curvature spaces, *Celestial Mech. Dynam. Astronom.* 54 (1992), 393-399.
- [21] H. Montanelli, N.I. Gushterov, Computing Planar and Spherical Choreographies, *SIAM J. Appl. Dyn. Syst.*, 15(1), 235-256 (2016).
- [22] E. Pérez-Chavela and J.G. Reyes Victoria, An intrinsic approach in the curved N-body problem. The positive curvature case, *Trans. Amer. Math. Soc.* 364, 7 (2012), 3805-3827.
- [23] E. Pérez-Chavela, Juan Manuel Sánchez-Cerritos, On the non-existence of hyperbolic polygonal relative equilibria for the negative curved n -body problem with equal masses, arXiv:1612.09270v1, 11p.
- [24] J.M. Sánchez-Cerritos, S. Zhang, Choreography solutions of the n-body problem on S^2 , arXiv:1806.03032, 10p.
- [25] P. Tibboel, Polygonal homographic orbits in spaces of constant curvature, *Proc. Amer. Math. Soc.* **141** (2013), 1465–1471.
- [26] P. Tibboel, Existence of a class of rotopulsators, *J. Math. Anal. Appl.* **404** (2013), 185–191.
- [27] P. Tibboel, Existence of a lower bound for the distance between point masses of relative equilibria in spaces of constant curvature, *J. Math. Anal. Appl.* **416** (2014), 205–211.

- [28] P. Tibboel, Existence of a lower bound for the distance between point masses of relative equilibria for generalised quasi-homogeneous n -body problems and the curved n -body problem, J. Math. Phys. (2015) **56**, 032901.
- [29] P. Tibboel, Finiteness of polygonal relative equilibria for generalised quasi-homogeneous n -body problems and n -body problems in spaces of constant curvature, J. Math. Anal. Appl. **441** (2016), 183-193.
- [30] P. Tibboel, Polygonal rotopulsators of the curved n -body problem, J. Math. Phys. **59** (2018), 022901.
- [31] P. Tibboel, Circular non-collision orbits for a large class of n -body problems, J. Dyn. Diff. Equat. (2018), DOI: 10.1007/s10884-018-9714-7.
- [32] S. Zhu, Eulerian relative equilibria of the curved 3-body problems in $\mathbf{S}^2$, Proc. Amer. Math. Soc. **142** (2014), 2837–2848.
- [33] X. Yu, S. Zhu, Regular Polygonal Equilibria on $\mathbb{S}^1$ and Stability of the Associated Relative Equilibria, J. Dyn. Diff. Equat. (2020), <https://doi.org/10.1007/s10884-020-09848-1>.
- [34] S. Zhao, S. Zhu, Three-dimensional central configurations in $\mathbb{H}^3$ and $\mathbb{S}^3$, J. Math. Phys. **58** (2017), no. 2, 022901.