

$\gamma N \rightarrow \Delta^+(1600)$ transition form factors in light-cone sum rules

T. M. Aliev,^{1,*} T. Barakat,^{2,†} S. Bilmis,^{1,‡} and M. Savci^{1,§}

¹*Department of Physics, Middle East Technical University, Ankara, 06800, Turkey*

²*Department of Physics, King Saud University, Riyadh, 11451, Saudi Arabia*

(Dated: May 18, 2022)

Abstract

The form factors of $\gamma^* N \rightarrow \Delta(1600)$ transition is calculated within the light-cone sum rules assuming that $\Delta^+(1600)$ is the first radial excitation of $\Delta(1232)$. The Q^2 dependence of the magnetic dipole $\tilde{G}_M(Q^2)$, electric quadrupole $\tilde{G}_E(Q^2)$, and Coulomb quadrupole $\tilde{G}_c(Q^2)$ form factors are investigated. Moreover, the Q^2 dependence of the ratios $R_{EM} = -\frac{\tilde{G}_E(Q^2)}{\tilde{G}_M Q^2}$ and $R_{SM} = -\frac{1}{4m_{\Delta(1600)}^2} \sqrt{4m_{\Delta(1600)}^2 Q^2 + (m_{\Delta(1600)}^2 - Q^2 - m_N^2)^2} \frac{\tilde{G}_c(Q^2)}{\tilde{G}_M(Q^2)}$ are studied. Finally, our predictions on $\tilde{G}_M(Q^2)$, $\tilde{G}_E(Q^2)$, and $\tilde{G}_C(Q^2)$ are compared with the results of other theoretical approaches.

* taliev@metu.edu.tr

† tbarakat@ksu.edu.sa

‡ sbilmis@metu.edu.tr

§ savci@metu.edu.tr

I. INTRODUCTION

Advance technologies in accelerators enabled to search of high energy regions as well as improving the precision of the experiments by collecting data with high luminosity. This reveals new possibilities to study the electromagnetic structures of the baryon resonances above the ground state region. The facilities like CLAS (Jefferson Lab), BATES (MIT), MAMI (Mainz), Spring-8 (Japan), and ESSA (Bonn) have the potential to measure the electromagnetic structures of baryons around their first excitations. These experimental possibilities stimulated theoretical studies for deeply understanding the properties of the baryon resonances. $\Delta(1600)$ baryon, which is the first excitation of $\Delta(1200)$ one, may be one of the resonances which deserves special attention. Theoretically, this resonance has not been studied comprehensively yet.

The electroproduction of $\Delta(1600)$ is studied within the quark model [1], and the effects of $\Delta(1600)$ in baryon-meson reactions is studied in [2, 3]. However, the existing data [2, 4] can be used for a more detailed analysis of this resonance. The $\gamma^*p \rightarrow \Delta^+(1232)$, ($\Delta^+(1600)$) transitions are computed using a diquark-quark picture and a covariant spectator constituent quark model in [5] and [6], respectively.

The form factors of the $\gamma^*N \rightarrow \Delta(1232)$ and γ^* octet \rightarrow decuplet baryon transitions within the same framework was studied in [7] and [8], respectively. And in this work, we study the transition form factors for the electroproduction of the $\Delta(1600)$ resonance within light-cone sum rules method.

The article is organized as follows. In section II, the sum rules for the transition form-factors of $\gamma^*N \rightarrow \Delta(1600)$ within the light-cone sum rules (LCSR) is derived. The numerical analysis of the obtained LCSRs is carried out in section III. This section also contains discussion and summary.

II. DERIVATION OF LCSR FOR $\gamma^*N \rightarrow \Delta(1600)$ TRANSITION FORM FACTORS

The transition $\gamma^*N \rightarrow \Delta(1200)$ and $\Delta(1600)$ is described by the matrix element of the electromagnetic current $j_\mu = e_u \bar{u} \gamma_\mu u + e_d \bar{d} \gamma_\mu d$ between the nucleon, ground, and first radial excitation of Δ baryon $\langle \Delta_i(p') | j_\nu(0) | N(p) \rangle$.

By using the Lorentz invariance and current conservation, this matrix element is deter-

mined in terms of the following form factors [9]:

$$\begin{aligned} \langle \Delta_i(p' | j_\mu^{el}) | N(p) \rangle = & \bar{u}_\alpha(p') \left\{ G_1^{(i)}(Q^2) (-q_\alpha \gamma_\mu + \not{q} g_{\alpha\mu}) + G_2^{(i)}(Q^2) (-q_\alpha \mathcal{P}_\mu + (q\mathcal{P}) g_{\alpha\mu}) \right. \\ & \left. + G_3^{(i)}(q_\alpha q_\mu - q^2 g_{\alpha\mu}) \right\} \gamma_5 u(p) \end{aligned} \quad (1)$$

where $i = 0$, and $i = 1$ corresponds to the ground and first radial excitation of Δ baryon, G_1^i , G_2^i , and G_3^i are the corresponding form-factors, and $\mathcal{P}_\alpha = \frac{1}{2}(p + p')_\alpha = \frac{1}{2}(2p - q)_\alpha$. However, the multipole form factors are more useful than the form factors G_1 , G_2 , and G_3 for the experimental point of view. The relations among the form factors $G_1(Q^2)$, $G_2(Q^2)$, and $G_3(Q^2)$ and multipole form factors (magnetic dipole G_M , electric quadrupole G_E , and Coulomb quadrupole G_C) form factors are given in [10]:

$$\begin{aligned} G_M^{(i)}(Q^2) &= \frac{m_N}{3(m_N + m_{\Delta_i})} \left[((3(m_{\Delta_i} + m_N)(m_{\Delta_i} + m_N) + Q^2) \frac{G_1^{(i)}(Q^2)}{m_{\Delta_i}} \right. \\ &\quad \left. + (m_{\Delta_i}^2 - m_N^2) G_2^{(i)}(Q^2) - 2Q^2 G_3^{(i)}(Q^2) \right], \\ G_E^{(i)}(Q^2) &= \frac{m_N}{3(m_N + m_{\Delta_i})} \left[(m_{\Delta_i} - m_N^2 - Q^2) \frac{G_1^{(i)}(Q^2)}{m_{\Delta_i}} \right. \\ &\quad \left. + (m_{\Delta_i}^2 - m_N^2) G_2^{(i)}(Q^2) - 2Q^2 G_3^{(i)}(Q^2) \right], \\ G_C^{(i)}(Q^2) &= \frac{2m_N}{3(m_N + m_{\Delta_i})} \left[2m_{\Delta_i} G_1^{(i)}(Q^2) + \frac{1}{2}(3m_{\Delta_i}^2 + m_N^2 + Q^2) G_2^{(i)}(Q^2) \right. \\ &\quad \left. + (m_{\Delta_i}^2 - m_N^2 - Q^2) G_3^{(i)}(Q^2) \right]. \end{aligned} \quad (2)$$

After these preliminary remarks, we can proceed with the determination of these form factors for $\gamma^* N \rightarrow \Delta(1600)$ transitions within the light-cone QCD sum rules. For this purpose, we consider the following correlator function

$$\Pi_{\alpha\mu} = i \int d^4x e^{iqx} \langle T \{ \eta_\alpha(0) j_\mu^{el}(x) \} N(p) \rangle, \quad (3)$$

where η_α is the interpolating current with the same quantum numbers of $\Delta(1232)$ and $\Delta(1600)$, and j_μ^{el} is the electromagnetic current.

Since $\Delta^+(1232)$ and $\Delta^+(1600)$ states have the same quantum numbers, the interpolating current for these states is also same and it is given by the following expression

$$\eta_\alpha = \frac{1}{\sqrt{3}} \epsilon^{abc} [2(u^a C \gamma_\alpha d^b) u^c + (u^a C \gamma_\alpha u^b) d^c], \quad (4)$$

where a, b, c are color indices, and C is the charge conjugation operator. According to the sum rules method approach, the correlation function should be calculated in two different

regions. In one domain, the correlation function is saturated by the full tower of states carrying the quantum numbers of Δ baryon in the region $p^2 \simeq m_{\Delta_i}^2$. On the other hand, the correlation function is calculated in the deep Euclidean region where $(p+q)^2 \ll 0$ by using the operator product expansion (OPE) in terms of the nucleon distribution amplitudes with an increasing twist. The sum rules for the relevant physical quantities are obtained by matching these results of the representations of the correlation functions via the dispersion relation.

Following the mentioned prescription above and for the hadronic part of the correlation function after isolating the contributions of the ground $\Delta(1232)$, and its first radial excitation $\Delta(1600)$ state we get

$$\Pi_{\alpha\mu} = - \sum_{i=1}^2 \frac{\langle 0 | \eta_\alpha(0) | \Delta_i \rangle \langle \Delta_i | j_\mu^{el} | N \rangle}{m_{\Delta_i}^2 - p'^2} + \int_{s_0}^{\infty} ds \frac{\Pi_{\alpha\mu}^{\text{had}}(s)}{s - p'^2}, \quad (5)$$

where i corresponds to the ground and first excited states. Parameterizing the matrix element

$$\langle 0 | \eta_\alpha | \Delta_i(p') \rangle = \lambda_i u_\alpha(p'), \quad (6)$$

where $u_\alpha(p')$ is the Rarita-Schwinger spinor and $p' = p - q$. Performing summation over the spins of Rarita-Schwinger spinors with the help of the formula

$$\sum_s u_\alpha^{(s)}(p') \bar{u}_\beta^{(s)}(p') = -(\not{p}' + m_{\Delta_i}) \left\{ g_{\alpha\beta} - \frac{1}{3} \gamma_\alpha \gamma_\beta - \frac{2p'_\alpha p'_\beta}{3m_{\Delta_i}^2} + \frac{p'_\alpha \gamma_\beta - p'_\beta \gamma_\alpha}{3m_{\Delta_i}} \right\} \quad (7)$$

we get the following result for the correlation function

$$\begin{aligned} \Pi_{\alpha\mu} = & \sum_i \frac{\lambda_i}{m_{\Delta_i}^2 - p'^2} (\not{p}' + m_{\Delta_i}) \left\{ g_{\alpha\beta} - \frac{1}{3} \gamma_\alpha \gamma_\beta - \frac{2p'_\alpha p'_\beta}{3m_{\Delta_i}^2} + \frac{p'_\alpha \gamma_\beta - p'_\beta \gamma_\alpha}{3m_{\Delta_i}} \right\} \\ & \{ G_1^i(-q_\beta \gamma_\mu + g_{\beta\mu} \not{q}) + G_2^i(-q_\beta \mathcal{P}_\mu + g_{\beta\mu} \not{q} \mathcal{P}) + \widetilde{G}_3^i(q_\beta q_\mu - g_{\beta\mu} q^2) \} \gamma_5 u_N(p). \end{aligned} \quad (8)$$

At this point, we would like to make the following remark. In general, the interpolating current, η_α , interacts not only with spin-3/2 states, but also with the spin-1/2 ones. For the generic spin-1/2 states, the matrix element of the η_μ current between the vacuum and spin 1/2 state is determined as

$$\langle 0 | \eta_\alpha | \frac{1}{2}(p') \rangle = (m\gamma_\alpha - 4p'_\alpha) u(p'). \quad (9)$$

In other words, the terms with $\sim \gamma_\alpha$ and p'_α contain the contributions of the spin-1/2 states. Comparing Eqs. (8) and (9), it follows that only the terms with $\sim g_{\alpha\beta}$ contains the

information of purely spin-3/2 states. Hence, for our problem the terms containing spin-1/2 contributions should be removed. Retaining the contributions of spin-3/2 states only, we get

$$\begin{aligned}
\Pi_{\alpha\mu} = & \frac{\lambda_0}{m_0^2 - (p-q)^2} (\not{p} - \not{q} + m_0) [G_1(-q_\alpha\gamma_\mu + g_{\alpha\mu}\not{q}) \\
& + G_2[-q_\alpha(p-q/2)_\mu + q \cdot (p-q/2)g_{\alpha\mu}] + G_3[q_\alpha q_\mu - q^2 g_{\alpha\mu}] \gamma_5 u_N(p)] \\
& + \frac{\lambda_1}{m_1^2 - (p-q)^2} (\not{p} - \not{q} + m_1) [\widetilde{G}_1(-q_\alpha\gamma_\mu + g_{\alpha\mu}\not{q}) \\
& + \widetilde{G}_2[-q_\alpha(p-q/2)_\mu + q \cdot (p-q/2)g_{\alpha\mu}] + \widetilde{G}_3[q_\alpha q_\mu - q^2 g_{\alpha\mu}] \gamma_5 u_N(p)].
\end{aligned} \tag{10}$$

in which λ_0 , m_0 (λ_1 , m_1) are the residue and mass of the ground state, $\Delta(1232)$, $\Delta(1600)$ baryons and G_i (\widetilde{G}_i) are the form factors for $\gamma^*N \rightarrow \Delta(1232)$ and $\gamma^*N \rightarrow \Delta(1600)$ transitions, respectively. For simplicity, the mass of the $\Delta(1600)$ state we will be denote as m_1 from now on.

From Eq. (10), it follows that, for the description $\gamma^*N \rightarrow \Delta(1600)$ transition we have six form factors which should be determined. To determine the six form factors, we need six structures. It should be noted that all structures are not independent. To obtain the independent structures, the ordering procedure of the Dirac matrices is implemented. In this work, we choose the following order of Dirac matrices $\gamma_\alpha \not{p} \not{q} \gamma_\mu \gamma_5$. Taking into account this remark, the correlation function can be decomposed in terms of the following independent invariant functions as follows (see Eq.(8)):

$$\begin{aligned}
\Pi_{\alpha\mu} = & \Pi_1 \not{q} \gamma_\mu \gamma_5 q_\alpha + \Pi_2 \gamma_\mu \gamma_5 q_\alpha + \Pi_3 \not{q} \gamma_5 p_\mu q_\alpha + \Pi_4 \gamma_5 p_\mu q_\alpha + \Pi_5 \not{q} \gamma_5 q_\alpha q_\mu \\
& + \Pi_6 \gamma_5 q_\alpha q_\mu + \text{other structures}
\end{aligned} \tag{11}$$

From Eqs. (9) and (10), the following six structures are found to determine the six form

factors

$$\begin{aligned}
\Pi_1 &= \frac{\lambda_0 G_1}{m_0^2 - (p-q)^2} + \frac{\lambda_1 \widetilde{G}_1}{m_1^2 - (p-q)^2}, \\
\Pi_2 &= -\frac{\lambda_0 m_0 G_1}{m_0^2 - (p-q)^2} - \frac{\lambda_1 m_1 \widetilde{G}_1}{m_1^2 - (p-q)^2}, \\
\Pi_3 &= \frac{\lambda_0 G_2}{m_0^2 - (p-q)^2} + \frac{\lambda_1 \widetilde{G}_2}{m_1^2 - (p-q)^2}, \\
\Pi_4 &= -\frac{\lambda_0 m_0 G_2}{m_0^2 - (p-q)^2} - \frac{\lambda_1 m_1 \widetilde{G}_2}{m_1^2 - (p-q)^2}, \\
\Pi_5 &= -\frac{\lambda_0}{m_0^2 - (p-q)^2} \left[\frac{G_2}{2} + G_3 \right] - \frac{\lambda_1}{m_1^2 - (p-q)^2} \left[\frac{\widetilde{G}_2}{2} + \widetilde{G}_3 \right], \\
\Pi_6 &= \frac{\lambda_0 m_0}{m_0^2 - (p-q)^2} \left[\frac{G_2}{2} + G_3 \right] + \frac{\lambda_1 m_1}{m_1^2 - (p-q)^2} \left[\frac{\widetilde{G}_2}{2} + \widetilde{G}_3 \right].
\end{aligned} \tag{12}$$

Solving these equations for the form factors we get

$$\begin{aligned}
m_0 \Pi_1 + \Pi_2 &= -\frac{\lambda_1 \widetilde{G}_1}{m^{*2} - (p-q)^2} (m_1 - m_0), \\
m_0 \Pi_3 + \Pi_4 &= -\frac{\lambda_1 \widetilde{G}_2}{m^{*2} - (p-q)^2} (m_1 - m_0), \\
m_0 \Pi_5 + \Pi_6 &= \frac{\lambda_1}{m^{*2} - (p-q)^2} (m_1 - m_0) \left[\frac{\widetilde{G}_2}{2} + \widetilde{G}_3 \right].
\end{aligned} \tag{13}$$

From Eq.(13), it follows that to obtain the sum rules for the $\gamma^* N \rightarrow \Delta(1600)$ transition form factors, we need to know the invariant functions Π_i . According to the sum rules methodology, the invariant functions $\Pi_i (i = 1 \div 6)$ are calculated at deep Euclidean domain with virtuality $p'^2 = (p-q)^2 \ll 0$ in terms of the nucleon distribution amplitudes (DA's). The nucleon DA's are the main non-perturbative ingredients of LCSR and they are calculated up to twist-6 in [11]. For completeness, definition of the nucleon's DA's and their expressions are presented in Appendix.

Using the expressions of the nucleon DA's and applying the quark-hadron duality ansatz, the invariant functions, Π_i , can be written in the following form

$$\Pi_i((p-q)^2, q^2) = \sum_{n=1}^3 \int_0^1 \frac{\rho_i^{(n)}(x, q^2, (p-q)^2)}{((q-px)^2)^n} \tag{14}$$

Matching the representations of the correlation functions and performing Borel transformations with respect to the variable $-(p-q)^2$ in order to suppress the contributions of higher

states and continuum, the corresponding sum rules for the form-factors $\widetilde{G}_1(Q^2)$, $\widetilde{G}_2(Q^2)$ and $\frac{\widetilde{G}_2(Q^2)}{2} + \widetilde{G}_3(Q^2)$ can be obtained as

$$\begin{aligned}
-\lambda_1 \widetilde{G}_1(Q^2)(m_1 - m_0)e^{-m_1^2/M^2} &= m_0 I_1(M^2, Q^2, s_0) + I_2(Q^2, M^2, s_0), \\
-\lambda_1 \widetilde{G}_2(Q^2)(m_1 - m_0)e^{-m_1^2/M^2} &= m_0 I_3(M^2, Q^2, s_0) + I_4(Q^2, M^2, s_0), \\
\lambda_1 \left(\frac{\widetilde{G}_2}{2} + \widetilde{G}_3^* \right) (m_1 - m_0)e^{-m_1^2/M^2} &= m_0 I_5(M^2, Q^2, s_0) + I_6(Q^2, M^2, s_0).
\end{aligned} \tag{15}$$

The functions $I_i(M^2, Q^2, s_0)$ can be written in the form of a master formula (see [12] and [13])

$$\begin{aligned}
I_i &= \sum_{n=1}^{\infty} \int_{x_0}^1 dx e^{-s/M^2} \frac{1}{(n-1)!} \frac{\rho_i^{(n)}}{x^n (M^2)^{n-1}} \\
&- e^{-s_0/M^2} \left[\frac{(-1)^{n-1}}{(n-1)!} \sum_{j=1}^{n-1} \frac{1}{(M^2)^{n-j-1}} \frac{1}{s'} \left(\frac{d}{dx} \frac{1}{s'} \right)^{j-1} \frac{\rho_i^{(n)}}{x^n} \right]_{|x=x_0}
\end{aligned} \tag{16}$$

where $\bar{x} = 1 - x$, $s' = \frac{ds}{dx}$, $s = \frac{m_N^2 \bar{x} x + Q^2 \bar{x}}{x}$, and x_0 is the solution of $s_0 = s$ equation. The explicit forms of the functions, ρ_i^n are presented in the Appendix. From Eq.(13), we see that to determine the $\gamma^* N \rightarrow \Delta(1600)$ transition form factors, the residue of $\Delta(1600)$ is also needed. This value within QCD sum rules is already calculated in [14] and obtained as $\lambda_1 = (0.057 \pm 0.016) \text{ GeV}^3$.

At the end of this section, we present the ratios R_{EM} [10] and R_{SM} [11] that are more suitable for the experimental point of view

$$\begin{aligned}
R_{EM} &= -\frac{\widetilde{G}_E(Q^2)}{\widetilde{G}_M(Q^2)}, \\
R_{SM} &= -\sqrt{Q^2 + \frac{m_1^2 - m_N^2 - Q^2)^2}{4m_1^2}} \frac{1}{2m_1} \frac{\widetilde{G}_C(Q^2)}{\widetilde{G}_M(Q^2)}.
\end{aligned} \tag{17}$$

It should be noted that these ratios are identically zero in SU(6) symmetric constituent quark model. The nonzero values are the indications of the deformation of one or both hadrons.

III. NUMERICAL ANALYSIS

This section is devoted to the numerical analysis of the multipole form factors as well as R_{EM} and R_{SM} ratios. The main non-perturbative input parameters of LCSR are the

DA's. The nucleon DA's and the parameters entering in their expressions are determined in [11] (see also [15, 16]). In addition to these input parameters, the sum rules contain two auxiliary parameters; the Borel parameter M^2 and continuum threshold s_0 . The working region of M^2 is determined by using the dominance of the perturbative contributions over the nonperturbative and continuum ones. Besides, the values of continuum threshold s_0 is determined from the condition that the sum rules should reproduce the mass of $\Delta(1600)$ state with 10% accuracy. These conditions lead to the following working regions of M^2 and s_0 : $1.5 \text{ GeV}^2 \leq M^2 \leq 3 \text{ GeV}^2$, $s_0 = (4.5 \pm 0.5 \text{ GeV}^2)$.

Having specified all the input parameters and determined the working region of M^2 and s_0 , we are ready to perform the numerical calculations.

In Figures, 2, 3, and 4, we present the dependencies of $\tilde{G}_M(Q^2)$, $\tilde{G}_E(Q^2)$ and $\tilde{G}_C(Q^2)$ on Q^2 at fixed s_0 and for various M^2 values. Here, we would like to note that since LCSR predictions on the form factors are reliable only in the $Q^2 > 1 \text{ GeV}^2$ region, we present the results only for this domain. From these figures, it follows that all three form factors decrease with increasing Q^2 and saturates for high Q^2 values.

By comparing the form factors of $\gamma^*N \rightarrow \Delta(1232)$ obtained in [7] and $\gamma^*N \rightarrow \Delta(1600)$ transitions, we infer the following results :

- The electric quadrupole form factor is very small in magnitude compared to the form factors, $\tilde{G}_M(Q^2)$ and $\tilde{G}_c(Q^2)$ in both transitions.
- For the region, $Q^2 > 1 \text{ GeV}^2$, the transition form factors, $\tilde{G}_M(Q^2)$ for $\gamma^*N \rightarrow \Delta(1600)$ are larger than the ones for $\gamma^*N \rightarrow \Delta(1232)$.

Moreover, we also compared our results on the considered form factors with the predictions obtained by quark-diquark approximation to the Poincare-covariant three-body bound state problem in relativistic quantum field theory [5] and found out that our predictions on the form factors at the considered regions of Q^2 is around 2 times larger in magnitude than the one predicted in [5].

Furthermore in Figures 5 and 6, we present the Q^2 dependence of $R_{EM}(Q^2)$ and R_{SM} at fixed values M^2 and s_0 considering their working regions. From these figures, we observe that while $R_{SM}(Q^2)$ is negative, $R_{EM}(Q^2)$ is positive at all values of Q^2 .

Comparing our predictions on R_{SM} with the results obtained in [7], we observed similar qualitative behaviour considered in both works. However, behavior of R_{EM} in our case is

remarkably different than in $\gamma^*N \rightarrow \Delta(1232)$ transition, i.e. magnitude R_{EM} is larger than the one in $\gamma^*N \rightarrow \Delta(1232)$ transition case. Finally, we compare our predictions on R_{SM} and R_{EM} with the results obtained within light-front relativistic quark model[17]. From Fig. 5, it follows that when Q^2 varies in the region $Q^2 = 5 \text{ GeV}^2$ to 12 GeV^2 , R_{EM} practically does not change and the value is around 0.22. However, the prediction of [17] on R_{EM} grows from 0.06 to 0.1. In other words, our result on R_{EM} is larger than the predicted in [17]. Besides, comparing our result on R_{SM} , we deduce that with increasing Q^2 , R_{SM} in our case grows mild, but Q^2 dependence is considerable [17]. For example, in our case, when Q^2 varies between 2 to 10 GeV^2 region, the R_{SM} varies between (0.20 – 0.25), however it changes between (0.1 – 0.3) in [17].

Our final note is that the obtained results will shed light to the understanding the inner structures of the resonance $\Delta(1600)$, and can be checked in ongoing and planning experiments.

IV. CONCLUSION

In this article, we studied the LCSR to evaluate the magnetic dipole $\tilde{G}_M(Q^2)$ electric quadrupole $\tilde{G}_E(Q^2)$ and Coulomb quadrupole $\tilde{G}^*(Q^2)$ form factors as well as the ratios $R_{EM} = -\frac{\tilde{G}_E}{\tilde{G}_M}$ and $R_{SM} = -\frac{1}{4m_1^2} \sqrt{4m_1^2 Q^2 + (m_1^2 - Q^2 - m_N^2)^2} \frac{\tilde{G}_c(Q^2)}{\tilde{G}_M(Q^2)}$ on Q^2 when Q^2 varies in the region $1 \text{ GeV}^2 \leq Q^2 \leq 10 \text{ GeV}^2$. This domain may be covered in the incoming CLAS-12 at the Jefferson Lab. Appearance of experimental information would be very useful to establish the nature of $\Delta^+(1600)$ resonance by assuming it as radial excitation of $\Delta(1232)$ in the $\gamma^*N \rightarrow \Delta(1600)$ transition. We also compared our predictions on the form factors $\tilde{G}_M(Q^2)$, $\tilde{G}_E(Q^2)$, and $\tilde{G}_C(Q^2)$, as well as R_{EM} and R_{SM} with results of results of other theoretical approaches.

ACKNOWLEDGMENT

One of the authors, T.Barakat extends his appreciation to the Deanship of Scientific Research at King Saud University for funding his work through Research Grant No: RG-1440-090.




FIG. 1.

FIG. 2. The dependency of the $G_M(Q^2)$ on Q^2 at a fixed values of s_0 and M^2 .




FIG. 3. The same as in Fig. 2, but for $G_E(Q^2)$ form factor.




FIG. 4. The same as in Fig. 2, but for $G_C(Q^2)$ form factor.




FIG. 5. The dependency of R_{EM} on Q^2 at the fixed values of s_0 and M^2 .




FIG. 6. The same as in Fig. 5, but for R_{SM} .

-
- [1] S. Capstick and B. D. Keister, Phys. Rev. **D51**, 3598 (1995), arXiv:nucl-th/9411016 [nucl-th].
- [2] V. D. Burkert, V. I. Mokeev, and B. S. Ishkhanov, Moscow Univ. Phys. Bull. **74**, 243 (2019), [Vestn. Mosk. Univ. Ser. III Fiz. Astron.74,no.3,28(2019)], arXiv:1901.09709 [nucl-ex].
- [3] B. Golli and S. Sirca, Eur. Phys. J. **A38**, 271 (2008), arXiv:0708.3759 [hep-ph].
- [4] A. Trivedi, *Proceedings, 11th International Workshop on the Physics of Excited Nucleons (NSTAR 2017): Columbia, SC, USA, August 20-23, 2017*, Few Body Syst. **60** (2019), 10.1007/s00601-018-1471-y.
- [5] Y. Lu, C. Chen, Z.-F. Cui, C. D. Roberts, S. M. Schmidt, J. Segovia, and H. S. Zong, Phys. Rev. **D100**, 034001 (2019), arXiv:1904.03205 [nucl-th].
- [6] G. Ramalho and K. Tsushima, Phys. Rev. **D82**, 073007 (2010), arXiv:1008.3822 [hep-ph].
- [7] V. M. Braun, A. Lenz, G. Peters, and A. V. Radyushkin, Phys. Rev. **D73**, 034020 (2006), arXiv:hep-ph/0510237 [hep-ph].
- [8] T. M. Aliev, Y. Oktem, and M. Savci, Phys. Rev. **D88**, 036002 (2013), arXiv:1308.0697 [hep-ph].
- [9] R. C. E. Devenish, T. S. Eisanschitz, and J. G. Korner, Phys. Rev. **D14**, 3063 (1976).
- [10] H. F. Jones and M. D. Scadron, Annals Phys. **81**, 1 (1973).
- [11] V. Braun, R. J. Fries, N. Mahnke, and E. Stein, Nucl. Phys. **B589**, 381 (2000), [Erratum: Nucl. Phys.B607,433(2001)], arXiv:hep-ph/0007279 [hep-ph].
- [12] N. Gubernari, A. Kokulu, and D. van Dyk, JHEP **01**, 150 (2019), arXiv:1811.00983 [hep-ph].
- [13] T. M. Aliev, H. Dag, A. Kokulu, and A. Ozpineci, Phys. Rev. **D100**, 094005 (2019), arXiv:1908.00847 [hep-ph].
- [14] T. M. Aliev, K. Azizi, and H. Sundu, Eur. Phys. J. **C77**, 222 (2017), arXiv:1612.03661 [hep-ph].
- [15] V. M. Braun, A. Lenz, and M. Wittmann, Phys. Rev. **D73**, 094019 (2006), arXiv:hep-ph/0604050 [hep-ph].
- [16] I. V. Anikin, V. M. Braun, and N. Offen, Phys. Rev. **D88**, 114021 (2013), arXiv:1310.1375 [hep-ph].
- [17] I. G. Aznauryan and V. D. Burkert, Phys. Rev. **C92**, 035211 (2015), arXiv:1506.03183 [hep-ph].

APPENDIX

In this Appendix we present the explicit expressions of the functions ρ_i^n entering to the sum rules for the form factors $\tilde{G}_1(Q^2)$, $\tilde{G}_2(Q^2)$ and $\tilde{G}_3(Q^2)$ for the $\gamma^*N \rightarrow \Delta(1600)$ transition.

V. FUNCTIONS $\rho_i^{(n)}$ FOR THE FORM FACTOR \tilde{G}_1

$$\rho_1^{(3)}(x) = 16e_{q_2}m_N^2(1-x) \left[(x^2m_N^2 + Q^2)\tilde{B}_6 - m_{q_2}m_Nx\tilde{B}_8 \right](x) + 8e_{q_3}m_{q_3}m_N^3x(1-x)(\hat{\tilde{D}}_6 + 2\hat{\tilde{C}}_6)(x) ,$$

$$\rho_2^{(3)}(x) = 16e_{q_2}m_N^2(1-x)m_{q_2}(Q^2\tilde{B}_6 + m_N^2x\tilde{B}_6)(x) - 8e_{q_3}m_{q_3}m_N^2(1-x)(m_N^2x + Q^2)\hat{\tilde{B}}_6(x) ,$$

$$\begin{aligned} \rho_1^{(2)}(x) &= 4e_{q_2}m_N^2 \left[(1-2x)\tilde{B}_6 \right](x) + 4e_{q_3}m_N^2 \left[x\hat{\tilde{B}}_6 \right](x) + 4e_{q_1}m_N \left[m_N(1-x)x(\check{D}_5 + \check{C}_5) \right](x) \\ &\quad + \frac{4e_{q_2}m_N}{x} \left\{ (1-x) \left[m_Nx^2(\tilde{D}_5 + \tilde{C}_5 - 2\tilde{H}_1 - 2\tilde{E}_1) - 2m_Nx^2\tilde{B}_5 \right] \right. \\ &\quad \left. - 2m_{q_2}x \left[\tilde{B}_2 - (1-x)\tilde{B}_4 \right] \right\}(x) \\ &\quad + 2e_{q_3}m_N \left\{ (1-x) \left[m_Nx(\hat{D}_4 - \hat{D}_5 + 2\hat{C}_4 - 2\hat{C}_5) \right] + m_{q_3} \left[2x(\hat{D}_2 + 2\hat{C}_2) + 2(\hat{B}_2 - \hat{B}_4) \right] \right\}(x) \\ &\quad - 8e_{q_2}m_N^2 \int_0^{\bar{x}} dx_1 \left[A_1^M - (1+x)T_1^M \right](x_1, x, 1-x_1-x) \\ &\quad - 4e_{q_3}m_N^2 \int_0^{\bar{x}} dx_1 \left[xA_1^M + 2(xV_1^M + T_1^M) \right](x_1, 1-x_1-x, x) , \end{aligned}$$

$$\begin{aligned} \rho_2^{(2)}(x) &= 8e_{q_1}m_N^3(1-x)\check{C}_6(x) + 4e_{q_2}m_N^2 \left[+ 2m_N(1-x)(\tilde{C}_6 - \tilde{B}_8) + 3m_{q_2}\tilde{B}_6 \right](x) \\ &\quad + 4e_{q_3}m_N^2 \left[- 2m_N(1-x)\hat{\tilde{B}}_8 - 2m_{q_3}\hat{\tilde{B}}_6 \right](x) + 4e_{q_1}m_N \left[2(1+x)\check{D}_2 \right](x) \\ &\quad + \frac{4e_{q_2}m_N}{x} \left\{ (1-x) \left[2x(m_N^2x + Q^2)\tilde{D}_2 + m_N^2x^2(\tilde{B}_2 + \tilde{B}_4) + Q^2(\tilde{B}_2 - \tilde{B}_4) + 2Q^2x\tilde{B}_4 \right] \right. \\ &\quad \left. - 2m_{q_2}x \left[- (1-x)(m_N\tilde{B}_5) \right] \right\}(x) \\ &\quad + 2e_{q_3}m_N \left\{ (1-x) \left[2\hat{C}_4 - 2\hat{C}_5 \right] - 2(m_N^2x + Q^2)(\hat{B}_2 + \hat{B}_4) \right\} \\ &\quad + m_{q_3} \left[m_N(1-x)(\hat{D}_4 + \hat{D}_5 - 2\hat{C}_4 - 2\hat{C}_5) \right] \right\}(x) \end{aligned}$$

$$\rho_1^{(1)}(x) = -4e_{q_2} \int_0^{\bar{x}} dx_1 \left[2D_1 - 2(1+x)B_1 \right](x_1, x, 1-x_1-x)$$

$$\begin{aligned}
& -4e_{q_3} \int_0^{\bar{x}} dx_1 \left[xD_1 + 2xC_1 + 2B_1 \right] (x_1, 1 - x_1 - x, x) . \\
\rho_2^{(1)}(x) = & -4e_{q_1} m_N (\check{D}_2 + \check{C}_2)(x) + \frac{4e_{q_2} m_N}{x} \left[\tilde{B}_2 - \tilde{B}_4 + x(\tilde{D}_2 - \tilde{C}_2 + 2\tilde{B}_4) \right] (x) \\
& + 4e_{q_1} m_N (1-x) \int_0^{\bar{x}} dx_3 \left[D_3 - C_3 \right] (x, 1-x-x_3, x_3) \\
& - 4e_{q_2} \int_0^{\bar{x}} dx_1 \left[-m_N(1-x)(D_3 - C_3 + 2P_1 - 2S_1) + 2m_{q_2} B_1 \right] (x_1, x, 1-x_1-x) \\
& - 4e_{q_3} \int_0^{\bar{x}} dx_1 \left[m_N(1-x)D_3 - 2m_N(1-x)C_3 m_{q_3} (D_1 - 2C_1) \right] (x_1, 1-x_1-x, x) .
\end{aligned} \tag{1}$$

VI. FUNCTIONS ρ_i FOR THE FORM FACTOR \tilde{G}_2

$$\begin{aligned}
\rho_3^{(3)}(x) = & 64e_{q_1} m_N^3 (1-x)x^2 \check{C}_6(x) \\
& + 16e_{q_2} m_N^2 \left\{ 2(1-x) \left[m_N x^2 (2\tilde{C}_6 - 4\tilde{B}_8) \right] + m_{q_2} x \tilde{B}_6 \right\} (x) \\
& - 16e_{q_3} m_N^2 x \left\{ 2(1-x) \left[m_N x \hat{D}_6 - m_N x (2\hat{C}_6 - 2\hat{B}_8) \right] + m_{q_3} \hat{B}_6 \right\} (x) , \\
\rho_4^{(3)}(x) = & 16e_{q_2} m_N^2 \left\{ 2(1-x) \left[Q^2(1-2x) \tilde{B}_6 - m_N^2 x^2 \tilde{B}_6 \right] - m_{q_2} x \left[2m_N(1-x) \tilde{B}_8 \right] \right\} (x) \\
& - 16e_{q_3} m_N^2 x \left\{ 2(1-x) \left[Q^2 \hat{B}_6 + m_N^2 x \hat{B}_6 \right] - m_{q_3} \left[m_N(1-x) (\hat{D}_6 + 2\hat{C}_6) \right] \right\} (x) , \\
\rho_3^{(2)}(x) = & 8e_{q_1} m_N x \left[(1-2x) \check{C}_2 \right] (x) \\
& + 8e_{q_2} m_N \left\{ x \left[\tilde{D}_2 + (1-2x)(\tilde{C}_2 - 2\tilde{B}_4) + 2\tilde{B}_2 \right] \right\} (x) \\
& + 8e_{q_3} m_N x \left\{ x \hat{D}_2 - 2 \left[x \hat{C}_2 - (1-x)(-\hat{B}_4) \right] \right\} (x) , \\
\rho_4^{(2)}(x) = & 8e_{q_2} m_N^2 (3-4x) \tilde{B}_6(x) + 8e_{q_1} m_N x \left[m_N(1-x) \check{D}_5 - m_N(1-x) \check{C}_4 \right] (x) \\
& + 8e_{q_2} m_N \left\{ x \left[m_N(1-x) \tilde{D}_5 - m_N \tilde{C}_4 + m_N x \tilde{C}_4 - 2m_N(1-x) (\tilde{H}_1 + \tilde{E}_1 - \tilde{B}_5) \right] \right. \\
& \left. - m_{q_2} \left[\tilde{B}_2 - (1-2x) \tilde{B}_4 \right] \right\} (x) \\
& + 8e_{q_3} m_N x \left\{ -m_N(1-x) \hat{D}_5 - 2 \left[-(1-x)(m_N \hat{B}_5 - m_N \hat{C}_5) \right] \right. \\
& \left. + m_{q_3} (\hat{D}_2 + 2\hat{C}_2) \right\} (x) - 8e_{q_1} m_N^2 (1-2x) \int_0^{\bar{x}} dx_3 V_1^M(x, 1-x-x_3, x_3)
\end{aligned}$$

$$\begin{aligned}
& -8e_{q_2}m_N^2 \int_0^{\bar{x}} dx_1 \left[A_1^M + (1-2x)V_1^M - 2(1-x)T_1^M \right] (x_1, x, 1-x_1-x) \\
& -16e_{q_3}m_N^2 x \int_0^{\bar{x}} dx_1 T_1^M(x_1, 1-x_1-x, x) ,
\end{aligned}$$

$$\rho_3^{(1)}(x) = 0 ,$$

$$\begin{aligned}
\rho_4^{(1)}(x) &= -8e_{q_1}(1-2x) \int_0^{\bar{x}} dx_3 V_1(x, 1-x-x_3, x_3) \\
&+ 8e_{q_2} \int_0^{\bar{x}} dx_1 \left[2(1-x)B_1 - (1-2x)C_1 - D_1 \right] (x_1, x, 1-x_1-x) \\
&- 16e_{q_3}x \int_0^{\bar{x}} dx_1 B_1(x_1, 1-x_1-x, x) .
\end{aligned}$$

(2)

VII. FUNCTIONS ρ_i FOR THE FORM FACTOR $\frac{\widetilde{G}_2}{2} + \widetilde{G}_3$

$$\begin{aligned}
\rho_5^{(3)}(x) &= 64e_{q_1}m_N^3(2-x)(1-x)x\check{C}_6(x) \\
&- 16e_{q_2}m_N^2 \left\{ 2(1-x) \left[m_N x^2(2\widetilde{C}_6 - 4\widetilde{B}_8) - x(4m_N\widetilde{C}_6 - 8m_N\widetilde{B}_8) \right] - m_{q_2}x\widetilde{B}_6 \right\} (x) \\
&- 16e_{q_3}m_N^2 \left\{ 2(2-x)(1-x) \left[m_N x\widehat{D}_6 - m_N x(2\widehat{C}_6 - 2\widehat{B}_8) \right] + m_{q_3}x\widehat{B}_6 \right\} (x) ,
\end{aligned}$$

$$\begin{aligned}
\rho_6^{(3)}(x) &= -16e_{q_2}m_N^2 \left\{ 2(1-x) \left[3Q^2\widetilde{B}_6 + m_N x^2(-3m_N\widetilde{B}_6) \right. \right. \\
&\quad \left. \left. - x(-4m_N^2\widetilde{B}_6 + 2Q^2\widetilde{B}_6) \right] - m_{q_2}x \left[-2m_N(1-x)\widetilde{B}_8 \right] \right\} (x) \\
&- 16e_{q_3}m_N^2 \left\{ 2(2-x)(1-x) \left[Q^2\widehat{B}_6 + m_N^2 x\widehat{B}_6 \right] - m_{q_3}x \left[m_N(1-x)(\widehat{D}_6 + 2\widehat{C}_6) \right] \right\} (x) ,
\end{aligned}$$

$$\begin{aligned}
\rho_5^{(2)}(x) &= +8e_{q_1}m_N \left\{ -x \left[(3-2x)\check{C}_2 \right] \right\} (x) \\
&- 8e_{q_2}m_N \left\{ -x\widetilde{D}_2 + x(3-2x)\widetilde{C}_2 - 2(2-x)\widetilde{B}_2 + 2(2-5x+2x^2)\widetilde{B}_4 \right\} (x) \\
&- 8e_{q_3}m_N \left\{ -x(2-x)\widehat{D}_2 + 2x(2-x)\widehat{C}_2 - 2(1-x)\widehat{B}_2 + 2(1-2x+x^2)\widehat{B}_4 \right\} (x) ,
\end{aligned}$$

$$\begin{aligned}
\rho_6^{(2)}(x) &= -8e_{q_2}m_N^2\widetilde{B}_6(x) - 16e_{q_3}m_N^2(1-x)\widehat{C}_6(x) \\
&+ 8e_{q_1}m_N \left\{ m_N(1-x)x\check{D}_5 - 2m_N\check{C}_4 - x \left[-m_N(3-x)\check{C}_4 \right] - 2m_N(1-x)^2\check{C}_5 \right\} (x)
\end{aligned}$$

$$\begin{aligned}
& -8e_{q_2}m_N \left\{ -m_N(1-x)x\tilde{D}_5 + m_N(2-3x+x^2)\tilde{C}_4 \right. \\
& + 2m_N(1-2x+x^2)\tilde{C}_5 + 2m_Nx(1-x)(\tilde{E}_1 + \tilde{H}_1) \\
& \left. - 2m_N(1-x)(4-3x)\tilde{B}_5 + m_{q_2} \left[\tilde{B}_2 - (1-2x)\tilde{B}_4 \right] \right\} (x) \\
& -8e_{q_3}m_N \left\{ m_N(1-x)\hat{D}_5 + m_N(1-x)^2\hat{D}_4 + 2m_N(1-2x+x^2)\hat{C}_4 + 2m_N(1-x)\hat{C}_5 \right. \\
& \left. - 2m_N(2-x)(1-x)\hat{B}_5 - m_{q_3}x(\hat{D}_2 + 2\hat{C}_2) \right\} (x) \\
& + 8e_{q_1}m_N^2(3-2x) \int_0^{\bar{x}} dx_3 V_1^M(x, 1-x-x_3, x_3) \\
& - 8e_{q_2}m_N^2 \int_0^{\bar{x}} dx_1 \left[A_1^M - (3-2x)V_1^M + 6(1-x)T_1^M \right] (x_1, x, 1-x_1-x) \\
& + 16e_{q_3}m_N^2 \int_0^{\bar{x}} dx_1 \left[(1-x)(A_1^M + 2V_1^M) - (2-x)T_1^M \right] (x_1, 1-x_1-x, x) ,
\end{aligned}$$

$$\rho_5^{(1)}(x) = 0 ,$$

$$\begin{aligned}
\rho_6^{(1)}(x) &= 8e_{q_1}(3-2x) \int_0^{\bar{x}} dx_3 V_1^M(x, 1-x-x_3, x_3) \\
& - 8e_{q_2} \int_0^{\bar{x}} dx_1 \left[6(1-x)B_1 - (3-2x)C_1 + D_1 \right] (x_1, x, 1-x_1-x) \\
& - 16e_{q_3} \int_0^{\bar{x}} dx_1 \left[(2-x)B_1 - (1-x)(2C_1 + D_1) \right] (x_1, 1-x_1-x, x) ,
\end{aligned}$$

(3)

where $q_1 = u$, $q_2 = u$, and $q_3 = d$, respectively.

In the above expressions for ρ_2 , ρ_4 , and ρ_6 the functions $\mathcal{F}(x_i)$ are defined in the following way:

$$\begin{aligned}
\check{\mathcal{F}}(x_1) &= \int_1^{x_1} dx'_1 \int_0^{1-x'_1} dx_3 \mathcal{F}(x'_1, 1-x'_1-x_3, x_3) , \\
\check{\check{\mathcal{F}}}(x_1) &= \int_1^{x_1} dx'_1 \int_1^{x'_1} dx''_1 \int_0^{1-x''_1} dx_3 \mathcal{F}(x''_1, 1-x''_1-x_3, x_3) , \\
\tilde{\mathcal{F}}(x_2) &= \int_1^{x_2} dx'_2 \int_0^{1-x'_2} dx_1 \mathcal{F}(x_1, x'_2, 1-x_1-x'_2) , \\
\tilde{\tilde{\mathcal{F}}}(x_2) &= \int_1^{x_2} dx'_2 \int_1^{x'_2} dx''_2 \int_0^{1-x''_2} dx_1 \mathcal{F}(x_1, x''_2, 1-x_1-x''_2) , \\
\hat{\mathcal{F}}(x_3) &= \int_1^{x_3} dx'_3 \int_0^{1-x'_3} dx_1 \mathcal{F}(x_1, 1-x_1-x'_3, x'_3) ,
\end{aligned}$$

$$\widehat{\mathcal{F}}(x_3) = \int_1^{x_3} dx'_3 \int_1^{x'_3} dx''_3 \int_0^{1-x''_3} dx_1 \mathcal{F}(x_1, 1 - x_1 - x''_3, x''_3) .$$

Definitions of the functions B_i , C_i , D_i , E_1 and H_1 that appear in the expressions for $\rho_i(x)$ are given as follows:

$$\begin{aligned} B_2 &= T_1 + T_2 - 2T_3 , \\ B_4 &= T_1 - T_2 - 2T_7 , \\ B_5 &= -T_1 + T_5 + 2T_8 , \\ B_6 &= 2T_1 - 2T_3 - 2T_4 + 2T_5 + 2T_7 + 2T_8 , \\ B_7 &= T_7 - T_8 , \\ B_8 &= -T_1 + T_2 + T_5 - T_6 + 2T_7 + 2T_8 , \\ C_2 &= V_1 - V_2 - V_3 , \\ C_4 &= -2V_1 + V_3 + V_4 + 2V_5 , \\ C_5 &= V_4 - V_3 , \\ C_6 &= -V_1 + V_2 + V_3 + V_4 + V_5 - V_6 , \\ D_2 &= -A_1 + A_2 - A_3 , \\ D_4 &= -2A_1 - A_3 - A_4 + 2A_5 , \\ D_5 &= A_3 - A_4 , \\ D_6 &= A_1 - A_2 + A_3 + A_4 - A_5 + A_6 , \\ E_1 &= S_1 - S_2 , \\ H_1 &= P_2 - P_1 . \end{aligned}$$

The forms of these functions can be found in [11].