

Strong Coupling Constants of the Doubly Heavy Ξ_{QQ} Baryons with π Meson

A. R. Olamaei^{1,4}, K. Azizi^{2,3,4}, S. Rostami⁵

¹ Department of Physics, Jahrom University, Jahrom, P. O. Box 74137-66171, Iran

² Department of Physics, University of Tehran, North Karegar Ave. Tehran 14395-547, Iran

³ Department of Physics, Doğuş University, Acibadem-Kadiköy, 34722 Istanbul, Turkey

⁴ School of Particles and Accelerators, Institute for Research in Fundamental Sciences (IPM),
P. O. Box 19395-5531, Tehran, Iran

⁵ Department of Physics, Shahid Rajaei Teacher Training University, Lavizan, Tehran 16788, Iran

Abstract

The doubly charmed $\Xi_{cc}^{++}(ccu)$ state is the only listed baryon in PDG, which was discovered in the experiment. The LHCb collaboration gets closer to discovering the second doubly charmed baryon $\Xi_{cc}^+(ccd)$, hence the investigation of the doubly charmed/bottom baryons from many aspects is of great importance that may help us not only get valuable knowledge on the nature of the newly discovered states, but also in the search for other members of the doubly heavy baryons predicted by the quark model. In this context, we investigate the strong coupling constants among the $\Xi_{cc}^{+(+)}$ baryons and $\pi^{0(\pm)}$ mesons by means of light cone QCD sum rule. Using the general forms of the interpolating currents of the $\Xi_{cc}^{+(+)}$ baryons and the distribution amplitudes (DAs) of the π meson, we extract the values of the coupling constants $g_{\Xi_{cc}\Xi_{cc}\pi}$. We extend our analyses to calculate the strong coupling constants among the partner baryons with π mesons, as well, and extract the values of the strong couplings $g_{\Xi_{bb}\Xi_{bb}\pi}$ and $g_{\Xi_{bc}\Xi_{bc}\pi}$. The results of this study may help experimental groups in the analyses of the data related to the strong coupling constants among the hadronic multiplets.

1 INTRODUCTION

The search for doubly heavy baryons and determination of their properties constitute one of the main directions of the research in the experimental and theoretical high energy physics. There is only one doubly charmed baryon, Ξ_{cc}^{++} , listed in the PDG. The searches for other members of the doubly heavy baryons in the experiments, as the natural outcomes of the quark model, are in progress. Theoretical investigations on properties of the doubly heavy baryons, are necessary as their results can help us better understand their structure and the dynamics of the QCD as the theory of the strong interaction.

The search for doubly heavy baryons is a long-standing issue. First evidence was reported by the SELEX experiment for Ξ_{cc}^{++} decaying into $\Lambda_c^+ K^- \pi^+$ and $p D^+ K^-$ in final states [1, 2]. The mass measured by SELEX, averaged over the two decay modes, was found to be $3518.7 \pm 1.7 \text{ MeV}/c^2$. However, this has not been confirmed by any other experiments so far. The FOCUS [3], BaBar [4], LHCb [5] and Belle [6] experiments did not find any evidence up to 2017. In 2017, the doubly charmed baryon Ξ_{cc}^{++} was observed by the LHCb collaboration via the decay channel $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi$ [7], and confirmed via measuring another decay channel $\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$ [8]. The weighted average of its mass for the two decay modes was determined to be $3621.24 \pm 0.65(\text{stat.}) \pm 0.31(\text{syst.}) \text{ MeV}/c^2$. Recently, with a data sample corresponding to an integrated luminosity of 9 fb^{-1} at the centre-of-mass energies of 7, 8 and 13 TeV, the LHCb Collaboration published the results of a search for the doubly charmed baryon Ξ_{cc}^+ [9]. The upper limit of the ratio of the production cross-sections between the Ξ_{cc}^+ and Λ_c^+ baryons times the branching fraction of the $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+$ decay, was improved by an order of magnitude than the previous search. However, still no significant signal is observed in the mass range from 3.4 to 3.8 GeV/c^2 . Future LHCb searches with further improved trigger conditions, additional Ξ_{cc}^+ decay modes, and larger data samples should significantly increase the Ξ_{cc}^+ signal sensitivity.

Theoretical studies on the properties and nature of the doubly heavy baryons can play an important role in searching for other members and help us get useful knowledge on the internal structures of the observed resonances. There have been many theoretical efforts aimed at understanding the properties of the doubly-heavy baryon states, see e.g. Refs. [10–38]. However, most researches are focused on the mass and weak decays of the doubly heavy baryons and the number of studies dedicated to their strong decays and the strong couplings of these baryons with other hadrons is very limited. In this context, we investigate the strong coupling constants among the doubly heavy $\Xi_{cc}/\Xi_{bb}/\Xi_{bc}$ baryons and π mesons. We use the well established non-perturbative method of light cone QCD sum rule (LCSR) (for more about this method see, e.g., [39–43] and references therein) as a powerful theoretical tool to calculate the coupling constants under study. In the calculations, we use the general forms of the interpolating currents for Ξ_{cc} , Ξ_{bc} and Ξ_{bb} baryons and DAs of the pions.

The outline of the paper is as follows. In Sec. 2, the light cone sum rules for the coupling constants of the doubly heavy baryons with π mesons are obtained. In Sec. 3, we present the numerical results and discussions, and Sec. 4 is reserved for our conclusions.

2 THEORETICAL FRAMEWORK

The starting point is to choose a suitable correlation function (CF) in terms of hadronic currents sandwiched between the QCD vacuum and the on-shell pseudoscalar meson (here the pion). The QCD vacuum interacts via vacuum fluctuations with the initial and final states and leads to non-perturbative contributions to the final results via the quark-quark, quark-gluon and gluon-gluon condensates. The DAs of π meson also contain non-perturbative information.

To calculate the physical observables like strong coupling constants, we have to calculate the CF in the large timelike momenta by inserting the full set of hadronic states in the CF and isolating the ground state from the continuum and excited states. On the other hand, in QCD or theoretical side, we need to study the CF in the large space-like momenta via the operator product expansion (OPE) that separates the perturbative and non-perturbative contributions in terms of different operators and distribution amplitudes of the particles under consideration. Thus, in QCD side, by contracting the hadronic currents in terms of quark fields we write the CF in terms of the light and heavy quarks propagators as well as wavefunctions of the considered meson in x -space. To proceed, we perform the Fourier transformation to transfer the calculations to the momentum space. To suppress the unwanted contributions coming from the higher states and continuum, we apply the Borel transformation as well as continuum subtraction, supplied by the quark-hadron duality assumption. The two representations of the same CF are then connected to each other via dispersion integrals. These procedures introduce some auxiliary parameters to the calculations, which are then fixed based on the standard prescriptions of the method.

Correlation Function

In order to calculate the strong coupling constants among doubly heavy baryons and light pseudoscalar π meson, we start our discussion by considering the following light-cone correlation function:

$$\Pi(p, q) = i \int d^4x e^{ipx} \langle \mathcal{P}(q) | \mathcal{T} \{ \eta(x) \bar{\eta}(0) \} | 0 \rangle, \quad (1)$$

where $\mathcal{P}(q)$ denotes the pseudoscalar mesons of momentum q , \mathcal{T} is the time ordering operator, and η represents the interpolating currents of the doubly heavy baryons. The general expressions of the interpolating currents for the spin1/2 doubly heavy baryons in their symmetric and antisymmetric forms can be written as

$$\begin{aligned} \eta^S &= \frac{1}{\sqrt{2}} \epsilon_{abc} \left\{ (Q^{aT} C q^b) \gamma_5 Q^c + (Q'^{aT} C q^b) \gamma_5 Q^c + t(Q^{aT} C \gamma_5 q^b) Q^c + t(Q'^{aT} C \gamma_5 q^b) Q^c \right\}, \\ \eta^A &= \frac{1}{\sqrt{6}} \epsilon_{abc} \left\{ 2(Q^{aT} C Q^b) \gamma_5 q^c + (Q^{aT} C q^b) \gamma_5 Q^c - (Q'^{aT} C q^b) \gamma_5 Q^c + 2t(Q^{aT} C \gamma_5 Q^b) q^c \right. \\ &\quad \left. + t(Q^{aT} C \gamma_5 q^b) Q^c - t(Q'^{aT} C \gamma_5 q^b) Q^c \right\}, \end{aligned} \quad (2)$$

where t is an arbitrary auxiliary parameter and the case, $t = -1$ corresponds to the Ioffe current. Here $Q^{(\prime)}$ and q stand for the heavy and light quarks respectively; $a, b,$

and c are the color indices, C stands for the charge conjugation operator and T denotes the transposition. For the doubly heavy baryons with two identical heavy quarks, the antisymmetric form of the interpolating current is zero and we just need to employ the symmetric form, η^S . In the following, we calculate the correlation function in Eq. 1 in two different windows.

Physical Side

To obtain the physical side of correlation function, we insert complete sets of hadronic states with the same quantum numbers as the interpolating currents and isolate the ground state. After performing the integration over four- x , we get

$$\Pi^{\text{Phys.}}(p, q) = \frac{\langle 0|\eta|B_2(p)\rangle\langle B_2(p)\mathcal{P}(q)|B_1(p+q)\rangle\langle B_1(p+q)|\bar{\eta}|0\rangle}{(p^2 - m_1^2)[(p+q)^2 - m_2^2]} + \dots, \quad (3)$$

where dots in the above equation stand for the contribution of the higher states and continuum. To proceed we introduce the matrix elements for spin-1/2 baryons as

$$\begin{aligned} \langle 0|\eta|B_2(p, r)\rangle &= \lambda_{B_2}u(p, r), \\ \langle B_1(p+q, s)|\bar{\eta}|0\rangle &= \lambda_{B_1}\bar{u}(p+q, s), \\ \langle B_2(p, r)\mathcal{P}(q)|B_1(p+q, s)\rangle &= g_{B_1B_2\mathcal{P}}\bar{u}(p, r)\gamma_5u(p+q, s), \end{aligned} \quad (4)$$

where $g_{B_1B_2\mathcal{P}}$, representing the strong decay $B_1 \rightarrow B_2\mathcal{P}$, is the strong coupling constant among the baryons B_1 and B_2 as well as the \mathcal{P} meson, λ_{B_1} and λ_{B_2} are the residues of the corresponding baryons and $u(q, s)$ is Dirac spinor with spin s . Putting the above equations all together, and performing summation over spins, we get the following representation of the correlator for the phenomenological side:

$$\Pi^{\text{Phys.}}(p, q) = \frac{g_{B_1B_2\mathcal{P}}\lambda_{B_1}\lambda_{B_2}}{(p^2 - m_{B_2}^2)[(p+q)^2 - m_{B_1}^2]}[\not{p}\not{q}\gamma_5 + \dots], \quad (5)$$

where dots represent other structures come from spin summation as well as the contributions of higher states and continuum. We will use the explicitly presented structure to extract the value of the strong coupling constant, $g_{B_1B_2\mathcal{P}}$. We apply the double Borel transformation with respect to the variables $p_1^2 = (p+q)^2$ and $p_2^2 = p^2$:

$$\begin{aligned} \mathcal{B}_{p_1}(M_1^2)\mathcal{B}_{p_2}(M_2^2)\Pi^{\text{Phys.}}(p, q) &\equiv \Pi^{\text{Phys.}}(M^2) \\ &= g_{B_1B_2\mathcal{P}}\lambda_{B_1}\lambda_{B_2}e^{-m_{B_1}^2/M_1^2}e^{-m_{B_2}^2/M_2^2}\not{p}\not{q}\gamma_5 + \dots, \end{aligned} \quad (6)$$

where $M^2 = M_1^2M_2^2/(M_1^2 + M_2^2)$ and the Borel parameters M_1^2 and M_2^2 for the problem under consideration are chosen to be equal as the masses of the initial and final state baryons are the same. Hence $M_1^2 = M_2^2 = 2M^2$.

QCD Side

In QCD side, the correlation function is calculated in deep Euclidean region with the help of OPE. To proceed, we need to determine the correlation function using the quark propagators and distribution amplitudes of the π meson. The $\Pi^{QCD}(p, q)$ can be written in the following general form:

$$\Pi^{QCD}(p, q) = \Pi((p+q)^2, p^2)\not{p}\not{q}\gamma_5 + \dots, \quad (7)$$

where the $\Pi((p+q)^2, p^2)$ is an invariant function that should be calculated in terms of QCD degrees of freedom as well as the parameters inside the DAs.

Inserting, for instance, η^S in the CF and using the Wick theorem to contract all the heavy quark fields we get the following expression in terms of the heavy quark propagators and π meson matrix elements:

$$\begin{aligned}
\Pi_{\rho\sigma}^{\text{QCD}}(p, q) &= \frac{i}{2} \epsilon_{abc} \epsilon_{a'b'c'} \int d^4x e^{iq \cdot x} \langle \pi(q) | \bar{q}_\alpha^c(0) q_\beta^c(x) | 0 \rangle \left\{ \left[\left(\tilde{S}_Q^{aa'}(x) \right)_{\alpha\beta} \left(\gamma_5 S_{Q'}^{bb'}(x) \gamma_5 \right)_{\rho\sigma} \right. \right. \\
&+ \left. \left(\gamma_5 S_{Q'}^{bb'}(x) C \right)_{\rho\alpha} \left(C S_Q^{aa'}(x) \gamma_5 \right)_{\beta\sigma} + t \left\{ \left(\gamma_5 \tilde{S}_Q^{aa'}(x) \right)_{\alpha\beta} \left(\gamma_5 S_{Q'}^{bb'}(x) \right)_{\rho\sigma} \right. \right. \\
&+ \left. \left. \left(\tilde{S}_Q^{aa'}(x) \gamma_5 \right)_{\alpha\beta} \left(S_{Q'}^{bb'}(x) \gamma_5 \right)_{\rho\sigma} + \left(\gamma_5 S_{Q'}^{bb'}(x) C \gamma_5 \right)_{\rho\alpha} \left(C S_Q^{aa'}(x) \right)_{\beta\sigma} \right. \right. \\
&- \left. \left. \left(S_{Q'}^{bb'}(x) C \right)_{\rho\alpha} \left(\gamma_5 C S_Q^{aa'}(x) \gamma_5 \right)_{\beta\sigma} \right\} + t^2 \left\{ \left(\gamma_5 \tilde{S}_Q^{aa'}(x) \gamma_5 \right)_{\alpha\beta} \left(S_{Q'}^{bb'}(x) \right)_{\rho\sigma} \right. \right. \\
&- \left. \left. \left(S_{Q'}^{bb'}(x) C \gamma_5 \right)_{\rho\alpha} \left(\gamma_5 C S_Q^{aa'}(x) \right)_{\beta\sigma} \right\} \right] + \left(Q \longleftrightarrow Q' \right) \left. \right\}, \quad (8)
\end{aligned}$$

where $\tilde{S} = C S^T C$, and $\langle \pi(q) | \bar{q}_\alpha^b(0) q_\beta^b(x) | 0 \rangle$ are the matrix elements for the light quark contents of the doubly heavy baryons. To proceed, we need to know the explicit expression for the heavy quark propagator that is

$$\begin{aligned}
S_Q(x) &= \frac{m_Q^2}{4\pi^2} \frac{K_1(m_Q \sqrt{-x^2})}{\sqrt{-x^2}} - i \frac{m_Q^2 \not{x}}{4\pi^2 x^2} K_2(m_Q \sqrt{-x^2}) \\
&- i g_s \int \frac{d^4k}{(2\pi)^4} e^{-ikx} \int_0^1 du \left[\frac{\not{k} + m_Q}{2(m_Q^2 - k^2)^2} G^{\mu\nu}(ux) \sigma_{\mu\nu} + \frac{u}{m_Q^2 - k^2} x_\mu G^{\mu\nu} \gamma_\nu \right] + \dots, \quad (9)
\end{aligned}$$

where K_1 and K_2 are the modified Bessel functions of the second kind, and $G_{ab}^{\mu\nu} \equiv G_A^{\mu\nu} t_{ab}^A$ with $A = 1, 2 \dots 8$, and $t^A = \lambda^A/2$, where λ^A are the Gell-Mann matrices. The first two terms correspond to perturbative or free part and the rest belong to the interacting parts.

The next step is to use the heavy quark propagator and the matrix elements $\langle \pi(q) | \bar{q}_\alpha^b(0) q_\beta^b(x) | 0 \rangle$ in Eq. 8. This leads to different kinds of contributions to the CF. Figs. 1 and 2 are the Feynman diagrams correspond to the leading and next-to-leading order contributions, respectively which are considered in this work. Only matrix elements corresponding to these diagrams are available. To calculate the leading order contribution, the heavy quark propagators are replaced by just their perturbative parts. This contribution can be computed using π meson two-particle DAs of twist two and higher.

The next-to-leading order contributions can also be calculated by choosing the gluonic parts in Eq. 9 for one of the heavy quark propagators and leaving the other with its perturbative term. They can be expressed in terms of pion three particles DAs. The terms involving more than one gluon field that proportional to four-particle DAs or more are neglected as they are not available

Now, we concentrate on the strong decay $\Xi_{cc}^{++} \rightarrow \Xi_{cc}^{++} \pi^0$ with the aim of calculating the corresponding strong coupling constant $g_{\Xi_{cc}^{++} \Xi_{cc}^{++} \pi^0}$. The other channels have similar procedures. To proceed, we replace the heavy quark propagators in 8 by their




Figure 1: The leading order diagram contributing to QCD side.




Figure 2: The one-gluon exchange diagrams corresponding to the next-to-leading contributions.

explicit expression and perform the summation over the color indices by applying the replacement

$$\bar{u}_\alpha^a(x)u_\beta^{a'}(0) \rightarrow \frac{1}{3}\delta_{aa'}\bar{u}_\alpha(x)u_\beta(0). \quad (10)$$

Now, using the expression

$$\bar{u}_\alpha(u)u_\beta(0) \equiv \frac{1}{4}\Gamma_{\beta\alpha}^J\bar{u}(x)\Gamma^J u(0), \quad (11)$$

one can relate the CF to the DAs of the pion with different twists. Here the summation over J runs as

$$\Gamma^J = \mathbf{1}, \gamma_5, \gamma_\mu, i\gamma_5\gamma_\mu, \sigma_{\mu\nu}/\sqrt{2}. \quad (12)$$

Following the similar way one can calculate the contributions involving the gluon field.

As a result, the CF is found in terms of the QCD parameters as well as the matrix elements

$$\begin{aligned} &\langle\pi^0(q)|\bar{u}(x)\Gamma^J u(0)|0\rangle, \\ &\langle\pi^0(q)|\bar{u}(x)\Gamma^J G_{\mu\nu}(vx)u(0)|0\rangle, \end{aligned} \quad (13)$$

whose expressions in terms of the wave functions of the pion with different twists are given in the Appendix.

Inserting the expression of the above-mentioned matrix elements in term of wave functions of different twists we get the CF in x space. This is followed by the Fourier

and Borel transformations as well as continuum subtraction. To proceed we need to perform the Fourier transformation of the following kind:

$$T_{[\quad, \alpha, \alpha\beta]}(p, q) = i \int d^4x \int_0^1 dv \int \mathcal{D}\alpha e^{ip \cdot x} (x^2)^n [e^{i(\alpha_{\bar{q}} + v\alpha_g)q \cdot x} \mathcal{G}(\alpha_i), e^{iq \cdot x} f(u)] \\ \times [1, x_\alpha, x_\alpha x_\beta] K_\mu(m_Q \sqrt{-x^2}) K_\nu(m_Q \sqrt{-x^2}), \quad (14)$$

where the expressions in the brackets denote different possibilities arise in the calculations, the blank subscript in the left hand side indicates no indices regarding no x_α in the configuration, $\mathcal{G}(\alpha_i)$ and $f(u)$ represent wave functions coming from the three and two-particle matrix elements and n is a positive integer. The measure

$$\int \mathcal{D}\alpha = \int_0^1 d\alpha_q \int_0^1 d\alpha_{\bar{q}} \int_0^1 d\alpha_g \delta(1 - \alpha_q - \alpha_{\bar{q}} - \alpha_g),$$

is used in the calculations. To start the Fourier transformation, we use

$$(x^2)^n = (-1)^n \frac{d^n}{d\beta^n} (e^{-\beta x^2})|_{\beta=0}. \quad (15)$$

for positive integer n and

$$x_\alpha e^{iP \cdot x} = (-i) \frac{d}{dP_\alpha} e^{iP \cdot x}. \quad (16)$$

We also use the following representation of the Bessel functions K_ν (see also Ref. [44]):

$$K_\nu(m_Q \sqrt{-x^2}) = \frac{\Gamma(\nu + 1/2) 2^\nu}{\sqrt{\pi} m_Q^\nu} \int_0^\infty dt \cos(m_Q t) \frac{(\sqrt{-x^2})^\nu}{(t^2 - x^2)^{\nu+1/2}}. \quad (17)$$

As an example let us consider the following generic form:

$$\mathcal{Z}_{\alpha\beta}(p, q) = i \int d^4x \int_0^1 dv \int \mathcal{D}\alpha e^{i[p + (\alpha_{\bar{q}} + v\alpha_g)q] \cdot x} \mathcal{G}(\alpha_i) (x^2)^n \\ \times x_\alpha x_\beta K_\mu(m_Q \sqrt{-x^2}) K_\nu(m_Q \sqrt{-x^2}). \quad (18)$$

We substitute Eqs. 15, 16 and 17 into 18. Then to perform the x -integration we go to the Euclidean space by Wick rotation and get

$$\mathcal{Z}_{\alpha\beta}(p, q) = \frac{i\pi^2 2^{\mu+\nu-2}}{m_{Q_1}^{2\mu} m_{Q_2}^{2\nu}} \int \mathcal{D}\alpha \int_0^1 dv \int_0^1 dy_1 \int_0^1 dy_2 \frac{\partial}{\partial P_\alpha} \frac{\partial}{\partial P_\beta} \frac{\partial^n}{\partial \beta^n} \\ \times \frac{y_1^{\mu-1} y_2^{\nu-1}}{(y_1 + y_2 + \beta)^2} e^{-\frac{1}{4} \left(\frac{m_{Q_1}^2}{y_1} + \frac{m_{Q_2}^2}{y_2} - \frac{P^2}{y_1 + y_2 + \beta} \right)}, \quad (19)$$

where $P = p + q(v\alpha_g + \alpha_q)$. Changing variables from y_1 and y_2 to ρ and z as

$$\rho = y_1 + y_2, \quad z = \frac{y_1}{y_1 + y_2}, \quad (20)$$

and taking derivative with respect to P_α and P_β , we get

$$\begin{aligned}
\mathcal{Z}_{\alpha\beta}(p, q) &= \frac{i\pi^2 2^{\mu+\nu-4}}{m_{Q_1}^{2\mu} m_{Q_2}^{2\nu}} \int \mathcal{D}\alpha \int_0^1 dv \int_0^\infty d\rho \int_0^1 dz \frac{\partial^n}{\partial \beta^n} z^{\mu-1} \bar{z}^{\nu-1} \frac{\rho^{\nu+\mu-1}}{(\rho+\beta)^4} e^{-\frac{1}{4}\left(\frac{m_{Q_1}^2 \bar{z} + m_{Q_2}^2 z}{z\bar{z}\rho} + \frac{P^2}{\beta+\rho}\right)} \\
&\times \left[p_\alpha p_\beta + (v\alpha_g + \alpha_q)(p_\alpha q_\beta + q_\alpha p_\beta) + (v\alpha_g + \alpha_q)^2 q_\alpha q_\beta + 2(\rho + \beta)g_{\alpha\beta} \right]. \quad (21)
\end{aligned}$$

Now we perform the double Borel transformation using

$$\mathcal{B}_{p_1}(M_1^2) \mathcal{B}_{p_2}(M_2^2) e^{b(p+uq)^2} = M^2 \delta\left(b + \frac{1}{M^2}\right) \delta(u_0 - u) e^{\frac{-q^2}{M_1^2 + M_2^2}}, \quad (22)$$

where $u_0 = M_1^2/(M_1^2 + M_2^2)$. After integrating over ρ we have

$$\begin{aligned}
\mathcal{Z}_{\alpha\beta}(M^2) &= \frac{i\pi^2 2^{4-\mu-\nu} e^{\frac{-q^2}{M_1^2 + M_2^2}}}{M^2 m_{Q_1}^{2\mu} m_{Q_2}^{2\nu}} \int \mathcal{D}\alpha \int_0^1 dv \int_0^1 dz \frac{\partial^n}{\partial \beta^n} e^{-\frac{m_{Q_1}^2 \bar{z} + m_{Q_2}^2 z}{z\bar{z}(M^2 - 4\beta)}} z^{\mu-1} \bar{z}^{\nu-1} (M^2 - 4\beta)^{\mu+\nu-1} \\
&\times \delta[u_0 - (\alpha_q + v\alpha_g)] \left[p_\alpha p_\beta + (v\alpha_g + \alpha_q)(p_\alpha q_\beta + q_\alpha p_\beta) + (v\alpha_g + \alpha_q)^2 q_\alpha q_\beta \right. \\
&\quad \left. + \frac{M^2}{2} g_{\alpha\beta} \right]. \quad (23)
\end{aligned}$$

The next step is to perform the continuum subtraction in order to more suppress the contribution of the higher states and continuum. The subtraction procedures for different systems are described in Ref. [44] in details. When the masses of the initial and final baryonic states are equal, as we stated previously, we set $M_1^2 = M_2^2 = 2M^2$. In this case, the double spectral density is concentrated near the diagonal $s_1 = s_2$ and reduces to a single representation s (see also Ref. [45] and references therein) and for the continuum subtraction more simple expressions are derived, which are not sensitive to the shape of the duality region. For the case, $M_1^2 = M_2^2 = 2M^2$ and $u_0 = 1/2$, the subtraction procedure is explained in Ref. [45] in details, which we use in the calculations.

By calculation of all the Fourier integrals and applying the Borel transformation and continuum subtraction, for QCD side of the calculations in Borel Scheme, we get,

$$\Pi^{QCD}(M^2, s_0, t) = [\Pi^{(0)}(M^2, s_0, t) + \Pi^{(GG)}(M^2, s_0, t)] \not{p} \gamma_5 + \dots, \quad (24)$$

where the functions $\Pi^{(0)}(M^2, s_0, t)$ and $\Pi^{(GG)}(M^2, s_0, t)$ are obtained as

$$\begin{aligned}
\Pi^{(0)}(M^2, s_0, t) &= \frac{e^{-\frac{m_\pi^2}{4M^2}}}{96\pi^2\sqrt{2}} \int_0^1 dz \frac{1}{z} e^{-\frac{m_c^2}{M^2 z \bar{z}}} \left\{ 3f_\pi m_\pi^2 m_c^3 (t^2 - 1) \mathbb{A}(u_0) \right. \\
&+ 2e^{-\frac{4m_c^2}{M^2}} \int_{4m_c^2}^{s_0} ds e^{-\frac{s}{M^2}} z \left[3f_\pi m_\pi^2 m_c (t^2 - 1) \bar{z} \mathbb{A}(u_0) \right. \\
&- 6f_\pi m_c (s - 4m_c^2) (t^2 - 1) \bar{z} \varphi_\pi(u_0) \\
&- \left. \left. \mu_\pi (\tilde{\mu}_\pi^2 - 1) \left[-4m_c^2 t + 3(s - 4m_c^2)(t - 1)^2 z \bar{z} \right] \varphi_\sigma(u_0) \right] \right\} \\
&+ 6e^{-\frac{4m_c^2}{M^2}} (t^2 - 1) \int_{4m_c^2}^{s_0} ds e^{-\frac{s}{M^2}} \int_0^1 dv \int \mathcal{D}\alpha \\
&\times \left[f_\pi m_\pi^2 m_c \delta[u_0 - (\alpha_q + v\alpha_g)] \left(-2z(v - 1/2) \mathcal{A}_\parallel(\alpha_i) + 2\bar{z} \mathcal{V}_\perp(\alpha_i) \right. \right. \\
&+ \left. \left. (2z - 3) \mathcal{V}_\parallel(\alpha_i) \right) + 2\mu_\pi \delta'[u_0 - (\alpha_q + v\alpha_g)] (s - 4m_c^2) z \bar{z} (v - 1/2) \mathcal{T}(\alpha_i) \right] \left. \right\}, \tag{25}
\end{aligned}$$

and

$$\begin{aligned}
\Pi^{(GG)}(M^2, s_0, t) &= \frac{\langle g_s^2 G^2 \rangle e^{-\frac{m_\pi^2}{4M^2}}}{6912\sqrt{2}\pi^2 m_c M^6} \int_0^1 dz \frac{1}{z^2 \bar{z}^4} e^{-\frac{m_c^2}{M^2 z \bar{z}}} \\
&\times \left\{ \bar{z}^2 \left[-3f_\pi m_\pi^2 (1 - t^2) \left(6M^6 z^2 \bar{z}^4 + 6M^4 m_c^2 z \bar{z}^3 + 3M^2 m_c^4 \bar{z}^2 - 2m_c^6 \right) \mathbb{A}(u_0) \right. \right. \\
&+ 4M^2 m_c \bar{z} \left(-3f_\pi M^2 m_c (1 - t^2) z \left[2m_c^2 + M^2 \bar{z} (5z - 3) \right] \varphi_\pi(u_0) \right. \\
&+ \left. \left. (\tilde{\mu}_\pi^2 - 1) \mu_\pi \left[2m_c^4 (t^2 + 1) + [(1 + t^2)(1 - 4z) - 6t](M^2 m_c^2 z + M^4 z^2 \bar{z}) \right] \varphi_\sigma(u_0) \right) \right. \\
&+ \left. \left. 72f_\pi M^8 (1 - t^2) z^2 \bar{z}^4 \left(1 - e^{-\frac{(s_0 - 4m_c^2)}{M^2}} \right) \varphi_\pi(u_0) \right] \right\} \\
&+ 6M^2 (1 - t) \int_0^1 dv \int \mathcal{D}\alpha \left[f_\pi m_\pi^2 (1 + t) \bar{z}^3 \delta[u_0 - (\alpha_q + v\alpha_g)] \left(2m_c^4 \mathcal{V}_\perp(\alpha_i) \right. \right. \\
&+ \left. \left[-2m_c^4 + M^2 m_c^2 (1 + 2z) z + M^4 (1 + 2z) z^2 \bar{z} \right] \mathcal{V}_\parallel(\alpha_i) \right. \\
&- \left. \left. (2v - 1) \left[m_c^4 + M^2 m_c^2 (1 + 2z) z + M^4 (1 + 2z) z^2 \bar{z} \right] \mathcal{A}_\parallel(\alpha_i) \right) \right. \\
&+ \left. \left. \mu_\pi (1 - t) (1 + 2v) z \bar{z}^3 \delta'[u_0 - (\alpha_q + v\alpha_g)] (M^2 m_c^3 + M^4 m_c z \bar{z}) \mathcal{T}(\alpha_i) \right] \right\}. \tag{26}
\end{aligned}$$

The sum rule for the coupling constant under study is found by matching the coefficients of the structure $\not{q}\not{p}\gamma_5$ from both the physical and QCD sides. As a result, we get:

$$g_{B_1 B_2 \mathcal{P}}(M^2, s_0, t) = \frac{1}{\lambda_{\Xi_{cc}} \lambda_{\Xi_{cc}}} e^{\frac{m_{B_1}^2}{2M^2}} e^{\frac{m_{B_2}^2}{2M^2}} \left[\Pi^{(0)}(M^2, s_0, t) + \Pi^{(GG)}(M^2, s_0, t) \right]. \tag{27}$$

Parameters	Values
m_c	1.27 ± 0.02 GeV
m_b	$4.18^{+0.03}_{-0.02}$ GeV
m_{π^0}	134.98 MeV
m_{π^\pm}	139.57 MeV
Ξ_{cc}	3621.2 ± 0.7 MeV
Ξ_{bc}	6.72 ± 0.20 GeV [10]
Ξ_{bb}	9.96 ± 0.90 GeV [10]
f_π	130.2 ± 1.2 MeV

Table 1: Some input values used in the calculations. They are mainly taken from [46], except the ones that the references are given next to the numbers.

As is seen, the sum rules for coupling constants contain the residues of doubly heavy baryons, which are borrowed from Ref. [10]. Similarly, we obtain the sum rules for other strong coupling constants under consideration.

3 NUMERICAL ANALYSIS

In this section, we numerically analyze the sum rules for the strong coupling constants of the π mesons with Ξ_{cc} , Ξ_{bc} and Ξ_{bb} baryons and discuss the results. The sum rules for the couplings $g_{\Xi_{cc}\Xi_{cc}\pi}$, $g_{\Xi_{bc}\Xi_{bc}\pi}$ and $g_{\Xi_{bb}\Xi_{bb}\pi}$ contain some input parameters like, quark masses the mass and decay constant of the π meson and the masses and residues of doubly heavy baryons. They were extracted from experimental data or calculated from nonperturbative methods. The values of some of these parameters together with quark masses are given in Tables 1. As we previously mentioned the values of the residues of baryons are used from Ref. [10]. Another set of important input parameters are the π meson wavefunctions of different twists, entering the DAs. These wavefunctions are

given as [47, 48]:

$$\begin{aligned}
\phi_\pi(u) &= 6u\bar{u}\left(1 + a_1^\pi C_1(2u-1) + a_2^\pi C_2^{\frac{3}{2}}(2u-1)\right), \\
\mathcal{T}(\alpha_i) &= 360\eta_3\alpha_q\alpha_{\bar{q}}\alpha_g^2\left(1 + w_3\frac{1}{2}(7\alpha_g - 3)\right), \\
\phi_P(u) &= 1 + \left(30\eta_3 - \frac{5}{2}\frac{1}{\mu_\pi^2}\right)C_2^{\frac{1}{2}}(2u-1) \\
&\quad + \left(-3\eta_3w_3 - \frac{27}{20}\frac{1}{\mu_\pi^2} - \frac{81}{10}\frac{1}{\mu_\pi^2}a_2^\pi\right)C_4^{\frac{1}{2}}(2u-1), \\
\phi_\sigma(u) &= 6u\bar{u}\left[1 + \left(5\eta_3 - \frac{1}{2}\eta_3w_3 - \frac{7}{20}\mu_\pi^2 - \frac{3}{5}\mu_\pi^2a_2^\pi\right)C_2^{\frac{3}{2}}(2u-1)\right], \\
\mathcal{V}_\parallel(\alpha_i) &= 120\alpha_q\alpha_{\bar{q}}\alpha_g\left(v_{00} + v_{10}(3\alpha_g - 1)\right), \\
\mathcal{A}_\parallel(\alpha_i) &= 120\alpha_q\alpha_{\bar{q}}\alpha_g\left(0 + a_{10}(\alpha_q - \alpha_{\bar{q}})\right), \\
\mathcal{V}_\perp(\alpha_i) &= -30\alpha_g^2\left[h_{00}(1 - \alpha_g) + h_{01}(\alpha_g(1 - \alpha_g) - 6\alpha_q\alpha_{\bar{q}}) + h_{10}(\alpha_g(1 - \alpha_g) - \frac{3}{2}(\alpha_{\bar{q}}^2 + \alpha_q^2))\right], \\
\mathcal{A}_\perp(\alpha_i) &= 30\alpha_g^2(\alpha_{\bar{q}} - \alpha_q)\left[h_{00} + h_{01}\alpha_g + \frac{1}{2}h_{10}(5\alpha_g - 3)\right], \\
B(u) &= g_\pi(u) - \phi_\pi(u), \\
g_\pi(u) &= g_0C_0^{\frac{1}{2}}(2u-1) + g_2C_2^{\frac{1}{2}}(2u-1) + g_4C_4^{\frac{1}{2}}(2u-1), \\
\mathbb{A}(u) &= 6u\bar{u}\left[\frac{16}{15} + \frac{24}{35}a_2^\pi + 20\eta_3 + \frac{20}{9}\eta_4 + \left(-\frac{1}{15} + \frac{1}{16} - \frac{7}{27}\eta_3w_3 - \frac{10}{27}\eta_4\right)C_2^{\frac{3}{2}}(2u-1)\right. \\
&\quad + \left(-\frac{11}{210}a_2^\pi - \frac{4}{135}\eta_3w_3\right)C_4^{\frac{3}{2}}(2u-1)\left. \right] \\
&\quad + \left(-\frac{18}{5}a_2^\pi + 21\eta_4w_4\right)\left[2u^3(10 - 15u + 6u^2)\ln u\right. \\
&\quad \left. + 2\bar{u}^3(10 - 15\bar{u} + 6\bar{u}^2)\ln \bar{u} + u\bar{u}(2 + 13u\bar{u})\right], \tag{28}
\end{aligned}$$

where $C_n^k(x)$ are the Gegenbauer polynomials and

$$\begin{aligned}
h_{00} &= v_{00} = -\frac{1}{3}\eta_4, \\
a_{10} &= \frac{21}{8}\eta_4w_4 - \frac{9}{20}a_2^\pi, \\
v_{10} &= \frac{21}{8}\eta_4w_4, \\
h_{01} &= \frac{7}{4}\eta_4w_4 - \frac{3}{20}a_2^\pi, \\
h_{10} &= \frac{7}{4}\eta_4w_4 + \frac{3}{20}a_2^\pi, \\
g_0 &= 1, \\
g_2 &= 1 + \frac{18}{7}a_2^\pi + 60\eta_3 + \frac{20}{3}\eta_4, \\
g_4 &= -\frac{9}{28}a_2^\pi - 6\eta_3w_3. \tag{29}
\end{aligned}$$

The constants inside the wavefunctions are calculated at the renormalization scale of $\mu = 1 \text{ GeV}^2$ and they are given as $a_1^\pi = 0$, $a_2^\pi = 0.44$, $\eta_3 = 0.015$, $\eta_4 = 10$, $w_3 = -3$ and $w_4 = 0.2$ [47, 48].

Finally, the sum rules for the coupling constants contain three auxiliary parameters: Borel mass parameter M^2 , continuum threshold s_0 and the general parameter t entered the general spin1/2 currents. We should find working regions of these parameters, at which the results of coupling constants have relatively small variations with respect to the changes of these parameters. To restrict these parameters, we employ the standard prescriptions of the method such as the pole dominance, convergence of the OPE and mild variations of the physical quantities with respect to the auxiliary parameters. The upper limit of M^2 is determined from the pole dominance condition, i.e.,

$$\frac{\Pi^{QCD}(M^2, s_0, t)}{\Pi^{QCD}(M^2, \infty, t)} > \frac{1}{2}. \quad (30)$$

The lower limit of M^2 is fixed by the condition of OPE convergence: in our case, $\Pi^{(0)}(M^2, s_0, t) > \Pi^{(GG)}(M^2, s_0, t)$. The continuum threshold s_0 is not totally arbitrary and it depends on the mass of the first excited state in the same channel. One has to choose the range of s_0 such that it does not contain the energy for producing the first excited state. Unfortunately, there is no experimental information on the masses of the first excited states in the case of doubly heavy baryons. Based on our analyses and considering the experimental information on the single heavy baryons, we consider the interval $m_{QQ} + E_1 \leq \sqrt{s_0} \leq m_{QQ} + E_2$ for $\sqrt{s_0}$, where a energy from E_1 to E_2 is needed to excite the baryons, and impose that the Borel curves are flat and the requirements of the pole dominance and the OPE convergence are satisfied. With these criteria, we choose the s_0 to lie in the interval $(m_{\Xi_{QQ}} + 0.3)^2 \leq s_0 \leq (m_{\Xi_{QQ}} + 0.7)^2 (\text{GeV}^2)$.

As a result of the above requirements, we obtain the working region of the Borel parameter for the Ξ_{cc} channel as,

$$3 \text{ GeV}^2 \leq M^2 \leq 6 \text{ GeV}^2. \quad (31)$$

The continuum threshold for this channel is obtained as,

$$16 \text{ GeV}^2 \leq s_0 \leq 18 \text{ GeV}^2. \quad (32)$$

For Ξ_{bc} baryon, we get

$$8 \text{ GeV}^2 \leq M^2 \leq 12 \text{ GeV}^2, \quad 49 \text{ GeV}^2 \leq s_0 \leq 55 \text{ GeV}^2. \quad (33)$$

Finally, for Ξ_{bb} baryon, these parameters lie in the intervals:

$$18 \text{ GeV}^2 \leq M^2 \leq 24 \text{ GeV}^2, \quad 106 \text{ GeV}^2 \leq s_0 \leq 114 \text{ GeV}^2. \quad (34)$$

The working window for the parameter t is obtained by the consideration of the minimum variations of the results with respect to this parameter. By imposing this condition together with the conditions of the pole dominance and convergence of the mass sum rules the working window for t is obtained in Ref. [10] as $|t| \leq 2$, which we also use in our analyses. As examples, we display the dependence of the strong coupling constant $g_{\Xi_{cc}^{++}\Xi_{cc}^{++}\pi^0}$, which is obtained from the sum rule for the strong coupling form factor at $q^2 = m_\pi^2$, with respect to M^2 and s_0 in Figs. 3 and 4 at $t = -2$.

From these figures we see mild variations of $g_{\Xi_{cc}\Xi_{cc}\pi^0}$ with respect to the M^2 and s_0 , which appear as the main uncertainty in the numerical values of the strong coupling constants. We extract the numerical values of the strong couplings $g_{\Xi_{cc}\Xi_{cc}\pi^0}$, $g_{\Xi_{bc}\Xi_{bc}\pi^0}$ and $g_{\Xi_{bb}\Xi_{bb}\pi^0}$ as displayed in table 2. The presented errors are due to the changes




Figure 3: The strong coupling $g_{\Xi_{cc}^{++}\Xi_{cc}^{++}\pi^0}$ as a function of the Borel parameter M^2 at $t = -2$ for different values of s_0 .




Figure 4: The strong coupling $g_{\Xi_{cc}^{++}\Xi_{cc}^{++}\pi^0}$ as a function of s_0 at $t = -2$ for different values of M^2 .

with respect to the auxiliary parameters in their working regions as well as those which propagate from other input parameters as well as π meson DAs. The values of coupling constants to the charged mesons π^+ and π^- , which are exactly the same from the isospin symmetry, are found by the multiplications of the strong coupling constants in π^0 channel by $\sqrt{2}$. This coefficient is the only difference in the couplings to the quark contents of the π^\pm and π^0 mesons when the isospin symmetry is used. As it is also clear from table 2, the values of the strong coupling constants in double- b channel are roughly four times greater than those of the other channels. The big difference between the strong couplings to pseudoscalar mesons in b and c channels is evident in the case of single heavy baryons, as well [49]. As it can be seen from this reference, the difference factor in the case of single heavy baryons is two-three times.

Channel	strong coupling constant
$\Xi_{cc} \rightarrow \Xi_{cc}\pi^0$	$5.52^{+0.64}_{-0.53}$
$\Xi_{bc} \rightarrow \Xi_{bc}\pi^0$	$4.75^{+0.42}_{-0.50}$
$\Xi_{bb} \rightarrow \Xi_{bb}\pi^0$	$21.60^{+1.77}_{-2.09}$

Table 2: The numerical values for the strong coupling constants extracted from the analyses.

4 SUMMARY AND CONCLUSIONS

The doubly charmed $\Xi_{cc}^{++}(ccu)$ baryon is the only listed doubly heavy baryon in PDG discovered in the experiment so far. The LHCb collaboration gets closer to observing other member $\Xi_{cc}^+(ccd)$, as well. Therefore, the investigation of the doubly charmed/bottom baryons from many aspects is of great importance that may help us in the course of search for new members of the doubly heavy baryons predicted by the quark model. The strong coupling constants among the hadronic multiplets are fundamental objects that can help us to explore the nature and structure of the participating particles as well as the properties of QCD as the theory of strong interaction.

We calculated the strong coupling constants $g_{\Xi_{ccq}\Xi_{ccq}\pi^{0,\pm}}$, $g_{\Xi_{bcq}\Xi_{bcq}\pi^{0,\pm}}$ and $g_{\Xi_{bbq}\Xi_{bbq}\pi^{0,\pm}}$, with q being either u or d quark, in the framework of the light cone QCD sum rule and using the general form of the interpolating currents for the doubly heavy baryons and the π meson's DAs. Based on the standard prescriptions of the method, we fixed the auxiliary parameters entering the calculations. We extracted the values of the strong coupling constants at different channels. Our results may be checked via different theoretical models and approaches. The obtained results may help us in constructing the strong interaction potential among the doubly heavy baryons and the pseudoscalar mesons. Our results may also help experimental groups in analyses of the obtained related data in hadron colliders.

Appendix: The pion distribution amplitudes

In this appendix, we present explicit expressions for the DAs of the π meson. For more information see Refs. [47, 48].

$$\begin{aligned}
\langle \pi(p) | \bar{q}(x) \gamma_\mu \gamma_5 q(0) | 0 \rangle &= -i f_\pi p_\mu \int_0^1 du e^{i\bar{u} p x} \left(\varphi_\pi(u) + \frac{1}{16} m_\pi^2 x^2 \mathbb{A}(u) \right) \\
&\quad - \frac{i}{2} f_\pi m_\pi^2 \frac{x_\mu}{p x} \int_0^1 du e^{i\bar{u} p x} \mathbb{B}(u), \\
\langle \pi(p) | \bar{q}(x) i \gamma_5 q(0) | 0 \rangle &= \mu_\pi \int_0^1 du e^{i\bar{u} p x} \varphi_P(u), \\
\langle \pi(p) | \bar{q}(x) \sigma_{\alpha\beta} \gamma_5 q(0) | 0 \rangle &= \frac{i}{6} \mu_\pi (1 - \tilde{\mu}_\pi^2) (p_\alpha x_\beta - p_\beta x_\alpha) \int_0^1 du e^{i\bar{u} p x} \varphi_\sigma(u), \\
\langle \pi(p) | \bar{q}(x) \sigma_{\mu\nu} \gamma_5 g_s G_{\alpha\beta}(vx) q(0) | 0 \rangle &= i \mu_\pi \left[p_\alpha p_\mu \left(g_{\nu\beta} - \frac{1}{p x} (p_\nu x_\beta + p_\beta x_\nu) \right) \right. \\
&\quad - p_\alpha p_\nu \left(g_{\mu\beta} - \frac{1}{p x} (p_\mu x_\beta + p_\beta x_\mu) \right) \\
&\quad - p_\beta p_\mu \left(g_{\nu\alpha} - \frac{1}{p x} (p_\nu x_\alpha + p_\alpha x_\nu) \right) \\
&\quad \left. + p_\beta p_\nu \left(g_{\mu\alpha} - \frac{1}{p x} (p_\mu x_\alpha + p_\alpha x_\mu) \right) \right] \\
&\quad \times \int \mathcal{D}\alpha e^{i(\alpha_{\bar{q}} + v\alpha_g) p x} \mathcal{T}(\alpha_i), \\
\langle \pi(p) | \bar{q}(x) \gamma_\mu \gamma_5 g_s G_{\alpha\beta}(vx) q(0) | 0 \rangle &= p_\mu (p_\alpha x_\beta - p_\beta x_\alpha) \frac{1}{p x} f_\pi m_\pi^2 \int \mathcal{D}\alpha e^{i(\alpha_{\bar{q}} + v\alpha_g) p x} \mathcal{A}_\parallel(\alpha_i) \\
&\quad + \left[p_\beta \left(g_{\mu\alpha} - \frac{1}{p x} (p_\mu x_\alpha + p_\alpha x_\mu) \right) \right. \\
&\quad \left. - p_\alpha \left(g_{\mu\beta} - \frac{1}{p x} (p_\mu x_\beta + p_\beta x_\mu) \right) \right] f_\pi m_\pi^2 \\
&\quad \times \int \mathcal{D}\alpha e^{i(\alpha_{\bar{q}} + v\alpha_g) p x} \mathcal{A}_\perp(\alpha_i), \\
\langle \pi(p) | \bar{q}(x) \gamma_\mu i g_s G_{\alpha\beta}(vx) q(0) | 0 \rangle &= p_\mu (p_\alpha x_\beta - p_\beta x_\alpha) \frac{1}{p x} f_\pi m_\pi^2 \int \mathcal{D}\alpha e^{i(\alpha_{\bar{q}} + v\alpha_g) p x} \mathcal{V}_\parallel(\alpha_i) \\
&\quad + \left[p_\beta \left(g_{\mu\alpha} - \frac{1}{p x} (p_\mu x_\alpha + p_\alpha x_\mu) \right) \right. \\
&\quad \left. - p_\alpha \left(g_{\mu\beta} - \frac{1}{p x} (p_\mu x_\beta + p_\beta x_\mu) \right) \right] f_\pi m_\pi^2 \\
&\quad \times \int \mathcal{D}\alpha e^{i(\alpha_{\bar{q}} + v\alpha_g) p x} \mathcal{V}_\perp(\alpha_i), \tag{35}
\end{aligned}$$

where

$$\mu_\pi = f_\pi \frac{m_\pi^2}{m_u + m_d}, \quad \tilde{\mu}_\pi = \frac{m_u + m_d}{m_\pi}. \tag{36}$$

Here, $\varphi_\pi(u)$, $\mathbb{A}(u)$, $\mathbb{B}(u)$, $\varphi_P(u)$, $\varphi_\sigma(u)$, $\mathcal{T}(\alpha_i)$, $\mathcal{A}_\perp(\alpha_i)$, $\mathcal{A}_\parallel(\alpha_i)$, $\mathcal{V}_\perp(\alpha_i)$ and $\mathcal{V}_\parallel(\alpha_i)$ are wave functions of definite twists.

References

- [1] M. Mattson *et al.* [SELEX Collaboration], “First Observation of the Doubly Charmed Baryon Ξ_{cc}^+ ,” *Phys. Rev. Lett.* **89**, 112001 (2002).
- [2] A. Ocherashvili *et al.* [SELEX Collaboration], “Confirmation of the double charm baryon $\Xi_{cc}^+(3520)$ via its decay to $p D^+ K^-$,” *Phys. Lett. B* **628**, 18 (2005).
- [3] S. P. Ratti, “New results on c-baryons and a search for cc-baryons in FOCUS,” *Nucl. Phys. Proc. Suppl.* **115**, 33 (2003).
- [4] B. Aubert *et al.* [BaBar Collaboration], “Search for doubly charmed baryons Ξ_{cc}^+ and Ξ_{cc}^{++} in BABAR,” *Phys. Rev. D* **74**, 011103 (2006).
- [5] R. Aaij *et al.* [LHCb Collaboration], “Search for the doubly charmed baryon Ξ_{cc}^+ ,” *JHEP* **1312**, 090 (2013).
- [6] Y. Kato *et al.* [Belle Collaboration], “Search for doubly charmed baryons and study of charmed strange baryons at Belle,” *Phys. Rev. D* **89**, no. 5, 052003 (2014).
- [7] R. Aaij *et al.* [LHCb Collaboration], “Observation of the doubly charmed baryon Ξ_{cc}^{++} ,” *Phys. Rev. Lett.* **119**, no. 11, 112001 (2017).
- [8] R. Aaij *et al.* [LHCb Collaboration], “First Observation of the Doubly Charmed Baryon Decay $\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$,” *Phys. Rev. Lett.* **121**, no. 16, 162002 (2018).
- [9] R. Aaij *et al.* [LHCb Collaboration], “Search for the doubly charmed baryon Ξ_{cc}^+ ,” *Sci. China Phys. Mech. Astron.* **63**, no. 2, 221062 (2020).
- [10] T. M. Aliev, K. Azizi and M. Savci, “Doubly Heavy Spin-1/2 Baryon Spectrum in QCD,” *Nucl. Phys. A* **895**, 59 (2012).
- [11] T. M. Aliev, K. Azizi and M. Savci, “The masses and residues of doubly heavy spin-3/2 baryons,” *J. Phys. G* **40**, 065003 (2013).
- [12] K. Azizi, T. M. Aliev and M. Savci, “Properties of doubly and triply heavy baryons,” *J. Phys. Conf. Ser.* **556**, no. 1, 012016 (2014).
- [13] W. Wang, F. S. Yu and Z. X. Zhao, “Weak decays of doubly heavy baryons: the $1/2 \rightarrow 1/2$ case,” *Eur. Phys. J. C* **77**, no. 11, 781 (2017).
- [14] W. Wang, Z. P. Xing and J. Xu, “Weak Decays of Doubly Heavy Baryons: SU(3) Analysis,” *Eur. Phys. J. C* **77**, no. 11, 800 (2017).
- [15] T. Gutsche, M. A. Ivanov, J. G. Krner and V. E. Lyubovitskij, “Decay chain information on the newly discovered double charm baryon state Ξ_{cc}^{++} ,” *Phys. Rev. D* **96**, no. 5, 054013 (2017).
- [16] H. S. Li, L. Meng, Z. W. Liu and S. L. Zhu, “Radiative decays of the doubly charmed baryons in chiral perturbation theory,” *Phys. Lett. B* **777**, 169 (2018).
- [17] L. Y. Xiao, K. L. Wang, Q. f. Lu, X. H. Zhong and S. L. Zhu, “Strong and radiative decays of the doubly charmed baryons,” *Phys. Rev. D* **96**, no. 9, 094005 (2017).

- [18] N. Sharma and R. Dhir, “Estimates of W-exchange contributions to Ξ_{cc} decays,” *Phys. Rev. D* **96**, no. 11, 113006 (2017).
- [19] Y. L. Ma and M. Harada, “Chiral partner structure of doubly heavy baryons with heavy quark spin-flavor symmetry,” *J. Phys. G* **45**, no. 7, 075006 (2018).
- [20] X. H. Hu, Y. L. Shen, W. Wang and Z. X. Zhao, “Weak decays of doubly heavy baryons: ”decay constants”,” *Chin. Phys. C* **42**, no. 12, 123102 (2018).
- [21] Y. J. Shi, W. Wang, Y. Xing and J. Xu, “Weak Decays of Doubly Heavy Baryons: Multi-body Decay Channels,” *Eur. Phys. J. C* **78**, no. 1, 56 (2018).
- [22] X. Yao and B. Mller, “Doubly charmed baryon production in heavy ion collisions,” *Phys. Rev. D* **97**, no. 7, 074003 (2018).
- [23] D. L. Yao, “Masses and sigma terms of doubly charmed baryons up to $\mathcal{O}(p^4)$ in manifestly Lorentz-invariant baryon chiral perturbation theory,” *Phys. Rev. D* **97**, no. 3, 034012 (2018).
- [24] Z. X. Zhao, “Weak decays of doubly heavy baryons: the $1/2 \rightarrow 3/2$ case,” *Eur. Phys. J. C* **78**, no. 9, 756 (2018).
- [25] Z. G. Wang, “Analysis of the doubly heavy baryon states and pentaquark states with QCD sum rules,” *Eur. Phys. J. C* **78**, no. 10, 826 (2018).
- [26] M. Z. Liu, Y. Xiao and L. S. Geng, “Magnetic moments of the spin-1/2 doubly charmed baryons in covariant baryon chiral perturbation theory,” *Phys. Rev. D* **98**, no. 1, 014040 (2018).
- [27] Z. P. Xing and Z. X. Zhao, “Weak decays of doubly heavy baryons: the FCNC processes,” *Phys. Rev. D* **98**, no. 5, 056002 (2018).
- [28] R. Dhir and N. Sharma, “Weak decays of doubly heavy charm Ω_{cc}^+ baryon,” *Eur. Phys. J. C* **78**, no. 9, 743 (2018).
- [29] A. V. Berezhnoy, A. K. Likhoded and A. V. Luchinsky, “Doubly heavy baryons at the LHC,” *Phys. Rev. D* **98**, no. 11, 113004 (2018).
- [30] L. J. Jiang, B. He and R. H. Li, “Weak decays of doubly heavy baryons: $\mathcal{B}_{cc} \rightarrow \mathcal{B}_c V$,” *Eur. Phys. J. C* **78**, no. 11, 961 (2018).
- [31] Q. A. Zhang, “Weak Decays of Doubly Heavy Baryons: W-Exchange,” *Eur. Phys. J. C* **78**, no. 12, 1024 (2018).
- [32] T. Gutsche, M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij and Z. Tyulemissov, “Ab initio three-loop calculation of the W-exchange contribution to nonleptonic decays of double charm baryons,” *Phys. Rev. D* **99**, no. 5, 056013 (2019).
- [33] Y. J. Shi, Y. Xing and Z. X. Zhao, “Light-cone sum rules analysis of $\Xi_{QQ'q} \rightarrow \Lambda_{Q'}$ weak decays,” *Eur. Phys. J. C* **79**, no. 6, 501 (2019).
- [34] X. H. Hu and Y. J. Shi, “Light-Cone Sum Rules Analysis of $\Xi_{QQ'} \rightarrow \Sigma_{Q'}$ Weak Decays,” *Eur. Phys. J. C* **80**, no. 1, 56 (2020).

- [35] S. J. Brodsky, F. K. Guo, C. Hanhart and U.-G. Meißner, “Isospin splittings of doubly heavy baryons,” *Phys. Lett. B* **698**, 251 (2011).
- [36] M. J. Yan, X. H. Liu, S. Gonzalez-Solis, F. K. Guo, C. Hanhart, U.-G. Meißner and B. S. Zou, “New spectrum of negative-parity doubly charmed baryons: Possibility of two quasistable states,” *Phys. Rev. D* **98**, no. 9, 091502 (2018).
- [37] H. Y. Cheng, G. Meng, F. Xu and J. Zou, “Two-body weak decays of doubly charmed baryons,” *Phys. Rev. D* **101**, no. 3, 034034 (2020).
- [38] U. Özdem, “Magnetic moments of doubly heavy baryons in light-cone QCD,” *J. Phys. G* **46**, no. 3, 035003 (2019).
- [39] V. L. Chernyak and I. R. Zhitnitsky, “B meson exclusive decays into baryons,” *Nucl. Phys. B* **345**, 137 (1990).
- [40] A. Khodjamirian, R. Ruckl, S. Weinzierl and O. I. Yakovlev, “Perturbative QCD correction to the $B \rightarrow \pi$ transition form-factor,” *Phys. Lett. B* **410**, 275 (1997)
- [41] E. Bagan, P. Ball and V. M. Braun, “Radiative corrections to the decay $B \rightarrow \pi e \nu$ and the heavy quark limit,” *Phys. Lett. B* **417**, 154 (1998)
- [42] P. Ball, “ $B \rightarrow \pi$ and $B \rightarrow K$ transitions from QCD sum rules on the light cone,” *JHEP* **9809**, 005 (1998).
- [43] P. Ball, V. M. Braun and N. Kivel, *Nucl. Phys. B* **649**, 263 (2003).
- [44] K. Azizi, A. R. Olamaei and S. Rostami, “Beautiful mathematics for beauty-full and other multi-heavy hadronic systems,” *Eur. Phys. J. A* **54**, no. 9, 162 (2018).
- [45] S. S. Agaev, K. Azizi and H. Sundu, “Application of the QCD light cone sum rule to tetraquarks: the strong vertices $X_b X_{b\rho}$ and $X_c X_{c\rho}$,” *Phys. Rev. D* **93**, no. 11, 114036 (2016).
- [46] M. Tanabashi *et al.* [Particle Data Group], “Review of Particle Physics,” *Phys. Rev. D* **98**, no. 3, 030001 (2018).
- [47] P. Ball, “Theoretical update of pseudoscalar meson distribution amplitudes of higher twist: The Nonsinglet case,” *JHEP* **9901**, 010 (1999).
- [48] P. Ball and R. Zwicky, “New results on $B \rightarrow \pi, K, \eta$ decay formfactors from light-cone sum rules,” *Phys. Rev. D* **71**, 014015 (2005).
- [49] T. Aliev, K. Azizi and M. Savci, “Strong coupling constants of light pseudoscalar mesons with heavy baryons in QCD,” *Phys. Lett. B* **696**, 220-226 (2011).