

THE ERDŐS-HAJNAL PROPERTY FOR GRAPHS WITH NO FIXED CYCLE AS A PIVOT-MINOR

JAEHOON KIM AND SANG-IL OUM

ABSTRACT. We prove that for every integer k , there exists $\varepsilon > 0$ such that for every n -vertex graph G with no pivot-minor isomorphic to C_k , there exist disjoint sets $A, B \subseteq V(G)$ such that $|A|, |B| \geq \varepsilon n$, and A is either complete or anticomplete to B . This proves the analog of the Erdős-Hajnal conjecture for the class of graphs with no pivot-minor isomorphic to C_k .

1. INTRODUCTION

In this paper all graphs are simple, having no loops and no parallel edges. For a graph G , let $\omega(G)$ be the maximum size of a clique, that is a set of pairwise adjacent vertices and let $\alpha(G)$ be the maximum size of an independent set, that is a set of pairwise non-adjacent vertices. Erdős and Hajnal [9] proposed the following conjecture in 1989.

Conjecture 1.1 (Erdős and Hajnal [9]). *For every graph H , there is $\varepsilon > 0$ such that all graphs G with no induced subgraph isomorphic to H satisfies*

$$\max(\alpha(G), \omega(G)) \geq |V(G)|^\varepsilon.$$

This conjecture still remains open. See [5] for a survey on this conjecture. We can ask the same question for weaker containment relations. Recently Chudnovsky and Oum [6] proved that this conjecture holds if we replace “induced subgraphs” with “vertex-minors” as follows. This is weaker in the sense that every induced subgraph G is a vertex-minor of G but not every vertex-minor of G is an induced subgraph of G .

Theorem 1.1 (Chudnovsky and Oum [6]). *For every graph H , there exists $\varepsilon > 0$ such that every graph with no vertex-minors isomorphic to H satisfies*

$$\max(\alpha(G), \omega(G)) \geq |V(G)|^\varepsilon.$$

We ask whether Conjecture 1.1 holds if we replace “induced subgraphs” with “pivot-minors” as follows.

Conjecture 1.2. *For every graph H , there exists $\varepsilon > 0$ such that every graph G with no pivot-minor isomorphic to H satisfies*

$$\max(\alpha(G), \omega(G)) \geq |V(G)|^\varepsilon.$$

The detailed definition of pivot-minors will be presented in Section 3. For now, note that the analog for vertex-minors is weakest, the analog for pivot-minors is weaker than that for induced subgraphs but stronger than that for vertex-minors. This is because every induced subgraph of G is a pivot-minor of G , and every pivot-minor of G is a vertex-minor of G . In

Date: July 2, 2021.

J.K. was supported by the POSCO Science Fellowship of POSCO TJ Park Foundation and by the KAIX Challenge program of KAIST Advanced Institute for Science-X (J. Kim). S.O. was supported by the Institute for Basic Science (IBS-R029-C1).

other words, Conjecture 1.1 implies Conjecture 1.2 and Conjecture 1.2 implies Theorem 1.1. We verify Conjecture 1.2 for $H = C_k$, the cycle graph on k vertices as follows.

Theorem 1.2. *For every $k \geq 3$, there exists $\varepsilon > 0$ such that every graph with no pivot-minor isomorphic to C_k satisfies*

$$\max(\alpha(G), \omega(G)) \geq |V(G)|^\varepsilon.$$

We actually prove a stronger property, as Chudnovsky and Oum [6] did. Before stating this property, let us first state a few terminologies. A class \mathcal{G} of graphs closed under taking induced subgraphs is said to have *the Erdős-Hajnal property* if there exists $\varepsilon > 0$ such that every graph G in \mathcal{G} satisfies

$$\max(\alpha(G), \omega(G)) \geq |V(G)|^\varepsilon.$$

A class \mathcal{G} of graphs closed under taking induced subgraphs is said to have *the strong Erdős-Hajnal property* if there exists $\varepsilon > 0$ such that every n -vertex graph in \mathcal{G} with $n > 1$ has disjoint sets A, B of vertices such that $|A|, |B| \geq \varepsilon n$ and A is either complete or anti-complete to B . It is an easy exercise to show that the strong Erdős-Hajnal property implies the Erdős-Hajnal property, see [1, 10].

Chudnovsky and Oum [6] proved that the class of graphs with no vertex-minors isomorphic to H for a fixed graph H has the strong Erdős-Hajnal property, implying Theorem 1.1. We propose its analog for pivot-minors as a conjecture, which implies the theorem of Chudnovsky and Oum [6]. Note that this conjecture is not true if we replace the pivot-minor with induced graphs. For example, the class of triangle-free graphs does not have the strong Erdős-Hajnal property [10].

Conjecture 1.3. *For every graph H , there exists $\varepsilon > 0$ such that for all $n > 1$, every n -vertex graph with no pivot-minor isomorphic to H has two disjoint sets A, B of vertices such that $|A|, |B| \geq \varepsilon n$ and A is complete or anti-complete to B .*

We prove that this conjecture holds if $H = C_k$. In other words, the class of graphs with no pivot-minor isomorphic to C_k has the strong Erdős-Hajnal property as follows. This implies Theorem 1.2.

Theorem 1.3. *For every integer $k \geq 3$, there exists $\varepsilon > 0$ such that for all $n > 1$, every n -vertex graph with no pivot-minor isomorphic to C_k has two disjoint sets A, B of vertices such that $|A|, |B| \geq \varepsilon n$ and A is complete or anti-complete to B .*

This paper is organized as follows. In Section 2, we will introduce basic definitions and review necessary theorems of Rödl [20] and Bonamy, Bousquet, and Thomassé [2]. In Section 3, we will present several tools to find a pivot-minor isomorphic to C_k . In particular, it proves that a long anti-hole contains C_k as a pivot-minor. In Section 4, we will present the proof of the main theorem, Theorem 1.3. In Section 5, we will relate our theorem to the problem on χ -boundedness, and discuss known results and open problems related to polynomial χ -boundedness and the Erdős-Hajnal property.

2. PRELIMINARIES

Let \mathbb{N} be the set of positive integers and for each $n \in \mathbb{N}$, we write $[n] := \{1, 2, \dots, n\}$. For a graph $G = (V, E)$, let $\overline{G} = (V, \binom{V}{2} - E)$ be the complement of G . We write $\Delta(G)$ and $\delta(G)$ to denote the maximum degree of G and the minimum degree of G respectively.

Let T be a tree rooted at a specified node v_r , called the *root*. If the path from v_r to a node y in T contains $x \in V(T) - \{y\}$, we say that x is an *ancestor* of y , and y is a *descendant* of x . If one of x and y is an ancestor of the other, we say that x, y are *related*. We say that two disjoint sets X and Y of nodes of T are *unrelated* if no pairs of $x \in X$ and $y \in Y$ are related.

For disjoint vertex sets X and Y , we say X is *complete* to Y if every vertex of X is adjacent to all vertices of Y . We say X is *anti-complete* to Y if every vertex of X is non-adjacent to Y . A *pure pair* of a graph G is a pair (A, B) of disjoint subsets of $V(G)$ such that A is complete or anticomplete to B .

For a vertex u , let $N_G(u)$ denote the set of neighbors of u in G . For each $U \subseteq V(G)$, we write

$$N_G(U) := \bigcup_{u \in U} N_G(u) - U.$$

The following lemma is proved in Section 2 of [2].

Lemma 2.1 (Bonamy, Bousquet, and Thomassé [2]). *For every connected graph G and a vertex $v_r \in V(G)$, there exist an induced subtree T of G rooted at v_r and a function $r : V(G) \rightarrow V(T)$ satisfying the following.*

- (T1) $r(v_r) = v_r$ and for each $u \in V(G) - \{v_r\}$, the vertex $r(u)$ is a neighbor of u . In particular, T is a dominating tree of G .
- (T2) If $r(x)$ and $r(y)$ are not related, then $xy \notin E(G)$.

Rödl [20] proved the following theorem. Its weaker version was later proved by Fox and Sudakov [11] without using the regularity lemma. A set U of vertices of G is an ε -stable set of a graph G if $G[U]$ has at most $\varepsilon \binom{|U|}{2}$ edges. Similarly, U is an ε -clique of a graph G if $G[U]$ has at least $(1 - \varepsilon) \binom{|U|}{2}$ edges.

Theorem 2.2 (Rödl [20]). *For all $\varepsilon > 0$ and a graph H , there exists $\delta > 0$ such that every n -vertex graph G with no induced subgraph isomorphic to H has an ε -stable set or an ε -clique of size at least δn .*

We will use the following simple lemma. We present its proof for completeness.

Lemma 2.3. *Let G be a graph. Every ε -stable set U of G has a subset U' of size at least $|U|/2$ with $\Delta(G[U']) \leq 4\varepsilon|U'|$.*


Proof. Let U' be the set of vertices of degree at most $2\varepsilon|U|$ in $G[U]$. Because $\sum_{v \in U} \deg_{G[U]}(v) < \varepsilon|U|^2$, we have $|U'| \geq |U|/2$. Moreover, for each vertex $v \in U'$, we have $\deg_{G[U']}(v) \leq 2\varepsilon|U| \leq 4\varepsilon|U'|$. \square

Using Lemma 2.3, we can deduce the following corollary of Theorem 2.2.

Corollary 2.4. *For all $\alpha > 0$ and a graph H , there exists $\delta > 0$ such that every graph G with no induced subgraph isomorphic to H has a set $U \subseteq V(G)$ with $|U| \geq \delta|V(G)|$ such that either $\Delta(G[U]) \leq \alpha|U|$ or $\Delta(\overline{G}[U]) \leq \alpha|U|$.*

The following easy lemma will be used to find a connected induced subgraph inside the output of Corollary 2.4. We omit its easy proof.

Lemma 2.5. *A graph G has a pure pair (A, B) such that $|A|, |B| \geq |V(G)|/3$ or has a connected induced subgraph H such that $|V(H)| \geq |V(G)|/3$.*

FIGURE 1. Pivoting uv .

Lemma 2.6 (Bonamy, Bousquet, and Thomassé [2, Lemma 3]). *Let T be a tree rooted at v_r and $w : V(T) \rightarrow \mathbb{R}$ be a non-negative weight function on $V(T)$ with $\sum_{x \in V(T)} w(x) = 1$. Then there exists either a path P from v_r with weight at least $1/4$ or two unrelated sets A and B both with weight at least $1/4$.*

A *hole* is an induced cycle of length at least 5.

Lemma 2.7 (Bonamy, Bousquet, and Thomassé [2, Lemma 4]). *For given $k \geq 3$, there exist $\alpha = \alpha(k) > 0$ and $\varepsilon = \varepsilon(k) > 0$ such that for any n -vertex graph G with $n \geq 2$ and $\Delta(G) \leq \alpha n$, if G has no holes of length at least k and has a dominating induced path, then G contains a pair (A, B) of disjoint vertex sets such that A is anticomplete to B and $|A|, |B| \geq \varepsilon n$.*

3. FINDING A CYCLE AS A PIVOT-MINOR

For a given graph G and an edge uv , a graph $G \wedge uv$ obtained from G by *pivoting* uv is defined as follows. Let $V_1 = N_G(u) \cap N_G(v)$, $V_2 = N_G(u) - N_G(v)$, $V_3 = N_G(v) - N_G(u)$. Then $G \wedge uv$ is the graph obtained from G by complementing adjacency between vertices between V_i and V_j for all $1 \leq i < j \leq 3$ and swapping the label of u and v . See Figure 1 for an illustration. We say that H is a *pivot-minor* of G if H can be obtained from G by deleting vertices and pivoting edges. For this paper, we will also say that H is a pivot-minor of G , when G has a pivot-minor isomorphic to H . A pivot-minor H of G is *proper* if $|V(H)| < |V(G)|$.

We describe several scenarios for constructing C_k as a pivot-minor. The following proposition is an easy one; One can obtain a desired pivot-minor from a longer cycle of the same parity.

Proposition 3.1. *For $m \geq k \geq 3$ with $m \equiv k \pmod{2}$, the cycle C_m has a pivot-minor isomorphic to C_k .*

Proof. We proceed by induction on $m - k$. We may assume that $m > k$. Let xy be an edge of C_m . Then $(C_m \wedge xy) - x - y$ is isomorphic to C_{m-2} , which contains a pivot-minor isomorphic to C_k by the induction hypothesis. \square

Proposition 3.2. *For integers $k \geq 3$ and $m \geq \frac{3}{2}k + 6$, the graph $\overline{C_m}$ has a pivot-minor isomorphic to C_k .*

Before proving Proposition 3.2, we present a simple lemma on partial complements of the cycle graph. The *partial complement*¹ $G \oplus S$ of a graph G by a set S of vertices is a graph

¹We found this concept in a paper by Kamiński, Lozin, and Milanič [14], though it may have been studied previously, as it is a natural concept.




FIGURE 2. Obtaining an $(s-2, 3)$ -cycle from an $(s, 9)$ -cycle when $s > 9$ in the proof of Lemma 3.3.

obtained from G by changing all edges within S to non-edges and non-edges within S to edges.

For $s \geq t \geq 0$, we say that G is an (s, t) -cycle if G is isomorphic to a graph $C_s \oplus X$ for a set X of t consecutive vertices in the cycle C_s .

Lemma 3.3. *Let $s \geq t \geq 6$. An (s, t) -cycle contains a pivot-minor isomorphic to an $(s-2, t-6)$ -cycle.*

Proof. Let v_1, \dots, v_s be the vertices of C_s in the cyclic order where $X = \{v_1, \dots, v_t\}$. Then it is easy to check that $(C_s \oplus X) \wedge v_2 v_{t-1} - \{v_2, v_{t-1}\}$ is isomorphic to $C_{s-2} \oplus X'$ where X' consists of $t-6$ consecutive vertices on the cycle. See Figure 2. \square

Proof of Proposition 3.2. As $\overline{C_m}$ is an (m, m) -cycle, by Lemma 3.3, $\overline{C_m}$ contains a pivot-minor isomorphic to an $(m-2i, m-6i)$ -cycle for all $i \leq m/6$.

Let us fix $i = \lceil (k-2)/4 \rceil$. Then $m-6i \geq m-6 \cdot (k+1)/4 \geq 9/2$ and therefore $\overline{C_m}$ contains a pivot-minor H isomorphic to an $(m-2i, m-6i)$ -cycle and $m-6i \geq 5$. We may assume that $H = C_{m-2i} \oplus X$ where $C_{m-2i} = v_1 \cdots v_{m-2i}$ and $X = \{v_{4i+1}, \dots, v_{m-2i}\}$.

Note that H contains an induced cycle $C = v_1 \cdots v_{4i} v_{4i+1} v_{m-2i} v_1$ of length $4i+2 \geq k$. If k is even, then by Proposition 3.1, H contains a pivot-minor isomorphic to C_k . So we may assume that k is odd and therefore $|V(C)| = 4i+2 \geq k+1$.

Let $x = v_{m-2i}$, $y = v_{4i+1}$ be the two vertices in $V(C) \cap X$. Since $m-6i \geq 5$, there is a common neighbor z of x and y in X . Then z has exactly two neighbors x and y in $V(C)$. Then $H[V(C) \cup \{z\}] \wedge yz - y - z$ is a cycle of length $4i+1$. Since $4i+1 \geq k$, by Proposition 3.1, it contains a pivot-minor isomorphic to C_k . \square

A *generalized fan* is a graph G with a specified vertex c , called the *center*, such that $G - c$ is an induced path of length at least 1, called the *main path* of G and both ends of the main path are adjacent to c . If c is adjacent to all vertices of $G - c$, then G is called a *fan*.

An *interval* of a generalized fan with a center c is a maximal subpath of the main path having no internal vertex adjacent to c . The *length* of an interval is its number of edges. A generalized fan is an (a_1, \dots, a_s) -fan if the lengths of intervals are a_1, \dots, a_s in order. Note that an (a_1, \dots, a_s) -fan is also an (a_s, \dots, a_1) -fan. An (a_1, \dots, a_s) -fan is a *k-good fan* if $a_1 \geq k-2$ or $a_s \geq k-2$. An (a_1, \dots, a_s) -fan is a *strongly k-good fan* if $s \geq 2$ and either

$a_1 \geq k - 2$ and a_s is odd, or $a_s \geq k - 2$ and a_1 is odd. It is easy to observe that every k -good fan has a hole of length at least k . However, that does not necessarily lead to a pivot-minor isomorphic to C_k because of the parity issues. In the next proposition, we show that every strongly k -good fan has a pivot-minor isomorphic to C_k .

Proposition 3.4. *Let $k \geq 5$ be an integer. Every strongly k -good fan has a pivot-minor isomorphic to C_k .*

Proof. Let G be an (a_1, \dots, a_s) -fan such that $s \geq 2$, $a_1 \geq k - 2$, and a_s is odd. We proceed by the induction on $|V(G)|$. We may assume that G has no proper pivot-minor that is a strongly k -good fan. Note that C_{a_1+2} is an induced subgraph of G , hence if $a_1 \equiv k \pmod{2}$, then C_k is isomorphic to a pivot-minor of G by Proposition 3.1. Thus we may assume that $a_1 \not\equiv k \pmod{2}$ and so $a_1 \geq k - 1$.

If a_i is odd for some $1 < i < s$, then G contains a smaller strongly k -good fan by taking the first i intervals, contradicting our assumption. Thus a_i is even for all $1 < i < s$. If $a_i \geq 3$ for some $i > 1$, then let uv be an internal edge of the i -th interval. Then $G \wedge uv - u - v$ is a strongly k -good fan, contradicting our assumption. Thus, we may assume that $a_i \leq 2$ for all $i > 1$ and so G is an $(a_1, 2, \dots, 2, 1)$ -fan.

Let xy be the last interval of G with length 1. Then $G \wedge xy - x - y$ is a $(a_1, 2, \dots, 2, 1)$ -fan with $s - 1$ intervals. By the assumption, we may assume that $s = 2$ and $G \wedge xy - x - y$ is an $(a_1 - 1)$ -fan with one interval, which is a cycle with $a_1 + 1$ edges. As $a_1 + 1 \geq k$ and $a_1 + 1 \equiv k \pmod{2}$, Proposition 3.1 implies that G contains a pivot-minor isomorphic to C_k . \square

4. PROOF OF THEOREM 1.3

First we choose $\alpha > 0$ and $\varepsilon_0 > 0$ so that

$$(1) \quad 4\alpha \leq \alpha(\lceil \frac{3}{2}k + 6 \rceil) \text{ and } \varepsilon_0 = \varepsilon(\lceil \frac{3}{2}k + 6 \rceil) \text{ where } \alpha(\cdot), \varepsilon(\cdot) \text{ are specified in Lemma 2.7}$$

and in addition $\alpha < 1/(8k)$ as well. Let $\delta > 0$ be a constant obtained by applying Corollary 2.4 with $\alpha/3$ as α and C_k as H . Choose $\varepsilon > 0$ so that

$$\varepsilon < \min \left(\frac{\delta}{12}, (1 - 4(k + 3)\alpha) \frac{\delta}{240}, \frac{\varepsilon_0 \delta}{12} \right).$$

Let $n > 1$ be an integer and G be an n -vertex graph with no pivot-minor isomorphic to C_k . In particular, G does not have C_k as an induced subgraph. To derive a contradiction, we assume that G contains no pure pair (A, B) with $|A|, |B| \geq \varepsilon n$. We may assume that $\varepsilon n > 1$, because otherwise an edge or a non-edge of G gives a pure pair.

By Corollary 2.4, there exists a subset U of $V(G)$ such that $|U| \geq \delta |V(G)|$ and $\Delta(G^0)[U] \leq (\alpha/3)|U|$ for some $G^0 \in \{G, \overline{G}\}$. By the assumption on G , $G^0[U]$ has no pure pair (A, B) with $|A|, |B| \geq (\varepsilon/\delta)|U|$. As $\varepsilon/\delta < 1/3$, by Lemma 2.5, $G^0[U]$ has a connected induced subgraph G' such that $|V(G')| \geq |U|/3$. Let $n' = |V(G')|$.

Then $n' \geq (\delta/3)n$ and $\Delta(G') \leq (\alpha/3)|U| \leq \alpha n'$. By the assumption on G ,

$$(2) \quad G' \text{ contains no pure pair } (A, B) \text{ with } |A|, |B| \geq (3\varepsilon/\delta)n'.$$

By applying Lemma 2.1 with G' , we obtain a dominating induced tree T and $r : V(G') \rightarrow V(T)$ satisfying Lemma 2.1 (T1)–(T2) with G' . For each $u \in V(T)$, let

$$w(u) := \frac{|r^{-1}(\{u\})|}{n'}$$

be the *weight* of u . By applying Lemma 2.6 with the weight w , we obtain either an induced path P of T with weight at least $1/4$ or two unrelated sets A and B both with weight at least $1/4$.

In the latter case, Lemma 2.1 (T2) implies that $r^{-1}(A)$ is anticomplete to $r^{-1}(B)$ in G' and $|r^{-1}(A)|, |r^{-1}(B)| \geq n'/4 \geq (3\varepsilon/\delta)n'$, contradicting (2).

Hence, there exists an induced path P in G' with $|V(P) \cup N_{G'}(V(P))| \geq n'/4$. Let $W := V(P) \cup N_{G'}(V(P))$. Note that $n'/4 \geq \delta n/12 > \varepsilon n > 1$ and so $|W| \geq 2$.

Suppose that G' is an induced subgraph of \overline{G} . Using (1), we apply Lemma 2.7 to $G'[W]$ with 4α as α and $\lceil \frac{3}{2}k + 6 \rceil$ as k . Then we can deduce from (2) and $\varepsilon'|W| \geq \frac{12\varepsilon}{\delta} \frac{n'}{4} = (3\varepsilon/\delta)n'$ that the graph $G'[W]$ contains an induced cycle C_m with $m \geq \lceil \frac{3}{2}k + 6 \rceil$ and by Proposition 3.2, $\overline{G'}$ contains a pivot-minor isomorphic to C_k , and so does G , a contradiction.

Thus G' is an induced subgraph of G . Let $G^* := G'[W]$ and let $n^* = |W|$. Then G^* has no pivot-minor isomorphic to C_k , $n^* \geq n'/4$, and $\Delta(G^*) \leq 4\alpha n^*$. By (2), G^* contains no pure pair (A, B) with $|A|, |B| \geq (12\varepsilon/\delta)n^*$. Now the theorem follows from applying the following lemma with $G^*, n^*, 4\alpha, 12\varepsilon/\delta$ playing the roles of $G, n, \alpha, \varepsilon$ respectively in the statement of the lemma.

Lemma 4.1. *Let $k \geq 3$ be an integer. Let $0 < \alpha < 1/(2k)$, $0 < \varepsilon \leq (1 - (k + 3)\alpha)/20$. Let G be a graph on $n \geq 2$ vertices such that $\Delta(G) \leq \alpha n$ and G has no pure pair (A, B) with $|A|, |B| \geq \varepsilon n$. If G has a dominating induced path P , then G has a pivot-minor isomorphic to C_k .*

Proof. Suppose that G has no pivot-minor isomorphic to C_k . Note that $\varepsilon n > 1$ as otherwise we have a pure pair on two vertices since $n \geq 2$. Let us label vertices of P by $1, 2, \dots, s$ in the order.

As P is a dominating path of G and $1 \leq \Delta(G) \leq \alpha n$, we have $2\alpha ns \geq (\alpha n + 1)s \geq n$ and therefore

$$s \geq 1/(2\alpha).$$

Note that $s - k > 0$ because $\alpha < \frac{1}{2k}$. As P is an induced path, it contains a pure pair (A, B) with $|A|, |B| \geq \lfloor \frac{s-1}{2} \rfloor$ and so $\frac{s-2}{2} \leq \lfloor \frac{s-1}{2} \rfloor < \varepsilon n$. Because $\varepsilon n > 1$, we have $2\varepsilon n + 2 < 4\varepsilon n$ and so

$$(3) \quad s < 2\varepsilon n + 2 < 4\varepsilon n.$$

Now, for each $i \in [s - k + 1]$, let

$$U_i^- := \{1, \dots, i - 1\}, \quad U_i^0 := \{i, \dots, i + k - 1\}, \quad \text{and} \quad U_i^+ := \{i + k, \dots, s\}.$$

In other words, this partitions P into three (possibly empty) subpaths. Furthermore, for all $i \in [s - k + 1]$ and $u \in N_G(U_i^-) - V(P)$, let

$$m_i^-(u) := \max(N_G(u) \cap U_i^-)$$

and for all $i \in [s - k + 1]$ and $u \in N_G(U_i^+) - V(P)$, let

$$m_i^+(u) := \min(N_G(u) \cap U_i^+),$$

indicating the largest neighbor of u in U_i^- and the smallest neighbor of u in U_i^+ respectively. For each $i \in [s - k + 1]$, let

$$A_i := N_G(U_i^0) - V(P) \text{ and}$$

$$B_i := (N_G(U_i^-) \cap N_G(U_i^+)) - (A_i \cup V(P)).$$




FIGURE 3. $s' = i + k - 1$. Bold lines indicate $m_i^-(u)$ and $m_i^+(u)$.

Note that for each $u \in B_i$, we have

$$(4) \quad m_i^+(u) - m_i^-(u) \not\equiv k \pmod{2},$$

because otherwise $(u, m_i^-(u), m_i^-(u) + 1, \dots, m_i^+(u), u)$ forms an induced cycle of length at least k and Proposition 3.1 implies that G contains a pivot-minor isomorphic to C_k , a contradiction.

For each $i \in [s - k + 1]$, let

$$\begin{aligned} C_i^1 &:= \{u \in N_G(U_i^-) - (A_i \cup B_i \cup V(P)) : m_i^-(u) \equiv 1 \pmod{2}\}, \\ C_i^2 &:= \{u \in N_G(U_i^-) - (A_i \cup B_i \cup V(P)) : m_i^-(u) \equiv 0 \pmod{2}\}, \\ D_i^1 &:= \{u \in N_G(U_i^+) - (A_i \cup B_i \cup V(P)) : m_i^+(u) \equiv k \pmod{2}\}, \text{ and} \\ D_i^2 &:= \{u \in N_G(U_i^+) - (A_i \cup B_i \cup V(P)) : m_i^+(u) \equiv k + 1 \pmod{2}\}. \end{aligned}$$

Recall that P is dominating. Hence, for each i , the sets $\{A_i, B_i, C_i^1, C_i^2, D_i^1, D_i^2, V(P)\}$ forms a partition of $V(G)$ into 7 possibly empty sets.

If there exists an edge between $u \in C_i^j$ and $v \in D_i^j$ for some $j \in [2]$, then we obtain an induced cycle $(u, m_i^-(u), m_i^-(u) + 1, \dots, m_i^+(v), v, u)$ having length $m_i^+(v) - m_i^-(u) + 3 > k$ and $m_i^+(v) - m_i^-(u) + 3 \equiv k \pmod{2}$, contradicting our assumption that G has no pivot-minor isomorphic to C_k by Proposition 3.1. Thus C_i^j is anticomplete to D_i^j . Hence,

$$(5) \quad \min\{|C_i^j|, |D_i^j|\} < \varepsilon n.$$

for all $i \in [s - k + 1]$ and $j \in [2]$. Furthermore, we prove the following.

Claim 1. *Let $i \in [s - k + 1]$. For each $v \in B_i$, all integers in $N_G(v) \cap U_i^-$ have the same parity and all integers in $N_G(v) \cap U_i^+$ have the same parity.*

Proof of Claim 1. If $N_G(v) \cap U_i^+$ has two integers $a < b$ of the different parity, then G contains a strongly k -good generalized fan by taking a subpath of P from $m_i^-(v)$ to b as its main path and v as its center. Then by Proposition 3.4, G contains a pivot-minor isomorphic to C_k ,

contradicting the assumption. Thus all integers in $N_G(v) \cap U_i^+$ have the same parity and similarly all integers in $N_G(v) \cap U_i^-$ have the same parity. ■

Claim 2. For all $i \in [s - k + 1]$, $|B_i| < 2(\alpha + 2\varepsilon)n$.

Proof of Claim 2. Suppose $|B_i| \geq 2(\alpha + 2\varepsilon)n$ for some $i \in [s - k + 1]$. Then there exists $r_B \in \{0, 1\}$ such that

$$B' := \{u \in B_i : m_i^-(u) \equiv r_B \pmod{2}\}$$

has size at least $(\alpha + 2\varepsilon)n$. By (4), $m_i^+(u) \equiv k + r_B + 1 \pmod{2}$ for all $u \in B'$.

We claim that if uv is an edge in $G[B']$, then $(m_i^-(u), m_i^+(u)) = (m_i^-(v), m_i^+(v))$. Suppose not. Without loss of generality, we may assume that $m_i^-(u) < m_i^-(v)$, because otherwise we may reverse the ordering of P to ensure that $m_i^-(u) \neq m_i^-(v)$ and swap u and v if necessary.

If $m_i^+(u) \geq m_i^+(v)$, then by Claim 1, $\{m_i^-(v), m_i^-(v) + 1, \dots, m_i^+(u), u, v\}$ induces a strongly k -good generalized fan with v as a center and $(m_i^-(v), m_i^-(v) + 1, \dots, m_i^+(u), u)$ as its main path. This implies that G has a pivot-minor isomorphic to C_k by Proposition 3.4, contradicting our assumption.

If $m_i^+(u) < m_i^+(v)$, then $(m_i^-(v), m_i^-(v) + 1, \dots, m_i^+(u), u, v)$ is an induced cycle of length $m_i^+(u) - m_i^-(v) + 3 \geq k$, and $m_i^+(u) - m_i^-(v) + 3 \equiv (k + r_B + 1) - r_B + 3 \equiv k \pmod{2}$, a contradiction by Proposition 3.1.

Hence, $(m_i^-(u), m_i^+(u)) = (m_i^-(v), m_i^+(v))$ for all $uv \in E(G[B'])$. Let C_1, \dots, C_t be the connected components of $G[B']$. By the above observation, for each $j \in [t]$, there exist $a_j \in U_i^-$ and $b_j \in U_i^+$ such that $V(C_j) \subseteq N_G(a_j) \cap N_G(b_j)$. So, $|V(C_j)| \leq \alpha n$. As $|B'| \geq (\alpha + 2\varepsilon)n$, there exists a set $I \subseteq \{1, 2, \dots, t\}$ such that $\varepsilon n \leq |\bigcup_{i \in I} V(C_i)| \leq (\alpha + \varepsilon)n$. Let $A := \bigcup_{i \in I} V(C_i)$ and $B := B' - A$. Then (A, B) is a pure pair of G with $|A|, |B| \geq \varepsilon n$, a contradiction. ■

Claim 3. There exist $i_* \in [s - k + 1]$ and $j_* \in [2]$ such that

$$|C_{i_*}^{j_*}|, |D_{i_*}^{3-j_*}| \geq 3\varepsilon n.$$

Proof of Claim 3. First, since $\Delta(G) \leq \alpha n$, $|A_i| \leq k\alpha n$ for each $i \in [s - k + 1]$.

Let $f(i) := |C_i^1| + |C_i^2|$. Then

$$f(1) = 0,$$

$$\begin{aligned} f(s - k + 1) &= n - |A_{s-k+1}| - s \quad \text{because } U_{s-k+1}^+ = D_{s-k+1}^1 = D_{s-k+1}^2 = B_{s-k+1} = \emptyset, \\ &\geq n - k\alpha n - 4\varepsilon n \quad \text{by (3) and the assumption that } \Delta(G) \leq \alpha n, \\ &= (1 - k\alpha - 4\varepsilon)n \geq 6\varepsilon n, \end{aligned}$$

and for each $i \in [s - k]$, we have

$$f(i + 1) - f(i) \leq \deg_G(i) \leq \alpha n.$$

Hence, there exists $i_* \in [s - k + 1]$ such that $6\varepsilon n \leq f(i_*) < (6\varepsilon + \alpha)n$. As $|B_{i_*}| < 2(\alpha + 2\varepsilon)n$, we have

$$\begin{aligned} |D_{i_*}^1| + |D_{i_*}^2| &= n - |A_{i_*}| - |B_{i_*}| - (|C_{i_*}^1| + |C_{i_*}^2|) - |V(P)| \\ &\geq n - k\alpha n - 2(\alpha + 2\varepsilon)n - (6\varepsilon + \alpha)n - 4\varepsilon n \\ &= (1 - (k + 3)\alpha - 14\varepsilon)n \geq 6\varepsilon n. \end{aligned}$$

So, there exist $a, b \in \{1, 2\}$ such that $|C_{i_*}^a|, |D_{i_*}^b| \geq 3\varepsilon n$. By (5), $a \neq b$ and so we take $j_* := a$. This proves the claim. ■

Claim 4. *For each component C of $G[C_{i_*}^{j*}]$ and each component D of $G[D_{i_*}^{3-j*}]$, (C, D) is a pure pair of G .*

Proof of Claim 4. Assume not. By symmetry, we may assume that C has a vertex u having both a neighbor and a non-neighbor in D , because otherwise we swap C and D by reversing the order of P . As D is connected, there exist $v, v' \in V(D)$ such that $uv, vv' \in E(G)$ and $uv' \notin E(G)$.

Note that $m_{i_*}^+(v) \equiv m_{i_*}^+(v') \pmod{2}$ and

(6) *for every neighbor $\ell \in N_G(v) \cap U_{i_*}^+$, the number $\ell - m_{i_*}^+(v)$ is even,*

because otherwise for the minimum $\ell \in N_G(v) \cap U_{i_*}^+$ with odd $\ell - m_{i_*}^+(v)$, a vertex set $\{v, m_{i_*}^-(u), m_{i_*}^-(u) + 1, \dots, \ell, u\}$ induces a strongly k -good generalized fan with v as its center, a contradiction by Proposition 3.4.

If $m_{i_*}^+(v) \leq m_{i_*}^+(v')$, then $\{v, u, m_{i_*}^-(u), m_{i_*}^-(u) + 1, \dots, m_{i_*}^+(v'), v'\}$ induces a strongly k -good generalized fan with v as a center by (6).

If $m_{i_*}^+(v) > m_{i_*}^+(v')$, then simply $(u, m_{i_*}^-(u), m_{i_*}^-(u) + 1, \dots, m_{i_*}^+(v'), v', v, u)$ is an induced cycle whose length is at least k and is of the same parity with k . Hence Proposition 3.1 implies a contradiction. \blacksquare

By Claim 4, there exists $S \in \{C_{i_*}^{j*}, D_{i_*}^{3-j*}\}$ such that every component of $G[S]$ has less than εn vertices. By Claim 3, we can greedily find a set of components of $G[S]$ covering at least εn vertices and at most $2\varepsilon n$ vertices. Since $|S| \geq 3\varepsilon n$, the vertices of S covered by this set of components with the vertices of S not covered by this set of components give a pure pair (A, B) with $|A|, |B| \geq \varepsilon n$, a contradiction. This proves the lemma. \square

5. DISCUSSIONS

For a graph G , we write $\chi(G)$ to denote its chromatic number and $\omega(G)$ to denote its clique number, that is the maximum size of a clique. A class \mathcal{G} of graphs is called χ -bounded if there exists a function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that for every induced subgraph H of a graph in \mathcal{G} , $\chi(H) \leq f(\omega(H))$. In addition, we say \mathcal{G} is *polynomially χ -bounded* if f can be taken as a polynomial.

Every polynomially χ -bounded class of graphs has the strong Erdős-Hajnal property, but the converse does not hold; see the survey paper by Scott and Seymour [22]. So it is natural to ask whether the class of graphs with no pivot-minor isomorphic to C_k is polynomially χ -bounded, which is still open. So far Choi, Kwon, and Oum [4] showed that it is χ -bounded.

Theorem 5.1 (Choi, Kwon, and Oum [4, Theorem 4.1]). *For each $k \geq 3$, the class of graphs with no pivot-minor isomorphic to C_k is χ -bounded.*

They showed that $\chi(G) \leq 2(6k^3 - 26k^2 + 25k - 1)^{\omega(G)-1}$ holds for graphs G having no pivot-minor isomorphic to C_k , far from being a polynomial. Theorem 5.1 is now implied by a recent theorem of Scott and Seymour [21], solving three conjectures of Gyárfás [13] on χ -boundedness all at once.

Theorem 5.2 (Scott and Seymour [21]). *For all $k \geq 0$ and $\ell > 0$, the class of all graphs having no induced cycle of length k modulo ℓ is χ -bounded.*

To see why Theorem 5.2 implies Theorem 5.1, take $\ell := 2\lceil k/2 \rceil$ and apply Proposition 3.1. Still the bound obtained from Theorem 5.2 is far from being a polynomial.

And yet no one was able to answer the following problem of Esperet.

Problem 5.1 (Esperet; see [15]). *Is it true that every χ -bounded class of graphs polynomially χ -bounded?*

Thus it is natural to pose the following conjecture.

Conjecture 5.2. *For every graph H , the class of graphs with no pivot-minor isomorphic to H is polynomially χ -bounded.*

It is open whether Conjecture 5.2 holds when $H = C_k$. Conjecture 5.2 implies not only Conjectures 1.2, 1.3 but also the following conjecture of Geelen (see [8]) proposed in 2009 at the DIMACS workshop on graph colouring and structure held at Princeton University.

Conjecture 5.3 (Geelen; see [8]). *For every graph H , the class of graphs with no vertex-minor isomorphic to H is χ -bounded.*

Of course it is natural to pose the following conjecture, weaker than Conjecture 5.2 but stronger than Conjecture 5.3.

Conjecture 5.4 (Kim, Kwon, Oum, and Sivaraman [16]). *For every graph H , the class of graphs with no vertex-minor isomorphic to H is polynomially χ -bounded.*

For vertex-minors, more results are known. Kim, Kwon, Oum, and Sivaraman [16] proved that for each $k \geq 3$, the class of graphs with no vertex-minor isomorphic to C_k is polynomially χ -bounded. Their theorem is now implied by the following two recent theorems. To describe these theorems, we first have to introduce a few terms. A *circle graph* is the intersection graph of chords in a circle. In particular, C_k is a circle graph. The *rank-width* of a graph is one of the width parameters of graphs, measuring how easy it is to decompose a graph into a tree-like structure while keeping every cut to have a small ‘rank’. Rank-width was introduced by Oum and Seymour [19]. We will omit the definition of the rank-width.

Theorem 5.3 (Geelen, Kwon, McCarty, and Wollan [12]). *For each circle graph H , there is an integer $r(H)$ such that every graph with no vertex-minor isomorphic to H has rank-width at most $r(H)$.*

Theorem 5.4 (Bonamy and Pilipczuk [3]). *For each k , the class of graphs of rank-width at most k is polynomially χ -bounded.*

As noted in [6], it is easy to prove directly that the class of graphs of bounded rank-width has the strong Erdős-Hajnal property, without using Theorem 5.4. See Figure 4 for a diagram showing the containment relations between these properties.

So, one may wonder whether the class of graphs with no pivot-minor isomorphic to C_k has bounded rank-width. Unfortunately, if k is odd, then it is not true, because all bipartite graphs have no pivot-minor isomorphic to C_k for odd k and yet have unbounded rank-width, see [17]. If k is even, then it would be true if the following conjecture hold.

Conjecture 5.5 (Oum [18]). *For every bipartite circle graph H , there is an integer $r(H)$ such that every graph with no pivot-minor isomorphic to H has rank-width at most $r(H)$.*

Note. Chudnovsky, Scott, Seymour, and Spirkl [7] proved that for every graph H , the class of graphs G such that neither G nor \overline{G} has any subdivision of H as an induced subgraph has the strong Erdős-Hajnal property. This implies that when k is even, the class of graphs with no induced even hole of length at least k and no induced even anti-hole of length at least k has the strong Erdős-Hajnal property. This is because every subdivision of a large theta




FIGURE 4. Known implications between properties of classes of graphs.

graph² contains a large even hole. This implies Theorem 1.2 for even k but not for odd k by Propositions 3.1 and 3.2. The authors would like to thank the authors of [7] to share this observation.

Acknowledgement. The authors would like to thank anonymous reviewers for their careful reading and helpful suggestions.

REFERENCES

- [1] Noga Alon, János Pach, Rom Pinchasi, Radoš Radoičić, and Micha Sharir, *Crossing patterns of semi-algebraic sets*, J. Combin. Theory Ser. A **111** (2005), no. 2, 310–326. MR 2156215
- [2] Marthe Bonamy, Nicolas Bousquet, and Stéphan Thomassé, *The Erdős-Hajnal conjecture for long holes and antiholes*, SIAM J. Discrete Math. **30** (2016), no. 2, 1159–1164. MR 3507546
- [3] Marthe Bonamy and Michał Pilipczuk, *Graphs of bounded cliquewidth are polynomially χ -bounded*, Adv. Comb. **Paper No. 8, 21 pp.** (2020). MR 4125349
- [4] Ilkyoo Choi, O-jeong Kwon, and Sang-il Oum, *Coloring graphs without fan vertex-minors and graphs without cycle pivot-minors*, J. Combin. Theory Ser. B **123** (2017), 126–147. MR 3597097
- [5] Maria Chudnovsky, *The Erdős-Hajnal conjecture—a survey*, J. Graph Theory **75** (2014), no. 2, 178–190. MR 3150572
- [6] Maria Chudnovsky and Sang-il Oum, *Vertex-minors and the Erdős-Hajnal conjecture*, Discrete Math. **341** (2018), no. 12, 3498–3499. MR 3862646
- [7] Maria Chudnovsky, Alex Scott, Paul Seymour, and Sophie Spirkl, *Pure pairs. II. Excluding all subdivisions of a graph*, arXiv:1804.01060, 2020.
- [8] Zdeněk Dvořák and Daniel Král, *Classes of graphs with small rank decompositions are χ -bounded*, European J. Combin. **33** (2012), no. 4, 679–683. MR 3350076
- [9] Paul Erdős and András Hajnal, *Ramsey-type theorems*, Discrete Appl. Math. **25** (1989), no. 1-2, 37–52, Combinatorics and complexity (Chicago, IL, 1987). MR 1031262
- [10] Jacob Fox and János Pach, *Erdős-Hajnal-type results on intersection patterns of geometric objects*, Horizons of combinatorics, Bolyai Soc. Math. Stud., vol. 17, Springer, Berlin, 2008, pp. 79–103. MR 2432528

²A *theta graph* is a graph consisting of three internally disjoint paths of length at least 1 joining two fixed vertices.

- [11] Jacob Fox and Benny Sudakov, *Induced Ramsey-type theorems*, Adv. Math. **219** (2008), no. 6, 1771–1800. MR 2455625
- [12] Jim Geelen, O-joung Kwon, Rose McCarty, and Paul Wollan, *The grid theorem for vertex-minors*, J. Combin. Theory Ser. B (2020).
- [13] A. Gyárfás, *Problems from the world surrounding perfect graphs*, Proceedings of the International Conference on Combinatorial Analysis and its Applications (Pokrzywna, 1985), vol. 19, 1987, pp. 413–441 (1988). MR 951359
- [14] Marcin Kamiński, Vadim V. Lozin, and Martin Milanič, *Recent developments on graphs of bounded clique-width*, Discrete Appl. Math. **157** (2009), no. 12, 2747–2761. MR 2536473
- [15] T. Karthick and Frédéric Maffray, *Vizing bound for the chromatic number on some graph classes*, Graphs Combin. **32** (2016), no. 4, 1447–1460. MR 3514976
- [16] Ringi Kim, O-joung Kwon, Sang-il Oum, and Vaidy Sivaraman, *Classes of graphs with no long cycle as a vertex-minor are polynomially χ -bounded*, J. Combin. Theory Ser. B **140** (2020), 372–386. MR 4033105
- [17] Sang-il Oum, *Rank-width and vertex-minors*, J. Combin. Theory Ser. B **95** (2005), no. 1, 79–100. MR 2156341
- [18] ———, *Excluding a bipartite circle graph from line graphs*, J. Graph Theory **60** (2009), no. 3, 183–203. MR 2478228
- [19] Sang-il Oum and Paul Seymour, *Approximating clique-width and branch-width*, J. Combin. Theory Ser. B **96** (2006), no. 4, 514–528. MR 2232389
- [20] Vojtěch Rödl, *On universality of graphs with uniformly distributed edges*, Discrete Math. **59** (1986), no. 1-2, 125–134. MR 837962
- [21] Alex Scott and Paul Seymour, *Induced subgraphs of graphs with large chromatic number. X. Holes of specific residue*, Combinatorica **39** (2019), no. 5, 1105–1132. MR 4039603
- [22] ———, *A survey of χ -boundedness*, J. Graph Theory **95** (2020), no. 3, 473–504. MR 4174126

(J. Kim) DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON, SOUTH KOREA
Email address: jaehoon.kim@kaist.ac.kr

(S. Oum) DISCRETE MATHEMATICS GROUP, INSTITUTE FOR BASIC SCIENCE (IBS), DAEJEON, SOUTH KOREA

(S. Oum) DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON, SOUTH KOREA
Email address: sangil@ibs.re.kr