# SYMMETRIC POLYNOMIALS IN LEIBNIZ ALGEBRAS AND THEIR INNER AUTOMORPHISMS

## ŞEHMUS FINDIK AND ZEYNEP ÖZKURT

ABSTRACT. Let  $L_n$  be the free metabelian Leibniz algebra generated by the set  $X_n = \{x_1, \ldots, x_n\}$  over a field K of characteristic zero. This is the free algebra of rank n in the variety of solvable of class 2 Leibniz algebras. We call an element  $s(X_n) \in L_n$  symmetric if  $s(x_{\sigma(1)}, \ldots, x_{\sigma(n)}) = s(x_1, \ldots, x_n)$  for each permutation  $\sigma$  of  $\{1, \ldots, n\}$ . The set  $L_n^{S_n}$  of symmetric polynomials of  $L_n$  is the algebra of invariants of the symmetric group  $S_n$ . Let  $K[X_n]$  be the usual polynomial algebra with indeterminates from  $X_n$ . The description of the algebra  $K[X_n]^{S_n}$  is well known, and the algebra  $(L'_n)^{S_n}$  in the commutator ideal  $L'_n$  is a right  $K[X_n]^{S_n}$ -module. We give explicit forms of elements of the  $K[X_n]^{S_n}$ -module  $(L'_n)^{S_n}$ . Additionally, we determine the description of the group  $Inn(L_n^{S_n})$  of inner automorphisms of the algebra  $L_n^{S_n}$ . The findings can be considered as a generalization of the recent results obtained for the free metabelian Lie algebra of rank n.

### Introduction

Hilbert's fourteen problem is one of those famous twenty three problems suggested by German mathematician David Hilbert [H] in 1900 at the Paris conference of the International Congress of Mathematicians, and it is related with the finite generation of the algebra  $K[X_n]^G$  of invariants of  $G < GL_n(K)$ , where  $K[X_n] = K[x_1, \ldots, x_n]$  is the usual polynomial algebra over a field K, and  $GL_n(K)$  is the general linear group. Nagata [Na] showed that the problem is not true in general in 1959. Earlier in 1916, Noether [N] solved the problem in affirmative for finite groups. In particular let  $G = S_n$  be the symmetric group acting on the algebra  $K[X_n]$  by permuting the variables:  $\pi \cdot p(x_1, \ldots, x_n) = p(x_{\pi(1)}, \ldots, x_{\pi(n)}), p \in K[X_n], \pi \in S_n$ . The algebra  $K[X_n]^{S_n}$  is generated by the set  $\{\sum_{i=1}^n x_i^k \mid k=1,\ldots,n\}$ , by the fundamental theorem of symmetric polynomials. Elementary symmetric polynomials  $e_j = \sum x_{i_1} \cdots x_{i_j}, i_1 < \cdots < i_j, j=1,\ldots,n$ , form another generating set.

A noncommutative analogue of the problem is the algebra  $K\langle X_n\rangle^{S_n}$  of symmetric polynomials in the free associative algebra  $K\langle X_n\rangle$ . One may see the works [Wo, BRRZ, GKL] on the algebra  $K\langle X_n\rangle^{S_n}$ . Another analogue is working in relatively free Lie algebras, which are not associative and commutative. The algebras  $F_n^{S_n}$ , and  $M_n^{S_n}$  are not finitely generated via [Br] and [Dr], where  $F_n$  and  $M_n$  are the free Lie algebra and the free metabelian Lie algebra of rank n, respectively. One may see the papers [FO, DFO] for the explicit elements of the algebra  $M_n^{S_n}$ . See also [FO2] for the inner automorphisms of  $M_n^{S_n}$ .

<sup>2010</sup> Mathematics Subject Classification. 17A32; 17A36; 17A50; 17B01; 17B30.

Key words and phrases. Leibniz algebras; metabelian identity, automorphisms, symmetric polynomials.

We consider the Leibniz algebras which can be thought as a generalization of the Lie algebras. Leibniz algebras are defined by the identity [x, [y, z]] = [[x, y], z] - [[x, z], y], where the bracket is bilinear; however, non-necessarily skew-symmetric. In the case of skew-symmetry the identity turns into the Jacobi identity, and we obtain a Lie algebra. Leibniz algebras are related with many branches of mathematics. See the papers [L, LP, MU, AO, O] for more details.

In the present study, we consider the free metabelian Leibniz algebra  $L_n$  and we determine the algebra  $L_n^{S_n}$  of symmetric polynomials. Additionally, we describe the group  $\operatorname{Inn}(L_n^{S_n})$  of inner automorphisms of  $L_n^{S_n}$ .

#### 1. Preliminaries

Let K be a field of characteristic zero. A Leibniz algebra L over K is a vector space furnished with bilinear commutator [.,.] satisfying the Leibniz identity

$$[[x, y], z] = [[x, z], y] + [x, [y, z]],$$

or

$$[x, y]r_z = [xr_z, y] + [x, yr_z],$$

 $x,y,z\in L$ . Here  $r_z$  stands for the adjoint operator  $\operatorname{ad} z$  acting from right side by commutator multiplication. The Leibniz algebra L is nonassociative and noncommutative. Leibniz algebras are generalizations of Lie algebras. There is a natural way to obtain a Lie algebra from a given Leibniz algebra L. Consider the ideal  $\operatorname{Ann}(L)$ , called the annihilator of L, generated by all elements  $[x,x],\ x\in L$ . It is well known, see e.g.  $[\operatorname{LP}]$ , that  $r_a=0$  if and only if  $a\in\operatorname{Ann}(L)$ . Clearly the elements of the form  $[x,y]+[y,x]\in\operatorname{Ann}(L)$  by the equality

$$[x, y] + [y, x] = [x + y, x + y] - [x, x] - [y, y], x, y \in L.$$

Hence the factor algebra L/Ann(L) turns into a Lie algebra, since the anticommutativity [x, y] + [y, x] = 0 and the Jacobi identity

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = 0,$$

are satisfied in  $L/\mathrm{Ann}(L)$ .

Now consider the free algebra  $L_n$  of rank n generated by  $X_n = \{x_1, \ldots, x_n\}$  in the variety of metabelian Leibniz algebras over the base field K. The algebra  $L_n$  satisfies the metabelian identity [[x,y],[z,t]]=0, and is a solvable of class 2 Leibniz algebra. Hence every element in the commutator ideal  $L'_n = [L_n, L_n]$  of the free metabelian Leibniz algebra  $L_n$  can be expressed as a linear combination of left-normed monomials of the form

$$[[\cdots [[x_{i_1}, x_{i_2}], x_{i_3}], \ldots], x_{i_k}] = [x_{i_1}, x_{i_2}, x_{i_3}, \ldots, x_{i_k}]$$
$$= [x_{i_1}, x_{i_2}]r_{i_3} \cdots r_{i_k} = [x_{i_1}, x_{i_2}]r_{i_{\pi(3)}} \cdots r_{i_{\pi(k)}}$$

where  $\pi$  is a permutation of the set  $\{3, \ldots, k\}$ . In this way the commutator ideal  $L'_n$  can be considered as a right  $K[R_n] = K[r_1, \ldots, r_n]$ -module, where  $r_i = r_{x_i} = \operatorname{ad} x_i$ ,  $i = 1, \ldots, n$ . It is well known, see e.g. Proposition 3.1. of the paper [DP], that the elements

$$x_{i_1}, [x_{i_1}, x_{i_2}], [x_{i_1}, x_{i_2}], [x_{i_3}, \dots, x_{i_k}], 1 \le i_1, i_2 \le n, 1 \le i_3 \le \dots \le i_k,$$

form a basis for  $L_n$ . The next result is a direct consequence of this basis.

**Corollary 1.1.** The commutator ideal  $L'_n$  of the free metabelian Leibniz algebra  $L_n$  is a free right  $K[R_n]$ -module with generators  $[x_i, x_j]$ ,  $1 \le i, j \le n$ .

A polynomial  $s = s(x_1, \dots, x_n)$  in the free metabelian Leibniz algebra  $L_n$  is said to be *symmetric* if

$$\sigma s = s(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = s(x_1, \dots, x_n), \quad \sigma \in S_n.$$

The set  $L_n^{S_n}$  of symmetric polynomials forms a Leibniz subalgebra, which is the algebra of invariants of the symmetric group  $S_n$ . The  $K[R_n]$ -module structure of the commutator ideal  $L'_n$  implies that the algebra  $(L'_n)^{S_n}$  is a right  $K[R_n]^{S_n}$ -module. One of the set of generators of the algebra  $K[R_n]^{S_n}$  of symmetric polynomials is well known:  $\{r_1^k + \cdots + r_n^k \mid 1 \leq k \leq n\}$ , see e.g. [S].

#### 2. Main Results

2.1. Symmetric polynomials. In this section we determine the algebra  $L_n^{S_n}$  of symmetric polynomials in the free metabelian Leibniz algebra  $L_n$ . Clearly the linear symmetric polynomials are included in the K-vector space spanned on a single element  $x_1 + \cdots + x_n$ . Hence it is sufficient to work in the commutator ideal  $L'_n$  of  $L_n$ , and describe the algebra  $(L'_n)^{S_n}$ . Let us fix the notations  $a_i = [x_i, x_i]$ ,  $1 \le i \le n$ , and  $b_{jk} = [x_j, x_k]$ ,  $1 \le j \ne k \le n$ , which are the free generators of  $K[R_n]$ -module L'. We provide explicit elements of  $K[R_n]^{S_n}$ -module  $(L'_n)^{S_n}$ . For this purpose, we study in the  $K[R_n]$ -submodules

$$A_n = \left\{ \sum_{i=1}^n a_i p_i \mid p_i \in K[R_n] \right\} \text{ and } B_n = \left\{ \sum_{1 \le j \ne k \le n} b_{jk} q_{jk} \mid q_{jk} \in K[R_n] \right\}$$

of the  $K[R_n]$ -module  $L'_n = A_n \oplus B_n$ , generated by  $a_i$ ,  $1 \le i \le n$ , and  $b_{jk}$ , respectively,  $1 \le j \ne k \le n$ , due to the fact that they are invariant under the action of  $S_n$ ; i.e.,  $A_n^{S_n} \subset A_n$ , and  $B_n^{S_n} \subset B_n$ .

Let us denote the subgroups  $\Pi_i = \{\pi \in S_n \mid \pi(i) = i\}, 1 \leq i \leq n$ , and  $\Pi_{jk} = \{\pi \in S_n \mid \pi(j) = j, \pi(k) = k\}, 1 \leq j \neq k \leq n$ , of  $S_n$ . In the next theorems, we determine symmetric polynomials in the  $K[R_n]$ -modules  $A_n$ , and  $B_n$ , respectively.

**Theorem 2.1.** Let  $p = \sum_{i=1}^{n} a_i p_i$  be a polynomial in  $A_n$ , for some  $p_i \in K[R_n]$ ,  $1 \le i \le n$ . Then p is symmetric if and only if

$$p_1(r_1, r_2, \dots, r_n) = \pi p_1(r_1, r_2, \dots, r_n) = p_1(r_1, r_{\pi(2)}, \dots, r_{\pi(n)}), \quad \pi \in \Pi_1,$$

 $\sigma p_i = p_i, \ \sigma \in \Pi_i, \ and \ p_i = (1i)p_1 = p_1(r_i, r_2, \dots, r_{i-1}, r_1, r_{i+1}, \dots, r_n), \ for \ transpositions \ (1i) \in S_n, \ i = 2, \dots, n.$ 

*Proof.* Let  $p \in A_n$  be an element of the form

$$p = \sum_{i=1}^{n} a_i p_i(r_1, \dots, r_n), \quad p_i \in K[R_n].$$

If p is a symmetric polynomial, then  $p = \pi p$ ; i.e.,

$$\sum_{i=1}^{n} a_i p_i(r_1, \dots, r_n) = \sum_{i=1}^{n} a_{\pi(i)} p_i(r_{\pi(1)}, \dots, r_{\pi(n)})$$

for each  $\pi \in S_n$  by definition, and by Corollary 1.1 we may compare the coefficients of  $a_i$ , i = 1, ..., n, from  $K[R_n]$ , in the last equality. In particular,  $p = \pi p$  for each  $\pi \in \Pi_i$ , and comparing the coefficients of  $a_i$ , i = 1, ..., n, we obtain that

$$p_i(r_1,\ldots,r_n)=p_i(r_{\pi(1)},\ldots,r_{\pi(i-1)},r_i,r_{\pi(i+1)},\ldots,r_{\pi(n)}).$$

Now consider (1i)p = p for every transposition  $(1i) \in S_n$ , i = 2, ..., n. Then these equalities give that  $p_i = (1i)p_1$ , and thus

$$p_i(r_1, \dots, r_n) = (1i)p_1(r_1, \dots, r_n)$$
  
=  $p_1(r_i, r_2, \dots, r_{i-1}, r_1, r_{i+1}, \dots, r_n)$ 

Conversely consider the element  $p = \sum_{i=1}^{n} a_i p_i$  satisfying the conditions in the theorem. It is sufficient to show that (1k)p = p, k = 2, ..., n, since these transpositions generate the symmetric group  $S_n$ . Note that if  $i \neq 1, k$ , then  $(1k)p_i = p_i$ , since  $(1k) \in \Pi_i$ . The following computations complete the proof.

$$(1k)p = (1k) \left( a_1 p_1 + a_k p_k + \sum_{i \neq 1, k} a_i p_i \right)$$

$$= a_k ((1k)p_1) + a_1 ((1k)p_k) + \sum_{i \neq 1, k} a_i ((1k)p_i)$$

$$= a_k p_k + a_1 p_1 + \sum_{i \neq 1, k} a_i p_i = p.$$

**Theorem 2.2.** Let  $q = \sum b_{ij}q_{ij}$  be a polynomial in  $B_n$ , for some  $q_{ij} \in K[R_n]$ ,  $1 \le i \ne j \le n$ . Then q is symmetric if and only if  $q_{ij} = \sigma q_{kl}$  for every  $\sigma : i \to k, j \to l$ , in particular,

$$q_{1i} = (2i)q_{12}, \quad q_{i2} = (1i)q_{12}, \quad q_{2i} = (1i)q_{21}, \quad q_{i1} = (2i)q_{21},$$
  
 $q_{21} = (12)q_{12}, \quad q_{ij} = (1i)(2j)q_{12}, \quad 3 \le i \ne j \le n.$ 

and  $q_{ij} = \pi q_{ij}$ , for all  $\pi \in \Pi_{ij}$ .

*Proof.* Assume that a polynomial  $q = \sum b_{ij}q_{ij} \in B_n$ ,  $1 \le i \ne j \le n$ , is symmetric. Then  $\pi q = q$  for each  $\pi \in \Pi_{12}$  gives that

$$q_{12}(r_1, r_2, \dots, r_n) = q_{12}(r_1, r_2, r_{\pi(3)}, \dots, r_{\pi(n)}).$$

Relations on  $q_{ij}$ 's in the theorem are straightforward, by making use of Corollary 1.1, and comparing the coefficients of  $b_{12}$ ,  $b_{21}$ , and  $b_{ij}$  from the equalities q = (12)q = (1i)q = (2i)q = (1i)(2j)q = (ij)q, where  $3 \le i \ne j \le n$ .

Now let the polynomial  $q \in B_n$  satisfy the conditions of the theorem, and  $(1k) \in S_n$  be a transposition for a fixed  $k \in \{3, \ldots, n\}$ . We have to show that (1k)q = q. Let express q in the following form.

$$q = b_{1k}q_{1k} + b_{k1}q_{k1} + \sum_{i,j\neq 1,k} b_{ij}q_{ij} + \sum_{i\neq 1,k} (b_{1i}q_{1i} + b_{ki}q_{ki} + b_{ik}q_{ik} + b_{i1}q_{i1}).$$

Note that  $(1k) \in \Pi_{ij}$ , and hence  $(1k)q_{ij} = q_{ij}$ , for  $i, j \neq 1, k$ . Then we have that

$$\begin{split} (1k)q = &b_{k1}((1k)q_{1k}) + b_{1k}((1k)q_{k1}) + \sum_{i,j \neq 1,k} b_{ij}((1k)q_{ij}) \\ &+ \sum_{i \neq 1,k} (b_{ki}((1k)q_{1i}) + b_{1i}((1k)q_{ki}) + b_{i1}((1k)q_{ik}) + b_{ik}((1k)q_{i1})) \\ = &b_{k1}q_{k1} + b_{1k}q_{1k} + \sum_{i,j \neq 1,k} b_{ij}q_{ij} + \sum_{i \neq 1,k} (b_{ki}q_{ki} + b_{1i}q_{1i} + b_{i1}q_{i1} + b_{ik}q_{ik}) = q. \end{split}$$

We obtain the next corollary by combining Theorem 2.1 and Theorem 2.2.

Corollary 2.3. If s is a symmetric polynomial in the free metabelian Leibniz algebra  $L_n$ , then it is of the form

$$s = \sum_{1 \le i \le n} \alpha x_i + \sum_{1 \le i \le n} [x_i, x_i]((1i)f) + \sum_{3 \le i \ne j \le n} [x_i, x_j]((1i)(2j)g)$$

$$+ [x_1, x_2]g + \sum_{3 \le i \le n} ([x_1, x_i]((2i)g) + [x_i, x_2]((1i)g)$$

$$+ [x_2, x_1]h + \sum_{3 \le i \le n} ([x_i, x_1]((2i)h) + [x_2, x_i]((1i)h)),$$

where  $\alpha \in K$ ,  $f, g, h \in K[R_n]$ , such that  $\pi f = f$  for  $\pi \in \Pi_1$ ,  $\sigma g = g$  for  $\sigma \in \Pi_{12}$ , and h = (12)g.

**Example 2.4.** Let n=2 and the free metabelian Leibniz algebra  $L_2$  be generated by  $x_1, x_2$ . Then each symmetric polynomial  $s \in L_2^{S_2}$  is of the form

$$s = \alpha(x_1 + x_2) + [x_1, x_1]f(r_1, r_2) + [x_2, x_2]f(r_2, r_1) + [x_1, x_2]g(r_1, r_2) + [x_2, x_1]g(r_2, r_1),$$

where  $\alpha \in K$ ,  $f, g \in K[R_2]$ . Note that the Lie correspondence of this result (modulo the annihilator) is that if  $s(x_1, x_2)$  is a symmetric polynomial in the free metabelian Lie algebra generated by  $x_1, x_2$ , then

$$s = \alpha(x_1 + x_2) + [x_1, x_2]t(r_1, r_2),$$

such that  $t(r_1, r_2) = -t(r_2, r_1)$ , which is compatible with the recent result given in [FO].

2.2. Inner automorphisms. Let u be an element in the commutator ideal  $L'_n$  of the free metabelian Leibniz algebra  $L_n$ . The adjoint operator

$$adu: L_n \to L_n, \quad adu(v) = [v, u], \quad v \in L_n$$

is nilpotent since  $\operatorname{ad}^2 u = 0$ , and that  $\psi_u = \exp(\operatorname{ad} u) = 1 + \operatorname{ad} u$  is called an inner automorphism of  $L_n$  with inverse  $\psi_{-u}$ . Clearly the group  $\operatorname{Inn}(L_n)$  consisting of all inner automorphisms is abelian due to the fact that  $\psi_{u_1}\psi_{u_2} = \psi_{u_1+u_2}$ .

In the next theorem we determine the group  $Inn(L_n^{S_n})$  of inner automorphisms preserving symmetric polynomials.

**Theorem 2.5.** 
$$\operatorname{Inn}(L_n^{S_n}) = \{ \psi_{u_1 + u_2} \mid u_1 \in \operatorname{Ann}(L_n), u_2 \in (L'_n)^{S_n} \}.$$

*Proof.* Let  $v \in L_n^{S_n}$ ,  $u = u_1 + u_2$ , for some  $u_1 \in \text{Ann}(L_n)$ , and  $u_2 \in (L'_n)^{S_n}$ . Then clearly

$$\psi_u(v) = v + [v, u_1 + u_2] = v + [v, u_2] \in L_n^{S_n}.$$

Conversely, let  $\psi_u(v) \in L_n^{S_n}$  for  $v \in L_n^{S_n}$ , and  $u \in L_n'$ . The action of  $\psi_u$  is identical when  $v \in L_n'$ . Hence we assume that the linear counterpart  $v_l = \alpha(x_1 + \dots + x_n)$ ,  $\alpha \in K$ , of v is nonzero. We may express  $u = u_1 + u_2$ ,  $u_1 \in \text{Ann}(L_n)$ ,  $u_2 \in L_n'$ , where  $u_2 \notin \text{Ann}(L_n)$ . Hence we have  $\psi_u(v) \in L_n^{S_n}$ , which implies that  $[v_l, u_2]$  is a symmetric polynomial. Let  $\pi \in S_n$  be an arbitrary permutation. Then

$$[v_l, u_2] = \pi[v_l, u_2] = [\pi v_l, \pi u_2] = [v_l, \pi u_2]$$

or 
$$[x_1 + \cdots + x_n, u_2 - \pi u_2] = 0$$
, and thus  $u_2 - \pi u_2 = 0$ . Therefore  $u_2 \in (L'_n)^{S_n}$ .  $\square$ 

We complete the paper by releasing the following problem.

**Problem 2.6.** Determine the group  $Aut(L_n^{S_n})$  of all automorphisms preserving the symmetric polynomials.

#### References

- [AO] T.T. Adıyaman, Z. Özkurt, Automorphisms of free metabelian Leibniz algebras of rank three, Turkish Journal of Mathematics, 43 (2019) 5, 2262-2274.
- [BRRZ] N. Bergeron, C. Reutenauer, M. Rosas, M. Zabrocki, Invariants and coinvariants of the symmetric groups in noncommuting variables, Canad. J. Math. 60 (2008) 2, 266-296.
- [Br] R.M. Bryant, On the fixed points of a finite group acting on a free Lie algebra, J. London Math. Soc. (2) 43 (1991), 2, 215-224.
- [Dr] V. Drensky, Fixed algebras of residually nilpotent Lie algebras, Proc. Amer. Math. Soc. 120 (1994) 4, 1021-1028.
- [DFO] V. Drensky, Ş. Fındık, N.Ş. Öğüşlü, Symmetric polynomials in the free metabelian Lie algebras, (2019), arXiv:1912.01066, to appear in Med. J. Math.
- [DP] V. Drensky, G.M. Piacentini Cattanneo, Varieties of metabelian Leibniz algebras, J. Alg. Appl., 1 (2002) 1, 31-50.
- [FO] Ş. Fındık, N.Ş. Öğüşlü, Palindromes in the free metabelian Lie algebras, Internat. J. Algebra Comput., 29 (2019) 5, 885-891.
- [FO2] Ş. Fındık, N.Ş. Öğüşlü, Inner automorphisms of Lie algebras of symmetric polynomials, (2019), arXiv:2003.06818.
- [GKL] I.M. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V.S. Retakh, J.-Y. Thibon, Noncommutative symmetric functions, Adv. Math., 112 (1995) 2, 218-348.
- [H] D. Hilbert, Mathematische Probleme, Göttinger Nachrichten (1900), 253-297; Archiv der Mathematik und Physik, 3 (1901) 1, 44-63; Translation: Bull. Amer. Math. Soc., 8 (1902), No. 10, 437-479.
- [L] J.-L. Loday, Une version non commutative des algebres de Lie: les algebres de Leibniz, Enseign. Math., 39 (1993), 269-293.
- [LP] J.-L. Loday, T. Pirashvili, Universal enveloping algebras of Leibniz algebras and (co)homology, Math. Ann., 296 (1993), 139-158.
- [MU] A.A. Mikhalev, U.U. Umirbaev, Subalgebras of free Leibniz algebras, Comm. Algebra, 26 (1998), 435-446.
- [N] E. Noether, Der Endlichkeitssatz der Invarianten endlicher Gruppen, Math. Ann., 77 (1916), 89-92.
- [Na] M. Nagata, On the 14-th problem of Hilbert, Amer. J. Math., 81 (1959), 766-772.
- [O] Z. Özkurt, Orbits and test elements in free Leibniz algebras of rank two, Commun. Alg., 43 (2015) 8, pp.3534-3544.
- [S] B. Sturmfels, Algorithms in Invariant Theory, 2nd ed. Texts and Monographs in Symbolic Computation, Springer-Verlag, Wien, 2008.
- [Wo] M.C. Wolf, Symmetric functions of non-commutative elements, Duke Math. J. 2 (1936), No. 4, 626-637.

Department of Mathematics, Çukurova University, 01330 Balcali, Adana, Turkey

E-mail address: sfindik@cu.edu.tr E-mail address: zyapti@cu.edu.tr