FREDHOLM CONDITIONS AND INDEX FOR RESTRICTIONS OF INVARIANT PSEUDODIFFERENTIAL OPERATORS TO ISOTYPICAL COMPONENTS

ALEXANDRE BALDARE, RÉMI CÔME, MATTHIAS LESCH, AND VICTOR NISTOR

ABSTRACT. Let Γ be a compact group acting on a smooth, compact manifold M, let $P \in \psi^m(M; E_0, E_1)$ be a Γ -invariant, classical pseudodifferential operato acting between sections of two equivariant vector bundles $E_i \to M$, i = 0, 1,and let α be an irreducible representation of the group $\Gamma.$ Then P induces a map $\pi_{\alpha}(P): H^{s}(M; E_{0})_{\alpha} \to H^{s-m}(M; E_{1})_{\alpha}$ between the α -isotypical components of the corresponding Sobolev spaces of sections. When Γ is finite, we explicitly characterize the operators P for which the map $\pi_{\alpha}(P)$ is Fredholm in terms of the principal symbol of P and the action of Γ on the vector bundles E_i . When $\Gamma = \{1\}$, that is, when there is no group, our result extends the classical characterization of Fredholm (pseudo)differential operators on compact manifolds. The proof is based on a careful study of the symbol C^* -algebra and of the topology of its primitive ideal spectrum. We also obtain several results on the structure of the norm closure of the algebra of invariant pseudodifferential operators and their relation to induced representations. Whenever our results also hold for non-discrete groups, we prove them in this greater generality. As an illustration of the generality of our results, we provide some applications to Hodge theory and to index theory of singular quotient spaces.

Contents

1. Introduction	2
1.1. The setting and general notation	2
1.2. The α -principal symbol and α -elliptic	eity 3
1.3. Statement of the main result	4
1.4. Contents of the paper	5
2. Preliminaries	6
2.1. Group representations	6
2.2. Induction and Frobenius reciprocity	7
2.3. The primitive ideal spectrum of a C^*	-algebra 9
2.4. Group actions on manifolds	10
2.5. Pseudodifferential operators	11
3. The principal symbol	11
3.1. The primitive ideal spectrum of A_M^{Γ}	12
3.2. The restriction morphisms	13
3.3. Local calculations	14

 $Key\ words:$ Fredholm operator, C^* -algebra, primitive ideal, pseudodifferential operator, group actions, induced representations. $AMS\ Subject\ classification:\ 47A53,\ 58J40,\ 46L60,\ 47L80,\ 46N20.$ A.B., R.C., and V.N. have been partially supported by ANR-14-CE25-0012-01 (SINGSTAR). V.N. was also supported by the NSF grant DMS 1839515. M.L. was partially supported by the Hausdorff Center for Mathematics, Bonn.

1

3.4. Calculations for the principal orbit bundle	15
3.5. The non-principal orbit case	19
4. Applications and extensions	22
4.1. Reduction to the connected case and α -ellipticity	22
4.2. Fredholm conditions and Hodge and index theory	26
4.3. Special cases	27
4.4. Simonenko's localization principle	29
4.5. Compact (non-finite) groups	30
References	32

1. Introduction

Fredholm operators have been extensively studied and appear in many questions in Mathematical Physics, in Partial Differential Equations (linear and non-linear), in Algebraic and Differential Geometry, in Index Theory, and in other areas. On a compact manifold, a classical pseudodifferential operator is Fredholm between suitable Sobolev spaces if, and only if, it is elliptic. In this paper, we obtain an analogous result for the restrictions to isotypical components of a classical pseudodifferential operator P invariant with respect to the action of a finite group Γ using C^* -algebra methods. Namely, the restriction of P to the isotypical component corresponding to an irreducible representation α of Γ is Fredholm if, and only if, the operator is α -elliptic (Definition 1.1 and Theorem 1.2).

Let us now formulate and explain this result in more detail.

1.1. The setting and general notation. We shall work essentially in the same setting as the one considered in [12], but for a general finite group Γ . Thus, throughout this paper, Γ will be a finite group acting by diffeomorphisms on a smooth Riemannian manifold M. As our main result is only valid for a compact manifold, we assume for this introduction that M is compact. For the main result (Theorem 1.2), we do need Γ to be discrete and finite, so our main result is optimal. Again, see [12]. There is no loss of generality to assume that M is endowed with an invariant Riemannian metric, so we will assume that this is the case also.

As usual, $\widehat{\Gamma}$ denotes the finite set of equivalence classes of irreducible Γ -modules (or representations). Let $T: V_0 \to V_1$ be a Γ -equivariant linear map of Γ -modules and $\alpha \in \widehat{\Gamma}$. Then T induces by restriction a Γ -equivariant linear map

(1)
$$\pi_{\alpha}(T): V_{0\alpha} \to V_{1\alpha}$$

between the α -isotypical components of the Γ -modules V_i , i = 0, 1.

We are mostly interested in this restriction morphism π_{α} in the following case. Let $P \in \psi^m(M; E_0, E_1)$ be a classical, Γ -invariant pseudodifferential operator acting between sections of two Γ -equivariant vector bundles $E_i \to M$, i = 0, 1. Then we obtain the operator

(2)
$$\pi_{\alpha}(P): H^{s}(M; E_{0})_{\alpha} \to H^{s-m}(M; E_{1})_{\alpha},$$

which acts between the α -isotypical components of the corresponding Sobolev spaces of sections. Our main result concerns this operator $\pi_{\alpha}(P)$. For simplicity, we will consider only classical pseudodifferential operators in this article [45, 46, 64, 88, 89, 92].

1.2. The α -principal symbol and α -ellipticity. To put our result into the right perspective, recall that a classical, order m, pseudodifferential operator P is called *elliptic* if its principal symbol

(3)
$$\sigma_m(P) \in \mathcal{C}^{\infty}(T^*M \setminus \{0\}; \operatorname{Hom}(E_0, E_1)),$$

is invertible. Also, recall that a linear operator $T: X_0 \to X_1$ acting between Banach spaces is Fredholm if, and only if, the vector spaces

$$\ker(T) := T^{-1}(0)$$
 and $\operatorname{coker}(T) := X_1/TX_0$

are (both) finite dimensional. Since M is compact, a very well known and widely used result states that $P: H^s(M; E_0) \to H^{s-m}(M; E_1)$ is Fredholm if, and only if, P is elliptic, see for instance [45, 64, 80, 83, 89, 95] and the references therein. Consequently, if P is elliptic, then $\pi_{\alpha}(P)$ is also Fredholm. The converse is not true, however, in general.

To state our main result characterizing the Fredholm property of $\pi_{\alpha}(P)$ in terms of the " α -principal symbol" $\sigma_m^{\alpha}(P)$ of P, Theorem 1.2, we shall need to introduce $\sigma_m^{\Gamma}(P)$, the " Γ -equivariant principal symbol" of P, which is a refinement of the principal symbol $\sigma_m(P)$ of P that takes into account the action of the group Γ . The α -principal symbol $\sigma_m^{\alpha}(P)$ of P is a suitable restriction of the Γ -equivariant principal symbol $\sigma_m^{\Gamma}(P)$. Let us formulate now the precise definition of these concepts.

The main question that we answer in this paper is to determine when the induced operator $\pi_{\alpha}(P)$ of Equation (2) is Fredholm in terms of its Γ -equivariant principal symbol $\sigma_m^{\Gamma}(P)$, see Theorem 1.2 below for the precise statement.

The Γ -invariance of P implies that its principal symbol is also Γ invariant:

$$\sigma_m(P) \in \mathcal{C}^{\infty}(T^*M \setminus \{0\}; \operatorname{Hom}(E_0, E_1))^{\Gamma}$$
.

Let $\Gamma_{\xi} := \{ \gamma \in \Gamma \mid \gamma \xi = \xi \}$ denote the isotropy of a $\xi \in T_x^*M$, $x \in M$, as usual. The isotropy Γ_x of $x \in M$ is defined similarly. Then $\Gamma_{\xi} \subset \Gamma_x$ acts on E_{0x} and on E_{1x} , the fibers of $E_0, E_1 \to M$ at x. If $Q \in C^{\infty}(T^*M \setminus \{0\}; \text{Hom}(E_0, E_1))^{\Gamma}$, then $Q(\xi) \in \text{Hom}(E_{0x}, E_{1x})^{\Gamma_{\xi}}$. Let $\rho \in \widehat{\Gamma}_{\xi}$ be an irreducible representation of Γ_{ξ} , then

(4)
$$\widehat{Q}(\xi,\rho) := \pi_{\rho}[Q(\xi)] \in \operatorname{Hom}(E_{0x\rho}, E_{1x\rho})^{\Gamma_{\xi}}$$

denotes the restriction of Q to the isotypical component corresponding to ρ , with π_{ρ} defined in Equation (1). Let

(5)
$$X_{M,\Gamma} := \{ (\xi, \rho) \mid \xi \in T^*M \setminus \{0\} \text{ and } \rho \in \widehat{\Gamma}_{\xi} \}.$$

Thus Q defines a function on $X_{M,\Gamma}$. Applying this construction to $\sigma_m(P) \in \mathcal{C}^{\infty}(T^*M \setminus \{0\}; \operatorname{Hom}(E_0, E_1))^{\Gamma}$ we obtain a function, the Γ -principal symbol

(6)
$$\sigma_m^{\Gamma}(P): X_{M,\Gamma} \to \bigcup_{(x,\rho) \in X_{M,\Gamma}} \operatorname{Hom}(E_{0x\rho}, E_{1x\rho})^{\Gamma_{\xi}},$$
$$\sigma_m^{\Gamma}(P)(\xi,\rho) := \pi_{\rho}(\sigma_m(P)(\xi)) \in \operatorname{Hom}(E_{0x\rho}, E_{1x\rho})^{\Gamma_{\xi}}, \quad \xi \in T_x^*M.$$

That is $\sigma_m^{\Gamma}(P) := \widehat{\sigma_m(P)}$.

The α -principal symbol $\sigma_m^{\alpha}(P)$ of P, $\alpha \in \widehat{\Gamma}$, is defined in terms of $\sigma_m^{\Gamma}(P)$, but we need a crucial additional ingredient that takes α into account.

Recall that $\Gamma_{g\xi} = g\Gamma_{\xi}g^{-1}$ and that this defines an action of Γ on the set of stabilizer subgroups $\operatorname{Stab}_{\Gamma}(T^*M) := \{\Gamma_{\xi} \mid \xi \in T^*M\}$ given by $g \cdot \Gamma_{\xi} = \Gamma_{g\xi}$. For $\rho \in \widehat{\Gamma}_{\xi}$ define $g \cdot \rho \in \widehat{\Gamma}_{g\xi}$ by $(g \cdot \rho)(h) = \rho(g^{-1}hg)$, for all $h \in \Gamma_{g\xi}$. Let $\Gamma_0 \subset \Gamma$ be a minimal element (for inclusion) among the isotropy groups Γ_x of elements

 $x \in M$. Such a minimal element exists trivially, since Γ is finite. Moreover, if M/Γ is connected, then Γ_0 is unique up to conjugacy (see Subsection 2.4.3). We assume for a moment that this is the case, that is, that M/Γ is connected. Then we let

(7)
$$X_{M,\Gamma}^{\alpha} := \{ (\zeta, \rho) \in X_{M,\Gamma} \mid \exists g \in \Gamma, \operatorname{Hom}_{\Gamma_0}(g \cdot \rho, \alpha) \neq 0 \}.$$

(Note that it is implicit in the definition of $X_{M,\Gamma}^{\alpha}$ that $\Gamma_0 \subset \Gamma_{g\zeta} = g \cdot \Gamma_{\zeta}$.) In general (if M/Γ is not connected), we define $X_{M,\Gamma}^{\alpha}$ by taking the disjoint union of the corresponding spaces for each connected component of M/Γ , see Subsection 4.1.

Definition 1.1. The α-principal symbol $\sigma_m^{\alpha}(P)$ of P is the restriction of the Γ-principal symbol $\sigma_m^{\Gamma}(P)$ to $X_{M,\Gamma}^{\alpha}$:

$$\sigma_m^{\alpha}(P) := \sigma_m^{\Gamma}(P)|_{X_{M,\Gamma}^{\alpha}}.$$

We shall say that $P \in \psi^m(M; E_0, E_1)^{\Gamma}$ is α -elliptic if its α -principal symbol $\sigma_m^{\alpha}(P)$ is invertible everywhere on its domain of definition. Note that when $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$ is such that $E_{x\rho} = 0$, then $\sigma_m^{\Gamma}(P)(\xi, \rho) : 0 \to 0$ is always invertible.

1.3. Statement of the main result. An alternative formulation of Definition 1.1 is that P is α -elliptic if, and only if, σ_m^{Γ} is invertible on $X_{M,\Gamma}^{\alpha}$ (this is, of course, a condition only for those ρ such that $E_{i\rho} \neq 0$, because, otherwise, we get an operator acting on the zero spaces, which we admit to be invertible). We then have the following result extending the classical result (i.e. $\Gamma = \{1\}$) and the one from [12] (i.e. Γ finite abelian) to a general finite group Γ .

Theorem 1.2. Let Γ be a finite group acting on a smooth, compact manifold M and let $P \in \psi^m(M; E_0, E_1)^{\Gamma}$ be a Γ -invariant classical pseudodifferential operator acting between sections of two Γ -equivariant bundles $E_i \to M$, $i = 0, 1, m \in \mathbb{R}$, and $\alpha \in \widehat{\Gamma}$. We have that

$$\pi_{\alpha}(P): H^{s}(M; E_{0})_{\alpha} \rightarrow H^{s-m}(M; E_{1})_{\alpha}$$

is Fredholm if, and only if it is α -elliptic.

As in the abelian case, if Γ acts without fixed points on a *dense* open subset of M, then $X_{M,\Gamma} = X_{M,\Gamma}^{\alpha}$ for all $\alpha \in \widehat{\Gamma}$, by Corollary 4.15. Hence, in this case, P is α -elliptic if, and only if, it is elliptic. The ellipticity of P can thus be checked in this case simply by looking at the action of P on a single isotypical component. We stress, however, that if Γ is not discrete, this statement, as well as the statement of the above theorem, are no longer true. However, many intermediate results remain valid for compact Lie groups.

A motivation for our result comes from index theory. Let us assume that P is Γ -invariant and elliptic. Atiyah and Singer have determined, for any $\gamma \in \Gamma$, the value at γ of the character of $\operatorname{ind}_{\Gamma}(P) \in R(G)$. More precisely, they have computed $\operatorname{ind}_{\Gamma}(P)(\gamma) \in \mathbb{C}$ in terms of data at the fixed points of γ on M [7]. (Here $R(G) := \mathbb{Z}^{\widehat{G}}$ is the representation ring of G and is identified with a subalgebra of $C^{\infty}(G)^G$, the ring of conjugacy invariant functions on G via the characters of representations.) By contrast, the multiplicity of $\alpha \in \widehat{\Gamma}$ in $\operatorname{ind}_{\Gamma}(P)$ was much less studied. It did appear, however, implicitly in the work of Brüning [18], who initiated the program of studying the "isotypical heat trace" $\operatorname{tr}(p_{\alpha}e^{-t\Delta})$ and its short time asymptotic expansion. Its heat trace is nothing but the heat trace of $\pi_{\alpha}(\Delta)$. This

question was addressed also by Paradan and Vergne, who obtained several important related results, see [74] and the references therein. Brüning's program would lead, in particular, to a heat equation determination of the α -isotypical component of the Γ -equivariant index ind Γ (D) for Dirac type operators D. This program is one of the motivations of this paper.

Indeed, we obtain that the (Fredholm) index of $\pi_{\alpha}(P)$ depends only on the homotopy class of its α -principal symbol. Some results are contained, however, in Theorem 4.9 and in the remark following it. In particular, this yields results on the index theory of singular quotient spaces. We therefore expect our results to have applications to the Hodge theory of algebraic varieties [2, 3, 16, 27, 41], see Remark 4.11. In the case of a non discrete compact Lie group, the computation of this index is related to the index class of G-transversally elliptic operators initiated in [6, 86]. Since then, this has been studied in K-theory [9, 11, 47, 49] and in equivariant cohomology [10, 14, 15, 73].

Our proof makes heavy use of C^* -algebras by considering the natural C^* -algebra completions of algebras of pseudodifferential operators and related C^* -algebras. This point of view has been used and advocated of course by many people. C^* algebras were used very recently to obtain Fredholm conditions in [33, 53, 70], for example. Some of the algebras involved were groupoid algebras [5, 4, 26, 35, 67, 78]. Fredholm conditions play an important role in the study of the essential spectrum of Quantum Hamiltonians [13, 38, 37, 43, 52]. The technique of "limit operators" [51, 57, 58, 76] is related to groupoids. Some of the most recent papers using related ideas include [8, 25, 26, 28, 68, 69, 94], to which we refer for further references. Besides C^* -algebras, pseudodifferential operators were also used to obtain Fredholm conditions, see, for example, [34, 54, 42, 64, 61, 80, 81, 82] and the references therein. In addition to the works already mentioned, several general results on C^* and related algebras related to this work were obtained by Cordes and McOwen [32], Melo, Nest, and Schrohe [62], Melrose and Nistor [63], Rabinovich, Schulze, and Tarkhanov [77], Taylor [87], Voiculescu [96, 97, 98], and others. See [22, 23, 31, 50, 39, 40, 44, 65, 66] for some older, related results on singular integral operators.

1.4. Contents of the paper. We start in Section 2 with some preliminaries. We recall some facts about group actions, most notably the induction of representations and Frobenius reciprocity for finite groups. We also review some notions concerning the primitive spectrum of a C^* -algebra, as well as basic facts concerning (equivariant) pseudodifferential operators.

As in [12], we may assume $E_0 = E_1 = E$ and P to be of order zero. Let $A_M := \mathcal{C}(S^*M; \operatorname{End}(E))$. The most substantial technical results are in Section 3. There, we introduce the subset $\Omega_M := \{(\xi, \rho) \in S^*M \times \hat{\Gamma}_{\xi}, \ \rho \subset E_{\xi}\}$ of $X_{M,\Gamma}$ described above and identify the primitive spectrum of the C^* -algebra A_M^{Γ} of Γ -invariant symbols with the set Ω_M/Γ . Some care is taken to describe the corresponding topology on Ω_M/Γ . We then consider the canonical map from A_M^{Γ} to the Calkin algebra of $L^2(M; E)_{\alpha}$ and show that the closed subset of $\operatorname{Prim}(A_M^{\Gamma})$ associated to its kernel is $X_{M,\Gamma}^{\alpha}/\Gamma$.

These descriptions are used in Section 4 to prove the main result of the paper, Theorem 1.2. This section also addresses some particular cases of the Theorem and gives a few examples. We also explain the relation with previously known results, namely: the particular formulation in the abelian case, which was established in [12]; Fredholm conditions for transversally elliptic operators when the group Γ is not discrete; and Simonenko's local principle for Fredholm operators.

We thank Claire Debord, Paul-Emile Paradan, Elmar Schrohe, Georges Skandalis, and Andrei Teleman for useful discussions. The last named author thanks Max Planck Institute for support while this research was performed.

2. Preliminaries

This section is devoted to background material. For the most part, it will consist of a brief review of sections 2 and 3 of [12], where the reader will find more details, as well as definitions and results not repeated here. Note, however, that we need certain preliminary results for the case Γ non-commutative that were not needed in the abelian case. Nevertheless, the reader familiar with [12] can skip this section at a first reading.

For simplicity, let us from now on for the rest of this paper also assume that M/Γ is connected, except in Subsection 4.1, where we explain how the disconnected case reduces to the connected case.

2.1. **Group representations.** We follow the standard terminology and conventions. See, for instance, [12, 17, 84], where one can find further details. Most of the needed basic background material was recalled in greater detail in [12].

Throughout the paper, we denote by Γ a *finite* group acting by isometries on a smooth, Riemannian manifold M (without boundary). We use the standard notations, see [12, 17, 84], to which we refer for further details. If $x \in M$, then Γx is the Γ orbit of x and

(8)
$$\Gamma_x := \{ \gamma \in \Gamma \mid \gamma x = x \} \subset \Gamma$$

the isotropy group of the action at x.

We shall write $H \sim H'$ if the subgroups H and H' are conjugated in Γ . If $H \subset \Gamma$ is a subgroup, then $M_{(H)}$ will denote the set of elements of M whose isotropy Γ_x is conjugated to H (in Γ), that is, the set of elements $x \in M$ such that $\Gamma_x \sim H$.

Assuming that Γ acts on a space X, we denote by $\Gamma \times_H X$ the space

(9)
$$\Gamma \times_H X := (\Gamma \times X) / \sim,$$

where $(\gamma h, x) \sim (\gamma, hx), \forall \gamma \in \Gamma, h \in H \text{ and } x \in X.$

Let V be a normed complex vector space and $\mathcal{L}(V)$ be the set of bounded operators on V. A representation of Γ on V is a group morphism $\Gamma \to \mathcal{L}(V)$; in that case we also call V a Γ -module.

For any two Γ -modules \mathcal{H} and \mathcal{H}_1 , we shall denote by

$$\operatorname{Hom}_{\Gamma}(\mathcal{H}, \mathcal{H}_1) = \operatorname{Hom}(\mathcal{H}, \mathcal{H}_1)^{\Gamma} = \mathcal{L}(\mathcal{H}, \mathcal{H}_1)^{\Gamma}$$

the set of continuous linear maps $T: \mathcal{H} \to \mathcal{H}_1$ that commute with the action of Γ , that is, $T(\gamma \xi) = \gamma T(\xi)$ for all $\xi \in \mathcal{H}$ and $\gamma \in \Gamma$.

Let \mathcal{H} be a Γ -module and α an irreducible Γ -module. Then p_{α} will denote the Γ -invariant projection onto the α -isotypical component \mathcal{H}_{α} of \mathcal{H} , defined as the largest (closed) Γ submodule of \mathcal{H} that is isomorphic to a multiple of α . In other words, \mathcal{H}_{α} is the sum of all Γ -submodules of \mathcal{H} that are isomorphic to α . Notice that $\mathcal{H}_{\alpha} \simeq \alpha \otimes \operatorname{Hom}_{\Gamma}(\alpha, \mathcal{H})$.

Since Γ is finite, it is, in particular, compact, and hence we have

(10)
$$\mathcal{H}_{\alpha} \neq 0 \Leftrightarrow \operatorname{Hom}_{\Gamma}(\alpha, \mathcal{H}) \neq 0 \Leftrightarrow \operatorname{Hom}_{\Gamma}(\mathcal{H}, \alpha) \neq 0.$$

If $T \in \mathcal{L}(\mathcal{H})^{\Gamma}$ (i.e. T is Γ -equivariant), then $T(\mathcal{H}_{\alpha}) \subset \mathcal{H}_{\alpha}$ and we let

(11)
$$\pi_{\alpha}: \mathcal{L}(\mathcal{H})^{\Gamma} \to \mathcal{L}(\mathcal{H}_{\alpha}), \quad \pi_{\alpha}(T) := T|_{\mathcal{H}_{\alpha}},$$

be the associated morphism, as in Equation (1) of the Introduction. The morphism π_{α} will play an essential role in what follows.

2.2. **Induction and Frobenius reciprocity.** We now recall some definitions and results for induced representations mainly to set up notation and to obtain some intermediate results.

We let $V^{(I)} := \{f : I \to V\}$ for I finite. If $H \subset \Gamma$ is a subgroup (hence also finite) and V is a H-module, we define, as usual,

(12)
$$\operatorname{Ind}_{H}^{\Gamma}(V) := \mathbb{C}[\Gamma] \otimes_{\mathbb{C}[H]} V \\ \simeq \{ f : \Gamma \to V \mid f(gh^{-1}) = hf(g) \} \simeq V^{(\Gamma/H)}$$

to be the induced representation. The last isomorphism is obtained using a set of representatives of the right cosets Γ/H . The action of the group Γ on $\mathrm{Ind}_H^\Gamma(V)$ is obtained from the left multiplication on $\mathbb{C}[\Gamma]$ and the first isomorphism defines the Γ -module structure on $\mathrm{Ind}_H^\Gamma(V)$. The induction is a functor, that is, the Γ -module $\mathrm{Ind}_H^\Gamma(V)$ depends functorially on V.

Remark 2.1. Summarizing Remark 2.2 of [12], we have that

- (1) if V is a H-algebra, then $\operatorname{Ind}_H^{\Gamma}(V)$ is an algebra for the pointwise product,
- (2) if V is a left R-module (with compatible actions of Γ), then $\operatorname{Ind}_H^{\Gamma}(V)$ is a $\operatorname{Ind}_H^{\Gamma}(R)$ module, again with the pointwise multiplication,
- (3) the induction is compatible with morphisms of modules and algebras (change of scalars), again by the function representation of the induced representation.

See [12, Remark 2.2] for more details.

We shall use the $Frobenius\ reciprocity$ in the form that states that we have an isomorphism

$$\Phi = \Phi_{H,V}^{\Gamma,\mathcal{H}} : \operatorname{Hom}_{H}(\mathcal{H}, V) \to \operatorname{Hom}_{\Gamma}(\mathcal{H}, \operatorname{Ind}_{H}^{\Gamma}(V)),$$

$$(13) \qquad \Phi(f)(\xi) := \frac{1}{|H|} \sum_{g \in \Gamma} g \otimes_{\mathbb{C}[H]} f(g^{-1}\xi), \quad \xi \in \mathcal{H}, \ f \in \operatorname{Hom}_{H}(\mathcal{H}, V).$$

The version of the Frobenius reciprocity used in this paper is valid only for finite groups [17, 84] (although it can be suitably be generalized to the compact case). We note that a more precise notation would be to write $\operatorname{Hom}_H(\operatorname{Res}_H^{\Gamma}(\mathcal{H}), V)$ instead of our simplified notation $\operatorname{Hom}_H(\mathcal{H}, V)$.

Definition 2.2. Let A and B be finite groups and let H a subgroup of both A and B. Let $\alpha \in \widehat{A}$ and $\beta \in \widehat{B}$. We say that α and β are H-disjoint if $\operatorname{Hom}_H(\alpha, \beta) = 0$, otherwise we say that they are H-associated (to each other).

Let $\alpha \in \widehat{\Gamma}$, let $H \subset \Gamma$ be a subgroup, and $\beta \in \widehat{H}$. A useful consequence of the Frobenius reciprocity is that the multiplicity of α in $\operatorname{ind}_H^{\Gamma}(\beta)$ is the same as the multiplicity of β in the restriction of α to H. In particular, α is contained in $\operatorname{ind}_H^{\Gamma}(\beta)$ if, and only if, β is contained in the restriction of α to H, in which case recall that we say that α and β are H-associated (Definition 2.2). On the other

hand, recall that if β is *not* contained in the restriction of α to H, we say that α and β are H-disjoint.

Let V be a H-module and \mathcal{H} be the trivial Γ -module \mathbb{C} . Then we obtain, in particular, an isomorphism

(14)
$$\Phi: V^{H} = \operatorname{Hom}_{H}(\mathbb{C}, V) \simeq \operatorname{Hom}_{\Gamma}(\mathbb{C}, \operatorname{Ind}_{H}^{\Gamma}(V)) = \operatorname{Ind}_{H}^{\Gamma}(V)^{\Gamma},$$
$$\Phi(\xi) := \frac{1}{|H|} \sum_{g \in \Gamma} g \otimes_{\mathbb{C}[H]} \xi = \sum_{x \in \Gamma/H} x \otimes \xi.$$

If V is an algebra, then the map Φ is an isomorphism of algebras. In particular, we obtain the following consequences.

Remark 2.3. Let $H \subset \Gamma$ be a subgroup of Γ , β_j be non-isomorphic simple H-modules, $j = 1, \ldots, N$, and

$$\beta := \bigoplus_{j=1}^{N} \beta_j^{k_j}.$$

We then have that $\operatorname{Ind}_H^{\Gamma}(\beta) \simeq \bigoplus_{i=1}^N \operatorname{Ind}_H^{\Gamma}(\beta_i^{k_j})$ and the Frobenius isomorphism gives

(16)
$$\operatorname{Ind}_{H}^{\Gamma}(\operatorname{End}(\beta))^{\Gamma} \simeq \operatorname{End}(\beta)^{H} \simeq \bigoplus_{j=1}^{N} \operatorname{End}(\beta_{j}^{k_{j}})^{H} \simeq \bigoplus_{j=1}^{N} M_{k_{j}}(\mathbb{C}),$$

which is a semi-simple algebra and where the first isomorphism is induced by Φ of Equation (14).

We shall need the following refinement of the above remark.

Lemma 2.4. Let $\beta := \bigoplus_{j=1}^N \beta_j^{k_j}$ be as in Equation (15), let

$$T = (T_j) \in \operatorname{End}(\beta)^H \simeq \bigoplus_{j=1}^N \operatorname{End}(\beta_j^{k_j})^H,$$

with $T_i \in \text{End}(\beta_i^{k_j})^H$, and let $\xi_i \in \text{Ind}_H^{\Gamma}(\beta_i^{k_j})$. We let

$$\xi := (\xi_j) \in \bigoplus_{j=1}^N \operatorname{Ind}_H^{\Gamma}(\beta_j^{k_j}) \simeq \operatorname{Ind}_H^{\Gamma}(\beta).$$

Then $\Phi(T)(\xi) = (\Phi(T_j)\xi_j)_{j=1,...,N}$.

For the abelian case, the following elementary result was proved in [12] Proposition 2.5. That proof *does not* generalize to our case.

Proposition 2.5. Let $\beta := \bigoplus_{j=1}^{N} \beta_j^{k_j}$ be as in Equation (15). Let $J \subset \{1, 2, ..., N\}$ be the set of indices j such that α and β_j are H-disjoint (i.e. β_j is not contained in the restriction of α to H). Then the morphism

$$\pi_{\alpha}: \operatorname{Ind}_{H}^{\Gamma}(\operatorname{End}(\beta))^{\Gamma} \to \operatorname{End}(p_{\alpha}\operatorname{Ind}_{H}^{\Gamma}(\beta))$$

is such that

$$\ker(\pi_{\alpha}) = \bigoplus_{j \in J} \operatorname{Ind}_{H}^{\Gamma}(\operatorname{End}(\beta_{j}^{k_{j}}))^{\Gamma} \text{ and } \operatorname{Im}(\pi_{\alpha}) \simeq \bigoplus_{j \notin J} \operatorname{Ind}_{H}^{\Gamma}(\operatorname{End}(\beta_{j}^{k_{j}}))^{\Gamma}.$$

Proof. By Lemma 2.4, we can assume that N=1. Therefore the algebra $\operatorname{End}(\beta)^H$ is simple (more precisely, isomorphic to a matrix algebra $M_q(\mathbb{C})$, $q=k_1$). We shall use the isomorphism of Equation (16). The action of $\operatorname{Ind}_H^{\Gamma}(\operatorname{End}(\beta))^{\Gamma} \simeq \operatorname{End}(\beta)^H \simeq M_q(\mathbb{C})$ on $\operatorname{Ind}_H^{\Gamma}(\beta)$ is unital (i.e. non-degenerate), so the morphism

(17)
$$M_q(\mathbb{C}) \simeq \operatorname{Ind}_H^{\Gamma}(\operatorname{End}(\beta))^{\Gamma} \to \operatorname{End}(p_{\alpha}\operatorname{Ind}_H^{\Gamma}(\beta))$$

is injective if, and only if, $p_{\alpha} \operatorname{Ind}_{H}^{\Gamma}(\beta) \neq 0$. Notice the following equivalences

$$p_{\alpha}\operatorname{Ind}_{H}^{\Gamma}(\beta) \neq 0 \Leftrightarrow \operatorname{Hom}(\alpha, \operatorname{Ind}_{H}^{\Gamma}(\beta))^{\Gamma} \neq 0 \Leftrightarrow \operatorname{Hom}(\alpha, \beta)^{H} \neq 0.$$

The result then follows from Equation (10).

2.3. The primitive ideal spectrum of a C^* -algebra. We shall need a few basic concepts and facts about C^* -algebras. A general reference is [36]. Recall that a two-sided ideal $I \subset A$ of a C^* -algebra A is called *primitive* if it is the kernel of a non-zero, irreducible *-representation of A. Hence A is not a primitive ideal of itself. By Prim(A) we shall denote the set of primitive ideals of A, called the *primitive ideal spectrum* of A. If X is a locally compact space, then $C_0(X)$ denotes the space of continuous functions $X \to \mathbb{C}$ that vanish at infinity. The concept of primitive ideal spectrum is important for us since we have a natural homeomorphism

(18)
$$\operatorname{Prim}(\mathcal{C}_0(X)) \simeq X.$$

This identification lies at the heart of non-commutative geometry [29]. See also [24, 60].

If A is a type I C^* -algebra, then $\operatorname{Prim}(A)$ identifies with the set of isomorphism classes of irreducible representations of A. Any C^* -algebra with only finite dimensional irreducible representations is a type I algebra [36]. Most of the algebras considered in this paper (a notable exception are the algebras of compact operators), have this property.

The following example from [12] will be used several times.

Example 2.6. Let H be a finite group and $\beta = \bigoplus_{j=1}^N \beta_j^{k_j}$ be as in Remark 2.3. Then, as explained in that remark, $\mathcal{L}(\beta)^H \simeq \bigoplus_j M_{k_j}(\mathbb{C})$. The algebra $\mathcal{L}(\beta)^H = \operatorname{End}_H(\beta)$ is thus a C^* -algebra with only finite dimensional representations and we have natural homeomorphisms

$$Prim(End_H(\beta)) \leftrightarrow \{\beta_1, \beta_2, \dots, \beta_N\} \leftrightarrow \{1, 2, \dots, N\}.$$

The space Prim(A) is a topological space for the Jacobson topology: we refer to [36] for more details. We will recall some facts about this topology when we need it, see Lemma 3.2 below.

We shall need the following "central character" map.

Remark 2.7. Let Z be a commutative C^* -algebra and $\phi: Z \to M(A)$ be a *-morphism to the multiplier algebra M(A) of A [1, 21]. Assume that $\phi(Z)$ commutes with A and $\phi(Z)A = A$. Then Schur's lemma gives that every irreducible representation of A restricts to (a multiple of) a character of Z and hence there exists a natural continuous map

(19)
$$\phi^* : \operatorname{Prim}(A) \to \operatorname{Prim}(Z),$$

which we shall call also the *central character map* (associated to ϕ).

We conclude our discussion with the following simple result.

Lemma 2.8. We freely use the notation of Example 2.6. The inclusion of the unit $\mathbb{C} \to \operatorname{End}_H(\beta)$ induces a morphism $j : \mathcal{C}_0(X) \to \mathcal{C}_0(X; \operatorname{End}_H(\beta)) \simeq \mathcal{C}_0(X) \otimes \operatorname{End}_H(\beta)$. The resulting central character map is the first projection

$$(20) j^* : \operatorname{Prim}(\mathcal{C}_0(X; \operatorname{End}_H(\beta))) \simeq X \times \{1, 2, \dots, N\} \to X \simeq \operatorname{Prim}(\mathcal{C}_0(X)).$$

- 2.4. Group actions on manifolds. As before, we consider a finite group Γ acting by isometries on a compact Riemannian manifold M.
- 2.4.1. Slices and tubes. Given $x \in M$, the isotropy group Γ_x acts linearly and isometrically on T_xM . For r > 0, let $U_x := (T_xM)_r$ denote the set of vectors of length < r in T_xM . It is known then that, for r > 0 small enough, the exponential map gives a Γ -equivariant isometric diffeomorphism

(21)
$$W_x = \exp(\Gamma \times_{\Gamma_x} U_x) \simeq \Gamma \times_{\Gamma_x} U_x$$

where W_x is a Γ -invariant neighborhood of x in M and $\Gamma \times_{\Gamma_x} U_x$ is defined in equation (9). More precisely, W_x is the set of $y \in M$ at distance < r to the orbit Γx , if r > 0 is small enough. The set W_x is called a *tube* around x (or Γx) and the set U_x is called the *slice* at x. When M is compact, the injectivity radius is bounded from below, so we may assume that the constant r does not depend on x.

2.4.2. Equivariant vector bundles. Let us consider now a Γ -equivariant smooth vector bundle $E \to M$. Let us fix $x \in M$ and consider as above the tube $W_x \simeq \Gamma \times_{\Gamma_x} U_x$ around x, see Equation (21). We use this diffeomorphism to identify U_x to a subset of M, in which case, we can also assume the restriction of E to the slice U_x to be trivial. Therefore, there exists a Γ_x -module β such that

(22)
$$E|_{U_x} \simeq U_x \times \beta \text{ and }$$
$$E|_{W_x} \simeq \Gamma \times_{\Gamma_x} (U_x \times \beta),$$

The second isomorphism is Γ -equivariant.

Assume E is endowed with a Γ -invariant hermitian metric. We then have isomorphisms of Γ -modules:

(23)
$$L^{2}(W_{x}; E|_{W_{x}}) \simeq \operatorname{Ind}_{\Gamma_{x}}^{\Gamma}(L^{2}(U_{x}; \beta)) \text{ and}$$
$$C_{0}(W_{x}; E|_{W_{x}}) \simeq \operatorname{Ind}_{\Gamma_{x}}^{\Gamma}(C_{0}(U_{x}; \beta)).$$

In view of the previous isomorphism, we will often identify W_x and $\Gamma \times_{\Gamma_x} U_x$, making no distinction between them to simplify notations.

2.4.3. The principal orbit bundle. Recall that $M_{(H)}$ denotes the set of points of M whose stabilizer is conjugated in Γ to H. Recall that we have assumed that M/Γ is connected. It is known then [91] that there exists a minimal isotropy subgroup $\Gamma_0 \subset \Gamma$, in the sense that $M_{(\Gamma_0)}$ is a dense open subset of M, with measure zero complement in M.

In particular, the fact that M/Γ is connected gives that there exist minimal elements for the set of isotropy groups of points in M (with respect to inclusion) and all minimal isotropy groups are conjugated to a fixed subgroup $\Gamma_0 \subset \Gamma$. By the definition, the set $M_{(\Gamma_0)}$ consists of the points whose stabilizer is conjugated to that minimal subgroup. The set $M_{(\Gamma_0)}$ is called the *principal orbit bundle* of M. We will denote $M_{(\Gamma_0)}$ by M_0 in the sequel.

The principal orbit bundle $M_0 := M_{(\Gamma_0)}$ has the following useful property. If $x \in M_0$, then Γ_x acts trivially on the slice U_x at x, by the minimality of Γ_0 . Hence Γ_0 acts trivially on T_x^*M as well, which implies that $\Gamma_0 \subset \Gamma_\xi$ for any $\xi \in T_x^*M$. If, on the other hand, $x \in M$ is arbitrary (not necessarily in the principal orbit bundle), then the isotropy of Γ_x will contain a subgroup conjugated to Γ_0 .

2.5. **Pseudodifferential operators.** We continue to follow [12]. We also continue to assume that Γ is a finite Lie group that acts smoothly and isometrically on a smooth Riemannian manifold M. Let $\psi^m(M; E)$ denote the space of order m, classical pseudodifferential operators on M with compactly supported distribution kernel.

Let $\overline{\psi^0}(M; E)$ and $\overline{\psi^{-1}}(M; E)$ denote the respective norm closures of $\psi^0(M; E)$ and $\psi^{-1}(M; E)$. The action of Γ then extends to an action on $\psi^m(M; E)$, $\overline{\psi^0}(M; E)$, and $\overline{\psi^{-1}}(M; E)$. We shall denote by $\mathcal{K}(\mathcal{H})$ the algebra of compact operators acting on a Hilbert space \mathcal{H} . Of course, we have $\overline{\psi^{-1}}(M; E) = \mathcal{K}(L^2(M; E))$, since we have considered only pseudodifferential operators with compactly supported distribution kernels.

Let S^*M denote the *unit cosphere bundle* of a smooth manifold M, that is, the set of unit vectors in T^*M , as usual. We shall denote, as usual, by $C_0(S^*M; \operatorname{End}(E))$ the set of continuous sections of the *lift* of the vector bundle $\operatorname{End}(E) \to M$ to S^*M .

Corollary 2.9. We have an exact sequence

$$0 \to \mathcal{K}(L^2(M;E))^{\Gamma} \to \overline{\psi^0}(M;E)^{\Gamma} \xrightarrow{\sigma_0} \mathcal{C}_0(S^*M; \operatorname{End}(E))^{\Gamma} \to 0.$$

Proof. See, for instance, [12, Corollary 2.7].

2.5.1. The structure of regularizing operators. From now on, all our vector bundles will be Γ -equivariant vector bundles. We want to understand the structure of the algebra $\pi_{\alpha}(\overline{\psi^0}(M; E)^{\Gamma})$, for any fixed $\alpha \in \widehat{\Gamma}$ (see Equations (1) and (11) for the definition of the restriction morphism π_{α} and of the projectors $p_{\alpha} \in C^*(\Gamma)$).

We shall need the following standard result about negative order operators. Recall that, for $\alpha \in \widehat{\Gamma}$, we let π_{α} be the representation of $\overline{\psi^0}(M; E)^{\Gamma}$ on $L^2(M; E)_{\alpha}$ defined by restriction as before, Equations (1) and (11).

Proposition 2.10. We have the identifications

$$p_{\alpha}\psi^{-1}(M;E)^{\Gamma} \simeq \pi_{\alpha}(\overline{\psi^{-1}}(M;E)^{\Gamma})$$
$$= \pi_{\alpha}(\mathcal{K}(L^{2}(M;E))^{\Gamma}) = \mathcal{K}(L^{2}(M;E)_{\alpha})^{\Gamma},$$

where the first isomorphism map is simply π_{α} and

$$\mathcal{K}(L^2(M;E))^{\Gamma} = \overline{\psi^{-1}}(M;E)^{\Gamma} \simeq \bigoplus_{\alpha \in \widehat{\Gamma}} \mathcal{K}(L^2(M;E)_{\alpha})^{\Gamma}.$$

Proof. See, for example, [12, Section 3] for a proof.

3. The principal symbol

From now on we assume that M is **compact** and that M/Γ is connected. Let us fix an irreducible representation α of Γ and consider the restriction morphism π_{α} to the α -isotypical component of $L^2(M; E)$. Recall that this morphism was first introduced in Equation (1) and discussed in detail in Section 2.1. As in [12], we now turn to the identification of the quotient

$$\pi_{\alpha}(\overline{\psi^0}(M;E)^{\Gamma})/\pi_{\alpha}(\overline{\psi^{-1}}(M;E)^{\Gamma}).$$

The methods used in this paper diverge, however, drastically from those of [12].

Since $\pi_{\alpha}(\overline{\psi^{-1}}(M;E)^{\Gamma})$ was identified in the previous section, the promised identification of the quotient $\pi_{\alpha}(\overline{\psi^{0}}(M;E)^{\Gamma})/\pi_{\alpha}(\overline{\psi^{-1}}(M;E)^{\Gamma})$ will give further insight into the structure of the algebra $\pi_{\alpha}(\overline{\psi^{0}}(M;E)^{\Gamma})$ and will provide us, eventually,

with Fredholm conditions. Recall that, in this paper, we are assuming Γ to be finite. Nevertheless, a several intermediate results hold also in the case Γ compact.

3.1. The primitive ideal spectrum of A_M^{Γ} . As before, S^*M denotes the unit cosphere bundle of M. For the simplicity of the notation, we shall write

$$A_M := \mathcal{C}(S^*M; \operatorname{End}(E)),$$

as in the Introduction. Recall from Corollary 2.9 that we have an algebra isomorphism

(24)
$$\overline{\psi^0}(M;E)^{\Gamma}/\overline{\psi^{-1}}(M;E)^{\Gamma} \simeq A_M^{\Gamma}.$$

In our case, the inclusion $j: \mathcal{C}(S^*M/\Gamma) \subset A_M^{\Gamma}$ as a central subalgebra induces, as in Equation (19), a central character map

$$j^*: \operatorname{Prim}(A_M^{\Gamma}) \to S^*M/\Gamma$$
,

that underscores the local nature of the structure of the primitive ideal spectrum of A_M^{Γ} . We introduce the representation $\pi_{\xi,\rho}$ defined for any $f\in A_M^{\Gamma}$ by

$$\pi_{\xi,\rho}(f) = \pi_{\rho}(f(\xi)),$$

that is $\pi_{\xi,\rho}(f)$ is the restriction of $f(\xi) \in \text{End}(E_x)$ to the ρ -isotypical component of E_x . The central character map j^* was used in [12], Corollary 4.2, to obtain the following identification of $\text{Prim}(A_M^{\Gamma})$.

Proposition 3.1 ([12]). Let Ω_M be the set of pairs (ξ, ρ) , where $\xi \in S_x^*M$, $x \in M$, and $\rho \in \widehat{\Gamma}_{\xi}$ appears in E_x (i.e. $\operatorname{Hom}_{\Gamma_{\xi}}(\rho, E_x) \neq 0$).

- (1) The map $\Omega_M/\Gamma \ni \Gamma(\xi,\rho) \mapsto \ker(\pi_{\xi,\rho}) \in \text{Prim}(A_M^{\Gamma})$ is bijective.
- (2) The central character map $\Omega_M/\Gamma \simeq \operatorname{Prim}(A_M^\Gamma) \to S^*M/\Gamma$ maps $\Gamma(\xi, \rho) \in \Omega_M/\Gamma$ to $\Gamma\xi$ and is continuous and finite-to-one.

The space $\operatorname{Prim}(A_M^{\Gamma})$ is endowed with the Jacobson topology, which was recalled in Subsection 2.3; thus Proposition 3.1 allows us to use the central character map j^* to obtain a topology on Ω_M/Γ that will play a crucial role in what follows. We thus now turn to the study of this topology on Ω_M/Γ . We begin with the following standard lemma.

Lemma 3.2. Let A be a C^* -algebra. The family $(V_a)_{a\in A}$ defined by

$$V_a = \{ J \in \operatorname{Prim} A \mid a \notin J \},\$$

for any $a \in A$, is a basis of open sets for Prim(A).

Proof. Following [36], we know that the open, non-empty subsets of Prim(A) are exactly the sets

$${J \in \operatorname{Prim}(A) \mid I \not\subset J} \simeq \operatorname{Prim}(I)$$

where I ranges through the closed, non-zero, two-sided ideals of A. If $a \in A$, let us denote by $I_a := \overline{AaA}$ the closed, two-sided ideal generated by a. Then $a \notin J \Leftrightarrow I_a \not\subset J$, and hence $V_a = \operatorname{Prim}(I_a)$. This shows that V_a is open.

Next, let $V \subset \operatorname{Prim}(A)$ be a non-empty open subset and $J_0 \in V$. We know then that there exists a closed, two-sided ideal I, $0 \neq I \subset A$, such that $V = \operatorname{Prim}(I)$. We have $I \not\subset J_0$, and hence we can choose $a \in I \setminus J_0$. If $J \subset A$ is a primitive ideal such that $a \notin J$, then a fortiori $I \not\subset J$. Therefore $V_a \subset \operatorname{Prim}(I)$. This shows that $J_0 \in V_a \subset V$. Therefore the family $(V_a)_{a \in A}$ is a basis for the topology on $\operatorname{Prim}(A)$.

We shall use the bijection of Proposition 3.1 to conclude the following.

Corollary 3.3. A basis for the induced topology on $\Omega_M/\Gamma \simeq \operatorname{Prim}(A_M^{\Gamma})$ is given by the sets

$$V_f := \{ \Gamma(\xi, \rho) \in \Omega_M / \Gamma \mid \pi_{\xi, \rho}(f) \neq 0 \},$$

where f ranges through the non-zero elements of A_M^{Γ} .

3.2. The restriction morphisms. Let $\mathcal{O} \subset M$ be an open subset. Then $S^*\mathcal{O}$ is the restriction of S^*M to \mathcal{O} . We shall need the algebras

(25)
$$A_{\mathcal{O}} := \mathcal{C}_0(S^*\mathcal{O}; \operatorname{End}(E)) \quad \text{and} \quad B_{\mathcal{O}} := \overline{\psi^0}(\mathcal{O}; E).$$

Assume that $\mathcal{O} \subset M$ is Γ -invariant. The group Γ does not act, in general, as multipliers on the C^* -algebra $B_{\mathcal{O}} := \overline{\psi^0}(\mathcal{O}; E)$ (it does however act by conjugation), so the method used in [12] to compute $\overline{\psi^{-1}}(\mathcal{O}; E)^{\Gamma} \simeq \mathcal{K}(L^2(\mathcal{O}; E))^{\Gamma}$ does not extend to compute $B_{\mathcal{O}}^{\Gamma}$. We shall thus consider the natural, surjective map

(26)
$$\mathcal{R}_{\mathcal{O}}: A_{\mathcal{O}}^{\Gamma} := \mathcal{C}_{0}(S^{*}\mathcal{O}; \operatorname{End}(E))^{\Gamma} \simeq B_{\mathcal{O}}^{\Gamma} / \overline{\psi^{-1}}(\mathcal{O}; E)^{\Gamma}$$

 $\to \pi_{\alpha}(B_{\mathcal{O}}^{\Gamma}) / \pi_{\alpha}(\overline{\psi^{-1}}(\mathcal{O}; E)^{\Gamma}).$

Recall from Corollary 2.10 that $\pi_{\alpha}(\overline{\psi^{-1}}(M;E)^{\Gamma}) = \mathcal{K}(L^{2}(M;E)_{\alpha})^{\Gamma}$. Therefore, for a given $P \in \overline{\psi^{0}}(M;E)$, we have that $\pi_{\alpha}(P)$ is Fredholm if, and only if, the principal symbol of P is invertible in $A_{M}^{\Gamma}/\ker(\mathcal{R}_{M})$. This will be discussed in more detail in the next section.

We shall approach the computation of $\ker(\mathcal{R}_M) \subset A_M^{\Gamma}$ by determining the closed subset

(27)
$$\Xi := \operatorname{Prim}(A_M^{\Gamma}/\ker(\mathcal{R}_M)) \subset \operatorname{Prim}(A_M^{\Gamma})$$

of the primitive ideal spectrum of A_M^{Γ} corresponding to $\ker(\mathcal{R}_M)$. Once we will have determined Ξ , we will also have determined $\ker(\mathcal{R}_M)$, in view of the definitions recalled in Subsection 2.3 that put in bijection the closed, two-sided ideals of a C^* -algebra with the closed subsets of its primitive ideal spectrum.

Since $\mathcal{C}(M/\Gamma) \subset B_M$, it follows from the definition of \mathcal{R}_M that it is a $\mathcal{C}(M/\Gamma)$ -module morphism, and hence that $\ker(\mathcal{R}_M)$ is a $\mathcal{C}(M/\Gamma)$ -module. Let us also recall that

$$\mathcal{C}(M/\Gamma) \,=\, \mathcal{C}(M)^\Gamma \,\subset\, Z_M \,:=\, \mathcal{C}(S^*M)^\Gamma \,\subset\, Z(A_M^\Gamma) \,\subset\, A_M^\Gamma \,\subset\, A_M \,.$$

The local nature of $\ker(\mathcal{R}_M)$ and of the space Ξ is explained in the following remark.

Remark 3.4. Let $M/\Gamma = \bigcup V_k$ be an open cover and

$$\ker(\mathcal{R}_M)_{V_k} := \mathcal{C}_0(V_k) \ker(\mathcal{R}_M) = \ker(\mathcal{R}_{V_k}).$$

If we determine each $\ker(\mathcal{R}_M)_{V_k}$, then we determine $\ker(\mathcal{R}_M)$ using a partition of unity through:

(28)
$$\ker(\mathcal{R}_M) = \sum_{k}' \phi_k \ker(\mathcal{R}_{V_k}),$$

where \sum' refers to sums with only finitely many non-zero terms and (ϕ_k) is a partition of unity of M/Γ with continuous functions subordinated to the covering (V_k) (thus, in particular, supp $(\phi_k) \subset V_k$). Since M is compact, we can assume the covering to be finite. (If M was non-compact, then we would need to take the closure of the right hand side in Equation (28).) To determine \mathcal{R}_M , we can

therefore replace M by any of the open sets V_k in the covering and study $\ker(\mathcal{R}_{V_k})$. We shall do that for the covering of M/Γ with the tubes $W_x \simeq \Gamma \times_{\Gamma_x} U_x$ considered in 2.4.1, see Equation (21).

3.3. Local calculations. In view of Remark 3.4, we shall concentrate now on the local structure of $\ker(\mathcal{R}_M)$, that is, on the structure of $\ker(\mathcal{R}_{\mathcal{O}})$ for suitable ("small") open, Γ -invariant subsets $\mathcal{O} \subset M$. Let us fix then $x \in M$ and let $W_x \simeq \Gamma \times_{\Gamma_x} U_x$ be the tube around x, Equation (21). For simplicity, we shall write

(29)
$$A_x := A_{U_x} := \mathcal{C}_0(S^*U_x; \text{End}(E)) \text{ and } Z_x := Z(A_x^{\Gamma_x}).$$

For these algebras, the role of Γ will be played by Γ_x . For the statement of the following lemma, recall the definitions in Subsection (2.4), especially Equation (21).

Lemma 3.5. Let $W_x \simeq \Gamma \times_{\Gamma_x} U_x$. Then $S^*W_x \simeq \Gamma \times_{\Gamma_x} S^*U_x$ and we have Γ -equivariant algebra isomorphisms

$$A_{W_x} := \mathcal{C}_0(S^*W_x; \operatorname{End}(E)) \simeq \operatorname{Ind}_{\Gamma_x}^{\Gamma} \left(\mathcal{C}_0(S^*U_x; \operatorname{End}(E)) \right) =: \operatorname{Ind}_{\Gamma_x}^{\Gamma} (A_x).$$

Consequently, the Frobenius isomorphism Φ of Equation (14) induces an isomorphism

$$\Phi^{-1}: A_{W_-}^{\Gamma} \to A_x^{\Gamma_x}.$$

Proof. We have that $E|_{W_x} \simeq \Gamma \times_{\Gamma_x} (E|_{U_x})$, hence $\operatorname{End}(E)|_{W_x} \simeq \Gamma \times_{\Gamma_x} (\operatorname{End}(E)|_{U_x})$. Equation (23) then gives that $\mathcal{C}_0(W_x, \operatorname{End}(E)) \simeq \operatorname{Ind}_{\Gamma_x}^{\Gamma}(\mathcal{C}_0(U_x, \operatorname{End}(E)))$. The rest follows right away from the Frobenius reciprocity (more precisely, from Equation (14)) and from Equation (23), with β replaced with $\operatorname{End}(E)$.

Remark 3.6. In view of Equation (14), the isomorphism Φ of Lemma 3.5 can be written explicitly as follows. Let $f \in A_x^{\Gamma_x}$. Then, for any equivalence class $[\gamma, \xi] := \Gamma_x(\gamma, \xi) \in \Gamma \times_{\Gamma_x} S^*U_x \simeq S^*W_x$ we have

$$\Phi(f)([\gamma,\xi]) = [\gamma,f(\xi)],$$

where $[\gamma, f(\xi)] \in \Gamma \times_{\Gamma_x} (U_x \times \operatorname{End}(E_x))^{\Gamma_x} \simeq \Gamma \times_{\Gamma_x} \operatorname{End}(E|_{U_x})^{\Gamma_x} \simeq \operatorname{End}(E|_{W_x})^{\Gamma}$. This defines $\Phi(f) \in \mathcal{C}_0(S^*W_x; \operatorname{End}(E|_{W_x}))^{\Gamma} = A_{W_x}^{\Gamma}$.

Lemma 3.5 together with the following remark will allow us to reduce the study of the algebra A_M^{Γ} to that of its analogues defined for slices.

Remark 3.7. Let U be an open set of some euclidean space and $W = U \times \{1, 2, ..., N\}$, where the space on the second factor is endowed with the discrete topology. For simplicity, we identify $L^2(W)$ with $L^2(U)^N$ using the map $f \mapsto (f(i))_{i=1...N}$. Then

(30)
$$\psi^{-1}(W) = M_N(\psi^{-1}(U)) \simeq \psi^{-1}(U) \otimes M_N(\mathbb{C}) \text{ and hence}$$
$$\overline{\psi^{-1}(W)} = M_N(\overline{\psi^{-1}(U)}) \simeq \overline{\psi^{-1}(U)} \otimes M_N(\mathbb{C}).$$

On the other hand, if A^N denotes the direct sum of N-copies of the algebra A, then we have the following inclusions of algebras

(31)
$$\psi^{0}(U)^{N} \subset \psi^{0}(W) \subset M_{N}(\psi^{0}(U)) \simeq \psi^{0}(U) \otimes M_{N}(\mathbb{C}), \text{ and hence}$$
$$\overline{\psi^{0}}(U)^{N} \subset \overline{\psi^{0}}(W) \subset M_{N}(\psi^{0}(U)) \simeq \overline{\psi^{0}}(U) \otimes M_{N}(\mathbb{C}).$$

The following lemma makes explicit the group actions in the isomorphisms of the last remark. Thus, in analogy with the definitions of the algebras $A_{W_x} = \mathcal{C}_0(S^*W_x; \operatorname{End}(E))$ and $A_x = \mathcal{C}_0(S^*U_x; \operatorname{End}(E))$, we consider the algebras

(32)
$$B_{W_x} := \overline{\psi^0}(W_x; E) \quad \text{and} \quad B_x := \overline{\psi^0}(U_x; E).$$

We shall also use the standard notation $V^{(I)} := \{f : I \to V\}$ for I finite, as before.

Lemma 3.8. We keep the notation of Lemma 3.5 and of Equation (32) above. Then we have Γ -equivariant algebra isomorphisms

$$B_{W_x} \simeq \operatorname{Ind}_{\Gamma_x}^{\Gamma}(B_x) + \overline{\psi^{-1}}(W_x; E)$$
.

Consequently, $B_{W_x}^{\Gamma} \simeq \Phi(B_x^{\Gamma_x}) + \overline{\psi^{-1}}(W_x; E)^{\Gamma}$.

Proof. Since $B_y = B_{U_y} \subset B_{W_x}$ for all $y \in \Gamma x$ and since U_x and U_y are diffeomorphic through any $\gamma \in \Gamma$ such that $\gamma x = y$ we obtain the inclusion $B_x^{(\Gamma/\Gamma_x)} \subset B_{W_x}$, as in Remark 3.7. Similarly, since $B_x \to A_x$ is surjective, we obtain the equality $B_{W_x} = B_x^{(\Gamma/\Gamma_x)} + \overline{\psi^{-1}}(W_x; E)$ as in the same remark. From Equation (24) and Lemma 3.5 we know that $B_{W_x}/\overline{\psi^{-1}}(W_x; E) \simeq A_{W_x} \simeq A_{U_x}^{(\Gamma/\Gamma_x)} = \operatorname{Ind}_{\Gamma_x}^{\Gamma}(A_x)$, and hence we obtain $B_{W_x} \simeq \operatorname{Ind}_{\Gamma_x}^{\Gamma}(B_x) + \overline{\psi^{-1}}(W_x; E)$. The last isomorphism follows from the Frobenius reciprocity (more precisely, from Equation (23), with β replaced with B_x) and from the exactness of the functor $V \to V^{\Gamma}$.

To be able to make further progress, it will be convenient to look first at the case when $x \in M$ has minimal isotropy $\Gamma_x \sim \Gamma_0$, that is, when x belongs to the principal orbit bundle $M_0 := M_{(\Gamma_0)}$. The notation Γ_0 will remain fixed from now on.

3.4. Calculations for the principal orbit bundle. We assume as before that M/Γ is connected. Let Γ_0 be a minimal isotropy group (which, we recall, is unique up to conjugation). Let $x \in M$ be our fixed point and Γ_x its isotropy, as before. The case when Γ_x is conjugated to Γ_0 is simpler since, as noticed already, then Γ_x acts trivially on U_x .

Let us fix $x \in M$ with isotropy group $\Gamma_x = \Gamma_0$. As before, we let

$$W_x \simeq \Gamma \times_{\Gamma_0} U_x$$
 and $E_{|W_x} \simeq \Gamma \times_{\Gamma_0} (U_x \times \beta)$,

where β is some Γ_0 -module, as in Equations (21) and (22). We decompose β into a direct sum of representations of the form $\beta_j^{k_j}$ for some non-isomorphic irreducible module (or representation) β_j of Γ_0 , again as before:

$$E_x = \beta \simeq \oplus \beta_i^{k_j}$$
.

Remark 3.9. We have noticed earlier that Γ_0 acts trivially on U_x , hence on T_x^*M . In particular S^*M also has Γ_0 as minimal isotropy subgroup, and S^*M_0 is a dense subset of the principal bundle of S^*M .

Corollary 3.10. Let $x \in M$ be such that $\Gamma_x = \Gamma_0$ and $\beta = \bigoplus_{j=1}^N \beta_j^{k_j}$, for some non-isomorphic, irreducible Γ_0 -modules β_j . Then

$$A_{W_x}^{\Gamma} \simeq A_x^{\Gamma_x} \simeq \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta) \simeq \bigoplus_{j=1}^N M_{k_j} (\mathcal{C}_0(S^*U_x)).$$

In particular, the canonical central character map

$$\operatorname{Prim}(A_x^{\Gamma_0}) \to S^*U_x \simeq \operatorname{Prim}(\mathcal{C}_0(S^*U_x)^{\Gamma_0})$$

of Proposition 3.1 corresponds to the trivial finite covering $S^*U_x \times \text{Prim}(\text{End}_{\Gamma_0}(\beta)) \to$ S^*U_x .

Proof. The first isomorphism is repeated from Lemma 3.5. The second one is obtained from the following:

- (i) from the definition of $A_x = A_{U_x}$,
- (ii) from the assumption that $\Gamma_x = \Gamma_0$,
- (iii) from the fact that Γ_0 acts trivially on U_x , and
- (iv) from the identifications

$$A_x^{\Gamma_0} := \mathcal{C}_0(S^*U_x; \operatorname{End}(E))^{\Gamma_0} \simeq \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}(\beta)^{\Gamma_0}.$$

The last isomorphism follows from Example 2.6 and the isomorphism $M_n(\mathbb{C}) \otimes$ $A \simeq M_n(A)$, valid for any algebra A. The rest follows from Lemma 2.8.

Indeed, since both $C_0(S^*U_x)$ and $\operatorname{End}(\beta)^{\Gamma_0}$ have only finite dimensional irreducible representations, we obtain $Prim(A_x^{\Gamma_0}) = S^*U_x \times Prim(End_{\Gamma_0}(\beta)) \simeq S^*U_x \times Pr$ $\{1,2,\ldots,N\}$, where we use the identification $Prim(\mathcal{C}_0(S^*U_x)) \simeq S^*U_x$ and where the set $\{1, 2, ..., N\}$ is in natural bijection with the primitive ideal spectrum of the algebra $\operatorname{End}_{\Gamma_0}(\beta) \simeq \bigoplus_{j=1}^N M_{k_j}(\mathbb{C})$. The inclusion $\mathcal{C}_0(S^*U_x) = \mathcal{C}_0(S^*U_x)^{\Gamma_0} \to A_x^{\Gamma_0}$ is given by the unital inclusion $\mathbb{C} \to \bigoplus_{j=1}^N M_{k_j}(\mathbb{C})$. Hence the map $\operatorname{Prim}(A_x^{\Gamma_0}) \to S^*U_x$ identifies with the first projection in $S^*U_x \times \{1, 2, \dots, N\} \to S^*U_x$. That is, it is a trivial covering, as claimed.

The fibers of $Prim(A_{M_0}^{\Gamma}) \to M_0/\Gamma$ are thus the simple factors of $End(E_x)^{\Gamma_0}$, whose structure was determined in Example 2.6. We shall need the following remark similar to Remark 3.7, but simpler.

Remark 3.11. Let U be an open subset of a euclidean space, let V be a finite dimensional vector space and let V denote, by abuse of notation, also the trivial, vector bundle with fiber V. Then we have natural isomorphisms

$$\psi^{-1}(U;V) \simeq \psi^{-1}(U) \otimes \operatorname{End}(V)$$
 and $\psi^{0}(U;V) \simeq \psi^{0}(U) \otimes \operatorname{End}(V)$.

Consequently, we also have the analogous isomorphisms for the completions

$$\overline{\psi^{-1}}(U;V) \simeq \overline{\psi^{-1}}(U) \otimes \operatorname{End}(V)$$
 and $\overline{\psi^{0}}(U;V) \simeq \overline{\psi^{0}}(U) \otimes \operatorname{End}(V)$.

We are in position now to determine the kernel of \mathcal{R}_{W_x} , when x is in the principal orbit bundle. We will use the notation of Subsection 2.4 that was recalled at the beginning of this subsection as well as the notation of Subsection 2.2. In particular, recall that $\beta_i \in \widehat{\Gamma}_0$ and $\alpha \in \widehat{\Gamma}$ are said to be Γ_0 -disjoint if β_i is not contained in the restriction of α to Γ_0 . Also, Φ is the Frobenius isomorphism, Equations (13) and (14) and Corollary 3.10.

Proposition 3.12. Let $\Gamma_x = \Gamma_0$, let $E_x = \beta = \bigoplus_{j=1}^N \beta_j^{k_j}$, and $\Phi : \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta) \simeq A_x^{\Gamma_0} \to A_{W_x}^{\Gamma}$ be the Frobenius isomorphism of Corollary 3.10. Then

- (1) $C_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta_j^{k_j}) \subset \Phi^{-1}(\ker(\mathcal{R}_{W_x}))$ if β_j and α are Γ_0 -disjoint, and (2) $C_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta_j^{k_j}) \cap \Phi^{-1}(\ker(\mathcal{R}_{W_x})) = 0$ if β_j and α are Γ_0 -associated.

In particular, Also, let $J \subset \{1, 2, ..., N\}$ be the set of indices j such that β_j and α are Γ_0 -disjoint, then

$$\ker(\mathcal{R}_{W_x}) = \Phi\left(\bigoplus_{j \in J} \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta_j^{k_j})\right) \text{ and}$$
$$\pi_{\alpha}(B_M^{\Gamma})/\pi_{\alpha}(\overline{\psi^{-1}}(M; E)^{\Gamma}) \simeq \Phi\left(\bigoplus_{j \notin J} \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta_j^{k_j})\right).$$

Proof. The proof is essentially a consequence of Proposition 2.5 by including U_x as a parameter, using also Lemma 3.8. To see how this is done, we will use the notation of that lemma, in particular, $W_x \simeq \Gamma \times_{\Gamma_0} U_x \simeq (\Gamma/\Gamma_0) \times U_x$ and $E \simeq \Gamma \times_{\Gamma_0} (U_x \times \beta)$. We identify W_x with $\Gamma \times_{\Gamma_x} U_x$, i.e. we work with $W_x = \Gamma \times_{\Gamma_x} U_x$.

Let π_{α} the fundamental morphism of restriction to the α -isotypical component, see Equations (1) and (11). Recall that $B_x := \overline{\psi^0}(U_x; E)$. Since Γ_x acts trivially on U_x , Remark 3.11 yields the Γ -equivariant isomorphisms

(33)
$$\operatorname{Ind}_{\Gamma_0}^{\Gamma}(B_x) \simeq \overline{\psi^0}(U_x) \otimes \operatorname{Ind}_{\Gamma_0}^{\Gamma}(\operatorname{End}(\beta)) \subset B_{W_x}$$

where the last inclusion is modulo the trivial identification given by $P \otimes f(s)(\gamma, x) = P(f(\gamma)s(\gamma))(x)$, $P \in \overline{\psi^0}(U_x)$, $f \in \operatorname{Ind}_{\Gamma_0}^{\Gamma}(\operatorname{End}(\beta))$ and $s \in C_c(W_x, \operatorname{End}(E))$. Combining further Remark 3.11 with Remark 3.7, we further obtain the isomorphism

$$\overline{\psi^{-1}}(W_x; E) \simeq \overline{\psi^{-1}}(U_x) \otimes \operatorname{End}(\operatorname{Ind}_{\Gamma_0}^{\Gamma}(\beta)).$$

Lemma 3.8 and the exactness of the functor $V \to V^{\Gamma}$ give $\pi_{\alpha}(B_{W_x}^{\Gamma}) = \pi_{\alpha} \circ \Phi(B_x^{\Gamma_x}) + \pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma})$. Hence we obtain

$$\pi_{\alpha}(B_{W_x}^{\Gamma})/\pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma}) = \pi_{\alpha} \circ \Phi(B_x^{\Gamma_x})/\pi_{\alpha} \circ \Phi(B_x^{\Gamma_x}) \cap \pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma}).$$

Let \mathfrak{A} and \mathfrak{J} be the image and, respectively, the kernel of $\pi_{\alpha}: \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\operatorname{End}(\beta))^{\Gamma} \to \operatorname{End}(p_{\alpha}\operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta))$, which have been identified in Proposition 2.5 in terms of the set J. Recall next from Equation (23) that $L^{2}(W_{x}; E) = L^{2}(U_{x}) \otimes \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta)$, again Γ -equivariantly. Each time, the action is on the second component, since $\Gamma_{0} = \Gamma_{x}$ acts trivially on $\overline{\psi^{0}}(U_{x})$. The action of $\operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(B_{x}) \subset B_{W_{x}}$ on $L^{2}(W_{x}; E) = L^{2}(U_{x}) \otimes \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta)$ is compatible with the tensor product decomposition of Equation (33), in the sense that $\overline{\psi^{0}}(U_{x})$ acts on $L^{2}(U_{x})$ and $\operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\operatorname{End}(\beta))$ acts on $\operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta)$. Also, $\operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(B_{x})^{\Gamma} \simeq \overline{\psi^{0}}(U_{x}) \otimes \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\operatorname{End}(\beta))^{\Gamma}$, (we use this isomorphism to identify them). We obtain that

(34)
$$\pi_{\alpha} \circ \Phi(B_x^{\Gamma_x}) = \pi_{\alpha}(\operatorname{Ind}_{\Gamma_0}^{\Gamma}(B_x)^{\Gamma}) = \overline{\psi^0}(U_x) \otimes \mathfrak{A}.$$

On the other hand, Corollary 2.10 then gives that $\pi_{\alpha}(\overline{\psi^{-1}}(W_x; \operatorname{End}(E))^{\Gamma})$ is the algebra of Γ -invariant compact operators acting on the space $p_{\alpha}(L^2(W_x, \operatorname{End}(E)))$. Therefore, $\overline{\psi^{-1}}(U_x) \otimes \mathfrak{A} \subset \pi_{\alpha}(\overline{\psi^{-1}}(W_x; \operatorname{End}(E))^{\Gamma})$, since $\overline{\psi^{-1}}(U_x) \otimes \mathfrak{A}$ consists of compact, Γ -invariant operators acting on $p_{\alpha}(L^2(W_x, E))$. Consequently,

$$(35) \quad \overline{\psi^{-1}}(U_x) \otimes \mathfrak{A} \subset \pi_{\alpha}(\operatorname{Ind}_{\Gamma_0}^{\Gamma}(B_x)^{\Gamma}) \cap \pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma}) \\ \subset \overline{\psi^0}(U_x) \otimes \mathfrak{A} \cap \mathcal{K}(p_{\alpha}L^2(W_x; E))^{\Gamma} \subset \overline{\psi^{-1}}(U_x) \otimes \mathfrak{A},$$

and hence we have equalities everywhere.

Recall from Corollary 3.10 that $A_{W_x}^{\Gamma} \simeq A_x^{\Gamma_x}$. We obtain that the map

$$(36) \mathcal{R}_{W_x}: A_{W_x}^{\Gamma} \simeq B_{W_x}^{\Gamma} / \overline{\psi^{-1}}(W_x; E)^{\Gamma} \to \pi_{\alpha}(B_{W_x}^{\Gamma}) / \pi_{\alpha}(\overline{\psi^{-1}}(W_x; E)^{\Gamma})$$

becomes, up to the canonical isomorphisms above, the map

(37)
$$A_x^{\Gamma_x} \simeq \mathcal{C}_0(S^*U_x) \otimes \operatorname{End}_{\Gamma_0}(\beta) \to \pi_{\alpha}(B_{W_x}^{\Gamma})/\pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma})$$
$$= \pi_{\alpha} \circ \Phi(B_x^{\Gamma_x})/\pi_{\alpha} \circ \Phi(B_x^{\Gamma_x}) \cap \pi_{\alpha}(\overline{\psi^{-1}}(W_x)^{\Gamma})$$
$$\simeq \overline{\psi^0}(U_x) \otimes \mathfrak{A}/\overline{\psi^{-1}}(U_x) \otimes \mathfrak{A} \simeq \mathcal{C}_0(S^*U_x) \otimes \mathfrak{A},$$

with all maps being surjective and preserving the tensor product decompositions. This identifies the kernel of \mathcal{R}_{W_x} with $\mathcal{C}_0(S^*U_x) \otimes \mathfrak{J}$ and the image of \mathcal{R}_{W_x} with $\mathcal{C}_0(S^*U_x) \otimes \mathfrak{A}$. The rest of the statement follows from the identification of \mathfrak{J} and \mathfrak{A} in Proposition 2.5.

Proposition 3.12 above and its proof give the following corollary.

Corollary 3.13. We use the notation of Proposition 3.12 and we identify the space $\operatorname{Prim}(\operatorname{End}(\beta))$ with $\{1,2,\ldots,N\}$ as in Remark 2.6. Then the homeomorphism $\operatorname{Prim}(A_{W_x}^{\Gamma}) \simeq S^*U_x \times \{1,2,\ldots,N\}$ maps the set $\Xi \cap \operatorname{Prim}(A_{W_x}^{\Gamma})$ to $S^*U_x \times J$. In particular, the restriction $\Xi \cap \operatorname{Prim}(A_{W_x}^{\Gamma}) \to S^*U_x$ of the central character is a covering as well.

Proof. Using the notations of the proof of Proposition 3.12, we have that $\ker(\mathcal{R}_{W_x})$ has primitive ideal spectrum $S^*U_x \times \operatorname{Prim}(\mathfrak{J})$. We have $\Xi \cap \operatorname{Prim}(A_{W_x}^{\Gamma}) = S^*U_x \times \operatorname{Prim}(\mathfrak{A})$.

The same methods yield the following result (recall that $M_0 = M_{(\Gamma_0)}$ is the principal orbit bundle).

Corollary 3.14. Let $M_0 := M_{(\Gamma_0)}$, the principal orbit bundle. The central character map $\operatorname{Prim}(A_{M_0}^{\Gamma}) \to S^*M_0/\Gamma$ defined by the inclusion $C_0(S^*M_0/\Gamma) \subset Z(A_{M_0}^{\Gamma})$ is a covering with typical fiber $\operatorname{Prim}(\operatorname{End}(E_x)^{\Gamma_0})$ such that $\Xi \cap \operatorname{Prim}(A_{M_0}^{\Gamma}) \to S^*M_0/\Gamma$ is a subcovering, see (27) for the definition of Ξ . In particular, $\Xi \cap \operatorname{Prim}(A_{M_0}^{\Gamma})$ is open and closed in $\operatorname{Prim}(A_{M_0}^{\Gamma})$.

Proof. The first statement is true locally, by Corollary 3.10, and hence it is true globally. Indeed, let $x \in M_0$, let $\xi \in S_x^*M_0$, and let $\rho \in \widehat{\Gamma}_x$ that appears in E_x (so $(\xi,\rho) \in \Omega_M$). We let $W_x \subset M_0 \subset M$ be the typical tube with minimal isotropy $\Gamma_x = \Gamma_0$, as before. Let $Z_x := \mathcal{C}_0(S^*W_x)^\Gamma \subset Z_M = \mathcal{C}(S^*M)^\Gamma$. Then $\operatorname{Prim}(Z_xA_M^\Gamma)$ is an open neighborhood in $\operatorname{Prim}(A_{M_0}^\Gamma)$ of the primitive ideal $\ker(\pi_{\xi,\rho})$, see Proposition 3.1 for notation and details. We have that $Z_xA_M^\Gamma = A_{W_x}^\Gamma$ and hence, on $\operatorname{Prim}(Z_xA_M^\Gamma)$, the central character is a covering, by Corollary 3.10. Similarly, its restriction to $\Xi \cap \operatorname{Prim}(Z_xA_M^\Gamma)$ is a covering by Corollary 3.13.

Putting Corollary 3.14 and Proposition 3.12 together we obtain the following results.

Corollary 3.15. Let M_0 be the principal orbit type of M. The ideal $\ker(\mathcal{R}_{M_0}) = A_{M_0}^{\Gamma} \cap \ker(\mathcal{R}_M)$ is defined by the closed subset $\Xi_0 := \Xi \cap \operatorname{Prim}(A_{M_0}^{\Gamma})$ of $\operatorname{Prim}(A_{M_0}^{\Gamma})$ consisting of the sheets of $\operatorname{Prim}(A_{M_0}^{\Gamma}) \to S^*M_0/\Gamma$ that correspond to the simple factors $\operatorname{End}(E_{x\rho})^{\Gamma_0}$ of $\operatorname{End}(E_x)^{\Gamma_0}$ with ρ and α Γ_0 -associated.

If Γ is abelian, then ρ and α are characters and saying that they are Γ_0 -associated means, simply, that their restrictions to Γ_0 coincide: $\rho|_{\Gamma_0} = \alpha_{\Gamma_0}$. This is consistent with the definition given in [12].

3.5. The non-principal orbit case. As in the rest of the paper, we assume M/Γ to be connected. We will show in Theorem 3.17 that Ξ is the closure of Ξ_0 in $\operatorname{Prim}(A_M^{\Gamma})$. To that end, we first construct a suitable basis of neighborhoods of $\operatorname{Prim}(A_M^{\Gamma})$ using Lemma 3.2.

Remark 3.16. Let $\Gamma(\xi, \rho) \in \operatorname{Prim}(A_M^{\Gamma})$, where we have used the description of $\operatorname{Prim}(A_M^{\Gamma})$ provided in Proposition 3.1 as orbits of pairs $\xi \in S^*M$ and suitable $\rho \in \widehat{\Gamma}_{\xi}$. We construct a basis of neighborhoods $(V_{\xi,\rho,n})_{n\in\mathbb{N}}$ of $\Gamma(\xi,\rho)$ in $\operatorname{Prim}(A_M^{\Gamma})$ as follows. Let $\xi \in S_x^*M$ (that is, ξ sits above $x \in M$) and we use the notation U_x and W_x of Equation (21), as always.

First, by choosing a different point ξ in its orbit, if necessary, we may assume that $\Gamma_0 \subset \Gamma_{\xi}$. Now let $(\mathcal{O}_n)_{n \in \mathbb{N}}$ be a family of Γ_{ξ} -invariant neighborhoods of ξ in S^*U_x such that:

- for all n and $\gamma \in \Gamma \setminus \Gamma_{\mathcal{E}}$, we have $\gamma \mathcal{O}_n \cap \mathcal{O}_n = \emptyset$,
- $\mathcal{O}_{n+1} \subset \mathcal{O}_n$ and $\bigcap_{n \in \mathbb{N}} \mathcal{O}_n = \{\xi\}.$

For any $n \in \mathbb{N}$, we choose a function $\varphi_n \in \mathcal{C}_c(\mathcal{O}_n)^{\Gamma_{\xi}}$ such that $\varphi_n \equiv 1$ on \mathcal{O}_{n+1} . Let $p_{\rho} \in \operatorname{End}(E_x)^{\Gamma_{\xi}}$ be the projection onto $E_{x\rho}$. We can assume the bundle E to be trivial on U_x and, using that, we first extend p_{ρ} constantly on \mathcal{O}_n and then as an element $q_n \in \mathcal{C}_c(S^*U_x; \operatorname{End}(E_x))^{\Gamma_x}$ defined as

$$q_n := \begin{cases} \Phi_{\Gamma_{\xi}, \Gamma_x}(\varphi_n p_{\rho}) & \text{on } \Gamma_x \mathcal{O}_n \\ 0 & \text{on } S^* U_x \setminus \Gamma_x \mathcal{O}_n, \end{cases}$$

with $\Phi_{\Gamma_{\xi},\Gamma_{x}}$ the Frobenius isomorphism of Equation (14). Let us set $\tilde{q}_{n} := \Phi_{\Gamma_{x},\Gamma}(q_{n}) \in A_{M}^{\Gamma}$, where $\Phi_{\Gamma_{x},\Gamma}$ is the Frobenius isomorphism of Equation (14). Finally, we associate to \tilde{q}_{n} the open set

$$V_{\xi,\rho,n} \,:=\, \{J\in \operatorname{Prim}(A_M^\Gamma)\mid \tilde{q}_n\notin J\}.$$

Recall from 3.2 that $V_{\xi,\rho,n}$ is an open subset of $\operatorname{Prim}(A_M^{\Gamma})$. Moreover, it follows from our definition that $V_{\xi,\rho,n+1} \subset V_{\xi,\rho,n}$ and that $\bigcap_{n \in \mathbb{N}} V_{\xi,\rho,n} = \{\Gamma(\xi,\rho)\}$.

Recall that we are assuming that M/Γ is connected.

Theorem 3.17. Let $\Xi := \operatorname{Prim}(A_M^{\Gamma}/\ker(\mathcal{R}_M)) \subset \operatorname{Prim}(A_M^{\Gamma})$ be the closed subset defined by the ideal $\ker(\mathcal{R}_M)$. Then Ξ is the closure in $\operatorname{Prim}(A_M^{\Gamma})$ of the set $\Xi_0 := \Xi \cap \operatorname{Prim}(A_{M_0}^{\Gamma})$, where M_0 is the principal orbit bundle of M.

Proof. We have that $\overline{\Xi}_0 \subset \Xi$ since $\Xi_0 \subset \Xi$ and the latter is a closed set. Conversely, let $\mathfrak{P} \in \operatorname{Prim}(A_M^{\Gamma}) \setminus \overline{\Xi}_0$. We will show that $\mathfrak{P} \notin \Xi$. Let \mathfrak{P} correspond to $(\xi, \rho) \in \Omega_M$, as in Proposition 3.1. We may assume that $\Gamma_0 \subset \Gamma_{\xi}$. Let x be projection of ξ onto M. Since the problem is local, we may also assume that $U_x \subset T_xM$, that $M = W_x := \Gamma \times_{\Gamma_x} U_x$, and that $E := \Gamma \times_{\Gamma_x} (U_x \times \beta)$ for some Γ_x -module β .

Using the notations of Remark 3.16, there exists n > 0 such that $V_{\xi,\rho,n} \cap \Xi_0 = \emptyset$. Let $\tilde{q}_n = \Phi_{\Gamma_x,\Gamma}(q_n)$ be the symbol defined in Remark 3.16. The description of Ξ_0 provided in Corollary 3.15, the definition of $V_{\xi,\rho,n}$, and the definition of \tilde{q}_n imply that $\pi_{\zeta,\rho'}(\tilde{q}_n) = 0$ for any $\zeta \in S^*M_0$ and $\rho' \in \widehat{\Gamma}_0$ such that $\Gamma(\zeta,\rho') \in \Xi_0$, that is, such that ρ' and α are Γ_0 -associated.

We next "quantize \tilde{q}_n " in an appropriate way, that is, we construct an operator $\widetilde{Q}_n \in B_{W_x}^{\Gamma}$ with symbol \widetilde{q}_n and with other convenient properties as follows. First, let $\chi \in \mathcal{C}_c^{\infty}(U_x)^{\Gamma_x}$ be such that $\chi \varphi_n = \varphi_n$, which is possible since φ_n has compact support. Then let $\psi \in \mathcal{C}^{\infty}(T_x^*M)^{\Gamma_x}$ be such that $\psi(0) = 0$ if $|\eta| < 1/2$ and $\psi(\eta) = 1$

whenever $|\eta| \geq 1$. Recall that in this proof $U_x \subset T_x M$ is identified with its image in $M = \Gamma \times_{\Gamma_x} U_x$ through the exponential map. Let for any symbol a

$$Op(a)f(y) := \int_{T_*^*M} \int_{U_x} e^{i(y-z)\cdot\eta} a(y,z,\eta) f(z) dz d\eta.$$

We shall use this for $a_n(y,z,\eta):=\chi(y)\psi(\eta)\tilde{q}_n\Big(\frac{\eta}{|\eta|}\Big)\chi(z),$ then set

$$Q_n := Op(a_n)$$
, that is

$$Q_n f(y) \, := \, \int_{T_x^* M} \int_{U_x} e^{i(y-z)\cdot \eta} \, \chi(y) \psi(\eta) \tilde{q}_n \left(\frac{\eta}{|\eta|}\right) \chi(z) f(z) dz d\eta$$

to be the standard pseudodifferential operator on U_x , associated to the symbol $a_n(y,z,\eta):=\chi(y)\psi(\eta)\tilde{q}_n\Big(\frac{\eta}{|\eta|}\Big)\chi(z)$. The operator Q_n is Γ_x -invariant by construction. Using the Frobenius isomorphism of Equation (14), we extend Q_n to the operator $\widetilde{Q}_n:=\Phi(Q_n)$, which acts on $M=W_x=\Gamma\times_{\Gamma_x}U_x$ (see also Equation (23) with regards to this isomorphism). Then $\widetilde{Q}_n\in\Psi^0(M;E)^\Gamma$, that is, it is Γ -invariant, by construction, and has principal symbol $\sigma_0(\widetilde{Q}_n)=\widetilde{q}_n$.

Now let $x_0 \in M_0 \cap U_x$, where, we recall, $M_0 := M_{(\Gamma_0)}$ denotes the principal orbit bundle. We have

$$L^{2}(W_{x_{0}}; E) = \operatorname{Ind}_{\Gamma_{0}}^{\Gamma} (L^{2}(U_{x_{0}}; \beta)) = L^{2}(U_{x_{0}}; \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta)),$$

where $\beta = E_{x_0} = E_x$ by the assumption that $E := \Gamma \times_{\Gamma_x} (U_x \times \beta)$.

Let $\beta_j \in \Gamma_0$ be the isomorphism classes of the Γ_{ξ} -submodules of β and $k_j \geq 0$ is the dimension of the corresponding β_j -isotypical component in β , so that $\beta \simeq \bigoplus_{j=1}^N \beta_j^{k_j}$, as Γ_0 -modules, as before. Thus

$$L^{2}(W_{x_{0}}; E) \simeq \bigoplus_{i=1}^{N} L^{2}(U_{x_{0}}; \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta_{j}^{k_{j}})).$$

Recall that the α -isotypical component of $\operatorname{Ind}_{\Gamma_0}^{\Gamma}(\beta_j^{k_j})$ is $\alpha \otimes \operatorname{Hom}_{\Gamma}(\alpha, \operatorname{Ind}_{\Gamma_0}^{\Gamma}(\beta_j^{k_j}))$, which is non-zero if, and only if, α and β_j are Γ_0 -associated, by the Frobenius isomorphism. Hence, passing to the α -isotypical components, we have

(38)
$$L^{2}(W_{x_{0}}; E)_{\alpha} := p_{\alpha}L^{2}(W_{x_{0}}; E) = \bigoplus_{j \in J^{c}} L^{2}(U_{x_{0}}; \operatorname{Ind}_{\Gamma_{0}}^{\Gamma}(\beta_{j}^{k_{j}}))_{\alpha},$$

where $J \subset \{1, ..., N\}$ is the set of indices such that $\beta_j \in \widehat{\Gamma}_0$ and α are Γ_0 -disjoint; J^c is its complement (i.e. $\beta_j \in \widehat{\Gamma}_0$ and α are Γ_0 -associated).

Let $p_J \in \text{End}(\beta)^{\Gamma_0}$ be the projector onto $\bigoplus_{j \in J^c} \beta_j^{k_j}$. Recall that $\pi_{\zeta,\beta_j}(\tilde{q}_n) = 0$ for any $(\zeta,\beta_j) \in S^*M_0 \times \widehat{\Gamma}_0$ with $j \notin J$. Therefore $\tilde{q}_n(\zeta)p_J = 0$, for all $\zeta \in S^*M_0$. Since S^*M_0 is dense in S^*M , this implies that $\tilde{q}_n p_J = 0$. Thus

$$\widetilde{Q}_n p_J \,=\, Op(\chi \psi \widetilde{q}_n \chi) p_J \,=\, Op(\chi \psi \widetilde{q}_n \chi p_J) \,=\, 0 \,. \label{eq:QnpJ}$$

Hence for any $f \in L^2(W_{x_0}; E)_{\alpha}$, we have that $\widetilde{Q}_n f = 0$. This is true for any $x_0 \in M_0$, so we conclude that \widetilde{Q}_n is zero on $L^2(M_0; E)_{\alpha}$. Since M_0 has measure zero complement in M, we have $L^2(M_0; E)_{\alpha} = L^2(M; E)_{\alpha}$; therefore $\pi_{\alpha}(\widetilde{Q}_n) = 0$. This implies that $\mathcal{R}_M(\widetilde{q}_n) = 0$, while $\pi_{\xi,\rho}(\widetilde{q}_n) = 1$. Thus $\Gamma(\xi,\rho) \notin \Xi$, which concludes the proof.

Our question now is to decide whether some given $\Gamma(\xi, \rho)$ is in Ξ or not. Recall that ρ and α are said to be Γ_0 -associated if $\operatorname{Hom}_{\Gamma_0}(\rho, \alpha) \neq 0$. The set $X_{M,\Gamma}^{\alpha}$ was defined in the introduction as the set of pairs $(\xi, \rho) \in T^*M \setminus \{0\} \times \widehat{\Gamma}_{\xi}$ for which there is an element $g \in \Gamma$ such that $g \cdot \rho$ and α are Γ_0 -associated.

Remark 3.18. Let us highlight the following interesting fact, implied by the proof of Theorem 3.17. We have that $E_{x\rho} = 0$ for any $(\xi, \rho) \in X_{M_0, \Gamma}^{\alpha}$ (with x the projection of ξ on M_0) if, and only if, $L^2(M; E)_{\alpha} = 0$.

Indeed, for any $x \in M_0$ with $\Gamma_x = \Gamma_0$, we have noted in Equation (38) that

$$L^2(W_x; E)_{\alpha} = \bigoplus_{\rho} L^2(U_x; \operatorname{Ind}_{\Gamma_0}^{\Gamma}(E_{x\rho}))_{\alpha},$$

where the direct sum is indexed by the representations $\rho \in \widehat{\Gamma}_0$ that are Γ_0 -associated to α . If $E_{x\rho} = 0$ for any such representation, then $L^2(W_x; E)_{\alpha} = 0$. Such open sets W_x cover M_0 , so $L^2(M_0; E)_{\alpha} = 0$. Since M_0 has measure zero complement, we conclude that $L^2(M; E)_{\alpha} = 0$.

Proposition 3.19. We use the notation in the last two paragraphs. We have $\Gamma(\xi, \rho) \in \Xi$ if, and only if, there is a $g \in \Gamma$ such that $g \cdot \rho$ and α are Γ_0 -associated.

Proof. Let $\Gamma(\xi, \rho) \in \operatorname{Prim}(A_M^{\Gamma})$, with $x \in M$ the base point of ξ . We can assume (by choosing a different element in the orbit if needed) that $\Gamma_0 \subset \Gamma_{\xi}$. Let $\tilde{q}_n \in A_M^{\Gamma}$ be the element defined in Remark 3.16 and $V_{\xi,\rho,n}$ the corresponding neighbourhood of $\Gamma(\xi, \rho)$ in $\operatorname{Prim}(A_M^{\Gamma})$.

There is a Γ_x -equivariant isomorphism $E|_{U_x} \simeq U_x \times \beta$, where $\beta = E_x$ is a Γ_x -module. Since $\Gamma_0 \subset \Gamma_x$, we may decompose β into Γ_0 -isotypical components, i.e. $\beta = \bigoplus_{j=1}^N \beta_j^{k_j}$, with the usual notation. If $\eta \in \mathcal{O}_n$, then $\pi_{\eta,\beta_j}(\tilde{q}_n) = \varphi_n(\eta)\pi_{\beta_j}(p_\rho)$. Therefore, for any $\eta \in S^*M$, we have

$$\pi_{n,\beta_i}(\tilde{q}_n) = 0 \Leftrightarrow \operatorname{Hom}_{\Gamma_0}(\beta_i, \rho) = 0 \text{ or } \tilde{q}_n(\eta) = 0.$$

This implies that

$$V_{\xi,\rho,n} \cap \Xi_0 = \{\Gamma(\eta,\beta) \in \Xi_0 \mid \tilde{q}_n(\eta) \neq 0 \text{ and } \operatorname{Hom}_{\Gamma_0}(\beta,\rho) \neq 0\}$$

It follows from the determination of Ξ_0 in Corollary 3.15 that $V_{\xi,\rho,n} \cap \Xi_0 \neq \emptyset$ if, and only if, we have $\operatorname{Hom}_{\Gamma_0}(\rho,\alpha) \neq 0$. Now $\Xi = \overline{\Xi}_0$ by Theorem 3.17. Since the open sets $(V_{\xi,\rho,n})_{n\in\mathbb{N}}$ form a basis of neighborhoods of $\Gamma(\xi,\rho)$, we conclude that $\Gamma(\xi,\rho) \in \Xi$ if, and only if, we have $\operatorname{Hom}_{\Gamma_0}(\rho,\alpha) \neq 0$.

Remark 3.20. Our definition of α -ellipticity for an operator $P \in \overline{\psi^0}(M; E)^{\Gamma}$ was stated in terms of the set $X_{M,\Gamma}^{\alpha}$, defined in Equation (7). Proposition 3.19 establishes that $\Xi \simeq \tilde{X}_{M,\Gamma}^{\alpha}/\Gamma$, where $\tilde{X}_{M,\Gamma}^{\alpha}$ is the (possibly smaller) subset of pairs $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$ such that $E_{x\rho} \neq 0$ (with x the projection of ξ on M). Keeping in mind the fact that the null operator on a trivial vector space is invertible, we have that $\sigma_0^{\Gamma}(P)(\xi, \rho)$ is invertible for any $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$ if, and only if, it is invertible for any $(\xi, \rho) \in \tilde{X}_{M,\Gamma}^{\alpha}$. The pathological case $\Xi = \emptyset$, for which $E_{x\rho} = 0$ for any $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$, causes no problem: indeed, as noticed in Remark 3.18, we then have $L^2(M; E)_{\alpha} = 0$. In that case $\pi_{\alpha}(P)$ is Fredholm for any $P \in \overline{\psi^0}(M; E)^{\Gamma}$, which is consistent with the invertibility of $\sigma_0^{\alpha}(P)(\xi, \rho) : 0 \to 0$ for any $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$.

We summarize part of the above discussions in the following proposition.

Proposition 3.21. Let $\tilde{X}_{M,\Gamma}^{\alpha}$ be as in Remark 3.20. The primitive ideal spectrum $\Xi = \operatorname{Prim}(A_M^{\Gamma}/\ker(\mathcal{R}_M))$ is canonically homeomorphic to $\tilde{X}_{M,\Gamma}^{\alpha}/\Gamma$ via the restriction map from $A_M^{\Gamma} := \mathcal{C}(S^*M; \operatorname{End}(E))^{\Gamma}$ to sections over $X_{M,\Gamma}^{\alpha}$.

4. Applications and extensions

We now prove the main result of the paper, Theorem 1.2, on the characterization of Fredholm operators, and discuss some extensions of our results. We first explain how to reduce the proof to the case M/Γ connected and we discuss in more detail the Γ -principal symbol and α -ellipticity (this discussion can be skipped at a first lecture).

4.1. Reduction to the connected case and α -ellipticity. In this subsection, unlike most of the rest of the paper, we do <u>not</u> assume that M/Γ is connected in order to explain how to reduce the general case to the connected one. We do assume however, as always, that M is compact. We also provide some other reductions of our proof.

Let $\pi_{M,\Gamma}: M \to M/\Gamma$ be the quotient map and let us write then $M/\Gamma = \bigcup_{i=1}^N C_i$ as the *disjoint* union of its connected components. We let $M_i := \pi_{M,\Gamma}^{-1}(C_i)$ be the preimages of these connected components. Note that, in general, the submanifolds M_i are not connected, but, for each i, $M_i/\Gamma = C_i$ is connected. In particular, a similar discussion applies to yield the definition of the space

$$(39) X_{M,\Gamma}^{\alpha} := \sqcup_{i=1}^{N} X_{M_{i},\Gamma}^{\alpha}$$

as a disjoint union of the spaces $X_{M_i,\Gamma}^{\alpha}$, which makes sense since each of the spaces M_i is invariant for Γ and M_i/Γ is connected. (See Equation (7) of the Introduction for the definition of the spaces $X_{M_i,\Gamma}^{\alpha}$.)

We shall decorate with the index i the restrictions of objects on M to M_i . Thus, $E_i := E|_{M_i}$, and so on and so forth. This almost works for an operator $P \in \overline{\psi^0}(M; E)^{\Gamma}$. Indeed, we first notice that

$$(40) L^2(M;E) \simeq \bigoplus_{i=1}^N L^2(M_i;E_i) \text{and} \bigoplus_{i=1}^N \overline{\psi^0}(M_i;E_i) \subset \overline{\psi^0}(M;E).$$

Recall that $\mathcal{K}(V)$ denotes the algebra of compact operators on a Hilbert space V. The following proposition provides the desired reduction to the connected case.

Proposition 4.1. Let $p_i: L^2(M; E) \to L^2(M_i; E_i)$ be the canonical orthoghonal projection. For $P \in \overline{\psi^0}(M; E)$, we let $P_i:=p_iPp_i \in \overline{\psi^0}(M_i; E_i)$. Then $P-\sum_{i=1}^N P_i \in \mathcal{K}(L^2(M; E))$. If we regard $\sum_{i=1}^N P_i = \bigoplus_{i=1}^N P_i$ as an element of $\bigoplus_{i=1}^N \overline{\psi^0}(M_i; E_i)$, then we see that

$$\overline{\psi^0}(M;E) = \bigoplus_{i=1}^N \overline{\psi^0}(M_i;E_i) + \mathcal{K}(L^2(M;E)).$$

Moreover, $\pi_{\alpha}(P) - \bigoplus_{i=1}^{N} \pi_{\alpha}(P_i)$ is compact and hence $\pi_{\alpha}(P)$ is Fredholm if, and only if, each $\pi_{\alpha}(P_i)$ is Fredholm, for i = 1, ..., N.

Proof. If $i \neq j$, $p_i P p_j$ has zero principal symbol, and hence it is compact. Therefore $P - \sum_{i=1}^{N} P_i = \sum_{i \neq j} p_i P p_j$ is compact. The rest follows from Equation (40), its corollary $L^2(M; E)_{\alpha} \simeq \bigoplus_{i=1}^{N} L^2(M_i; E_i)_{\alpha}$, and the fact that π_{α} respects these direct sum decompositions.

Recall the algebras $A_{\mathcal{O}}$ of symbols of the previous section, see Equation (25), where \mathcal{O} is an open subset of M.

Remark 4.2. The Γ -principal symbol $\sigma_m^{\Gamma}(P)$ was defined in (6), and we stress that the definition of the space $X_{M,\Gamma}$ did not require that M/Γ be connected. The disjoint union definition of the space $X_{M,\Gamma} = \bigsqcup_{i=1}^{N} X_{M_i,\Gamma}$ means that

$$\sigma_m^{\Gamma}(P)|_{X_{M-\Gamma}} = \sigma_m^{\Gamma}(P_i)$$

for each $i=1,\ldots,N$. The analogous disjoint union decomposition of $X_{M,\Gamma}^{\alpha}:=\bigcup_{i=1}^{N}X_{M_{i},\Gamma}^{\alpha}$ gives that P is α -elliptic if, and only if, for each i, P_{i} is α -elliptic.

This allows us to reduce the proof of our main theorem, Theorem 1.2 to the connected case since, assuming that the connected case has been proved, we have

$$\pi_{\alpha}(P)$$
 is Fredholm $\Leftrightarrow \forall i, \pi_{\alpha}(P_i)$ is Fredholm $\Leftrightarrow \forall i, P_i$ is α -elliptic $\Leftrightarrow P$ is α -elliptic,

where the first equivalence is by Proposition 4.1, the second equivalence is by the assumption that our main theorem has been proved in the connected case, and the last equivalence is by the first part of this remark.

We now resume our assumption that M/Γ is connected, for convenience. In particular, Γ_0 will be a minimal isotropy group, which is unique up to conjugacy (since we are again assuming that M/Γ is connected). We shall take a closer look next at the Γ - and α -principal symbols, so the following simple discussion will be useful. Recall that if $K \subset \Gamma$, $\rho \in \widehat{\Gamma}$, and $g \in \Gamma$, then $g \cdot K := gKg^{-1}$ and $(g \cdot \rho)(\gamma) := \rho(g^{-1}\gamma g)$, so that $g \cdot \rho$ is an irreducible representation of $g \cdot K$ (i.e. $g \cdot \rho \in \widehat{g \cdot K}$).

Remark 4.3. Let $\xi \in T^*M \setminus \{0\}$ and $\rho \in \widehat{\Gamma}_{\xi}$ (that is, $(\xi, \rho) \in X_{M,\Gamma}$). Then the following three statements are equivalent:

- (i) the pair $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$;
- (ii) there is $g \in \Gamma$ such that $\Gamma_0 \subset g \cdot \Gamma_{\xi} = \Gamma_{g\xi}$ and such that $g \cdot \rho$ and α are Γ_0 -associated;
- (iii) There is $\gamma \in \Gamma$ such that $\gamma \cdot \Gamma_0 \subset \Gamma_{\xi}$ and $\operatorname{Hom}_{\gamma \cdot \Gamma_0}(\rho, \alpha) \neq 0$.

Indeed, if (i) is satisfied, then the definition of $X_{M,\Gamma}^{\alpha}$ (see Equation (7) and Definition 2.2) is equivalent to the existence of g, i.e. (i) \Leftrightarrow (ii). Recalling that $g \cdot \rho \in \widehat{\Gamma_{g\xi}}$, we stress then that we need $\Gamma_0 \subset g \cdot \Gamma_{\xi} = \Gamma_{g\xi}$ for α and $g \cdot \rho$ to be associated.

To prove (ii) \Leftrightarrow (iii), let $\gamma = g^{-1}$. We have $\Gamma_0 \subset g \cdot \Gamma_{\xi}$ and $\operatorname{Hom}_{\Gamma_0}(g \cdot \rho, \alpha) \neq 0$ if, and only if, $\gamma \cdot \Gamma_0 \subset \Gamma_{\xi}$ and $\operatorname{Hom}_{\gamma \cdot \Gamma_0}(\rho, \gamma \cdot \alpha) \neq 0$. The result follows from the fact that α and $\gamma \cdot \alpha$ are equivalent (since $\gamma \in \Gamma$ and α is a representation of Γ).

We include next below, in Proposition 4.5, a reformulation of our α -ellipticity condition in terms of the fixed point manifold $S^*M^{\Gamma_0}$, with Γ_0 a minimal isotropy subgroup as before. This result was suggested by some discussions with P.-E. Paradan, whom we thank for his useful input.

In the following, $\operatorname{Stab}_{\Gamma}(M)$ will denote the set of stabilizer subgroups K of Γ , that is, the set of subgroups $K \subset \Gamma$ such that there is $m \in M$ with $K = \Gamma_m$. It is a finite set, since Γ is finite. Similarly, we let

$$\operatorname{Stab}_{\Gamma^0}^{\Gamma_0}(M) := \{ K \in \operatorname{Stab}_{\Gamma}(M) \mid \Gamma_0 \subset K \}.$$

Note that $\operatorname{Stab}_{\Gamma}(T^*M) = \operatorname{Stab}_{\Gamma}(M)$. Recall also that $(T^*M)^K = T^*(M^K)$, where M^K is the submanifold of fixed points of M by K, as usual. For a Γ -space X and $K \subset \Gamma$ a subgroup, we shall let $X_K := \{x \in X \mid \Gamma_x = K\} \subset X^K$ denote the set of points of X with isotropy K. Note that, in general, $T^*(M_K) \neq (T^*M)_{(K)}$.

Lemma 4.4. The set $M_K := \{m \in M \mid \Gamma_m = K\}$ is a submanifold.

Proof. Let $x \in M_K$, that is, $\Gamma_x = K$. The problem is local, so, using, [91, Proposition 5.13], we see that it suffices to consider the case $M = \Gamma \times_K V$, where V is a K-representation. Then, if $z = (\gamma, y) \in \Gamma \times_K V$, we have $\Gamma_z = \gamma K_y \gamma^{-1}$ and hence, if $\Gamma_z = K$, we obtain $K = \gamma K_y \gamma^{-1}$, which, in turn, gives $K_y = K$ and $\gamma \in N(K) := \{g \in \Gamma \mid gKg^{-1} = K\}$. We thus obtain that

$$M_K = \{(\gamma, y) \in \Gamma \times_K M \mid K_y = \gamma^{-1} K \gamma \} = N(K) \times_K V^K$$

which is a submanifold of M.

Let $K \subset \Gamma$ be a subgroup and $\rho \in \widehat{K}$. Then $E_{\rho} := \bigsqcup_{x \in M^K} E_{x\rho}$ is a smooth vector bundle over M^K , the set of fixed points of M with respect to K. Similarly, $(E \otimes \rho)^K \to M^K$ is a smooth vector bundle (over M^K). Moreover, we have an isomorphism

(41)
$$\operatorname{End}(E_{\rho})^{K} \simeq \operatorname{End}((E \otimes \rho)^{K} \otimes \rho)^{K} \simeq \operatorname{End}((E \otimes \rho)^{K}),$$

of vector bundles over M^K , where the last isomorphism comes from the fact that $\operatorname{End}(\rho)^K = \mathbb{C}$. In view of this discussion, we choose to state the following result in terms of the vector bundle $(E \otimes \rho)^K$ over M^K rather than in terms of E_ρ . This discussion shows also that it is enough in our proofs to assume that α is the trivial (one-dimensional) representation.

Proposition 4.5. Let $\alpha \in \widehat{\Gamma}$ and $P \in \psi^m(M; E)$, for some $m \in \mathbb{R}$. Recall the vector bundle $(M \otimes \rho)^K \to M^K \supset M_K$. The following are equivalent:

- (1) P is α -elliptic (Definition 1.1).
- (2) For all $K \in \operatorname{Stab}_{\Gamma}^{\Gamma_0}(M)$ and all $\rho \in \widehat{K}$ that are Γ_0 -associated with α , we have that $(\sigma_m(P) \otimes id_{\rho})|_{(E \otimes \rho)^K}$ defines by restriction an invertible element of

$$C^{\infty}((T^*M \setminus \{0\})_K, \operatorname{End}((E \otimes \rho)^K))$$
.

(3) The principal symbol $(\sigma_m(P) \otimes id_{\alpha})|_{(E \otimes \alpha)^{\Gamma_0}}$ defines by restriction an invertible element in

$$\mathcal{C}^{\infty}(T^*M^{\Gamma_0} \setminus \{0\}; \operatorname{End}((E \otimes \alpha)^{\Gamma_0}).$$

Recall that for representations α and β to be H-associated, they have to be defined, after restriction, on H. See Definition 2.2.

Proof. Recall that P is α -elliptic if the restriction of $\sigma_m^{\Gamma}(P)$ to $X_{M,\Gamma}^{\alpha}$ is invertible (see Remark 4.3 for a detailed definition and discussion of the space $X_{M,\Gamma}^{\alpha}$ appearing in the definition of α -ellipticity).

Let $K \in \operatorname{Stab}_{\Gamma}^{\Gamma_0}(M)$ (so $\Gamma_0 \subset K$), $\rho \in \widehat{K}$, and $\xi \in T_x^*M \setminus \{0\}$ with $\Gamma_{\xi} = K$. We have that $(\sigma_m(P) \otimes id_{\rho})|_{(E \otimes \rho)^K}$ is invertible at $\xi \in (T^*M)_K$ if, and only if, the restriction of $\sigma_m(P)(\xi)$ to $E_{x\rho}$ is invertible, since they correspond to each other under the isomorphism of Equation (41). The relation (2) thus means that the restriction of the principal symbol $\sigma_m(P)$ is invertible on a *subset* of $X_{M,\Gamma}^{\alpha}$, so (1) implies (2) right away.

Let us show next that (2) implies (1), let $\xi \in T^*M \setminus \{0\}$ and let $K' := \Gamma_{\xi}$. By definition, ξ belongs to $(T^*M)_{K'}$. Assume that $(\xi, \rho) \in X^{\alpha}_{M,\Gamma}$. This means that there exists $g \in \Gamma$ such that $\rho' := g \cdot \rho$ and α are Γ_0 associated (see Equation (7) and Definition 2.2; alternatively, this is also recalled in Remark 4.3). For this to make sense, it is implicit that

$$\Gamma_0 \subset \Gamma_{g\xi} = g \cdot \Gamma_{\xi} = g \cdot K' =: K$$

(again, see Remark 4.3). Then $g:(T^*M)_{K'}\to (T^*M)_K$ is a diffeomorphism. Condition (2) for the group K gives that $\pi_{g\xi,\rho'}(\sigma_m(P))$ is invertible, since the irreducible representation ρ' of $\Gamma_{g\xi}$ is Γ_0 -associated to α (we have used here again the isomorphism (41)). Furthermore, $g:E_{\xi,\rho}\to E_{g\xi,\rho'}$ is an isomorphism. Now, by the Γ -invariance of $\sigma:=\sigma_m(P)$, we have $(g^{-1}\sigma)(\xi)=g^{-1}(\sigma(g\xi))g=\sigma(\xi)$ therefore $\pi_{\xi,\rho'}(\sigma)$ is invertible if, and only if, $\pi_{g\xi,\rho}(\sigma)$ is.

For the equivalence of (1) and (3), we can assume that m=0. Recall first that the density of Ξ_0 in Ξ established in Theorem 3.17 gives that the family of representations

$$\mathcal{F}_0 := \{ \pi_{\xi,\rho} \mid (\xi,\rho) \in X_{M,\Gamma}^{\alpha}, \, \Gamma_{\xi} = \Gamma_0 \}$$

is faithful for the C^* -algebra $A_M^{\Gamma}/\ker(\mathcal{R}_M)$ (see e.g. [79, Theorem 5.1]). In other words, the restriction morphism

$$A_M^{\Gamma}/\ker(\mathcal{R}_M) \to \bigoplus_{\substack{\rho \in \widehat{\Gamma}_0, \\ \rho \subset \alpha|_{\Gamma_0}}} \mathcal{C}((T^*M \setminus \{0\})_{\Gamma_0}, \operatorname{End}(E_\rho)^{\Gamma_0})$$

is injective. Since $(T^*M)_{\Gamma_0}$ is dense in $T^*M^{\Gamma_0}$, it follows that the restriction morphism

$$R_M: A_M^{\Gamma}/\ker(\mathcal{R}_M) \to \bigoplus_{\substack{\rho \in \widehat{\Gamma}_0, \\ \rho \subset \alpha|_{\Gamma_0}}} \mathcal{C}\big(T^*M^{\Gamma_0} \setminus \{0\}, \operatorname{End}(E_\rho)^{\Gamma_0}\big)$$

is also injective.

Let us write $\alpha_{|\Gamma_0} = \bigoplus_{\rho \in \widehat{\Gamma}_0} m_\rho \rho$, with multiplicities $m_\rho \geq 0$. By considering the representations ρ with $m_\rho > 0$, we see that there is an injective vector bundle morphism over the manifold M^{Γ_0} defined by

(42)
$$\Psi: \bigoplus_{\substack{\rho \in \widehat{\Gamma}_{0}, \\ \rho \subset \alpha|_{\Gamma_{0}}}} \operatorname{End}(E_{\rho})^{\Gamma_{0}} \simeq \bigoplus_{\substack{\rho \in \widehat{\Gamma}_{0}, \\ \rho \subset \alpha|_{\Gamma_{0}}}} \operatorname{End}((E \otimes \rho)^{\Gamma_{0}}) \hookrightarrow \operatorname{End}((E \otimes \alpha)^{\Gamma_{0}}),$$

where the last morphism maps any element $T \in \text{End}((E \otimes \rho)^{\Gamma_0})$ to a direct sum of copies of T acting on the direct summand $[(E \otimes \rho)^{\Gamma_0}]^{m_\rho} \subset (E \otimes \alpha)^{\Gamma_0}$.

Condition (3) amounts to the fact that

$$\Psi(R_M(\sigma_0^{\Gamma}(P))) \in \mathcal{C}^{\infty}(T^*M^{\Gamma_0} \setminus \{0\}; \operatorname{End}((E \otimes \alpha)^{\Gamma_0})$$

is invertible. To establish that $(1) \Leftrightarrow (3)$, we thus need to prove that P is α -elliptic if, and only if, $\Psi(R_M(\sigma_0^{\Gamma}(P)))$ is invertible.

Recall the definition of the symbol algebras A_M from Equation (25). We have that $P \in \overline{\psi^0}(M; E)$ is α -elliptic if, and only if, the image of $\sigma_0^{\Gamma}(P)$ in the quotient algebra $A_M^{\Gamma}/\ker(\mathcal{R}_M)$ is invertible (by the determination of $\ker(\mathcal{R}_M)$ in Remark 3.20 or Proposition 3.21). But since both Ψ and R_M are injective, $\Psi \circ R_M$ is injective on $A_M^{\Gamma}/\ker(\mathcal{R}_M)$. Thus $\sigma_0^{\Gamma}(P)$ is invertible in the quotient algebra $A_M^{\Gamma}/\ker(\mathcal{R}_M)$

if, and only if, $\Psi(R_M(\sigma_0^{\Gamma}(P)))$ is invertible. As we have seen above, this amounts to $(1) \Leftrightarrow (3)$.

4.2. Fredholm conditions and Hodge and index theory. We continue to assume that M is a compact smooth manifold. We have the following Γ -equivariant version of Atkinson's theorem. (Recall that K(V) denotes the algebra of compact operators acting on the Hibert space V.)

Proposition 4.6. Let V be a Hilbert space with a unitary action of Γ and $P \in \mathcal{L}(V)^{\Gamma}$ be a Γ -equivariant bounded operator on V. We have that P is Fredholm if, and only if, it is invertible modulo $\mathcal{K}(V)^{\Gamma}$, in which case, we can choose the parametrix (i.e. the inverse modulo the compacts) to also be Γ -invariant.

Proof. See for example [12, Proposition 5.1].

Since $\pi_{\alpha}(\mathcal{K}(L^2(M;E))^{\Gamma}) = \mathcal{K}(L^2(M;E)_{\alpha})^{\Gamma}$, we obtain the following corollary.

Corollary 4.7. Let $P \in \overline{\psi^0}(M; E)^{\Gamma}$ and $\alpha \in \widehat{\Gamma}$. We have that $\pi_{\alpha}(P)$ is Fredholm on $L^2(M; E)_{\alpha}$ if, and only if, $\pi_{\alpha}(P)$ is invertible modulo $\pi_{\alpha}(\mathcal{K}(L^2(M; E))^{\Gamma})$ in $\pi_{\alpha}(\overline{\psi^0}(M; E)^{\Gamma})$.

We are now in a position to prove the main result of this paper, Theorem 1.2.

Proof of Theorem 1.2. As in [12, Section 2.6], we may assume that $P \in \overline{\psi^0}(M; E)^{\Gamma}$. Corollary 4.7 then states that $\pi_{\alpha}(P)$ is Fredholm if, and only if, the image of its symbol $\sigma(P)$ is invertible in the quotient algebra

$$\mathcal{R}_M(A_M^{\Gamma}) = \pi_{\alpha}(\overline{\psi^0}(M; E)^{\Gamma}) / \pi_{\alpha}(\mathcal{K}(L^2(M; E))^{\Gamma}).$$

According to Proposition 3.19 and Remark 3.20 following it, the primitive spectrum Ξ of $\mathcal{R}_M(A_M^{\Gamma})$ identifies with $X_{M,\Gamma}^{\alpha}$. Therefore $\mathcal{R}_M(\sigma(P))$ is invertible if, and only if, the endomorphism $\pi_{\xi,\rho}(\sigma(P))$ is invertible for all $(\xi,\rho) \in X_{M,\Gamma}^{\alpha}$, i.e. if, and only if, P is α -elliptic (see Definition 1.1).

Remark 4.8. Let $P: H^s(M; E) \to H^{s-m}(M; E)$ be an order m, classical pseudodifferential operator. Since the index of Fredholm operators is invariant under small perturbations and under compact perturbations, we obtain that the index of $\pi_{\alpha}(P)$ depends only on the homotopy class of its α -principal symbol $\sigma_m^{\alpha}(P)$.

An alternative approach to the Fredholm property (Theorem 1.2) can be obtained from the following theorem. Recall that $X_{M,\Gamma}^{\alpha}$ was defined in (7). Below, by ∂ we shall denote the connecting morphism in the six-term K-theory exact sequence associated to a short exact sequence of C^* -algebras. Recall that σ_0^{α} is the α -principal symbol map, see Definition 1.1.

Theorem 4.9. Let $X_{M,\Gamma}^{\alpha}(E) := X_{M,\Gamma}^{\alpha} \cap \Omega_M$ and let us denote by $\mathcal{C}(X_{M,\Gamma}^{\alpha}(E)/\Gamma)$ the algebra of restrictions of $A_M^{\Gamma} := \mathcal{C}(S^*M; \operatorname{End}(E))^{\Gamma}$ to $X_{M,\Gamma}^{\alpha}(E)/\Gamma$. Using the notation of Corollary 4.7, we have an exact sequence

$$0 \to \mathcal{K} \to \pi_{\alpha}(\overline{\psi^{0}}(M; E)^{\Gamma}) \xrightarrow{\sigma_{0}^{\alpha}} \mathcal{C}(X_{M,\Gamma}^{\alpha}(E)/\Gamma) \to 0.$$

Let $\partial: K_1(\mathcal{C}(X_{M,\Gamma}^{\alpha}(E)/\Gamma)) \to \mathbb{Z} \simeq K_0(\mathcal{K})$ be the associated connecting morphism and let $P \in \overline{\psi^0}(M; E)^{\Gamma}$ be such that $\pi_{\alpha}(P)$ is Fredholm. Then

$$\operatorname{ind}(\pi_{\alpha}(P)) = \dim(\alpha)\partial[\sigma_0^{\alpha}(P)].$$

Proof. The exactness of the sequence follows from the proof of Corollary 4.7 and the fact that $\mathcal{K}(L^2(M;E)_{\alpha})^{\Gamma} \simeq \mathcal{K}$, the algebra of compact operators on a model separable Hilbert space \mathcal{H} . Under this isomorphism, the resulting representation of \mathcal{K} on $L^2(M;E)_{\alpha}$ is isomorphic to $\dim(\alpha)$ times the standard representation of \mathcal{K} on \mathcal{H} . This justifies the factor $\dim(\alpha)$. The rest follows from the fact that the index is the connecting morphism in K-theory for the Calkin exact sequence. See [71] for more details.

Remark 4.10. As in [71], it follows that the index of $\pi_{\alpha}(P)$ with $P \in \psi^0(M; E)^{\Gamma}$ is the pairing between a cyclic cocycle ϕ on $\mathcal{C}^{\infty}(X_{M,\Gamma})$ (the algebra of principal symbols of operators in $\psi^0(M; E)^{\Gamma}$) and the K-theory class of the α -principal symbol of P [29]. See also [24, 48, 59, 55, 60, 93]. Lemma 3.8 gives that the restriction of this cyclic cocycle to the principal orbit bundle is the usual Atiyah-Singer cocycle (i.e. the cocycle that yields the Atiyah-Singer index theorem in cyclic homology [30, 56, 71, 75], which thus corresponds, after suitable rescaling, to the Todd class). The full determination of the class of the index cyclic cocycle ϕ require, however, a non-trivial use of cyclic homology, since the quotient algebra $\mathcal{C}^{\infty}(X_{M,\Gamma})$ is non-commutative, in general.

Remark 4.11. As for the case of compact complex varieties [41, 99], we can consider complexes of operators [19] and the corresponding notion of α -ellipticity. In particular, we obtain the finiteness of the corresponding cohomology groups if the complex is α -elliptic. This is related to the Hodge theory of singular spaces [2, 3, 16, 20, 27, 90].

- 4.3. Special cases. We now specialize our main result to some particular cases.
- 4.3.1. The abelian group case [12]. Many statements and definitions become easier in the case of abelian groups. In particular, if Γ_i , i=1,2, are both abelian, then the irreducible representations $\alpha_i \in \widehat{\Gamma}_i$ are characters, that is, morphisms $\alpha_i : \Gamma_i \to \mathbb{C}^*$, and we have that they are H-associated for some subgroup H if, and only if, $\alpha_1|_H = \alpha_2|_H$.

Let α be an irreducible representation of Γ . When Γ is abelian, the conjugacy class of isotropy subgroups corresponding to the principal orbit type of the action has only one element, namely Γ_0 . In that case, the set $X_{M,\Gamma}^{\alpha}$ defined in Equation (7) of the introduction has the simpler expression:

$$X^\alpha_{M,\Gamma} = \{(\xi,\rho) \mid \, \xi \in T^*M \smallsetminus \{0\}, \, \rho \in \widehat{\Gamma}_\xi, \rho|_{\Gamma_0} = \alpha|_{\Gamma_0} \, \}.$$

As a consequence, it is easier to check the α -ellipticity for an operator P in the abelian case. Let E, F be Γ -equivariant vector bundles over M and set $\alpha_0 := \alpha|_{\Gamma_0}$. We then recover the main result of [12]. Indeed, Theorem 1.2 can then be stated as follows:

Theorem 4.12. [12, Theorem 1.2] Let Γ be a finite, abelian group acting on a smooth, compact manifold M and let $P \in \psi^m(M; E, F)^{\Gamma}$. Then, for any $s \in \mathbb{R}$, the following are equivalent:

- (1) the operator $\pi_{\alpha}(P): H^{s}(M; E)_{\alpha} \to H^{s-m}(M; F)_{\alpha}$ is Fredholm,
- (2) for all $(x,\xi) \in T^*M \setminus \{0\}$, $\rho \in \widehat{\Gamma}_{\xi}$ such that $\rho_{|\Gamma_0} = \alpha_0$, the restriction of $\sigma(P)(x,\xi)$ defines an isomorphism

$$\pi_{\rho}(\sigma(P)(x,\xi)): E_{x\rho} \to F_{x\rho}$$
.

4.3.2. Scalar operators. Our main theorem becomes quite explicit when we are dealing with scalar operators, i.e. when the vector bundles $E_i = M \times \mathbb{C}$, where \mathbb{C} denotes the trivial representation of Γ .

Proposition 4.13. Let $P: H^s(M) \to H^{s-m}(M)$ be a Γ -invariant pseudodifferential operator. Let $\alpha \in \widehat{\Gamma}$. Then P is α -elliptic if, and only if, $\sigma(P)(\xi)$ is invertible for all $\xi \in T^*M \setminus \{0\}$ such that α is Γ_0 -associated to the trivial (constant 1) representation of Γ_{ξ} .

Proof. Let $\widehat{1}_{\Gamma_{\xi}}$ denote the trivial representation of Γ_{ξ} and let $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$. If $\rho \neq \widehat{1}_{\Gamma_{\xi}}$ then $\mathbb{C}_{\rho} = 0$ and then $\pi_{\rho}(\sigma(P)(\xi)) : 0 \to 0$ is invertible. Now if $\rho = \widehat{1}_{\Gamma_{\xi}}$ then $(\xi, \rho) \in X_{M,\Gamma}^{\alpha}$ if, and only if, α is Γ_{0} -associated to $\widehat{1}_{\Gamma_{\xi}}$.

4.3.3. Trivial actions. Assume that Γ acts trivially on M (in particular, M is then also connected). Our assumption implies that $\Gamma_0 = \Gamma_\xi = \Gamma$, for all $\xi \in T^*M \setminus \{0\}$. It follows that $\rho \in \widehat{\Gamma}_\xi$ is Γ_0 -associated to $\alpha \in \widehat{\Gamma}$ if, and only if, $\alpha = \rho$.

Let $E \to M$ be a Γ -equivariant vector bundle. For any $x \in M$, recall that we denote $E_{x\alpha}$ the α -isotypical component of E_x . Assuming M to be connected, we have that $E_{\alpha} = \bigcup_{x \in M} E_{x\alpha}$ is a Γ -equivariant sub-vector bundle of E. Our main result then becomes the following statement.

Proposition 4.14. Assume that Γ acts trivially on M and let $\alpha \in \widehat{\Gamma}$. Let E, F be two Γ -equivariant vector bundles over M and let $P \in \psi^m(M; E, F)^{\Gamma}$. Then for any $s \in \mathbb{R}$, the following are equivalent

- (1) $\pi_{\alpha}(P): H^{s}(M; E_{\alpha}) \to H^{s-m}(M; F_{\alpha})$ is Fredholm,
- (2) for all $(x,\xi) \in T^*M \setminus \{0\}$, the morphism

$$\pi_{\alpha}(\sigma(P)(x,\xi)): E_{x\alpha} \to F_{x\alpha}$$

is invertible,

(3) for all $(x,\xi) \in T^*M \setminus \{0\}$, the morphism

$$\sigma_m(P) \otimes \mathrm{id}_{\alpha^*}(x,\xi) : \mathrm{Hom}_{\Gamma}(\alpha,E_x) \to \mathrm{Hom}_{\Gamma}(\alpha,F_x)$$

is invertible.

Of course, the above result is nothing but the classical condition that the elliptic operator $p_{F_{\alpha}}Pp_{E_{\alpha}} \in \psi^{m}(M; E_{\alpha}, F_{\alpha})$ be Fredholm.

Proof. The equivalence between (1) and (2) is a direct consequence of Theorem 1.2. Let us check the equivalence of (1) and (3). First note that

$$(H^s(M, E) \otimes \alpha)^{\Gamma} = H^s(M, (E \otimes \alpha)^{\Gamma}),$$

since the action of Γ on M is trivial. The operator $\pi_{\alpha}(P)$ is Fredholm if, and only if, the pseudodifferential operator $P_{\alpha}: H^s(M, \operatorname{Hom}(\alpha, E)^{\Gamma}) \to H^{s-m}(M, \operatorname{Hom}(\alpha, F)^{\Gamma})$ defined for any $v^* \in \alpha^*$ and $s \in \mathcal{C}^{\infty}(M, E)$ by $P_{\alpha}(v^*s) = v^*Ps$ is Fredholm. Furthermore, the operator P_{α} is Fredholm if, and only if, it is elliptic, that is if, and only if, $\sigma_m(P) \otimes \operatorname{id}_{\alpha^*}(x, \xi) : \operatorname{Hom}_{\Gamma}(\alpha, E_x) \to \operatorname{Hom}_{\Gamma}(\alpha, F_x)$ is invertible for any $(x, \xi) \in T^*M \setminus \{0\}$. Note that the invertibility of $\sigma_m(P) \otimes \operatorname{id}_{\alpha^*}(x, \xi)$ is equivalent to the invertibility of $\pi_{\alpha}(\sigma_m(P)(x, \xi))$ by definition, which is consistent with (2). \square

4.3.4. Free action on a dense subset. As in the previous sections, the group Γ is finite and acts continuously on the manifold M. We consider vector bundles $E, F \to M$.

We have following corollary of the last few results in Section 3.

Corollary 4.15. Let us assume that Γ acts freely on a dense open subset of M. Then $\Xi = \operatorname{Prim}(A_M^{\Gamma})$.

Proof. The assumption on the action implies that $\Gamma_0 = \{1\}$. If $\xi \in T^*M \setminus \{0\}$ and $\rho \in \widehat{\Gamma}_{\xi}$, then ρ and α are always $\{1\}$ -associated. The Corollary then follows from Proposition 3.19.

Similarly, we have the following result.

Proposition 4.16. Assume that Γ acts freely on a dense subset in M, and let $P \in \psi^m(M; E, F)^{\Gamma}$. For any $\alpha \in \widehat{\Gamma}$, we have that P is α -elliptic if, and only if, P is elliptic.

Proof. It follows from Corollary 4.15 that $X_{M,\Gamma}^{\alpha} = X_{M,\Gamma}$. Thus the operator P_{α} is α -elliptic if, and only if, the sum $\bigoplus_{\rho \in \widehat{\Gamma}_{\xi}} \pi_{\rho}(\sigma_{m}(P)(\xi)) = \sigma_{m}(P)(\xi)$ is invertible for all $\xi \in T^{*}M \setminus \{0\}$, that is, if, and only if, P is elliptic. \square

- 4.4. Simonenko's localization principle. In this section, we obtain an equivariant version of Simonenko's principle [85]. In this subsection and the rest of the paper, we consider a compact Lie group G instead of Γ .
- 4.4.1. Simonenko's general principle. Let A be a unital C^* -algebra and $Z \simeq \mathcal{C}(\Omega_Z)$ a unital sub- C^* -algebra in A, i.e. $1_Z = 1_A$. An element $a \in A$ is said to have the strong Simonenko local property with respect to Z if, for every $\phi, \psi \in Z$ with compact disjoint supports, $\phi a \psi = 0$.

Lemma 4.17. The set $B \subset A$ of elements a satisfying the strong Simonenko local property is the set of elements of A commuting with Z.

Proof. We are going to show that the set of elements $a \in A$ with the strong Simonenko local type property is a C^* -algebra B containing Z and that every irreducible representation of B restricts to a scalar valued representation on Z, and hence that Z commutes with B.

Let us show first that B is a sub- C^* -algebra of A. Note that B is not empty since $Z \subset B$. To show that B is a sub- C^* -algebra, the only fact that is non-trivial to prove is that $ab \in B$, for all $a,b \in B$. Let ϕ and $\psi \in Z$ with disjoint compact supports and let θ be a function equal to 1 on $\operatorname{supp}(\psi)$ and 0 on $\operatorname{supp}(\phi)$, which exists by Urysohn's lemma. Then we have

(43)
$$\phi ab\psi = \phi a(\theta + 1 - \theta)b\psi = \phi a\theta b\psi + \phi a(1 - \theta)b\psi = 0,$$

since $\phi a\theta = 0$ and $(1 - \theta)b\psi = 0$.

Let $\pi: B \to \mathcal{L}(H)$ be an irreducible representation of B. First, let us show that for any $\phi, \psi \in Z$ with disjoint support, we either have $\pi(\phi) = 0$ or $\pi(\psi) = 0$. Indeed we have $\pi(\phi)\pi(a)\pi(\psi) = 0$ since $\phi a\psi = 0$, for any $a \in B$. Assume that $\pi(\psi) \neq 0$ then there is $\eta \in H$ such that $\pi(\psi)\eta \neq 0$. Now, π is irreducible so we get that the set $\{\pi(a)\pi(\psi)\eta, \ a \in B\}$ is dense in H. Thus $\pi(\phi) = 0$ on a dense subspace of H and so on H.

Assume now that $\pi(Z) \neq \mathbb{C}1_H$. Then there exist two distinct characters $\chi_0, \chi_1 \in \operatorname{Sp}(\pi(Z))$. Denote by $h_{\pi} : \operatorname{Sp}(\pi(Z)) \to \operatorname{Sp}(Z)$ the injective map adjoint to π , and choose $\phi, \psi \in \mathcal{C}(\operatorname{Sp}(Z))$ with disjoint supports such that $\phi(h_{\pi}(\chi_0)) = 1$ and $\psi(h_{\pi}(\chi_1)) = 1$. Then $\pi(\phi)(\chi_0) = 1$ and $\pi(\psi)(\chi_1) = 1$, which contradicts the fact that either $\pi(\phi) = 0$ or $\pi(\psi) = 0$.

Recall that a family $(\varphi_i)_{i\in I}$ of morphisms of a C^* -algebra A is said to be *exhaustive* if any primitive ideal contains some $\ker(\varphi_i)$, for a suitable $i \in I$ [72]. Then Remark 2.7 gives that the family of morphisms

$$\chi_{\omega}: A \to A/\omega A,$$

for $\omega \in \Omega_Z$, is exhaustive for A.

Definition 4.18. Denote by $\mathcal{H} = L^2(M)$. An operator $P \in \mathcal{L}(\mathcal{H})$ is said to be locally invertible at $x \in M$ if there are a neighbourhood V_x of x and operators Q_1^x and $Q_2^x \in \mathcal{L}(\mathcal{H})$ such that

(45)
$$Q_1^x P \phi = \phi \text{ and } \phi P Q_2^x = \phi, \text{ for any } \phi \in \mathcal{C}_c(V_x).$$

The operator P is said to be *locally invertible* if it is locally invertible at any $x \in M$.

Let $\Psi_M \subset \mathcal{L}(\mathcal{H})$ be the C^* -algebra of all $P \in \mathcal{L}(\mathcal{H})$ such that $\phi P \psi \in \mathcal{K}(\mathcal{H})$, for all $\phi, \psi \in \mathcal{C}(M)$ with disjoint support. We denote by \mathcal{B}_M the image of Ψ_M in the Calkin algebra $\mathcal{Q}(\mathcal{H}) := \mathcal{L}(\mathcal{H})/\mathcal{K}(\mathcal{H})$. We know by Lemma 4.17 that

$$\mathcal{B}_M = \{ P \in \mathcal{Q}(H) \mid \phi P = P\phi \text{ for all } \phi \in \mathcal{C}(M) \}.$$

Simonenko's principle is then [85]:

Proposition 4.19 (Simonenko's principle). If $P \in \Psi_M$, then P is locally invertible if, and only if, it is Fredholm.

We shall prove, in fact, a stronger version of this result, see Proposition 4.21.

4.5. Compact (non-finite) groups. We now allow for compact groups and try to see to what extent our results remain valid. To that end, we turn to an analog of Simonenko's principle. Let then G be a compact Lie group acting smoothly on M. We continue to study Fredholm conditions for $\pi_{\alpha}(P)$, $\alpha \in \widehat{G}$. Denote by $\mathcal{H} := L^2(M, E)$ and by \mathcal{H}_{α} the α -isotypical component associated to $\alpha \in \widehat{G}$.

Definition 4.20. We shall say that $P \in \mathcal{L}(\mathcal{H})$ is locally α -invertible at $x \in M$ if there are a G-invariant neighbourhood V_x of Γx and operators Q_1^x and $Q_2^x \in \mathcal{L}(\mathcal{H}_{\alpha})$ such that

(46)
$$Q_1^x \pi_\alpha(P) \phi = \phi \quad \text{and} \quad \phi \pi_\alpha(P) Q_2^x = \phi,$$

as operators on \mathcal{H}_{α} , for any $\phi \in \mathcal{C}(M)^G$ supported in V_x .

We denote by Ψ_M^G the G-invariant elements in the C^* -algebra Ψ_M , which was defined in the previous subsection.

Proposition 4.21 (Simonenko's equivariant principle). Let $P \in \Psi_M^G$. Then P is locally α -invertible if, and only if, $\pi_{\alpha}(P)$ is Fredholm.

Proof. We now use the results of the last section for $Z = \mathcal{C}(M)^G = \mathcal{C}(M/G)$. Let \mathcal{B}_M^{α} be the image of Ψ_M^G in the Calkin algebra $\mathcal{Q}(\mathcal{H}_{\alpha})$. We know from Lemma 4.17 that

$$\mathcal{B}_{M}^{\alpha} = \{ P \in \mathcal{Q}(\mathcal{H}_{\alpha}) \mid \phi P = P\phi, \ \forall \phi \in \mathcal{C}(M)^{G} \}.$$

Assume that P is locally α -invertible, i.e. $\forall x \in M$, there are a neighborhood V_x of Gx and operators $Q_1^x, Q_2^x \in \mathcal{L}(\mathcal{H}_{\alpha})$ such that $Q_1^x \pi_{\alpha}(P)\phi = \phi$ and $\phi \pi_{\alpha}(P)Q_2^x = \phi$, for any $\phi \in \mathcal{C}(M)^G$ supported in V_x . Let χ_x , be the family of representations of \mathcal{B}_G^{α} introduced in Equation (44). We use the same notation for $\pi_{\alpha}(P)$ and its image in $\mathcal{Q}(\mathcal{H}_{\alpha})$. We have that

$$\chi_x(Q_1^x \pi_\alpha(P)\phi) = \chi_x(Q_1^x)\chi_x(\pi_\alpha(P))\chi_x(\phi) = \chi_x(\phi).$$

Since $\chi_x(\phi) = 1$, we get:

$$\chi_x(Q_1^x)\chi_x(\pi_\alpha(P)) = 1.$$

And similarly,

$$\chi_x(\pi_\alpha(P))\chi_x(Q_2^x) = 1.$$

Therefore, $\chi_x(\pi_\alpha(P))$ is invertible for all x. Since the family χ_x is exhaustive, it follows that that $\pi_\alpha(P)$ is invertible in \mathcal{B}_M^α and so it is Fredholm.

Now assume that $\pi_{\alpha}(P)$ is Fredholm and let Q be an inverse modulo $\mathcal{K}(\mathcal{H}_{\alpha})$ for $\pi_{\alpha}(P)$, i.e. $\pi_{\alpha}(P)Q = id + K$ and $Q\pi_{\alpha}(P) = id + K'$, with $K, K' \in \mathcal{K}(\mathcal{H}_{\alpha})$. Using Proposition 4.6 and Lemma 2.10, we can assume that $K = \pi_{\alpha}(k)$ and $K' = \pi_{\alpha}(k') \in \mathcal{K}(\mathcal{H}_{\alpha}) = \pi_{\alpha}(\mathcal{K}(\mathcal{H})^{G})$, with $k, k' \in \mathcal{K}(\mathcal{H})^{G}$. Let $\chi \in \mathcal{C}(M)^{G}$ be equal to 1 on a G-invariant neighbourhood V_{x} of Gx and let $\phi \in \mathcal{C}(M)^{G}$ be supported in V_{x} then

$$\phi \chi \pi_{\alpha}(P)Q\chi = \phi \chi^2 + \phi \chi K \chi$$
 and $\chi \pi_{\alpha}(P)Q\chi \phi = \chi^2 \phi + \chi K' \chi \phi$.

Since ϕ is supported in V_x , we have $\phi \chi = \phi$ and so

$$\phi \pi_{\alpha}(P)Q\chi = \phi(1 + \chi K\chi)$$
 and $\pi_{\alpha}(P)Q\chi\phi = (1 + \chi K'\chi)\phi$.

As V_x becomes smaller and smaller, we have that χ converges strongly to 0. Since K is compact, we obtain that $\|\chi K\chi\| \to 0$. Thus, by choosing V_x small enough, we may assume that $\|\chi K\chi\| < 1$ and $\|\chi K'\chi\| < 1$.

It follows that $(1 + \chi K \chi)$ and $(1 + \chi K' \chi)$ are invertible and this implies

$$\phi \pi_{\alpha}(P) (Q\chi(1+\chi K\chi)^{-1}) = \phi$$
 and $((1+\chi K'\chi)^{-1}\chi Q)\pi_{\alpha}(P)\phi = \phi$,

i.e. P is locally α -invertible.

Corollary 4.22. Assume that M is compact, Γ is a finite group and M/Γ connected. Let $P \in \psi(M; E, F)^{\Gamma}$ and $\alpha \in \widehat{\Gamma}$. Then the following are equivalent:

- (1) $\pi_{\alpha}(P): H^{s}(M; E)_{\alpha} \to H^{s-m}(M; F)_{\alpha}$ is Fredholm for any $s \in \mathbb{R}$,
- (2) P is α -elliptic,
- (3) P is locally α -invertible.

Proof. The first equivalence is given by Theorem 1.2. Now since a finite group is compact Proposition 4.21 implies that (1) is equivalent to (3).

4.5.1. Transversally elliptic operators. Assume that M is a compact smooth manifold and that G is a compact Lie group acting on M. Denote by $\mathfrak g$ the Lie algebra of G. Then any $X \in \mathfrak g$ defines as usual the vector field X_M^* given by $X_M^*(m) = \frac{d}{dt}_{|t=0} e^{tX} \cdot m$. Denote by $\pi: T^*M \to M$ the canonical projection and let us introduce as in [6] the G-transversal space

$$T_G^*M := \{ \alpha \in T^*M \mid \alpha(X_M^*(\pi(\alpha))) = 0, \forall X \in \mathfrak{g} \}.$$

Recall that a G-invariant classical pseudodifferential operator P of order m is said G-transversally elliptic if its principal symbol is invertible on $T_G^*M \setminus \{0\}$ [6, 73].

We may now state the classical result of Atiyah and Singer [6, Corollary 2.5].

Theorem 4.23 (Atiyah-Singer [6]). Assume P is G-transversally elliptic. Then, for every irreducible representation $\alpha \in \widehat{G}$, $\pi_{\alpha}(P) : H^{s}(M; E_{0})_{\alpha} \to H^{s-m}(M; E_{1})_{\alpha}$, is Fredholm.

Note that this implies that Theorem 1.2 is not true anymore for if G is non-discrete. In particular, we obtain the following consequence of the localization principle.

Corollary 4.24. Assume that M is compact and that G is a compact Lie group and let $P \in \psi^m(M; E)^G$ be a G-transversally elliptic operator. Then P is locally α -invertible for any $\alpha \in \widehat{G}$, as in Definition 4.20.

Proof. Using Theorem 4.23 we obtain that $\pi_{\alpha}(P)$ is Fredholm. Therefore by Proposition 4.21 P is α -invertible.

References

- C. Akemann, G. Pedersen, and J. Tomiyama. Multipliers of C*-algebras. J. Functional Analysis, 13:277–301, 1973.
- [2] P. Albin. On the Hodge theory of stratified spaces. In Hodge theory and L²-analysis, volume 39 of Adv. Lect. Math. (ALM), pages 1–78. Int. Press, Somerville, MA, 2017.
- [3] P. Albin, E. Leichtnam, R. Mazzeo, and P. Piazza. Hodge theory on Cheeger spaces. J. Reine Angew. Math., 744:29–102, 2018.
- [4] I. Androulidakis and G. Skandalis. The analytic index of elliptic pseudodifferential operators on a singular foliation. J. K-Theory, 8(3):363–385, 2011.
- [5] I. Androulidakis and G. Skandalis. Pseudodifferential calculus on a singular foliation. J. Noncommut. Geom., 5(1):125-152, 2011.
- [6] M. Atiyah. Elliptic operators and compact groups. Lecture Notes in Mathematics, Vol. 401. Springer-Verlag, Berlin-New York, 1974.
- [7] M. Atiyah and I Singer. The index of elliptic operators III. Ann. of Math., 87:546–604, 1968.
- [8] K. Austin and J. Zhang. Limit operator theory for groupoids. arXiv preprint, to appear in Trans. AMS, 2018.
- [9] A. Baldare. The index of G-transversally elliptic families I. To appear in J. noncomm. Geometry. arXiv:1803.11148, 2018.
- [10] A. Baldare. The index of G-transversally elliptic families II. To appear in J. noncomm. Geometry. arXiv:1809.04819v2, 2018.
- [11] A. Baldare and M.-T. Benameur. The index of leafwise G-transversally elliptic operators on foliations. Arxiv Preprint, 2020.
- [12] A. Baldare, R. Côme, M. Lesch, and V. Nistor. Fredholm conditions for invariant operators: finite abelian groups and boundary valuel problems. Max Planck and Arxiv Preprint, 2019.
- [13] S. Beckus, D. Lenz, M. Lindner, and Ch. Seifert. On the spectrum of operator families on discrete groups over minimal dynamical systems. *Math. Z.*, 287(3-4):993–1007, 2017.
- [14] N. Berline and M. Vergne. The Chern character of a transversally elliptic symbol and the equivariant index. *Inventiones mathematicae*, 124(1):11–49, 1996.
- [15] N. Berline and M. Vergne. L'indice équivariant des opérateurs transversalement elliptiques. Inventiones mathematicae, 124(1):51–101, 1996.
- [16] M. Brion and M. Vergne. An equivariant Riemann-Roch theorem for complete, simplicial toric varieties. J. Reine Angew. Math., 482:67–92, 1997.
- [17] T. Bröcker and T. tom Dieck. Representations of compact Lie groups, volume 98 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995. Translated from the German manuscript, Corrected reprint of the 1985 translation.
- [18] J. Brüning and E. Heintze. Representations of compact Lie groups and elliptic operators. Invent. Math., 50(2):169–203, 1978/79.
- [19] J. Brüning and M. Lesch. Hilbert complexes. J. Funct. Anal., 108(1):88-132, 1992.
- [20] J. Brüning and M. Lesch. Kähler-Hodge theory for conformal complex cones. Geom. Funct. Anal., 3(5):439–473, 1993.
- [21] R. Busby. Double centralizers and extensions of C*-algebras. Trans. Amer. Math. Soc., 132:79–99, 1968.

- [22] A. P. Calderon and A. Zygmund. On the existence of certain singular integrals. Acta Math., 88:85–139, 1952.
- [23] A. P. Calderón and A. Zygmund. On a problem of Mihlin. Trans. Amer. Math. Soc., 78:209–224, 1955.
- [24] A. L. Carey, V. Gayral, A. Rennie, and F. A. Sukochev. Index theory for locally compact noncommutative geometries. *Mem. Amer. Math. Soc.*, 231(1085):vi+130, 2014.
- [25] C. Carvalho, R. Côme, and Yu Qiao. Gluing action groupoids: Fredholm conditions and layer potentials. Rev. Roumaine Math. Pures Appl., 64(2-3):113–156, 2019.
- [26] C. Carvalho, V. Nistor, and Yu Qiao. Fredholm conditions on non-compact manifolds: theory and examples. In Operator theory, operator algebras, and matrix theory, volume 267 of Oper. Theory Adv. Appl., pages 79–122. Birkhäuser/Springer, Cham, 2018.
- [27] J. Cheeger. On the Hodge theory of Riemannian pseudomanifolds. In Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), Proc. Sympos. Pure Math., XXXVI, pages 91–146. Amer. Math. Soc., Providence, R.I., 1980.
- [28] R. Côme. The Fredholm Property for Groupoids is a Local Property. Results in Mathematics, 74(4):160, August 2019.
- [29] A. Connes. Noncommutative geometry. Academic Press, San Diego, 1994.
- [30] A. Connes and H. Moscovici. The local index formula in noncommutative geometry. Geom. Funct. Anal., 5(2):174–243, 1995.
- [31] H. O. Cordes. The algebra of singular integral operators in Rⁿ. J. Math. Mech., 14:1007–1032, 1965.
- [32] H. O. Cordes and R. McOwen. The C^* -algebra of a singular elliptic problem on a noncompact Riemannian manifold. $Math.\ Z.,\ 153(2):101-116,\ 1977.$
- [33] A. de Monvel-Berthier and V. Georgescu. Graded C*-algebras in the N-body problem. J. Math. Phys., 32(11):3101–3110, 1991.
- [34] C. Debord, J.-M. Lescure, and F. Rochon. Pseudodifferential operators on manifolds with fibred corners. *Ann. Inst. Fourier (Grenoble)*, 65(4):1799–1880, 2015.
- [35] C. Debord and G. Skandalis. Pseudodifferential extensions and adiabatic deformation of smooth groupoid actions. Bull. Sci. Math., 139(7):750-776, 2015.
- [36] J. Dixmier. Les C*-algèbres et leurs représentations. Les Grands Classiques Gauthier-Villars. [Gauthier-Villars Great Classics]. Éditions Jacques Gabay, Paris, 1996. Reprint of the second (1969) edition.
- [37] V. Georgescu. On the structure of the essential spectrum of elliptic operators on metric spaces. J. Funct. Anal., 260(6):1734–1765, 2011.
- [38] V. Georgescu and A. Iftimovici. Crossed products of C*-algebras and spectral analysis of quantum Hamiltonians. Comm. Math. Phys., 228(3):519–560, 2002.
- [39] I.'T. Gokhberg. On an application of the theory of normed rings to singular integral equations. Uspekhi Matematicheskikh Nauk, 7(2):149–156, 1952.
- [40] I.'T. Gokhberg. On the theory of multidimensional singular integral equations. In *Doklady Akademii Nauk*, volume 133, pages 1279–1282. Russian Academy of Sciences, 1960.
- [41] Ph. Griffiths and J. Harris. Principles of algebraic geometry. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.
- [42] L. Hartmann, M. Lesch, and B. Vertman. On the domain of Dirac and Laplace type operators on stratified spaces. J. Spectr. Theory, 8(4):1295–1348, 2018.
- [43] B. Helffer and A. Mohamed. Caractérisation du spectre essentiel de l'opérateur de Schrödinger avec un champ magnétique. Ann. Inst. Fourier (Grenoble), 38(2):95–112, 1988.
- [44] L. Hörmander. Pseudo-differential operators. Comm. Pure Appl. Math., 18:501-517, 1965.
- [45] L. Hörmander. The analysis of linear partial differential operators. III. Classics in Mathematics. Springer, Berlin, 2007. Pseudo-differential operators, Reprint of the 1994 edition.
- [46] L. Jager. Pseudodifferential operators in infinite dimensional spaces: a survey of recent results. Rev. Roumaine Math. Pures Appl., 64(2-3):251–282, 2019.
- [47] P. Julg. Induction holomorphe pour le produit croisé dune C*-algebre par un groupe de Lie compact. CR Acad. Sci. Paris Sér. I Math, 294(5):193–196, 1982.
- [48] M. Karoubi. K-theory, volume 226 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin-New York, 1978.
- [49] G. Kasparov. Elliptic and transversally elliptic index theory from the viewpoint of KK-theory. Journal of Noncommutative Geometry, 10(4):1303–1378, 2016.

- [50] J. J. Kohn and L. Nirenberg. An algebra of pseudo-differential operators. Comm. Pure Appl. Math., 18:269–305, 1965.
- [51] B. Lange and V. Rabinovich. Pseudodifferential operators on \mathbb{R}^n and limit operators. Mat.Sb.(N.S.), 129(171(2)):175–185, 1986.
- [52] Y. Last and B. Simon. The essential spectrum of Schrödinger, Jacobi, and CMV operators. J. Anal. Math., 98:183–220, 2006.
- [53] R. Lauter, B. Monthubert, and V. Nistor. Pseudodifferential analysis on continuous family groupoids. Doc. Math., 5:625–655 (electronic), 2000.
- [54] R. Lauter and S. Moroianu. Fredholm theory for degenerate pseudodifferential operators on manifolds with fibered boundaries. Comm. Partial Differential Equations, 26:233–283, 2001.
- [55] M. Lesch, H. Moscovici, and M. Pflaum. Relative pairing in cyclic cohomology and divisor flows. J. K-Theory, 3(2):359–407, 2009.
- [56] M. Lesch, H. Moscovici, and M. Pflaum. Connes-Chern character for manifolds with boundary and eta cochains. Mem. Amer. Math. Soc., 220(1036):viii+92, 2012.
- [57] M. Lindner. Infinite matrices and their finite sections. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006. An introduction to the limit operator method.
- [58] M. Lindner and M. Seidel. An affirmative answer to a core issue on limit operators. J. Funct. Anal., 267(3):901–917, 2014.
- [59] J.-L. Loday and D. Quillen. Cyclic homology and the Lie homology of matrices. Comment. Math. Helv., 59:565–591, 1984.
- [60] Y. Manin. Quantum groups and noncommutative geometry. CRM Short Courses. Centre de Recherches Mathématiques, [Montreal], QC; Springer, Cham, second edition, 2018. With a contribution by Theo Raedschelders and Michel Van den Bergh.
- [61] R. Mazzeo and R. Melrose. Pseudodifferential operators on manifolds with fibred boundaries. Asian J. Math., 2:833–866, 1998.
- [62] S. T. Melo, R. Nest, and E. Schrohe. C*-structure and K-theory of Boutet de Monvel's algebra. J. Reine Angew. Math., 561:145–175, 2003.
- [63] R. Melrose and V. Nistor. K-theory of C*-algebras of b-pseudodifferential operators. Geom. Funct. Anal., 8(1):88–122, 1998.
- [64] R. B. Melrose. Geometric scattering theory. Stanford Lectures. Cambridge University Press, Cambridge, 1995.
- [65] S. G. Mihlin. Singular integral equations. Uspehi Matem. Nauk (N.S.), 3(3(25)):29-112, 1948.
- [66] S. G. Mihlin. On the theory of multidimensional singular integral equations. Vestnik Leningrad. Univ., 11(1):3–24, 1956.
- [67] B. Monthubert. Pseudodifferential calculus on manifolds with corners and groupoids. Proc. Amer. Math. Soc., 127(10):2871–2881, 1999.
- [68] J. Mougel. Essential spectrum, quasi-orbits and compactifications: application to the Heisenberg group. Rev. Roumaine Math. Pures Appl., 64(2-3):325–343, 2019.
- [69] M. Măntoiu and V. Nistor. Spectral theory in a twisted groupoid setting: Spectral decompositions, localization and fredholmness. to appear in Muenster J. Math, preprint 2018.
- [70] M. Măntoiu, R. Purice, and S. Richard. Spectral and propagation results for magnetic Schrödinger operators; a C^* -algebraic framework. J. Funct. Anal., 250(1):42–67, 2007.
- [71] V. Nistor. Higher index theorems and the boundary map in cyclic cohomology. Doc. Math., 2:263–295 (electronic), 1997.
- [72] V. Nistor and N. Prudhon. Exhausting families of representations and spectra of pseudodifferential operators. arXiv preprint arXiv:1411.7921, 2014.
- [73] P.-É. Paradan and M. Vergne. Index of transversally elliptic operators. Astérisque, (328):297–338 (2010), 2009.
- [74] P.-É. Paradan and M. Vergne. Equivariant Dirac operators and differentiable geometric invariant theory. Acta Math., 218(1):137–199, 2017.
- [75] D. Perrot. Pseudodifferential extension and Todd class. Adv. Math., 246:265–302, 2013.
- [76] V. Rabinovich, S. Roch, and B. Silbermann. Limit operators and their applications in operator theory, volume 150 of Operator Theory: Advances and Applications. Birkhäuser, 2004.
- [77] V. Rabinovich, B.-W. Schulze, and N. Tarkhanov. C*-algebras of singular integral operators in domains with oscillating conical singularities. Manuscripta Math., 108(1):69–90, 2002.
- [78] J. Renault. A groupoid approach to C*-algebras, volume 793 of LNM. Springer, 1980.

- [79] S. Roch. Algebras of approximation sequences: Structure of fractal algebras. In Singular Integral Operators, Factorization and Applications, volume 142 of Oper. Theory Adv. Appl., pages 287–310. Birkhäuser, Basel, 2003.
- [80] E. Schrohe. The symbols of an algebra of pseudodifferential operators. Pacific J. Math., 125(1):211–224, 1986.
- [81] E. Schrohe. Fréchet algebra techniques for boundary value problems on noncompact manifolds: Fredholm criteria and functional calculus via spectral invariance. *Math. Nachr.*, 199:145–185, 1999.
- [82] B.-W. Schulze. Boundary value problems and singular pseudo-differential operators. Wiley-Interscience Series in Pure and Applied Mathematics. Chichester: John Wiley & Sons., 1998.
- [83] R.T. Seeley. The index of elliptic systems of singular integral operators. Journal of Mathematical Analysis and Applications, 7(2):289 309, 1963.
- [84] J.-P. Serre. Linear representations of finite groups. Springer-Verlag, New York-Heidelberg, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.
- [85] I.B. Simonenko. A new general method for investigating linear operator equations of the singular integral operator type. i, ii. Izv. Akad. Nauk SSSR, Ser. Mat. 29:567–586, 1965.
- [86] I.M. Singer. Recent applications of index theory for elliptic operators. In Partial differential equations (Proc. Sympos. Pure Math., Vol. XXIII, Univ. California, Berkeley, Calif., 1971), pages 11–31, 1973.
- [87] M. Taylor. Gelfand theory of pseudo differential operators and hypoelliptic operators. Trans. Amer. Math. Soc., 153:495–510, 1971.
- [88] M. Taylor. Pseudodifferential operators, volume 34 of Princeton Mathematical Series. Princeton University Press, Princeton, N.J., 1981.
- [89] M. Taylor. Partial differential equations II. Qualitative studies of linear equations, volume 116 of Applied Mathematical Sciences. Springer, New York, second edition, 2011.
- [90] N. Teleman. Combinatorial Hodge theory and signature operator. Invent. Math., 61(3):227–249, 1980.
- [91] T. tom Dieck. Transformation groups, volume 8 of De Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, 1987.
- [92] F. Trèves. Introduction to pseudodifferential and Fourier integral operators. Vol. 1. Plenum Press, New York-London, 1980. Pseudodifferential operators, The University Series in Mathematics.
- [93] B. Tsygan. Homology of matrix Lie algebras over rings and Hochschild homology. Uspekhi Math. Nauk., 38:217–218, 1983.
- [94] E. Van Erp and R. Yuncken. A groupoid approach to pseudodifferential operators. http://arxiv.org/abs/1511.01041 [math.DG], 2015.
- [95] A. Vasy. Propagation of singularities in three-body scattering. Astérisque, (262):vi+151, 2000.
- [96] D. Voiculescu. A non-commutative Weyl-von Neumann theorem. Rev. Roumaine Math. Pures Appl., 21(1):97–113, 1976.
- [97] D. Voiculescu. Some results on norm-ideal perturbations of Hilbert space operators. J. Operator Theory, 2(1):3–37, 1979.
- [98] D.-V. Voiculescu. K-theory and perturbations of absolutely continuous spectra. Comm. Math. Phys., 365(1):363–373, 2019.
- [99] R. Wells, Jr. Differential analysis on complex manifolds, volume 65 of Graduate Texts in Mathematics. Springer, New York, third edition, 2008. With a new appendix by Oscar Garcia-Prode

E-mail address: alexandre.baldare@math.uni-hannover.de

Institut fur Analysis, Welfengarten 1, 30167 Hannover, Germany

 $E ext{-}mail\ address: remi.come@univ-lorraine.fr}$

Université Lorraine, 57000 Metz, France

Mathematisches Institut, Universität Bonn, Endenicher Allee 60, 53115 Bonn, Germany

 $E{-}mail~address{:}~ \verb|ml@matthias| esch.de, lesch@math.uni-bonn.de \\ URL{:}~ \verb|www.matthias| esch.de, www.math.uni-bonn.de/people/lesch \\ |$

 $E ext{-}mail\ address: nistor@univ-lorraine.fr}$

Université Lorraine, 57000 Metz, France

 $\mathit{URL} : \mathtt{http://www.iecl.univ-lorraine.fr/} \texttt{`Victor.Nistor}$