Dark solitons in the unitary Bose gas

M. Calzavara and L. Salasnich

Dipartimento di Fisica e Astronomia "Galileo Galilei", Universita di Padova, Via Marzolo 8, 35131, Padova, Italy

Abstract. We study the dilute and ultracold unitary Bose gas, characterized by a universal equation of state due to the diverging s-wave scattering length, under a transverse harmonic confinement. From the hydrodynamic equations of superfluids we derive an effective one-dimensional nonpolynomial Schrödinger equation (1D NPSE) for the axial wavefunction which, however, takes also into account the transverse wavefunction. Finally, by solving the 1D NPSE we obtain meaningful analytical formulas for the dark (gray and black) solitons of the bosonic system.

Keywords: Bose-Einstein condensate, unitary Bose gas, dark solitons, nonpolynomial Schrödinger equation.

1. Introduction

The possibility of tuning the interaction between atoms near Feshbach resonances [1] almost at will, by precision control of external magnetic fields, is allowing the experimenters to explore the physics of ultracold gases in a wide range of values of the coupling constant characterized by the s-wave scattering length. The possibility to control the scattering length via Feshbach resonances makes ultracold atomic gases an excellent setting for studies of strongly correlated behavior. In particular it made possible to observe first a Fermi gas [2–6] and then also a Bose gas at unitarity [7–10], that is the limit in which the s-wave scattering length is infinitely large. In this so-called unitary regime, the interactions are as strong as allowed by quantum mechanics, and the physics cannot explicitly depend on the scattering length, leading to the possibility of new types of universal behavior. In the case of a Bose gas, the presence of three body recombination processes [12] complicates the experimental realization of the unitary limit, which can be held for a limited amount of time. Nevertheless, the peculiar properties of a unitary gas, which behaves in a universal manner and exhibits scale invariance, encourage us to study the physics of such systems more in depth.

In this paper we investigate solitonic configurations of the unitary Bose gas. More specifically, we analyze axial and transverse density profiles of dark solitons in bosonic alkali-metal atoms at unitarity under the action of a transverse harmonic confinement. A dark soliton is a self-bound solitary wave, whose existence is made possible by the interplay between the repulsive interaction and the presence of a phase gradient. A dark soliton is characterized by a local decrease in density with respect to a uniform

background. The depth of the dark soliton crucially depends on the phase angle of the complex field which describes the Bose gas.

The realizability of black and gray solitons in ultracold gases is well documented in the weak-coupling limit [13,14] and in a unitary Fermi gas [15]. Moreover, the properties of black solitons have been theoretically investigated for superfluid fermions also in the BCS-BEC crossover [16–19]. It is then a natural question to ask, whether black and gray solitons can be realized also in the case of Bose gases at unitarity, and what are their specific properties. It is important to stress that dark (black and gray) solitons of the unitary Bose gas have not yet been observed experimentally nor theoretically analyzed. Our theoretical paper can help experimental groups because we are giving analytical and semi-analytical formulas for density profiles, widths, and velocities of these dark solitons. All these quantities are universal because, contrary to the dark solitons of weakly-interacting bosonic gases, they do not depend on the s-wave scattering length.

We first derive the equations which govern the dynamics of these systems. The resulting equation is a nonlinear Schrödinger equation for a complex order parameter $\psi(\mathbf{r},t) = |\psi(\mathbf{r},t)|e^{i\alpha(\mathbf{r},t)}$ [27], where $n(\mathbf{r},t) = |\psi(\mathbf{r},t)|^2$ is the local number density of the bosonic system and $\mathbf{v}(\mathbf{r},t) = (\hbar/m)\nabla\alpha(\mathbf{r},t)$ is the local velocity with $\alpha(\mathbf{r},t)$ the phase angle of the complex field. Then, we find the solutions which are in the form of dark solitons, specifically as stationary objects in a moving frame.

Mimicking the experimental practice of confining ultracold gases in the minimum of a potential, we study the case in which the Bose gas is placed in a harmonic potential with cylindrical symmetry, which is an approximation to a more realistic cigar shaped potential [13, 15]. A Gaussian variational approach, whose effectiveness in similar cases has already been proved [20–25], is then deployed to obtain an effective one-dimensional nonpolynomial Schrödinger equation (1D NPSE) for the axial wavefunction. The relevance of using the 1D NPSE in the study of solitons is due to the fact that the obtained solutions are a reliable generalization of familiar strictly one-dimensional results. These generalized solutions take properly into account that the transverse width of the cigar-shaped bosonic cloud depends on the axial coordinate.

In particular, we focus on the case of weak transverse confinement, in which more naive attempts to reduce the dimensionality of the problem fail. We integrate analytically the 1D NPSE thanks to the presence of a constant of motion, leaving behind a solution in terms of an integral, which we finally evaluate by means of numerical techniques. We are then able to find the axial density profile, transverse width and phase of the bosonic system. Moreover, we find the relation between density at the minimum and velocity of the soliton. While doing so, we also find that the theory actually depends on just one free parameter. Finally, we comment on the limits of the weak-coupling approximation.

2. From Euler to Schrödinger

Euler equations describe the dynamics of a non-viscous and irrotational fluid, such as a superfluid Bose gas at zero temperature [32]. In presence of an external potential $U(\mathbf{r})$, these equations are given by

$$\frac{\partial n}{\partial t} + \nabla \cdot (n\mathbf{v}) = 0 \tag{1}$$

$$m\frac{\partial \mathbf{v}}{\partial t} + \nabla(U(\mathbf{r}) + An^{\frac{2}{3}} + \frac{1}{2}m\mathbf{v}^{2}) = 0.$$
(2)

The equation of state of the unitary Bose gas at zero temperature is assumed to be

$$P(n) = \frac{2}{5} A n^{\frac{5}{3}},\tag{3}$$

where n is the density of particles and $A = u \frac{\hbar^2}{m}$, with u a universal and adimensional (positive) coefficient. In fact, we expect this to be the case in the unitary limit $a_s \to \infty$ because of dimensional analysis, taking into account that the only important length scale is the mean distance between atoms $\approx n^{-\frac{1}{3}}$. The value of $u = \xi(6\pi^2)^{\frac{2}{3}} \frac{5}{6}$ has been derived theoretically in many ways [26,28–31] and, depending on the procedure, values in the range $\xi \approx 0.4 \div 1.75$ have been reported.

We shall now derive a nonlinear Schrödinger equation (NLSE) with a 4/3 nonlinearity exponent [27], starting from equations (1)-(2) and adding a term, which will account for quantum effects. This is done via the mapping

$$n(\mathbf{r},t) = |\psi(\mathbf{r},t)|^2 \tag{4}$$

$$\mathbf{v}(\mathbf{r},t) = \frac{\hbar}{m} \nabla \alpha(\mathbf{r},t), \tag{5}$$

where $\alpha(\mathbf{r}, t)$ is a scalar field. Equation (5) is indeed fully justified by the fact the fluid is irrotational, i.e. $\nabla \wedge \mathbf{v} = \mathbf{0}$.

If we substitute (4)-(5) into (1)-(2) we obtain

$$\frac{\partial |\psi|}{\partial t} + \frac{\hbar}{2m} (\nabla^2 \alpha) |\psi| + \frac{\hbar}{m} \nabla \alpha \cdot \nabla |\psi| = 0$$
 (6)

$$\hbar \nabla \frac{\partial \alpha}{\partial t} + \nabla (U(\mathbf{r}) + An^{\frac{2}{3}} + \frac{\hbar^2}{2m} |\nabla \alpha|^2) = 0.$$
 (7)

Let us integrate equation (7) and multiply both sides by $|\psi|$

$$-|\psi|\hbar\frac{\partial\alpha}{\partial t} = |\psi|(U(\mathbf{r}) + An^{\frac{2}{3}} + \frac{\hbar^2}{2m}|\nabla\alpha|^2 + C). \tag{8}$$

The integration constant C appeared, but we shall see that it is not going to affect the dynamics. In order to get the Schrödinger equation we now add the term $-\frac{\hbar^2}{2m}\nabla^2|\psi(\boldsymbol{r},t)|$ on the right hand side of equation (8). This is chosen to obtain the same kinetic term which appears in the weak-coupling limit $P \propto n$. In that case we would obtain the Gross-Pitaevskii equation, whose validity for weak coupling is well established [33,34].

We can express the system of real equations as a single complex equation by adding equation (6) multiplied by $i\hbar$ to equation (8). Then we multiply every term by $e^{i\alpha}$, to

get

$$\left[i\hbar\frac{\partial|\psi|}{\partial t} - \hbar|\psi|\frac{\partial\alpha}{\partial t}\right]e^{i\alpha} = \left[-\frac{\hbar^2}{2m}(-|\psi||\nabla\alpha|^2 + \nabla^2|\psi| + i\nabla^2\alpha|\psi| + 2i\nabla\alpha\cdot\nabla|\psi|)\right]e^{i\alpha} + \left[(U(\mathbf{r}) + An^{\frac{2}{3}} + C)|\psi|\right]e^{i\alpha}.$$
(9)

Finally, by defining

$$\psi(\mathbf{r},t) = |\psi(\mathbf{r},t)| e^{i\alpha(\mathbf{r},t)}$$
(10)

last equation can be written as a NLSE

$$i\hbar\frac{\partial\psi}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\psi + (U(\mathbf{r}) + C)\psi + A|\psi|^{\frac{4}{3}}\psi. \tag{11}$$

C results in a potential energy offset. It can be removed by substituting $\psi \to e^{-\frac{\imath}{\hbar}Ct}\psi$. It is important to stress that equation (10) implies that the scalar field $\alpha(\mathbf{r},t)$ is an angle and the circulation of the velocity field $\mathbf{v}(\mathbf{r},t) = (\hbar/m)\nabla\alpha(\mathbf{r},t)$ is quantized.

3. Nonpolynomial Schrödinger equation

Equation (11) can also be derived minimizing the action functional $S = \int dt L$

$$S[\psi, \psi^{\star}] = \int_{t_1}^{t_2} dt \int d^3 \mathbf{r} \, \frac{i\hbar}{2} \left(\psi^{\star} \frac{\partial \psi}{\partial t} - \psi \frac{\partial \psi^{\star}}{\partial t} \right) - \frac{\hbar^2}{2m} |\nabla \psi|^2 - U(\mathbf{r}) |\psi|^2 - A|\psi|^{\frac{10}{3}}$$
(12)

with the constraint

$$\int d^3 \mathbf{r} |\psi(\mathbf{r}, t)|^2 = N. \tag{13}$$

Let us suppose that the confinement potential is harmonic, with cylindrical symmetry along the z axis,

$$U(\mathbf{r}) = \frac{1}{2}m\omega_{\perp}^2(x^2 + y^2) \tag{14}$$

and consider the following variational ansatz for ψ

$$\psi(\mathbf{r},t) = \phi(z,t) \frac{exp\left(-\frac{x^2+y^2}{2\sigma^2(z,t)}\right)}{\sqrt{\pi\sigma^2(z,t)}},$$
(15)

where the variational parameters are the axial wavefunction $\phi(z,t)$ and the width $\sigma(z,t)$ of the transverse Gaussian wavefunction. The choice of a Gaussian function in the (x,y) plane is justified by the presence of a harmonic potential in the (x,y) plane. Only in the strictly one-dimensional case one has $\sigma = a_{\perp}$ with $a_{\perp} = \sqrt{\hbar/(m\omega_{\perp})}$ the characteristic length of the transverse harmonic confinement.

By plugging equation (15) into the action in equation (12) we obtain

$$S[\phi, \sigma] = \int_{t_1}^{t_2} dt \int dz \, \frac{i\hbar}{2} \left(\phi^* \frac{\partial \phi}{\partial t} - \phi \frac{\partial \phi^*}{\partial t} \right) - \frac{\hbar^2}{2m} \left| \frac{\partial \phi}{\partial z} \right|^2$$

$$+ \int_{t_1}^{t_2} dt \int dz \, - \left[\frac{\hbar^2}{2m\sigma^2} + \frac{1}{2} m\omega_\perp^2 \sigma^2 \right] |\phi|^2 - \frac{3}{5} \frac{A}{(\sqrt{\pi}\sigma)^{\frac{4}{3}}} |\phi|^{\frac{10}{3}}$$

$$(16)$$

under the approximation of neglecting the space derivative of $\sigma(z,t)$. In the case of weakly-interacting bosons, which are very well described in three dimensions by the Gross-Pitaevskii equation (i.e. the 3D Schrödinger equation with cubic nonlinearity) this approximation has been found to be quite good also when $\sigma(z,t)$ depends strongly but monotonically on z [20, 25].

Minimizing the action with respect to ϕ with the constraint (13), we obtain

$$i\hbar\frac{\partial\phi}{\partial t} = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial z^2} + \frac{\hbar^2}{2m\sigma^2} + \frac{1}{2}m\omega_\perp^2\sigma^2 + \frac{A}{(\sqrt{\pi}\sigma)^{\frac{4}{3}}}|\phi|^{\frac{4}{3}} - \mu \right]\phi \tag{17}$$

and by doing the same with respect to σ

$$\frac{\sigma^4}{a_\perp^4} = 1 + \frac{4}{5} \frac{A}{\pi^{\frac{2}{3}}} \frac{m}{\hbar^2} |\phi|^{\frac{4}{3}} \sigma^{\frac{2}{3}},\tag{18}$$

where $a_{\perp} = \sqrt{\hbar/(m\omega_{\perp})}$. We shall refer to equation (17) endowed with (18) as the nonpolynomial Schrödinger equation [20–25].

Let us rewrite the equations in terms of adimensional quantities. We can do that with the substitutions

$$\begin{split} z &\to z a_\perp \\ t &\to \frac{t}{\omega_\perp} \\ \sigma(z,t) &\to \sigma(z,t) a_\perp \\ \phi(z,t) &\to \sqrt{\frac{N}{a_\perp}} \phi(z,t) \\ \mu &\to \hbar \omega_\perp (\mu+1) \\ A &\to A \frac{\hbar^2}{2m}. \end{split}$$

Equation (17) then becomes

$$i\frac{\partial\phi}{\partial t} = \left[-\frac{1}{2}\frac{\partial^2}{\partial z^2} + \frac{1}{2\sigma^2} + \frac{1}{2}\sigma^2 - 1 + \frac{A}{2}\frac{N^{\frac{2}{3}}}{\pi^{\frac{2}{3}}}\frac{|\phi|^{\frac{4}{3}}}{\sigma^{\frac{4}{3}}} - \mu \right]\phi \tag{19}$$

while equation (18) becomes

$$\sigma^4 = 1 + \frac{2}{5} \frac{A}{\pi^{\frac{2}{3}}} N^{\frac{2}{3}} |\phi|^{\frac{4}{3}} \sigma^{\frac{2}{3}}.$$
 (20)

Remember that in this equation σ is adimensional: it is in units of the characteristic length a_{\perp} of the transverse harmonic confinement. In equation (20) one can identify two regimes: a regime of strong transverse confinement where $\sigma \simeq 1$ and consequently the system is truly one-dimensional, but also a regime of weak transverse confinement where $\sigma \gg 1$. Thus, the regime of weak transverse confinement is obtained by neglecting the first addend on the right hand side of equation (20), supposing that $AN^{\frac{2}{3}}|\phi|^{\frac{4}{3}} \gg 1$. This gives rise to

$$\sigma = \left(\frac{2}{5}A\right)^{\frac{3}{10}} \left(\frac{N}{\pi}\right)^{\frac{1}{5}} |\phi|^{\frac{2}{5}},\tag{21}$$

which plugged back in equation (19) yields the following equation for ϕ

$$i\frac{\partial\phi}{\partial t} = \left[-\frac{1}{2}\frac{\partial^2}{\partial z^2} + \frac{1}{2}\left(\frac{5}{2A}\right)^{\frac{3}{5}} \left(\frac{\pi}{N}\right)^{\frac{2}{5}} |\phi|^{-\frac{4}{5}} + \frac{7}{10}\left(A\right)^{\frac{3}{5}} \left(\frac{5}{2}\frac{N}{\pi}\right)^{\frac{2}{5}} |\phi|^{\frac{4}{5}} - \mu - 1 \right] \phi.(22)$$

In the same approximation we can neglect the term $\propto (AN^{\frac{2}{3}}|\phi|^{\frac{4}{3}})^{-\frac{3}{5}}$, and obtain

$$i\frac{\partial\phi}{\partial t} = \left[-\frac{1}{2}\frac{\partial^2}{\partial z^2} + \gamma|\phi|^{\frac{4}{5}} - \mu \right]\phi,\tag{23}$$

where we defined

$$\gamma = \frac{7}{10} A^{\frac{3}{5}} \left(\frac{5}{2} \frac{N}{\pi} \right)^{\frac{2}{5}}.$$
 (24)

Finally, let us express also σ in terms of γ

$$\sigma = \sqrt{\frac{4}{7}}\sqrt{\gamma}|\phi|^{\frac{2}{5}}.\tag{25}$$

It will prove useful to study also the limit of strong confinement, that is instead given by $\sigma = 1$:

$$i\frac{\partial\phi}{\partial t} = \left[-\frac{1}{2}\frac{\partial^2}{\partial z^2} + \gamma_{SC}|\phi|^{\frac{4}{3}} - \mu \right]\phi. \tag{26}$$

$$\gamma_{SC} = \frac{A}{2} \left(\frac{N}{\pi} \right)^{\frac{2}{3}} = \frac{5}{4} \left(\frac{4}{7} \gamma \right)^{\frac{5}{3}}.$$
 (27)

4. Dark solitons in the 1D NPSE

A soliton is a solution to equation (23) of the form

$$\phi(z,t) = f(z - vt)e^{i\alpha(z,t)}, \tag{28}$$

where $f(\zeta) \geq 0$ and $\alpha(z,t)$ are real functions. Once we substitute (28) in (23) we obtain the equations for f and α

$$f'' = 2\left[f\left(\frac{\partial\alpha}{\partial t} + \frac{1}{2}\left(\frac{\partial\alpha}{\partial z}\right)^2 - \mu\right) + \gamma f^{\frac{9}{5}}\right],\tag{29}$$

$$f'v = \left(f'\frac{\partial\alpha}{\partial z} + \frac{1}{2}f\frac{\partial^2\alpha}{\partial z^2}\right). \tag{30}$$

Upon inspection of equation (30), we find that the right hand side should depend only on ζ , so $\frac{\partial \alpha}{\partial z}$ is only function of ζ . Let us assume the following

$$\alpha(z,t) = \theta(z - vt) + \beta(t). \tag{31}$$

If we plug this back into equation (29)-(30) we get

$$f'' = 2\left[f\left(\frac{d\beta}{dt} - v\theta' + \frac{1}{2}(\theta')^2 - \mu\right) + \gamma f^{\frac{9}{5}}\right],\tag{32}$$

$$f'v = \left(f'\theta' + \frac{1}{2}f\theta''\right). \tag{33}$$

4.1. Phase

Let us solve equation (33) in order to obtain θ' in terms of f. We can rewrite it as

$$v\frac{(f^2)'}{f} = \frac{(f^2\theta')'}{f},$$
 (34)

for $f(\zeta) \neq 0$. After integration with respect to ζ we have

$$f^{2}(\zeta)\left(v - \theta'(\zeta)\right) = D,\tag{35}$$

where D is an integration constant. If we assume that

$$\lim_{\zeta \to +\infty} |f(\zeta)| = f_{\infty},\tag{36}$$

$$\lim_{\zeta \to \pm \infty} \theta'(\zeta) = 0, \tag{37}$$

and we get $D = v f_{\infty}^2$. The expression for θ' we are looking for is then

$$\theta'(\zeta) = v \left(1 - \frac{f_{\infty}^2}{f^2(\zeta)} \right), \tag{38}$$

valid for $f(\zeta) \neq 0$.

4.2. Modulus

Let us plug equation (38) back into equation (32)

$$f'' = 2\frac{d\beta}{dt}f - v^2f + v^2\frac{f_\infty^4}{f^3} - 2\mu f + 2\gamma f^{\frac{9}{5}}.$$
 (39)

We can choose β in order to get simpler equations. Let us suppose

$$\beta(t) = \frac{v^2}{2}t,\tag{40}$$

so that equation (39) becomes

$$f'' = v^2 \frac{f_\infty^4}{f^3} - 2\mu f + 2\gamma f^{\frac{9}{5}}.$$
 (41)

Equation (41) can be written in the form of a Newton equation for a conservative force field

$$f'' = -\frac{dW}{df},\tag{42}$$

$$W = \frac{1}{2}v^2 \frac{f_\infty^4}{f^2} + \mu f^2 - \frac{5}{7}\gamma f^{\frac{14}{5}}.$$
 (43)

The total energy K is then a conserved quantity for all ζ

$$K = \frac{1}{2}f^{2} + W, (44)$$

and energy balance gives us the following first order equation

$$K = \frac{1}{2}f'^2 + \frac{1}{2}v^2\frac{f_\infty^4}{f^2} + \mu f^2 - \frac{5}{7}\gamma f^{\frac{14}{5}},\tag{45}$$

which is satisfied by the two branches

$$f' = \pm \frac{1}{|f|} \sqrt{2Kf^2 - v^2 f_\infty^4 - 2\mu f^4 + \frac{10}{7} \gamma f^{\frac{24}{5}}}.$$
 (46)

This equation can be put in the integral form with the separation of variables, giving

$$|\zeta - \zeta_0| = \int_{f(\zeta_0)}^{f(\zeta)} df \frac{f}{\sqrt{2Kf^2 - v^2 f_\infty^4 - 2\mu f^4 + \frac{10}{7}\gamma f^{\frac{24}{5}}}},$$
(47)

where, since the integral is positive, we have chosen the + sign when $\zeta - \zeta_0 \ge 0$ and the - sign otherwise. Therefore, $f(\zeta) = f(|\zeta - \zeta_0|)$ has even parity with respect to reflections around ζ_0 , and we just need to study the case $\zeta - \zeta_0 \ge 0$.

We have now to find the values of the parameters μ, K, f_{∞} . This can be done by choosing boundary conditions for f.

4.3. Black solitons

Let us find an odd parity solution (which features a node in the origin for the density n) with a positive horizontal asymptote at $+\infty$. We set $\zeta_0 = 0$, and since f is even, the phase θ will account for the change of sign. Then, the following boundary conditions are needed

$$f(0) = 0, \quad f(+\infty) = f_{\infty}, \quad f'(+\infty) = f''(+\infty) = 0.$$
 (48)

This is possible only if v = 0 ($\zeta = z$) in order to get rid of the term in equation (45) which diverges as $f \to 0$. The condition $f''(+\infty) = 0$ applied to equation (41) gives

$$\mu = \gamma f_{\infty}^{\frac{4}{5}},\tag{49}$$

while $f'(+\infty) = 0$ applied to (45) gives

$$K = \frac{2}{7} \gamma f_{\infty}^{\frac{14}{5}}.\tag{50}$$

Equation (47) then becomes

$$z = \int_0^{f(z)} \frac{df}{\sqrt{\frac{4}{7}\gamma f_{\infty}^{\frac{14}{5}} - 2\gamma f_{\infty}^{\frac{4}{5}} f^2 + \frac{10}{7}\gamma f_{\infty}^{\frac{14}{5}}}}.$$
 (51)

It is possible to get rid of f_{∞} with the rescaling

$$h(z) = \frac{f(z)}{f_{\infty}} \tag{52}$$

and the parameter

$$\delta = \gamma f_{\infty}^{4/5},\tag{53}$$

while σ does not change: $\sigma(z) = \sqrt{\frac{4}{7}} \sqrt{\gamma} f(z)^{\frac{2}{5}} = \sqrt{\frac{4}{7}} \sqrt{\delta} h(z)^{\frac{2}{5}}$. We obtain

$$z = \frac{1}{\sqrt{\delta}} \int_0^{h(z)} \frac{dh}{\sqrt{\frac{4}{7} - 2h^2 + \frac{10}{7}h^{\frac{14}{5}}}}$$
 (54)




Figure 1. Black soliton. Normalized axial density profile $\rho(z) = h^2(z) = (f(z)/f_{\infty})^2$ and transverse width profile $\sigma(z)$ vs axial coordinate z, for three values of the parameter $\delta = \gamma |f_{\infty}|^{4/5}$, with γ given by equation (24) and f_{∞} the bulk value of the soliton wavefunction.

Plots of the numerical integration of equation (54) are shown in figure 1, for three values of δ . Accordingly with those integrations, also $\sigma(z)$ is computed and shown in the plot.

We may define $\tilde{z} = \sqrt{\delta}z$ and $\tilde{\sigma} = \frac{\sigma}{\sqrt{\delta}}$, so that the theory has no free parameters. This way there is only one plot to make, but we want to explicitly show in the plots the effects of changing the interaction, so for the moment we are keeping z and σ .

4.4. Gray solitons

Let us find an even parity solution with a positive horizontal asymptote. We set $\zeta_0 = 0$ once again, and require the following boundary conditions

$$f'(0) = 0, \quad f(+\infty) = f_{\infty}, \quad f'(+\infty) = f''(+\infty) = 0.$$
 (55)

The condition $f''(+\infty) = 0$ applied to equation (41) gives

$$\mu = \frac{1}{2}v^2 + \gamma f_{\infty}^{\frac{4}{5}},\tag{56}$$

 $f'(+\infty) = 0$ applied to equation (45) gives

$$K = f_{\infty}^2 \left[v^2 + \frac{2}{7} \gamma f_{\infty}^{\frac{4}{5}} \right], \tag{57}$$

while by equating K inside equation (45) evaluated at 0 and at ∞ , applying f'(0) = 0 we get a relation between v, γ and f_0

$$\frac{v^2}{\gamma} = \frac{2}{7} f_0^2 \frac{7f_\infty^{\frac{4}{5}} f_0^2 - 5f_0^{\frac{14}{5}} - 2f_\infty^{\frac{14}{5}}}{2f_\infty^2 f_0^2 - f_\infty^4 - f_0^4}.$$
 (58)

Equation (47) then becomes

$$\zeta = \frac{1}{\sqrt{\gamma}} \int_{f_0}^{f(\zeta)} \frac{df \ f}{\sqrt{\frac{10}{7} f^{\frac{24}{5}} - \frac{v^2}{\gamma} f^4 - 2f_{\infty}^{\frac{4}{5}} f^4 + \frac{v^2}{\gamma} 2f_{\infty}^2 f^2 + \frac{4}{7} f_{\infty}^{\frac{14}{5}} f^2 - \frac{v^2}{\gamma} f_{\infty}^4}}. (59)$$

Also in this case we can get rid of f_{∞} in the same way as before, obtaining

$$\left(\frac{v}{v_s}\right)^2 = \frac{5}{7}h_0^2 \frac{5h_0^{\frac{14}{5}} - 7h_0^2 + 2}{(1 - h_0^2)^2}.$$
(60)

$$\zeta = \frac{1}{\sqrt{\delta}} \sqrt{\frac{7}{2}} \int_{h_0}^{h(\zeta)} \frac{dh \ h}{\sqrt{2h^2 - 7h^4 + 5h^{\frac{24}{5}} - (2h_0^2 - 7h_0^4 + 5h_0^{\frac{24}{5}})\frac{(1-h^2)^2}{(1-h_0^2)^2}}}$$
(61)

where of course $h_0 = \frac{f_0}{f_{\infty}}$ and $v_s = \sqrt{\frac{2}{5}\delta}$ is the speed of sound, i.e. the propagation velocity of a infinitesimal perturbation in the fluid density.

Now the theory has two free parameters, namely h_0 and δ , while v is given in terms of them. In order to be more concise, we now define $\tilde{\zeta} = \sqrt{\delta}\zeta$, $\tilde{\sigma} = \frac{\sigma}{\sqrt{\delta}}$ and

$$\tilde{v} = \frac{v}{v_s},\tag{62}$$

so that we are left with a theory that depends only on h_0 (or \tilde{v}). The scaled velocity \tilde{v} ranges from 0 (black soliton) to 1 (sound wave).

Plots of the numerical integration of equation (61) are shown in figure 2, for three values of \tilde{v} . Accordingly with those integrations, also $\tilde{\sigma}(\tilde{\zeta})$ is computed and shown in the plot.

4.5. Phase

In the case of the black soliton, equation (38) becomes $\theta' = 0$. Since we want the solution to be odd, we choose

$$\theta(z) = -sgn(z)\frac{\pi}{2}. (63)$$

In the general case, we can now compute h and substitute inside equation (38) to obtain θ' . Let us express also θ in terms of $\tilde{\zeta}$:

$$\theta(\zeta) = \theta(\frac{\tilde{\zeta}}{\sqrt{\delta}}) =: \tilde{\theta}(\tilde{\zeta}) \tag{64}$$

it follows that

$$\theta'(\zeta) = \sqrt{\delta}\tilde{\theta}'(\tilde{\zeta}),\tag{65}$$




Figure 2. Gray soliton. Upper panel: Scaled axial density profile $\rho(\tilde{\zeta}) = h(\tilde{\zeta})^2 = (f(z)/f_{\infty})^2$ vs scaled comoving axial coordinate $\tilde{\zeta} = (z - vt)\sqrt{\delta}$. Lower panel: Scaled transverse width $\tilde{\sigma}(\tilde{\zeta})$ vs $\tilde{\zeta}$. Notice that $\delta = \gamma |f_{\infty}|^{4/5}$, with γ given by equation (24) and f_{∞} the bulk value of the soliton wavefunction. $\tilde{v} = v/v_s$ is the velocity rescaled by the sound velocity v_s .

therefore

$$\tilde{\theta}'(\tilde{\zeta}) = \sqrt{\frac{2}{5}}\tilde{v}\left(1 - \frac{1}{h^2(\tilde{\zeta})}\right). \tag{66}$$

Last equation upon integration yields $\tilde{\theta}$. A plot of the numerical integration of equation (66) is shown in figure 3 for four values of \tilde{v} , including $\tilde{v} = 0$, which is the black soliton case.

5. Weak vs strong transverse confinement

In the previous sections we have mainly investigated the regime of weak transverse confinement where $\sigma \gg 1$. See the discussion of equation (20). Let us now briefly analyze some properties of the regime of strong transverse confinement where $\sigma \simeq 1$. In this case the system is truly one dimensional and the equations are much simpler.

By performing computations which are very close to the ones we have already shown in the weak confinement case, one can derive from (26) that in the case of a black soliton the following relation holds

$$z = \frac{1}{\sqrt{\delta_{SC}}} \int_0^{h(z)} \frac{dh}{\sqrt{\frac{4}{5} - 2h^2 + \frac{6}{5}h^{\frac{10}{3}}}},\tag{67}$$




Figure 3. Scaled phase $\tilde{\theta}(\tilde{\zeta})$ of gray $(\tilde{v}>0)$ and black $(\tilde{v}=0)$ solitons vs the scaled comoving axial coordinate $\tilde{\zeta}$.

where $\delta_{SC} = \gamma_{SC} f_{\infty}^{\frac{4}{3}}$. Instead for a gray soliton we obtain

$$\left(\frac{v}{v_s^{SC}}\right)^2 = \frac{3}{5}h_0^2 \frac{3h_0^{\frac{10}{3}} - 5h_0^2 + 2}{(1 - h_0^2)^2}.$$
(68)

$$\left(\frac{v}{v_s^{SC}}\right)^2 = \frac{3}{5}h_0^2 \frac{3h_0^{\frac{10}{3}} - 5h_0^2 + 2}{(1 - h_0^2)^2}.$$

$$\zeta = \frac{1}{\sqrt{\delta_{SC}}} \sqrt{\frac{5}{2}} \int_{h_0}^{h(\zeta)} \frac{dh \ h}{\sqrt{2h^2 - 5h^4 + 3h_0^{\frac{16}{3}} - (2h_0^2 - 5h_0^4 + 3h_0^{\frac{16}{3}})\frac{(1 - h^2)^2}{(1 - h_0^2)^2}}}, (69)$$

where $v_s^{SC} = \sqrt{\frac{2}{3}}\delta_{SC}$. In dimensional units, taking into account equation (18), denoting $\rho_0 = \rho(0)$ the minimal axial density of the dark soliton and $\rho_{\infty} = \rho(\pm \infty)$ the bulk axial density of the dark soliton, one finds that under the condition $\rho_{\infty}a_{\perp}\ll 1$ the dark soliton is surely in the strictly 1D regime of strong transverse confinement at any point of the axial coordinate. Instead, under the condition $\rho_0 a_{\perp} \gg 1$ the dark soliton is surely in the 3D regime of weak transverse confinement at any point of the axial coordinate. Clearly, in the case of the black soliton, where $\rho_0 = 0$, near the minimum the bosonic cloud cannot be in the regime of weak transverse confinement.

It is important to stress that the comparison of our theoretical results with future experiments is constrained not only by the short lifetime of unitary bosonic gas due to fast three-body recombinations but also by the lifetime due to the snake instability [16–19]. A dark soliton, that is not strictly 1D, has a snaking transverse oscillation which breaks the axial symmetry and eventually destroys the soliton. Since the NPSE preserves axial symmetry, it is not suitable for investigating snaking oscillations and the dynamics of the dissolving dark soliton. However, it is possible to write a triaxial NPSE by using two transverse width $\sigma_x(z,t)$ and $\sigma_y(z,t)$ in the ansatz of equation (15); see [35] for details in the case of weakly-interacting bosons.

6. Conclusion

We have derived the dynamical equations that govern the motion of a unitary Bose gas at zero temperature. Using these equations, we have studied dark solitons in a unitary Bose gas, confined in a cylindrically symmetric potential. We have reduced the dimensionality of the 3D nonlinear Schrödinger problem by means of the 1D nonpolynomial Schrödinger equation approach, which keeps dynamically into account also the transverse width. We have solved the equations employing for the most analytical techniques, in order to find axial density, transverse width, phase and velocity of both black and gray solitons, in the weak and strong confinement regimes. We have found that the weak confinement approximation breaks down in the case of a black soliton.

Our theoretical predictions could be tested experimentally employing Bose gases of alkali-metal atoms at ultra-low temperatures, whose scattering length can be tuned by means of Feshbach resonance techniques to reach the unitary limit. Despite the difficulties which presently limit the time during which such systems can be observed due to three-body losses but also to the snake instability, we are confident that the recent developments in the experimental techniques are going to allow in the future to put our predictions at test.

References

- [1] Inouye S, Andrews MR, Stenger J, Miesner H-J, Stamper-Kurn DM and Ketterle W 1998 Observation of Feshbach resonances in a Bose-Einstein condensate *Nature* **392** 151
- [2] O'Hara KM, Hemmer SL, Gehm ME, Granade SR and Thomas JE 2002 Observation of a Strongly Interacting Degenerate Fermi Gas of Atoms Science 298 2179
- [3] Regal CA, Ticknor C, Bohn JL and Jin DS 2003 Creation of ultracold molecules from a Fermi gas of atoms Nature 424 47
- [4] Strecker KE, Partridge GB and Hulet RG 2003 Conversion of an Atomic Fermi Gas to a Long-Lived Molecular Bose Gas Phys. Rev. Lett. 91 080406
- [5] Jochim S, Bartenstein M, Altmeyer A, Hendl G, Riedl S, Chin C, Hecker Denschlag J and Grimm R 2003 Bose-Einstein Condensation of Molecules Science 302 2101
- [6] Gehm ME, Hemmer SL, Granade RS, O'Hara KM and Thomas JE 2003 Mechanical stability of a strongly interacting Fermi gas of atoms Phys. Rev. A 68 011401
- [7] Li W and Ho T-L 2012 Bose Gases near Unitarity Phys. Rev. Lett. 108 195301
- [8] Rem BS, Grier AT, Ferrier-Barbut I, Eismann U, Langen T, Navon N, Khaykovich L, Werner F, Petrov DS, Chevy F and Salomon C 2013 Lifetime of the Bose Gas with Resonant Interactions Phys. Rev. Lett. 110 163202
- [9] Makotyn P, Klauss CE, Goldberger DL, Cornell EA and Jin DS 2014 Universal dynamics of a degenerate unitary Bose gas Nature Physics 10 116
- [10] Fletcher RJ, Gaunt AL, Navon N, Smith RP and Hadzibabic Z 2013 Stability of a Unitary Bose Gas Phys. Rev. Lett. 111 125303
- [11] Giorgini S, Pitaevskii LP and Stringari S 2008 Rev. Mod. Phys. 80 1215
- [12] Roberts JL, Claussen NR, Cornish SL and Wieman CE 2000 Magnetic Field Dependence of Ultracold Inelastic Collisions near a Feshbach Resonance *Phys. Rev. Lett.* **85** 728
- [13] Burger S, Bongs K, Dettmer S, Ertmer W, Sengstock K, Sanpera A, Shlyapnikov GV and Lewenstein M 1999 Dark Solitons in Bose-Einstein Condensates Phys. Rev. Lett. 83 5198
- [14] Denschlag J, Simsarian JE, Feder DL, Clark CW, Collins LA, Cubizolles J, Deng L, Hagley EW,

- Helmerson K, Reinhardt WP, Rolston SL, Schneider BI and Phillips WD 2000 Generating Solitons by Phase Engineering of a Bose-Einstein Condensate Science 287 97
- [15] Ku MJH, Mukherjee B, Yefsah T, and Zwierlein MW 2016 Cascade of Solitonic Excitations in a Superfluid Fermi gas: From Planar Solitons to Vortex Rings and Lines Phys. Rev. Lett. 116 045304
- [16] Antezza M, Dalfovo F, Pitaevskii LP and Stringari S 2007 Dark solitons in a superfluid Fermi gas Phys. Rev. A 76 043610
- [17] Cetoli A, Brand J, Scott RG, Dalfovo F, and Pitaevskii LP 2013 Snake instability of dark solitons in fermionic superfluids Phys. Rev. A 88 043639
- [18] Lombardi G, Van Alphen W, Klimin SN and Tempere J 2017 Soliton core filling in superfluid Fermi gases with spin-imbalance *Phys. Rev. A* **96** 033609
- [19] Van Alphen W, Lombardi G, Klimin SN and Tempere J 2018 Dark soliton collisions in superfluid Fermi gases New J. Phys. **20** 053052
- [20] Salasnich L, Parola A and Reatto L 2002 Effective wave equations for the dynamics of cigar-shaped and disk-shaped Bose condensates Phys. Rev. A 65 043614
- [21] Salasnich L, Parola A and Reatto L 2004 Transition from three dimensions to one dimension in Bose gases at zero temperature Phys. Rev. A 70 013606
- [22] Salasnich L, Parola A and Reatto L 2005 Quasi-one-dimensional bosons in three-dimensional traps: From strong-coupling to weak-coupling regime Phys. Rev. A 72 025602
- [23] Salasnich L and Malomed BA 2006 Vector solitons in nearly one-dimensional Bose-Einstein condensates Phys. Rev. A 74 053610
- [24] Salasnich L, Cetoli A, Malomed BA, Toigo F and Reatto L 2007 Bose-Einstein condensates under a spatially modulated transverse confinement Phys. Rev. A 76 013623
- [25] Salasnich L, Malomed BA and Toigo F 2007 Matter-wave vortices in cigar-shaped and toroidal waveguides Phys. Rev. A 76 063614
- [26] Cowell S, Heiselberg H, Mazets IE, Morales J, Pandharipande VR and Pethick CJ 2002 Cold Bose Gases with Large Scattering Lengths Phys. Rev. Lett. 88 210403
- [27] Adhikari SK and Salasnich L 2008 Nonlinear Schrödinger equation for a superfluid Bose gas from weak coupling to unitarity: Study of vortices Phys. Rev. A 77 033618
- [28] Song JL and Zhou F 2009 Ground State Properties of Cold Bosonic Atoms at Large Scattering Lengths *Phys. Rev. Lett.* **103** 025302
- [29] Lee Y-L and Lee Y-W 2010 Universality and stability for a dilute Bose gas with a Feshbach resonance *Phys. Rev. A* 81 063613
- [30] Diederix JM, van Heijst TCF and Stoof HTC 2011 Universality and stability for a dilute Bose gas with a Feshbach resonance *Phys. Rev. A* 84 033618
- [31] Rossi M, Salasnich L, Ancilotto F, and Toigo F 2014 Monte Carlo simulations of the unitary Bose gas *Phys. Rev. A* 89 041602
- [32] Pitaevskii LP and Stringari S 2003 Bose-Einstein Condensation Oxford Univ Press
- [33] Gross EP 1961 Structure of a quantized vortex in boson systems Nuovo Cimento 20 454
- [34] Pitaevskii LP 1961 Vortex lines in an imperfect Bose gas Sov. Phys. JETP 13 451
- [35] Salasnich L 2009 Generalized nonpolynomial Schrödinger equations for matter waves under anisotropic transverse confinement J. Phys. A: Math. Theor. 42 335205