

EXISTENCE OF CSCK METRICS ON SMOOTH MINIMAL MODELS

ZAKARIAS SJÖSTRÖM DYREFELT

ABSTRACT. Given a compact Kähler manifold X it is interesting to ask whether it admits a constant scalar curvature Kähler (cscK) metric. In this short note we show that there always exist cscK metrics on compact Kähler manifolds with nef canonical bundle, thus on all smooth minimal models, and also on the blowup of any such manifold. This confirms an expectation of Jian-Shi-Song [19] and extends their main result from K_X semi-ample to K_X nef, with a direct proof that does not appeal to the Abundance conjecture. As a byproduct we obtain that the connected component $\text{Aut}_0(X)$ of a compact Kähler manifold with K_X nef is either trivial or a complex torus.

1. INTRODUCTION

Let X be a compact Kähler manifold of dimension $n \geq 2$, let K_X be the canonical bundle and $c_1(X) := c_1(-K_X)$ the associated canonical $(1, 1)$ -cohomology class. Going back to the work of Calabi [5], existence of constant scalar curvature (cscK) metrics has been a much studied question in Kähler geometry. In particular, by the classical work of Yau [30] and Aubin [1] it has been known for decades that cscK metrics always exist on compact Kähler manifolds with ample canonical bundle. Much more recently it was shown by Jian-Shi-Song [19] that cscK metrics always exist also when K_X is only semi-ample. The step from ample to semi-ample is of special interest thanks to the minimal model program, and especially in view of the Abundance conjecture. Indeed, by MMP and Abundance for surfaces, every compact Kähler surface of Kodaira dimension $\kappa \geq 0$ is birational to a surface which is minimal. In particular, it follows from [19] that all such surfaces admit cscK metrics. There are some other situations when the Abundance conjecture is known, but in general the gap K_X semi-ample to K_X nef is still conjecture.

In this short note we confirm the expectation of [19] that their existence result should hold in general for $-c_1(X)$ nef, with a proof not relying on the Abundance conjecture, and which is valid for arbitrary (projective or non-projective) compact Kähler manifolds in any dimension:

Theorem 1. *Suppose that X is a compact Kähler manifold with $-c_1(X)$ nef. Then for any Kähler class $[\omega] \in H^{1,1}(X, \mathbb{R})$, there is $\epsilon_{X, [\omega]} > 0$, such that for all $0 < \epsilon < \epsilon_{X, [\omega]}$, there exists a unique cscK metric in the Kähler class $-c_1(X) + \epsilon[\omega]$. In particular, every smooth minimal model admits a cscK metric.*

In fact, we prove the stronger statement that the Mabuchi K-energy functional is proper in the above classes $-c_1(X) + \epsilon[\omega]$ (see Definition 1 for explanations of the terminology). Since the K-energy is then invariant under automorphisms, this implies that the automorphism group is compact, and hence the connected component $\text{Aut}_0(X)$ is either a singleton or a complex torus:

Corollary 2. *Suppose that X is a compact Kähler manifold with $-c_1(X)$ nef. Then $\text{Aut}_0(X)$ is either trivial or a complex torus.*

This result adds to the literature on automorphism groups for projective algebraic varieties with K_X semi-ample, in which case the corresponding result is well-known (see e.g. [20, Theorem 8.1] and references therein).

As a further application of Theorem 1 we note that if $-c_1(X)$ is nef, then X cannot support any non-trivial Hamiltonian vector fields, and hence the machinery of Arezzo-Pacard [2] yields existence of cscK metrics also on blowups of smooth minimal models:

Corollary 3. *Blowups of compact Kähler manifolds with $-c_1(X)$ nef admit cscK metrics.*

A few comments on the proof of Theorem 1 are in order: Just as in [19] our proof partly relies on the work of Chen-Cheng [7], which is concerned with coercivity of energy functionals on the closure \mathcal{E}^1 of the space of Kähler metrics. The new ingredient is to combine the machinery in [7] with Collins-Székelyhidi [10] and an observation about stability thresholds, namely the following numerical quantity

$$\delta([\omega]) := \delta_1([\omega]) - \delta_2([\omega])$$

where

$$\delta_1([\omega]) := \sup\{\delta_1 \in \mathbb{R} : \exists C > 0, M(\varphi) \geq \delta_1 \|\varphi\| - C, \forall \varphi \in \mathcal{E}^1\}$$

and

$$\delta_2([\omega]) := \sup\{\delta_2 \in \mathbb{R} : \exists C > 0, E_{\omega}^{-\rho\omega}(\varphi) \geq \delta_2 \|\varphi\| - C, \forall \varphi \in \mathcal{E}^1\}.$$

Here M is the K-energy functional, $E_{\omega}^{-\rho\omega}$ is the energy part in its Chen-Tian decomposition, see [9]. Note that the above thresholds are well-defined real numbers, so in particular they cannot take the value minus infinity (see e.g. [25, Proposition 12]). By coercivity of entropy we also observe that $\delta([\omega]) \geq \frac{n+1}{n} \alpha_X([\omega]) > 0$ where the latter denotes the classical alpha invariant of Tian [27].

In order to streamline the proof of Theorem 1 we introduce the following shorthand terminology related to the J-equation and J-stability: Let θ be any smooth $(1,1)$ -form on X and write $[\theta] \in H^{1,1}(X, \mathbb{R})$ for the associated cohomology class. We say that $((X, [\omega]); [\theta])$ is *J-coercive* (or *J-stable*, see [10, 6]) if and only if the E_{ω}^{θ} -functional is coercive on the space $\mathcal{E}^1 := \mathcal{E}^1(X, \omega)$ of finite energy potentials, i.e. there are constants $\delta, C > 0$ such that

$$E_{\omega}^{\theta}(\varphi) \geq \|\varphi\| - C$$

for all $\varphi \in \mathcal{E}^1$.

The strategy of proof is then based on the following elementary observation, which characterizes properness of the Mabuchi K-energy via a condition that in view of [25] can be interpreted as a geometric condition on the Kähler cone:

Theorem 4. *Suppose that $-c_1(X)$ is nef. Then the K-energy is proper on the space of Kähler metrics cohomologous to ω precisely if $((X, [\omega]); [\theta])$ is J-coercive, where*

$$[\theta] := -c_1(X) + \delta([\omega])[\omega].$$

In particular, if either of the above conditions hold, then X admits a cscK metric in the Kähler class $[\omega] \in H^{1,1}(X, \mathbb{R})$.

Given a compact Kähler manifold with $-c_1(X)$ nef, the idea is then to show that the J-coercivity condition of the above theorem is always satisfied for any Kähler class close enough to $-c_1(X)$. In fact a bit more can be said, as shown by the proofs of Theorem 1 and Theorem 4 in the next section.

Remark 1. By a minor alteration of the proof of Theorem 1 we can more generally obtain a version of the above for the twisted cscK metrics that play a central role in the continuity method of Chen-Cheng [7]: Suppose that $\eta \in H^{1,1}(X, \mathbb{R})$ is a $(1, 1)$ -form such that $-c_1(X) + \eta$ is nef. Then for any Kähler class $[\omega]$ on X , there is $\epsilon_{X, [\omega]} > 0$ such that for all $0 < \epsilon < \epsilon_{X, [\omega]}$, there exists a unique η -twisted cscK metric, that is, a solution to $\text{Tr}_\beta(-\text{Ric}(\omega) + \eta) = \text{constant}$, in the Kähler class $[\beta] = -c_1(X) + \eta + \epsilon[\omega]$. This follows by simply replacing $-c_1(X)$ by $-c_1(X) + \eta$ everywhere in the proof.

Acknowledgements. The author is grateful to Jian Song, Yalong Shi and Claudio Arezzo for helpful comments and discussions, as well as to the anonymous referee for improving the paper, in particular by pointing out Corollary 2.

2. PROOF

We quickly recall the notation and terminology for energy functionals that we will use, which is the standard variational setup frequently used throughout the Kähler geometry literature. To introduce our notation, let (X, ω) be a compact Kähler manifold of complex dimension $n \geq 2$ and write $\gamma := [\omega] \in H^{1,1}(X, \mathbb{R})$ for the associated Kähler class. Let

$$V_\gamma := \int_X \frac{\omega^n}{n!}$$

be the Kähler volume of (X, ω) . Let ρ_ω be the Ricci curvature form, normalized such that $[\rho_\omega] = c_1(X)$, and write $S(\omega) := \text{Tr}_\omega \rho_\omega$ for the scalar curvature of (X, ω) . By a standard abuse of terminology we say that ω is a constant scalar curvature Kähler (cscK) metric if $S(\omega)$ is constant. Denote the automorphism group of X by $\text{Aut}(X)$ and its connected component of the identity by $\text{Aut}_0(X)$. Write $\mathcal{C}_X \subset H^{1,1}(X, \mathbb{R})$ for the cone of Kähler cohomology classes on X . Let $\overline{\mathcal{C}}_X$ be the nef cone, $\partial\mathcal{C}_X$ its boundary, and let Big_X be the cone of big $(1, 1)$ -classes on X .

We write $(\mathcal{H}_\omega, d_1)$ for the space of Kähler potentials on X endowed with the L^1 -Finsler metric d_1 , and denote by (\mathcal{E}^1, d_1) its metric completion (see [11, 12, 13, 14] and references therein). Write $\text{PSH}(X, \omega) \cap L^\infty(X)$ for the space of bounded ω -psh functions on X .

Now consider $\varphi \in \text{PSH}(X, \omega) \cap L^\infty(X)$. We may define well-known energy functionals

$$\begin{aligned} \mathbb{I}_\omega(\varphi) &:= \frac{1}{V_\gamma n!} \int_X \varphi(\omega^n - \omega_\varphi^n) \\ \mathbb{J}_\omega(\varphi) &= \frac{1}{V_\gamma n!} \int_X \varphi \omega^n - \frac{1}{V_\gamma(n+1)!} \int_X \varphi \sum_{j=0}^n \omega^j \wedge \omega_\varphi^{n-j} \\ \mathbb{E}_\omega^\theta(\varphi) &:= \frac{1}{V_\gamma n!} \int_X \varphi \sum_{j=0}^{n-1} \theta \wedge \omega^j \wedge \omega_\varphi^{n-j-1} - \frac{1}{V_\gamma(n+1)!} \int_X \varphi \underline{\theta} \sum_{j=0}^n \omega^j \wedge \omega_\varphi^{n-j} \end{aligned}$$

where θ is any smooth closed $(1,1)$ -form on X and $\underline{\theta}$ is the topological constant given by

$$\underline{\theta} := \frac{\int_X \theta \wedge \frac{\omega^{n-1}}{(n-1)!}}{\int_X \frac{\omega^n}{n!}}.$$

By the Chen-Tian formula [9] the K-energy functional can be written as the sum of an energy/pluripotential part and an entropy part as

$$\mathbb{M}_\omega = \mathbb{E}_\omega^{-\rho_\omega} + \mathbb{H}_\omega \quad (1)$$

where

$$\mathbb{H}_\omega(\varphi) := \frac{1}{V_\gamma n!} \int_X \log \left(\frac{\omega_\varphi^n}{\omega^n} \right) \omega_\varphi^n$$

is the relative entropy of the probability measures $\omega_\varphi^n/V_\gamma$ and ω^n/V_γ . In particular, it is well known that $\mathbb{H}_\omega(\varphi)$ is always non-negative.

For any given smooth closed $(1,1)$ -form θ on X we also consider the θ -twisted K-energy functional

$$\mathbb{M}_\omega^\theta := \mathbb{M}_\omega + \mathbb{E}_\omega^\theta.$$

As explained in [25], it will also in this paper be convenient to measure properness/coercivity of the K-energy against the functional

$$\|\varphi\| := (\mathbb{I}_\omega - \mathbb{J}_\omega)(\varphi) = \frac{1}{V_\gamma(n+1)!} \int_X \varphi \sum_{j=0}^n \omega^j \wedge \omega_\varphi^{n-j} - \frac{1}{V_\gamma n!} \int_X \varphi \omega_\varphi^n$$

rather than against the Aubin J-functional or the d_1 -distance.

Definition 1. Let $F : \mathcal{E}^1 \rightarrow \mathbb{R}$ be any of the above considered energy functionals. We then say that F is coercive if

$$F(\varphi) \geq \delta \|\varphi\| - C$$

for some $\delta, C > 0$ and all $\varphi \in \mathcal{E}^1$.

In the rest of this note we will use the notation of projective polarized manifolds (X, L) , while emphasizing that this is only cosmetic, and all the arguments below hold for arbitrary Kähler classes on arbitrary (possibly non-projective) compact Kähler manifolds.

Proof of Theorem 4. The proof of Theorem 4 is a simple reformulation of the fact that existence of cscK metrics in a Kähler class $c_1(L)$ is connected to J-stability/J-coercivity of the triple $((X, L); K)$ as in the introduction, where $K := K_X$. We will use the following necessary and sufficient existence criteria due to [28, 29] and [16]: A polarised manifold $((X, L); K)$ is J-stable if

$$n \frac{K \cdot L^{n-1}}{L^n} L - (n-1)K > 0, \quad (2)$$

that is this divisor is ample, and a necessary condition for J-stability of $((X, L); K)$ is that

$$n \frac{K \cdot L^{n-1}}{L^n} L - K > 0. \quad (3)$$

In dimension $n = 2$ the condition thus becomes necessary and sufficient.

These conditions can equivalently be expressed in terms of stability thresholds, following [25]. Indeed, in the notation of that paper we can reformulate the above conditions as a double inequality

$$n \frac{K \cdot L^{n-1}}{L^n} - (n-1)\sigma(K, L) \leq \delta_{2,K}(L) \leq n \frac{K \cdot L^{n-1}}{L^n} - \sigma(K, L), \quad (4)$$

where

$$\sigma(K, L) := \inf\{\lambda \in \mathbb{R} : K - \lambda L < 0\}$$

is a Seshadri type constant and

$$\delta_{2,K}(L) := \sup\{\delta \in \mathbb{R} : \exists C > 0, E_\omega^{-\rho\omega}(\varphi) \geq \delta\|\varphi\| - C, \forall \varphi \in \mathcal{E}^1\}$$

is the optimal coercivity constant for the *energy part* $E_\omega^{-\rho\omega}$ of the Mabuchi K-energy, which by [10] is positive if and only if the triple $((X, L); K)$ is J-stable.

The relationship with the cscK equation follows by considering in the same way the optimal coercivity constant

$$\delta_1(L) := \sup\{\delta \in \mathbb{R} : \exists C > 0, M(\varphi) \geq \delta\|\varphi\| - C, \forall \varphi \in \mathcal{E}^1\}$$

associated to the K-energy, appealing to the results of [7], and then relating it to J-stability of $((X, L); K)$ by observing that

$$\delta_1(L) = \delta_{2, K + (\delta_1(L) - \delta_{2,K}(L))L}(L).$$

In other words the K-energy is proper on the space of Kähler metrics in $c_1(L)$ if and only if $((X, L); \theta)$ is J-stable with respect to the smooth $(1, 1)$ -form defined by

$$\theta := \theta_L := K + (\delta_1(L) - \delta_{2,K}(L))L.$$

Moreover, if any of the above equivalent conditions hold, then (X, L) is a cscK manifold, which is precisely the statement of Theorem 4. Note moreover that θ is not necessarily assumed positive here, since while stability requires positivity to make sense, coercivity can be defined more generally with respect to any smooth $(1, 1)$ -form θ on X .

Proof of Theorem 1. Motivated by Theorem 4 we now study the $(1, 1)$ -form $\theta := \theta_L$ more closely. The first thing to comment on is that we cannot expect to compute θ_L in any easy manner, since then we would know essentially everything about the existence problem for cscK metrics. However, it is immediate to observe the following inequality

$$\delta_1(L) - \delta_{2,K}(L) \geq \frac{n+1}{n} \alpha_X(L) > 0, \quad (5)$$

which follows directly from the fact that entropy is coercive (in fact, with the bound $H(\varphi) \geq \epsilon \|\varphi\| - C$ for every $\epsilon < \frac{n+1}{n} \alpha_X(L)$).

Suppose now once and for all that the canonical bundle $K := K_X$ is nef. Then by (5) it follows that θ is ample. By adding $\delta_1(L) - \delta_{2,K}(L)$ to each side of the double inequality (4), we deduce a similar double inequality also for $\delta_1(L)$:

$$n \frac{\theta \cdot L^{n-1}}{L^n} - (n-1) \sigma(\theta, L) \leq \delta_1(L) \leq n \frac{\theta \cdot L^{n-1}}{L^n} - \sigma(\theta, L). \quad (6)$$

In order to make use of this observation, it will be enough to show that there exists an ample line bundle L on X for which the left hand side of (6) is strictly positive. This is achieved by a simple application of the methods in [25]. In case $n = 2$ the subcone of the Kähler cone where the E_ω^θ -functional is coercive with respect to a given θ is described explicitly in [25]. In higher dimension a complete description was not given, but all that is needed for the proof of our main result are the criteria (2), (3) and the following lemma, whose proof is identical to that in [25]:

Lemma 5. *Let Λ, T be line bundles such that $c_1(\Lambda) \in \mathcal{C}_X$ and $c_1(T) \in \partial \mathcal{C}_X$, that is, Λ is ample and T is nef but not ample. Then $L_t := (1-t)T + t\Lambda$, $t \in [0, 1]$ satisfies*

$$\sigma(\Lambda, L_t) = \frac{1}{t}.$$

The same result holds if $\Lambda \in \partial \mathcal{C}_X$, as long as $c_1(L_t) \in \mathcal{C}_X$ for all $t \in (0, 1)$.

Proof. The first part is [25, Lemma 16]. For the second part, if we write $\sigma'(\beta, \gamma) := \sup\{\delta > 0 : \beta - \delta\gamma > 0\}$ then $\sigma(\Lambda, L_t) = \sigma'(L_t, \Lambda)^{-1}$. We moreover have $\sigma'(L_t, \Lambda) = (1-t)\sigma'(L_0, \Lambda) + t\sigma'(L_1, \Lambda)$. If $L_0, L_1 \in \partial \mathcal{C}_X$ are boundary classes such that $L_t \in \mathcal{C}_X$ for all $t \in (0, 1)$, then $\sigma'(L_0, \Lambda) = 0$ and $\sigma'(L_1, \Lambda) = 1$. \square

We are finally ready to prove the main result, extending [19] and replacing the hypothesis of K_X semi-ample with K_X nef, as predicted by the Abundance conjecture. We recall that while it is here presented in the terminology of projective polarized manifolds (X, L) , it is valid also for arbitrary compact Kähler manifolds with the same proof.

Theorem 6. (Theorem 1) *Suppose that K_X is nef. Then for any ample line bundle L on X , there is $\epsilon_{X,L} > 0$, such that for all $0 < \epsilon < \epsilon_{X,L}$, there exists a unique cscK metric in the Kähler class $-c_1(X) + \epsilon c_1(L)$. In particular, every smooth minimal model X admits a cscK metric.*

Proof. Suppose that $K := K_X$ is nef but not ample (if it is ample then we already know that cscK metrics exist, by Aubin, Yau [1, 30]). As explained

above it is a consequence of [7, 10, 16, 28, 29] that cscK metrics must exist if the Kähler class

$$\theta := K + (\delta_1(L) - \delta_{2,K}(L))L$$

satisfies

$$n \frac{\theta \cdot L^{n-1}}{L^n} - (n-1)\sigma(\theta, L) > 0.$$

Now fix an auxiliary line bundle T which is nef but not ample, in such a way that $L_t := (1-t)T + tK$ is ample for all $t \in (0, 1)$. Consider

$$R_\epsilon(t) := n \frac{(K + \epsilon L_t) \cdot L_t^{n-1}}{L_t^n} - (n-1)\sigma((K + \epsilon L_t), L_t)$$

as a function of $\epsilon \in \mathbb{R}$ and $t \in (0, 1)$. By the last part of Lemma 5 we then have

$$R_0(t) = n \frac{K \cdot L_t^{n-1}}{L_t^n} - \frac{n-1}{t},$$

so that in particular $R_0(1) = 1$. By continuity we may therefore fix $t_0 \in (0, 1)$ such that $R_0(t) > 0$ for all $t \in (t_0, 1)$. Moreover, it can be easily checked that

$$\sigma((K + \epsilon L_t), L_t) = \sigma(K, L_t) + \epsilon$$

and hence also

$$R_\epsilon(t) = R_0(t) + \epsilon.$$

It follows that $R_\epsilon(t) > 0$ for all $t > t_0$ and all $\epsilon > 0$. Taking $\epsilon_{L_t} := \delta_1(L_t) - \delta_{2,K}(L_t)$ and noting that this quantity is always positive, it follows from Theorem 4 and (4) that $c_1(L_t)$ admits cscK metrics for all $t > t_0$. By possibly rescaling L if necessary, it follows that for every ample line bundle L on X , there is $\epsilon_{X,L} > 0$ such that for all $0 < \epsilon < \epsilon_{X,L}$, there exists a unique cscK metric in the Kähler class $-c_1(X) + \epsilon c_1(L)$. This is what we wanted to prove. \square

Since it is also clear that the line bundles can be replaced everywhere with arbitrary nef or Kähler $(1, 1)$ -cohomology classes, we obtain precisely the statement of Theorem 1 in the introduction.

2.1. Proof of Corollary 2 and Corollary 3. By Theorem 1 there exists a Kähler class $[\omega] \in H^{1,1}(X, \mathbb{R})$ (as described in the statement of the theorem) such that the associated Mabuchi K-energy functional M is d_1 -proper on the space \mathcal{H}_ω of Kähler potentials. In particular, by [7] there exists a cscK metric in $[\omega]$, and we may fix once and for all a cscK potential $v \in \mathcal{H}_\omega$. Let moreover $G := \text{Aut}_0(X)$ be the connected complex Lie group given by the identity component of the group of biholomorphisms of X . Denote by \mathfrak{g} its Lie algebra, consisting of holomorphic vector fields on X . The key step of the proof of Corollary 2 is showing that properness of the K-energy forces G to be compact. To see this, we recall the arguments of [15]: Let $K := \text{Isom}_0(X, \omega_v)$ be the identity component of the isometry group of (X, ω_v) , and let \mathfrak{k} be its Lie algebra. Note that K is compact, since X is compact (see [24, Proposition 29.4]). By a result of Matsushima and Lichnerowicz (see [18, Theorem 3.6.1]) and by [15, Proposition 6.2], it follows that if $g_j \in G$, then there are $k_j \in K$ and $V_j \in \mathfrak{k}$ such that $g_j = k_j \exp_I(JV_j)$. Recall moreover that G acts on the space of normalized potentials $\mathcal{H}_\omega \cap E^{-1}(0)$ (where $E := E_\omega$ is

the Monge-Ampère energy functional) such that $g \cdot \varphi \in \mathcal{H}_\omega \cap E^{-1}(0)$ is the unique potential defined by the relation

$$\omega_{g \cdot \varphi} = g^* \omega_\varphi.$$

With this setup, it was shown in [15, Section 7.1] that

$$[0, +\infty) \ni t \mapsto \exp_I(-tJV_j) \cdot v \in \mathcal{H}_\omega \cap E^{-1}(0)$$

is a d_1 -geodesic ray. Since the K-energy is invariant under the action of G , properness then implies that $d_1(g_j \cdot v, v) \leq C$, with a bound independent of j . Arguing as in [4, Lemma 3.7] it then follows that $d_1(\exp_I(-JV_j) \cdot v, v) \leq C$, and hence $\|V_j\|$ is uniformly bounded in $\mathfrak{k} := \mathfrak{isom}(X, \omega_v)$. By compactness we can then, up to relabeling, choose $V \in \mathfrak{k}$ and $k \in K$ such that $k_j \rightarrow k$ and $V_j \rightarrow V$ smoothly. Hence also $g_j \rightarrow g := k \exp_I(JV)$ smoothly, and properness implies compactness. Finally, it is a standard exercise (see e.g. [17, Chapter 8]) to see that a connected compact complex Lie group is abelian, and in fact a complex torus. This proves Corollary 2.

In order to prove Corollary 3 we finally note that the machinery of Arezzo-Pacard [2] applies in case there are no non-trivial holomorphic vector fields with zeros, that is no non-trivial Hamiltonian vector fields. To see that we are in this case, we once again use properness of the K-energy functional: Let as before v be a cscK potential and suppose for contradiction that $V \in \mathfrak{isom}(X, \omega_v)$ is a non-trivial Hamiltonian vector field. By the work of [12] and [15, Section 7.1] it follows that $[0, +\infty) \ni t \mapsto \exp_I(-tJV) \cdot v$ defines a constant speed geodesic ray in the space of normalized Kähler potentials $\mathcal{H}_\omega \cap E^{-1}(0)$. In particular, if V is non-trivial then well-known convexity properties imply that $d_1(v, \exp_I(-tJV) \cdot v) \rightarrow +\infty$ as $t \rightarrow +\infty$ (recall that d_1 is comparable to the norm $\|\cdot\|$ and to Aubin's J-functional, which is strictly convex along geodesics). On the other hand, since (X, ω) admits a cscK metric, the K-energy is invariant under the action of G , contradicting the properness assumption. In other words X cannot support any non-trivial Hamiltonian vector fields, and [2] combined with Theorem 1 yields the statement of Corollary 3.

REFERENCES

- [1] T. Aubin, *Equations du type Monge-Ampère sur les variétés kähleriennes compactes*, Bull. Sci. Math. (2) 102 (1978), no. 1, 63-95
- [2] C. Arezzo and F. Pacard, *Blowing up and desingularizing constant scalar curvature Kähler manifolds*, Acta Math. **196** (2006), no. 2, 179-228.
- [3] R. Berman, S. Boucksom, V. Guedj and A. Zeriahi, *A variational approach to complex Monge-Ampère equations*, Publ. Math. de l'IHES **117** (2013), 179-245.
- [4] R. Berman and T. Darvas and C. Lu, *Regularity of weak minimizers of the K-energy and applications to properness and K-stability*, Preprint arXiv:1602.03114v1, (2016).
- [5] E. Calabi, *Extremal Kähler metrics*, In: "Seminars on Differential Geometry", S.T. Yau (ed.), Princeton Univ. Press and Univ. of Tokyo Press, Princeton, New York, 1982, 259-290.
- [6] G. Chen, *On J-equation*, Preprint arXiv:1905.10222 (2019).
- [7] X.X. Chen and J. Cheng, *On the constant scalar curvature Kähler metrics, existence results*, Preprint arXiv:1801.00656 (2018).
- [8] X.X. Chen and J. Cheng, *On the constant scalar curvature Kähler metrics, general automorphism group*, arXiv:1801.05907 (2018).

- [9] X.X. Chen, *On the lower bound of the Mabuchi K-energy and its application*, Int. Math. Res. Not. **12** (2000), 607-623.
- [10] T. Collins and G. Székelyhidi, *Convergence of the J-flow on toric manifolds*, J. Diff. Geom. **107** (2017) no. 1, 47-81.
- [11] T. Darvas, *The Mabuchi completion of the space of Kähler potentials*, Amer. J. Math **139** (2017), 1275-1313.
- [12] T. Darvas, *The Mabuchi Geometry of finite energy classes*, Adv. Math. **285** (2015), 182-219.
- [13] T. Darvas, *Geometric pluripotential theory on Kähler manifolds*, Survey article (2017).
- [14] T. Darvas, *Weak geodesic rays in the space of Kähler potentials and the class $E(X, \omega_0)$* , J. Inst. Math. Jussieu **16** (2017), no. 4, 837-858.
- [15] T. Darvas and Y.A. Rubinstein, *Tian's properness conjecture and Finsler geometry of the space of Kähler metrics*, J. Amer. Math. Soc. **30** (2017), no. 2, 347-387.
- [16] S.K. Donaldson, *Moment maps and diffeomorphisms*, Asian J. Math., **3** (1999) no. 1, pp. 1-15.
- [17] W. Fulton and J. Harris, *Representation theory: A first course*, Springer-Verlag New York, 2004.
- [18] P. Gauduchon, *Calabi's extremal metrics: an elementary introduction*, manuscript, 2010.
- [19] W. Jian, Y. Shi and J. Song, *A remark on constant scalar curvature Kähler metrics on minimal models*, Proc. Amer. Math. Soc. **147** (2019), 3507-3513.
- [20] S. Kobayashi, *Transformation groups in differential geometry*, Springer-Verlag Berlin Heidelberg New York, 1972.
- [21] T. Mabuchi, *A functional integrating Futaki invariants*, Proc. Japan Acad. **61** (1985), 119-120.
- [22] T. Mabuchi, *K-energy maps integrating Futaki invariants*, Tohoku Math. J. **38** (1986), no. 4, 575-593.
- [23] T. Mabuchi, *Some symplectic geometry on compact Kähler manifolds I*, Osaka J. Math. **24** (1987), 227-252.
- [24] M.M. Postnikov, *Geometry VI. Riemannian geometry*, Springer, 2001.
- [25] Z. Sjöström Dyrefelt, *Optimal lower bounds for Donaldson's J-functional*, Adv. Math. DOI: 10.1016/j.aim.2020.107271.
- [26] J. Song and B. Weinkove, *On the convergence and singularities of the J-flow with applications to the Mabuchi energy*, Comm. Pure Appl. Math., **61** (2008), 210 - 229.
- [27] G. Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), no. 1, 1-37.
- [28] B. Weinkove, *Convergence of the J-flow on Kähler surfaces*, Comm. Anal. Geom. **12** (2004), no. 4, 949-965.
- [29] B. Weinkove, *On the J-flow in higher dimensions and the lower boundedness of the Mabuchi energy*, J. Diff. Geom. **73** (2006), no. 2, 351-358.
- [30] S.-T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*, Comm. Pure Appl. Math. **31** (1978), 339 - 411.

THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS (ICTP),
 STR. COSTIERA, 11, 34151 TRIESTE TS, ITALY.
 Email address: zsjostro@ictp.it