

Completeness of Wronskian Bethe equations for rational $\mathfrak{gl}_{m|n}$ spin chains

Dmitry Chernyak^{a,b} Sébastien Leurent^c Dmytro Volin^{a,d,e}

^a*School of Mathematics & Hamilton Mathematics Institute, Trinity College Dublin, College Green, Dublin 2, Ireland*

^b*Département de Physique, École Normale Supérieure, Université PSL, 24 rue Lhomond, 75005 Paris, France*

^c*Université de Bourgogne-Franche-Comté, Institut de Mathématiques de Bourgogne, UMR 5584 du CNRS, 9 avenue Alain Savary, 21000 Dijon, France*

^d*Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden*

^e*Department of Physics and Astronomy, Uppsala University, Box 516, SE-751 20 Uppsala, Sweden*

E-mail: dmitry.chernyak@ens.fr, sebastien.leurent@u-bourgogne.fr, dmytro.volin@physics.uu.se

ABSTRACT: We consider rational integrable supersymmetric $\mathfrak{gl}_{m|n}$ spin chains in the defining representation and prove the isomorphism between a commutative algebra of conserved charges (the Bethe algebra) and a polynomial ring (the Wronskian algebra) defined by functional relations between Baxter Q-functions that we call Wronskian Bethe equations. These equations, in contrast to standard nested Bethe equations, admit only physical solutions for any value of inhomogeneities and furthermore we prove that the algebraic number of solutions to these equations is equal to the dimension of the spin chain Hilbert space (modulo relevant symmetries).

Both twisted and twist-less periodic boundary conditions are considered, the isomorphism statement uses, as a sufficient condition, that the spin chain inhomogeneities θ_ℓ , $\ell = 1, \dots, L$ satisfy $\theta_\ell + \hbar \neq \theta_{\ell'}$ for $\ell < \ell'$. Counting of solutions is done in two independent ways: by computing a character of the Wronskian algebra and by explicitly solving the Bethe equations in certain scaling regimes supplemented with a proof that the algebraic number of solutions is the same for any value of θ_ℓ . In particular, we consider the regime $\theta_{\ell+1}/\theta_\ell \gg 1$ for the twist-less chain where we succeed to provide explicit solutions and their systematic labelling with standard Young tableaux.

Notations

Typical values of indices

a, b	1 to m
i, j	$\hat{1}$ to \hat{n} (hat is omitted sometimes)
α, β	from the set $\{1, \dots, m, \hat{1}, \dots, \hat{n}\}$
ℓ	1 to L

Parameters

z_α	twist eigenvalues, $z_a \equiv x_a, z_{\hat{i}} \equiv y_i$
θ_ℓ	inhomogeneities (as variables)
$\bar{\theta}_\ell$	inhomogeneities (fixed number)
χ_ℓ	elementary symmetric polynomials
$\bar{\chi}_\ell$	$= \chi_\ell(\bar{\theta}_1, \dots, \bar{\theta}_L)$

Lie algebra

$\mathfrak{g}_{m n}$	symmetry of the system (broken to Cartan in the twisted case)
$E_{\alpha\beta}$	abstract generators and defining representation
$\mathcal{E}_{\alpha\beta}$	global spin chain action
$\Lambda^+ = (\underline{\lambda}_1, \underline{\lambda}_2, \dots)$	Young diagram \equiv integer partition (typically of L)
$(\underline{\lambda}'_1, \underline{\lambda}'_2, \dots)$	transposed partition, $h_{\Lambda^+} := \underline{\lambda}'_1$.
$\Lambda = [\lambda_1, \dots, \lambda_m \nu_1, \dots, \nu_n]$	fundamental weight (eigenvalues of $\mathcal{E}_{\alpha\alpha}$)
$(\hat{\lambda}_1, \dots, \hat{\lambda}_{m'} \hat{\nu}_1, \dots, \hat{\nu}_{n'})$	shifted weight (describes Λ^+ with marked point)

Spin chain

V	Hilbert space of the spin chain ($\simeq (\mathbb{C}^{m n})^{\otimes L}$)
V_Λ	subspace of V spanned by states of weight Λ
V_Λ^+	subspace of V spanned by highest weight states of irreps Λ^+
U_Λ	either V_Λ or V_Λ^+
d_Λ	dimension of U_Λ

Bethe and Wronskian algebras

$\hat{c}_k^{(d)}, \hat{c}_\ell$	operators acting on spin chain, coefficients in Baxter Q-operators, <i>e.g.</i> $Q_k = u^{M_k} (1 + \frac{\hat{c}_k^{(1)}}{u} + \dots)$
$c_k^{(d)}, c_\ell$	abstract variables and/or eigenvalues of $\hat{c}_k^{(d)}, \hat{c}_\ell$
\mathcal{B}_Λ	Bethe algebra restricted to U_Λ (generated by \hat{c}_ℓ), a $\mathbb{C}[\chi]$ -module
$\mathcal{B}_\Lambda(\bar{\theta})$	specialised Bethe algebra (for spin chain representation at point $\bar{\theta}$)
$\mathcal{B}_\Lambda(\bar{\chi})$	specialised Bethe algebra (for symmetrised representation at point $\bar{\chi}$)
\mathcal{W}_Λ	Wronskian algebra (generated by c_ℓ subject to Wronskian Bethe equations)
$\mathcal{W}_\Lambda(\bar{\chi})$	specialised Wronskian algebra

Functional relations conventions

u	spectral parameter
\hbar	Unit of discrete shift in <i>e.g.</i> Baxter equation, typically $\hbar = \pm i, \pm 1, \pm 2$
$f^{[n]}$	$f^{[n]} \equiv f(u + \frac{\hbar}{2}n), f^\pm \equiv f^{[\pm 1]}$
$f \propto g$	f and g , as functions of u , are equal up to a normalisation

Contents

1	Introduction	4
2	Definitions and basic properties	8
2.1	$\mathfrak{gl}_{m n}$ Lie superalgebra, shifted and fundamental weights	9
2.2	Yangian	11
2.3	Bethe algebra	12
2.4	Q-operators	14
2.5	Quantisation condition (Wronskian Bethe equations)	15
3	Completeness	21
3.1	Analytic description	21
3.2	How to count solutions with multiplicity	22
3.3	Wronskian algebra is a free $\mathbb{C}[\chi]$ -module	25
3.4	Number of solutions <i>via</i> Hilbert series	25
4	Faithfulness	27
4.1	Isomorphism between Wronskian and Bethe algebra	28
4.2	What can happen upon specialisation	30
4.3	Specialisation of the isomorphism	32
4.4	Construction of Bethe vectors	35
5	Various parameterisations of the Bethe algebra	36
5.1	Restriction and extension of Q-systems	36
5.2	Isomorphism of twist-less \mathcal{B}_Λ across $\mathfrak{gl}_{m n}$ algebras of various ranks	37
5.3	Q-system on Young diagrams	39
5.4	Relation to nested Bethe equations and quantum eigenvalues	41
6	Labelling solutions	45
6.1	Twisted case, labelling with a multinomial expansion	45
6.2	Twist-less case, labelling with standard Young tableaux	46
7	Summary and applications	48
7.1	Completeness, faithfulness, and maximality of the Bethe algebra	48
7.2	Simplicity of the spectrum and controlled numerical solution	50
7.3	Gaudin model	51
7.4	Separation of variables	52
7.5	Geometric representation theory and Bethe/Gauge correspondence	53
A	Wronskian algebra - a pedestrian approach	54
A.1	Some facts from commutative algebra and algebraic geometry	54
A.2	$\mathbb{C}(\chi)$ -module and invariance of solutions multiplicity	56
A.3	Freeness of \mathcal{W}_Λ and trivialisation of a vector bundle	59
A.4	Non-symmetric functions	60

B	Cyclicity of representations	61
B.1	Symmetrised Yangian representation	62
B.2	Symmetrised Bethe modules and their characters	65
B.3	Cyclicity of symmetrised Bethe modules	67
B.4	Specialisations	68
B.5	An explicit case study	70
C	Q-operators belong to the Bethe algebra	73
D	Details about structural study of Bethe equations	74
D.1	Properness - twisted case	75
D.2	Properness - twist-less case	75
D.3	Labelling solutions with standard Young tableaux – technical details	78

1 Introduction

Rational integrable spin chains are one of the first quantum integrable systems that were discovered and studied. In fact, their simplest $SU(2)$ representative was introduced and solved, by means of coordinate Bethe Ansatz, in the seminal paper of Hans Bethe [1].

In this article we consider periodic integrable spin chains of length L constructed using the $\mathfrak{gl}_{m|n}$ -invariant rational R-matrix, and with spin chain nodes being in fundamental (defining) representation of $\mathfrak{gl}_{m|n}$. The parameters defining the model are the twist matrix G and inhomogeneities $\theta_1, \dots, \theta_L$. We cover the cases when G is either equal to the identity (twist-less case) or is diagonalisable with distinct eigenvalues $x_1, \dots, x_m, y_1, \dots, y_n$ (generic twisted case).

Spectrum of the commuting charges that form the so-called Bethe algebra \mathcal{B} can be encoded into rational symmetric combinations of the Bethe roots $u_k^{(\alpha)}$. Equations defining the values of $u_k^{(\alpha)}$ shall be called Bethe equations, and their most known presentation is given by nested Bethe Ansatz equations (NBAE) which is the following relation between fractions [2–5]

$$\prod_{\ell=1}^L \frac{u_k^{(\alpha)} - \theta_\ell + \frac{c_{1,2} + c_{1,1}}{2} \hbar \delta_{\alpha,1}}{u_k^{(\alpha)} - \theta_\ell - \frac{c_{1,2} + c_{1,1}}{2} \hbar \delta_{\alpha,1}} = \frac{z_{\alpha+1}}{z_\alpha} \prod_{\substack{1 \leq \beta \leq m+n-1 \\ 1 \leq l \leq M_\beta \\ (\beta,l) \neq (\alpha,k)}} \frac{u_k^{(\alpha)} - u_l^{(\beta)} + \frac{\hbar}{2} c_{\alpha,\beta}}{u_k^{(\alpha)} - u_l^{(\beta)} - \frac{\hbar}{2} c_{\alpha,\beta}}. \quad (1.1)$$

Here $\alpha \in \{1, 2, \dots, m+n-1\}$ and all $k \in \{1, \dots, M_\alpha\}$, we denote $z_\alpha = x_\alpha$ for $1 \leq \alpha \leq m$, and $z_\alpha = y_{\alpha-m}$ for $m+1 \leq \alpha \leq m+n$, and \hbar is a non-zero complex number (typical choices are $i, 1, 2$). Finally $c_{\alpha,\beta}$ is the Cartan matrix of the $\mathfrak{sl}_{m|n}$ subalgebra of $\mathfrak{gl}_{m|n}$. It is equal *e.g.* to $\begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}$ for $\mathfrak{sl}(2|2)$, the expression for other ranks should be obvious from this example. This expression of the Cartan matrix is written in the so-called distinguished grading of $\mathfrak{gl}_{m|n}$ but other gradings are also possible [6, 7], the corresponding equations are obtained *via* duality transformations, and we briefly mention them in Section 5.4.

Obvious questions arising are whether each solution of the Bethe equations describes some physical state, which we call the *faithfulness* property, and whether all physical states can be described in this way, which we call the *completeness* property. In particular, one often asks whether the number of solutions to the Bethe equations is equal to the dimension of the Hilbert space, probably after some obvious symmetries are factored out. In the literature, these properties are typically covered by the name of completeness, however the precise meaning of the word varies.

Quite surprisingly, despite the fundamental nature of these questions, they were properly resolved only in 2009 for the \mathfrak{gl}_2 case and in 2013 for the \mathfrak{gl}_m case by Mukhin, Tarasov and Varchenko [8, 9]. Completeness and faithfulness were also recently proven for $\mathfrak{gl}_{1|1}$ by Huang, Lu, and Mukhin [10, 11]. Proving completeness and faithfulness for an arbitrary rank $\mathfrak{gl}_{m|n}$ case is the subject of the current paper. For formal proofs, we build on ideas of [9] and add several new insights, even for the bosonic \mathfrak{gl}_m subcase, to achieve the result. Besides formal proofs, we also give a recipe to explicitly label solutions.

Counting of solutions was first time addressed already in [1] using the so-called string hypothesis, and later on this approach was extended to \mathfrak{gl}_m [12], $\mathfrak{gl}_{2|1}$ [13] and $\mathfrak{gl}_{2|2}$ [14] cases. Further study of combinatorics implied by string hypothesis for \mathfrak{gl}_m spin chains led to the formulation of the Kerov-Kirillov-Reshetikhin bijection [15, 16] between rigged configurations of Bethe strings and (in case of spin chains in the defining representation) standard Young tableaux. Although counting assuming string hypothesis leads to correct numbers, the hypothesis is strictly speaking wrong as one can show by a more detailed analysis and explicit counter-examples, see *e.g.* [17, 18]. Hence this approach, after all, did not accomplish its original thought application – proving the completeness of rational Bethe equations. Instead, ideas of [15, 16] became extremely fruitful and were further generalised in various applications of algebraic combinatorics, in particular in the context of “combinatorial” integrability that can be viewed as the crystal limit $q \rightarrow 0$ of q -deformed (XXZ-type) spin chains, see *e.g.* [19, 20] and references therein.

Apart from the combinatorial challenge, analysing solutions of (1.1) has clear technical complications. First, solutions with coinciding Bethe roots generically do not correspond to physical states and then they should be discarded. However, there are cases when such solutions should be kept [21–23]. Second, the so-called exceptional solutions with $u = u' \pm \hbar$ and/or $u = \theta \pm \frac{\hbar}{2}$ (case *a* in [21]) render relation (1.1) singular. Some of the exceptional solutions are physical and some of them are not, and, for instance, their behaviour upon change of twist or inhomogeneities can decide for their physicality. Tracing this behaviour becomes a burden, especially at higher rank. To our knowledge, only the homogeneous \mathfrak{gl}_2 case was properly understood [24]. At higher ranks, non-physicality can be also hidden in non-physical exceptional solutions of dual Bethe equations even if the original Bethe equations appear as being free from any singularities [25].

It is then not surprising that one should look for a different set of equations instead of (1.1) to prove completeness [26], and it is indeed the case for the proof in [9] where a very elegant Wronskian condition was used. Define the finite-difference Wronskian between any

number of k functions as

$$W(F_1, \dots, F_k) \equiv \det_{1 \leq i, j \leq k} F_i(u + \hbar(\frac{k+1}{2} - j)), \quad (1.2)$$

introduce m monic polynomials¹ $q_{a|\emptyset} = u^{M_{a|\emptyset}} \left(1 + \sum_{k=1}^{M_{a|\emptyset}} \frac{c_{a|\emptyset}^{(k)}}{u^k} \right)$, $a = 1, \dots, m$ of degree $M_{a|\emptyset}$. Then the eigenstates of the Bethe algebra of the \mathfrak{gl}_m spin chain are in a one-to-one correspondence with the solutions of

$$\frac{W(x_1^{u/\hbar} q_{1|\emptyset}, \dots, x_m^{u/\hbar} q_{m|\emptyset})}{W(x_1^{u/\hbar}, \dots, x_m^{u/\hbar})} = \prod_{\ell=1}^L (u - \theta_\ell) \quad (1.3)$$

which should be considered as equations on the coefficients $c_{a|\emptyset}^{(k)}$. As formulated, the statement holds for the case when all x_a are pairwise distinct, and it also applies to the twist-less case after arrangements discussed in Section 2.5.

We shall call (1.3) Wronskian Bethe equations (WBE). They are equivalent to NBAE (with coinciding Bethe roots solutions being discarded) for generic values of θ_ℓ but, in contrast to (1.1), smoothly work at *any* values of inhomogeneities, and this includes an important physical case of homogeneous spin chain with all $\theta_\ell = 0$. Relation to Bethe roots of (1.1) is given by $\frac{W(x_1^{u/\hbar} q_{1|\emptyset}, \dots, x_a^{u/\hbar} q_{a|\emptyset})}{W(x_1^{u/\hbar}, \dots, x_a^{u/\hbar})} = \prod_{k=1}^{M_a} (u - u_k^{(a)})$, $M_a = \sum_{b=1}^a M_{b|\emptyset}$.

WBE are natural in the logic of the analytic Bethe Ansatz [27, 28] (though in early works in this formalism their significance was not recognised), while NBAE are often associated with the (nested) coordinate or algebraic Bethe Ansatz [29, 30] (though they are derived *via* analytic Bethe Ansatz as well [31, 32]). The details of the completeness and faithfulness questions depend on the chosen approach.

The nested coordinate/algebraic Bethe Ansatz is indeed an ansatz to build an eigenfunction and Bethe equations appear as the necessary consistency conditions for the ansatz to succeed. Faithfulness is then the question of sufficiency of these conditions, while completeness is the question whether there exist eigenfunctions not described by this Ansatz. In physics literature, faithfulness is often considered as granted reducing completeness to counting the number of solutions to the Bethe equations. The generic position faithfulness indeed follows rather straightforwardly from the ansatz itself, but a full systematic proof covering all exceptional situations would be much harder to achieve. To our knowledge, such a proof using a direct algebraic Bethe Ansatz framework was done only very recently for the \mathfrak{gl}_2 homogeneous case [33]. In the context of separation of variables, a new type of ansatz to build eigenstates in higher-rank systems emerged [34–36]. Its faithfulness for \mathfrak{gl}_m chains is proven in the case of non-degenerate twist and assuming $\theta_\ell - \theta_{\ell'} \neq \hbar\mathbb{Z}$.

The analytic Bethe Ansatz is actually not an ansatz to build wave functions. It often starts by considering the Bethe algebra – a set of commuting operators which satisfy various functional relations as functions of the spectral parameter u that were extensively studied

¹We use notations suited for our generalisation to supersymmetric case. They are different from those in [9].

[37–41]. WBE is one of (or a consequence of) these relations with $Q_{a|\emptyset} := x_a^{u/\hbar} q_{a|\emptyset}$ being the renowned Q-operators, the first example of such an operator is due to Baxter [42, 43]. In the analytic Bethe Ansatz approach, faithfulness is non-trivial to demonstrate even in generic position, and it is an important part of our paper to prove it. On the other hand, one is certain that each of the Q-operators eigenvalues satisfy WBE. Hence the question of completeness becomes equivalent to the question of whether the Bethe algebra generated by Q-operators contains the full set of commuting charges, *i.e.* whether the eigenvalues of Q-operators are sufficient to fully parametrise the Hilbert space. We shall resolve this question positively by explicitly counting the number of solutions of WBE. Hence we still face the question of counting as in the algebraic/coordinate Bethe Ansatz scenario, however the statement that is proven as a consequence of counting is a bit different.

In physical applications, inhomogeneities are often set to $\theta_\ell = 0$. However, keeping inhomogeneities as parameters that we are going to vary is decisive for the approach discussed in this paper ². To start with, inhomogeneities are regulators that put our system to a generic position. One can even explore regimes where Bethe equations can be solved explicitly which provides a constructive way to count solutions. In the twisted case, such a regime is $\left| \frac{\theta_{\ell+1} - \theta_\ell}{\hbar} \right| \gg 1$ when we label solutions using binomial expansions, and in the twist-less case such a regime is $\left| \frac{\theta_{\ell+1}}{\theta_\ell} \right| \gg 1$ when we can label solutions using standard Young tableaux (SYT), a result first time stated in [44]. Moreover if θ_ℓ are positive, we will demonstrate that there is an unambiguous analytic continuation from this regime to the point $\theta_\ell = 0$.

Furthermore, we show that the Bethe algebra can be generated by only L generators. There are also L inhomogeneities which allows us to prove generic position faithfulness statements using a rigorous version of a “number of variables equals number of parameters” argument.

Finally, it can be demonstrated that all properties can be made *polynomial* in χ_ℓ – elementary symmetric polynomials in inhomogeneities. Algebraically, this shall be formalised by proving that certain properly designed objects are free $\mathbb{C}[\chi_1, \dots, \chi_L]$ -modules. This implies that general position completeness and faithfulness statements can be specialised to *any* numerical value of χ_ℓ .

The paper is organised as follows.

In Section 2, we recall all the necessary known results about the Yangian, the Bethe algebra, and Q-operators. The fact that Q-operators belong to the Bethe algebra on the level of representation is proven in Appendix C. We conclude the section with the formulation of the supersymmetric twisted and twist-less versions of WBE (1.3).

Sections 3 and 4 provide proofs of completeness and faithfulness.

Section 3 proves completeness. We introduce the concept of Wronskian algebra (a polynomial ring defined by WBE), prove that it is a free $\mathbb{C}[\chi]$ -module, and explicitly count its rank using Hilbert series (a.k.a. character, index, partition function) confirming

²The twist values x_a, y_i are always kept fixed however.

that it coincides with the dimension of the (corresponding) Hilbert space. By a standard argument, the rank of the Wronskian algebra is the number of solutions to WBE counted with multiplicities. The proof of freeness essentially uses the so-called properness of WBE which is proven in Appendix D.1 and Appendix D.2.

Section 4 proves faithfulness. More accurately, faithfulness means that the Bethe algebra is a faithful representation of the Wronskian algebra, which will immediately imply that these two algebras are isomorphic. The proof is first done for generic values of inhomogeneities (over the polynomial ring $\mathbb{C}[\chi]$) and then is specialised to numerical values of χ_ℓ . Important results allowing one to specialise at any numerical value are covered in Appendix B which builds substantially on the approach of [9].

Sections 5 and 6 aim to make the obtained results more practical.

Section 5 discusses various ways to parametrise the Bethe algebra in the twist-less case. In particular, we demonstrate that all the restricted Bethe algebras \mathcal{B}_Λ for $\mathfrak{gl}_{m|n}$ with fixed Young diagram Λ^+ and different m, n are isomorphic to one-another and to the Q-system on this diagram. We also explain how this formalism is mapped to NBAE.

Section 6 considers regimes $\left| \frac{\theta_{\ell+1} - \theta_\ell}{\hbar} \right| \gg 1$, $\left| \frac{\theta_{\ell+1}}{\theta_\ell} \right| \gg 1$ and shows how to explicitly find solutions of WBE in these regimes. Some technical questions are postponed to Appendix D.3.

In Section 7, we summarise the results and then discuss their immediate applications. This includes an algorithm to solve Bethe equations (including at $\theta_\ell = 0$) with solutions being labelled with SYT, and applications to the Gaudin model and to the separation of variables program. We conclude the section with a review of a relation between the restricted Bethe algebras and certain quantum cohomology rings.

The paper uses substantially results and terminology from algebraic geometry and commutative algebra while the target audience includes researchers with no appropriate background. To alleviate the issue, we illustrate the discussion with numerous examples, including a comprehensive case study in Appendix B.5, and supplement the paper with Appendix A containing mostly textbook material applied to the concrete problem that we consider. Section 3.2 also summarises textbook knowledge about multiplicity of solutions but we decided to keep it in the main text given its importance for the paper.

2 Definitions and basic properties

As often happens in mathematical physics, it will be useful to recast a physical question into a problem in representation theory. A spin chain should be considered as a representation of the $\mathfrak{gl}_{m|n}$ Yangian. Commuting Hamiltonians belong to its certain commutative subalgebra known as [45] the Bethe algebra \mathcal{B} , and the completeness question is closely related to the explicit realisation of this algebra by Baxter polynomials subjected to constraints.

This section collects definitions of the above-mentioned objects. For the supersymmetric Yangian, our sign conventions are the same as in [46, 47], however we define quantum Berezinian with a different overall shift of the spectral parameter. Many hidden subtleties in a consistent usage of signs are nicely reviewed in [48].

2.1 $\mathfrak{gl}_{m|n}$ Lie superalgebra, shifted and fundamental weights

Let us recall some essential facts about the $\mathfrak{gl}_{m|n}$ Lie superalgebra [49]. Assign parity $\bar{a} = 0$ for any “bosonic” index $a \in \{1, \dots, m\}$ and parity $\bar{i} = 1$ for any “fermionic” index $i \in \{\hat{1}, \dots, \hat{n}\}$. The $\mathfrak{gl}_{m|n}$ algebra is spanned by the generators $E_{\alpha\beta}$, $\alpha, \beta \in \{1, \dots, m, \hat{1}, \dots, \hat{n}\}$ whose graded Lie bracket is

$$[E_{\alpha\beta}, E_{\alpha'\beta'}] = \delta_{\beta\alpha'} E_{\alpha\beta'} - (-1)^{(\bar{\alpha}+\bar{\beta})(\bar{\alpha}'+\bar{\beta}')} \delta_{\beta'\alpha} E_{\alpha'\beta}. \quad (2.1)$$

The Hilbert space³ comprising states of the spin chain of length L is $V := (\mathbb{C}^{m|n})^{\otimes L}$. Each spin chain site $\mathbb{C}^{m|n}$ transforms under the defining (vector) representation of $\mathfrak{gl}_{m|n}$. The global action of $\mathfrak{gl}_{m|n}$ on V shall be denoted as $\mathcal{E}_{\alpha\beta}$. It is induced from the single site action by using the standard graded product rule, *e.g.* if $E_{\alpha\beta} \mathbf{e}_\gamma = \delta_{\beta\gamma} \mathbf{e}_\alpha$ then, for $L = 2$

$$\mathcal{E}_{\alpha\beta} \mathbf{e}_\gamma \otimes \mathbf{e}_{\gamma'} = \delta_{\beta\gamma} \mathbf{e}_\alpha \otimes \mathbf{e}_{\gamma'} + (-1)^{(\bar{\alpha}+\bar{\beta})\bar{\gamma}} \delta_{\beta\gamma'} \mathbf{e}_\gamma \otimes \mathbf{e}_\alpha. \quad (2.2)$$

Overall, the sign rule $(A \otimes B)(C \otimes D) = (-1)^{\bar{B}\bar{C}}(AC) \otimes (BD)$ for tensor products of graded algebras and modules shall be always assumed.

When describing spin chains with periodic boundary conditions (*twist-less* case), $\mathfrak{gl}_{m|n}$ is the symmetry of the system in the sense that it commutes with the Hamiltonians we are interested to diagonalise. Correspondingly, we shall encounter covariant representations – the irreps appearing in the tensor powers of the defining $\mathfrak{gl}_{m|n}$ representation. These representations are of the highest-weight type, with highest-weight vector v being defined by the property $\mathcal{E}_{\alpha\beta} v = 0$ for $\alpha < \beta$.

The highest-weight property depends on a choice of the total order $<$ in the set $\{1, \dots, m, \hat{1}, \dots, \hat{n}\}$, and different choices related to permutation of bosonic and fermionic indices lead to non-equivalent parameterisations of the system. Most of the properties that we shall discuss do not depend on such an order, and so we will often use an invariant description based on labelling of the irreps with Young diagrams which is possible due to the supersymmetric version of the Schur-Weyl duality [50, 51].

Young diagrams Λ^+ with a marked point (m', n') on the boundary are in bijection with tuples of shifted⁴ weights $(\hat{\lambda}_1, \dots, \hat{\lambda}_{m'} | \hat{\nu}_1, \dots, \hat{\nu}_{n'})$. We choose the marked point to be $m' = m - r, n' = n - r$ for $r \in \mathbb{Z}_{\geq 0}$, the role of r is to reduce diagonally the rank of the $\mathfrak{gl}_{m|n}$ algebra such that the inner corner of the fat hook attains the Young diagram boundary, see Figure 1. The explicit relation between the shifted weights and the shape of Λ^+ is

$$\begin{aligned} \hat{\lambda}_a &= \underline{\lambda}_a - a - n + m, & a = 1, 2, \dots, m - r, \\ \hat{\nu}_i &= \underline{\lambda}'_i - i - m + n, & a = 1, 2, \dots, n - r, \end{aligned} \quad (2.3)$$

where $(\lambda_1, \lambda_2, \dots)$ is the integer partition forming the shape Λ^+ , $(\lambda'_1, \lambda'_2, \dots)$ is the integer partition of the transposed diagram, and $r = \min_k (\lambda_{m-k} + k - n \geq 0)$.

³Usage of terminology “Hilbert space” is customary for quantum systems, however we do not use any scalar products in this work, except in Section 7.2.

⁴by the Weyl vector and with the additional shift by -1 of $\hat{\nu}_i$ to get a symmetric description




Figure 1: One-to-one correspondence between shifted weights $(9, 6, 4, 0|3, 2)$ and a Young diagram with a marked point (red dot). Other points on the boundary (black dots) and hence other sets of shifted weights can be chosen. The outlined option is diagonally shifted from the corner of the $m|n$ -hook that is linked to the $\mathfrak{gl}_{m|n}$ representation theory [49]: the Young diagrams that fit into the hook exactly describe finite-dimensional irreps. The Young diagrams that touch the hook corner correspond to the so-called long (typical) irreps, and the diagrams that do not touch the corner correspond to the so-called short (atypical) irreps.

Most of the results *do not depend* on the choice of the marked point, this is demonstrated in Section 5. We made the choice of the diagonal reduction only for easier connections with the results already known in the literature.

For spin chains with twisted boundary conditions, and for generic diagonal twist, only the Cartan subalgebra of $\mathfrak{gl}_{m|n}$ is the symmetry of the system and states are then described using the fundamental weight. We define it as the tuple $\Lambda = [\lambda_1, \dots, \lambda_m | \nu_1, \dots, \nu_n]$, where $\mathcal{E}_{aa} v = \lambda_{aa} v$, $\mathcal{E}_{ii} v = \nu_i v$.

Dictated by the symmetry of the problem, we introduce restrictions of the Hilbert space to the weight subspaces

$$V \supset V_\Lambda \supset V_\Lambda^+. \quad (2.4)$$

V_Λ is defined as the space of all vectors with fundamental weight Λ . Its dimension is given by the multinomial coefficient $L! / (\prod_{1 \leq a \leq m} \lambda_a! \prod_{1 \leq i \leq n} \nu_i!)$. V_Λ^+ is defined as the space of the highest-weight vectors for all irreps with Young diagram Λ^+ inside V . Its dimension is equal to the number of standard Young tableaux of shape Λ^+ . For concreteness we choose the standard order $1 < 2 < \dots < m < \hat{1} < \dots < \hat{n}$ in which case the fundamental weight Λ of the highest-weight vectors of the irrep Λ^+ is given by the rule

$$\lambda_a = \underline{\lambda}_a, \quad \nu_i = \max(0, \underline{\lambda}'_i - m). \quad (2.5)$$

As is explained on page 67, any other choice of the total order would lead to an isomorphic description and to the same conclusions albeit explicit realisation of V_Λ^+ and certain related objects will be modified.

We shall use the notation U_Λ to denote either V_Λ^+ or V_Λ when discussion equally applies to both subspaces V_Λ^+ and V_Λ .

2.2 Yangian

The Yangian $Y(\mathfrak{gl}_{m|n})$ is a quasi-triangular Hopf algebra with rational R-matrix. We summarise below its properties which will be relevant for us, see *e.g.* [47, 52] for a more detailed discussion.

Let $\alpha, \beta \in \{1, \dots, m\} \cup \{\hat{1}, \dots, \hat{n}\}$ and $k \in \{1, 2, \dots\}$. The Yangian's generators $t_{\alpha\beta}^{(k)}$ are collected, *via* formal series in \hbar/u

$$t_{\alpha\beta}(u) = \delta_{\alpha\beta} \mathbb{1} + \frac{\hbar}{u} t_{\alpha\beta}^{(1)} + \dots \quad (2.6)$$

into ‘‘monodromies’’ $t_{\alpha\beta}(u)$ whose parity is equal to $\bar{\alpha} + \bar{\beta}$.

Quasi-triangularity is an RTT-type relation that reads in component form as

$$[t_{\alpha\beta}(u), t_{\gamma\delta}(v)] = \frac{\hbar(-1)^{\bar{\alpha}\bar{\beta} + \bar{\alpha}\bar{\gamma} + \bar{\beta}\bar{\gamma}}}{u - v} (t_{\gamma\beta}(u)t_{\alpha\delta}(v) - t_{\gamma\beta}(v)t_{\alpha\delta}(u)). \quad (2.7)$$

From Hopf algebra structures, we will only need the co-product $\Delta(t_{\alpha\beta}(u)) = \sum_\gamma t_{\alpha\gamma} \otimes t_{\gamma\beta}$.

To realise the Yangian representation on the spin chain, consider first the evaluation homomorphism

$$ev_{\theta_\ell} : t_{\alpha\beta}(u) \mapsto \delta_{\alpha\beta} \mathbb{1} + \hbar \frac{(-1)^{\bar{\alpha}} \mathbf{E}_{\alpha\beta}}{u - \theta_\ell}, \quad (2.8)$$

where $\mathbf{E}_{\alpha\beta}$ are the $\mathfrak{gl}_{m|n}$ generators in the defining representation. Then, for $\theta := (\theta_1, \dots, \theta_L)$, combine L such maps by repetitively using the co-product

$$ev_\theta : t_{\alpha\beta} \mapsto ev_{\theta_1} \otimes \dots \otimes ev_{\theta_L} \left(\sum_{\gamma_1, \dots, \gamma_{L-1}} t_{\alpha\gamma_1} \otimes t_{\gamma_1\gamma_2} \otimes \dots \otimes t_{\gamma_{L-1}\beta} \right). \quad (2.9)$$

We shall call (2.9) the *spin chain representation* of the Yangian.

This representation contains, in the first non-trivial coefficient of the \hbar/u expansion, the global action of $\mathfrak{gl}_{m|n}$ on the spin chain defined in Section 2.1:

$$ev_\theta(t_{\alpha\beta}(u)) = \delta_{\alpha\beta} \mathbb{1} + (-1)^{\bar{\alpha}} \frac{\hbar}{u} \mathcal{E}_{\alpha\beta} + \dots \quad (2.10)$$

For $T_{\alpha\beta} \equiv Q_\theta(u)t_{\alpha\beta}(u)$, where $Q_\theta(u) = \prod_{\ell=1}^L (u - \theta_\ell)$, $ev_\theta(T_{\alpha\beta}) \equiv Q_\theta ev_\theta(t_{\alpha\beta})$ are *polynomials* in u of degree at most L . Note that $ev_\theta(T_{\alpha\beta})$ are also polynomials in θ_ℓ . Construction (2.9) corresponds to the graphics commonly used to define the monodromy matrix of a spin chain from Lax operators: $ev_\theta(T_{\alpha\beta}) = \alpha \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \vdots \\ \bullet \end{array} \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \vdots \\ \bullet \end{array} \dots \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \vdots \\ \bullet \end{array} \begin{array}{c} \bullet \\ \vdots \\ \bullet \end{array} \beta$, and θ_ℓ , $\ell = 1, \dots, L$ are commonly known as the spin chain *inhomogeneities*.

We will use $T_{\alpha\beta} = Q_\theta t_{\alpha\beta}$ to denote both the Yangian generators and their images $ev_\theta(T_{\alpha\beta})$. The context shall make it clear which meaning is being used. Note that ev_θ is

not a faithful map and hence not all algebra-level results subdue the representation-level properties.

For the discussion of this paper, it will be often important to consider θ_ℓ as unevaluated commuting variables. When it is the case, the image of the map ev_θ are endomorphisms with polynomial coefficients ⁵:

$$ev_\theta : Y(\mathfrak{gl}_{m|n}) \longrightarrow (\text{End}(\mathbb{C}^{m|n}))^{\otimes L} \otimes \mathbb{C}[\theta], \quad (2.11)$$

where $\mathbb{C}[\theta] \equiv \mathbb{C}[\theta_1, \theta_2, \dots, \theta_L]$ is the polynomial ring in variables θ_ℓ . Such operators naturally act on $\mathcal{V} := V \otimes \mathbb{C}[\theta]$. Such a description also appears in the context of a Hecke algebra, see page 64.

If we are interested in inhomogeneities having particular numerical values – in which case we typically denote them by $\bar{\theta}_\ell$ – then we get a representation of the Yangian in a more standard sense

$$ev_{\bar{\theta}} : Y(\mathfrak{gl}_{m|n}) \longrightarrow (\text{End}(\mathbb{C}^{m|n}))^{\otimes L}. \quad (2.12)$$

If we need to emphasise that (2.12) but not (2.11) is being used, we shall refer to (2.12) as the *spin chain representation at point $\bar{\theta} \equiv (\bar{\theta}_1, \dots, \bar{\theta}_L)$* .

2.3 Bethe algebra

The below-defined Bethe algebra \mathcal{B} is a commutative subalgebra of $Y(\mathfrak{gl}_{m|n})$ which depends on a constant $\text{GL}(m|n)$ group matrix G dubbed twist. We restrict ourselves to the case when G is diagonalisable and furthermore choose a reference frame that diagonalises G , so \mathcal{B} actually depends only on the eigenvalues ⁶ $x_1, \dots, x_m, y_1, \dots, y_n$ of G . We will consider only two opposing cases: of generic twist, when all x_a, y_i are distinct, and of no-twist when $G = \mathbf{1}$. Considering intermediate cases is possible but combinatorially bulky ⁷.

The Bethe algebra \mathcal{B} is defined as the algebra that is polynomially generated by the transfer matrices \mathbb{T}_μ in covariant representations of $\mathfrak{gl}_{m|n}$ labelled with integer partitions or equivalently Young diagrams μ . By “polynomially generated” we mean that elements of \mathcal{B} are finite degree polynomials in \hat{d}_k – coefficients of the (*a priori* formal) \hbar/u expansion $\mathbb{T}_\mu = \chi_\mu(G) u^{L|\mu|} (\mathbf{1} + \hat{d}_1 \frac{\hbar}{u} + \dots)$, where $\chi_\mu(G)$ is the character of G in representation μ .

Transfer matrices \mathbb{T}_μ can be constructed using fusion from $T_{\alpha\beta}$ [53, 54] and hence are defined on the level of the Yangian as well as its representation ev_θ . When we descend to the representation level, $\mathbb{T}_\mu(u)$ is a degree- $L|\mu|$ polynomial in u , so the \hbar/u expansion truncates.

⁵Here we allow freedom of speech and consider $ev_\theta(Y(\mathfrak{gl}_{m|n}))$ in the sense of $ev_\theta(T_{\alpha\beta})$.

⁶Matrices of the $\text{GL}(m|n)$ group have entries belonging to a Grassmann algebra, hence their eigenvalues are in principle not complex numbers. Our discussion will assume that twists are complex numbers nevertheless. One can then check that the results still hold for any twists of type $x_a = A_a + n_a, y_i = B_i + n_i$ where $A, B \in \mathbb{C}$ and n - even nilpotent elements of the Grassmann algebra, assuming that A_a, B_i are pairwise distinct.

⁷Analytic structure of Q-functions for partially degenerate twists, which is an essential ingredient for the completeness statements, was explored in detail in [41].

Let (1^a) denote the Young diagram consisting of one column of height a , and (s) – the Young diagram consisting of one row of width s . To avoid discussing fusion in detail, we note that the \mathbb{T}_μ , and hence the Bethe algebra, can be polynomially generated from $\mathbb{T}_{(1^a)}$ or from $\mathbb{T}_{(s)}$, $a, s = 1, 2, \dots$ using the determinant Cherednik-Bazhanov-Reshetikhin (CBR) formula [55–57] – the Yangian version of Jacobi-Trudi identities for characters $\chi_\mu(G)$, while $\mathbb{T}_{(1^a)}$ and $\mathbb{T}_{(s)}$ are compactly defined through monodromies $T_{\alpha\beta}$ as [58]

$$\text{Ber} [\mathbb{1} - \mathcal{D} T(u) G \mathcal{D}] = \sum_{a=0}^{\infty} (-1)^a \mathcal{D}^a \mathbb{T}_{(1^a)}(u) \mathcal{D}^a, \quad (2.13a)$$

$$\frac{1}{\text{Ber} [\mathbb{1} - \mathcal{D} T(u) G \mathcal{D}]} = \sum_{s=0}^{\infty} \mathcal{D}^s \mathbb{T}_{(s)}(u) \mathcal{D}^s, \quad (2.13b)$$

where $\mathcal{D} \equiv e^{-\frac{1}{2} \hbar \partial_u}$.

The l.h.s. of (2.13) is defined as follows. For $\mathbf{M} = \mathbb{1} - \mathcal{D} T G \mathcal{D}$, introduce the notation $\mathbf{M} = \begin{pmatrix} [\mathbf{M}]_A & [\mathbf{M}]_B \\ [\mathbf{M}]_C & [\mathbf{M}]_D \end{pmatrix}$ and $\mathbf{M}^{-1} = \begin{pmatrix} [\mathbf{M}^{-1}]_A & [\mathbf{M}^{-1}]_B \\ [\mathbf{M}^{-1}]_C & [\mathbf{M}^{-1}]_D \end{pmatrix}$, where $[\mathbf{M}]_A$ is the $m \times m$ block of \mathbf{M} , $[\mathbf{M}^{-1}]_D$ is the $n \times n$ block of \mathbf{M}^{-1} etc. Then the Berezinian is defined as $\text{Ber}(\mathbf{M}) = \det [\mathbf{M}]_A \det [\mathbf{M}^{-1}]_D$, where the determinant of the block “A” (resp. “D”) is defined through a column-ordered (resp. line-ordered) expansion e.g. $\det [\mathbf{M}]_A = \epsilon_{a_1 \dots a_m} ([\mathbf{M}]_A)_{a_1 1} \dots ([\mathbf{M}]_A)_{a_m m}$. Although \mathbf{M} is a matrix with non-commutative entries $\mathbf{M}_{\alpha\beta}$, the entries satisfy the (supersymmetric version of) Manin relations $[\mathbf{M}_{\alpha\beta}, \mathbf{M}_{\gamma\delta}] = (-1)^{\bar{\alpha}\bar{\beta} + \bar{\alpha}\bar{\gamma} + \bar{\beta}\bar{\gamma}} [\mathbf{M}_{\gamma\beta}, \mathbf{M}_{\alpha\delta}]$ which ensure that the above-defined Berezinian can only change sign if columns/rows of \mathbf{M} are permuted. From earlier works, we mention that Berezinians in the context of $Y(\mathfrak{gl}_{m|n})$ were introduced in [46], and generalise similar constructions in the $Y(\mathfrak{gl}_m)$ case [45, 59, 60], see also [52].

The physical Hamiltonian of the system is an element of the Bethe algebra and it is usually chosen to be $\mathbb{H} = \partial_u \log \mathbb{T}_{(1)}(u)|_{u=0}$. The algebraic equivalent of the statement that the Bethe algebra contains all commuting charges is the statement that it is a maximal commutative subalgebra that contains \mathbb{H} . The question about maximality can be asked on the level of the Yangian algebra or of its spin chain representation. On the level of the algebra, in the bosonic $Y(\mathfrak{gl}_m)$ case, polynomial combinations of $\mathbb{T}_{(1^a)}$ indeed generate, for non-degenerate twist, a maximal commutative subalgebra of $Y(\mathfrak{gl}_m)$ [45, 52] but it seems an equivalent statement was not proved for the supersymmetric case. To our knowledge, a comprehensive study of the Bethe algebra on the Yangian algebra level is still lacking.

However, our goal is to describe the Bethe algebra represented on the spin chain in which case it can be understood much better. In particular, the quantum Berezinian defined

$$\text{qBer} \equiv \mathcal{D}^{(n-m)} \text{Ber} [\mathcal{D} T(u) G \mathcal{D}] \mathcal{D}^{(n-m)} \quad (2.14)$$

and which is known to generate the center of the Yangian [46, 61] can be expressed, at least on the level of representation, as a ratio of transfer matrices $\text{qBer} \propto \frac{\mathbb{T}_{((n+1)^m)}}{\mathbb{T}_{(n^{m+1})}}$ [48] and hence, by Hamilton-Cayley, belongs to the Bethe algebra.

It will be one of our results that, under a mild sufficient assumption on θ ($\theta_\ell + \hbar \neq \theta_{\ell'}$ for $\ell < \ell'$), the Bethe algebra is a maximal commutative subalgebra of $\text{End}(V)$ – the algebra of all linear transformations of the spin chain Hilbert space.

2.4 Q-operators

Although there are infinitely many $\mathbb{T}_{(1^a)}$ and $\mathbb{T}_{(s)}$ in the expansions (2.13), we need finitely many functions of u to generate the Bethe algebra. This can be seen for instance by analysing the CBR formula. On the level of representation, probably the most economic way is to express transfer matrices through Baxter Q-operators that were explicitly constructed as operators acting on the supersymmetric spin chain in [62–66]⁸. The Q-operators are not elements of the Yangian, but they do belong to the Bethe algebra in the representation ev_θ , in particular they are matrices whose coefficients are polynomials in θ_ℓ , see Appendix C.

The Q-operators generate the Bethe algebra as follows⁹ [37, 71, 72]

$$\begin{aligned} \mathbb{T}_{(s^a)} &= u^{asL} \chi_{a,s}(G) \left(\mathbb{1} + \hat{d}_1 \frac{\hbar}{u} + \dots \right) \\ &\propto \frac{1}{Q_{\emptyset|\emptyset}^{[a-s]}} \prod_{k=1}^a \prod_{l=1}^s \frac{Q_{\emptyset|\emptyset}^{[a+s+2-2k-2l]}}{(\text{Ber } G)^{u/\hbar}} \times \begin{cases} \epsilon^{b_1 \dots b_m} Q_{b_1 \dots b_a|\emptyset}^{[m-n+s]} Q_{b_{a+1} \dots b_m|\emptyset}^{[-s]}, & s \geq a - m + n \\ \epsilon^{i_1 \dots i_n} Q_{\emptyset|i_1 \dots i_s}^{[m-n-a]} Q_{\emptyset|i_{s+1} \dots i_n}^{[+a]}, & a \geq s + m - n \end{cases}, \end{aligned} \quad (2.15)$$

where ϵ denotes the Levi-Cevita antisymmetric tensor, summation over repeated indices is performed, and the \propto symbol involves a proportionality factor which is identified by imposing that the coefficient of the highest degree of $\mathbb{T}_{(s^a)}$ (as a polynomial in u) is the character $\chi_{(s^a)}(G)$.

In total, there are 2^{m+n} Q-operators. They are labelled as $Q_{A|I}$, where A is a multi-index from $\{1, \dots, m\}$ and I is a multi-index from $\{1, \dots, n\}$. $Q_{A|I}$ are anti-symmetric w.r.t. permutations in A and I , and polynomial up to an exponential prefactor (as in 1.3):

$$Q_{A|I} \propto \frac{\prod_{a \in A} x_a^{u/\hbar}}{\prod_{i \in I} y_i^{u/\hbar}} q_{A|I}, \quad (2.16)$$

where the proportionality factor¹⁰ in “ \propto ” is fixed by the condition that each $q_{A|I}$ is a monic polynomial in the variable u .

The Q-operators satisfy the following QQ-relations

$$Q_{Aab|I} Q_{A|I} = W(Q_{Aa|I}, Q_{Ab|I}), \quad (2.17a)$$

$$Q_{Aa|I} Q_{A|Ii} = W(Q_{Aa|Ii}, Q_{A|I}), \quad (2.17b)$$

$$Q_{A|Iij} Q_{A|I} = W(Q_{A|Ii}, Q_{A|Ij}). \quad (2.17c)$$

⁸For constructions of Q-operators for bosonic spin chains, see eg [38, 67–70].

⁹Equation (2.15) expresses \mathbb{T} for the so-called rectangular representations, where the Young diagram $(s^a) \equiv \underbrace{(s, s, s, \dots, s)}_{a \text{ times}}$ is of rectangular shape; representations (1^a) , (s) are special subcases. Generalisation to arbitrary representations is known [71].

¹⁰One can set the proportionality factor to be equal to one when $|A| + |I| \leq 1$, which fixes this factor for other values of A and I due to the relations (2.17). It is explicitly spelled out in *e.g.* [41].

Furthermore, the two operators $Q_{\emptyset|\emptyset}$ and $Q_{\bar{\emptyset}|\bar{\emptyset}}$ are central: on the one hand $Q_{\bar{\emptyset}|\bar{\emptyset}}$ is by construction a multiple of identity operator (hence a central element), and one can note that (2.15) gives¹¹

$$\text{qBer} \propto \frac{\mathbb{T}((n+1)^m)}{\mathbb{T}(n^{m+1})} \propto \begin{cases} \frac{Q_{\bar{\emptyset}|\bar{\emptyset}}^{[-m+n+1]} Q_{\bar{\emptyset}|\bar{\emptyset}}^{[-m+n+3]} \dots Q_{\bar{\emptyset}|\bar{\emptyset}}^{[+m-n+1]}}{(\text{Ber } G)^{(m-n)u/\hbar} Q_{\bar{\emptyset}|\bar{\emptyset}}^{[m-n-1]}} & \text{if } m \geq n, \\ \frac{(\text{Ber } G)^{(n-m)u/\hbar} Q_{\bar{\emptyset}|\bar{\emptyset}}^{[m-n+1]}}{Q_{\bar{\emptyset}|\bar{\emptyset}}^{[-m+n-1]} Q_{\bar{\emptyset}|\bar{\emptyset}}^{[-m+n-3]} \dots Q_{\bar{\emptyset}|\bar{\emptyset}}^{[+m-n-1]}} & \text{if } n \geq m. \end{cases} \quad (2.18)$$

The statement that qBer is in the center of the Yangian hence reflects the property that $Q_{\bar{\emptyset}|\bar{\emptyset}}$ itself is central.

On the other hand $Q_{\emptyset|\emptyset} = 1$, which allows to write all Q -operators explicitly in terms of $Q_{a|\emptyset}$, $Q_{\emptyset|i}$, and $Q_{a|i}$ [40, 41]:

$$Q_{(a|a+c)} \propto (Q_{(1|1)}^{[c]})^a Q_{(0|1)}^{[c-1]} \dots Q_{(0|1)}^{[1-c]}, \quad Q_{(a+c|a)} \propto (Q_{(1|1)}^{[c]})^a Q_{(1|0)}^{[c-1]} \dots Q_{(1|0)}^{[1-c]}, \quad (2.19)$$

For compactness, we used exterior forms $Q_{(a|b)} = \frac{1}{a!b!} Q_{a_1 \dots a_a | i_1 \dots i_b} \psi_0^{a_1} \dots \psi_0^{a_a} \psi_1^{i_1} \dots \psi_1^{i_b}$ with ψ_0^a, ψ_1^i being auxiliary Grassmann variables.

We shall need the expression for $Q_{\bar{\emptyset}|\bar{\emptyset}} \equiv Q_{1 \dots m | 1 \dots n}$ which explicitly is the following determinant

$$Q_{\bar{\emptyset}|\bar{\emptyset}}(u) = (-1)^{n(m-n)} \begin{vmatrix} Q_{1|1}^{[m-n]} & \dots & Q_{1|n}^{[m-n]} & Q_{1|\emptyset}^{[m-n-1]} & Q_{1|\emptyset}^{[m-n-3]} & \dots & Q_{1|\emptyset}^{[-(m-n)+1]} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ Q_{m|1}^{[m-n]} & \dots & Q_{m|n}^{[m-n]} & Q_{m|\emptyset}^{[m-n-1]} & Q_{m|\emptyset}^{[m-n-3]} & \dots & Q_{m|\emptyset}^{[-(m-n)+1]} \end{vmatrix}, \quad (2.20)$$

for the case $m \geq n$; a similar expression can be written also for $m > n$.

One should also note that the coefficients $Q_{a|\emptyset}$, $Q_{\emptyset|i}$, and $Q_{a|i}$ are related by (2.17b):

$$Q_{a|i}^+ - Q_{a|i}^- = Q_{a|\emptyset} Q_{\emptyset|i}. \quad (2.21)$$

2.5 Quantisation condition (Wronskian Bethe equations)

The essential property for the description of the Bethe algebra is the explicit analytic structure of Q -operators (a.k.a. rational analytic Bethe Ansatz) which is known from results of [63, 64, 69], and can be also derived using the logic of Appendix C.

The definition of Q -operators depends on a gauge choice, see *e.g.* [73] for details. Below we write expressions in one particular gauge which is suitable for our goals¹².

For the central element $Q_{\bar{\emptyset}|\bar{\emptyset}}$ it is possible to directly compute its explicit value which in the gauge of our choice becomes

$$Q_{\bar{\emptyset}|\bar{\emptyset}}(u) \propto (\text{Ber } G)^{u/\hbar} Q_{\theta}(u). \quad (2.22)$$

¹¹There exists a different choice of normalizations that simplifies this expression to $\text{qBer} = Q_{\bar{\emptyset}|\bar{\emptyset}}^- / Q_{\bar{\emptyset}|\bar{\emptyset}}^+$.

¹²Apart from the gauge choices, there also exists several other discrepancies in labelling conventions across the literature. First, an arrangement in the spectral parameter can be present. Second, the role of the Q -operators and their Hodge duals $Q^{A|J} \propto \varepsilon^{AA'} \varepsilon^{JJ'} Q_{JJ'}$ can be swapped. Third, a permutation of indices $1 \dots m | 1 \dots n$ can be used.

This property is an important aspect of the Bethe algebra and it is essentially equivalent to the set of Bethe equations as it will become clear below.

We set $\text{Ber } G = 1$ for convenience. It only affects the overall normalisation of transfer matrices and hence is inessential.

To write expressions for the other Q-operators, we need to restrict the representation space to a certain subspace. The generic twist and twist-less cases should be treated separately.

Twisted case

The Cartan generators $\mathcal{E}_{\alpha\alpha}$ of the global $\mathfrak{gl}_{m|n}$ action (2.10) commute with \mathcal{B} (and belong to it). To describe analytic properties of the Q-operators in a useful manner, we need to restrict to an eigenspace of $\mathcal{E}_{\alpha\alpha}$ which is the weight space V_Λ defined after (2.4). The Bethe algebra restricted to this subspace shall be denoted as

$$\mathcal{B}_\Lambda := \mathcal{B}|_{V_\Lambda}. \quad (2.23)$$

Upon restriction to V_Λ , the polynomial operators $q_{A|I}$ of (2.16) read

$$q_{A|J} = u^{M_{A|J}} + \sum_{k=1}^{M_{A|J}} \hat{c}_{A|J}^{(k)} u^{M_{A|J}-k}. \quad (2.24)$$

These are monic polynomials of degree $M_{A|J}$ with operator-valued coefficients $\hat{c}_{A|J}^{(k)}$. The diagonalisation of $\hat{c}_{A|J}^{(k)}$ is the subject of a Bethe Ansatz. The degree $M_{A|J}$ has fixed value on each V_Λ , which can for instance be identified by explicit computations following Appendix C (see [64]), they have the following expression in terms of the fundamental weight $\Lambda = [\lambda_1, \dots, \lambda_m | \nu_1, \dots, \nu_n]$

$$M_{A|J} = \sum_{a \in A} \lambda_a + \sum_{j \in J} \nu_j, \quad (2.25)$$

and can be interpreted as the ‘‘magnon’’ numbers of the nested Bethe Ansätze.

For generic twist, (2.21) is a non-degenerate system of linear equations in the coefficients $\hat{c}_{a|i}^{(k)}$ that fixes $\hat{c}_{a|i}^{(k)}$ and thus $Q_{a|i}$ uniquely. Hence all the Q-operators are generated by the single-index Q-functions $Q_{a|\emptyset}$, $Q_{\emptyset|i}$. There are precisely L coefficients $\hat{c}_{a|\emptyset}^{(k)}$ and $\hat{c}_{\emptyset|i}^{(k)}$ as one can quickly conclude from (2.25) and the invariant value of the total charge $\sum_{a=1}^m \lambda_a + \sum_{j=1}^n \nu_j = L$.

We can use C_Λ – the set of all coefficients $\hat{c}_{a|\emptyset}^{(k)}$ and $\hat{c}_{\emptyset|i}^{(k)}$ – to polynomially generate $Q_{\bar{\emptyset}|\bar{\emptyset}}$ using (2.20). This operation is a supersymmetric generalisation of the Wronskian determinant in (1.3) and shall be denoted as $\text{SW}(C_\Lambda)(u)$,

$$\boxed{\text{SW}(C_\Lambda)(u) = \prod_{\ell=1}^L (u - \theta_\ell)}. \quad (2.26)$$

Note that we chose the normalisation of SW such that the leading u^L term is monic.

We call (2.26) the quantisation condition or the Wronskian Bethe equations (WBE). Its important feature is to provide exactly L equations on L variables (elements of the set C_Λ) and to contain L free parameters θ_ℓ . We shall denote this system of L equations as

$$\text{SW}_\ell(c) = \chi_\ell, \quad \ell = 1, 2, \dots, L, \quad (2.27)$$

where χ_ℓ are elementary symmetric polynomials of $\theta_1, \theta_2, \dots, \theta_L$. Dependence on inhomogeneities only through their symmetric combinations χ_ℓ will be very important in our studies. Quite often, we will consider χ_ℓ as independent variables instead of inhomogeneities.

We shall consider the quantisation condition as an equation both on the level of operators denoted uniformly as \hat{c}_ℓ , $\ell = 1, \dots, L$ and on the level of abstract variables denoted as c_ℓ . We shall show eventually that *any* c_ℓ solving (2.26) provides eigenvalues for \hat{c}_ℓ 's.

Example:

Consider a $GL(3)$ spin chain of length $L = 3$, and the weight subspace V_Λ with $\Lambda = [2, 1, 0]$. The Q-system is parameterised by

$$\begin{aligned} Q_1 &\propto x_1^{u/\hbar} \times (u^2 + \hat{c}_1^{(1)}u + \hat{c}_1^{(2)}), \\ Q_2 &\propto x_2^{u/\hbar} \times (u + \hat{c}_2^{(1)}), \\ Q_3 &\propto x_3^{u/\hbar}. \end{aligned} \quad (2.28)$$

$\text{SW} \propto W(Q_1, Q_2, Q_3) = \det_{1 \leq a, b \leq 3} Q_a(u + \hbar(2-b))$. Set for simplicity $x_3 = 3, x_2 = 2, x_1 = 1$, then (2.27) becomes explicitly

$$\begin{aligned} (\hat{c}_1^{(2)}\hat{c}_2^{(1)} - \hbar(\hat{c}_1^{(2)} + \frac{5}{2}\hat{c}_1^{(1)}\hat{c}_2^{(1)}) + \frac{\hbar^2}{2}(9\hat{c}_1^{(1)} + 7\hat{c}_2^{(1)}) - \frac{15}{2}\hbar^3) & - \theta_1 \theta_2 \theta_3 \\ + u \times (\hat{c}_1^{(2)} + \hat{c}_1^{(1)}c_2^{(1)} + \hbar(-\frac{7}{2}\hat{c}_1^{(1)} - 5\hat{c}_2^{(1)}) + \frac{25}{2}\hbar^2) & = +u \times (\theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3) \\ + u^2 \times (\hat{c}_1^{(1)} + \hat{c}_2^{(1)} - 6\hbar) & + u^2 \times (-\theta_1 - \theta_2 - \theta_3) \\ + u^3 & + u^3 \end{aligned} \quad (2.29)$$

which yields us three equations satisfied by $c_1^{(1)}, c_1^{(2)}, c_2^{(1)}$, both on the level of operators and their eigenvalues.

Counting the number of solutions is easy in this example: one can derive a cubic equation on $c_2^{(1)}$ with a cubic term that never vanishes and furthermore we observe that $c_1^{(1)}, c_1^{(2)}$ follow uniquely if we fix the value of $c_2^{(1)}$. So there are always three solutions which is the dimension of V_Λ . For generic values of θ_ℓ , all solutions are distinct.

Example:

Consider a $\mathfrak{gl}_{1|1}$ spin chain of length $L = 3$, and V_Λ with $\Lambda = [\lambda|\nu] = [2|1]$. We

parameterise the Q-system by

$$\begin{aligned} Q_{1|\emptyset} &\propto x^{+u/\hbar} \times (u^2 + \hat{c}_{1|\emptyset}^{(1)}u + \hat{c}_{1|\emptyset}^{(2)}), \\ Q_{\emptyset|1} &\propto y^{-u/\hbar} \times (u + \hat{c}_{\emptyset|1}^{(1)}). \end{aligned} \quad (2.30)$$

In this case $Q_\theta = \text{SW} \propto Q_{1|1}$, and one needs to compute $Q_{1|1}$ from the finite-difference equation $Q_{1|1}(u + \hbar/2) - Q_{1|1}(u - \hbar/2) = Q_{1|\emptyset}Q_{\emptyset|1}$ which supplies the equations on $c_{1|\emptyset}^{(1)}$, $c_{1|\emptyset}^{(2)}$, and $c_{\emptyset|1}^{(1)}$. For $x = 3, y = 1$ they are

$$\begin{aligned} &u^3 - u^2(\chi_1 - 3\hbar) + u(\chi_2 - 2\chi_1\hbar + \frac{3}{4}\hbar^2) - (\chi_3 - \chi_2\hbar + \frac{1}{4}\chi_1\hbar^2 - \frac{1}{4}\hbar^3) \\ &= (u^2 + c_{1|\emptyset}^{(1)}u + c_{1|\emptyset}^{(2)})(u + c_{\emptyset|1}^{(1)}), \end{aligned} \quad (2.31)$$

where $\chi_1 = \theta_1 + \theta_2 + \theta_3$, $\chi_2 = \theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3$, $\chi_3 = \theta_1\theta_2\theta_3$.

Counting solutions for this example is even simpler, and it is a good demonstration of when a supersymmetric system can be advantageous for finding the spectrum of the Bethe algebra. The l.h.s. of (2.31) is a degree-three polynomial with all coefficients known through the parameters of the theory. It has three roots, and one chooses which of these roots is $-c_{\emptyset|1}^{(1)}$ which fixes $c_{\emptyset|1}^{(1)}$. Values for other c 's follow. Hence there are three solutions which is indeed the dimension of the weight subspace.

Twist-less case

For the twist-less case, the symmetry of the system is enhanced as all the generators $\mathcal{E}_{\alpha\beta}$ commute with the Bethe algebra. Now, the Cartan subalgebra of $\mathfrak{gl}_{m|n}$ does not belong to the Bethe algebra and so the latter acting on the spin chain is definitely not maximal commutative. However, if we restrict ourselves to the weight subspace V_Λ^+ , maximal commutativity on this subspace will follow from completeness.

$$\mathcal{B}_\Lambda^+ := \mathcal{B}|_{V_\Lambda^+}. \quad (2.32)$$

We will typically drop the superscript $+$ and denote the restricted Bethe algebra as \mathcal{B}_Λ .

The Q-operators were constructed in [62–66] for the case of generic twist. Taking the twist-less limit is quite a tricky procedure [68] which was analysed substantially in sections 3.3 and 3.4 of [41]. The result of this analysis is that the below-presented properties that define the twist-less Q-system remain true at the level of operators.

Long representations Consider first a situation when Λ is a long representation of $\mathfrak{gl}_{m|n}$. The Young diagram of such a representation touches the corner of the fat hook, consider for instance the situation with $m \rightarrow m - r, n \rightarrow n - r$ in Figure 1.

In this case, the Q-operators $Q_{A|J}$ are polynomials in u of degree $M_{A|J}$ ¹³,

$$M_{A|J} = \sum_{a \in A} \hat{\lambda}_a + \sum_{j \in J} \hat{\nu}_j - \frac{|A|(|A| - 1)}{2} - \frac{|J|(|J| - 1)}{2} + |A||J|, \quad (2.33)$$

where $\hat{\lambda}_a, \hat{\nu}_i$ are the shifted weights defined as in Figure 1.

We will need mostly

$$M_{a|\emptyset} = \hat{\lambda}_a, \quad M_{\emptyset|i} = \hat{\nu}_i, \quad M_{a|i} = \hat{\lambda}_a + \hat{\lambda}_i + 1. \quad (2.34)$$

A few modifications have to be made to obtain an equivalent of (2.26). First, we notice that (2.21) fixes $Q_{a|i}$ only up to an additive constant and hence $\hat{c}_{a|i}^{(M_{a|i})}$ are new independent parameters used in the computation of $Q_{\bar{\emptyset}|\bar{\emptyset}}$. Second, the computation of $Q_{\bar{\emptyset}|\bar{\emptyset}}(u)$ and of other physically relevant quantities such as transfer matrices is invariant under the transformations

$$Q_{\emptyset|i} \rightarrow Q_{\emptyset|i} + \alpha Q_{\emptyset|j}, \quad Q_{a|\emptyset} \rightarrow Q_{a|\emptyset} + \beta Q_{b|\emptyset}. \quad (2.35)$$

We impose inequalities $i \geq j$ and $a \geq b$ to preserve the polynomial degrees (2.34). We fix these symmetry transformations by putting to zero all the coefficients $c_{\emptyset|i}^{(\hat{\nu}_j)}$ for $j \leq i$ and $c_{a|\emptyset}^{(\hat{\lambda}_b)}$ for $b \leq a$.

A straightforward counting shows that the set consisting of all non-zero $c_{a|\emptyset}^{(k)}, c_{\emptyset|i}^{(k)}$ combined together with $c_{a|i}^{(M_{a|i})}$ gives us exactly L variables. Denote this set by C_Λ . $Q_{\bar{\emptyset}|\bar{\emptyset}} \propto Q_\theta$ is unambiguously and polynomially reconstructed from C_Λ according to (2.20) supplemented with (2.21), we denote the corresponding operation again as $\text{SW}(C_\Lambda)(u)$ though explicit polynomial realisation of SW is different now.

In this modified setting, (2.26) holds.

Example:

Consider a \mathfrak{gl}_3 spin chain of length $L = 3$, and consider states in the representation $\Lambda^+ = \boxplus$. By the recipe of Figure 1, $\hat{\lambda}_1 = 4, \hat{\lambda}_2 = 2, \hat{\lambda}_3 = 0$. Then one generates the Bethe algebra by three Q-functions

$$\begin{aligned} Q_1 &= u^4 + \hat{c}_1^{(1)} u^3 + \hat{c}_1^{(3)} u, \\ Q_2 &= u^2 + \hat{c}_2^{(1)} u, \\ Q_3 &= 1. \end{aligned} \quad (2.36a)$$

We fixed $c_1^{(2)} = c_1^{(4)} = c_2^{(1)} = 0$ using symmetries of the system. The Wronskian condition (2.26) which is $\det_{1 \leq a, b \leq 3} Q_a(u + (2 - b)\hbar) \propto Q_\theta$ provides three equations to be

¹³For comparison with other literature, it might be needed to relabel Q-functions using the maps $a \rightarrow m + 1 - a, i \rightarrow n + 1 - i$. One checks the notation by asking for which a, i $Q_{a|i}$ is a polynomial of the smallest degree. In our conventions, it is $Q_{m|n}$.

satisfied by $c_1^{(1)}, c_1^{(3)}, c_2^{(1)}$:

$$c_1^{(3)} - \hbar^2(c_2^{(1)} - c_1^{(1)}) = 8\chi_3, \quad 3c_1^{(1)}c_2^{(1)} - 2\hbar^2 = 8\chi_2, \quad 3c_1^{(1)} + 6c_2^{(1)} = -8\chi_1, \quad (2.37)$$

it has two solutions.

Example:

Consider a $\mathfrak{gl}_{1|1}$ spin chain of length $L = 3$, again in the representation $\Lambda^+ = \boxplus$. By the recipe of Figure 1, $\hat{\lambda} = 1, \hat{\nu} = 2$. Then we use (2.33) to deduce the degree of Q-functions and get

$$\begin{aligned} Q_{1|0} &= u + \hat{c}_{1|0}, \\ Q_{0|1} &= u + \hat{c}_{0|1}. \end{aligned} \quad (2.38a)$$

Equation $Q_{1|1}^+ - Q_{1|1}^- = Q_{1|0}Q_{0|1}$ provides $Q_{1|1}$ up to an additive constant $c_{1|1}$,

$$Q_{1|1} \propto u^3 + \frac{3}{2}(\hat{c}_{1|0} + \hat{c}_{0|1})u^2 + (3\hat{c}_{1|0}\hat{c}_{0|1} - \frac{1}{4}\hbar^2)u + \hat{c}_{1|1}, \quad (2.39)$$

which together with $c_{1|0}$ and $c_{0|1}$ yields three variables that generate the Bethe algebra. The Wronskian condition (2.26) is $Q_{1|1} \propto Q_\theta$, it implies the equations on c 's:

$$c_{1|0} + c_{0|1} = -\frac{2}{3}\chi_1, \quad c_{1|0}c_{0|1} = \frac{1}{3}\chi_2 + \frac{1}{12}\hbar^2, \quad c_{1|1} = -\chi_3 \quad (2.40)$$

which have two solutions.

Short representations The Young diagram of a short representation does not touch the internal corner of the $m|n$ fat hook. Define r according to Figure 1. Introduce sets $\mathbf{A} = \{m - r + 1, m - r + 2, \dots, m\}$ and $\mathbf{J} = \{n - r + 1, n - r + 2, \dots, n\}$, and label all Q-functions as $Q_{AA_0|JJ_0}$, where A_0 is a multi-index from \mathbf{A} and J_0 is a multi-index from \mathbf{J} . Then the properties of the Q-functions can be described as follows [41]: If $|A_0| = |J_0|$ then $Q_{AA_0|JJ_0} = Q_{A|J}$ and, if $|A_0| \neq |J_0|$ then $Q_{AA_0|JJ_0}$ are not uniquely defined in the twist-less limit but also such Q-operators appear in the physically-relevant quantities, such as transfer matrices and $Q_{\bar{\emptyset}|\bar{\emptyset}}$, in combinations that *vanish* in the twist-less limit.

The described property allows us to restrict the $\mathfrak{gl}_{m|n}$ Q-system to the $\mathfrak{gl}_{m-r|n-r}$ Q-system defined as $Q_{A|J}^{\text{rest}} = Q_{AA|JJ}$ which has the property $Q_{\bar{\emptyset}|\bar{\emptyset}}^{\text{rest}} = Q_{\mathbf{A}|\mathbf{J}} = Q_{\bar{\emptyset}|\bar{\emptyset}} = 1$ and $Q_{\bar{\emptyset}|\bar{\emptyset}}^{\text{rest}} = Q_{1\dots m-r|1\dots n-r}^{\text{rest}} = Q_{\bar{\emptyset}|\bar{\emptyset}}$. This subsystem is sufficient to generate the Bethe algebra. Since an originally short representation becomes long from the point of view of $\mathfrak{gl}_{m-r|n-r}$ subalgebra and since the polynomial degrees are correctly captured by (2.33), we can use the same logic as for the long representations and formulate the supersymmetric Wronskian condition (2.26) using C_Λ of the $\mathfrak{gl}_{m-r|n-r}$ Q-system.

Example:

The representation \boxplus can be considered as a short one of the $\mathfrak{gl}_{2|2}$ algebra. Then $\mathbf{A} = \{2\}, \mathbf{J} = \{2\}$, and so all the physical information is contained in the functions

$Q_{12|12} = Q_{1|1}$, $Q_{12|2} = Q_{1|\emptyset}$, $Q_{2|12} = Q_{\emptyset|1}$, $Q_{2|2} = Q_{\emptyset|\emptyset}$. The Wronskian is given by

$$\text{SW} = Q_{1|1}Q_{2|2} - Q_{1|2}Q_{2|1}. \quad (2.41)$$

While $Q_{1|2}$ and $Q_{2|1}$ are not uniquely defined in the twist-less limit, any prescription would imply that at least either $Q_{1|2}$ or $Q_{2|1}$ vanish and so their product vanishes as well. Given that $Q_{2|2} = Q_{\emptyset|\emptyset} = 1$, $\text{SW} = Q_\theta$ implies $Q_{1|1} = Q_\theta$ which fully parallels the above-described $\mathfrak{gl}_{1|1}$ example.

3 Completeness

So far we introduced \mathcal{B}_Λ – the restriction of the Bethe algebra to the weight subspace U_Λ (which is V_Λ or V_Λ^+). It is generated by the restriction of the Q-operators who in turn are (twisted) polynomials of the spectral parameter. We also selected precisely L coefficients of these polynomials assembled into the set C_Λ and explained how they are used to generate the whole \mathcal{B}_Λ .

From now on, the elements of C_Λ are labelled in a uniform manner as c_ℓ , $\ell = 1, \dots, L$. It is important to articulate what notation c_ℓ means exactly. If it is an explicit matrix acting on U_Λ then we denote it as \hat{c}_ℓ . In contrast, we agree to denote by c_ℓ without hat abstract commuting variables that have, by definition, only one property: they satisfy Wronskian Bethe equations (2.27).

A one-to-one correspondence between c_ℓ solving WBE and eigenvalues of \hat{c}_ℓ will be established later, in Section 4. In the current section, we show that WBE have precisely the right number of solutions $d_\Lambda = \dim_{\mathbb{C}} U_\Lambda$. This property is usually referred to as *completeness*, why this naming is justified was discussed in the introduction.

3.1 Analytic description

First, we develop some intuition about analytic description of the Wronskian Bethe equations $\text{SW}_\ell(c) = \chi_\ell$. Think about them as a polynomial map

$$\begin{aligned} \text{SW} : \quad \mathbb{C}^L &\longrightarrow \mathbb{C}^L, \\ (c_1, \dots, c_L) &\longmapsto (\chi_1 = \text{SW}_1(c), \dots, \chi_L = \text{SW}_L(c)). \end{aligned} \quad (3.1)$$

Denote by $\mathcal{C} \simeq \mathbb{C}^L$ the domain of definition of the map and by $\mathcal{X} := \text{SW}(\mathcal{C})$ its image.

Surjectivity. The map SW is in fact surjective, that is $\mathcal{X} \simeq \mathbb{C}^L$ which means that the Wronskian relations (2.27) have at least one solution for any complex value $\bar{\chi}_\ell$ of χ_ℓ . Indeed, matrix coefficients of \hat{c}_ℓ are polynomials in θ_ℓ , *e.g.* by construction of Baxter Q-operators, and so they are defined for any numerical value $\bar{\theta}_\ell$. Furthermore \hat{c}_ℓ commute and so they have at least one common eigenvector $\mathbf{u}(\theta)$. Eigenvalues of \hat{c}_ℓ on this vector satisfy (2.27) and so they provide a solution to $\text{SW}_\ell(c) = \bar{\chi}_\ell$ for $\bar{\chi}_\ell = \chi_\ell(\bar{\theta})$.

Critical and regular points. Denote by $\mathcal{C}_{\text{crit}}$ the set of all the critical (degeneration) points c where the differential of SW is not invertible. Its image $\mathcal{X}_{\text{crit}} \equiv \text{SW}(\mathcal{C}_{\text{crit}})$ shall be called the set of critical points χ . Using *e.g.* Sard's theorem one states that $\mathcal{X}_{\text{crit}}$ is of measure zero

in \mathcal{X} . The complement to $\mathcal{C}_{\text{crit}}$, resp. $\mathcal{X}_{\text{crit}}$, shall be called domain of regular points c , resp. χ . Restricted to the regular points, the map SW is locally a diffeomorphism, *i.e.* for each point $c \notin \mathcal{C}_{\text{crit}}$ there is a neighbourhood of SW(c) where SW can be smoothly inverted. This implies that all solutions to the Bethe equations are *distinct* in a neighbourhood around a regular point χ . This also shows that in such a neighbourhood the fibers of SW are all finite and of the same cardinality (SW is polynomial and so it cannot have infinite discrete fibers).

Properness. All solutions c_ℓ are bounded at any finite value of θ_ℓ 's or, in more abstract terms, the inverse image of a compact set is compact. SW is then said to be proper. This very important technical point is proved in two independent ways: using the fact that Q -operators have bounded spectrum, as is explained in the remark on page 32; and by a direct analysis of the equations themselves, in Appendix D.1 for the twisted case and Appendix D.2 for the twist-less case.

Path-connectivity $\mathcal{C}_{\text{crit}}$ can be easily described as $\det \frac{\partial \text{SW}_\ell}{\partial c_\ell} = 0$ which is just a polynomial equation on c_ℓ that, obviously, defines a domain of (complex) co-dimension 1.

This implies that any two solutions c_ℓ and c'_ℓ can be connected by a smooth path γ that avoids the singular domain $\mathcal{C}_{\text{crit}}$. We can always choose γ such that its image SW(γ) also passes only through regular points of \mathcal{X} . Note that one or both points c_ℓ and c'_ℓ can actually belong to $\mathcal{C}_{\text{crit}}$. So any singular solution can be obtained as a limit of regular solutions. Sporadic solutions, defined as solutions that exist only for some subspace of points in \mathcal{X} cannot exist either by the same argument ¹⁴.

We in particular conclude that for *any* choice of χ_ℓ , the number of solutions of Bethe equations is less or equal to d_Λ , where d_Λ is defined as the number of solutions at regular points of \mathcal{X} (this number does not depend on $\chi_\ell \notin \mathcal{X}_{\text{crit}}$ since the regular domain of \mathcal{X} is path-connected).

Finiteness By definition, a map is called finite if it is proper and its fibers at all points are finite ¹⁵, so SW is an example of such a map. This property will be used later.

As we have established, all solutions to the Bethe equations (2.27) are distinct for χ_ℓ being in the regular domain of \mathcal{X} . Some solutions coincide if $\chi_\ell \in \mathcal{X}_{\text{crit}}$, and so the number of distinct solutions is smaller. It is typical to count solutions with multiplicities in such a case. When we deal with equations in several variables, the notion of multiplicity requires an appropriate formalism to be introduced which is our next goal.

3.2 How to count solutions with multiplicity

Starting from now, we will gradually introduce an algebraic formalism to analyse the Wronskian Bethe equations. We will be using standard terminology from commutative algebra

¹⁴For an example of equations with sporadic solutions, consider $x(x-1) = 0, \theta x = 0$. For all $\theta \neq 0$ there is only one solution $x = 0$. However, for $\theta = 0$ there is one extra sporadic solution $x = 1$.

¹⁵The concept of finite morphism is usually defined in a more general set-up using a rather abstract algebraic formalism. Here we are working with analytic varieties when the general ‘‘algebraic’’ definition is equivalent to the ‘‘topological’’ definition that we are using, see [74, 75].

which is briefly summarised in Appendix A.1.

Let us introduce a polynomial ring \mathcal{W}_Λ that shall be called the *Wronskian algebra* and which is defined as follows. Consider $\mathbb{C}[\chi][c]$ – the algebra of polynomials in variables $\chi_1, \chi_2, \dots, \chi_L, c_1, \dots, c_L$; and $\mathcal{I}_\Lambda = \langle \chi_1 - \text{SW}_1(c), \chi_2 - \text{SW}_2(c), \dots, \chi_L - \text{SW}_L(c) \rangle$ – the ideal generated by the equations (2.27). Then

$$\mathcal{W}_\Lambda := \mathbb{C}[\chi][c]/\mathcal{I}_\Lambda. \quad (3.2)$$

Over \mathbb{C} , \mathcal{W}_Λ is obviously isomorphic to $\mathbb{C}[c]$. However, additionally, it is also naturally a $\mathbb{C}[\chi]$ -module. Namely, one defines the action of χ_ℓ on $\mathcal{W}_\Lambda \simeq \mathbb{C}[c]$ as follows: we multiply elements of \mathcal{W}_Λ by χ_ℓ and then replace χ_ℓ with $\text{SW}_\ell(c)$.

To link the $\mathbb{C}[\chi]$ -module structure with the Wronski map from the previous section we note that $\mathbb{C}[c]$ is the coordinate ring of \mathcal{C} and $\mathbb{C}[\chi]$ is the coordinate ring of \mathcal{X} . The map

$$\text{SW}^* : \mathbb{C}[\chi] \rightarrow \mathbb{C}[c], \quad \chi_\ell \mapsto \text{SW}_\ell(c) \quad (3.3)$$

used in the definition of the $\mathbb{C}[\chi]$ -action on \mathcal{W}_Λ is a pullback of (3.1).

The number of solutions to the Wronskian Bethe equations appears as follows in the algebraic context. We *specialise* the Wronskian algebra to the complex point $\bar{\chi}$ where we would like to count the solutions. Specialisation is defined as

$$\mathcal{W}_\Lambda(\bar{\chi}) := \mathcal{W}_\Lambda / \langle \chi_\ell - \bar{\chi}_\ell \rangle \simeq \mathbb{C}[c] / \langle \text{SW}_\ell(c) - \bar{\chi}_\ell \rangle. \quad (3.4)$$

Then, it is a standard result that the number of solutions of a polynomial set of equations $P_i(x_1, \dots, x_n) = 0$, $i = 1, \dots, m$ is equal to the dimension of the quotient ring $\mathcal{R} = \mathbb{C}[x] / \langle P_1, \dots, P_m \rangle$ (as a vector space over \mathbb{C}). Moreover, in the case when solutions degenerate, the dimension of the quotient ring is used as a *definition*¹⁶ of the algebraic number of solutions (*i.e.* counted with multiplicity). In our case, the quotient ring in question is $\mathcal{R} = \mathcal{W}_\Lambda(\bar{\chi})$, and so the algebraic number of solutions of the Wronskian Bethe equation at point $\bar{\chi}$ is equal to $\dim_{\mathbb{C}} \mathcal{W}_\Lambda(\bar{\chi})$. Since at all points $\bar{\chi}$ the number of solutions is finite $\dim_{\mathbb{C}} \mathcal{W}_\Lambda(\bar{\chi}) < +\infty$.

To see how this definition comes about in practice, consider the regular representation of the (finite-dimensional) quotient ring \mathcal{R} which is a map from elements of the ring to the ring endomorphisms defined by the ring multiplication. We can describe this map in terms of explicit matrices. Let $\mathbf{b}_1, \dots, \mathbf{b}_r$ be some basis elements of \mathcal{R} . Then, for any $X \in \mathcal{R}$, one has $X \mathbf{b}_i = \sum_{j=1}^r \check{X}_{ij} \mathbf{b}_j$, where $\check{X}_{ij} \in \mathbb{C}$. The regular representation maps X to the matrix \check{X} whose components are \check{X}_{ij} .

Example:

Consider $\mathcal{R} = \mathbb{C}[x] / \langle x^2 - ax + b \rangle$. Elements x and 1 span \mathcal{R} , choose them as basis

¹⁶Again, the general definition of multiplicity in the full formalism of algebraic geometry is much more intricate but in our case it is equivalent to the one we use.

elements. Then one has $x \cdot x = x^2 = ax - b$, $x \cdot 1 = x$ and so

$$\check{x} = \begin{pmatrix} 0 & 1 \\ -b & a \end{pmatrix}. \quad (3.5)$$

It is easy to prove that the image of the regular representation is isomorphic to the algebra \mathcal{R} . This allows one to understand properties of a polynomial ring in a more familiar setting of a matrix algebra that we denote as $\check{\mathcal{R}}$.

By the isomorphism, $P_i(\check{x}_1, \dots, \check{x}_n) = 0$ for $i = 1, 2, \dots, m$. So all joint eigenvalues of $\check{x}_1, \dots, \check{x}_n$ are solutions of the set of equations. And each solution should be one of the joint eigenvalues (to see this, take $\sum_j (\check{x}_k)_{ij} \mathbf{b}_j = x_k \mathbf{b}_i$ and evaluate x_k and \mathbf{b}_i , who are polynomials in x_k , to numerical values corresponding to the solution of interest).

Hence, when \check{x}_ℓ are diagonalisable then it is clear that the number of solutions is equal to the size of the matrix which is the same as the dimension of the quotient ring. Moreover, all solutions should be distinct (otherwise, isomorphism between $\check{\mathcal{R}}$ and \mathcal{R} won't hold).

When \check{x}_ℓ are not diagonalisable, one could expect that the solutions degenerate and that the multiplicity of degeneration is the size of the corresponding Jordan block, as in (3.5) when $a = b = 0$. But since commuting matrices are not always simultaneously jordanisable, this intuitive picture should be slightly updated: Any commuting set of matrices, $\check{\mathcal{R}}$ in our case, admits a Dunford-Jordan-Chevalley decomposition, namely there is a basis where all the matrices take the form $D + N$, where D is diagonal and N is upper-triangular with zeros on the diagonal. Moreover, all elements of N form a subalgebra in $\check{\mathcal{R}}$ known as the nil-radical $\text{Nil}(\check{\mathcal{R}})$ which is the ideal of all nilpotent elements of the ring. The quotient algebra $\text{diag}(\check{\mathcal{R}}) \equiv \check{\mathcal{R}}/\text{Nil}(\check{\mathcal{R}})$ is isomorphic to the algebra of matrices from D .

Resorting to the regular representation was of course optional, the concepts of the nil-radical $\text{Nil}(\mathcal{R})$ and the quotient $\text{diag}(\mathcal{R})$ exist for any (commutative) ring. In summary, one has a short exact sequence

$$0 \longrightarrow \text{Nil}(\mathcal{R}) \longrightarrow \mathcal{R} \longrightarrow \text{diag}(\mathcal{R}) \longrightarrow 0. \quad (3.6)$$

$\dim_{\mathbb{C}} \text{diag}(\mathcal{R})$ is precisely the number of *distinct* solutions to the polynomial equations. $\dim_{\mathbb{C}} \text{Nil}(\mathcal{R})$ counts the amount of degeneration in solutions, and

$$\dim_{\mathbb{C}} \mathcal{R} = \dim_{\mathbb{C}} \text{diag}(\mathcal{R}) + \dim_{\mathbb{C}} \text{Nil}(\mathcal{R}) \quad (3.7)$$

is the total number of solutions counted with multiplicity.

We would like to emphasise that $\dim_{\mathbb{C}} \mathcal{R}$ is both the dimension of the quotient ring and the dimension of its regular representation (size of matrices). Eigenspaces of $\check{\mathcal{R}}$ are all of dimension one¹⁷ and $\check{\mathcal{R}}$ is a maximal commutative subalgebra of $\text{End}(\mathcal{R})$. This remark will become important in our study of the Bethe algebra.

¹⁷Not to confuse with degeneration of solutions of polynomial equations. By definition, eigenspaces of a matrix \check{X} are those that are annihilated by $\lambda - \check{X}$. In contrast, degenerate solutions correspond to existence of vectors that are annihilated by $(\lambda - \check{X})^n$ for $n > 1$.

3.3 Wronskian algebra is a free $\mathbb{C}[\chi]$ -module

The following very powerful result can be proven about the Wronskian algebra:

Proposition 3.1. \mathcal{W}_Λ is a free $\mathbb{C}[\chi]$ -module.

Proof. SW^* (3.3) is a ring morphism from $\mathbb{C}[\chi]$ to $\mathbb{C}[c]$ making \mathcal{W}_Λ a $\mathbb{C}[\chi]$ -algebra and therefore a $\mathbb{C}[\chi]$ -module. On the other hand, we can view SW (3.1) as an algebraic morphism from the variety $\mathcal{C} \simeq \mathbb{C}^L$ (for the current discussion, the affine space \mathbb{A}^L) to $\mathcal{X} \simeq \mathbb{A}^L$. We know that all the fibres of SW are finite sets and are therefore of dimension 0. Moreover, \mathbb{A}^L , as an algebraic variety, is regular and (therefore) Cohen-Macaulay. Then, by a general result (sometimes called “miracle flatness theorem”) SW^* is a flat ring morphism and so \mathcal{W}_Λ is a flat $\mathbb{C}[\chi]$ -module, see for example [76, 77]. Since $\mathbb{C}[\chi]$ is Noetherian and \mathcal{W}_Λ is finitely-generated as a $\mathbb{C}[\chi]$ -module (because SW is finite) it is actually projective [78]. Finally, by the Quillen–Suslin theorem [79] it is free. \square

The above proof looks very short, however it is based on several abstract results from algebraic geometry. In appendix A we provide an elementary study of the Wronskian algebra which helps in understanding the logic behind the above proof.

By definition of a free module, \mathcal{W}_Λ has a basis – a collection of elements $\mathbf{b}_1, \dots, \mathbf{b}_r$ such that any element $\mathbf{a} \in \mathcal{W}_\Lambda$ is represented in a unique way as

$$\mathbf{a} = k_1 \mathbf{b}_1 + \dots + k_r \mathbf{b}_r, \quad (3.8)$$

where $k_i \in \mathbb{C}[\chi]$. It is easy to prove that $\mathbf{b}_1, \dots, \mathbf{b}_r$ remains a basis after specialisation (3.4) which leads to an immediate corollary of Proposition 3.1:

Corollary 3.2. *The algebraic number of solutions of the Wronskian Bethe equations is the same for any value of $\bar{\chi}_\ell$.*

This number is equal to the number d_Λ of solutions to Bethe equations at regular points. Next section shows that this number is $\dim U_\Lambda$.

3.4 Number of solutions via Hilbert series

Let us find the value of d_Λ explicitly. With the help of Proposition 3.1, counting is reduced to a simple dimensional analysis as we shall describe now.

If one chooses a rule by which we assign degree 0 to the identity element and some positive integer degrees to other elements of a ring \mathcal{R} then we can define the ring filtration

$$\mathcal{F}_0 \subset \dots \mathcal{F}_k \subset \mathcal{F}_{k+1} \subset \dots,$$

where \mathcal{F}_k is the vector subspace of \mathcal{R} (over \mathbb{C}) spanned by all elements of degree not exceeding k . Grading assignment should be compatible with the ring structure meaning that for any k, k' and any ring element \mathbf{r}_k of degree k one has $\mathbf{r}_k \mathcal{F}_{k'} \subset \mathcal{F}_{k+k'}$.

A useful characterisation of a graded ring is given by its Hilbert series defined as

$$\text{ch}_{\mathcal{R}}(t) = \sum_{k=0}^{\infty} \dim_{\mathbb{C}}(\mathcal{F}_k/\mathcal{F}_{k-1}) t^k. \quad (3.9)$$

Since the Wronskian ring \mathcal{W}_{Λ} is \mathbb{C} -isomorphic to $\mathbb{C}[c_1, \dots, c_L]$, computing its Hilbert series is particularly simple. It is just given by

$$\text{ch}_{\mathcal{W}_{\Lambda}}(t) = \prod_{\ell=1}^L \frac{1}{1 - t^{\deg c_{\ell}}}. \quad (3.10)$$

Recall that c_{ℓ} is a selected subset of $c_{A|J}^{(k)}$. Define $\deg c_{A|J}^{(k)} = k$. With the labeling $c_{A|J}^{(k)}$ of (2.24), this grading is consistent with the ring structure because the latter follows from the relations between Q-functions which are polynomials in u . Then we have

Lemma 3.3. *The Hilbert series of the Wronskian algebra \mathcal{W}_{Λ} of the twisted system for $\Lambda = [\lambda_1, \dots, \lambda_m; \nu_1, \dots, \nu_n]$ is*

$$\text{ch}_{\Lambda}(t) := \text{ch}_{\mathcal{W}_{\Lambda}}(t) = \prod_{a=1}^m \prod_{k=1}^{\lambda_a} \frac{1}{1 - t^k} \prod_{i=1}^n \prod_{k=1}^{\nu_i} \frac{1}{1 - t^k}. \quad (3.11a)$$

The Hilbert series of the Wronskian algebra \mathcal{W}_{Λ} of the $\mathfrak{gl}_{m|n}$ twist-less system with $\Lambda = \Lambda^+$ being a Young diagram depends on the Young diagram Λ^+ alone and is given by

$$\text{ch}_{\Lambda^+}(t) := \text{ch}_{\mathcal{W}_{\Lambda}}(t) = \prod_{(a,s) \in \Lambda^+} \frac{1}{1 - t^{h_{a,s}}}, \quad (3.11b)$$

where the product runs over all boxes of Λ^+ and $h_{a,s}$ is the hook length at box (a, s) .

Proof. The result in the twisted case is immediately obvious from (2.25) and the fact that c_1, \dots, c_L are precisely all $c_{a|\emptyset}^{(k)}$ and $c_{\emptyset|i}^{(k)}$. For the twist-less case, one needs to perform a little analysis on precisely what $c_{A|J}^{(k)}$ generate the Wronskian algebra. One can do it by filling the boxes of Λ^+ with degrees of the variables c_{ℓ} in a special way. This procedure outlined in Figure 2 clearly establishes a bijection with the lengths of the corresponding hooks. \square

To make the above-introduced filtration compatible with the $\mathbb{C}[\chi]$ -action on \mathcal{W}_{Λ} , one needs to define $\deg \chi_{\ell} = \ell$. This is a simple reflection of the fact that the Wronskian relations $\chi_{\ell} = \text{SW}_{\ell}(c)$ come from equating coefficients between polynomials in u .

Because \mathcal{W}_{Λ} is a free $\mathbb{C}[\chi]$ -module, $\dim \mathcal{W}_{\Lambda}(\bar{\chi}) = \frac{\text{ch}_{\mathcal{R}}(t)}{\text{ch}_{\mathbb{C}[\chi]}(t)}|_{t=1}$ for any $\bar{\chi}$. We then compute $\text{ch}_{\mathbb{C}[\chi]}(t) = \prod_{\ell=1}^L \frac{1}{1-t^{\ell}}$ and easily conclude




Figure 2: Degrees of c_α that generate the twist-less Wronskian algebra. On the left: The blue numbers show degrees of $c_{a|\emptyset}^{(k)}$, where $a = 1, 2, \dots$ is the corresponding row of the Young diagram. The red numbers show degrees of $c_{\emptyset|i}^{(k)}$ where $i = 1, \dots$ is the corresponding column. Crosses mean the terms $c_{a|\emptyset}^{(k)}$ and $c_{\emptyset|i}^{(k)}$ that are excluded by symmetry reasons, see the discussion after (2.34). For instance, the term $c_{1|\emptyset}^{(9)}$ (the constant term of $Q_{1|\emptyset}$) is cancelled in the linear combination $Q_{1|\emptyset} + \alpha Q_{4|\emptyset}$. On the right: the same blue/red numbers compressed to the right/bottom after the crosses are removed. Note that this arrangement corresponds to the hook lengths. The green rectangular area has exactly as many boxes as number of physically-relevant functions $Q_{a|i}$. The constant terms of these functions, $c_{a|i}^{(k)}$ with $k = \deg Q_{a|i} = \hat{\lambda}_a + \hat{\lambda}_i + 1$, are independent variables used in generation of the Wronskian algebra. Their degrees, shown in green, match the hook lengths as well.

Theorem 3.4. *The number of solutions of the Wronskian Bethe equations counted with algebraic multiplicity is equal to*

$$\dim \mathcal{W}_\Lambda(\bar{\chi}) = \dim V_\Lambda = \frac{L!}{\prod_{a=1}^m \lambda_a! \prod_{i=1}^n \nu_i!} \quad \text{for the twisted case,} \quad (3.12a)$$

$$\dim \mathcal{W}_\Lambda(\bar{\chi}) = \dim V_\Lambda^+ = \frac{L!}{\prod_{(a,s) \in \Lambda} h_{a,s}} \quad \text{for the twist-less case.} \quad (3.12b)$$

□

In summary, the number of solutions to the Wronskian Bethe equations is the correct one and hence the equations are complete for any numerical choice of the inhomogeneities.

4 Faithfulness

Until now, we have concentrated on the Wronskian algebra \mathcal{W}_Λ to study the properties of the Bethe equations (2.27). In particular we have shown that they have the expected number of solutions counted with multiplicity. Now we will establish a bijective correspondence between these solutions and eigenvalues (in degenerate cases, trigonal blocks) of the Bethe algebra. Mathematically, this correspondence is formulated as an isomorphism between the Wronskian algebra and the Bethe algebra. The map $\varphi : c_\ell \mapsto \hat{c}_\ell$ from the Wronskian algebra to the Bethe algebra is tautologically surjective, and it is its injectivity (faithfulness) that we need to prove.

In this section, we shall first establish the isomorphism over the polynomial ring $\mathbb{C}[\chi]$ (which in practice means “in general position”) and then prove that the isomorphism also

holds for any numerical value $\bar{\chi}_\ell$ of χ_ℓ . A sufficient condition for the isomorphism to hold is that inhomogeneities that solve $\bar{\chi}_\ell = \chi_\ell(\bar{\theta}_1, \dots, \bar{\theta}_L)$ satisfy $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$.


4.1 Isomorphism between Wronskian and Bethe algebra

Recall that \mathcal{B}_Λ – the Bethe algebra restricted to U_Λ – can be viewed as an algebra of operators generated by \hat{c}_ℓ . As we learned from the previous section, it is beneficial to first keep inhomogeneities as indeterminates, and this is what we are going to do with the Bethe algebra as well. Then \mathcal{B}_Λ is naturally a subalgebra of $\text{End}(U_\Lambda) \otimes \mathbb{C}[\theta]$, however one should be careful because we define \mathcal{B}_Λ as the algebra generated from \hat{c}_ℓ by considering any polynomials in \hat{c}_ℓ with coefficients from \mathbb{C} , and not from $\mathbb{C}[\theta]$. This is a non-trivial remark because of the following

Lemma 4.1. *If $p \in \mathbb{C}[\theta]$ and $p \times \mathbb{1} \in \mathcal{B}_\Lambda$ then p is a symmetric polynomial in inhomogeneities.*

Proof. Recall that the spin chain vector space is a tensor product of L copies of $\mathbb{C}^{m|n}$. Introduce $r_\ell(\theta) = (\theta_\ell - \theta_{\ell+1})\mathcal{P}_{\ell,\ell+1} + \hbar\mathbb{1}$, where $\mathcal{P}_{\ell,\ell+1}$ is the graded permutation of two copies of $\mathbb{C}^{m|n}$ at the ℓ 'th and the $(\ell + 1)$ 'th position of $(\mathbb{C}^{m|n})^{\otimes L}$. This is an intertwining operator that satisfies

$$r_\ell \text{ev}_{(\theta_1, \dots, \theta_\ell, \theta_{\ell+1}, \dots, \theta_L)}(T_{ij}) = \text{ev}_{(\theta_1, \dots, \theta_{\ell+1}, \theta_\ell, \dots, \theta_L)}(T_{ij}) r_\ell \quad (4.1)$$

which can be graphically represented as 

is essentially the Yang-Baxter equation in the physical channel.

If we also introduce Π_ℓ – permutation of inhomogeneities θ_ℓ and $\theta_{\ell+1}$ in $\mathbb{C}[\theta]$ then $\Pi_\ell r_\ell$ commutes with the Yangian action and hence with the Bethe algebra. Therefore, if there is any equation $P(\hat{c}_1, \dots, \hat{c}_L) = p(\theta_1, \dots, \theta_L) \times \mathbb{1}$ that holds so will hold $P(\hat{c}_1, \dots, \hat{c}_L) = p(\theta_{\sigma(1)}, \dots, \theta_{\sigma(L)}) \times \mathbb{1}$ for any $\sigma \in \mathcal{S}_L$. Since inhomogeneities are independent, this is only consistent if p is a symmetric polynomial.

To be prudent, we notice that the derivation of $P(\hat{c}_1, \dots, \hat{c}_L) = p(\theta_{\sigma(1)}, \dots, \theta_{\sigma(L)}) \times \mathbb{1}$ emerges from the following argument:

$$\begin{aligned} 0 &= \Pi_\ell r_\ell (P(\hat{c}_1, \dots, \hat{c}_L) - p(\theta_1, \dots, \theta_\ell, \theta_{\ell+1}, \dots, \theta_L) \times \mathbb{1}) \Pi_\ell r_\ell \\ &= (P(\hat{c}_1, \dots, \hat{c}_L) - p(\theta_1, \dots, \theta_{\ell+1}, \theta_\ell, \dots, \theta_L) \times \mathbb{1}) (\Pi_\ell r_\ell)^2 =: A \times (\Pi_\ell r_\ell)^2. \\ (\Pi_\ell r_\ell)^2 &= (\hbar^2 - (\theta_\ell - \theta_{\ell+1})^2) \times \mathbb{1}. \end{aligned}$$

Then, because components of the matrix A are polynomials in θ_ℓ , $0 = A \times (\hbar^2 - (\theta_\ell - \theta_{\ell+1})^2)$ is only possible if $A = 0$. \square

We see that for instance $\theta_\ell \times \mathbb{1}$ does not belong to the Bethe algebra, except for $L = 1$. But any symmetric polynomial in inhomogeneities (times the identity operator) is an element of \mathcal{B}_Λ because of (2.27). So \mathcal{B}_Λ is naturally a $\mathbb{C}[\chi]$ -algebra.

Recall now also the definition of the Wronskian algebra \mathcal{W}_Λ (3.2) which is a polynomial algebra generated by c_ℓ and which is also a $\mathbb{C}[\chi]$ -algebra.

There is a potential difference between \mathcal{W}_Λ and \mathcal{B}_Λ . The generators of the Wronskian algebra, by definition, satisfy only (2.27). The generators of the Bethe algebra are certain

explicit operators and they could in principle satisfy some other additional constraints. However, we can show that they do not.

Theorem 4.2. *The map φ defined as*

$$\varphi : \mathcal{R} \longrightarrow \mathcal{B}_\Lambda, \quad \varphi : c_\ell \longmapsto \hat{c}_\ell \quad (4.2)$$

is an isomorphism of $\mathbb{C}[\chi]$ -algebras.

In other words, \hat{c}_ℓ not only satisfy (2.27) but any polynomial relation between \hat{c}_ℓ with coefficients in $\mathbb{C}[\chi]$ should follow from (2.27).

The proof below makes precise the following argument: as there are as many variables c_ℓ as the parameters χ_ℓ in (2.27), there cannot exist an extra relation between the variables because it would imply a relation between the parameters which are known to be independent.

Proof. We need to show that the exhibited map φ is a well-defined (consistent) morphism and that it is surjective and injective. It is well-defined because Q-operators form a commutative algebra and they satisfy (2.27) from the very derivation of this relation. It is surjective because \hat{c}_ℓ generate \mathcal{B}_Λ .

The non-trivial part is the injectivity (faithfulness). To prove this we take an element $P \in \mathcal{W}_\Lambda$ (so just a polynomial in the variables c_ℓ and χ_ℓ modulo relations in the ideal) and show that $\varphi(P) = 0$ implies $P = 0$.

Note that P can be viewed as a polynomial in c_ℓ with constant coefficients as all occurrences of χ_ℓ can be replaced by $\text{SW}_\ell(c)$. Since $\varphi(P(c)) = P(\hat{c})$, P has to vanish every time when c_ℓ are eigenvalues of \hat{c}_ℓ on a joint eigenvector. Then it suffices to construct enough of such eigenvalues to conclude that $P = 0$.

To this end consider $\chi \notin \mathcal{X}_{\text{crit}}$. There exists a neighbourhood \mathcal{O}_χ where all the solutions of (2.27) are distinct and can be parameterised by d_Λ diffeomorphisms SW_i^{-1} from \mathcal{O}_χ to d_Λ non-intersecting open sets U_i in \mathcal{C} .

We know that for all points of \mathcal{O}_χ the Bethe algebra has at least one common eigenvector and that the corresponding eigenvalues of \hat{c}_ℓ provide a solution of (2.27). By choosing in some way exactly one eigenvector at each point of \mathcal{O}_χ we create a disjoint partition of \mathcal{O}_χ into d_Λ sets \mathcal{O}_i corresponding to points of \mathcal{O}_χ where the common eigenvector gives the i -th solution. The closure (in \mathcal{O}_χ) of one of the \mathcal{O}_i 's, say $\bar{\mathcal{O}}_1$, contains an L -dimensional ball \mathcal{O} . This is proved as follows¹⁸. Consider λ the Lebesgue measure on \mathcal{O}_χ normalised to 1. Then either $\lambda(\bar{\mathcal{O}}_1) = 1$ and therefore $\bar{\mathcal{O}}_1 = \mathcal{O}_\chi$ since its complementary in \mathcal{O}_χ is an open set of measure zero or $\lambda(\bar{\mathcal{O}}_1) < 1$ in which case by restricting to its complementary in \mathcal{O}_χ we are brought back to the same problem but with $d_\Lambda - 1$ sets. We conclude by induction.

Then P vanishes on $\text{SW}_1^{-1}(\mathcal{O}_1)$ and since the zeros of a polynomial form a closed set and SW_1^{-1} is a diffeomorphism on \mathcal{O}_χ it will also vanish on $\text{SW}^{-1}(\bar{\mathcal{O}}_1)$ which contains an L -dimensional ball. P being a polynomial in L variables thus implies $P = 0$. \square

¹⁸This can be also proven by arguing that the common eigenvector can be chosen continuously in which case taking closure is unnecessary as well.

One may ask whether there are some additional polynomial relations between \hat{c}_ℓ with coefficients being non-symmetric polynomials of inhomogeneities. This is not possible either as can be shown using a slightly updated version of Lemma 4.1, see Appendix A.4.

4.2 What can happen upon specialisation

Now we shall consider what happens with this isomorphism when inhomogeneities θ_ℓ get concrete numerical values. We call this procedure specialisation at point $\bar{\theta}$.

Specialisation of the Bethe algebra $\mathcal{B}_\Lambda(\bar{\theta})$ is replacing θ_ℓ with $\bar{\theta}_\ell$ in *all matrix entries* of the operators \hat{c}_ℓ . On the other hand, specialisation of the Wronskian algebra is

$$\mathcal{W}_\Lambda(\bar{\theta}) \equiv \mathcal{W}_\Lambda(\bar{\chi}_\ell = \chi_\ell(\bar{\theta})) \simeq \mathcal{W}_\Lambda / \langle \chi - \bar{\chi} \rangle. \quad (4.3)$$

Its image under the map φ is $\mathcal{B}_\Lambda / \langle (\chi(\theta) - \bar{\chi}) \times \mathbf{1} \rangle$ which explicitly means the following: replace χ_ℓ with its numerical value each time it *multiplies* some matrix belonging to \mathcal{B}_Λ . This operation is less restrictive than specialisation of the Bethe algebra and hence one can state that the morphism

$$\varphi_{\bar{\theta}} : \mathcal{W}_\Lambda(\bar{\theta}) \longrightarrow \mathcal{B}_\Lambda(\bar{\theta}) \quad (4.4)$$

is surjective but may have a non-zero kernel. We denote by Θ_{not} the set of $\bar{\theta}$ when $\varphi_{\bar{\theta}}$ is not an isomorphism.

Example:

Consider a Wronskian algebra \mathcal{W} realised by relations $c_1 + c_2 = \chi_1$ and $c_1 c_2 = \chi_2$ ^a. It is a free $\mathbb{C}[\chi_1, \chi_2]$ -module, for the basis one can choose $1, c_1$, and the ring multiplication rule follows from $c_1^2 - \chi_1 c_1 + \chi_2 = 0$. Then $\check{c}_1 = \begin{pmatrix} 0 & 1 \\ -\chi_2 & \chi_1 \end{pmatrix}$.

Consider two ‘‘Bethe algebras’’ $\mathcal{B}^{\text{good}}$ and \mathcal{B}^{bad} , with, respectively,

$$\check{c}_1^{\text{good}} = \begin{pmatrix} \theta_1 & 1 \\ 0 & \theta_2 \end{pmatrix}, \quad \text{and} \quad \check{c}_1^{\text{bad}} = \begin{pmatrix} \theta_1 & 0 \\ 0 & \theta_2 \end{pmatrix}. \quad (4.5)$$

They are both $\mathbb{C}[\chi_1, \chi_2]$ -isomorphic, as algebras, to \mathcal{W} . Note however that they realise non-isomorphic representations over $\mathbb{C}[\chi_1, \chi_2]$, *i.e.* there is no intertwiner matrix mapping $\check{c}_1^{\text{good}}$ to \check{c}_1^{bad} whose coefficients are polynomial in χ_1, χ_2 .

If we specialise at any point where $\bar{\theta}_1 \neq \bar{\theta}_2$, the corresponding $\varphi_{\bar{\theta}}$ would be an algebra isomorphism both for $\mathcal{B}^{\text{good}}(\bar{\theta})$ and $\mathcal{B}^{\text{bad}}(\bar{\theta})$, also there would be obviously an intertwiner over \mathbb{C} between ‘‘good’’ and ‘‘bad’’ representations making them isomorphic.

Now, let us specialise to a point $\bar{\theta}_1 = \bar{\theta}_2$. The specialised Wronskian algebra becomes a two-dimensional algebra over \mathbb{C} generated by $1, c_1$ and relation $(c_1 - \bar{\theta}_1)^2 = 0$. It is isomorphic to the algebra generated by $\check{c}_1 = \begin{pmatrix} 0 & 1 \\ -\bar{\theta}_1^2 & 2\bar{\theta}_1 \end{pmatrix}$ which cannot be diagonalised, *cf.* (3.5).

The Bethe algebra $\mathcal{B}^{\text{good}}(\bar{\theta})$ is also two-dimensional and isomorphic to $\mathcal{W}(\bar{\theta})$ whereas $\mathcal{B}^{\text{bad}}(\bar{\theta})$ is one-dimensional since $c_1 - \bar{\theta}_1$ is in the kernel of $\varphi_{\bar{\theta}}$.

^aUp to an isomorphism, it is the Wronskian algebra \mathcal{W}_{\square} with $\hbar = 0$ which is partially specialised

to e.g. $\theta_3 = 0$, cf. Appendix B.5

The difference between “good” and “bad” cases is in the presence of the nilpotent piece $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ in $\hat{\mathcal{C}}_1^{\text{good}}$ which becomes an element of $\mathcal{B}(\bar{\theta})$ each time $\bar{\theta}_1 = \bar{\theta}_2$. This well illustrates what happens in the general situation. As $\mathcal{B}_\Lambda(\bar{\theta})$ is a commutative algebra, we can define the short exact sequence (3.6) for it. Then we can state the following

Theorem 4.3. *In the map between the two sequences*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Nil}(\mathcal{W}_\Lambda(\bar{\theta})) & \longrightarrow & \mathcal{W}_\Lambda(\bar{\theta}) & \longrightarrow & \text{diag}(\mathcal{W}_\Lambda(\bar{\theta})) \longrightarrow 0 \\ & & \varphi_{\bar{\theta}}^{\text{nil}} \downarrow & & \varphi_{\bar{\theta}} \downarrow & & \varphi_{\bar{\theta}}^{\text{diag}} \downarrow \\ 0 & \longrightarrow & \text{Nil}(\mathcal{B}_\Lambda(\bar{\theta})) & \longrightarrow & \mathcal{B}_\Lambda(\bar{\theta}) & \longrightarrow & \text{diag}(\mathcal{B}_\Lambda(\bar{\theta})) \longrightarrow 0 \end{array}, \quad (4.6)$$

$\varphi_{\bar{\theta}}^{\text{diag}}$ is an isomorphism for any $\bar{\theta}$.

The isomorphism $\varphi_{\bar{\theta}}^{\text{diag}}$ literally means that the distinct solutions of the Wronskian Bethe equations are in one-to-one correspondence with the eigenspaces of the Bethe algebra (there are no non-physical solutions).

In the regular case when $\chi(\bar{\theta}) \notin \mathcal{X}_{\text{crit}}$, $\text{Nil}(\mathcal{W}_\Lambda(\bar{\theta})) = 0$ and so the theorem implies that the Wronskian and the Bethe algebras are isomorphic. Hence $\chi(\Theta_{\text{not}}) \subset \mathcal{X}_{\text{crit}}$.

In the degenerate case $\chi(\bar{\theta}) \in \mathcal{X}_{\text{crit}}$, the non-isomorphism between the Wronskian and Bethe algebras, if present, can be only due to $\varphi_{\bar{\theta}}^{\text{nil}}$ having non-zero kernel. Roughly speaking, one can only lose information about Jordan block structure. The Bethe algebra could in principle have eigenspaces of dimension higher than 1 while this never happens with $\hat{\mathcal{W}}_\Lambda$, see Section 3.2.

Proof. First note that $\varphi_{\bar{\theta}}^{\text{nil}}$ and hence $\varphi_{\bar{\theta}}^{\text{diag}}$ are well-defined because the nil-radical is an ideal and the image of a nilpotent element is nilpotent.

Take a regular point $\bar{\theta}$ ($\chi(\bar{\theta}) \notin \mathcal{X}_{\text{crit}}$) and consider a linear combination X of c_ℓ 's taking pairwise distinct values at the d solutions¹⁹ of (2.27) at $\bar{\theta}$ (this is possible since all the solutions are distinct). Then $(X^i)_{0 \leq i \leq d-1}$ is a basis of $\mathcal{W}_\Lambda(\bar{\theta})$ and by continuity it will remain a basis upon specialisation to any θ in some open neighbourhood $\mathcal{O}_{\bar{\theta}}$. Other choices of local bases are possible but for convenience we will work with this one.

Suppose φ_θ is not an isomorphism for all $\theta \in \mathcal{O}_{\bar{\theta}}$. Then $(\hat{X}^i)_{0 \leq i \leq d-1}$ are not linearly dependent for all $\theta \in \mathcal{O}_{\bar{\theta}}$. Construct columns from the d^2 components of the matrices \hat{X}^i and combine the columns into a $d^2 \times d$ matrix. Linear dependence implies that all of the $d \times d$ minors of this matrix vanish on $\mathcal{O}_{\bar{\theta}}$. Since these minors are polynomials in θ_ℓ this means that they are zero as polynomials. This in turn provides a non-trivial relation $\sum_{i=1}^{d-1} p_i(\theta_\ell) \hat{X}^i = 0$ with $p_i \in \mathbb{C}[\theta]$.

Now we would like to be able to take $p_i \in \mathbb{C}[\chi]$. To this end we use the braiding property (4.1) which implies $\sum_{i=1}^{d-1} p_i(\theta_{\sigma(\ell)}) \hat{X}^i = 0$ for any $\sigma \in \mathbf{S}_L$. Thus we can replace the

¹⁹We do not assume $d = d_\Lambda$ to make this proof independent of the counting result of Section 3.4. We also note that this proof does not rely on Proposition 3.1.

p_i by their symmetric part. To ensure that it is non-zero for at least one of them we can multiply the relation we started with by $\prod_{\sigma \in S_L \setminus \text{Id}} p_k(\theta_{\sigma(\ell)})$ for some non-zero p_k and then take the symmetric part.

In the end we obtain a non-zero polynomial P with coefficients in $\mathbb{C}[\chi]$ of degree smaller or equal to $d - 1$ such that $P(\hat{X}) = 0$. But by the isomorphism (4.2) this implies that $P(X) \in \mathcal{I}_\Lambda$. Specialising \mathcal{W}_Λ at a point of $\mathcal{O}_{\bar{\theta}}$ where one of the coefficients of P does not vanish we obtain a contradiction with the fact that $(X^i)_{0 \leq i \leq d-1}$ must be a basis at that point.

Therefore $\varphi_{\bar{\theta}'}$ is an isomorphism for at least one regular point $\bar{\theta}' \in \mathcal{O}_{\bar{\theta}}$. By path-connectivity this immediately propagates to all regular points. Indeed, $\varphi_{\bar{\theta}}$ can cease to be an isomorphism only if the dimension of the Bethe algebra drops but since the spectrum of Q-operators (the set of roots of their characteristic polynomials) is continuous in θ this can only happen when two solutions cross, that is, at singular points.

At singular points, by continuity of the spectrum, solutions of the Wronskian Bethe equations are still in bijection with the spectrum of Q-operators. The only information that can be lost is the multiplicity of the solutions. Then considering $\varphi_{\bar{\theta}}$ up to nilpotent parts restores the isomorphism. \square

Note that the set of $\bar{\theta}$ where $\varphi_{\bar{\theta}}(X^i)$ for $0 \leq i \leq d-1$ do not form a basis is *a priori* not related to the set of $\bar{\theta}$ for which $\chi(\bar{\theta}) \in \mathcal{X}_{\text{crit}}$. Hence we can typically expect that $\chi(\Theta_{\text{not}})$ is of measure zero inside $\mathcal{X}_{\text{crit}}$.

Remark The continuity of the spectrum of Q-operators combined with the above theorem provides an immediate proof that SW is proper as was previously announced. There is no circular argument as we did not use properness in the proof above.

4.3 Specialisation of the isomorphism

Although the set Θ_{not} where the Wronskian and the Bethe algebras are not isomorphic is constrained to be, most likely, in a measure zero subset of critical points $\bar{\theta}$, we still do not have means to locate Θ_{not} . This is unsatisfactory because we cannot guarantee to be outside Θ_{not} for physically interesting cases, for instance when all inhomogeneities coincide. In this section we will provide an explicit constraint on Θ_{not} . Since the required formalism is quite heavy we will only present the logic behind it and the final results. The technical details are postponed to Appendix B.

The main conceptual step is the following. Although the Wronskian and the Bethe algebras were shown to be isomorphic in Theorem 4.2, there is an important qualitative difference between them. Namely, the Bethe algebra is represented by matrices and so it naturally acts on a vector space (the spin chain Hilbert space), whereas the Wronskian algebra is abstractly defined by generators and relations and does not admit such a representation. The only natural space on which \mathcal{W}_Λ could possibly act is itself (*i.e.* by the regular representation). In addition to an isomorphism between algebras, we would like to build an isomorphism between this representation of \mathcal{W}_Λ and the physical representation of \mathcal{B}_Λ .

To build such an isomorphism, a standard procedure is to try to find a cyclic vector. By definition, for a given algebra \mathcal{A} that acts on some vector space V , a vector ω is said to be cyclic if the action of \mathcal{A} on ω spans V . Then V is said to be a cyclic \mathcal{A} -module. Equivalently, ω is cyclic if and only if the map $\psi_\omega : \mathcal{A} \rightarrow V$, $A \mapsto A \cdot \omega$ is surjective. If moreover it is injective, it is an isomorphism identifying V with the regular representation of \mathcal{A} .

In the case of \mathcal{B}_Λ acting on $U_\Lambda \otimes \mathbb{C}[\theta]$, it turns out that ψ_ω is injective for *any* nonzero vector ω as shown in Lemma B.7. Unfortunately, the image of ψ_ω is not $U_\Lambda \otimes \mathbb{C}[\theta]$ except probably for $L = 1$. This can already be seen by the fact that \mathcal{B}_Λ as an algebra involves only symmetric polynomials χ_ℓ whereas the matrix coefficients of \hat{c}_ℓ are from $\mathbb{C}[\theta]$. Nevertheless, for a specific and unique (up to normalisation) choice of ω one can explicitly identify the image of ψ_ω as a subspace $\mathcal{U}_\Lambda^S \subset U_\Lambda \otimes \mathbb{C}[\theta]$ invariant under an action of the symmetric group S_L commuting with the Yangian (Lemma B.8). Thus the regular representation of \mathcal{W}_Λ can be identified with the representation \mathcal{U}_Λ^S of \mathcal{B}_Λ .

The above remarks are the tools to prove a powerful and explicit constraint on Θ_{not} . Since it is a central result to us, we first recall the definitions of all the objects.

$\mathcal{W}_\Lambda(\bar{\chi})$ is the specialised Wronskian algebra at point $\bar{\chi} \equiv (\bar{\chi}_1, \dots, \bar{\chi}_L) \in \mathcal{X} \simeq \mathbb{C}^L$. It is defined as $\mathcal{W}_\Lambda(\bar{\chi}) := \mathbb{C}[c_1, \dots, c_L]/\mathcal{I}$, where $\mathcal{I} := \langle \text{SW}_1 - \bar{\chi}_1, \dots, \text{SW}_L - \bar{\chi}_L \rangle$ is an ideal in $\mathbb{C}[c_1, \dots, c_L]$. For the definition of SW_ℓ see (2.27). Also denote $\hat{\mathcal{I}} := \varphi(\mathcal{I})$, where φ is the map (4.2), and $\mathcal{J} := \langle \chi_1 - \bar{\chi}_1, \dots, \chi_L - \bar{\chi}_L \rangle$ considered as a (maximal) ideal of $\mathbb{C}[\chi]$.

$\mathcal{B}_\Lambda(\bar{\theta})$ is the Bethe subalgebra of the Yangian in the spin chain representation at point $\bar{\theta} = (\bar{\theta}_1, \dots, \bar{\theta}_L)$ restricted to the weight subspace U_Λ . It is generated by operators $\hat{c}_1, \dots, \hat{c}_L$. Matrix entries of these operators are polynomials in inhomogeneities θ_ℓ that are being set to values $\bar{\theta}_\ell$.

$\chi_\ell(\theta) := \sum_{1 \leq i_1 < \dots < i_\ell \leq L} \theta_{i_1} \dots \theta_{i_\ell}$ are the elementary symmetric polynomials of degree ℓ , $\ell = 1, \dots, L$.

Let us first state an immediate consequence of Theorem 4.2 and the discussion above.

Lemma 4.4. *i) φ induces an isomorphism of \mathbb{C} -algebras $\mathcal{W}_\Lambda(\bar{\chi}) \simeq \mathcal{B}_\Lambda/\hat{\mathcal{I}}$,*

ii) ψ_ω induces an isomorphism of representations $\mathcal{W}_\Lambda(\bar{\chi}) \simeq \mathcal{U}_\Lambda^S/\mathcal{J} \cdot \mathcal{U}_\Lambda^S$.

Proof. One easily checks that indeed $\psi_\omega(\hat{\mathcal{I}}) = \hat{\mathcal{I}} \cdot \omega = \mathcal{J} \cdot \mathcal{B}_\Lambda \cdot \omega = \mathcal{J} \cdot \mathcal{U}_\Lambda^S$. □

This does not seem to be a very helpful statement since it is not clear how one should interpret the abstract quotients $\mathcal{B}_\Lambda/\hat{\mathcal{I}}$ and $\mathcal{U}_\Lambda^S/\mathcal{J} \cdot \mathcal{U}_\Lambda^S$. However, Theorem B.9 implies the following

Theorem 4.5. *If $\bar{\chi}_\ell = \chi_\ell(\bar{\theta}_1, \dots, \bar{\theta}_L)$ and $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$ then*

$$\text{ev}_{\bar{\theta}} : \mathcal{U}_\Lambda^S/\mathcal{J} \cdot \mathcal{U}_\Lambda^S \longrightarrow U_\Lambda, \quad [v] \longmapsto v(\bar{\theta}_1, \dots, \bar{\theta}_L) \quad (4.7)$$

is an isomorphism of the Bethe algebra representations. □

Here $\text{ev}_{\bar{\theta}}$ denotes the map²⁰ induced by $\text{Ev}_{\bar{\theta}}$, the evaluation of vectors of $\mathcal{U}_{\Lambda}^{\mathbb{S}} \subset U_{\Lambda} \otimes \mathbb{C}[\theta]$ at $\bar{\theta}$. As $\text{Ev}_{\bar{\theta}}(\mathcal{J} \cdot \mathcal{U}_{\Lambda}^{\mathbb{S}}) = 0$, (4.7) is well-defined. Since the Bethe algebra is represented by $\mathcal{B}_{\Lambda}/\hat{\mathcal{I}} \simeq \mathcal{W}_{\Lambda}(\bar{\chi})$ in $\text{End}(\mathcal{U}_{\Lambda}^{\mathbb{S}}/\mathcal{J} \cdot \mathcal{U}_{\Lambda}^{\mathbb{S}})$ and by $\mathcal{B}_{\Lambda}(\bar{\theta})$ in $\text{End}(U_{\Lambda})$ we thus obtain

Theorem 4.6. *If $\bar{\chi}_{\ell} = \chi_{\ell}(\bar{\theta}_1, \dots, \bar{\theta}_L)$ and $\bar{\theta}_{\ell} + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$ then*

$$\varphi_{\bar{\theta}} : \mathcal{W}_{\Lambda}(\bar{\chi}) \longrightarrow \mathcal{B}_{\Lambda}(\bar{\theta}), \quad c_{\ell} \longmapsto \hat{c}_{\ell} \quad (4.8)$$

is an algebra isomorphism over \mathbb{C} . □

This result improves Theorem 4.3 by giving an explicit condition under which not only the diagonal but also the nilpotent parts of the Wronskian and Bethe algebras are isomorphic. Note that values of $\bar{\chi}_{\ell}$ are not restricted in any way, there is only a restriction on which solution of $\bar{\chi}_{\ell} = \chi_{\ell}(\bar{\theta}_1, \dots, \bar{\theta}_L)$ can be taken.

Assuming the condition $\bar{\theta}_{\ell} + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$ is satisfied, the construction above has several immediate consequences. Composing $\text{ev}_{\bar{\theta}} \circ \psi_{\omega}$ or, equivalently, acting with $\mathcal{W}_{\Lambda}(\bar{\chi})$ on $\omega(\bar{\theta})$ via $\varphi_{\bar{\theta}}$ we obtain

Corollary 4.7. *The spin chain representation of the Bethe algebra restricted to U_{Λ} is isomorphic to the regular representation of $\mathcal{W}_{\Lambda}(\bar{\chi})$.* □

This result is important for separation of variables, as is discussed in Section 7.4.

Corollary 4.8. *$\mathcal{B}_{\Lambda}(\bar{\theta})$ is a maximal commutative subalgebra of $\text{End}(U_{\Lambda})$.*

Proof. $\check{\mathcal{W}}_{\Lambda}(\bar{\chi})$ – the image of generators $\mathcal{W}_{\Lambda}(\bar{\chi})$ under the regular representation – is a maximal commutative subalgebra of $\text{End}_{\mathbb{C}}(\mathcal{W}_{\Lambda}(\bar{\chi}))$ (this is true for any regular representation, see Section 3.2). □

In physical terms, this means that the Bethe algebra contains all commuting charges of the system.

Corollary 4.9. *If $\mathcal{B}_{\Lambda}(\bar{\theta})$ is diagonalisable then its spectrum is simple.*

Proof. A diagonalisable commutative algebra of matrices whose spectrum is not simple is not maximal commutative. □

In particular, the Bethe algebra has simple spectrum every time it is invariant under a Hermitian conjugation which is often the case in physical applications. Simplicity of the spectrum allows us to introduce a new way of classifying solutions by continuously deforming inhomogeneities, see Section 7.2.

Example:

Let us anticipate on the example detailed in Appendix B.5 corresponding to a \mathfrak{gl}_2 non-twisted spin chain of length $L = 3$. The Bethe algebra on the two-dimensional highest-weight subspace V_{\square} is generated (as a $\mathbb{C}[\chi]$ -module) by the identity and the

²⁰Not to confuse with $\text{ev}_{\bar{\theta}}$ in (2.9)

non-trivial operator

$$c = \begin{pmatrix} 2\chi_1 - \sqrt{3}(\theta_1 - \theta_3) & \chi_1 - 3\theta_2 + \sqrt{3}\hbar \\ \chi_1 - 3\theta_2 - \sqrt{3}\hbar & 2\chi_1 + \sqrt{3}(\theta_1 - \theta_3) \end{pmatrix}. \quad (4.9)$$

As long as c is not proportional to the identity, the evaluated Bethe algebra will be of dimension 2 and therefore isomorphic to the Wronskian algebra. We see that $c \propto \mathbb{1}$ if and only if $\hbar = 0$, that is in the Gaudin limit, and $\bar{\theta}_1 = \bar{\theta}_2 = \bar{\theta}_3$.

This example demonstrates that the hypothesis $\theta_{\ell'} \neq \theta_{\ell} + \hbar$ in Theorem 4.6 is not a necessary condition.

4.4 Construction of Bethe vectors

Let us finally comment on how to use the established isomorphism to construct in a bijective way eigenstates of the Bethe algebra from solutions of the Bethe equations. This gives us the practical meaning of the words ‘‘completeness’’ and ‘‘faithfulness’’.

We can always take an element of the Wronskian algebra X such that $(X^k)_{0 \leq k \leq d_{\Lambda}-1}$ forms a basis. In other words the polynomial of smallest degree such that $P(X) = 0$ in $\mathcal{W}_{\Lambda}(\bar{\chi})$ is of degree d_{Λ} ²¹. Its roots x_i , $i = 1, \dots, d_{\Lambda}$ is a way to encode the solutions of WBE (2.26).

By the established isomorphism, P is both the characteristic and minimal polynomial of the matrix $\hat{X} \in \mathcal{B}_{\Lambda}(\bar{\theta})$ which means that any smaller-degree polynomial of \hat{X} yields a non-zero matrix. Let x_i be an eigenvalue of \hat{X} . Then $\frac{\det(\lambda - \hat{X})}{\lambda - x_i}$ is a polynomial in λ . Take a cyclic vector ω ²² and define

$$v_{x_i} = \frac{\det(\lambda - \hat{X})}{\lambda - x_i} \Big|_{\lambda = \hat{X}} \omega. \quad (4.10)$$

Since $\det(\lambda - \hat{X})|_{\lambda = \hat{X}} = 0$, one has $(\hat{X} - x_i)v_{x_i} = 0$ and so v_{x_i} is an eigenvector of \hat{X} with eigenvalue x_i .

We emphasise that as $B_{\Lambda}(\bar{\theta})$ is isomorphic to the regular representation of the Wronskian algebra all eigenspaces are one-dimensional and so all eigenvectors with eigenvalue x_i are collinear with v_{x_i} which guarantees the bijection.

In case of degeneration of solutions, different Jordan blocks of \hat{X} must have distinct eigenvalues. If x_i is a root of multiplicity n then $v_{x_i}^{(m)} \equiv \frac{\det(\lambda - \hat{X})}{(\lambda - x_i)^m} \Big|_{\lambda = \hat{X}} \omega$, $m \leq n$, provide a Jordan basis for \hat{X} . Since \hat{X} generates $B_{\Lambda}(\bar{\theta})$ this basis will also trigonalise the Bethe algebra.

²¹The existence of such an X is obvious at non-degenerate points $\bar{\theta}$ and otherwise follows from the analysis in Appendix A.2. We introduced X for clarity, but the discussed construction of the Bethe algebra eigenstates can be also formulated in a way that does not rely on the existence of X .

²²Corollary 4.7 implies that under the usual assumption on $\bar{\theta}$ such a vector always exists. By continuity, this implies that any generic vector will also be cyclic.

5 Various parameterisations of the Bethe algebra

Although we proved that the number of solutions of the Wronskian Bethe equations is the correct one, we did not develop any intuition about how these solutions are organised. We address this issue in the next two sections by proposing techniques to systematically label solutions.

5.1 Restriction and extension of Q-systems

We introduced a restricted Q-system on page 20 to cover the case of short representations: for special sets \mathbf{A}, \mathbf{J} , $Q_{\mathbf{A}|\mathbf{J}}^{\text{rest}} := Q_{AA|JJ}$. Now, we remark that the Q-functions $Q_{\mathbf{A}|\mathbf{J}}^{\text{rest}} = Q_{AA|JJ}$ satisfy the QQ-relations (2.17) and $Q_{\emptyset|\emptyset}^{\text{rest}} = Q_{\emptyset|\emptyset}$ for *any* sets \mathbf{A}, \mathbf{J} .

If $Q_{\mathbf{A}|\mathbf{J}} = 1$ then one can interpret the restricted Q-system as a Q-system of a smaller $\mathfrak{gl}_{m'|n'}$ algebra, where $m' = m - |\mathbf{A}|$, $n' = n - |\mathbf{J}|$. The condition $Q_{\mathbf{A}|\mathbf{J}} = 1$ is of course non-trivial to demand. By counting degrees of polynomials according to (2.33) we see that this is possible if (m, n) is outside of the Young diagram or if (m, n) is situated on the boundary of the Young diagram such that (m', n') is also on the boundary. We note that the degrees of the restricted Q-functions come out to be given by (2.33) for the $\mathfrak{gl}_{m'|n'}$ Q-system.

Let us now construct an opposite to the restriction procedure. For simplicity consider an “elementary” move. Define an extension from the $\mathfrak{gl}_{m|n-1}$ Q-system to the $\mathfrak{gl}_{m|n}$ Q-system as follows:

$$Q_{\mathbf{A}|Jn}^{\text{ext}} = Q_{\mathbf{A}|J}, \quad Q_{\emptyset|n}^{\text{ext}} = Q_{\emptyset|\emptyset}^{\text{ext}} = 1, \quad (5.1)$$

supplemented with the requirement that the Q-functions that do not contain n are fixed by consistency of the QQ-relations. An example of this extension is depicted in Figure 3 using a Hasse diagram [40].

Lemma 5.1. *The extension (5.1) is always possible and moreover it defines all the Q-functions $Q_{\mathbf{A}|J}^{\text{ext}}$ uniquely up to symmetries.*

Proof. Let us find $Q_{a|\emptyset}^{\text{ext}}, Q_{\emptyset|j}^{\text{ext}}, Q_{a|j}^{\text{ext}}$ by the prescribed construction. Since $Q_{\emptyset|n} = 1$, one computes $Q_{a|\emptyset}^{\text{ext}} = (Q_{a|n}^{\text{ext}})^+ - (Q_{a|n}^{\text{ext}})^-$ using the known value of $Q_{a|n}^{\text{ext}} = Q_{a|\emptyset}$. To find $Q_{\emptyset|j}^{\text{ext}}$ one solves $(Q_{\emptyset|j}^{\text{ext}})^+ - (Q_{\emptyset|j}^{\text{ext}})^- = Q_{\emptyset|jn}^{\text{ext}}$. The polynomial solution is fixed up to an additive constant, but we remind that $Q_{\emptyset|j}^{\text{ext}} \rightarrow Q_{\emptyset|j}^{\text{ext}} + \alpha Q_{\emptyset|n}^{\text{ext}}$ is a symmetry of the twist-less Q-systems. Finally, we use $(Q_{a|j}^{\text{ext}})^- (Q_{\emptyset|n}^{\text{ext}}) - (Q_{a|n}^{\text{ext}})^- Q_{\emptyset|j}^{\text{ext}} = Q_{a|jn}^{\text{ext}}$ which is a consequence of (2.17) to uniquely fix $Q_{a|j}^{\text{ext}}$ from the already identified quantities. One can now check that the above-constructed Q's satisfy $(Q_{a|j}^{\text{ext}})^+ - (Q_{a|j}^{\text{ext}})^- = Q_{a|\emptyset}^{\text{ext}} Q_{\emptyset|j}^{\text{ext}}$ and so they properly generate the whole Q-system. \square

A caveat of the extension procedure is that $Q_{a|\emptyset}^{\text{ext}} = 0$ if $Q_{a|n}^{\text{ext}}$ is a constant. This does not happen however if both points $(m, n - 1)$ and (m, n) belong to the boundary of the Young diagram.

The analogous definitions and statements can be made also for the extension from the $\mathfrak{gl}_{m-1|n}$ to the $\mathfrak{gl}_{m|n}$ Q-system.




Figure 3: Hasse diagram for extension of the \mathfrak{gl}_2 Q-system (blue square) to the $\mathfrak{gl}_{2|1}$ Q-system (blue and red squares). One considers nodes of the blue square as known, supplements this data with $Q_{\bar{\emptyset}|\bar{\emptyset}}^{\text{ext}} = 1$ condition, and finds the rest by QQ-relations. Note that if $Q_2 = 1$ then $Q_{2|\emptyset}^{\text{ext}} = 0$. In this case, $Q_{\bar{\emptyset}}$ contains all physical information and we can restrict both the \mathfrak{gl}_2 and the $\mathfrak{gl}_{2|1}$ systems to the \mathfrak{gl}_1 system (green line) that consists of $Q_{\bar{\emptyset}}^{\text{rest}} = Q_{\bar{\emptyset}} = Q_{\bar{\emptyset}|\bar{\emptyset}}^{\text{ext}}$ and $Q_{\emptyset}^{\text{rest}} = Q_2 = Q_{2|1}^{\text{ext}} = 1$.

5.2 Isomorphism of twist-less \mathcal{B}_Λ across $\mathfrak{gl}_{m|n}$ algebras of various ranks

An important conclusion from the made observations is that if $\mathfrak{gl}_{m'|n'} \subset \mathfrak{gl}_{m|n}$ and both points (m', n') and (m, n) belong to the boundary of the Young diagram then the restriction and extension procedures of the Q-system are inverse of one another and hence both Q-systems contain precisely the same physical information.

By performing a sequence of restrictions and extensions, we conclude that all the $\mathfrak{gl}_{m'|n'}$ Q-systems with (m', n') being on the boundary of Λ^+ (black and red dots in Figure 1) are bijectively related. Moreover, this is done by using only polynomial operations. Hence we conclude the following.

Lemma 5.2. *In the twist-less case, all the Bethe algebras \mathcal{B}_Λ generated by the c_ℓ of $\mathfrak{gl}_{m|n}$ Q-systems, where (m, n) is any point on the boundary of the Young diagram Λ^+ , are isomorphic as $\mathbb{C}[\chi]$ -algebras. \square*

We note that arguments leading to this conclusion do not require isomorphism of the Wronskian and Bethe algebra.

Example:

The \mathfrak{gl}_3 and $\mathfrak{gl}_{1|1}$ Q-systems for representation \boxplus from the previous examples are related by the prescribed procedure:

$$\mathfrak{gl}_{3|0} \xrightarrow{\text{rest}} \mathfrak{gl}_{2|0} \xrightarrow{\text{ext}} \mathfrak{gl}_{2|1} \xrightarrow{\text{rest}} \mathfrak{gl}_{1|1}, \quad (5.2)$$

the extension bit of which is outlined in Figure 3.

Hence $Q_1^{\mathfrak{gl}_3}, Q_2^{\mathfrak{gl}_3}$ (and $Q_3^{\mathfrak{gl}_3} = 1$) should contain the same physical information as $Q_{1|\emptyset}^{\mathfrak{gl}_{1|1}}, Q_{\emptyset|1}^{\mathfrak{gl}_{1|1}}$ (and $Q_{1|1}^{\mathfrak{gl}_{1|1}} = Q_\theta$) and expressible through one another.

By performing the restriction and extension transformations as is described above one finds

$$Q_2^{\mathfrak{gl}_3} \propto \Psi(Q_{\emptyset|1}^{\mathfrak{gl}_{1|1}}), \quad (5.3a)$$

$$Q_1^{\mathfrak{gl}_3} \propto \Psi \left(\frac{W(Q_{1|1}^{\mathfrak{gl}_{1|1}}, \Psi(Q_{1|\emptyset}^{\mathfrak{gl}_{1|1}}))}{Q_{1|\emptyset}^{\mathfrak{gl}_{1|1}}} \right), \quad (5.3b)$$

where $\Psi(A) = B$ if $B^+ - B^- = A$ and the ratio in (5.3b) is a polynomial (Bethe equations ensure that the Euclidean remainder of the numerator and the denominator is the zero polynomial). Ψ is defined up to addition of a constant, there are three such constants in (5.3) which corresponds to three symmetries in the \mathfrak{gl}_3 system restricted to V_{\square} . We fixed the symmetry in (2.36) by setting certain $c_i^{(k)}$ to zero, and we should set the constants of integration to the values that reproduce this choice.

Explicitly in terms of c 's, the transformation (5.3) becomes

$$c_1^{(3)} = 4c_{1|\emptyset}, \quad (5.4a)$$

$$c_1^{(1)} = (2c_{\emptyset|1} - 4c_{1|\emptyset})\hbar^2 - 8c_{1|1}, \quad (5.4b)$$

$$c_2^{(1)} = 2c_{\emptyset|1}. \quad (5.4c)$$

By reversing (5.2), we find that

$$Q_{\emptyset|1}^{\mathfrak{gl}_{1|1}} \propto Q_{23}^{\mathfrak{gl}_3}, \quad (5.5a)$$

$$Q_{1|\emptyset}^{\mathfrak{gl}_{1|1}} \propto W(Q_1^{\mathfrak{gl}_3}, Q_2^{\mathfrak{gl}_3}, Q_3^{\mathfrak{gl}_3}, u), \quad (5.5b)$$

$$Q_{1|1}^{\mathfrak{gl}_{1|1}} \propto Q_{123}^{\mathfrak{gl}_3}. \quad (5.5c)$$

In terms of c 's, this becomes

$$c_{\emptyset|1} = \frac{c_2^{(1)}}{2}, \quad (5.6a)$$

$$c_{1|\emptyset} = \frac{c_1^{(3)}}{4}, \quad (5.6b)$$

$$c_{1|1} = \frac{(c_2^{(1)} - c_1^{(3)})\hbar^2 - c_1^{(1)}}{8}. \quad (5.6c)$$

Transformation (5.6) is the inverse of (5.4).

Remark In general, the isomorphism between Bethe algebras for different (m, n) is a non-linear polynomial map in c_ℓ , and it is remarkable that its inverse is also polynomial.

By Lemma 5.2, we can use the Q-system which corresponds to $\mathfrak{gl}_{m=h_{\Lambda^+}|n=0}$, where

h_{Λ^+} is the height of the Young diagram, and is therefore purely bosonic. Hence we can use in principle results from [9] for bosonic Bethe algebras to formulate and prove the completeness and faithfulness statements for supersymmetric spin chains²³, but we choose to not rely on this relation to [9], highlight instead novel important features of the system, and to produce results in a way that does not use the nested Bethe Ansatz.

5.3 Q-system on Young diagrams

One of the interesting features is that the Bethe algebra can be also generated from the so-called Q-system on a Young diagram which was introduced in [80], extensively used in application to the AdS/CFT spectral problem [81, 82] and more recently in several other studies, see *e.g.* [83, 84].

The Q-system on a Young diagram is a collection of monic polynomials $Q_{a,s}$ defined as

$$Q_{a,s} \propto Q_{a+1,a+2,\dots,m|s+1,s+2,\dots,n}, \quad (5.7)$$

where the Q-function on the r.h.s. is a member of the $\mathfrak{gl}_{m|n}$ Q-system and (m, n) belongs to the boundary of the Young diagram. It is clear from the extension-restriction procedure that $Q_{a,s}$ does not depend on m, n . In particular, $Q_{0,0} = Q_\theta$.

$Q_{a,s}$ are naturally assigned to the nodes of the \mathbb{Z}^2 -lattice which belong to the Young diagram shape, $Q_{a,s} = 1$ on the boundary of the diagram (black/red dots in Figure 1). The QQ-relations between $Q_{a,s}$ are

$$Q_{a+1,s+1}Q_{a,s} \propto W(Q_{a+1,s}, Q_{a,s+1}), \quad (5.8)$$

this follows from (2.17b).

Example:

For the Young diagram \boxplus , $Q_{a,s}$ that are not equal to 1 are $Q_{1,0} = Q_{\emptyset|1}^{\mathfrak{gl}_{1|1}} = Q_2^{\mathfrak{gl}_2} = Q_{23}^{\mathfrak{gl}_3}$, $Q_{0,1} = Q_{1|\emptyset}^{\mathfrak{gl}_{1|1}}$, and $Q_{0,0} = Q_\theta$. There is only one non-trivial relation $Q_{0,1}Q_{1,0} = Q_{0,0}^+ - Q_{0,0}^-$, so there is no significant difference between this simple Q-system and the above-discussed example of a $\mathfrak{gl}_{1|1}$ Q-system. Systems for larger Young diagrams are more interesting of course, there are a plenty of examples in [81, 82] and we give one explicit example on page 43.

The Q-system on a Young diagram Λ^+ is polynomially generated from the Bethe algebra \mathcal{B}_Λ as it follows directly from (5.7). The converse is also true (the polynomiality lemma of [80]):

Lemma 5.3. *Consider the Q-system on a Young Diagram Λ^+ , and choose any m, n that lie on the boundary of Λ^+ . Then, up to symmetry, one can uniquely construct a solution of the $\mathfrak{gl}_{m|n}$ QQ-relations such that (5.7) holds and that the coefficients of the $Q_{A|J}$ are polynomial functions of the coefficients of the $Q_{a,s}$.*

²³It is also possible to establish a bijection between bosonic and supersymmetric Q-systems in the presence of twist. To this end one first extends the original Q-system to a larger one with a partially degenerate twist where an analog of the Young diagram boundary and hence a possibility to move along it emerges.

So the Q-system on a Young diagram is yet another description of the Bethe algebra restricted to V_Λ^+ . We shall benefit from it for counting purposes.

We will now prove Lemma 5.3 in a bit different way compared to [80]. The techniques introduced in this proof are used in Appendix D.

Proof. It suffices to find all $Q_{a|\varnothing}^{\mathfrak{gl}_m}$ for the $\mathfrak{gl}_{m|0}$ Q-system, where $m = h_{\Lambda^+}$ is the height of the Young diagram. Then we can use extensions and restrictions to get all other $\mathfrak{gl}_{m'|n'}$ Q-systems. To this end, let us extend $\mathfrak{gl}_{m|0}$ to the $\mathfrak{gl}_{m|n}$ Q-system, where $n = \lambda_1$ is the width of the Young diagram. Upon extension $Q_{a|\varnothing}^{\mathfrak{gl}_m} = Q_{a|\varnothing}^{\mathfrak{gl}_{m|n}}$. If (m, n) lies outside of the Young diagram boundary, the extended Q-system contains non-physical Q-functions making the extension procedure non-unique but this will not induce an ambiguity in fixing $Q_{a|\varnothing}^{\mathfrak{gl}_m}$.

In the following, the superscript $\mathfrak{gl}_{m|n}$ will be omitted.

One has $Q_{\varnothing|\bar{\varnothing}} = Q_{m,0} = 1$ which is very suitable for applying the so-called bosonisation trick [72]. The bosonisation trick is the observation that the Q-functions $B_{AJ} := Q_{A|\bar{J}}$, where \bar{J} means the complementary set to J , satisfy the QQ-relations of the bosonic \mathfrak{gl}_{n+m} system:

$$B_{\Sigma\alpha\beta}B_\Sigma \propto W(B_{\Sigma\alpha}, B_{\Sigma\beta}), \quad (5.9)$$

where Σ is a multi-index and α, β are indices from the set $\{1, \dots, m, \hat{1}, \dots, \hat{n}\}$. This immediately follows from (2.17) and the definition of B .

Since $B_\varnothing = Q_{\varnothing|\bar{\varnothing}} = 1$, the relations (5.9) are solved by $B_\Sigma = W(B_{\alpha_1}, B_{\alpha_2}, \dots, B_{\alpha_k})$, where $\Sigma = \{\alpha_1, \alpha_2, \dots, \alpha_k\}$ and so (5.7) becomes

$$\mathbb{Q}_{a,s} \propto W(B_{a+1}, \dots, B_m, B_{\hat{1}}, \dots, B_{\hat{s}}). \quad (5.10)$$

We are going to solve these equations for B_α in a unique way modulo admissible symmetry transformations. Note that our main interest is $B_a = Q_{a|\varnothing}^{\mathfrak{gl}_m}$.

From (2.34), $\deg B_a = \deg Q_{a|\varnothing}^{\mathfrak{gl}_m} = \lambda_a + m - a$. Furthermore, remark the following property of the Wronskian determinant: if $W(P_1, \dots, P_k) \propto 1$ for some polynomials P_1, \dots, P_k then, modulo permutations, $\deg P_r = r - 1$ for $r = 1, \dots, k$. And so, by examining (5.10) for (a, s) being on the boundary of Young diagram, where $\mathbb{Q}_{a,s} = 1$, we conclude that all B_α for $\alpha = 1, \dots, n + m$ should have distinct degrees from 0 till $m + n - 1$. The degrees $d_a \equiv \deg B_a$'s satisfy $d_1 \leq d_2 \leq \dots \leq d_m$. Their assignment rule is explained in Figure 4.

We can actually set $B_i = u^{d_i}$ because any subleading orders in polynomials B_i do not affect physically relevant Q-functions (*i.e.* those that survive restrictions that make short representations long, see page 20), in particular $\mathbb{Q}_{a,s}$. Also, using (2.35), we can restrict B_a to the form ²⁴

$$B_a = u^{\lambda_a + m - a} + \sum_{s=1}^{\lambda_a} c_{a-1, s-1} u^{d_s}. \quad (5.11)$$

²⁴If we assign grading to $c_{a,s}$ as is done in Section 3.4 then $\deg c_{a,s}$ is equal to $h_{a+1, s+1}$ – the hook length, *cf.* Figure 2.

Nested Bethe equations are the equations on zeros of $Q_{\leftarrow k}$, $k = 1, \dots, m+n-1$, where we mean *e.g.* $Q_{\leftarrow 3} = Q_{2|\hat{1}\hat{2}}$. Note also that, independently of the path choice, $Q_{\leftarrow(m+n)} = Q_\theta$ – this is the fixed Q-function which plays the role of the source term.

The equations are derived as follows. Let a, b denote some indices from the set $\{1, \dots, m\}$, and i, j – some indices from the set $\{\hat{1}, \dots, \hat{n}\}$. Then (2.17) imply

$$Q_{\leftarrow(k+1)}Q_{\leftarrow(k-1)} \propto W(Q_{\leftarrow k}, Q_{\leftarrow(k-1)b}), \quad \text{for } \leftarrow(k+1) = \leftarrow(k-1)ab, \quad (5.12a)$$

$$Q_{\leftarrow(k-1)a}Q_{\leftarrow(k-1)j} \propto W(Q_{\leftarrow(k+1)}, Q_{\leftarrow(k-1)}), \quad \begin{array}{l} \text{for } \leftarrow(k+1) = \leftarrow(k-1)aj \\ \text{or } \leftarrow(k+1) = \leftarrow(k-1)ja \end{array}, \quad (5.12b)$$

$$Q_{\leftarrow(k+1)}Q_{\leftarrow(k-1)} \propto W(Q_{\leftarrow k}, Q_{\leftarrow(k-1)j}), \quad \text{for } \leftarrow(k+1) = \leftarrow(k-1)ij. \quad (5.12c)$$

Take (5.12a) (or (5.12c)), make a shift $u \rightarrow u + \frac{\hbar}{2}$ and evaluate it at a zero of $Q_{\leftarrow k}$. Do the same for $u \rightarrow u - \frac{\hbar}{2}$ and divide the two evaluated equations by one another. One gets “bosonic” nested Bethe equations

$$\frac{Q_{\leftarrow(k+1)}^+ Q_{\leftarrow(k-1)}^+}{Q_{\leftarrow(k+1)}^- Q_{\leftarrow(k-1)}^-} = -\frac{Q_{\leftarrow k}^{++}}{Q_{\leftarrow k}^{--}} \quad \text{at zeros of } Q_{\leftarrow k}. \quad (5.13a)$$

By evaluating (5.12b) at zeros of $Q_{\leftarrow k}$, one gets $W(Q_{\leftarrow k}, Q_{\leftarrow(k-1)j}) = 0$ which can be written in the “fermionic” nested Bethe equations form

$$\frac{Q_{\leftarrow(k+1)}^+ Q_{\leftarrow(k-1)}^-}{Q_{\leftarrow(k+1)}^- Q_{\leftarrow(k-1)}^+} = 1 \quad \text{at zeros of } Q_{\leftarrow k}. \quad (5.13b)$$

For $\leftarrow k = \leftarrow(k-1)\alpha_k$, we remind that Q-functions are actually (twisted) polynomials and denote them as $Q_{\leftarrow k} \propto \prod_{\beta \in \leftarrow k} z_\beta^{(-1)^\beta u/\hbar} \prod_{l=1}^{M_k} (u - u_l^{(k)})$. Equations (5.13) read

$$\frac{z_{\alpha_k}}{z_{\alpha_{k+1}}} \prod_{(k', l') \neq (k, l)} \frac{u_l^{(k)} - u_{l'}^{(k')} + \frac{\hbar}{2} c_{k, k'}}{u_l^{(k)} - u_{l'}^{(k')} - \frac{\hbar}{2} c_{k, k'}} = 1, \quad (5.14)$$

where k' runs from 0 to $m+n$ (whereas $1 \leq k \leq m+n-1$), and the matrix $c_{k, k'}$ has coefficients $c_{k, k} = (-1)^{\bar{\alpha}_k} + (-1)^{\bar{\alpha}_{k+1}}$, $c_{k, k+1} = c_{k+1, k} = -(-1)^{\bar{\alpha}_{k+1}}$ and all other coefficients equal to zero. In the particular case $\alpha_k = k$, if we rename the label (k) as (α) where $\alpha := m+n-k$ and we remember that $Q_{\leftarrow 0} = 1$ and $Q_{\leftarrow(m+n)} \propto Q_\theta$, then equation (5.13) becomes precisely (1.1)²⁵.

In particular we see that the choice of the Cartan matrix, or equivalently of the Kac-Dynkin diagram, see *e.g.* [86], is implied by the choice of the nesting path [85]. Namely, the Kac-Dynkin diagram should be a chain of $m+n-1$ nodes where the k 'th node is fermionic (crossed) if the k 'th and the $(k+1)$ 'th letters have different grading and is bosonic (blank) otherwise.

For twist-less systems, two comments are due. First, a twist-less Q-system is invariant under symmetry transformations (2.35) and this ambiguity can propagate to the nested

²⁵The matrix $c_{\alpha, \beta}$ of (1.1) is obtained from the matrix $c_{k, k'}$ by restricting to $1 \leq \alpha, \beta < m+n$ and re-ordering of the rows and columns.

double zero, *i.e.* coinciding Bethe roots, we are losing information when passing from (5.12) to (5.13). Indeed, we should consider also a derivative of (5.12) at a double zero which provides an extra constraint in addition to the nested Bethe equations. Double zeros can indeed exist as physical solutions since we can always collide roots by fine-tuning values of inhomogeneities or twist [22]. Moreover, the fine-tuned points coincide sometimes with the physically-relevant case of $\theta_\ell = 0$ [21, 23]²⁶.

Based on the above comments, (5.12) look as more appropriate equations than (5.13). The QQ-relations (5.12) along a nesting path are closely related to the description in terms of quantum eigenvalues²⁷ [6, 31, 37, 92]. For clarity, we introduce them in the example of $\mathfrak{gl}_{2|2}$ spin chain in a concrete grading choice - $2\hat{2}\hat{1}1$. Consider a generating functional [85]

$$\sum_{a=0}^{\infty} (-1)^a \mathcal{D}^a \mathbb{T}_{(1^a)}(u) \mathcal{D}^a = \frac{1}{1 - \mathcal{D}\Lambda_1\mathcal{D}} (1 - \mathcal{D}\Lambda_1\mathcal{D})(1 - \mathcal{D}\Lambda_2\mathcal{D}) \frac{1}{1 - \mathcal{D}\Lambda_2\mathcal{D}} \quad (5.16)$$

that can be viewed as a way to factorise $\text{Ber}[\mathbb{1} - \mathcal{D}T(u)G\mathcal{D}]$ (see (2.13)), and thus Λ_α are sometimes called quantum eigenvalues (of the monodromy matrix). Λ_α commute between themselves and are expressed in terms of the Q -functions as

$$\Lambda_2 = \frac{Q_2^+}{Q_2^-} Q_\theta, \quad \Lambda_{\hat{2}} = \frac{Q_{2|2}^{[-2]} Q_2^+}{Q_{2|2} Q_2^-} Q_\theta, \quad \Lambda_{\hat{1}} = \frac{Q_{2|12}^- Q_{2|2}^{[+2]}}{Q_{2|12}^+ Q_{2|2}^-} Q_\theta, \quad \Lambda_1 = \frac{Q_{2|12}^- Q_{12|12}^{[+2]}}{Q_{2|12}^+ Q_{12|12}^-} Q_\theta. \quad (5.17)$$

The general rule is

$$\Lambda_\alpha = \frac{Q_{\leftarrow k}^{[\pm 2 + m - n - |A| + |J|]} Q_{\leftarrow (k-1)}^{[\mp 2 + m - n - |A'| + |J'|]}}{Q_{\leftarrow k}^{[m - n - |A| + |J|]} Q_{\leftarrow (k-1)}^{[m - n - |A'| + |J'|]}} Q_\theta, \quad (5.18)$$

where α is on the k 'th position of the nesting path, and $\leftarrow k = A|J$, $\leftarrow (k-1) = A'|J'$, the upper sign corresponds to $\bar{\alpha} = 0$ and the lower sign - to $\bar{\alpha} = 1$. While Λ_α depend on the choice of the nesting path, the generating functional (5.16) does not which follows from the QQ-relations (2.17)²⁸.

Given that (5.16) generates transfer matrices, $\Lambda_2 - \Lambda_{\hat{2}}$ should not have poles at zeros of Q_2^- , $\Lambda_{\hat{1}} + \Lambda_{\hat{2}}$ should not have poles at zeros of $Q_{2|2}$ *etc.*, these conditions are another way to generate nested Bethe equations, in the same spirit as they are derived from Baxter equation, see *e.g.* [70].

Both QQ-relations along the path and no-poles conditions for combinations of quantum eigenvalues (which are also path-dependent) are less constraining than the Wronskian condition (2.26). What happens is that polynomiality should be ensured for *all* choices of paths, that is for all Q -functions. This requirement is achieved by (2.26) or equivalent

²⁶the observed cases are however for \mathfrak{gl}_2 chains in a higher spin representation

²⁷The name, to our knowledge, first time appears in the work of Sklyanin [91]. Historically, the name “analytic Bethe Ansatz” (for transfer matrices as sum over Λ 's) was more often in use following the work of Reshetikhin [28].

²⁸In [10], the same concepts and statements are expressed more formally. There, *population* is the same as Q -system reviewed on page 14 and onwards, and *reproduction procedure* is the same as the above-mentioned duality transformations.

formulations. We remark that even if a solution of (5.12) looks normal (*i.e.* it has no coinciding Bethe roots or roots separated by $\hbar, \frac{1}{2}\hbar$) it might be still not physical because of the problems with polynomiality happening when we try to change the path ²⁹.

6 Labelling solutions

In this section we solve Wronskian Bethe equations explicitly in special regimes, when all θ_ℓ are far away from one another. Then one can continuously deform θ_ℓ to any desired values thus obtaining a way to label solutions. A practical application of this labelling approach is demonstrated in Section 7.2. The labelling approach provides an alternative physics-style proof of the completeness.

6.1 Twisted case, labelling with a multinomial expansion

Consider the regime when $|\theta_\ell - \theta_{\ell'}| \sim \Lambda$, and Λ is large. This limit appeared previously in the literature including for counting purposes, see *e.g.* [9, 93].

By rescaling

$$u \rightarrow u/\Lambda, \quad \theta_\ell \rightarrow \theta_\ell/\Lambda, \quad \hbar \rightarrow \hbar/\Lambda, \quad (6.1)$$

which is a symmetry of the Wronskian equations, we can consider the $\hbar \rightarrow 0$ limit with all θ_ℓ being finite and distinct instead of $\Lambda \rightarrow \infty$. We will show below that when $\hbar \rightarrow 0$, we can neglect shifts of the spectral parameter in the polynomial piece of the Baxter function $Q = z^{-u/\hbar}q(u)$, $q(u + \hbar) \simeq q(u) + \mathcal{O}(\hbar)$. So all the QQ-relations (2.17) become of type $QQ = QQ$ and equation (2.26) simplifies to

$$\prod_{a=1}^n q_{a|\emptyset}(u) \prod_{i=1}^m q_{\emptyset|i}(u) = \prod_{\ell=1}^L (u - \theta_\ell). \quad (6.2)$$

The number of solutions to the last equation which is an equation on $c_{a|\emptyset}, c_{\emptyset|i}$ is easily counted to be

$$\frac{L!}{\prod_{a=1}^n \lambda_a! \prod_{i=1}^m \nu_i!} = \dim V_\Lambda. \quad (6.3)$$

The value $\hbar = 0$ is quite special, therefore let us ensure that the conclusion about number of solutions holds also for $\hbar \neq 0$.

Lemma 6.1. *For distinct θ_ℓ , the number of solutions of (2.26) in some neighbourhood of $\hbar = 0$ is given by (6.3) and the solutions are in one-to-one correspondence, by analytic continuation in \hbar , with solutions at $\hbar = 0$.*

Proof. Let us treat \hbar as a parameter and consider SW as a smooth map from $\mathcal{C} \times \mathcal{C}$ to \mathcal{X} . For our discussion, it will be safe to use θ_ℓ instead of χ_ℓ to (locally) parameterise \mathcal{X} . Take a solution \bar{c} of (6.2) corresponding to a point $\bar{\theta}$ with pairwise distinct coordinates. It is easy

²⁹This was observed by C. Marboe and one of the authors [25] while computing the AdS/CFT spectrum for [81, 82]. Curiously, attempts to mitigate this issue led to the formulation of the Q-system on a Young diagram [80].

to check that the differential of SW with respect to the first L coordinates (the c_ℓ variables) at the point $\bar{c}, \hbar = 0$ is an invertible $L \times L$ matrix. Indeed it just reduces to the differential of the smooth map from \mathbb{C}^L to \mathbb{C}^L defined by (6.2) which is clearly non-degenerate if all $\bar{\theta}_\ell$ are distinct. Therefore we can apply the analytic implicit function theorem to conclude that all the solutions at $\hbar = 0$ can be analytically extended to solutions in a neighbourhood of $\hbar = 0$.

It remains to show that all solutions for some $\hbar \neq 0$ can be obtained by extending from $\hbar = 0$. This is equivalent to establishing a version of properness, namely that all sequences of solutions $\bar{c}^{(n)}$ corresponding to $\bar{\theta}$ and $\hbar^{(n)} \rightarrow 0$ as $n \rightarrow \infty$ remain bounded. Assuming the contrary and rewriting (2.26) in terms of the roots u_i of the Q-functions, we can extract a subsequence of solutions $\bar{u}^{(n)}$ such that every root either converges to a finite limit or diverges to infinity. But then one of the equations of (2.26) will contain a monomial with all the diverging roots and only those. This can be seen by the fact that at $\hbar = 0$, (2.26) reduces to (6.2). This monomial will therefore grow faster than any other possible monomial as $n \rightarrow \infty$. Since $\hbar^{(n)} \rightarrow 0$ and $\bar{\theta}$ is finite we arrive at a contradiction. \square

6.2 Twist-less case, labelling with standard Young tableaux

Using the $\hbar \rightarrow 0$ limit is not sufficient in the absence of twist as there are no longer fast-oscillating terms $z_\alpha^{u/\hbar}$ in Q-functions and so the QQ relations won't simplify to the structure $QQ = QQ$ ³⁰. Hence we use a stronger regime when

$$\theta_L \gg \theta_{L-1} \gg \dots \gg \theta_1 \gg \hbar. \quad (6.4)$$

The technical analysis of this limit is given in Appendix D, and here we describe its combinatorial outcome. Similarly to the twisted case, each root of each Q-function $\mathbb{Q}_{a,s}$ “sticks to” a specific inhomogeneity in the sense that, at the leading order of (6.4), it is proportional to this inhomogeneity. However, in contrast to the twisted case, the coefficients of proportionality are not equal to one and an arbitrary distribution of the roots between the inhomogeneities is not allowed. Namely, when θ_L is large compared to the other inhomogeneities, the Q-functions must exhibit the behaviour³¹

$$\mathbb{Q}_{a,s}(u) \sim (u - N_{a,s}^{(L)} \theta_L) \tilde{\mathbb{Q}}_{a,s}(u) \quad (6.5)$$

for certain (a, s) and $\mathbb{Q}_{a,s}(u) \sim \tilde{\mathbb{Q}}_{a,s}(u)$ for the other (a, s) , such that $\tilde{\mathbb{Q}}_{a,s}$ form a Q-system on a Young diagram which is obtained from Λ^+ by removing one box. $N_{a,s}^{(L)}$ are numerical coefficients fixed below. It is clear that (6.5) will hold precisely for those (a, s) for which $\deg \mathbb{Q}_{a,s} = \deg \tilde{\mathbb{Q}}_{a,s} + 1$, and these are the points satisfying $a < \bar{a}, s < \bar{s}$ if the box (\bar{a}, \bar{s}) is being removed.

After removing one box, we end up with a Q-system for a spin chain of length $L - 1$. Now we repeat the argument with $\theta_{L-1} \rightarrow \infty$ and so on and recursively fully disentangle the Young diagram. We associate a number $\mathcal{T}_{\bar{a}, \bar{s}}$ to each box of the Young diagram which

³⁰They will reduce to those of the Gaudin model, see Section 7.3

³¹ \sim designates an equality at the leading order of the corresponding expansion, for this case – the large- θ_L expansion. Equality is verified by comparing coefficients of polynomials in u .




Figure 5: Description of a solution of twist-less Q-system *via* a standard Young tableau: in the regime $\theta_L \gg \theta_{L-1} \gg \dots \gg \theta_1 \gg \hbar$, the scaling of the roots of the Q function at any given node is given by the set of boxes to the bottom right of this node (for instance the red resp. green boxes for $\mathbb{Q}_{0,3}$ resp. $\mathbb{Q}_{2,1}$).

is equal to the length of the spin chain at which the box (\bar{a}, \bar{s}) decouples. These numbers range from 1 to L and increase across each row and each column, *i.e.* they form a standard Young tableau (SYT) \mathcal{T} of shape Λ^+ . An equivalent statement on the level of NBAE was made for \mathfrak{gl}_m in [44], although the $\hbar \rightarrow 0$ regime was suggested (which we know is not sufficient) and details were not given for $n > 2$.

For a given SYT \mathcal{T} , the solution $\mathbb{Q}_{a,s}$ at the leading order of (6.4) is then given by

$$\mathbb{Q}_{a,s} \sim \mathbb{Q}_{a,s}^{\text{lead}} = \prod_{\bar{a} > a, \bar{s} > s} \left(u - N_{a,s}^{(\mathcal{T}_{\bar{a},\bar{s}})} \theta_{\mathcal{T}_{\bar{a},\bar{s}}} \right), \quad (6.6)$$

see Figure 5.

Let us now fix the coefficients $N_{a,s}^{(L)}$. To this end recall that $\mathbb{Q}_{a,s}$ are defined as monic polynomials and restore the normalisation in (5.8)

$$\mathbb{Q}_{a+1,s} \mathbb{Q}_{a,s+1} = \frac{\mathbb{Q}_{a,s}^+ \mathbb{Q}_{a+1,s+1}^- - \mathbb{Q}_{a,s}^- \mathbb{Q}_{a+1,s+1}^+}{\hbar (\deg \mathbb{Q}_{a,s} - \deg \mathbb{Q}_{a+1,s+1})}. \quad (6.7)$$

The key identity we will need is the relation of the polynomial degrees of $\mathbb{Q}_{a,s}$ to the hook length $h_{a,s}$ of the Young diagram box (a, s)

$$\deg \mathbb{Q}_{a,s} - \deg \mathbb{Q}_{a+1,s+1} = h_{a+1,s+1}. \quad (6.8)$$

Note that $h_{a,s}$ will change when we remove certain boxes from the Young diagram. Denote therefore by $h_{a,s}^{(\ell)}$ the hook length in the diagram with ℓ boxes which appears in the recursive procedure. Also, denote by $\mathbb{Q}_{a,s}^{(\ell)}$ the Q-functions on this diagram.

Let us remove the box (\bar{a}, \bar{s}) in the recursive procedure. This means that there are currently $\ell = \mathcal{T}_{\bar{a},\bar{s}}$ boxes in the diagram, and one has $\mathbb{Q}_{a,s}^{(\ell)} \sim (u - N_{a,s}^{(\ell)} \theta_\ell) \mathbb{Q}_{a,s}^{(\ell-1)}$ for all pairs (a, s) with $a < \bar{a}, s < \bar{s}$ and $\mathbb{Q}_{a,s}^{(\ell)} \sim \mathbb{Q}_{a,s}^{(\ell-1)}$ otherwise. Consider the regime $u \ll \theta_\ell$ for which $(u + \kappa \hbar - N_{a,s}^{(\ell)} \theta_\ell) \simeq -N_{a,s}^{(\ell)} \theta_\ell$ for any finite κ and so (6.7) simplifies providing consistency relations between $N_{a,s}^{(\ell)}$:

$$\begin{aligned} N_{a+1,s}^{(\ell)} N_{a,s+1}^{(\ell)} &= N_{a,s}^{(\ell)} N_{a+1,s+1}^{(\ell)}, & a < \bar{a} - 1, & s < \bar{s} - 1, \\ N_{a,s+1}^{(\ell)} &= N_{a,s}^{(\ell)} \frac{h_{a+1,s+1}^{(\ell)} - 1}{h_{a+1,s+1}^{(\ell)}}, & a = \bar{a} - 1, & s < \bar{s} - 1, \\ N_{a+1,s}^{(\ell)} &= N_{a,s}^{(\ell)} \frac{h_{a+1,s+1}^{(\ell)} - 1}{h_{a+1,s+1}^{(\ell)}}, & a < \bar{a} - 1, & s = \bar{s} - 1. \end{aligned} \quad (6.9)$$

There is no constraint at the point $a = \bar{a} - 1, s = \bar{s} - 1$ but instead we have to set $N_{0,0}^{(\ell)} = 1$ since $\mathbb{Q}_{0,0}^{(L)} = Q_\theta$ and so $\mathbb{Q}_{0,0}^{(\ell)} = (u - \theta_\ell) \mathbb{Q}_{0,0}^{(\ell-1)}$. This normalisation allows finding all $N_{a,s}^{(\ell)}$ with no ambiguities:

$$N_{a,s}^{(\ell)} = \prod_{a'=1}^a \frac{h_{a',\bar{s}}^{(\ell)} - 1}{h_{a',\bar{s}}^{(\ell)}} \prod_{s'=1}^s \frac{h_{\bar{a},s'}^{(\ell)} - 1}{h_{\bar{a},s'}^{(\ell)}}, \quad a < \bar{a}, s < \bar{s}. \quad (6.10)$$

We emphasise that this solution depends on the choice of \mathcal{T} through the condition $\mathcal{T}_{\bar{a},\bar{s}} = \ell$. Therefore, two distinct $\mathbb{Q}_{a,s}^{\text{lead}}$ differ by the scaling of at least one Bethe root.

We hence confirmed that $\mathbb{Q}_{a,s}^{\text{lead}}$ is explicitly and bijectively fixed by standard Young tableaux. We remind that the number of SYT is the dimension over \mathbb{C} of V_Λ^+ on which the Bethe algebra is restricted.

Since inhomogeneities are not bounded in the regime (6.4) one still needs to perform work to show that solutions in this regime are bijectively linked to solutions at finite values of inhomogeneities. This is done in Appendix D.3. In summary, we have the following result

Lemma 6.2. *For $\Lambda_\ell := \frac{\bar{\theta}_{\ell+1}}{\theta_\ell}$, $\ell = 1, \dots, L-1$, being large enough but finite, solutions of the Q -system on a Young diagram Λ^+ and hence of the Wronskian Bethe equations (2.26) at point $\bar{\theta}$ are bijectively labelled with standard Young tableaux, where the solution associated with a tableau \mathcal{T} approaches*

$$\mathbb{Q}_{a,s} \sim \prod_{\bar{a} > a, \bar{s} > s} \left(u - \bar{\theta}_{\mathcal{T}_{\bar{a},\bar{s}}} \prod_{a'=1}^a \frac{h_{a',\bar{s}}^{(\mathcal{T}_{\bar{a},\bar{s}})} - 1}{h_{a',\bar{s}}^{(\mathcal{T}_{\bar{a},\bar{s}})}} \prod_{s'=1}^s \frac{h_{\bar{a},s'}^{(\mathcal{T}_{\bar{a},\bar{s}})} - 1}{h_{\bar{a},s'}^{(\mathcal{T}_{\bar{a},\bar{s}})}} \right) \quad (6.11)$$

when Λ_ℓ approaches infinity. □

7 Summary and applications

7.1 Completeness, faithfulness, and maximality of the Bethe algebra

In this paper we proved completeness of the Wronskian Bethe equations (WBE) and faithfulness of the map from the Wronskian to the Bethe algebra, for the case of both twisted and twist-less supersymmetric spin chains.

Completeness on the level of equations is the statement that the algebraic number of solutions of the WBE is the "right one", *i.e.* it is equal to the dimension of the weight space U_Λ (as a vector space over \mathbb{C}). We proved the statement for arbitrary numerical values of χ_ℓ – elementary symmetric polynomials in inhomogeneities θ_ℓ . The paper actually contains two independent proofs. The first one is based on a character computation presented in Section 3.4 which is valid because the Wronskian algebra \mathcal{W}_Λ is a free $\mathbb{C}[\chi]$ -module, by Lemma 3.1. The second proof is based on the explicit solution counting in the limits $\left| \frac{\theta_{\ell+1} - \theta_\ell}{h} \right| \gg 1$ (twisted case) and $\frac{\theta_{\ell+1}}{\theta_\ell} \gg 1$ (twist-less case). The fact that this counting is valid for finite (but probably large) values of inhomogeneities is summarised in Lemma 6.1 and Lemma 6.2; the fact that the algebraic number of solutions remains the same for any

values of inhomogeneities is a consequence of freeness but we also show this using more elementary arguments in Lemma A.2.

Faithfulness and hence bijectivity of the map established in Theorem 4.2 allows one to transfer algebraic properties of the Wronskian algebra \mathcal{W}_Λ to the Bethe algebra \mathcal{B}_Λ . The Bethe algebra over $\mathbb{C}[\chi]$ and restricted to the weight subspace U_Λ can then be viewed as a polynomial ring defined by WBE. Furthermore, for the twist-less case, \mathcal{B}_Λ depends on the Young diagram alone and does not depend on the rank of $\mathfrak{gl}_{m|n}$. Its description in terms of a Q-system on a Young diagram directly follows from the results of [41, 80] although this fact was not explained there and we filled in the gap in Section 5. Using the bosonisation trick, we also found a novel very explicit way (5.10) to parameterise functions $Q_{a,s}$ using Wronskian determinants which is the main technical tool for analysing the $\frac{\theta_{\ell+1}}{\theta_\ell} \gg 1$ regime.

The faithfulness property holds also for specialisation of χ_ℓ to any numerical value $\bar{\chi}_\ell = \chi_\ell(\bar{\theta}_1, \dots, \bar{\theta}_L)$, whereas inhomogeneities should probably satisfy the constraint $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$ ³². For almost any values of $\bar{\chi}_\ell$, this follows already from Theorem 4.3 which uses very general properties of the WBE. However, to get really arbitrary values of $\bar{\chi}_\ell$, a more refined analysis is performed in Appendix B which relies on properties of the Yangian and its representations. This analysis builds on ideas of [9] generalising them to the supersymmetric case, with notable exception of Lemma B.7.

Completeness and faithfulness combined ensure that \mathcal{B}_Λ is a maximal commutative subalgebra of $\text{End}(U_\Lambda)$ which should be viewed as the completeness property on the Bethe algebra level. Each *distinct* solution of WBE bijectively corresponds to a joint eigenstate of commuting charges. The word “distinct” means that even in the case when solutions degenerate, the eigenspace corresponding to the coinciding solutions is still one-dimensional. This does not contradict maximality of the Bethe algebra as the latter becomes non-diagonalisable in the degenerate case with the size of the corresponding trigonal block equal to the degree of degeneration.

The results of this paper are likely to be generalisable for spin chains in an arbitrary highest-weight representation of $\mathfrak{gl}_{m|n}$ ³³. Given our preliminary studies, an analog of the quantisation condition (2.26) will not be sufficient but we can build on a generalisation of Lemma 5.3 for Q-systems on Young diagrams. Q-systems on Young diagrams can be also defined for non-compact spin chains [81] and they suggest an explicit isomorphism map between restricted Bethe algebras \mathcal{B}_Λ for non-compact spin chains and compact spin chains. The isomorphism class of \mathcal{B}_Λ should depend only on the extended Young diagram introduced in [81, 96]. Performing the suggested program should prove completeness of the quantum spectral curve for $\mathcal{N}=4$ SYM [73, 97] which is confirmed so far by an extensive

³²If solutions of WBE are non-degenerate this constraint is not needed. For $\chi \in \mathcal{X}_{\text{crit}}$, it might be needed but we did not analyse precisely when, so we keep it as a sufficient requirement. Analysing when it is necessary would probably require exploration of Yangian representation theory beyond techniques developed in the paper.

³³A unified approach for twisted and non-twisted Bethe subalgebras of $Y(\mathfrak{gl}_m)$ has been put forward in [94]. Using these ideas and representation-theoretic arguments, powerful completeness-type results have been proven for the \mathfrak{gl}_2 case in [95].

analysis of its solutions in [81, 82].

7.2 Simplicity of the spectrum and controlled numerical solution

Choose normalisation $\hbar = i$ and consider the twist-less case and real values of inhomogeneities θ_ℓ . The Bethe algebra is invariant under Hermitian conjugation in this case and hence is diagonalisable. On the other hand, diagonalisation is impossible if there are coinciding solutions which immediately implies that restriction of the Bethe algebra to a weight subspace V_Λ^+ has *simple spectrum*, cf. Corollary 4.9.

The considered scenario contains both the homogeneous spin chain with $\theta_\ell = 0$ and the spin chain decoupling limit $|\theta_{\ell+1}/\theta_\ell| \gg 1$ where WBE can be solved explicitly and labelled with standard Young tableaux. We can continuously connect the homogeneous spin chain and the spin chain in the decoupling limit while keeping θ_ℓ real and in this way *unambiguously* label solutions of homogeneous Bethe equations by SYT ³⁴.


We have realised this idea numerically. Parameterise inhomogeneities as $\bar{\theta}_\ell = \Lambda^{\ell-1} - 1$, $\ell = 1, \dots, L$. For a chosen tableau \mathcal{T} , start with the solution (6.11) in the decoupling regime $\Lambda \gg 1$ and then incrementally decrease Λ until it reaches the point $\Lambda = 1$. While changing Λ we require that (6.7) are always satisfied. The numerical realisation turned out to be very stable for any choice of \mathcal{T} that we tried. For $L \lesssim 20$, we are able to produce, for a given \mathcal{T} , a numerical solution with an 80-digit precision in less than three minutes, the speed is obviously much faster for shorter chains. Further substantial optimisation of the code should be possible.

The details of implementation and the code will be published in a separate work. Here we give one illustration. For the example of Q-system on Young diagram on page 43, choose a standard Young tableau


1	2	3	8	9	10	14	18
4	6	7	13	16	17		
5	11						
12	15						

Then we get the following numerical solution

³⁴If we allow θ_ℓ to be complex then we can connect any two solutions of WBE by varying θ . This follows from the path connectivity argument of Section 3.1 and allows in turn to connect any two SYT proving a conjecture made in [44].



Roots of $\mathbb{Q}_{1,0}$ (blue), $\mathbb{Q}_{2,0}$ (green), $\mathbb{Q}_{3,0}$ (red) are shown



Roots of $\mathbb{Q}_{0,1}$ (blue), $\mathbb{Q}_{1,1}$ (green), $\mathbb{Q}_{1,2}$ (red) are shown

To our knowledge, the proposed approach is the first example when we have a systematic and proven to be unambiguous way to control all solutions of the Bethe equations for systems of size where direct brute-forcing (*e.g.* by numerical diagonalisation of Hamiltonian matrix) is unlikely to be practical. For instance, there are 2 148 120 different standard Young tableaux of the same shape (and hence distinct solutions of the Bethe equations with the same magnon numbers) as in the above example. Overall, the length $L = 18$ \mathfrak{gl}_4 chain has a Hilbert space of dimension $\sim 6.8 \times 10^{10}$ comprising 81 662 152 irreps of \mathfrak{gl}_4 with one solution of WBE per irrep.

To compare with other approaches, all solutions of the \mathfrak{gl}_2 chain for $L = 14$ were reported in [98]. Yet, at this length the Hilbert space is of dimension 16384 and the Hamiltonian being a sparse matrix [99] can be diagonalised numerically. In [100] solutions with large magnon numbers were studied quite systematically but only particular classes of solutions were controlled. For spin chains of rather large length, the low-energy excitations around the antiferromagnetic vacuum are also of numerical interest, see *e.g.* [101]. It would be interesting to explore whether we can apply the proposed techniques in this regime and improve the existing tools which rely on the string hypothesis.

7.3 Gaudin model

The \mathfrak{gl}_m Gaudin model [29, 102] can be obtained as the $\hbar \rightarrow 0$ limit of the non-twisted \mathfrak{gl}_m spin chain. Formally, this just amounts to replacing the discrete Wronskian by an actual Wronskian. On the representation theory side the spin chain will no longer be a representation of the Yangian $Y(\mathfrak{gl}_m)$ but of the current algebra $\mathfrak{gl}_m[u]$ (*e.g.* in the terminology of [103]). Completeness of the Bethe Ansatz for the \mathfrak{gl}_m Gaudin model has been proven in [103] under the assumption that all inhomogeneities are pairwise distinct. The philosophy of the proof and the end result are similar to [9]. As far as we know, for the supersymmetric $\mathfrak{gl}_{m|n}$ Gaudin model, completeness is proven for generic values of θ_ℓ [104] in the twist-less case, and for any pair-wise distinct θ_ℓ for the twisted case [105].

In our construction, the Bethe equations and Q-operators admit a well-defined $\hbar \rightarrow 0$ limit and an analogue of Theorem 4.2 can be proven along the same lines. Similarly, all the results relying solely on the analytic properties of the map SW will also be true in the Gaudin case. In particular, the algebraic number of solutions will not depend on $\bar{\chi}$ and

will be equal to d_Λ . Specialisation of the isomorphism will also hold generically using the same arguments as in Theorem 4.3.

To prove further constraints on specialisation as in Theorem 4.6, one way would be to investigate the representation theory of $\mathfrak{gl}_{m|n}[u]$. But instead of doing that, we can reformulate statements for the Bethe algebra \mathcal{B}_Λ of the $\mathfrak{gl}_{m|n}$ system as statements for the non-supersymmetric $\mathfrak{gl}_{h_{\Lambda^+}}$ system. This is based on the results of Section 5 for isomorphisms of Bethe algebras and the discussion on page 67 for the existence of a cyclic vector. They apply in the $\hbar \rightarrow 0$ limit as well. Using [103], one then confirms that the specialisation of the isomorphism also holds in the case of the supersymmetric Gaudin model for pairwise distinct inhomogeneities which is a naive $\hbar \rightarrow 0$ limit of Theorem 4.6 and related statements.

7.4 Separation of variables

To construct a basis that factorises wavefunctions of eigenstates of the Bethe algebra, Maillet and Niccoli proposed [106] to repeatedly act with transfer matrices on a reference state. One can reach factorisation also by choosing other Bethe algebra elements that depend on the spectral parameter u . This idea was fruitfully used recently alongside with other related tools in application to rational spin chains [36, 48, 93, 107–110]. Currently, an SoV basis was constructed for \mathfrak{gl}_m spin chains in arbitrary finite-dimensional representation as [109]

$$\langle \mathbf{x} | = \langle 0 | \prod_{\ell=1}^L \prod_{k=1}^{m-1} \det_{1 \leq i, j \leq k} Q_i(x_{kj}^\ell) \quad (7.1)$$

and for $\mathfrak{gl}_{m|n}$ spin chains in the defining representation as [48]

$$\langle \mathbf{x} | = \langle 0 | \prod_{\ell=1}^L \mathbb{T}_{(1)}(\theta_\ell)^{d_\ell}. \quad (7.2)$$

Here $x_{kj}^\ell = \theta_\ell + \hbar m_{kj}^\ell$, where m_{kj}^ℓ are integers forming Gelfand-Tsetlin patterns and defining what is $\langle \mathbf{x} |$, and d_ℓ are integers from $0 \leq d_\ell \leq m + n - 1$ also defining what is $\langle \mathbf{x} |$. There are exactly as many choices for m_{kj}^ℓ and d_ℓ as the dimension of the corresponding Hilbert space.

An important technical challenge of this approach is to prove that a construction as above indeed produces a basis of the Hilbert space. It was resolved in the mentioned works, however only spin chains with generic twist were considered and certain restrictions on admissible values of inhomogeneities were used.

We can now give an alternative insight on resolving this challenge. By Theorem 4.5, the representation of the Bethe algebra is isomorphic to the regular representation of the Wronskian algebra. In particular there is always a cyclic vector. One can choose the cyclic vector as a reference state $\langle 0 |$. Then $\langle \mathbf{x} | = \langle 0 | \varphi(\mathbf{b}_x)$ form a basis as long as \mathbf{b}_x form a basis in the Wronskian algebra. The Wronskian algebra is a polynomial algebra with Plücker-type relations. Then the basis question reduces to questions very similar to those of projective geometry. This naturally links to the last topic we would like to review.

7.5 Geometric representation theory and Bethe/Gauge correspondence

So far we have concentrated on the isomorphism between the restricted Bethe algebra \mathcal{B}_Λ and the polynomial ring $\mathcal{W}_\Lambda = \mathbb{C}[\chi][c]/\mathcal{I}_\Lambda$. It turns out that for \mathfrak{gl}_m spin chains the Bethe algebra can be realised in a third, purely geometric way [111–114]. Consider the manifold \mathcal{F}_Λ of all the partial flags associated to the weight $\Lambda = [\lambda_1, \dots, \lambda_m]$, that is chains of vector spaces

$$\{0\} = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_{m-1} \subsetneq V_m = \mathbb{C}^L \quad (7.3)$$

such that $\dim V_a/V_{a-1} = \lambda_a$ for all $1 \leq a \leq m$. \mathcal{F}_Λ admits a natural action of $\mathrm{GL}(L)$. Now consider its cotangent bundle $T^*\mathcal{F}_\Lambda$ with an action of $\mathrm{GL}(L) \times \mathbb{C}^*$, where \mathbb{C}^* acts on the cotangent spaces by multiplication. It is known [114] that the equivariant cohomology ring of $T^*\mathcal{F}_\Lambda$ with this action of $\mathrm{GL}(L) \times \mathbb{C}^*$ is given by

$$H_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(T^*\mathcal{F}_\Lambda, \mathbb{C}) = \mathbb{C}[c, \chi, \hbar] / \left\langle \prod_{a=1}^m q_a(u) - Q_\theta(u) \right\rangle. \quad (7.4)$$

In this identification the $L + 1$ parameters $(\chi_\ell)_{1 \leq \ell \leq L}$ and \hbar come from the standard $H_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(\mathrm{pt}, \mathbb{C}) = \mathbb{C}[\chi, \hbar]$ -module structure of $H_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(T^*\mathcal{F}_\Lambda, \mathbb{C})$. Treating \hbar as an additional parameter now, $H_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(T^*\mathcal{F}_\Lambda, \mathbb{C})$ is thus $\mathbb{C}[\chi, \hbar]$ -isomorphic to \mathcal{B}_Λ in the singular twist limit $x_1 \ll x_2 \ll \dots \ll x_m$. In general, when x_a are arbitrary pairwise distinct complex numbers, the twisted Bethe algebra \mathcal{B}_Λ can be identified with a quantum deformation of $H_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(T^*\mathcal{F}_\Lambda, \mathbb{C})$, the so-called equivariant quantum cohomology ring $QH_{\mathrm{GL}(L) \times \mathbb{C}^*}^\bullet(T^*\mathcal{F}_\Lambda, \mathbb{C})$.

This connection is actually a particular case of a more general construction [111] which first appeared in the context of Bethe/Gauge correspondence [112, 113]. Starting from any so-called Nakajima quiver variety one can build a Yangian action³⁵ on its equivariant quantum cohomology ring considered as a Hilbert space. Moreover, the action of the Yangian generators can be expressed as some geometric operations on the classes of the variety. In particular the Baxter Q-operator can be constructed in a purely geometric way [115].

It is still unclear how to properly extend this construction to supersymmetric Yangians [116]. We hope that results of our paper, in particular the isomorphism between the bosonic and supersymmetric case elucidated in Section 5, will be useful for the advancement of the subject.

Acknowledgments

We are grateful to Luca Cassia, Vladimir Dotsenko, Antoine Ducros, Arnaud Eteve, David Hernandez, Atsuo Kuniba, Jules Lamers, Fedor Levkovich-Maslyuk, Maksim Maydanskiy, Lucy Moser, Sergey Mozgovoy, Rafael Nepomechie, Antoine Picard, Paul Ryan, Peter Schauenburg, Didina Serban, Samson Shatashvili, Pedro Tamaroff, Ronan Terpereau, and Emmanuel Wagner for useful discussions.

³⁵The Yangian in question is not necessarily $Y(\mathfrak{gl}_m)$. An identification of these geometrically-realised Yangians with known integrable systems and solutions of the Yang-Baxter equation is an open question.

The work of D.V. and S.L. was partially supported by the Knut and Alice Wallenberg Foundation under grant “Exact Results in Gauge and String Theories” Dnr KAW 2015.0083. The work of S.L. was partially supported by the European Union (through the PO FEDER-FSE Bourgogne 2014/2020 program) and the EIPHI Graduate School (contract ANR-17-EURE-0002) as part of the ISA 2019 project, and by the région Bourgogne-Franche-Comté as part of the MolQuan project.

D.V. is very grateful to Institut de Mathématiques de Bourgogne and Département de Physique de l’Ecole Normale supérieure for hospitality where a part of this work was done. S.L. is very grateful to Nordiska institutet för teoretisk fysik, where a part of this work was done.

A Wronskian algebra - a pedestrian approach

A.1 Some facts from commutative algebra and algebraic geometry

Basic definitions

The study of the polynomial equations can be done in an analytic (geometric) or in an algebraic way. In the analytic approach, to a set of m polynomial equations in n variables $P_i(x_1, \dots, x_n) = 0$, $i = 1, \dots, m$ is assigned an algebraic variety \mathcal{A} - a set of points $x \equiv (x_1, \dots, x_n) \in \mathbb{C}^n$ where these equations hold. The algebraic approach attaches to the equations an ideal generated by P_ℓ , $\mathcal{I} = \langle P_\ell \rangle$ which is the set of all possible polynomials in n variables $Q \in \mathbb{C}[x_1, \dots, x_n]$ that can be written in the form $Q = \sum_\ell q_\ell P_\ell$ for some polynomials q_ℓ .

The relation between the two approaches is established by Hilbert’s Nullstellensatz: if Q vanishes on \mathcal{A} then $Q^r \in \mathcal{I}$ for some integer r . The ideal constructed by all polynomials that vanish on \mathcal{A} is called the radical of \mathcal{I} and is denoted by $\sqrt{\mathcal{I}}$.

The algebraic description is more abstract and is less used in physics but it allows one to more accurately formulate some of the properties of the Wronskian algebra. In particular, we can work over fields different to \mathbb{C} , *e.g.* the field of fractions $\mathbb{C}(\theta)$.

The next concept is to consider functions on \mathcal{A} formalised as the quotient ring

$$\mathcal{R} = \mathbb{C}[x_1, \dots, x_n]/\mathcal{I}. \tag{A.1}$$

If $\mathcal{I} = \sqrt{\mathcal{I}}$ then \mathcal{R} is called the coordinate ring of \mathcal{A} . Note that in this paper not all ideals are equal to their radicals.

Finally, recall that a ring is said to be an integral domain if $ab = 0$ implies $a = 0$ or $b = 0$. The corresponding ideal is then called prime ($ab \in \mathcal{I}$ implies $a \in \mathcal{I}$ or $b \in \mathcal{I}$).

Polynomial division

Easiness of the study of polynomials in *one variable* exists mainly due to the unambiguous polynomial division procedure. Recall how it works: let P be a polynomial in x of degree b which is one (of those polynomials) that generates the ideal \mathcal{I} in $\mathbb{C}[x]$. Let Q be any polynomial in x . If Q contains a monomial cx^a with $a \geq b$, we represent Q as a combination $Q = cx^{a-b}P + (Q - cx^{a-b}P)$ in which the first term is divisible by P and the second term

has no monomial of degree a . One performs the same procedure with $P' = Q - c x^{a-b} P$ and continue it recursively until no monomials divisible by x^b remain. So one obtains a representation $Q = qP + r$, where the degree of r is strictly smaller than b . Both q and r are fixed uniquely. Furthermore, one can guarantee the Bézout's lemma, that is, one can find such $\alpha, \beta \in \mathbb{C}[x]$ that $\alpha P_1 + \beta P_2 = \text{GCD}(P_1, P_2)$ for any polynomials P_1, P_2 , and hence conclude that any ideal in one variable is principal, *i.e.* it is generated by a single polynomial – the GCD of polynomials P_1, \dots, P_m that generate the ideal.

A practical application in our case would be: if a Bethe algebra is generated by a single operator \hat{x} then this algebra is guaranteed to be isomorphic to a quotient $\mathbb{C}[x]/\mathcal{I}$ where \mathcal{I} is the ideal generated by the minimal polynomial of \hat{x} . As we have L generating operators \hat{c}_ℓ , things are not that simple.

Gröbner bases

Many problems in systems with multiple variables arise from difficulties with the polynomial division. First, to even define a division algorithm one needs to introduce a total order on the set of monomials $x^d \equiv \prod_{i=1}^n x_i^{d_i}$ that should be an order in which 1 is the smallest monomial and $a < b$ implies $ac < bc$ for any a, b, c . A diversity of monomial orders is available in contrast to only one option for the single-variable case. We shall use below only lexicographic orders which form a small subset of all possibilities.

After fixing a monomial order and denoting by P_i the generators of the ideal \mathcal{I} , one can perform long polynomial division (exclusion of all monomials that are divisible by leading monomials of P_i) to represent any polynomial Q as

$$Q = \sum_i q_i P_i + r. \quad (\text{A.2})$$

Unfortunately, neither the procedure nor its result are unique if P_i are arbitrary generators, so the division is essentially meaningless.

However, if P_i form a special set called Gröbner basis then r is uniquely defined by the polynomial division³⁶. Hence $Q \in \mathcal{I}$ iff $r = 0$. q_i are not unique though, but uniqueness of r suffices for the study of the quotient ring (A.1).

A set of polynomials P_i forms a Gröbner basis of an ideal \mathcal{I} if *i*) they generate \mathcal{I} , *ii*) the set is closed under computation of S-polynomials, see *e.g.* [117] for further explanations. If moreover, for all $i \neq i'$, P_i does not contain monomials divisible by the leading monomial of $P_{i'}$ then such a Gröbner basis is called the reduced one and it is unique for the given choice of a monomial order. By a Gröbner basis we mean the reduced basis in the following.

Monomial basis

Let us fix a Gröbner basis. The set of monomials that can arise in the remainders of polynomial divisions forms a basis in the quotient ring \mathcal{R} considered as a vector space. This basis shall be called the monomial basis.

³⁶Note however that, in contrast to the one-dimensional case, monomials comprising r can be still bigger than the leading monomials of P_i and yet not divisible by the latter. Hence we might be unable to perform a chain of divisions that leads to the Bézout's lemma and it generically does not hold in the multivariable case.

We can use the monomial basis to realise the regular representation of an algebra in terms of explicit matrices, see the example on page 23. Such a basis has an important advantage – all computations in it are performed in the original field, and so the coefficients of \check{x} will belong to the same field, where \check{x} is a matrix in the regular representation, see Section 3.2.

A.2 $\mathbb{C}(\chi)$ -module and invariance of solutions multiplicity

Consider the ring of polynomials in $2L$ variables $\mathbb{C}[\chi][c] \equiv \mathbb{C}[\chi_1, \dots, \chi_L][c_1, \dots, c_L]$. We define the Wronskian algebra as $\mathcal{W}_\Lambda := \mathbb{C}[\chi][c]/\mathcal{I}_\Lambda$, where $\mathcal{I}_\Lambda = \langle \text{SW}_\ell(c) - \chi_\ell \rangle$ is the ideal generated by Wronskian relations. As we can simply exclude χ_ℓ using equations $\chi_\ell = \text{SW}_\ell(c)$, \mathcal{W}_Λ is isomorphic over \mathbb{C} to $\mathbb{C}[c]$ – the ring of polynomials in L variables. Hence, in particular, \mathcal{W}_Λ is an integral domain and \mathcal{I}_Λ is a prime ideal.

In the case of prime ideals, it is quite easy to promote rings to fields. In this subsection, we shall consider $\chi_\ell = \text{SW}_\ell(c)$ as an equation on c_ℓ in the field of fractions $\mathbb{C}(\chi)$ and \mathcal{W}_Λ as a ring over $\mathbb{C}(\chi)$. “Easiness” of promotion lies in the following statement: any polynomial in variables c_ℓ and χ_ℓ that belongs to \mathcal{W}_Λ considered as an object in a ring over $\mathbb{C}(\chi)$ would also belong to \mathcal{W}_Λ considered as an object in a ring over $\mathbb{C}[\chi]$.

When we work over a field of fractions, we can compute a Gröbner basis. Simply, instead of conventional computation in $\mathbb{C}[c_1, \dots, c_L]/\langle \text{SW}_\ell(c) - \bar{\chi}_\ell \rangle$ with numerical $\bar{\chi}_\ell \in \mathbb{C}$, we do a computation in $\mathbb{C}(\chi)[c_1, \dots, c_L]/\langle \text{SW}_\ell(c) - \chi_\ell \rangle$ with symbolic $\chi_\ell \in \mathbb{C}(\chi)$. When the Gröbner basis is computed, we can construct the corresponding monomial basis and conclude what is the dimension of \mathcal{W}_Λ (as a vector space over $\mathbb{C}(\chi)$) and hence what is the number of solutions of the Wronskian equations. Note that the solutions themselves would typically only exist in an algebraic closure of $\mathbb{C}(\chi)$. However, computation of the monomial basis can be performed directly in $\mathbb{C}(\chi)$ and this is the only thing needed.

Working over $\mathbb{C}(\chi)$ is equivalent to considering χ_ℓ in generic position, when no accidental relations happen. When we specialise to a concrete numerical value $\bar{\chi}_\ell$ of χ_ℓ , we are interested whether the number of solutions changes. We can formulate (a bit stronger) question from the point of view of the Gröbner basis: does it remain a Gröbner basis upon specialisation?

Lemma A.1. *Let the Gröbner basis of the ideal $\mathcal{I}_\Lambda = \langle \text{SW}_\ell - \chi_\ell \rangle$ in $\mathbb{C}(\chi)[c]$ w.r.t. some monomial order $<$ be given by polynomials*

$$s_m = c^m + \sum_{m' < m} p_{mm'}(\chi) c^{m'}, \quad m \in M, \quad (\text{A.3})$$

where $m := (m_1, \dots, m_L)$, $c^m := \prod_{\ell=1}^L c_\ell^{m_\ell}$, M is a set of tuples m , and $p_{mm'} \in \mathbb{C}(\chi)$.

Let $p_{mm'}$ be finite numbers when evaluated at $\chi_\ell = \bar{\chi}_\ell \in \mathbb{C}$. Then $\bar{s}_m = c^m + \sum_{m' < m} p_{mm'}(\bar{\chi}) c^{m'}$, $m \in M$, form the Gröbner basis of the ideal $\mathcal{I}_\Lambda(\bar{\chi}) = \langle \text{SW}_\ell - \bar{\chi}_\ell \rangle$ in $\mathbb{C}[c]$ for the same monomial order.

In other words, it is safe to specialise a Gröbner basis at those values of χ_ℓ where denominators of $p_{mm'}$ do not vanish.

Proof. To verify the statement first we check that the declared set of \bar{s}_m generates $\mathcal{I}_\Lambda(\bar{\chi})$. To this end, use long division in $\mathbb{C}(\chi)[c_1, \dots, c_L]$ to write $\text{SW}_\ell - \chi_\ell = \sum_m q_m(\chi) s_m$. From the algorithm of long division it is clear that $q_m(\chi)$ are not singular at $\chi = \bar{\chi}$ if $p_{mm'}(\chi)$ are not singular which is the case by the condition of the theorem. Hence $\text{SW}_\ell - \chi_\ell = \sum_m q_m(\chi) s_m$ can be evaluated and still holds at $\chi = \bar{\chi}$. To check that \bar{s}_m form a Gröbner basis we need to *e.g.* compute S-polynomials but this is combinatorially the same exercise as for s_m since the leading monomials are not affected by specialisation. \square

Wronskian equations can be obviously specialised at arbitrary point $\bar{\chi}$ and so the ring $\mathcal{W}_\Lambda(\bar{\chi})$ is always a well-defined object. Now we would like to show that, for a given $\bar{\chi}$, one can find a Gröbner basis that can be specialised at this point and its vicinity. This is the key point to prove the following theorem:

Theorem A.2. $d_\Lambda := \dim_{\mathbb{C}} \mathcal{W}_\Lambda(\bar{\chi})$ does not depend on $\bar{\chi}$.

In other words, number of solutions of Wronskian equations counted with multiplicities is always the same, even on the degeneration set $\mathcal{X}_{\text{crit}}$.

Proof. We know that the theorem holds for all points $\bar{\chi} \notin \text{SW}(D)$ since all the solutions of the Wronskian equations are distinct there and so the dimension of the quotient ring coincides with the number of solutions that we denote as d_Λ . We can path-connect any two regular points and the number of solutions cannot change along the path, see Section 3.1.

Take L linearly independent constant vectors $w_\ell = (w_{\ell 1}, \dots, w_{\ell L})$ and define $x_\ell = \sum_{\ell'} w_{\ell \ell'} c_{\ell'}$. For almost any choice of w_ℓ , the Gröbner basis of \mathcal{I}_Λ in $\mathbb{C}(\chi)$ w.r.t. the monomial order $x_1 < x_2 < \dots < x_L$ should have the form

$$x_1^{d_\Lambda} + a_1^{(d_\Lambda-1)}(\chi) x_1^{d_\Lambda-1} + \dots + a_1^{(0)}, \quad (\text{A.4a})$$

$$x_2 - \sum_{k=0}^{d_\Lambda-1} b_{2k}(\chi) x_1^k, \quad (\text{A.4b})$$

...

$$x_L - \sum_{k=0}^{d_\Lambda-1} b_{Lk}(\chi) x_1^k.$$

Indeed, take a point $\bar{\chi} \notin \mathcal{X}_{\text{crit}}$ for which the conditions of Lemma A.1 hold. At such a point, leading monomials of the Gröbner basis are the same before and after specialisation, and so we can judge about the Gröbner basis from its specialised version. Since $\bar{\chi} \notin \mathcal{X}_{\text{crit}}$, \check{x}_ℓ (regular representation of x_ℓ , written as a matrix in the monomial basis) should have d_Λ distinct eigenvalues for almost any choice of ω_ℓ , and therefore the minimal polynomial equation it satisfies is of degree d_Λ which is (A.4a). In the chosen lexicographic order this equation should belong to the Gröbner basis. Other variables x_2, \dots, x_L should satisfy (A.4b) (*i.e.* they are uniquely fixed if x_1 is fixed) otherwise dimension of $\mathcal{W}_\Lambda(\bar{\chi})$ would exceed d_Λ .

By the properness of WBE $a_1^{(a)}(\chi)$ cannot have singularities, hence they are simply polynomials in χ_ℓ . Coefficients $b_{\ell k}$ however are rational functions of χ_ℓ that can contain

poles. Everywhere outside of these poles, the conditions of Lemma A.1 hold and we can perform specialisation asserting that the dimension of the specialised polynomial ring is d_Λ .

It remains to show that for any $\bar{\chi} \in \mathbb{C}^L$, one can choose ω_ℓ such that $b_{\ell k}$ are not singular at $\bar{\chi}$. To this end, we can actually explicitly express $b_{\ell k}$ in terms of solutions of the Wronskian system. Let $x_\ell = x_\ell^{(i)}$ be the i -th solution. Then polynomials (A.4b) can be rewritten as

$$x_\ell - \sum_{k=0}^{d_\Lambda-1} b_{\ell k}(\chi) x_1^k = \frac{\det \begin{vmatrix} x_\ell & 1 & x_1 & x_1^2 & \dots \\ x_\ell^{(1)} & 1 & x_1^{(1)} & (x_1^{(1)})^2 & \dots \\ x_\ell^{(2)} & 1 & x_1^{(2)} & (x_1^{(2)})^2 & \dots \\ x_\ell^{(3)} & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \end{vmatrix}}{\det \begin{vmatrix} 1 & x_1^{(1)} & (x_1^{(1)})^2 & \dots \\ 1 & x_1^{(2)} & (x_1^{(2)})^2 & \dots \\ 1 & \dots & \dots & \dots \end{vmatrix}}. \quad (\text{A.5})$$

Indeed, equality of the above polynomials to zero implies $x_\ell = x_\ell^{(i)}$ precisely when $x_1 = x_1^{(i)}$.

While $x_\ell^{(i)}$ belong to an algebraic closure of $\mathbb{C}(\chi)$, the above ratio of determinants is symmetric under permutations $x_\ell^i \rightarrow x_\ell^{\sigma(i)}$ and hence should be a polynomial in x_1 with coefficients in the base field, *i.e.* $\mathbb{C}(\chi)$. This follows for instance from $\sum_{i=1}^{d_\Lambda} f(x^{(i)}) = \text{Tr } f(\check{x})$ and basic combinatorial arguments. Of course, one can conclude the same from the fact that (A.4b) are obtained in the process of computation of the Gröbner basis.

At points $\bar{\chi}$ where all $x_1^{(i)}$ are distinct, the denominator of (A.5) is non-zero and hence $b_{\ell k}$ are non-singular. As discussed, for a given regular $\bar{\chi}$, we can adjust ω_ℓ in a way that x_1 has non-degenerate solutions.

When $\bar{\chi} \in \text{SW}(D)$ all x_ℓ degenerate. Then consider a one-parametric smooth path $\chi(t)$ in the space of parameters such that $\chi(t=0) = \bar{\chi}$ is the degeneration point of interest and $\bar{\chi}(t \neq 0) \notin \mathcal{X}_{\text{crit}}$. Moreover, one chooses such a path that all $x_\ell^{(i)}$ are distinct along the path for sufficiently small t , except for the point $t=0$ itself.

The value of the ratio of determinants in (A.5) is not well-defined at $t=0$ but it can be computed as the limit $t \rightarrow 0$. Since this ratio is a rational function of χ_ℓ , the limit, if finite, should produce polynomials (A.4b) specialised at $t=0$.

To compute the limit, note that all x_ℓ , for generic enough choice of ω_ℓ , satisfy one-variable equations $x_\ell^{d_\Lambda} + a_\ell^{(d_\Lambda)} x_\ell^{d_\Lambda-1} + \dots = 0$, where $a_\ell^{(k)}$ are polynomials in χ and hence are well-defined even at $t=0$. Define $x_\ell^i(t)$ as solutions of these equations that coincide with solutions of Wronskian equations for $t \neq 0$; their $t=0$ value is then defined as the continuation $t \rightarrow 0$. If $\mu_{i\ell}$ is the degree of degeneration of solution $x_\ell^{(i)}$ at $t=0$ (*i.e.* $\mu_{i\ell}$ solutions of the one-variable equation on x_ℓ coincide at this point) then $x_\ell^{(i)}(t)$ is expanded in the Puiseux series

$$x_\ell^{(i)}(t) = x_\ell^{(i)}(0) + r_{\ell i,1} t^{1/\mu} + r_{\ell i,2} t^{2/\mu} + \dots, \quad (\text{A.6})$$

where $\mu = \text{LCM}(\mu_{11}, \dots, \mu_{LL})$.

One should know finitely many terms in the series (A.6) to compute the determinants ratio in (A.5) in the limit $t \rightarrow 0$. We require that for these finitely many terms, for each k and i , if at least one $r_{\ell i, k}$ is non-zero then all $r_{\ell i, k}$, $\ell = 1, \dots, L$, are non-zero. It is sufficient to guarantee that the ratio is finite, while imposing such a requirement excludes a measure zero subspace from acceptable values of ω_ℓ . Recall that we already excluded the space of ω_ℓ where the degree of the minimal polynomial of \check{x}_ℓ is less than d_Λ which is of measure zero as well. The majority of ω_ℓ are outside of the stated restrictions, and we can choose any valid option to guarantee the regularity of $b_{\ell k}$ at $\chi = \chi^{(0)}$ and hence the possibility to specialise the Gröbner basis (A.4) at this point thus concluding that $\dim_{\mathbb{C}} \mathcal{W}_\Lambda(\bar{\chi}) = d_\Lambda$. \square

The proposed proof shows that there is a close analogy between WBE and a polynomial equation in a single variable. Indeed, for any point $\bar{\chi} \in \mathcal{X}$, regular or not, we can choose a variable x_1 that satisfies (A.4a) and such that there is a neighbourhood $\mathcal{O}_{\bar{\chi}}$ where $b_{\ell, k}$ are non-singular which allows one to compute all elements of \mathcal{W}_Λ using (A.4b). So a single-variable equation (A.4a) contains all information about \mathcal{W}_Λ in the selected neighbourhood.

A.3 Freeness of \mathcal{W}_Λ and trivialisation of a vector bundle

For each $\mathcal{O}_{\bar{\chi}}$, we have a basis generated by powers of x_1 . Two different bases constructed at $\bar{\chi}$ and $\bar{\chi}'$ are related by a transition matrix which is regular together with its inverse on the intersection of $\mathcal{O}_{\bar{\chi}}$ and $\mathcal{O}_{\bar{\chi}'}$. Hence we get a structure of a holomorphic bundle with fibers being d_Λ -dimensional vector spaces over the field \mathbb{C} and with base \mathcal{X} . The existence of this holomorphic bundle is the same as saying that \mathcal{W}_Λ is a projective $\mathbb{C}[\chi]$ -module. This is the so-called Serre-Swan correspondence [118]³⁷.

Because the base $\mathcal{X} \simeq \mathbb{C}^L$ is contractible, this bundle must be topologically trivial, that is, we can find d_Λ global holomorphic sections forming a basis of the fiber at each point. A much more complicated question, already asked by Serre [118], is whether we can choose these global sections to be polynomials of \mathcal{W}_Λ . A positive answer was given by the Quillen-Suslin theorem [79]. This theorem requires that \mathcal{W}_Λ is a finitely generated $\mathbb{C}[\chi]$ -module, *i.e.* that there exist finitely many elements $\tilde{\mathbf{b}}_1, \dots, \tilde{\mathbf{b}}_{\tilde{d}}$ such that any element of \mathcal{W}_Λ is their linear combination with coefficients from $\mathbb{C}[\chi]$. This is easy to see to be the case. Take for instance the finite set of $\tilde{d} = L \times d_\Lambda$ monomials $x^n := x_1^{n_1}, x_2^{n_2}, \dots, x_L^{n_L}$ with $0 \leq n_i < d_\Lambda$, where x_i are the ones from the proof of Theorem A.2. Due to properness, x_i satisfy a degree- d_Λ equations with polynomial coefficients, *cf.* (A.4a), and hence any higher powers of x_i are expressible as linear combinations of the first d_Λ powers.

The Quillen-Suslin theorem establishes that there are no non-trivial algebraic vector bundles over \mathbb{C}^L or equivalently by the Serre-Swan correspondence, that any finitely generated projective $\mathbb{C}[\chi]$ -module is free, with a basis given by the aforementioned global sections. Applied to \mathcal{W}_Λ , this is precisely the statement that it is a free module over $\mathbb{C}[\chi]$, see Section 3.3.

³⁷We are grateful to L. Cassia for pointing out this relation to us

A.4 Non-symmetric functions

Most of the time we work with only symmetric combinations χ_ℓ of inhomogeneities. However, the Baxter operators as explicit matrices acting on the spin chain have coefficients from $\mathbb{C}[\theta] \equiv \mathbb{C}[\theta_1, \dots, \theta_L]$. This prompts us to understand some properties of $\mathbb{C}[\theta]$ -modules as compared to $\mathbb{C}[\chi]$ -modules. Also, one can consider equations

$$\chi_\ell(\theta_1, \dots, \theta_L) = \chi_\ell \quad (\text{A.7})$$

as a toy model for (2.27) with $c_\ell = \theta_\ell$ and $\text{SW}_\ell(c) = \chi_\ell$.

First, we demonstrate how to use the Gröbner basis techniques to conclude that the polynomial ring $\mathbb{C}[\theta]$ is a free $\mathbb{C}[\chi]$ -module and count the number of solutions to (A.7). To this end denote the elementary symmetric polynomial of degree ℓ in k variables $\theta_1, \dots, \theta_k$ as $\chi_\ell^{(k)}$. Being roots of $\prod_{\ell=1}^k (u - \theta_\ell)$, inhomogeneities satisfy the characteristic equations

$$s_k := \theta_k^k + \sum_{\ell=1}^k (-1)^\ell \chi_\ell^{(k)} \theta_k^{k-\ell} = 0, \quad k = 1, \dots, L. \quad (\text{A.8})$$

Now note that the polynomials $\chi_\ell^{(k)}$ can be rewritten as polynomials in $\chi_{\ell'} \equiv \chi_{\ell'}^{(L)}$, with $\ell' \leq \ell$, and θ_m , $m > k$. As a result, s_k become

$$s_1 = \theta_1 + (-\chi_1 + \sum_{i=2}^L \theta_i), \quad (\text{A.9a})$$

$$s_2 = \theta_2^2 + \theta_2(-\chi_1 + \sum_{i=3}^L \theta_i) + (\chi_2 - (\chi_1 - \sum_{i=3}^L \theta_i) \sum_{i=3}^L \theta_i - \sum_{3 \leq i < j \leq L} \theta_i \theta_j), \quad (\text{A.9b})$$

...

$$s_L = \theta_L^L + \sum_{\ell=1}^{L-1} (-1)^\ell \chi_\ell \theta_L^{L-\ell} + (-1)^L \chi_L. \quad (\text{A.9c})$$

Now let's make a small formalisation: treat θ_ℓ and χ_ℓ as independent variables and consider an ideal $\mathcal{I} = \langle s_1, \dots, s_L \rangle$ as an ideal in $\mathbb{C}[\chi][\theta]$. In the quotient ring $\mathbb{C}[\chi][\theta]/\mathcal{I}$, excluding χ_ℓ in favour of θ_ℓ is easy. However, we are interested in the opposite – to solve for θ_ℓ in terms of χ_ℓ . We won't do this explicitly because this requires an algebraic closure but compute a Gröbner basis instead.

Lemma A.3. *The above-introduced polynomials s_1, \dots, s_L form the Gröbner basis of the ideal $\mathcal{I} = \langle s_1, \dots, s_L \rangle$ w.r.t. a lexicographic order for which $\theta_1 > \theta_2 > \dots > \theta_L > \chi_\ell$, for any ℓ .*

Proof. First, by definition, s_k generate the ideal \mathcal{I} . Then, s_k has θ_k^k as its leading monomial. Indeed, the other monomials are products of $\theta_{k'}^{k'}$ with $k' < k$, powers of $\theta_{k'}$ with $k' > k$, and χ_ℓ which hence are lexicographically smaller than θ_k^k . Finally, as the leading monomials enjoy the property $\text{GCD}(\theta_k^k, \theta_{k'}^{k'}) = 1$, the S-polynomials between s_k and $s_{k'}$ do not produce new relations and so this set of ideal generators is indeed a Gröbner basis. \square

Conceptually the Gröbner basis tells us how to algorithmically find θ_α from the values of their symmetric combinations χ_ℓ . First one needs to solve (A.9c) for fix θ_L (L solutions), then one needs to substitute the found value of θ_L to the equation $s_{L-1} = 0$ and solve it for θ_{L-1} ($L - 1$ solutions) *etc.*

Very similarly to the analysis of the Wronskian algebra, we note that the ring $\mathbb{C}[\chi][\theta]/\mathcal{I}$ is isomorphic (over \mathbb{C}) to $\mathbb{C}[\theta]$, but it is also naturally endowed with the structure of a $\mathbb{C}[\chi]$ -module. The computation of the Gröbner basis above immediately implies that this module is free and of rank $L!$. Indeed, the corresponding monomial basis are given by monomials $\theta_2^{n_2} \theta_3^{n_3} \dots \theta_L^{n_L}$, with $n_\ell < \ell$. Any relations between these monomials is impossible precisely because s_k form a Gröbner basis and leading monomials of s_k do not belong to the monomial basis. Of course we know that $L!$ is an expected number, if to count with multiplicities: the equation $\theta^L - \chi_1 \theta^{L-1} + \dots + (-1)^L \chi_L = 0$ has L solutions, and any permutation of solutions is allowed as well.

Finally, let us extend Theorem 4.2 to the case of non-symmetric polynomials in θ_ℓ . Consider first the following example

Example:

Let the generators of a Wronskian algebra \mathcal{W} satisfy equations $c_1 + c_2 = \chi_1$, $c_1 c_2 = \chi_2$ ^a, and (a hypothetical) Bethe algebra \mathcal{B} is generated by 2×2 diagonal matrices $\hat{c}_1 = \theta_1 \times \mathbf{1}_2$, $\hat{c}_2 = \theta_2 \times \mathbf{1}_2$. These Wronskian and Bethe algebras are isomorphic as $\mathbb{C}[\chi]$ -modules. Let us now consider the extension of the Wronskian algebra $\mathcal{W}^\theta \simeq \mathcal{W} \otimes_{\mathbb{C}[\chi]} \mathbb{C}[\theta]$, *i.e.* consider generators satisfying $c_1 + c_2 = \theta_1 + \theta_2$, $c_1 c_2 = \theta_1 \theta_2$ and treat this algebra as a $\mathbb{C}[\theta]$ -module. This is a rank-two $\mathbb{C}[\theta]$ -module. In contrast, the Bethe algebra considered as a $\mathbb{C}[\theta]$ -module is of rank one.

^aUp to normalisations, it is the $\hbar = 0$ version of (2.40)

By Lemma 4.1 we actually know that generators of type $\theta_\ell \times \mathbf{1}$ cannot appear in polynomial combinations of c_ℓ , and so the hypothetical Bethe algebra in the above example cannot exist.

More generally, we can show that all polynomial relations satisfied \hat{c}_ℓ , even with non-symmetric coefficients, should follow from the Wronskian algebra in the following sense. We can add non-symmetric polynomials by hand to the Wronskian algebra by considering $\mathcal{W}_\Lambda^\theta \simeq \mathcal{W}_\Lambda \otimes_{\mathbb{C}[\chi]} \mathbb{C}[\theta]$. Likewise, non-symmetric polynomials (times the identity operator) are not elements of the Bethe algebra by Lemma 4.1, and hence appending them as extra generators is also realised as $\mathcal{B}_\Lambda^\theta \simeq \mathcal{B}_\Lambda \otimes_{\mathbb{C}[\chi]} \mathbb{C}[\theta]$. Isomorphism between $\mathcal{W}_\Lambda^\theta$ and $\mathcal{B}_\Lambda^\theta$ as $\mathbb{C}[\theta]$ -algebras is then obvious from the isomorphism between \mathcal{B}_Λ and \mathcal{W}_Λ as $\mathbb{C}[\chi]$ -algebras. We also note that $\mathcal{W}_\Lambda^\theta$ and $\mathcal{B}_\Lambda^\theta$ are free as $\mathbb{C}[\theta]$ -modules and $\mathbb{C}[\chi]$ -modules as follows *e.g.* from Lemma A.3.

B Cyclicity of representations

The goal of this appendix is to build all the formalism necessary for the proof of Theorems 4.5 and 4.6. There are two reasons why proving isomorphism of the specialised map

$\varphi_{\bar{\theta}}$ (4.4) is problematic. First, setting θ_ℓ to numerical values, which is done for the Bethe algebra, is more restrictive than setting their symmetric combinations χ_ℓ to numerical values, which is done for the Wronskian algebra. Second, the specialisation procedure is actually native to the representation of an algebra, not to the algebra alone. Namely, we set to numerical values coefficients of a matrix which is more restrictive than setting to numerical values only the factors that multiply this matrix as a whole.

To overcome these difficulties, we want to “rigidify” the algebra isomorphism (4.2) by also proving isomorphism between certain representations of these algebras. As was already mentioned in Section 4.3, the only natural choice of a representation for the Wronskian algebra \mathcal{W}_Λ is its regular representation. As for the Bethe algebra, it acts on the *a priori* unrelated physical space $\text{End}(U_\Lambda) \otimes \mathbb{C}[\theta]$. These two representations are not isomorphic. This is why we need to introduce an alternative Yangian representation dubbed symmetrised representation. Using a cyclic vector argument we prove that its weight subspaces \mathcal{U}_Λ^S are indeed isomorphic to \mathcal{W}_Λ as representations of $\mathcal{B}_\Lambda \simeq \mathcal{W}_\Lambda$, which resolves the second difficulty. This symmetrised representation has the virtue to manifestly depend only on symmetric combinations of inhomogeneities. We show that, under some explicit restriction on $\bar{\theta}$, its specialisation at a point $\bar{\chi}$ is isomorphic to the spin chain representation at a point $\bar{\theta}$ which resolves the first difficulty.

The discussed approach was developed in [9] for \mathfrak{gl}_m spin chains. The below-presented generalisation to the supersymmetric case is conceptually very straightforward. The only difference, apart from the way we present the results, is in the proof of Lemma B.7 which is in line with the ideas of Theorem 4.2.

B.1 Symmetrised Yangian representation

Consider the Yangian spin chain representation at point $\bar{\theta} = (\bar{\theta}_1, \dots, \bar{\theta}_L)$ defined in Section 2.2. We note that the order of inhomogeneities in $\bar{\theta}$ is often superfluous. Indeed, the operator $r_\ell(\bar{\theta}) = (\bar{\theta}_\ell - \bar{\theta}_{\ell+1})\mathcal{P}_{\ell, \ell+1} + \hbar \mathbf{1}$ satisfies (4.1) evaluated at $\theta = \bar{\theta}$. If $\bar{\theta}_\ell \neq \bar{\theta}_{\ell+1} \pm \hbar$, it is invertible and hence an intertwiner between the two representations that differ by the permutation of $\bar{\theta}_\ell$ and $\bar{\theta}_{\ell+1}$. From here we conclude that the isomorphism class of the representation at point $\bar{\theta}$ is decided only by $\bar{\chi}_\ell$ if there is no ℓ, ℓ' such that $\bar{\theta}_\ell - \bar{\theta}_{\ell'} = \pm \hbar$.

More generally, the following facts hold for supersymmetric representations of Yangians:

Proposition B.1. *If $(\bar{\theta}_1, \dots, \bar{\theta}_L)$ satisfy $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$ then the spin chain representation of $Y(\mathfrak{gl}_{m|n})$ at point $\bar{\theta}$ is cyclic with cyclic vector $\mathbf{e}^+ = \mathbf{e}_1 \otimes \dots \otimes \mathbf{e}_1$, where \mathbf{e}_1 is the highest-weight vector of the defining $\mathfrak{gl}_{m|n}$ representation* ³⁸. \square

This statement follows from Theorem 5.2 of [119].

\mathbf{e}^+ is obviously a highest-weight vector of the Yangian representation, *i.e.* it satisfies the condition $T_{ij}\mathbf{e}^+ = 0$ for $i < j$. Its weight is given by

$$T_{ii}\mathbf{e}^+ = Q_\theta(u + \hbar \delta_{i,1})\mathbf{e}^+. \quad (\text{B.1})$$

³⁸In the choice of ordering when bosonic indices are considered smaller than fermionic indices.

Proposition B.2. *The spin chain representation at point $\bar{\theta}$ is irreducible if there are no ℓ, ℓ' such that $\bar{\theta}_\ell - \bar{\theta}_{\ell'} = \hbar$.*

For the $Y(\mathfrak{gl}_m)$ case, this is a standard result appearing in the study of Kirillov-Reshetikhin modules [120]. For $Y(\mathfrak{gl}_{1|1})$ it was proven in [121], Theorem 5. For the $Y(\mathfrak{gl}_{m|n})$ case, it apparently follows from [119], Proposition 5.4. But as this was not stated explicitly we give an alternative argument for irreducibility in the style of statistical lattice models.

Proof. Let $\mathbb{C}^{m|n}$ be the Hilbert space of the ℓ -th node of the spin chain, consider also the auxiliary space $\mathbb{C}^{m|n}$ with basis vectors $\mathbf{e}_\alpha^{\text{aux}}$, $\alpha = 1, 2, \dots, m+n$. The dual basis vectors of $\mathbf{e}_\alpha^{\text{aux}}$ shall be denoted $\mathbf{e}_{\text{aux}}^\alpha$. Define the Lax matrix acting on the tensor product of the mentioned spaces as $\mathcal{L}(u - \theta_\ell) = (u - \theta_\ell) \mathbb{1} + \hbar P$, where P is the graded permutation. In the notations of (2.8), $\mathcal{L}(u - \theta_\ell) := (u - \theta_\ell) \sum_{\alpha, \beta} ev_{\theta_\ell}(t_{\alpha\beta}) \otimes (\mathbf{e}_\beta^{\text{aux}} \otimes \mathbf{e}_{\text{aux}}^\alpha)$. The key property we use is that the Lax matrix becomes, up to normalisation, the graded permutation if $u = \theta_\ell$.

Introduce V^{aux} – the tensor product of L auxiliary spaces spanned by $\mathbf{e}_A^{\text{aux}} = \mathbf{e}_{\alpha_1}^{\text{aux}} \otimes \dots \otimes \mathbf{e}_{\alpha_L}^{\text{aux}}$ and define $B = \sum_A \mathbf{e}_A^{\text{aux}} T_{1\alpha_L}(\bar{\theta}_L) \times \dots \times T_{1\alpha_2}(\bar{\theta}_2) \times T_{1\alpha_1}(\bar{\theta}_1)$. Given the above-mentioned key property, B maps V to $\mathbf{e}^+ \otimes V^{\text{aux}}$ which is easiest to see by a graphical representation of how B acts:

(B.2)

Here each vertical direction corresponds to a node $\mathbb{C}^{m|n}$ of the spin chain and each horizontal direction corresponds to a tensor factor $\mathbb{C}^{m|n}$ of V^{aux} . Intersections are the places where the Lax matrices should be applied (considered as maps from South-West to North-East spaces). Red crosses are the places where the corresponding Lax matrix becomes the permutation.

The map B is also invertible. Indeed, up to a non-zero factor it reduces to an ordered product of $\frac{L(L-1)}{2}$ Lax matrices, *e.g.* these are the three Lax matrices marked by the encircled numbers in the image above. Each of these Lax matrices is invertible since $\bar{\theta}_\ell - \bar{\theta}_{\ell'} \neq \hbar$.

Because B is invertible, for any $v \in V$ one can find a vector $v^* \in (V^{\text{aux}})^*$ such that $(v^*, B)v = \mathbf{e}^+$. Then irreducibility follows from Proposition B.1. \square

Remark We expect that a further refinement of argument (B.2) can be used to prove Proposition B.1 about cyclicity.

For finite-dimensional irreducible $Y(\mathfrak{gl}_{m|n})$ representations, the highest-weight vector exists and is unique and the representation is fully determined, up to an isomorphism, by the vector's weight [122]. Note that the weight of \mathbf{e}^+ depends only on symmetric combinations of inhomogeneities according to (B.1). Then we can consider the induced representation from \mathbf{e}^+ which is isomorphic to the spin chain one by the above-mentioned uniqueness but, in contrast to the spin chain realisation is manifestly invariant under permutations of inhomogeneities.

We shall now introduce a different permutation-invariant realisation which does not require the irreducibility argument and will be formulated for inhomogeneities being abstract variables.

Yangian centraliser Consider the vector space $\mathcal{V} \simeq (\mathbb{C}^{m|n})^{\otimes L} \otimes \mathbb{C}[\theta_1, \dots, \theta_L]$ on which the Yangian representation ev_θ (2.11) is realised. An interesting question is what is the centraliser of the Yangian action on \mathcal{V} .

Define operators S_ℓ acting on \mathcal{V} by

$$S_\ell = \mathcal{P}_{\ell, \ell+1} \Pi_{\ell, \ell+1} - \frac{\hbar}{\theta_\ell - \theta_{\ell+1}} (\Pi_{\ell, \ell+1} - \mathbb{1}), \quad (\text{B.3})$$

where $\mathcal{P}_{\ell, \ell+1}$ is the graded permutation in $(\mathbb{C}^{m|n})^{\otimes L}$, and $\Pi_{\ell, \ell+1}$ permutes variables θ_ℓ and $\theta_{\ell+1}$. These permutations were already used in the proof of Lemma 4.1 on page 28.

Although S_ℓ contains θ_ℓ in denominator, its action on polynomials in θ_ℓ yields again polynomials and hence its action on \mathcal{V} is well-defined.

Lemma B.3. *For $\ell = 1, \dots, L-1$, S_ℓ commutes with the $Y(\mathfrak{gl}_{m|n})$ action, (i.e. $[S_\ell, T_{\alpha\beta}] = 0$) and they form a representation of the symmetric group S_L on \mathcal{V} .*

Proof. The commutativity follows from $(\theta_\ell - \theta_{\ell+1})S_\ell = -\Pi_{\ell, \ell+1} r_\ell + \hbar \mathbb{1}$ and (4.1). Then, by explicit computation one checks $S_\ell^2 = \mathbb{1}$ and $(S_\ell S_{\ell+1})^3 = \mathbb{1}$ – the defining relations of S_L . \square

dAHA In the limit $\hbar \rightarrow 0$, S_L becomes an explicit permutation defined on a graded space that commutes with the action of $\mathfrak{gl}_{m|n}$. Hence $\hbar \neq 0$ should be considered as a generalisation of the Schur-Weyl duality to the case of the Yangian algebra. This statement was made mathematically precise for the bosonic \mathfrak{gl}_m case [123–125]: $(S_\ell)_{1 \leq \ell \leq L-1}$ together with $(\theta_\ell \times \mathbb{1})_{1 \leq \ell \leq L}$ form a representation of \mathcal{H}_L , the *degenerate affine Hecke algebra* (dAHA) on L sites. Moreover, the dAHA and the Yangian form a dual pair – they are maximal mutual centralisers of one another when acting on \mathcal{V} . More formally, one can view \mathcal{V} as the tensor product of S_L -modules $\mathcal{H}_L \otimes_{S_L} (\mathbb{C}^m)^{\otimes L}$ where S_L acts on \mathcal{H}_L as a subalgebra and on $(\mathbb{C}^m)^{\otimes L}$ by permutation of tensor factors. This point of view is conceptually interesting because to generalise Schur-Weyl duality to supersymmetric Yangians, one does not need to change the defining relations of the dAHA \mathcal{H}_L but simply to replace the usual action of S_L on $(\mathbb{C}^n)^{\otimes L}$ by the graded action on $(\mathbb{C}^{m|n})^{\otimes L}$ as in (B.3). The full mathematical treatment (for the affine Hecke algebra³⁹) can be found in [126].

³⁹Recall that the degenerate affine Hecke algebra and the Yangian can be obtained as $q \simeq 1 + \hbar$ expansions of respectively the affine Hecke algebra $\mathcal{H}_L(q)$ and $U_q(\hat{\mathfrak{gl}}_{m|n})$.

We are only going to use the slightly weaker statement of Lemma B.3 that S_ℓ commute with the Yangian action.

$\mathbb{C}[\chi]$ -Yangian module Define $\mathcal{V}^S \subset \mathcal{V}$ as the subspace of S_ℓ -invariant vectors. As $[S_\ell, T_{AB}] = 0$, the Yangian action is well-defined on \mathcal{V}^S . Multiplication by symmetric polynomials is also well-defined on \mathcal{V}^S and, moreover, $\mathbb{C}[\chi] \times \mathbb{1}$ belongs to $ev_\theta(Y(\mathfrak{gl}_{m|n}))$ due to (2.27). Hence we shall call this action of the Yangian on \mathcal{V}^S the *symmetrised Yangian representation*. Note that \mathcal{V}^S is also naturally a $\mathbb{C}[\chi]$ -module.

B.2 Symmetrised Bethe modules and their characters

Since S_L commutes with the global $\mathfrak{gl}_{m|n}$ action it is consistent to define $\mathcal{V}_\Lambda^S = \mathcal{V}^S \cap (V_\Lambda \otimes \mathbb{C}[\theta_1, \dots, \theta_L])$ – the weight Λ subspace of \mathcal{V}^S – and $\mathcal{V}_\Lambda^{S+} = \mathcal{V}^S \cap (V_\Lambda^+ \otimes \mathbb{C}[\theta_1, \dots, \theta_L]) \subset \mathcal{V}_\Lambda^S$ – the subspace of $\mathfrak{gl}_{m|n}$ highest-weight vectors corresponding to the Young diagram Λ^+ . These spaces are also naturally $\mathbb{C}[\chi]$ -modules.

Characters For an element v of \mathcal{V}^S , define its degree as the maximal degree of the monomials in θ_ℓ 's which occur in v . Define $\mathcal{F}_k \mathcal{V}^S$ as the space of all vectors of degree less or equal to k . Finally, define the character

$$\text{ch}(\mathcal{V}^S) = \sum_{k=0}^{\infty} (\dim \mathcal{F}_k / \mathcal{F}_{k-1}) t^k. \quad (\text{B.4})$$

Since the $\mathfrak{gl}_{m|n}$ action does not change the degree, we can also define in an analogous way $\text{ch}(\mathcal{V}_\Lambda^S)$ and $\text{ch}(\mathcal{V}_\Lambda^{S+})$.

Proposition B.4. \mathcal{V}_Λ^S is a free $\mathbb{C}[\chi]$ -module of rank $\binom{L}{\lambda_1, \dots, \lambda_n}$ and its character is given by

$$\text{ch}(\mathcal{V}_\Lambda^S) = t^{\Upsilon_\Lambda} \prod_{a=1}^m \prod_{k=1}^{\lambda_a} \frac{1}{1-t^k} \prod_{i=1}^n \prod_{k=1}^{\nu_i} \frac{1}{1-t^k}, \quad (\text{B.5})$$

where $\Upsilon_\Lambda := \sum_{i=1}^n \frac{\nu_i(\nu_i-1)}{2}$.

Proof. \mathcal{V}_Λ^S is the image of $V_\Lambda \otimes \mathbb{C}[\theta]$ by the projector $p := \frac{1}{L!} \sum_{\sigma \in S_L} S_\sigma$, where the symmetry group acts with S_σ on $V_\Lambda \otimes \mathbb{C}[\theta]$ as generated from S_ℓ (B.3). Since the construction is polynomial in \hbar and the \hbar -term of S_ℓ lowers degrees of polynomials the proposition is true iff it is true for $\hbar = 0$. Hence we will consider only the $\hbar = 0$ case. Every element $\sigma \in S_L$ is then represented as $S_\sigma = \mathcal{P}_\sigma \Pi_\sigma$, where \mathcal{P}_σ is the graded permutation acting on V_Λ and Π_σ is the ordinary permutation acting on $\mathbb{C}[\theta]$.

Denote by \mathbf{e}_α the standard basis vectors of $\mathbb{C}^{m|n}$ defined by $\mathbf{E}_{\beta\beta} \mathbf{e}_\alpha = \delta_{\alpha\beta} \mathbf{e}_\alpha$. The standard spin basis of V_Λ is indexed by tuples $I = (i_1, \dots, i_L)$ such that $|\{k : i_k = a\}| = \lambda_a$ and $|\{k : i_k = i\}| = \nu_i$ corresponding to vectors $\mathbf{e}_I := \otimes_{\ell=1}^L \mathbf{e}_{i_\ell}$.

To get \mathcal{V}_Λ^S , it is enough to consider the image of $\mathbf{w} \otimes \mathbb{C}[\theta]$ by p with

$$\mathbf{w} = \underbrace{\mathbf{e}_1 \otimes \dots \otimes \mathbf{e}_1}_{\lambda_1} \otimes \dots \otimes \underbrace{\mathbf{e}_m \otimes \dots \otimes \mathbf{e}_m}_{\lambda_m} \otimes \underbrace{\mathbf{e}_{m+1} \otimes \dots \otimes \mathbf{e}_{m+1}}_{\nu_1} \otimes \dots \otimes \underbrace{\mathbf{e}_{m+n} \otimes \dots \otimes \mathbf{e}_{m+n}}_{\nu_n}. \quad (\text{B.6})$$

Indeed, for all I we can always find $\sigma \in \mathcal{S}_L$ such that $\mathcal{P}_\sigma \cdot \mathbf{e}_I = \mathbf{e}_{\sigma(I)} = \pm \mathbf{w}$.

Denote by $\mathcal{S}_\Lambda := \prod_{a=1}^m \mathcal{S}_{\lambda_a} \prod_{i=1}^n \mathcal{S}_{\nu_i}$ the stabiliser of \mathbf{w} and by $H := \mathcal{S}_L / \mathcal{S}_\Lambda$ the space of orbits with respect to the right group multiplication. Then the projection by p can be represented as

$$\mathcal{V}_\Lambda^{\mathcal{S}} \simeq p(\mathbf{w} \otimes \mathbb{C}[\theta]) = \frac{1}{L!} \sum_{[\sigma] \in H} S_{[\sigma]} \cdot (\mathbf{w} \otimes R_B \cdot C_F \cdot \mathbb{C}[\theta]), \quad (\text{B.7})$$

where $R_B := \sum_{\sigma \in \prod_{a=1}^m \mathcal{S}_{\lambda_a}} \Pi_\sigma$ and $C_F := \sum_{\sigma \in \prod_{i=1}^n \mathcal{S}_{\nu_i}} (-1)^{|\sigma|} \Pi_\sigma$.

To decide about linear independence in $\mathcal{V}_\Lambda^{\mathcal{S}}$, it is enough to consider one term in the sum $\sum_{[\sigma] \in H} S_{[\sigma]}$, e.g. $[\sigma] = [\mathbf{1}]$ since different terms would be proportional to $\mathcal{P}_\sigma \mathbf{w}$ which are linearly independent in V_Λ . Also, $R_B C_F$ commutes with symmetric polynomials and hence we conclude that $\mathcal{V}_\Lambda^{\mathcal{S}}$ and $R_B C_F \cdot \mathbb{C}[\theta]$ are isomorphic as $\mathbb{C}[\chi]$ -modules.

It is easy to describe $R_B C_F \cdot \mathbb{C}[\theta]$: it is spanned by $Q_\Lambda \times \text{Sym}_{\lambda_1} \times \dots \times \text{Sym}_{\nu_m}$, where $Q_\Lambda := \prod_{i=1}^n \prod_{1 \leq k < l \leq \nu_i} (\theta_{j_i+k} - \theta_{j_i+l})$ with $j_i := \sum_{a=1}^m \lambda_a + \sum_{s=1}^{i-1} \nu_s$, and Sym_k is the space of symmetric polynomials in k variables. $Q_\Lambda \times \text{Sym}_{\lambda_1} \times \dots \times \text{Sym}_{\nu_m}$ is free under the action of $\mathbb{C}[\chi]$, see Appendix A.4, and its character is (B.5). Rank is computed from $\frac{\text{ch}(\mathcal{V}_\Lambda^{\mathcal{S}})}{\text{ch}(\mathbb{C}[\chi])}|_{t=1}$. \square

Corollary B.5. $\mathcal{V}^{\mathcal{S}} = \bigoplus_\Lambda \mathcal{V}_\Lambda^{\mathcal{S}}$ is a free $\mathbb{C}[\chi]$ -module of rank $(\mathbf{m} + \mathbf{n})^L$. \square

Proposition B.6. $\mathcal{V}_\Lambda^{\mathcal{S}^+}$ is a free $\mathbb{C}[\chi]$ -module of rank $\frac{L!}{\prod_{(a,s) \in \Lambda^+} h_{a,s}} = \dim V_\Lambda^+$, where $h_{a,s}$ is the hook length at box (a, s) . Its character is given by

$$\text{ch}(\mathcal{V}_\Lambda^{\mathcal{S}^+}) = t^{\Upsilon_\Lambda^+} \prod_{(a,s) \in \Lambda^+} \frac{1}{1 - t^{h_{a,s}}}, \quad (\text{B.8})$$

where $\Upsilon_\Lambda^+ = \sum_{s=1}^{\lambda_1} \frac{h_s(h_s-1)}{2}$ with h_s being the height of the s -th column of Λ^+ .

Note that $\Upsilon_\Lambda^+ \geq \Upsilon_\Lambda$ which is consistent with $\mathcal{V}_\Lambda^{\mathcal{S}^+} \subset \mathcal{V}_\Lambda^{\mathcal{S}}$.

Proof. As in the previous proof, it is enough to consider $\hbar = 0$.

The space $\mathcal{V}_\Lambda^{\mathcal{S}^+}$ can be constructed from $V \otimes \mathbb{C}[\theta]$ as follows. Take the standard Young tableau \mathcal{T} which is obtained by filling the shape Λ^+ first by filling the boxes defining the weight λ_1 , then λ_2 , ..., then ν_n . For instance, for $[\lambda_1 | \nu_1, \nu_2] = [4 | 2, 2]$, $\mathcal{T} = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 7 & & \\ \hline 6 & 8 & & \\ \hline \end{array}$.

Take the normalised⁴⁰ Young symmetriser $S_{\mathcal{T}} \propto R_{\mathcal{T}} C_{\mathcal{T}}$, where $R_{\mathcal{T}}$ is the symmetrisation over rows and $C_{\mathcal{T}}$ is the antisymmetrisation over columns. Then $\mathcal{V}_\Lambda^{\mathcal{S}^+}$ is the image by the projector p of $S_{R_{\mathcal{T}} C_{\mathcal{T}}} \mathbf{w} \otimes \mathbb{C}[\theta]$. Singling out a special vector \mathbf{w} and (correlated to it) tableau \mathcal{T} is enough to construct $\mathcal{V}_\Lambda^{\mathcal{S}^+}$ because p sums over all permutations. Using this feature of p again, and by repeating the same construction as (B.7), one gets

$$\begin{aligned} \mathcal{V}_\Lambda^{\mathcal{S}^+} &\simeq p(S_{R_{\mathcal{T}} C_{\mathcal{T}}} \mathbf{w} \otimes \mathbb{C}[\theta]) = p(\mathbf{w} \otimes \Pi_{C_{\mathcal{T}} R_{\mathcal{T}}} \mathbb{C}[\theta]) \\ &= \frac{1}{L!} \sum_{[\sigma] \in H} S_{[\sigma]} \cdot (\mathbf{w} \otimes R_B C_F \Pi_{C_{\mathcal{T}} R_{\mathcal{T}}} \cdot \mathbb{C}[\theta]), \end{aligned} \quad (\text{B.9})$$

⁴⁰We normalise all symmetrisations/antisymmetrisations such that they are projectors.

and so \mathcal{V}_Λ^{S+} , as a $\mathbb{C}[\chi]$ -module, is isomorphic to $R_B C_F \Pi_{C_\tau R_\tau} \cdot \mathbb{C}[\theta]$. We can omit $R_B C_F$ as it has zero kernel when acting on $\Pi_{C_\tau R_\tau} \cdot \mathbb{C}[\theta]$. Indeed, $\Pi_{C_\tau R_\tau} R_B C_F \Pi_{C_\tau R_\tau} = \Pi_{C_\tau R_\tau}$. But the module $\Pi_{C_\tau R_\tau} \cdot \mathbb{C}[\theta]$ is the standard application of (reversed) Young symmetriser to a polynomial ring which is well understood, see *e.g.* [103, 127–129]. It is a free $\mathbb{C}[\chi]$ -module with character given by (B.8)⁴¹. Again, the rank is computed from $\frac{\text{ch}(\mathcal{V}_\Lambda^{S+})}{\text{ch}(\mathbb{C}[\chi])}|_{t=1}$. \square

Young diagram dependence One may wonder how comes that the character (B.8) and, in fact, the $\mathbb{C}[\chi]$ -isomorphism class of \mathcal{V}_Λ^{S+} do not depend on $\mathfrak{gl}_{m|n}$ but only on the Young diagram Λ^+ . To understand this property, let us extend the underlying symmetry algebra from $\mathfrak{gl}_{m|n}$ to $\mathfrak{gl}_{m'|n'}$, where $m' = \max(m, h_{\Lambda^+})$ and $n' = \max(n, \lambda_1)$. In addition to $a = 1, \dots, m, i = \hat{1}, \dots, \hat{n}$, introduce also \mathfrak{z} to label the new indices that appeared due to the extension. In the order $a < i < \mathfrak{z}$, the highest-weight vectors do not involve $v_{\mathfrak{z}}$ and hence \mathcal{V}_Λ^{S+} for the extended system is isomorphic to the one of the $\mathfrak{gl}_{m|n}$ system. Now we perform a chain of fermionic duality transformations (odd Weyl reflections) to get any other order in the set a, i, \mathfrak{z} (the order between separately bosonic and fermionic indices will be preserved though). The procedure is described for instance in [96]. It changes the highest-weight vectors, *i.e.* the actual embedding of V_Λ^+ inside V is modified, but this change is performed by acting with elements of the global $\mathfrak{gl}_{m'|n'} \subset Y(\mathfrak{gl}_{m'|n'})$ that commutes with the dAHA and in particular with the action of $\mathbb{C}[\chi]$. Since the procedure is invertible it establishes a $\mathbb{C}[\chi]$ -isomorphism between two spaces \mathcal{V}_Λ^{S+} that differ by the choice of the order defining the highest-weight vector. The order is bijected to a Manhattan-type path (*e.g.* the one in the Example on page 43), and only those indices that belong to the Young diagram part of the path participate in the highest-weight vectors. This last observation allows us to choose m', n' to be any pair such that (m', n') lies on the boundary of Λ^+ (black/red dots of Figure 1) or outside of Λ^+ .

The dependence of \mathcal{V}_Λ^{S+} , as a $\mathbb{C}[\chi]$ -module, on the Young diagram alone parallels results of Section 5 that show that the isomorphism class of the twist-less \mathcal{B}_Λ , as a $\mathbb{C}[\chi]$ -algebra, only depends on the Young diagram. This is of course not a coincidence because the twist-less \mathcal{B}_Λ and \mathcal{V}_Λ^{S+} are isomorphic as $\mathbb{C}[\chi]$ -modules which follows from the results of the next subsection.

B.3 Cyclicity of symmetrised Bethe modules

We shall use the notation \mathcal{U}_Λ^S to cover both \mathcal{V}_Λ^S and \mathcal{V}_Λ^{S+} in the discussion below.

We know that \mathcal{W}_Λ and \mathcal{B}_Λ algebras are isomorphic as $\mathbb{C}[\chi]$ -algebras. The goal of this subsection is to show that the regular representation \mathcal{W}_Λ of the Wronskian algebra and the symmetrised Bethe module \mathcal{U}_Λ^S are isomorphic as $\mathbb{C}[\chi]$ -modules. To do so we need a map from \mathcal{W}_Λ to \mathcal{U}_Λ^S commuting with the action of $\mathcal{W}_\Lambda \equiv \mathcal{B}_\Lambda$. A standard approach is to take a vector $\omega \in U_\Lambda \otimes \mathbb{C}[\theta]$ and to consider the morphism of representations

⁴¹This character is an important combinatorial object: $\frac{\text{ch}(\mathcal{V}_\Lambda^{S+})}{\text{ch}(\mathbb{C}[\chi])}$ is the Kostka-Foulkes polynomial $K_{\mu\nu}(t)$ with $\mu = \Lambda^{+'}$ and $\nu = (1^L)$, see *e.g.* [127, 128].

$$\begin{aligned} \psi_\omega : \mathcal{W}_\Lambda &\longrightarrow U_\Lambda \otimes \mathbb{C}[\theta], \\ c_\ell &\longmapsto \hat{c}_\ell \omega. \end{aligned} \tag{B.10}$$

Of course this map has no reason to be an isomorphism. Nevertheless we can prove the following.

Lemma B.7. *For any non-zero vector $\omega \in U_\Lambda \otimes \mathbb{C}[\theta]$, ψ_ω is injective.*

Proof. Take $P \in \mathcal{W}_\Lambda$ such that $\hat{P}\omega = 0$. As before, use WBE to write P as a polynomial in c_ℓ only. Around a regular point θ , we know from our previous results that the Bethe algebra can be fully diagonalised and that the spectrum of the operators \hat{c}_ℓ are exactly the solutions of the Bethe equations. Denote by $W(\theta)$ the corresponding θ -dependent change of basis that diagonalises \hat{c}_ℓ . At least one of the components of $W(\theta)\omega$ has to be non-zero at θ and hence in some L -dimensional ball \mathcal{O} around θ . Then $\hat{P}\omega = 0$ implies that for one of the solutions $c_\ell(\theta)$, $P(c_\ell(\theta)) = 0$ for $\theta \in \mathcal{O}$. Since c_ℓ is a local diffeomorphism, P vanishes on a L -dimensional ball and thus $P = 0$. \square

We hence see that for all ω , ψ_ω is an isomorphism on its image. Let us take a very precise ω – the vector of the smallest degree ⁴² (as a polynomial in θ_ℓ) that belongs to \mathcal{U}_Λ^S .

Lemma B.8. *$\psi_\omega(\mathcal{W}_\Lambda) = \mathcal{U}_\Lambda^S$. Therefore ψ_ω is an isomorphism of $\mathbb{C}[\chi]$ -modules between \mathcal{W}_Λ and \mathcal{U}_Λ^S .*

Proof. Injectivity is proven by Lemma B.7. The fact that ψ_ω preserves the action of $\mathbb{C}[\chi]$ is obvious from the definition. Note also that ψ_ω just increases the degree by Υ_Λ (by Υ_Λ^+ in the non-twisted case) and otherwise preserves the natural filtrations on \mathcal{W}_Λ and \mathcal{U}_Λ^S . Therefore surjectivity of ψ_ω follows from the comparison of the corresponding characters computed in Sections 3.4 and B.2. \square

B.4 Specialisations

Let us summarise what has been done so far. On one side, we have the Wronskian algebra \mathcal{W}_Λ acting on itself *via* the regular representation. On the other side, we have the Bethe algebra \mathcal{B}_Λ acting on the space \mathcal{U}_Λ^S . Moreover the two couples $(\mathcal{W}_\Lambda, \mathcal{W}_\Lambda)$ and $(\mathcal{B}_\Lambda, \mathcal{U}_\Lambda^S)$ are isomorphic *via* (φ, ψ_ω) .

Now consider the ideal $\mathcal{I} := \langle \text{SW}_1 - \bar{\chi}_1, \dots, \text{SW}_L - \bar{\chi}_L \rangle$ of \mathcal{W}_Λ and its image $\hat{\mathcal{I}}$ in \mathcal{B}_Λ under φ . Automatically, the isomorphisms (φ, ψ_ω) will induce isomorphisms between $(\mathcal{W}_\Lambda/\mathcal{I}, \mathcal{W}_\Lambda/\mathcal{I})$ and $(\mathcal{B}_\Lambda/\hat{\mathcal{I}}, \mathcal{U}_\Lambda^S/\psi_\omega(\mathcal{I}))$. Moreover, as shown in Lemma 4.4 $\psi_\omega(\mathcal{I}) = \mathcal{J} \cdot \mathcal{U}_\Lambda^S$, where $\mathcal{J} := \langle \chi - \bar{\chi} \rangle \subset \mathbb{C}[\chi]$. Denote $\mathcal{B}_\Lambda(\bar{\chi}) := \mathcal{B}_\Lambda/\hat{\mathcal{I}}$ and recall that $\mathcal{W}_\Lambda(\bar{\chi}) := \mathcal{W}_\Lambda/\mathcal{I}$.

We also have a third pair in this correspondence, namely $\mathcal{B}_\Lambda(\bar{\theta})$, the Bethe algebra evaluated at $\bar{\theta}$, acting on U_Λ . Our final goal is to show that $\varphi_{\bar{\theta}} : \mathcal{W}_\Lambda(\bar{\chi}) \simeq \mathcal{B}_\Lambda(\bar{\theta})$. This is equivalent to showing that $\mathcal{B}_\Lambda(\bar{\theta}) \simeq \mathcal{B}_\Lambda(\bar{\chi})$. Let us emphasise that *a priori* $\mathcal{B}_\Lambda(\bar{\chi})$ and $\mathcal{B}_\Lambda(\bar{\theta})$ are two different objects.

⁴²It is unique up to a normalisation as follows from (B.5) and (B.8).

Example:

Consider \mathcal{W} and \mathcal{B}^{bad} from the example on page 30. \mathcal{B}^{bad} acts on the space $\mathcal{V} = \mathbb{C}^2 \otimes \mathbb{C}[\theta_1, \theta_2]$. \mathcal{V} is a $\mathbb{C}[\chi_1, \chi_2]$ -module of rank four, we can take $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\theta_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, $\theta_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ as its basis elements. In this basis

$$\hat{c}_1^{\text{bad}} = \begin{pmatrix} 0 & -\chi_2 & 0 & 0 \\ 1 & \chi_1 & 0 & 0 \\ 0 & 0 & 0 & -\chi_2 \\ 0 & 0 & 1 & \chi_1 \end{pmatrix}. \quad (\text{B.11})$$

Take a vector $\omega = A \begin{pmatrix} 1 \\ 0 \end{pmatrix} + B \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, where A, B are some polynomials in χ_1, χ_2 . Then $\mathcal{U}^{\text{S}} := \psi_\omega(\mathcal{W})$ is a $\mathbb{C}[\chi_1, \chi_2]$ -module of rank two spanned by $\xi_1 := \omega$, and $\xi_2 := A \theta_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + B \theta_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. $\hat{c}_1^{\text{bad}} \in \mathcal{B}^{\text{bad}}$ acting on \mathcal{U}^{S} is $\begin{pmatrix} 0 & -\chi_2 \\ 1 & \chi_1 \end{pmatrix}$ in the basis ξ_1, ξ_2 .

Specialisation $\mathcal{B}^{\text{bad}}(\bar{\chi})$ is two-dimensional for any $\bar{\chi}$ and is clearly \mathbb{C} -isomorphic to $\mathcal{W}(\bar{\chi})$, in particular $\hat{c}_1^{\text{bad}}(\bar{\chi}) = \begin{pmatrix} 0 & -\bar{\chi}_2 \\ 1 & \bar{\chi}_1 \end{pmatrix}$. Note that the statement is completely independent of the choice of ω . It holds even if $A(\bar{\chi}_1, \bar{\chi}_2) = B(\bar{\chi}_1, \bar{\chi}_2) = 0$ because $\xi_i \notin \psi_\omega(\mathcal{I}) := \langle \chi - \bar{\chi} \rangle \mathcal{U}^{\text{S}}$, $i = 1, 2$.

We chose \mathcal{B}^{bad} in the example above to explicitly demonstrate that $\mathcal{B}_\Lambda(\bar{\theta})$ and $\mathcal{B}_\Lambda(\bar{\chi})$ can be in principle non-isomorphic.

Instead of showing isomorphism between $\mathcal{B}_\Lambda(\bar{\chi})$ and $\mathcal{B}_\Lambda(\bar{\theta})$ directly, let us show isomorphism between $\mathcal{U}_\Lambda^{\text{S}}/\mathcal{J} \cdot \mathcal{U}_\Lambda^{\text{S}}$ and U_Λ . Since these spaces both carry representations of the Bethe algebra, if one can find an isomorphism commuting with these actions, it would automatically imply $\mathcal{B}_\Lambda(\bar{\chi}) \equiv \mathcal{B}_\Lambda(\bar{\theta})$ as is argued in Section 4.3. The advantage of this strategy is that we can leverage Yangian representation theory to prove such an isomorphism.

To relate the symmetrised Yangian representation at point $\bar{\chi}$ and the spin chain Yangian representation at point $\bar{\theta}$, recall that \mathcal{V}^{S} is a subspace of $\mathcal{V} := (\mathbb{C}^{m|n})^{\otimes L} \otimes \mathbb{C}[\theta_1, \dots, \theta_L]$ and so we can define a map $\text{Ev}_{\bar{\theta}} : \mathcal{V}^{\text{S}} \rightarrow (\mathbb{C}^{m|n})^{\otimes L}$ simply by evaluating all vectors at $\bar{\theta}$. Since $\mathcal{J} \cdot \mathcal{V}^{\text{S}} \subset \text{Ker Ev}_{\bar{\theta}}$, this induces a well-defined map

$$\text{ev}_{\bar{\theta}} : \mathcal{V}^{\text{S}}(\bar{\chi}) \rightarrow (\mathbb{C}^{m|n})^{\otimes L}, \quad (\text{B.12})$$

where $\mathcal{V}^{\text{S}}(\bar{\chi}) := \mathcal{V}^{\text{S}}/\mathcal{J} \cdot \mathcal{V}^{\text{S}}$. Concretely this just means the following: take a class $[v] \in \mathcal{V}^{\text{S}}(\bar{\chi})$, represent it by some $v \in \mathcal{V}$ and evaluate it at $\bar{\theta}$.

Note that $\mathcal{V}^{\text{S}}(\bar{\chi})$ is the space where the symmetrised Yangian representation at point $\bar{\chi}$ is realised and $(\mathbb{C}^{m|n})^{\otimes L}$ is the Hilbert space of the spin chain. We can realise on it the spin chain Yangian representation at point $\bar{\theta}$.

We are ready to formulate the main conceptual result of this appendix which is Proposition 3.5 of [9] generalised to the supersymmetric case.

Theorem B.9. Let $\bar{\theta} = (\bar{\theta}_1, \dots, \bar{\theta}_L)$ be a solution of equations $\chi_\ell(\theta) = \bar{\chi}_\ell$ such that $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$. Then $\text{ev}_{\bar{\theta}}$ is an isomorphism of $Y(\mathfrak{gl}_{\mathbf{m}|\mathbf{n}})$ representations.

Proof. Since $\text{ev}_{\bar{\theta}}$ commutes with the Yangian action, it defines a homomorphism from the symmetrised representation to the spin chain representation. Assume $\mathbf{m} \geq 1$ and consider the vector $\mathbf{e}^+ := \mathbf{e}_1^{\otimes L} \in \mathcal{V}^S(\bar{\chi})$. Since $\text{ev}_{\bar{\theta}} : \mathbf{e}^+ \mapsto \mathbf{e}^+$, and \mathbf{e}^+ is a cyclic vector of the spin chain module by Theorem B.1, $\text{ev}_{\bar{\theta}}$ is surjective. As $\mathcal{V}^S(\bar{\chi})$ and $(\mathbb{C}^{\mathbf{m}|\mathbf{n}})^{\otimes L}$ are of the same dimension by Corollary B.5, $\text{ev}_{\bar{\theta}}$ is an isomorphism.

If $\mathbf{m} = 0$, one can check that \mathcal{V}^S contains the vector $\prod_{1 \leq \ell < \ell' \leq L} (\theta_\ell - \theta_{\ell'} + \hbar) \mathbf{e}^+$, whose image under $\text{ev}_{\bar{\theta}}$ is nonzero as long as $\bar{\theta}_\ell + \hbar \neq \bar{\theta}_{\ell'}$ for $\ell < \ell'$. The rest of the proof is the same. \square

B.5 An explicit case study

Finally, we provide a concrete comprehensive example to illustrate the above-discussed ideas.

Explicit Q-operators Consider the $L = 3$ \mathfrak{gl}_2 spin chain in the absence of twist. By convention, we use the following basis of $(\mathbb{C}^2)^{\otimes 3}$

$$\{|\downarrow\downarrow\downarrow\rangle, |\uparrow\downarrow\downarrow\rangle, |\downarrow\uparrow\downarrow\rangle, |\downarrow\downarrow\uparrow\rangle, |\downarrow\uparrow\uparrow\rangle, |\uparrow\downarrow\uparrow\rangle, |\uparrow\uparrow\downarrow\rangle, |\uparrow\uparrow\uparrow\rangle\}. \quad (\text{B.13})$$

In this basis, the periodic (“twist-less”) limit of the Q -operators is:

$$Q_\emptyset = 1, \quad Q_1 = \begin{pmatrix} 1 & & & \\ & M_1 & & \\ & & M_1 & \\ & & & 1 \end{pmatrix}, \quad Q_{12} = \prod_{i=1}^3 (u - \theta_i), \quad (\text{B.14})$$

where M_1 is the following 3×3 block matrix

$$M_1 = \underbrace{\frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}}_{P_s} + \left(u - 2 \frac{\theta_1 + \theta_2 + \theta_3}{3} \right) \underbrace{\frac{1}{3} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}}_{P_h} + \frac{1}{6} \times \underbrace{\begin{pmatrix} 2\theta_2 + 2\theta_3 & -\hbar - 2\theta_3 & \hbar - 2\theta_2 \\ \hbar - 2\theta_3 & 2\theta_1 + 2\theta_3 & -\hbar - 2\theta_1 \\ -\hbar - 2\theta_2 & \hbar - 2\theta_1 & 2\theta_1 + 2\theta_2 \end{pmatrix}}_{c_0}. \quad (\text{B.15})$$

Notice that the projector to the symmetric irrep $\square\square$ is $\begin{pmatrix} 1 & & \\ & P_s & \\ & & P_s & \\ & & & 1 \end{pmatrix}$, whereas the projector to hook irreps $2 \times \square$ is $\begin{pmatrix} 0 & & \\ & P_h & \\ & & P_h & \\ & & & 0 \end{pmatrix}$. Also notice that $c_0 P_s = 0$, *i.e.* c_0 only affects the hook irreps \square .

In addition to these notations, use $\chi_1 = \theta_1 + \theta_2 + \theta_3$, $\chi_2 = \theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3$, $\chi_3 = \theta_1\theta_2\theta_3$, and get

$$Q_2 = \frac{-6u^4 + 8u^3\chi_1 + u^2(3\hbar^2 - 12\chi_2) + u(-2\hbar^2\chi_1 + 24\chi_3)}{24\hbar} \begin{pmatrix} 1 & P_s \\ & P_s \\ & & 1 \end{pmatrix} + \left(-\frac{u^3}{2\hbar} + \frac{\hbar\chi_1}{12} - \frac{\chi_3}{\hbar}\right) \begin{pmatrix} 0 & P_h \\ & P_h \\ & & 0 \end{pmatrix} + \left(\frac{u^2}{4\hbar} - \frac{\hbar}{12}\right) \begin{pmatrix} 0 & c_0 \\ & c_0 \\ & & 0 \end{pmatrix}, \quad (\text{B.16})$$

$$\mathbb{Q}_{0,1} = \left(3\hbar u^2 - 2\hbar\chi_1 u + \hbar\chi_2 + \frac{\hbar^3}{4}\right) \begin{pmatrix} 1 & P_s \\ & P_s \\ & & 1 \end{pmatrix} + 3\hbar u \begin{pmatrix} 0 & P_h \\ & P_h \\ & & 0 \end{pmatrix} - \frac{\hbar}{2} \begin{pmatrix} 0 & c_0 \\ & c_0 \\ & & 0 \end{pmatrix}. \quad (\text{B.17})$$

Restriction to the hook irreps If we restrict to the subspace $2 \times \boxplus$, we obtain 2×2 matrices written for instance in the basis ⁴³

$$\left\{ \frac{\sqrt{3}+3}{6} |\uparrow\downarrow\downarrow\rangle - \frac{\sqrt{3}}{3} |\downarrow\uparrow\downarrow\rangle + \frac{\sqrt{3}-3}{6} |\downarrow\downarrow\uparrow\rangle, \frac{\sqrt{3}-3}{6} |\uparrow\downarrow\downarrow\rangle - \frac{\sqrt{3}}{3} |\downarrow\uparrow\downarrow\rangle + \frac{\sqrt{3}+3}{6} |\downarrow\downarrow\uparrow\rangle \right\}. \quad (\text{B.18})$$

In this basis, c_0 becomes the 2×2 matrix

$$c := c_0 = \begin{pmatrix} 2\chi_1 - \sqrt{3}(\theta_1 - \theta_3) & \chi_1 - 3\theta_2 + \sqrt{3}\hbar \\ \chi_1 - 3\theta_2 - \sqrt{3}\hbar & 2\chi_1 + \sqrt{3}(\theta_1 - \theta_3) \end{pmatrix} \quad (\text{B.19})$$

and equations (B.15), (B.16) and (B.17) become respectively

$$Q_1 \Big|_{2 \times \boxplus} = (u - 2\frac{\chi_1}{3})\mathbb{I} + \frac{c}{6} = \begin{pmatrix} u - \frac{\chi_1}{3} - \frac{\sqrt{3}}{6}(\theta_1 - \theta_3) & \frac{\chi_1}{6} - \frac{\theta_2}{2} + \frac{\sqrt{3}}{6}\hbar \\ \frac{\chi_1}{6} - \frac{\theta_2}{2} - \frac{\sqrt{3}}{6}\hbar & u - \frac{\chi_1}{3} + \frac{\sqrt{3}}{6}(\theta_1 - \theta_3) \end{pmatrix} \quad (\text{B.20})$$

$$Q_2 \Big|_{2 \times \boxplus} = \left(-\frac{u^3}{2\hbar} + \frac{\hbar\chi_1}{12} - \frac{\chi_3}{\hbar}\right)\mathbb{I} + \left(\frac{u^2}{4\hbar} - \frac{\hbar}{12}\right)c \quad (\text{B.21})$$

$$\mathbb{Q}_{0,1} \Big|_{2 \times \boxplus} = 3\hbar u - \frac{\hbar}{2}c = 3\hbar \begin{pmatrix} u - \frac{\chi_1}{3} + \frac{\sqrt{3}}{6}(\theta_1 - \theta_3) & -\frac{\chi_1}{6} + \frac{\theta_2}{2} - \frac{\sqrt{3}}{6}\hbar \\ -\frac{\chi_1}{6} + \frac{\theta_2}{2} + \frac{\sqrt{3}}{6}\hbar & u - \frac{\chi_1}{3} - \frac{\sqrt{3}}{6}(\theta_1 - \theta_3) \end{pmatrix} \quad (\text{B.22})$$

Symmetrised modules Let us now compute these operators in the symmetrised Yangian representation. This results in presenting a θ -dependent change of basis such that all the matrix coefficients of c , the only non-trivial operator of the Bethe algebra, are symmetric polynomials in θ_ℓ . We will explicitly compute this basis by using the proof of B.8.

Let us first consider the case $\hbar = 0$. The normalised Young symmetriser for $\mathcal{T} = \boxed{\frac{1}{3} \frac{2}{3}}$ is given by

$$S_{\mathcal{T}} = \frac{1}{3} R_{\mathcal{T}} C_{\mathcal{T}} = \frac{1}{3} (1 + (2 \ 1 \ 3) - (3 \ 2 \ 1) - (3 \ 1 \ 2)). \quad (\text{B.23})$$

⁴³This basis is an orthogonal basis of the 2D subspace, the expression of which is ‘‘quite symmetric’’.

We now have to compute $S_{\mathcal{T}} \cdot \mathbb{C}[\theta]$ and moreover to find a $\mathbb{C}[\chi]$ -basis for it. Start by picking a $\mathbb{C}[\chi]$ -basis of $\mathbb{C}[\theta]$, for example the Schubert polynomials (in the case of S_3 they are given by $\{1, \theta_1, \theta_1 + \theta_2, \theta_1^2, \theta_1\theta_2, \theta_1^2\theta_2\}$)⁴⁴. Since $S_{\mathcal{T}}$ commutes with multiplication by symmetric polynomials we just have to compute its action on Schubert polynomials. Eliminating obvious redundancies we obtain only two $\mathbb{C}[\chi]$ -independent basis elements

$$\eta_1 := \theta_1 + \theta_2 - 2\theta_3 \quad \eta_2 := 2\theta_1\theta_2 - \theta_1\theta_3 - \theta_2\theta_3. \quad (\text{B.24})$$

At this stage we can already check the character formula (B.8). Indeed

$$\frac{\text{ch}(\mathcal{V}_{(2,1)}^{\mathcal{S}^+})}{\text{ch}(\mathbb{C}[\chi])} = \frac{t(1-t)(1-t^2)(1-t^3)}{(1-t)^2(1-t^3)} = t(1+t) = t^{\deg \eta_1} + t^{\deg \eta_2}. \quad (\text{B.25})$$

To obtain a $\mathbb{C}[\chi]$ -basis of $\mathcal{V}_{(2,1)}^{\mathcal{S}^+}$ it remains to compute $p(\mathbf{w} \otimes \eta_1)$ and $p(\mathbf{w} \otimes \eta_2)$ which can be done straightforwardly.

Now assume $\hbar \neq 0$. This case is more complicated because now we have to take \hbar corrections into account. In particular now $p(\mathbf{w} \otimes \eta_1), p(\mathbf{w} \otimes \eta_2) \notin \mathcal{V}_{(2,1)}^{\mathcal{S}^+}$. Nevertheless $p(\mathbf{w} \otimes \eta_1), p(\mathbf{w} \otimes \eta_2) \in \mathcal{V}_{(2,1)}^{\mathcal{S}}$ and we have to correct them by some vectors of lower degree such that they belong to $\mathcal{V}_{(2,1)}^{\mathcal{S}^+}$. Since $\deg p(\mathbf{w} \otimes \eta_1) = \deg \eta_1 = 1$ it can only be corrected by a vector of degree zero. There is only one such vector in $\mathcal{V}_{(2,1)}^{\mathcal{S}}$: the totally symmetric combination $|\uparrow\downarrow\downarrow\rangle + |\downarrow\uparrow\downarrow\rangle + |\downarrow\downarrow\uparrow\rangle$. Its coefficient can be uniquely fixed by requiring that the corrected vector is highest-weight. By a similar argument we can compute the \hbar corrections to $p(\mathbf{w} \otimes \eta_2)$. In the end we obtain the following $\mathbb{C}[\chi]$ -basis of $\mathcal{V}_{(2,1)}^{\mathcal{S}^+}$

$$\begin{aligned} \xi_1 &:= \frac{1}{6}(-2\theta_1 + \theta_2 + \theta_3 - 3\hbar) |\uparrow\downarrow\downarrow\rangle \\ &\quad + \frac{1}{6}(\theta_1 - 2\theta_2 + \theta_3) |\downarrow\uparrow\downarrow\rangle \\ &\quad + \frac{1}{6}(\theta_1 + \theta_2 - 2\theta_3 + 3\hbar) |\downarrow\downarrow\uparrow\rangle \\ \xi_2 &:= \frac{1}{18}(-3\theta_1\theta_2 - 3\theta_1\theta_3 + 6\theta_2\theta_3 - \hbar(\theta_1 + 4\theta_2 + 4\theta_3)) |\uparrow\downarrow\downarrow\rangle \\ &\quad + \frac{1}{18}(-3\theta_1\theta_2 + 6\theta_1\theta_3 - 3\theta_2\theta_3 - \hbar(4\theta_1 + \theta_2 - 5\theta_3) - 3\hbar^2) |\downarrow\uparrow\downarrow\rangle \\ &\quad + \frac{1}{18}(6\theta_1\theta_2 - 3\theta_1\theta_3 - 3\theta_2\theta_3 + \hbar(5\theta_1 + 5\theta_2 - \theta_3) + 3\hbar^2) |\downarrow\downarrow\uparrow\rangle \end{aligned} \quad (\text{B.26})$$

that is, $\mathcal{V}_{(2,1)}^{\mathcal{S}^+} = \mathbb{C}[\chi]\xi_1 \oplus \mathbb{C}[\chi]\xi_2$. Changing c to this basis we finally obtain its symmetrised representative

$$c^{\mathcal{S}} := \begin{pmatrix} -\hbar & -2\chi_2 - \frac{2}{3}\hbar(\chi_1 + \hbar) \\ 6 & 4\chi_1 + \hbar \end{pmatrix}. \quad (\text{B.27})$$

A few remarks are in order. First, note that c is homogeneous of degree 1 (if we consider $\deg \hbar = 1$) whereas $c^{\mathcal{S}}$ is non homogeneous of degree 2. This has to do with the fact that

⁴⁴It is also a good example to check equation (B.5). Indeed $\frac{\text{ch}(\mathbb{C}[\theta])}{\text{ch}(\mathbb{C}[\chi])} = \frac{(1-t)(1-t^2)(1-t^3)}{(1-t)^3} = 1 + 2t + 2t^2 + t^3$ which is exactly the character of the Schubert basis.

ξ_1 and ξ_2 are θ -dependent and of different degrees. Second, one can check that c and c^S have the same characteristic polynomial and therefore the same spectrum. However there is a crucial difference: if we set $\hbar = 0$ and take $\bar{\theta}_1 = \bar{\theta}_2 = \bar{\theta}_3$ (a potentially “bad” point by Theorem B.9), c becomes proportional to the identity whereas c^S does not, so they cannot be related by a change of basis. In particular the symmetrised Bethe algebra $\mathcal{B}_\Lambda(\bar{\chi})$ is still maximal, whereas the evaluated Bethe algebra $\mathcal{B}_\Lambda(\bar{\theta})$ is not. Again this has to do with the fact that the basis (ξ_1, ξ_2) is θ -dependent: at this particular point it becomes degenerate *as a basis of the physical vector space*, but not *as a basis of the quotient* $\mathcal{V}_{(2,1)}^{S+}(\bar{\chi})$. Actually, the determinant of the matrix of change of basis between (B.18) and (ξ_1, ξ_2) is given by $\frac{-1}{4\sqrt{3}}(\theta_1 - \theta_2 + \hbar)(\theta_1 - \theta_3 + \hbar)(\theta_2 - \theta_3 + \hbar)$ and it equates to zero precisely at the “bad” points of Theorem B.9. Note however that even at most “bad” points $\mathcal{B}_\Lambda(\bar{\theta})$ and $\mathcal{B}_\Lambda(\bar{\chi})$ are still isomorphic and maximal.

This example shows that a $\mathbb{C}[\chi]$ -basis of the symmetrised Yangian representation is quite difficult to compute. As long as the hypothesis of Theorem B.9 is satisfied, the traditional physical frame is perfectly equivalent to the symmetrised one and can be used without trouble for all practical applications.

C Q-operators belong to the Bethe algebra

In order to show that the Q functions/operators belong to the Bethe algebra we will show how to find them from T functions/operators. When $a = 1$, the contraction with the Levi-Civita tensor in the Wronskian expression (2.15) reduces to $(m-1)! \sum_b (-1)^b Q_b^{[m-n+s]} Q_b^{[-s]}$, hence we can express the t -dependent sum (where t is a free parameter)

$$S(t) := \sum_b \left(\sum_{s \geq 1} Q_b^{[-m+n-2s]} t^s \right) Q_b \quad (\text{C.1})$$

as an infinite linear combination of the $\mathbb{T}_{(s^1)}$: this linear combination looks like $\sum_{s \geq 1} \mathbb{T}_{(s^1)}^{[-m+n-s]} t^s$, up to the first terms (when $s < a - m + n$) and up to factors that have no impact on the present argument and would make expressions extremely bulky. These are the factors $Q_{\bar{\theta}|\bar{\theta}}$, $\text{Ber } G$ (which we set to 1), $(m-1)!$ and the proportionality⁴⁵ factor denoted by the symbol \propto in (2.15).

This infinite sum (where s runs from 0 to $+\infty$) converges in the disk $|t| < \min \left| \frac{1}{x_b} \right|$ and is then analytically continued to $t \in \mathbb{C} \setminus \left\{ \frac{1}{x_b} \mid 1 \leq b \leq m \right\}$. Indeed, if we denote $Q_{\bar{b}} =$

⁴⁵This proportionality factor is a supersymmetric version of a Vandermonde determinant of the eigenvalues z_α , as can be found from requiring that q_θ, q_a and q_i are monic.

$\left(\frac{1}{x_b}\right)^{u/\hbar} \sum_{k=0}^{M_{\bar{b}}} c_{\bar{b}}^{(k)} u^k$ then for $|t| < \min \left| \frac{1}{x_b} \right|$ we have:

$$\sum_{s \geq 1} Q_{\bar{b}}^{[-m+n-2s]} t^s = \left(\frac{1}{x_b}\right)^{\frac{-m+n}{2}} \sum_{j=0}^{M_{\bar{b}}} (-1)^j \left(\sum_{k=j}^{M_{\bar{b}}} \binom{k}{j} (u^{[-m+n]})^{k-j} c_{\bar{b}}^{(k)} \right) \underbrace{\sum_{s \geq 1} s^j (t x_b)^s}_{\frac{\sum_{k=0}^{j-1} A(j,k) (t x_b)^{k+1}}{(1-t x_b)^{j+1}}}, \quad (\text{C.2})$$

where the combinatorial factors $A(j, k)$ are the positive integers known as the Eulerian numbers, and where the sum $\sum_{k=0}^{j-1} A(j, k) (t x_b)^{k+1}$ should be replaced by 1 in the ill-defined case $j = 0$.

If the eigenvalues x_b are pairwise distinct we deduce that

$$Q_b \propto \lim_{t \rightarrow \frac{1}{x_b}} (1 - t x_b)^{M_{\bar{b}}+1} S(t). \quad (\text{C.3})$$

This expression allows one to conclude that, at the level of representations, it belongs to the Bethe algebra. Indeed, the Bethe Algebra forms a linear subspace of the space of operators on the Hilbert space, which is finite-dimensional. It is hence topologically closed, so that the sum $S(t)$ belongs to the Bethe algebra, not only when $|t| < \min \left| \frac{1}{x_b} \right|$ but even for arbitrary t by analytic continuation. Then, by taking the limit (C.3) Q_b belongs to the Bethe Algebra.

In addition to the Q-operators Q_b ($1 \leq b \leq m$), the same approach also allows producing the operators Q_i ($\hat{1} \leq i \leq \hat{n}$) by focusing on a sum of the form $\sum_{a \geq 0} \mathbb{T}_{(1^a)}^{[-m+n+a]} t^a$. Hence it allows expressing all Q-operators using (2.21) and (2.19).

In [64], explicit computations of such infinite sums and of their limit were performed combinatorially at the level of the representation ev_θ (for twist with pairwise-distinct eigenvalues, as in the above discussion). This explicit construction of the Q-operators shows that their matrix coefficients are indeed polynomial functions of u and of the inhomogeneities θ_ℓ , and that they are rational functions of the twist eigenvalues z_α . It also shows that the degree $M_{A|I}$ of $q_{A|I}$ is indeed given by (2.25).

For comparison purposes, we note that the infinite sum which was computed in [64] is actually of the form $\sum_{s \geq 1} \mathbb{T}_{(s^1)}^{[+s]} t^s$, by contrast with the opposite shifts in the above discussion. Consequently the explicit combinatorial description gives an expression of $Q_{\bar{b}}$ instead of Q_b , and Q_b was extracted after a few more steps – after $m - 1$ successive limits – and the whole Q-system is expressed explicitly.

D Details about structural study of Bethe equations

In this section we will adopt the analytic point of view of Section 3.1 on inhomogeneities. Namely we will think them as complex numbers that we are going to vary. Then c_ℓ – the coefficients of (twisted) Baxter polynomials – turn out to be algebraic functions of

inhomogeneities. The main purpose of this section is to prove properness of WBE – that all solutions c_ℓ are bounded if θ_ℓ (and hence χ_ℓ) are bounded. A natural consequence of this analysis will be the behaviour of c_ℓ when θ_ℓ tend to infinity (in the twist-less case) which we analyse in D.3 to provide the necessary technical results for Section 6.2.

D.1 Properness - twisted case

The feature that ensures that c_α are bounded at finite θ_α is the fact that c_α are coefficients of polynomials in u and the quantization condition (2.26) is an equation on these polynomials.

Assume that there is a point $\bar{\chi} \in \mathcal{X}$ by approaching which some of c_α diverge (become unbounded). If c_α is a coefficient of a twisted polynomial $Q(u)$ then divergence of c_α implies divergence of some of the roots of $Q(u)$. Factorise $Q(u)$ in the form $Q = Q^{\gg} Q^{\lessgtr}$, where Q^{\gg} is a polynomial containing all diverging roots, and Q^{\lessgtr} is the function containing the twist prefactor and all finite roots.

Consider first the equation $Q_{a|i}^+ - Q_{a|i}^- = Q_{a|\emptyset} Q_{\emptyset|i}$. One can rewrite $Q_{a|i}^+ - Q_{a|i}^- = Q_{a|i}^{\gg} ((Q_{a|i}^{\lessgtr})^+ - (Q_{a|i}^{\lessgtr})^-) + H$, where H is the function that ensures equality. By taking a point χ sufficiently close to the point $\bar{\chi}$, one can ensure that H is small in the following sense: there exists R such that all roots of $(Q_{a|i}^{\lessgtr})^+ - (Q_{a|i}^{\lessgtr})^-$ lie inside the circle $|u| = R$, all roots of Q^{\gg} lie outside it, and that the absolute value of H is smaller than the absolute value of $Q_{a|i}^{\gg} ((Q_{a|i}^{\lessgtr})^+ - (Q_{a|i}^{\lessgtr})^-)$ when $|u| = R$. Then by Rouché's theorem, the number of zeros of $Q_{a|i}^+ - Q_{a|i}^-$ inside and outside of the circle is the same as that of $Q_{a|i}^{\gg} ((Q_{a|i}^{\lessgtr})^+ - (Q_{a|i}^{\lessgtr})^-)$. Hence existence of large zeros of $Q_{a|i}$ imply existence of large zeros, in the same amount, in the product $Q_{a|\emptyset} Q_{\emptyset|i}$. Their distribution between $Q_{a|\emptyset}$ and $Q_{\emptyset|i}$ depends on the solution we consider.

Consider now the WBE (2.26) and recall that SW is explicitly the determinant (2.20). Applying the same logic, we write

$$\text{SW}(C_\Lambda) = \prod_{a=1}^m Q_{a|\emptyset}^{\gg} \prod_{i=1}^n Q_{\emptyset|i}^{\gg} \text{SW}(Q^{\lessgtr}) + H, \quad (\text{D.1})$$

and then conclude using Rouché's theorem that $Q_\theta = \text{SW}(Q)$ has large zeros, *i.e.* the point $\bar{\chi}$ cannot have $\bar{\chi}_\ell$ all finite. For the argument to work, one needs to ensure that $\text{SW}(Q^{\lessgtr})$ is not vanishing but it is straightforward as the presence of twist prefactors $z_\alpha^{u/\hbar}$ ensures that already the leading- u term in $\text{SW}(Q^{\lessgtr})$ is non-vanishing.

D.2 Properness - twist-less case

To study the twist-less case, we will focus on the bosonised parameterisation of the Q-system on a Young diagram (5.10). In particular, one has

$$Q_\theta = \mathbb{Q}_{0,0} \propto W(B_1, \dots, B_m), \quad (\text{D.2})$$

where $m = h_{\Lambda^+}$. Equation (D.2) contains in principle the full information since the $\mathfrak{gl}_{m|0}$ Q-system is a possible way to parameterise the Bethe algebra \mathcal{B}_Λ .

To prove properness, we would like to use an argument similar to that of (D.1), however cancellations in the Wronskian determinant make things more subtle.

Example:

Take $B_1 = u$, $B_2 = (u - \Lambda)(u + 1)$, $B_3 = (u - \Lambda)^3$. Let $\Lambda \rightarrow \infty$ if $\chi \rightarrow \bar{\chi}$. Formally there are four divergent roots when $\Lambda \rightarrow \infty$. However $W(B_1, B_2, B_3) = u^3 + u(3\Lambda - \hbar^2) - \Lambda^2(\Lambda + 3)$ which has three divergent roots.

The issue in the example comes (at least) from the fact that in the decomposition $Q = Q^{\gg S} Q^{\lesssim S}$, $B_1^{\lesssim S}$ and $B_2^{\lesssim S}$ are polynomials of the same degree (equal to one).

To continue, we do a couple of formalisations.

Parametric factorisation Let Λ be a parameter, and we intend to consider the $\Lambda \rightarrow \infty$ behaviour of Q -functions that are algebraic functions of Λ . Define a scale function $S = \Lambda^\beta$ for some real β . We say that $Q = Q^{\gg S} Q^{\lesssim S}$ is the parametric factorisation of the polynomial Q at scale S if all roots of the monic polynomial $Q^{\gg S}$ are much larger than S , and all roots of the monic polynomial $Q^{\lesssim S}$ are comparable to or smaller than S . More precisely, for each u that satisfies $Q^{\gg S}(u) = 0$ one has $\lim_{\Lambda \rightarrow \infty} S/u = 0$, and for each u that satisfies $Q^{\lesssim S}(u) = 0$ the $\Lambda \rightarrow \infty$ limit of u/S is finite.

Then the argument around (D.1) can be formalised by the following lemma.

Lemma D.1. *For some polynomials Q_1, \dots, Q_a , let $Q_{12\dots a} = W(Q_1, \dots, Q_a)$ and let $Q = Q^{\gg S} Q^{\lesssim S}$ be the parametric factorisation at scale S .*

If degrees $\deg Q_1^{\lesssim S}$, $\deg Q_2^{\lesssim S}$, $\deg Q_3^{\lesssim S}$, \dots are pairwise distinct, and degrees $\deg Q_1$, $\deg Q_2$, $\deg Q_3$, \dots are also pairwise distinct then

$$\deg Q_{12\dots a}^{\gg S} = \sum_{a'=1}^a \deg Q_{a'}^{\gg S}. \quad (\text{D.3})$$

Proof. Perform an equivalent of decomposition (D.1) and apply Rouché's theorem. An important ingredient is that $W(u^{\deg Q_1}, \dots, u^{\deg Q_a})$ (resp. $W(u^{\deg Q_1^{\lesssim S}}, \dots, u^{\deg Q_a^{\lesssim S}})$) is not zero and hence provides the term of the highest degree of $W(Q_1, \dots, Q_a)$ (resp. $W(Q_1^{\lesssim S}, \dots, Q_a^{\lesssim S})$) which is why the restriction on the degrees is imposed. \square

Now we recall that not all coefficients of B_a bear physical information as they are subject to the symmetry transformation (2.35). We shall benefit from (2.35) to ensure that a parametric factorisation of B_a satisfies the conditions of the above lemma.

Lemma D.2. *Let B_1, \dots, B_m be monic polynomials with $\deg B_1 < \dots < \deg B_m$. For any scale S , one can find a "rotation"*

$$B_a \rightarrow B_a + \sum_{b < a} h_{ab} B_b, \quad (\text{D.4})$$

where the h_{ab} 's are complex-valued functions⁴⁶ of Λ , such that, after the rotation, the degrees of $B_1^{\lesssim S}$, $B_2^{\lesssim S}$, \dots , $B_m^{\lesssim S}$ are pairwise distinct.

We note that the proof below is constructive and it provides an algorithm to find h_{ab} explicitly.

⁴⁶More accurately, they are algebraic functions of the inhomogeneities θ_ℓ whose values depend on Λ .

Proof. Without loss of generality one can set $S = 1$ in which case we denote the parametric factorisation as $Q = Q^{\gg} Q^{\lessdot}$. Indeed, we can always perform the rescaling $u \rightarrow uS$.

In the labelling of polynomials $B_a = u^{\lambda_a + m - a} + \dots + b_k^{(a)} u^k + \dots + b_1^{(a)} u + b_0^{(a)}$, consider all $b_k^{(a)}$ that have the largest exponent when $\Lambda \rightarrow \infty$ and choose $b_{(k)}^{(a)}$ with the largest k among them. For instance, in $u^3 + \Lambda u^2 + \Lambda^3 u + 2\Lambda^3$, it is $b_{(1)} = \Lambda^3$. Then $\deg B_a^{\lessdot} = k$.

If there exist such a, b , $b < a$ that $\deg B_a^{\lessdot} = \deg B_b^{\lessdot} = k$ then perform the transformation $B_a \rightarrow B_a - \frac{b^{(a)}}{b^{(k)}} B_b$. This transformation will affect the parametric factorisation of B_a . Two things can happen. First, $\deg B_a^{\lessdot}$ becomes smaller. Second, all terms with the largest exponent are cancelled out from B_a in which case one gets a new (smaller) largest exponent and a new value for $\deg B_a^{\lessdot}$ (in principle arbitrarily large, only bounded by the degree of B_a).

We repeat recursively the procedure of comparison between all available pairs of a, b and terminate when $\deg B_a^{\lessdot}$ become pairwise distinct. The recursion will terminate in a finite number of steps and produce a meaningful result for the following reasons: there are finitely many polynomials of finite degree to operate with, the maximal exponents can only decrease in the procedure and they are bounded by zero from below, and B_a cannot vanish entirely as $\deg B_a$ are pairwise distinct and so the leading monomial is never affected by the performed transformations. \square

Example:

For $S = 1$ and the system in the Example on page 76, the rotation is done as follows. First, transformation $B_2 \rightarrow B_2 + \Lambda B_1 = u^2 + u - \Lambda$ drops the degree of B_2^{\lessdot} to zero. Now both degrees of B_2^{\lessdot} and B_3^{\lessdot} are zero. We perform transformation $B_3 \rightarrow B_3 - \Lambda^2 B_2$. This drops the maximal exponent in B_3 from three to two, and $\deg B_3^{\lessdot}$ computed with respect to the new maximal exponent is two. Now all $\deg B_a^{\lessdot}$ are pairwise distinct. In summary, we get the rotated values $B_1 = u, B_2 = u^2 + u - \Lambda, B_3 = u^3 - \Lambda(\Lambda + 3)u^2 + 2\Lambda^2 u$. B_2 has two divergent roots, B_3 has one divergent root, $W(B_1, B_2, B_3)$ remains unchanged by the performed rotation and it has three divergent roots.

Now we are ready to prove properness as declared on page 22. Assume that there is a finite point $\bar{\chi} \in \mathcal{X}$ by approaching which some coefficients of B_a diverge. We follow some path parameterised by Λ , and $\Lambda \rightarrow \infty$ corresponds to the approach of the point. Choose $S = 1$ and perform the transformation (D.4) to get $\deg B_a^{\lessdot}$ pairwise distinct. If there are still divergent coefficients after this transformation, we get $\deg Q_{\theta}^{\gg} > 0$ by Lemma D.1 and (D.2) and hence reach a contradiction. Thus all B_a have finite coefficients. Compute $\mathbb{Q}_{a,s}$ following (5.10) and use the procedure in the proof of Lemma 5.3 to compute the set C_{Λ} introduced after (2.35). c_{ℓ} appearing in this set are hence non-divergent when we approach $\bar{\chi}$ which is the properness in the sense of Section 3.

D.3 Labelling solutions with standard Young tableaux – technical details

All solutions approach (6.6)

The key step is to justify the formula (6.5). Consider the situation when $\theta_L = \Lambda$, $\Lambda \rightarrow \infty$, and all other θ_ℓ are finite. In any scaling $S = \Lambda^\beta$ with $0 < \beta < 1$, we rotate B_a to a frame where $\deg B_a^\lessgtr$ are pair-wise distinct. By Lemma D.1, there is precisely one $a = a_0$ for which precisely one root of B_{a_0} diverges, and all roots of $B_{a \neq a_0}$ stay finite.

Recall that $\deg B_a = \lambda_a + \mathbf{m} - a$ and $\deg B_i = d_i$. The assignment rules are explained in Figure 4. Since B_a are in the frame with pair-wise distinct $\deg B_a^\lessgtr$, if $a_0 \neq \mathbf{m}$, it must be that $\lambda_{a_0} > \lambda_{a_0+1}$ (equality is impossible). Then, there exists s_0 such that $d_{s_0} = \deg B_{a_0} - 1$ and so the box (a_0, s_0) is a corner box of the Young diagram.

Finally, we consider (5.10) to decide which $\mathbb{Q}_{a,s}$ have a divergent root. If $a > a_0$ then $\mathbb{Q}_{a,s}$ does not depend on B_{a_0} and hence has no divergent roots. If $s > s_0$ then all polynomials of degree from 0 to $\deg B_{a_0} - 1$ appear in the Wronskian determinant. Then the polynomial structure of B_{a_0} is irrelevant, we can replace it with the leading monomial and so $\mathbb{Q}_{a,s}$ cannot have divergent roots. Finally if $a \leq a_0$ and $s \leq s_0$ then B_{a_0} is present in the Wronskian and there is no polynomial of degree $\deg B_{a_0} - 1$ in the Wronskian, so the conditions of the Lemma D.1 are satisfied, hence $\mathbb{Q}_{a,s}$ has precisely one divergent root.

Now we can deduce that roots scale exactly as Λ . Indeed, for $\beta' \in]\beta, 1[$, the rotation of Lemma D.2 may change but the \mathbb{Q} functions are invariant under this triangular rotation. If the value of a_0 changes at scale $\Lambda^{\beta'}$ compared to scale Λ^β , then there would be another corner-box $(a'_0, s'_0) \neq (a_0, s_0)$ such that at scale $\Lambda^{\beta'}$ the \mathbb{Q} -functions with diverging roots are the nodes with $a \leq a'_0$ and $s \leq s'_0$. This is impossible because all \mathbb{Q} functions that diverge at scale $\Lambda^{\beta'}$ also diverge at scale Λ^β . Therefore a_0 is independent of $\beta \in]0, 1[$, the number of diverging roots is thus also independent of β and the diverging roots scale exactly as Λ . Finally, we get (6.5), and also that $\tilde{\mathbb{Q}}_{a,s}$ introduced alongside (6.5) is a \mathbb{Q} -system on the Young diagram with the box (a_0, s_0) removed.

Unambiguous continuation of the limiting solution (6.6) for each SYT to finite inhomogeneities

To discuss this question, we should not send inhomogeneities one after another to infinity, but do a more smooth realisation of the limit (6.4). Namely, for $\alpha_1, \alpha_2, \dots, \alpha_L \in \mathbb{C}^*$ and $\beta_L > \beta_{L-1} > \dots > \beta_1 > 0$, we parameterise $\theta_1 = \alpha_1 \Lambda^{\beta_1}$, $\theta_2 = \alpha_2 \Lambda^{\beta_2}$, \dots , $\theta_L = \alpha_L \Lambda^{\beta_L}$. All roots and all coefficients of all \mathbb{Q} -polynomials are algebraic functions⁴⁷ of the parameter Λ and hence have a large- Λ behavior of the form $\alpha \Lambda^\beta$ which allows applying scaling argumentation from previous sections.

If β_ℓ are spaced apart well, we can recover the same results as if inhomogeneities are sent to infinity one by one. But now, after we know that all solutions of the \mathbb{Q} -system approach (6.6), a sharper judgement about possible β_ℓ can be made by observing that the

⁴⁷For a \mathbb{Q} -system on a Young diagram, this is an immediate consequence of the QQ-relations. For a \mathbb{Q} -system on a Hasse diagram, we should restrict symmetry transformations (2.35) to algebraically depend on θ_ℓ .

leading order of the large Λ expansion (6.6) is solved by ⁴⁸

$$\forall a \leq m, \quad B_a \sim u^{m-a} \prod_{s=1}^{\lambda_a} \left(u - (m - a + s) N_{a-1, s-1}^{(\mathcal{T}_{a,s})} \theta_{\mathcal{T}_{a,s}} \right). \quad (\text{D.5})$$

Let us now *parameterise* the Q-system by $\kappa_1, \dots, \kappa_L$ as follows

$$\forall a \leq m, \quad B_a = u^{m-a} \prod_{s=1}^{\lambda_a} \left(u - (m - a + s) N_{a-1, s-1}^{(\mathcal{T}_{a,s})} \kappa_{\mathcal{T}_{a,s}} \Lambda^{\beta_{\mathcal{T}_{a,s}}} \right). \quad (\text{D.6})$$

Recall that $\theta_\ell = \alpha_\ell \Lambda^{\beta_\ell}$. Then $Q_\theta = W(B_1, \dots, B_m)$ realises a map from κ_ℓ to α_ℓ which analytically depends on $1/\Lambda$ for β_ℓ being integers. When $1/\Lambda = 0$ this map is simply an identity map with obviously non-zero Jacobian. Hence we can apply the analytic implicit function theorem to invert the map. By the theorem, for some neighbourhood of the point $1/\Lambda = 0$, κ_ℓ are analytic functions of $\alpha_1, \dots, \alpha_L$ and $1/\Lambda$ and hence each limiting solution (6.6) can be continued to finite values of inhomogeneities.

References

- [1] H. Bethe, “Zur Theorie der Metalle,” *Z. Phys.* **71** (1931) 205–226.
- [2] B. Sutherland, “A General Model for Multicomponent Quantum Systems,” *Phys. Rev. B* **12** (1975) 3795–3805.
- [3] P. Kulish and N. Reshetikhin, “Generalized Heisenberg ferromagnet and the Gross-Neveu model,” *Sov. Phys. JETP* **53** (1981) 108–114.
- [4] P. Kulish, “Integrable graded magnets,” *J. Sov. Math.* **35** (1986) 2648–2662.
- [5] E. Ogievetsky and P. Wiegmann, “Factorized S Matrix and the Bethe Ansatz for Simple Lie Groups,” *Phys. Lett. B* **168** (1986) 360–366.
- [6] Z. Tsuboi, “Analytic Bethe Ansatz And Functional Equations Associated With Any Simple Root Systems Of The Lie Superalgebra $sl(r + 1|s + 1)$,” *Physica A* **252** (1998) 565–585, 0911.5387.
- [7] E. Ragoucy and G. Satta, “Analytical Bethe Ansatz for closed and open $\mathfrak{gl}(M|N)$ super-spin chains in arbitrary representations and for any Dynkin diagrams,” *JHEP* **09** (2007) 001, 0706.3327.
- [8] E. Mukhin, V. Tarasov, and A. Varchenko, “Bethe algebra of homogeneous XXX Heisenberg model has simple spectrum,” *Communications in Mathematical Physics* **288** (May, 2009) 1–42, 0706.0688.
- [9] E. Mukhin, V. Tarasov, and A. Varchenko, “Spaces of quasi-exponentials and representations of the Yangian $Y(\mathfrak{gl}_N)$,” *ArXiv e-prints* (Mar., 2013) 1303.1578.
- [10] C. Huang, K. Lu, and E. Mukhin, “Solutions of $\mathfrak{gl}_{m|n}$ XXX Bethe ansatz equation and rational difference operators,” *Journal of Physics A: Mathematical and Theoretical* **52** (Aug, 2019) 375204, 1811.11225.

⁴⁸The computation reduces to the fact that $W(1, u, \dots, u^{k-1}, u^k(u - (k + 1)\alpha\theta)) \propto (u - \alpha\theta)$.

- [11] K. Lu and E. Mukhin, “On the supersymmetric XXX spin chains associated to $\mathfrak{gl}_{1|1}$,” 1910.13360.
- [12] A. N. Kirillov, “Completeness of states of the generalized Heisenberg magnet,” *J. Sov. Math.* **36** (1987), no. 1 115–128.
- [13] A. Foerster and M. Karowski, “Algebraic properties of the Bethe ansatz for an $\text{spl}(2,1)$ supersymmetric t-J model,” *Nucl. Phys. B* **396** (1993) 611–638.
- [14] K. Schoutens, “Complete solution of a supersymmetric extended Hubbard model,” *Nucl. Phys. B* **413** (1994) 675–688, [cond-mat/9310006](#).
- [15] S. V. Kerov, A. N. Kirillov, and N. Y. Reshetikhin, “Combinatorics, Bethe Ansatz, and representations of the symmetric group,” *J. Sov. Math.* **41** (1988), no. 2 916–924.
- [16] A. N. Kirillov and N. Y. Reshetikhin, “The Bethe Ansatz and the combinatorics of Young tableaux,” *J. Sov. Math.* **41** (1988), no. 2 925–955.
- [17] A. Vladimirov, “Non-string two-magnon configurations in the isotropic Heisenberg magnet,” *Physics Letters A* **105** (1984), no. 8 418 – 420.
- [18] K. Isler and M. B. Paranjape, “Violations of the string hypothesis in the solutions of the Bethe ansatz equations in the XXX Heisenberg model,” *Phys. Lett. B* **319** (1993) 209–214, [hep-th/9304078](#).
- [19] A. Kuniba, T. Nakanishi, and Z. Tsuboi, “The Bethe equation at $q=0$, the Mobius inversion formula, and weight multiplicities. 3. The $X(N)^{**}(r)$ case,” *Lett. Math. Phys.* **59** (2002) 19–31, [math/0105146](#).
- [20] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Z. Tsuboi, “Paths, crystals and fermionic formulae,” *Prog. Math. Phys.* **23** (2002) 205–272, [math/0102113](#).
- [21] L. V. Avdeev and A. A. Vladimirov, “Exceptional solutions to the Bethe ansatz equations,” *Theor. Math. Phys.* **69** (nov, 1986) 1071–1079.
- [22] D. Volin, “String hypothesis for $\mathfrak{gl}(n|m)$ spin chains: a particle/hole democracy,” *Lett. Math. Phys.* **102** (2012) 1–29, [1012.3454](#).
- [23] W. Hao, R. I. Nepomechie, and A. J. Sommese, “Singular solutions, repeated roots and completeness for higher-spin chains,” *J. Stat. Mech.* **1403** (2014) P03024, [1312.2982](#).
- [24] R. I. Nepomechie and C. Wang, “Algebraic Bethe ansatz for singular solutions,” *J. Phys. A* **46** (2013) 325002, [1304.7978](#).
- [25] C. Marboe and D. Volin, “Unpublished.”
- [26] R. J. Baxter, “Completeness of the Bethe ansatz for the six and eight vertex models,” *J. Stat. Phys.* **108** (2002) 1–48, [0111188](#).
- [27] R. Baxter, “Generalized ferroelectric model on a square lattice,” *Stud. Appl. Math.* **50** (1971) 51–69.
- [28] N. Reshetikhin, “The functional equation method in the theory of exactly soluble quantum systems,” *ZhETF* **84** (1983) 1190–1201.
- [29] M. Gaudin, *La fonction d’onde de Bethe*. Masson, 1983.
- [30] L. Faddeev, “How algebraic Bethe ansatz works for integrable model,” in *Relativistic gravitation and gravitational radiation. Proceedings, School of Physics, Les Houches, France, September 26-October 6, 1995*, pp. pp. 149–219, 5, 1996. [hep-th/9605187](#).

- [31] Z. Tsuboi, “Analytic Bethe ansatz and functional equations for Lie superalgebra $sl(r+1|s+1)$,” *J. Phys.* **A30** (1997) 7975–7991, 0911.5386.
- [32] A. Arnaudon, N. Crampe, A. Doikou, L. Frappat, and E. Ragoucy, “Analytical Bethe Ansatz for closed and open $gl(n)$ -spin chains in any representation,” *J. Stat. Mech.* **0502** (2005) P02007, [math-ph/0411021](#).
- [33] E. Granet and J. L. Jacobsen, “On zero-remainder conditions in the Bethe ansatz,” *JHEP* **03** (2020) 178, 1910.07797.
- [34] N. Gromov, F. Levkovich-Maslyuk, and G. Sizov, “New Construction of Eigenstates and Separation of Variables for $SU(N)$ Quantum Spin Chains,” *JHEP* **09** (2017) 111, 1610.08032.
- [35] A. Liashyk and N. A. Slavnov, “On Bethe vectors in gl_3 -invariant integrable models,” *JHEP* **06** (2018) 018, 1803.07628.
- [36] P. Ryan and D. Volin, “Separated variables and wave functions for rational $gl(N)$ spin chains in the companion twist frame,” *J. Math. Phys.* **60** (2019), no. 3 032701, 1810.10996.
- [37] I. Krichever, O. Lipan, P. Wiegmann, and A. Zabrodin, “Quantum integrable systems and elliptic solutions of classical discrete nonlinear equations,” *Commun. Math. Phys.* **188** (1997) 267–304, [hep-th/9604080](#).
- [38] V. V. Bazhanov, S. L. Lukyanov, and A. B. Zamolodchikov, “Integrable structure of conformal field theory. 3. The Yang-Baxter relation,” *Commun. Math. Phys.* **200** (1999) 297–324, [hep-th/9805008](#).
- [39] G. P. Pronko and Yu. G. Stroganov, “Bethe equations ‘on the wrong side of equator’,” *J. Phys.* **A32** (1999) 2333–2340, [hep-th/9808153](#).
- [40] Z. Tsuboi, “Solutions of the T-system and Baxter equations for supersymmetric spin chains,” *Nucl. Phys.* **B826** (2010) 399–455, 0906.2039.
- [41] V. Kazakov, S. Leurent, and D. Volin, “T-system on T-hook: Grassmannian Solution and Twisted Quantum Spectral Curve,” *JHEP* **12** (2016) 044, 1510.02100.
- [42] R. J. Baxter, “Partition function of the eight vertex lattice model,” *Ann. Phys.* **70** (1972) 193–228.
- [43] R. J. Baxter, *Exactly solved models in statistical mechanics*. 1982.
- [44] A. N. Kirillov and N. Y. Reshetikhin, “The Yangians, Bethe Ansatz and combinatorics,” *Lett. Math. Phys.* **12** (oct, 1986) 199–208.
- [45] M. Nazarov and G. Olshanski, “Bethe subalgebras in twisted Yangians,” *Communications in Mathematical Physics* **178** (May, 1996) 483–506, [arXiv:q-alg/9507003](#).
- [46] M. L. Nazarov, “Quantum Berezinian and the classical Capelli identity,” *Letters in Mathematical Physics* **21** (1991) 123–131.
- [47] L. Gow, “Yangians of lie superalgebras,” *PhD thesis* (2007).
- [48] J. Maillet, G. Niccoli, and L. Vignoli, “Separation of variables bases for integrable $gl_{M|N}$ and Hubbard models,” *SciPost Phys.* **9** (2020) 060, 1907.08124.
- [49] V. Kac, “Lie superalgebras,” *Advances in Mathematics* **26** (1977), no. 1 8 – 96.
- [50] A. N. Sergeev, “The tensor algebra of the identity representation as a module over the Lie superalgebras $gl(n|m)$ and $Q(n)$,” *Math. USSR-Sbornik* **51** (feb, 1985) 419–427.

- [51] A. Berele and A. Regev, “Hook young diagrams with applications to combinatorics and to representations of Lie superalgebras,” *Advances in Mathematics* **64** (1987), no. 2 118 – 175.
- [52] A. Molev, *Yangians and classical Lie algebras*. No. 143. American Mathematical Soc., 2007.
- [53] P. P. Kulish and N. Y. Reshetikhin, “GL3-invariant solutions of the Yang-Baxter equation and associated quantum systems,” *Journal of Soviet Mathematics* **34** (1986), no. 5 1948–1971.
- [54] A. Zabrodin, “Discrete Hirota’s equation in quantum integrable models,” *Int. J. Mod. Phys. B* **11** (1997) 3125, [hep-th/9610039](#).
- [55] I. V. Cherednik, “An analogue of the character formula for Hecke algebras,” *Functional Analysis and Its Applications* **21** (1987), no. 2 172–174.
- [56] V. Bazhanov and N. Reshetikhin, “Restricted Solid on Solid Models Connected With Simply Based Algebras and Conformal Field Theory,” *J. Phys. A* **23** (1990) 1477.
- [57] V. Kazakov and P. Vieira, “From characters to quantum (super)spin chains via fusion,” *JHEP* **10** (2008) 050, [0711.2470](#).
- [58] A. I. Molev and E. Ragoucy, “The MacMahon Master Theorem for right quantum superalgebras and higher Sugawara operators for $\widehat{\mathfrak{gl}}(m|n)$,” *Moscow Mathematical Journal* **14** (2014) 83–119, [0911.3447](#).
- [59] D. Talalaev, “Quantization of the Gaudin system,” [hep-th/0404153](#).
- [60] A. Chervov and G. Falqui, “Manin matrices and Talalaev’s formula,” *Journal of Physics A: Mathematical and Theoretical* **41** (Apr, 2008) 194006.
- [61] L. Gow, “Gauss Decomposition of the Yangian $Y(\mathfrak{gl}(m|n))$,” *Communications in Mathematical Physics* **276** (Oct, 2007) 799–825, [math/0605219](#).
- [62] A. Belitsky, S. E. Derkachov, G. Korchemsky, and A. Manashov, “Baxter Q-operator for graded $SL(2|1)$ spin chain,” *J. Stat. Mech.* **0701** (2007) P01005, [hep-th/0610332](#).
- [63] V. V. Bazhanov and Z. Tsuboi, “Baxter’s Q-operators for supersymmetric spin chains,” *Nucl. Phys. B* **805** (2008) 451–516, [0805.4274](#).
- [64] V. Kazakov, S. Leurent, and Z. Tsuboi, “Baxter’s Q-operators and operatorial Backlund flow for quantum (super)-spin chains,” *Commun. Math. Phys.* **311** (2012) 787–814, [1010.4022](#).
- [65] R. Frassek, T. Lukowski, C. Meneghelli, and M. Staudacher, “Oscillator Construction of $\mathfrak{su}(n|m)$ Q-Operators,” *Nucl. Phys. B* **850** (2011) 175–198, [1012.6021](#).
- [66] Z. Tsuboi, “A note on q -oscillator realizations of $U_q(\mathfrak{gl}(M|N))$ for Baxter Q-operators,” *Nucl. Phys. B* **947** (2019) 114747, [1907.07868](#).
- [67] S. E. Derkachov and A. N. Manashov, “Factorization of R-matrix and Baxter Q-operators for generic $\mathfrak{sl}(N)$ spin chains,” *J. Phys. A* **42** (2009) 075204, [0809.2050](#).
- [68] V. V. Bazhanov, T. Lukowski, C. Meneghelli, and M. Staudacher, “A Shortcut to the Q-Operator,” *J. Stat. Mech.* **1011** (2010) P11002, [1005.3261](#).
- [69] V. V. Bazhanov, R. Frassek, T. Lukowski, C. Meneghelli, and M. Staudacher, “Baxter Q-Operators and Representations of Yangians,” *Nucl. Phys. B* **850** (2011) 148–174, [1010.3699](#).

- [70] E. Frenkel and D. Hernandez, “Baxter’s relations and spectra of quantum integrable models,” *Duke Math. J.* **164** (2015), no. 12 2407–2460, 1308.3444.
- [71] Z. Tsuboi, “Wronskian solutions of the T, Q and Y-systems related to infinite dimensional unitarizable modules of the general linear superalgebra $gl(M|N)$,” *Nucl. Phys.* **B870** (2013) 92–137, 1109.5524.
- [72] N. Gromov, V. Kazakov, S. Leurent, and Z. Tsuboi, “Wronskian Solution for AdS/CFT Y-system,” *JHEP* **01** (2011) 155, 1010.2720.
- [73] N. Gromov, V. Kazakov, S. Leurent, and D. Volin, “Quantum spectral curve for arbitrary state/operator in AdS₅/CFT₄,” *JHEP* **09** (2015) 187, 1405.4857.
- [74] M. Raynaud and A. Grothendieck, *Revêtements Étales et Groupe Fondamental: Séminaire de Géométrie Algébrique du Bois Marie 1960/61 (SGA 1)*. Lecture Notes in Mathematics. Springer Berlin Heidelberg, 2006. Proposition 3.2.
- [75] A. Grothendieck, “Éléments de géométrie algébrique : IV. Étude locale des schémas et des morphismes de schémas, Quatrième partie,” *Publications Mathématiques de l’IHÉS* **32** (1967) 182. Corollaire 18.12.4.
- [76] H. Matsumura and M. Reid, *Commutative Ring Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1989. Theorem 23.1 and Corollary.
- [77] R. Hartshorne, *Algebraic Geometry*. Graduate Texts in Mathematics. Springer New York, 2013. Exercise 10.9.
- [78] P. Cohn, *Further Algebra and Applications*. Springer London, 2002. Corollary 4.6.4.
- [79] D. Quillen, “Projective modules over polynomial rings,” *Inventiones Mathematicae* **36** (Dec, 1976) 167–171.
- [80] C. Marboe and D. Volin, “Fast analytic solver of rational Bethe equations,” *J. Phys.* **A50** (2017), no. 20 204002, 1608.06504.
- [81] C. Marboe and D. Volin, “The full spectrum of AdS₅/CFT₄ I: Representation theory and one-loop Q-system,” *J. Phys.* **A51** (2018), no. 16 165401, 1701.03704.
- [82] C. Marboe and D. Volin, “The full spectrum of AdS₅/CFT₄ II: Weak coupling expansion via the quantum spectral curve,” 1812.09238.
- [83] J. Lykke Jacobsen, Y. Jiang, and Y. Zhang, “Torus partition function of the six-vertex model from algebraic geometry,” *JHEP* **03** (2019) 152, 1812.00447.
- [84] Z. Bajnok, E. Granet, J. L. Jacobsen, and R. I. Nepomechie, “On Generalized Q-systems,” *JHEP* **03** (2020) 177, 1910.07805.
- [85] V. Kazakov, A. S. Sorin, and A. Zabrodin, “Supersymmetric Bethe ansatz and Baxter equations from discrete Hirota dynamics,” *Nucl. Phys.* **B790** (2008) 345–413, hep-th/0703147.
- [86] L. Frappat, P. Sorba, and A. Sciarrino, “Dictionary on Lie superalgebras,” hep-th/9607161.
- [87] F. Woynarovich, “Low-energy excited states in a hubbard chain with on-site attraction,” *Journal of Physics C: Solid State Physics* **16** (dec, 1983) 6593–6604.
- [88] P.-A. Bares, J. M. P. Carmelo, J. Ferrer, and P. Horsch, “Charge-spin recombination in the one-dimensional supersymmetric t-j model,” *Phys. Rev. B* **46** (Dec, 1992) 14624–14654.

- [89] E. Mukhin and A. Varchenko, “Solutions to the XXX type Bethe ansatz equations and flag varieties,” *Central European Journal of Mathematics* **1** (nov, 2003) 238–271, 0211321.
- [90] N. Gromov and P. Vieira, “Complete 1-loop test of AdS/CFT,” *JHEP* **04** (2008) 046, 0709.3487.
- [91] E. K. Sklyanin, “Separation of variables in the quantum integrable models related to the Yangian $Y[\mathfrak{sl}(3)]$,” *J. Math. Sci.* **80** (1996) 1861–1871, hep-th/9212076. [Zap. Nauchn. Semin.205,166(1993)].
- [92] A. Kuniba and J. Suzuki, “Analytic Bethe Ansatz for fundamental representations of Yangians,” *Commun. Math. Phys.* **173** (1995) 225–264, hep-th/9406180.
- [93] N. Gromov, F. Levkovich-Maslyuk, P. Ryan, and D. Volin, “Dual Separated Variables and Scalar Products,” *Phys. Lett. B* **806** (2020) 135494, 1910.13442.
- [94] A. Ilin and L. Rybnikov, “Degeneration of Bethe subalgebras in the Yangian of \mathfrak{gl}_n ,” *Letters in Mathematical Physics* (Dec, 2017) 1703.04147.
- [95] I. Mashanova-Golikova, “Simplicity of spectra for Bethe subalgebras in $Y(\mathfrak{gl}_2)$,” 1906.09049.
- [96] M. Günaydin and D. Volin, “The Complete Unitary Dual of Non-compact Lie Superalgebra $\mathfrak{su}(p, q|m)$ via the Generalised Oscillator Formalism, and Non-compact Young Diagrams,” *Commun. Math. Phys.* **367** (2019), no. 3 873–939, 1712.01811.
- [97] N. Gromov, V. Kazakov, S. Leurent, and D. Volin, “Quantum Spectral Curve for Planar $\mathcal{N} = 4$ Super-Yang-Mills Theory,” *Phys. Rev. Lett.* **112** (2014), no. 1 011602, 1305.1939.
- [98] W. Hao, R. I. Nepomechie, and A. J. Sommese, “Completeness of solutions of Bethe’s equations,” *Phys. Rev. E* **88** (2013), no. 5 052113, 1308.4645.
- [99] A. W. Sandvik, “Computational Studies of Quantum Spin Systems,” *AIP Conf. Proc.* **1297** (2010), no. 1 135, 1101.3281.
- [100] T. Bargheer, N. Beisert, and N. Gromov, “Quantum Stability for the Heisenberg Ferromagnet,” *New J. Phys.* **10** (2008) 103023, 0804.0324.
- [101] J.-S. Caux and J.-M. Maillet, “Computation of dynamical correlation functions of Heisenberg chains in a field,” *Phys. Rev. Lett.* **95** (2005) 077201, cond-mat/0502365.
- [102] M. Gaudin, “Diagonalisation d’une classe d’hamiltoniens de spin,” *Journal de Physique* **37** (1976), no. 10 1087–1098.
- [103] E. Mukhin, V. Tarasov, and A. Varchenko, “Schubert calculus and representations of the general linear group,” *Journal of the American Mathematical Society* **22** (Apr, 2009) 909–940, 0711.4079.
- [104] E. Mukhin, B. Vicedo, and C. Young, “Gaudin models for $\mathfrak{gl}(m|n)$,” *Journal of Mathematical Physics* **56** (May, 2015) 051704, 1404.3526.
- [105] C. Huang and E. Mukhin, “The duality of $\mathfrak{gl}(m|n)$ and $\mathfrak{gl}(k)$ Gaudin models,” *Journal of Algebra* **548** (2020) 1 – 24.
- [106] J. Maillet and G. Niccoli, “On quantum separation of variables,” *J. Math. Phys.* **59** (2018), no. 9 091417, 1807.11572.
- [107] J. Maillet and G. Niccoli, “Complete spectrum of quantum integrable lattice models associated to $Y(\mathfrak{gl}(n))$ by separation of variables,” *SciPost Phys.* **6** (2019), no. 6 071, 1810.11885.

- [108] J. Maillet and G. Niccoli, “On quantum separation of variables beyond fundamental representations,” 1903.06618.
- [109] P. Ryan and D. Volin, “Separation of variables for rational $\mathfrak{gl}(n)$ spin chains in any compact representation, via fusion, embedding morphism and Backlund flow,” 2002.12341.
- [110] J. Maillet, G. Niccoli, and L. Vignoli, “On Scalar Products in Higher Rank Quantum Separation of Variables,” 2003.04281.
- [111] D. Maulik and A. Okounkov, “Quantum groups and quantum cohomology,” 2012.
- [112] N. A. Nekrasov and S. L. Shatashvili, “Supersymmetric vacua and Bethe ansatz,” *Nuclear Physics B - Proceedings Supplements* **192-193** (Jul, 2009) 91–112, 0901.4744.
- [113] N. A. Nekrasov and S. L. Shatashvili, “Quantum integrability and supersymmetric vacua,” *Prog. Theor. Phys. Suppl.* **177** (2009) 105–119, 0901.4748.
- [114] V. Gorbounov, R. Rimányi, V. Tarasov, and A. Varchenko, “Quantum cohomology of the cotangent bundle of a flag variety as a Yangian Bethe algebra,” *Journal of Geometry and Physics* **74** (Dec, 2013) 56–86.
- [115] P. P. Pushkar, A. Smirnov, and A. M. Zeitlin, “Baxter Q-operator from quantum K-theory,” *Adv. Math.* **360** (2020) 106919, 1612.08723.
- [116] N. Nekrasov, “Superspin chains and supersymmetric gauge theories,” *JHEP* **03** (2019) 102, 1811.04278.
- [117] T. Becker, H. Kredel, and V. Weispfenning, *Gröbner Bases: A Computational Approach to Commutative Algebra*. Graduate Texts in Mathematics. Springer New York, 1998.
- [118] J.-P. Serre, “Faisceaux algébriques cohérents,” *Annals of Mathematics* **61** (1955), no. 2 197–278. §50.
- [119] H. Zhang, “RTT realization of quantum affine superalgebras and tensor products,” *Int. Math. Res. Not.* **2016** (2016), no. 4 1126–1157, 1407.7001.
- [120] A. N. Kirillov and N. Y. Reshetikhin, “Representations of Yangians and multiplicities of occurrence of the irreducible components of the tensor product of representations of simple Lie algebras,” *J. Sov. Math.* **52** (nov, 1990) 3156–3164.
- [121] R.-b. Zhang, “Representations of superYangian,” *J. Math. Phys.* **36** (1995) 3854–3865, hep-th/9411243.
- [122] R.-b. Zhang, “The $\mathfrak{gl}(m|n)$ superYangian and its finite dimensional representations,” *Lett. Math. Phys.* **37** (1996) 419–434, q-alg/9507029.
- [123] V. G. Drinfel’d, “Degenerate affine Hecke algebras and Yangians,” *Functional Analysis and Its Applications* **20** (1986), no. 1 58–60.
- [124] T. Arakawa, “Drinfeld functor and finite-dimensional representations of Yangian,” *Communications in Mathematical Physics* **205** (Aug, 1999) 1–18, math/9807144.
- [125] A. Davydov and A. Molev, “A categorical approach to classical and quantum Schur-Weyl duality,” *Groups, Algebras and Applications* (2011) 143–171.
- [126] Y. Z. Flicker, “Affine quantum super Schur-Weyl duality,” *Algebras and Representation Theory* **23** (Dec, 2018) 135–167.
- [127] I. Macdonald, *Symmetric Functions and Hall Polynomials*. Oxford classic texts in the physical sciences. Clarendon Press, 1998.

- [128] J. Désarménien, B. Leclerc, and J.-Y. Thibon, “Hall-Littlewood functions and Kostka-Foulkes polynomials in representation theory,” *Séminaire Lotharingien de Combinatoire [electronic only]* **32** (1994) 38 p.–38 p.
- [129] V. Chari and S. Loktev, “Weyl, Demazure and fusion modules for the current algebra of \mathfrak{sl}_{r+1} ,” *Advances in Mathematics* **207** (2006), no. 2 928 – 960, [math/0502165](#).