Riemann-Hilbert method and N-soliton solutions for the mixed Chen-Lee-Liu derivative nonlinear Schrödinger equation

Fang Fang, Beibei Hu[∗] , Ling Zhang

School of Mathematics and Finance, Chuzhou University, Anhui 239000 China

Abstract

In this paper, we aim to investigate the mixed Chen-Lee-Liu derivative nonlinear Schrödinger(CLL-NLS) equation via the Riemann-Hilbert(RH) method. we construct a RH problem base on the Jost solution of the Lax pair. By solving this RH problem corresponding to the non reflection case, the N-soliton solution of CLL-NLS equation is obtained, which expression is the ratio of $(2N + 1) \times (2N + 1)$ determinant and $2N \times 2N$ determinant.

Keywords:

Riemann-Hilbert method; Chen-Lee-Liu derivative nonlinear Schrödinger equation; soliton solutions; boundary conditions

1. Introduction

Soliton theory is an important branch of nonlinear science. In physics, solitons are used to describe solitary waves with elastic scattering characteristics. In mathematics, soliton theory provides a series of methods for solving nonlinear partial differential equations, which attracts the attention of mathematical physicists. With the development of soliton theory, many methods to solve soliton equations with important physical background have been proposed, for example, inverse scattering transform (IST) [\[1,](#page-9-0) [2](#page-9-1), [3\]](#page-9-2), Darboux transformation(DT) [\[4\]](#page-9-3), Bäcklund transformation [\[5\]](#page-9-4), Hirota bilinear method [\[6,](#page-9-5) [7,](#page-9-6) [8\]](#page-9-7), Wroskian technique [\[9,](#page-9-8) [10,](#page-9-9) [11\]](#page-9-10) and so on.

The derivative nonlinear schrödinger(DNLS) equation describing Alfvén waves in magnetic field is as follows [\[12\]](#page-9-11):

$$
i u_t = u_{xx} + i(|u|^2 u)_x = 0,
$$
\n(1.1)

which is one of the most significant equations in physics. Kaup and Newell obtained the solution of Eq.[\(1.1\)](#page-0-0) by using IST method[\[13](#page-9-12)]. The IST method has obvious advantages in solving the initial value of the soliton equation, but its calculation is large. Fortunately, on the basis of IST method, Riemann-Hilbert(RH) method, a relatively simple and direct method for solving soliton equation, was proposed by Novikovet et al.[\[14\]](#page-9-13). This method is similar to the IST method, the RH method first considers the direct scattering problem, that is, the RH problem is constructed, from the initial data of the soliton equation, the scattering data at the initial time is obtained, and the scattering data at any time is obtained by using its time evolution law. Finally, the exact solution is established by using the IST method[\[15\]](#page-9-14). Since the RH method was proposed, many solutions of soliton equations have been discussed, such as,

[∗]Corresponding author. Email: hu chzu@163.com; hu chzu@shu.edu.cn

Preprint submitted to Elsevier April 21, 2020

the coupled DNLS equation [\[16](#page-9-15)], the coupled higher-order NLS equation [\[17](#page-9-16)], the short pulse(SP) equation [\[18](#page-9-17)], the coupled modified Korteweg-de Vries (mKdV) equation [\[19\]](#page-9-18), the generalized Sasa-Satsuma equation [\[20](#page-10-0)], the twocomponent Gerdjikov-Ivanov(GI) equation [\[21](#page-10-1)], the modified SP equation [\[22\]](#page-10-2), and so on [\[23](#page-10-3), [24](#page-10-4), [25](#page-10-5)]. In particular, RH method is an effective way to working the initial-boundary value problems(IBVPs) of the integrable systems [\[26,](#page-10-6) [27,](#page-10-7) [28,](#page-10-8) [29,](#page-10-9) [30,](#page-10-10) [31\]](#page-10-11).

Recently, Chan et al.[\[32](#page-10-12)] reported the the mixed Chen-Lee-Liu derivative nonlinear Schödinger(CLL-NLS) equation as follows

$$
ir_t + r_{xx} + |r|^2 r - i|r|^2 r_x = 0,
$$
\n(1.2)

which is a completely integrable model, and a large number of solutions of Eq.[\(1.2\)](#page-1-0) are discussed, such as, the soliton solution by Hirota bilinear method [\[33\]](#page-10-13), the higher-order soliton, breathers, and rogue wave solutions by DT method[\[34\]](#page-10-14). In particular, the IBVPs of Eq.[\(1.2\)](#page-1-0) to be investigated by Fokas method[\[35\]](#page-10-15).

The design structure of this paper is as follows. Section 2, we will construct a basic RH problem based on the Jost solution of Lax pair. Section 3, we will give the reconstruction of potential function and the law of scattering data evolution with time. Section 4, the formula of N-soliton solution expressed by determinant ratio is proposed. Section 5 is the conclusions.

2. The Riemann-Hilbert problem

In this section, we shall construct a RH problem for the CLL-NLS equation, by using IST method. First of all, we introduce the coupled CLL-NLS equation as following:

$$
\begin{cases}\nr_t - ir_{xx} + ir^2q + rqr_x = 0, \\
q_t + iq_{xx} - iq^2r + qrq_x = 0.\n\end{cases}
$$
\n(2.1)

which reduces to the CLL-NLS equation while $q = -r^*$ and the $*$ denotes complex conjugation. These two equations in [\(2.1\)](#page-1-1) are the compatibility condition of the following Lax pair[\[34,](#page-10-14) [35,](#page-10-15) [36,](#page-10-16) [37\]](#page-10-17)

$$
\Phi_x = U\Phi = \left(i(\lambda^2 - \frac{1}{2})\sigma + \lambda Q + \frac{1}{4}iQ^2\sigma\right)\Phi,
$$
\n
$$
\Phi_t = V\Phi = [-2i(\lambda^2 - \frac{1}{2})^2\sigma - 2\lambda^3Q - i\lambda^2Q^2\sigma + \lambda\left(Q + i\sigma Q_x - \frac{1}{2}Q^3\right) - \frac{1}{8}iQ^4\sigma + \frac{1}{4}(QQ_x - Q_xQ)]\Phi,
$$
\n(2.2b)

with

$$
\sigma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, Q = \begin{pmatrix} 0 & r \\ -r^* & 0 \end{pmatrix}.
$$
 (2.3)

where $\Phi = \Phi(x, t; \lambda)$ is a matrix function of the complex spectral parameter, λ is the spectral parameter. *Q* is named potential function.

For the sake of convenience, we introduce a new matrix function $J = J(x, t; \lambda)$ defined by

$$
\Phi = Je^{i(\lambda^2 - \frac{1}{2})\sigma x - 2i(\lambda^2 - \frac{1}{2})^2 \sigma t}
$$
\n(2.4)

Obviously, we can check that $Je^{i(\lambda^2-\frac{1}{2})\sigma_x-2i(\lambda^2-\frac{1}{2})^2\sigma t}$ satisfies Eqs.[\(2.2a\)](#page-1-2),[\(2.2b\)](#page-1-3). Inserting [\(2.4\)](#page-1-4) into [\(2.2a\)](#page-1-2)-(2.2b), the form of the Lax pair [\(2.2a\)](#page-1-2)-[\(2.2b\)](#page-1-3) becomes

$$
J_x = i(\lambda^2 - \frac{1}{2})[\sigma, J] + U_1 J,\tag{2.5a}
$$

$$
J_t = -2i(\lambda^2 - \frac{1}{2})^2[\sigma, J] + V_1 J.
$$
 (2.5b)

where $[\sigma, J] = \sigma J - J\sigma$ is the commutator.

$$
U_1 = \lambda Q + \frac{1}{4}iQ^2\sigma
$$
\n
$$
V_1 = -2\lambda^3 Q - i\lambda^2 Q^2\sigma + \lambda \left(Q + i\sigma Q_x - \frac{1}{2}Q^3\right) - \frac{1}{8}iQ^4\sigma + \frac{1}{4}(QQ_x - Q_xQ)
$$

As usual, in the following scattering process, we only concentrate on the *x*-part of the Lax pair [\(2.2a\)](#page-1-2). Indeed the *x*-part of the Lax pair [\(2.2a\)](#page-1-2) allows us to take use of the existing symmetry relations of the potential Q. Consequently, we shall treat the time t as a dummy variable and omit it. Now, we calculate two Jost solutions $J_{\pm} = J_{\pm}(x, \lambda)$ of Eq.[\(2.5a\)](#page-2-0) for $\lambda \in \mathbb{R}$. Based on the properties of the Jost solutions $J_{\pm} = J_{\pm}(x, \lambda)$

$$
J_{+} = ([J_{+}]_{1}, [J_{+}]_{2}), \tag{2.7}
$$

$$
J_{-} = ([J_{-}]_1, [J_{-}]_2), \tag{2.8}
$$

with the boundary conditions

$$
J_+ \to I, \ x \to -\infty,\tag{2.9a}
$$

$$
J_- \to I, \ x \to +\infty. \tag{2.9b}
$$

the subscripts in $J(x, \lambda)$ represent which end of the *x*-axis the boundary conditions are set. Where $[J_{\pm}]_n$ $(n = 1, 2)$ denote the *n*-th column vector of J_+ , $I = diag\{1, 1\}$ is the 2×2 unit matrix. According to the method of variation of parameters as well as the boundary conditions, we can turn Eq.[\(2.5a\)](#page-2-0) for $\lambda \in \mathbb{R}$ into the Volterra integral equations.

$$
J_{+}(x,\lambda) = I - \int_{x}^{+\infty} e^{i(\lambda^2 - \frac{1}{2})\hat{\sigma}(x-\xi)} U_1 J_{+}(\xi,\lambda) d\xi,
$$
 (2.10)

$$
J_{-}(x,\lambda) = I + \int_{-\infty}^{x} e^{i(\lambda^2 - \frac{1}{2})\hat{\sigma}(x-\xi)} U_1 J_{-}(\xi,\lambda) d\xi,
$$
 (2.11)

where $\hat{\sigma}$ represents a matrix operator acting on 2 × 2 matrix *X* by $\hat{\sigma}X = [\sigma, X]$ and by $e^{x\hat{\sigma}}X = e^{x\sigma}Xe^{-x\sigma}$. Moreover, after simple analysis, we find that $[J_+]_1$, $[J_-]_2$ are analytic for $\lambda \in D_+$ and continuous for $\lambda \in D_+ \cup \mathbb{R} \cup i\mathbb{R}$. Similarly $[J_1], [J_1]_2$ are analytic for $\lambda \in D_-$ and continuous for $\lambda \in D_- \cup \mathbb{R} \cup i\mathbb{R}$, here

$$
D_{+} = \left\{ \lambda | arg \lambda \in \left(0, \frac{\pi}{2}\right) \bigcup \left(\pi, \frac{3\pi}{2}\right) \right\}
$$

$$
D_{-} = \left\{ \lambda | arg \lambda \in \left(\frac{\pi}{2}, \pi\right) \bigcup \left(\frac{3\pi}{2}, 2\pi\right) \right\}.
$$

Secondly, let us investigate the properties of J_{\pm} . We deduce the determinants of J_{\pm} are constants for all *x* on the basis of the Abel's identity and $Tr(U_1) = 0$. Furthermore, due to the boundary conditions Eq.[\(2.9a\)](#page-2-1),[\(2.9b\)](#page-2-2), we have

$$
\det J_{\pm} = 1, \quad \lambda \in \mathbb{R} \bigcup i\mathbb{R}.
$$
 (2.13)

By introducing a new function $E(x, \lambda) = e^{i(\lambda^2 - \frac{1}{2})\sigma x}$, we find that spectral problem Eq.[\(2.5a\)](#page-2-0) exists two fundamental matrix solutions J_+E and J_-E , which are linearly related by a 2 × 2 scattering matrix $S(\lambda)$

$$
J_{-}E = J_{+}E \cdot S(\lambda), \quad \lambda \in \mathbb{R} \bigcup i\mathbb{R}.
$$
 (2.14)

From Eq. (2.13) and (2.14) , we know that

$$
\det S(\lambda) = 1. \tag{2.15}
$$

Furthermore, let *x* go to + ∞ , the 2 × 2 scattering matrix *S*(λ) is given as

$$
S(\lambda) = \lim_{x \to +\infty} E^{-1} J_{-} E = I + \int_{-\infty}^{+\infty} E^{-1} U_{1} J_{-} E d\xi, \ \lambda \in \mathbb{R} \bigcup i \mathbb{R}.
$$

From the analytic property of *J*−, we find that *s*²² can be analytically extended to *D*+, *s*¹¹ allows analytic extensions to *D*_−. Generally speaking, s_{12} , s_{21} can only be defined in the $\mathbb{R} \cup i\mathbb{R}$.

In what follows, we shall construct a RH problem for the CLL-NLS equation. Firstly, using the analytic properties of J_{\pm} , we define a new Jost solution $P_1 = P_1(x, \lambda)$ as

$$
P_1 = ([J_+]_1, [J_-]_2). \tag{2.16}
$$

which is obviously analytic for $\lambda \in D_+$. In addition, from Eqs. [\(2.10\)](#page-2-4) and [\(2.11\)](#page-2-5), we have

$$
P_1 \to I, \qquad \lambda \to +\infty, \ \lambda \in D_+.
$$
 (2.17)

Next, we introduce the limit of *P*¹

$$
P^+ = \lim_{\lambda \to \Gamma} P_1 \qquad \qquad \Gamma = \mathbb{R} \bigcup i\mathbb{R}.\tag{2.18}
$$

From [\(2.14\)](#page-3-0) and [\(2.16\)](#page-3-1), we can get

$$
P^{+} = J_{+} \left(\begin{array}{cc} 1 & e^{i(2\lambda^{2}-1)x} s_{12} \\ 0 & s_{22} \end{array} \right). \tag{2.19}
$$

To obtain the analytic counterpart of P^+ in D^- , denoted by P_2 , we consider the inverse matrices J_{\pm}^{-1} defined as

$$
J_{+}^{-1} = \left(\begin{array}{c} [J_{+}^{-1}]_{1} \\ [J_{+}^{-1}]_{2} \end{array}\right), \quad J_{-}^{-1} = \left(\begin{array}{c} [J_{-}^{-1}]_{1} \\ [J_{-}^{-1}]_{2} \end{array}\right), \tag{2.20}
$$

here $[J_{\pm}^{-1}]_n$ ($n = 1, 2$) denote the *n*-th row vector of J_{\pm}^{-1} . Then we can see that $[J_{+}^{-1}]_1$, $[J_{-}^{-1}]_2$ are analytic for $\lambda \in D_-$ and continuous for $\lambda \in D_- \cup \mathbb{R} \cup i\mathbb{R}$, whereas $[J_-^{-1}]_1$, $[J_+^{-1}]_2$ are analytic for $\lambda \in D_+$ and continuous for $\lambda \in D_+ \cup \mathbb{R} \cup i\mathbb{R}$. Obviously, J_{\pm}^{-1} satisfy the adjoint scattering equation of Eq.[\(2.5a\)](#page-2-0):

$$
K_x = i(\lambda^2 - \frac{1}{2})[\sigma, K] - K U_1.
$$
 (2.21)

In addition, it is not difficult to find that the inverse matrices J_+^{-1} and J_-^{-1} satisfy the following boundary conditions.

$$
J_+^{-1} \to I, \ x \to -\infty,\tag{2.22a}
$$

$$
J_{-}^{-1} \to I, \ x \to +\infty. \tag{2.22b}
$$

Taking the similar procedure as above, a matrix function *P*² which is analytic in *D*[−]

$$
P_2 = \left(\begin{array}{c} [J_+^{-1}]_1 \\ [J_-^{-1}]_2 \end{array}\right). \tag{2.23}
$$

and *P* −

$$
P^- = \lim_{\lambda \to \Gamma} P_2 \qquad \qquad \Gamma = \mathbb{R} \bigcup i\mathbb{R}.\tag{2.24}
$$

are expressed. Moreover, we can get that

$$
P_2 \to I, \qquad \lambda \to -\infty, \lambda \in D_-.
$$
 (2.25)

and

$$
P^{-} = \begin{pmatrix} 1 & 0 \\ e^{-i(2\lambda - 1)x} r_{21} & r_{22} \end{pmatrix} J_{+}^{-1}.
$$
 (2.26)

with $R(\lambda) \equiv (r_{kj})_{2 \times 2} = S^{-1}(\lambda)$. Similar to the scattering coefficients s_{ij} above, it is easy to know that r_{11} allows an analytic extension to D_+ and r_{22} is analytically extendible to D_- . Generally speaking, r_{12} , r_{21} can only be defined in the $\mathbb{R} \cup i\mathbb{R}$. In addition, from [\(2.14\)](#page-3-0), we get

$$
E^{-1}J_{-}^{-1} = R(\lambda) \cdot E^{-1}J_{+}^{-1}, \qquad \lambda \in \mathbb{R} \bigcup i\mathbb{R}.
$$
 (2.27)

Summarizing the above the results, we find that two matrix functions P^+ and P^- which are analytic in D_+ and D_- , respectively, are related by

$$
P^{-}(x,\lambda)P^{+}(x,\lambda) = \begin{pmatrix} 1 & s_{12}e^{i(2\lambda-1)x} \\ r_{21}e^{-i(2\lambda-1)x} & 1 \end{pmatrix}, \lambda \in \mathbb{R} \bigcup i\mathbb{R}.
$$
 (2.28)

Eq.[\(2.28\)](#page-4-0) is just the RH problem for the CLL-NLS equation. In order to obtain solution for this RH problem, we assume that the RH problem is non-regular when $\det P_1$ and $\det P_2$ can be zero for $\lambda_k \in D_+$, and $\lambda_k \in D_-$, respectively. $1 \leq k \leq N$. where *N* is the number of these zeros. Recalling the definitions of P_1 and P_2 , we can see that

$$
\det P_1(x,\lambda) = s_{22}(\lambda), \qquad \lambda \in D_+, \tag{2.29}
$$

$$
\det P_2(x,\lambda) = r_{22}(\lambda), \qquad \lambda \in D_-.
$$
 (2.30)

To specify these zeros, we first can take use of a symmetry relation for *U*¹

$$
U_1^{\dagger} = \sigma U_1 \sigma,
$$

where the superscript \dagger means the Hermitian of a matrix. Therefore from Eq. [\(2.21\)](#page-3-2), we arrive at

$$
J_{\pm}^{\dagger}(x,t,\lambda^*) = \sigma J_{\pm}^{-1}(x,t,\lambda)\sigma,
$$
\n(2.31)

Then from [\(2.14\)](#page-3-0), we also gain

$$
S^{\dagger}(\lambda^*) = \sigma S^{-1}(\lambda)\sigma,\tag{2.32}
$$

which implies the following relations

$$
r_{11}(\lambda) = s_{11}^*(\lambda^*) \qquad \lambda \in D_+, \tag{2.33}
$$

$$
r_{22}(\lambda) = s_{22}^*(\lambda^*) \qquad \lambda \in D_-, \tag{2.34}
$$

$$
r_{12}(\lambda) = -s_{21}^*(\lambda^*) \qquad \lambda \in \mathbb{R} \bigcup i\mathbb{R},\tag{2.35}
$$

 \mathbf{I} \mathbf{I}

$$
r_{21}(\lambda) = -s_{12}^*(\lambda^*) \qquad \lambda \in \mathbb{R} \bigcup i\mathbb{R}.\tag{2.36}
$$

Moreover from Eq. [\(2.32\)](#page-4-1) and the definitions of P_1 , P_2 , we point out that the analytic solutions P_1 , P_2 satisfy the involution property

$$
P_1^{\dagger}(x,\lambda^*) = \sigma P_2(x,\lambda)\sigma, \qquad \lambda \in D_- \tag{2.37}
$$

The similar analysis shows that the potential matrix *Q* also satisfies another symmetry relations

 $Q = -\sigma Q \sigma$

It follows that

$$
J_{\pm}(-\lambda) = \sigma J_{\pm}(\lambda)\sigma. \tag{2.38}
$$

and

$$
P_1(-\lambda) = \sigma P_1(\lambda)\sigma. \tag{2.39}
$$

thus

$$
s_{11}(-\lambda) = -s_{11}(\lambda) \qquad \lambda \in D_{-},
$$
\n(2.40)

$$
s_{22}(-\lambda) = -s_{22}(\lambda) \qquad \lambda \in D_+, \tag{2.41}
$$

$$
s_{12}(-\lambda) = -s_{12}(\lambda) \qquad \lambda \in \mathbb{R} \bigcup i\mathbb{R}, \tag{2.42}
$$

$$
s_{21}(-\lambda) = -s_{21}(\lambda) \qquad \lambda \in \mathbb{R} \left(\begin{array}{c} \end{array} \right) i\mathbb{R}.
$$
 (2.43)

Therefore, from [\(2.34\)](#page-4-2), we find that if λ_j is a zero of det P_1 , then $\hat{\lambda}_j = \lambda_j^*$ is a zero of det P_2 . Moreover, in view of [\(2.41\)](#page-5-0), we know that $-\lambda$ is also a zero of det P_1 . Hence we suppose that det P_1 has 2*N* simple zeros { λ_j }^{2*N*}</sup> satisfying $\lambda_{N+l} = -\lambda_l$ (1 ≤ *l* ≤ *N*), which all lie in *D*₊. From the zeros of det*P*₁, we see that det*P*₂ possesses 2*N* simple zeros ${\lbrace \hat{\lambda}_j \rbrace}_1^{2N}$ satisfying $\hat{\lambda}_j = \lambda_j^*$, (1 ≤ *j* ≤ 2*N*), which are all in *D*−. Obviously the zeros of det*P*₁ and det*P*₂ always appear in quadruples. To solve the RH problem, we need the scattering data including the continuous scattering data {*s*₁₂, *s*₂₁} and the discrete scattering data $\{\lambda_j, \hat{\lambda}_j, v_j, \hat{v}_j\}$ which a single column vector v_j and row vector \hat{v}_j satisfying

$$
P_1(\lambda_j)v_j = 0, \quad \hat{v}_j P_2(\hat{\lambda}_j) = 0.
$$
\n(2.44)

On the one hand, by taking the Hermitian conjugate of $P_1(\lambda_i)v_i = 0$, we can construct the relationship between each pair of v_j and \hat{v}_j .

$$
v_j = \sigma v_{j-N} \qquad N+1 \le j \le 2N,
$$

$$
\hat{v}_j = v_j^{\dagger} \sigma \qquad 1 \le j \le N.
$$
 (2.45)

On the other hand, in order to obtain the spatial evolutions for vectors $v_i(x)$, taking the *x*-derivative to equation $P_1v_j = 0$ and using [\(2.5a\)](#page-2-0), we obtain

$$
v_j = e^{i(\lambda_j^2 - \frac{1}{2})\sigma x} v_{j_0} \qquad \quad 1 \le j \le N,\tag{2.46}
$$

where $v_{j_0} = v_j |_{x} = 0$.

Using these vectors, the RH problem which corresponds to the reflection-less case, that is to say, we set the vanishing cofficient $s_{12} = s_{21} = 0$ in the RH problem [\(2.28\)](#page-4-0), can be solved exactly, and the result is

$$
P_1(\lambda) = I - \sum_{k=1}^{2N} \sum_{j=1}^{2N} \frac{\nu_k (M^{-1})_{kj} \hat{\nu}_j}{\lambda - \hat{\lambda}_j},
$$
\n(2.47a)

$$
P_2(\lambda) = I + \sum_{k=1}^{2N} \sum_{j=1}^{2N} \frac{\nu_k (M^{-1})_{kj} \hat{\nu}_j}{\lambda - \lambda_k}.
$$
 (2.47b)

where $M = (m_{ki})_{2N \times 2N}$ is a matrix whose entries are

$$
m_{kj} = \frac{\hat{v}_k v_j}{\lambda_j - \hat{\lambda}_k}, \ 1 \le k, \ j \le 2N. \tag{2.48}
$$

3. Inverse scattering transform

In this section, with the help of P_1 in [\(2.47a\)](#page-6-0), we can write out explicitly the potential *Q*. Owing to $P_1(\lambda)$ is the solution of spectral problem [\(2.5a\)](#page-2-0), we assume that the asymptotic expansion of $P_1(\lambda)$ at large λ as

$$
P_1 = I + \frac{P_1^{(1)}}{\lambda} + \frac{P_1^{(2)}}{\lambda^2} + O(\lambda^{-3}) \quad \lambda \to \infty,
$$
\n(3.1)

Then by substituting the above expansion into $(2.5a)$ and comparing $O(\lambda)$ terms, we obtains

$$
Q = -i[\sigma, P_1^{(1)}] = \begin{pmatrix} 0 & -2i(P_1^{(1)})_{12} \\ 2i(P_1^{(1)})_{21} & 0 \end{pmatrix}.
$$
 (3.2)

which implies that *r* can be reconstructed as

$$
r = -2i(P_1^{(1)})_{12}.\tag{3.3}
$$

where $P_1^{(1)}$ $\binom{1}{1} = (P_1^{(1)}$ $\binom{1}{1}$ _{2×2} and $\binom{P_1^{(1)}}{1}$ $1^{(1)}$ ₁)_{*i*j} is the (*i*; *j*)-entry of $P_1^{(1)}$ $1_1^{(1)}$, *i*, *j* = 1, 2. Here, the matrix function $P_1^{(1)}$ $_1^{(1)}$ can be found from [\(2.47a\)](#page-6-0)

$$
P_1^{(1)} = \sum_{k=1}^{2N} \sum_{j=1}^{2N} v_k (M^{-1})_{kj} \hat{v}_j.
$$
 (3.4)

4. The soliton solutions

To derive the solutions for the CLL-NLS equation, we also need the scattering data at time *t*, which need investigate the time evolution of scattering data. In fact, by using [\(2.5b\)](#page-2-6) and [\(2.14\)](#page-3-0), making the limit $x \to +\infty$, and taking into account the boundary condition [\(2.9a\)](#page-2-1) for *J*₊ as well as $V_1 \rightarrow 0$ as $x \rightarrow \pm \infty$, we arrive at

$$
s_{11}|_t = s_{22}|_t = 0, \quad s_{12}|_t = -4i(\lambda^2 - \frac{1}{2})^2 s_{12}, \quad s_{21}|_t = -4i(\lambda^2 - \frac{1}{2})^2 s_{21},
$$

\n
$$
\frac{d\lambda_j}{dt} = 0, \quad v_j|_t = -2i(\lambda^2 - \frac{1}{2})^2 v_j.
$$
\n(4.1)

Combining [\(2.45\)](#page-5-1) with [\(2.46\)](#page-5-2), we can derive the column vectors v_j and the row vector \hat{v}_j explicitly,

$$
v_j = \begin{cases} e^{\theta_j \sigma} v_{j_0}, & 1 \le j \le N \\ \sigma e^{\theta_{j-N}\sigma} v_{j-N,0}, & N+1 \le j \le 2N \end{cases}
$$
(4.2)

and

$$
\hat{\nu}_j = \begin{cases}\n\nu_{j_0}^{\dagger} e^{\theta_j^* \sigma} \sigma & 1 \le j \le N \\
\nu_{j - N, 0}^{\dagger} e^{\theta_{j - N}^* \sigma}, & N + 1 \le j \le 2N\n\end{cases}
$$
\n(4.3)

where $\theta_j = i(\lambda_j^2 - \frac{1}{2})x - 2i(\lambda_j^2 - \frac{1}{2})^2t$ ($\lambda_j \in D+$), v_{j_0} is a constant vector.

we have chosen $v_{j_0} = (c_j, 1)^T$, it follows from [\(3.4\)](#page-6-1) that the N-soliton solutions for the CLL-NLS equation reads

$$
r = 2i \sum_{k=1}^{N} \sum_{j=1}^{N} c_k e^{\theta_k - \theta_j^*} (M^{-1})_{kj} + 2i \sum_{k=N+1}^{2N} \sum_{j=1}^{N} c_{k-N} e^{\theta_{k-N} - \theta_j^*} (M^{-1})_{kj}
$$

$$
-2i \sum_{k=1}^{N} \sum_{j=N+1}^{2N} c_k e^{\theta_k - \theta_{j-N}^*} (M^{-1})_{kj} - 2i \sum_{k=N+1}^{2N} \sum_{j=N+1}^{2N} c_{k-N} e^{\theta_{k-N} - \theta_{j-N}^*} (M^{-1})_{kj}.
$$
 (4.4)

and $M = (m_{kj})_{2N \times 2N}$ is given by

$$
m_{kj} = \begin{cases} \frac{c_j c_k^* e^{\theta_k^* + \theta_j} - e^{-(\theta_k^* + \theta_j)}}{\lambda_k^* - \lambda_j}, & 1 \le k, j \le N, \\ \frac{c_{j-N} c_k^* e^{\theta_k^* + \theta_{j-N}} + e^{-(\theta_k^* + \theta_{j-N})}}{\lambda_k^* - \lambda_j}, & 1 \le k \le N, N+1 \le j \le 2N, \\ \frac{c_j c_{k-N}^* e^{\theta_k^* - N + \theta_j} + e^{-(\theta_k^* - N + \theta_j)}}{-\lambda_k^* - \lambda_j}, & 1 \le j \le N, N+1 \le k \le 2N, \\ \frac{c_{j-N} c_{k-N}^* e^{\theta_k^* - N + \theta_{j-N}} - e^{-(\theta_k^* - N + \theta_{j-N})}}{-\lambda_{k-N}^* + \lambda_{j-N}}, & N+1 \le k, j \le 2N. \end{cases} (4.5)
$$

with $\theta_j = i(\lambda_j^2 - \frac{1}{2})x - 2i(\lambda_j^2 - \frac{1}{2})^2t$ $(\lambda_j \in D+)$

The simplest situation occurs when $N = 1$ in formula [\(4.4\)](#page-7-0). the single-soliton solution is

$$
r = -\frac{2ic_1e^{\theta_1 - \theta_1^*}(m_{11} + m_{21} - m_{12} - m_{22})}{det M}
$$
(4.6)

where $M = (m_{kj})_{2 \times 2}$ is given by

$$
m_{11} = \frac{|c_1| \left(e^{\theta_1^* + \theta_1 + \xi_1} - e^{-\theta_1^* - \theta_1 - \xi_1}\right)}{\lambda_1^* - \lambda_1}, m_{12} = \frac{|c_1| \left(e^{\theta_1^* + \theta_1 + \xi_1} + e^{-\theta_1^* - \theta_1 - \xi_1}\right)}{\lambda_1^* + \lambda_1},
$$

\n
$$
m_{21} = \frac{|c_1| \left(e^{\theta_1^* + \theta_1 + \xi_1} + e^{-\theta_1^* - \theta_1 - \xi_1}\right)}{-\lambda_1^* - \lambda_1}, m_{22} = \frac{|c_1| \left(e^{\theta_1^* + \theta_1 + \xi_1} - e^{-\theta_1^* - \theta_1 - \xi_1}\right)}{-\lambda_1^* + \lambda_1}.
$$

Letting $\lambda_1 = \lambda_{11} + i\lambda_{12}$, $|c_1| = e^{\xi_1}$, then the single-soliton solution can be written as

$$
r = \frac{4\lambda_{11}\lambda_{12}e^{iY}(\lambda_{11}sinh X + i\lambda_{12}cosh X)}{\lambda_{12}^{2}cosh^{2} X + \lambda_{11}^{2}sinh^{2} X}.
$$
 (4.7)

with

$$
X = 8\lambda_{11}\lambda_{12}(\lambda_{11}^2 - \lambda_{12}^2)t - 2\lambda_{11}\lambda_{12}x - 4\lambda_{11}\lambda_{12}t + ln|c_1|,
$$

$$
Y = 2(\lambda_{11}^2 - \lambda_{12}^2)x - 4(\lambda_{11}^4 - 6\lambda_{11}^2\lambda_{12}^2 + \lambda_{12}^4 - \lambda_{11}^2 - \lambda_{12}^2)t.
$$

In order to understand the properties of the resulting soliton solution, we can take the model

$$
|r| = \frac{4|\lambda_{11}\lambda_{12}|}{\sqrt{\lambda_{12}^2 \cosh^2 X + \lambda_{11}^2 \sinh^2 X}}.
$$
\n(4.8)

Now we investigate the case for $N = 2$ in formula [\(4.4\)](#page-7-0). we arrive at the two-soliton solution as follows:

$$
r = -2i \sum_{k=1}^{2} \sum_{j=1}^{2} c_k e^{\theta_k - \theta_j^*} (M^{-1})_{kj} - 2i \sum_{k=3}^{4} \sum_{j=1}^{2} c_{k-2} e^{\theta_{k-2} - \theta_j^*} (M^{-1})_{kj}
$$

+2i $\sum_{k=1}^{2} \sum_{j=3}^{4} c_k e^{\theta_k - \theta_{j-2}^*} (M^{-1})_{kj} + 2i \sum_{k=3}^{4} \sum_{j=3}^{4} c_{k-2} e^{\theta_{k-2} - \theta_{j-2}^*} (M^{-1})_{kj}.$ (4.9)

and $M = (m_{kj})_{4\times4}$ is given by

$$
m_{11} = \frac{c_1 c_1^* e^{\theta_1^* + \theta_1} - e^{-\theta_1^* - \theta_1}}{\lambda_1^* - \lambda_1}, m_{12} = \frac{c_1 c_2^* e^{\theta_1^* + \theta_2} - e^{-\theta_1^* - \theta_2}}{\lambda_1^* - \lambda_2},
$$
\n
$$
m_{13} = \frac{c_1 c_1^* e^{\theta_1^* + \theta_1} + e^{-\theta_1^* - \theta_1}}{\lambda_1^* + \lambda_1}, m_{14} = \frac{c_1^* c_2 e^{\theta_1^* + \theta_2} + e^{-\theta_1^* - \theta_2}}{\lambda_1^* + \lambda_2},
$$
\n
$$
m_{21} = \frac{c_1 c_2^* e^{\theta_2^* + \theta_1} - e^{-\theta_2^* - \theta_1}}{\lambda_2^* - \lambda_1}, m_{22} = \frac{c_2 c_2^* e^{\theta_2^* + \theta_2} - e^{-\theta_2^* - \theta_2}}{\lambda_2^* - \lambda_2},
$$
\n
$$
m_{23} = \frac{c_1 c_2^* e^{\theta_2^* + \theta_1} + e^{-\theta_2^* - \theta_1}}{\lambda_2^* + \lambda_1}, m_{24} = \frac{c_2 c_2^* e^{\theta_2^* + \theta_2} + e^{-\theta_2^* - \theta_2}}{\lambda_2^* + \lambda_2},
$$
\n
$$
m_{31} = \frac{c_1 c_1^* e^{\theta_1^* + \theta_1} + e^{-\theta_1^* - \theta_1}}{-\lambda_1^* - \lambda_1}, m_{32} = \frac{c_2 c_1^* e^{\theta_1^* + \theta_2} + e^{-\theta_1^* - \theta_2}}{-\lambda_1^* - \lambda_2},
$$
\n
$$
m_{33} = \frac{c_1 c_1^* e^{\theta_1^* + \theta_1} - e^{-\theta_1^* - \theta_1}}{-\lambda_1^* + \lambda_1}, m_{34} = \frac{c_2 c_1^* e^{\theta_1^* + \theta_2} - e^{-\theta_1^* - \theta_2}}{-\lambda_2^* - \lambda_2},
$$
\n<math display="block</math>

Further, we can also write the obtained N-soliton solutions [\(4.4\)](#page-7-0) into the form of the determinant ratio

$$
r = 2i \frac{det \widetilde{M}_{12}}{det M}.
$$
\n(4.11)

where \widetilde{M}_{12} is

$$
\widetilde{M}_{12} = \begin{pmatrix} 0 & f_1 \\ \hat{g}_2 & M \end{pmatrix} . \tag{4.12}
$$

where the matrix M is defined by Eq.[\(4.5\)](#page-7-1), and $f_1 = (v_{11}, v_{21} \cdots v_{2N,1}), \hat{g}_2 = (\hat{v}_{11}, \hat{v}_{21} \cdots \hat{v}_{2N,1})^T$

5. Conclusions

In this work, by applying RH method, we establish the N-soliton solutions for the CLL-NLS equation. First of all, the Lax pair of the coupled CLL-NLS equation are transformed to obtain the corresponding Jost solution. Then we study spectrum analysis and construct the particular RH problem, which is non-regular case. Then we derive the scattering data including the continuous scattering data { s_{12} , s_{21} } and the discrete scattering data { λ_j , λ_j , v_j , \hat{v}_j }. With aid of reconstructing the potential, we solve the N-soliton solutions for the CLL-NLS. Finally we obtain a simple and compact N-soliton solutions formula.

Acknowledgements

This work is supported by the National Natural Science Foundation of China under the grant No. 11601055, Natural Science Foundation of Anhui Province under the grant No. 1408085QA06, Natural Science Research Projects of Anhui Province under Grant Nos. KJ2019A0637 and gxyq2019096.

References

References

- [1] Gardner, CS, Green, JM, Kruskal, MD, Miüra, RM. Method for solving the Korteweg-de Vries equation. *Phys. Rev. Lett.* 1967; 19: 1095-1097.
- [2] Constantin, A, Ivanov, RI, Lenells, J. Inverse scattering transform for the Degasperis-Procesi equation. *Nonlinearity*. 2010; 23: 2559-2575.
- [3] Ji, JL, Zhu, ZN. Soliton solutions of an integrable nonlocal modified Korteweg-de Vries equation through inverse scattering transform. *J. Math. Anal. Appl.* 2017; 453: 973-984.
- [4] Matveev, VB, Salle, MA. Darboux transformations and solitons, (*Springer-Verlag, Berlin.* 1991)
- [5] Olver, PJ. Application of Lie group to differential equation, (*Springer-Verlag, New York.* 1986)
- [6] Hirota, R. Exact solution of the Korteweg-de Vries equation for multiple collisions of solitons. *Phys. Rev. Lett.* 1971; 27: 1192-1194.
- [7] Ma, WX, Yong, XL, Zhang, HQ. Diversity of interaction solutions to the (2+1)-dimensional Ito equation. *Comput. Math. Appl.* 2018; 75: 289-295.
- [8] Han, Z, Chen, Y, Chen, JC. General N-dark soliton solutions of the multi-component Mel'nikov system.*J. Phys. Soc. Japan.* 2017; 86: 074005.
- [9] Lou, SY. A note on the new similarity reductions of the Boussinesq equation. *Phys. Lett. A* 1990; 151: 133-135.
- [10] Li, JB, Liu, ZR. Smooth and non-smooth traveling waves in a nonlinearly dispersive equation. *Appl. Math. Model.* 2000; 25: 41-56.
- [11] Fan, EG. Uniformly constructing a series of explicit exact solutions to nonlinear equations in mathematical physics. *Chaos Solitons Fract.* 2003; 16: 819-839.
- [12] Mio, K, Ogino, T, Minami, K, Takeda, S. Modified nonlinear Schrödinger equation for Alfvén waves propagating along the magnetic field in cold plasmas. *J. Phys. Soc. Japan.* 1976; 41: 265-271.
- [13] Kaup, D, Newell, AC. An exact solution for a derivative nonlinear Schrödinger equation. *J. Math. Phys.* 1978; 19: 798-801.
- [14] Novikov, S, Manakov, S, Pitaevskii, L, Zakharov, V. Theory of solitons: the inverse scattering method, (*New York and London*, *Consultants Bureau.* 1984)
- [15] Shchesnovich, VS, Yang, JK. General soliton matrices in the Riemann-Hilbert problem for integrable nonlinear equation. *J. Math. Phys.* 2003; 44: 4604-4639.
- [16] Guo, B, Ling, L. Riemann-Hilbert approach and N-soliton formula for coupled derivative Schrödinger equation. *J. Math. Phys.* 2012; 53: 073506.
- [17] Wang, DS, Yin, SJ, Liu, YF. Integrability and bright soliton solutions to the coupled nonlinear Schrödinger equation with higher-order effects. *Appl. Math. Comput.* 2014; 229: 296-309.
- [18] De Monvel, AB, Shepelsky, D, Zielinski, L. The short pulse equation by a Riemann-Hilbert approach. Arxiv ID: 1608.02249.
- [19] Wu, JP, Geng, XG. Inverse scattering transform and soliton classification of the coupled modified Korteweg-de Vries equation. *Commun. Nonlinear Sci. Numer. Simulat.* 2017; 53: 83-93.
- [20] Geng, XG, Wu, JP. Riemann-Hilbert approach and N-soliton solutions for a generalized Sasa-Satsuma equation. *Wave Motion*. 2016; 60: 62-72.
- [21] Zhang, YS, Cheng, Y, He, JS. Riemann-Hilbert method and N-soliton for two-component Gerdjikov-Ivanov equation. *J. Nonlinear Math. Phy.* 2017; 24: 210-223.
- [22] Guo, BL, Liu, N. A Riemann-Hilbert approach for the modified short pulse equation. *Appl. Anal.* 2018; 1: 1-14.
- [23] Ma, WX. Riemann-Hilbert problems and N-soliton solutions for a coupled mKdV system. *J. Geom. Phys.* 2018; 132: 45-54.
- [24] Hu, J, Xu, J, Yu, GF. Riemann-Hilbert approach and N-soliton formula for a higher-order Chen-Lee-Liu equation. *J. Nonlinear Math. Phys.* 2018; 25: 633-649.
- [25] Ai, LP, Xu, J. On a Riemann-Hilbert problem for the Fokas-Lenells equation. *Appl. Math. Lett.* 2019; 87: 57-63.
- [26] Zhang, N, Xia, TC, Hu, BB. A Riemann-Hilbert approach to the complex Sharma-Tasso-Olver equation on the half line. *Commun. Theor. Phys.* 2017; 68: 580-594.
- [27] Xia, BQ, Fokas, AS. Initial-boundary value problems associated with the Ablowitz-Ladik system. *Physica D.* 2018; 364: 27-61.
- [28] Hu, BB, Xia, TC, Zhang, N, Wang, JB. Initial-boundary value problems for the coupled higher-order nonlinear Schrödinger equations on the half-line. *Int. J. Nonlin. Sci. Num.* 2018; 19: 83-92.
- [29] Hu, BB, Xia, TC, Ma, WX. Riemann-Hilbert approach for an initial-boundary value problem of the two-component modified Korteweg-de Vries equation on the half-line. *Appl. Math. Comput.* 2018; 332: 148-159.
- [30] Hu, BB, Xia, TC, Ma, WX. The Riemann-Hilbert approach to initial-boundary value problems for integrable coherently coupled nonlinear schrödinger systems on the half-line. *E. Asian J. Appl. Math.* 2018; 8: 531-548.
- [31] Manakov, SV. On the theory of two-dimensional stationary self-focusing of electromagenic waves. *Sov. Phys. JETP* 1974; 38: 248-253.
- [32] Kundu, A. Landau-Lifshitz and higher order nonlinear systems gauge generated from nonlinear Schrödinger type equations. *J. Math. Phys.* 1984; 25: 3433-3438.
- [33] Chan, HN, Chow, KW, Kedziora, DJ, Grimshaw, RHJ. Rogue wave modes for a derivative nonlinear Schrödinger model. *Phys. Rev. E* 2014; 89: 032914.
- [34] Zhang, YS, Guo, LJ, Amin, C, He, JS. Higher-order rogue wave dynamics for derivative nonlinear Schrödinger equation. [arXiv:1409.7923.](http://arxiv.org/abs/1409.7923)
- [35] Hu, BB, Zhang, L, Zhang, N. On the Riemann-Hilbert problem for the Chen-Lee-Liu derivative nonlinear Schrödinger equation, [arXiv:2004.07608.](http://arxiv.org/abs/2004.07608)
- [36] Vijayajayanthi, M, Kanna, T, Lakshmanan, M. Bright-dark solitons and their collisions in mixed N-coupled nonlinear Schrödinger equations. *Phys. Rev. A* 2008; 77 : 013820.
- [37] Clarkson, PA, Cosgrove, CM. Painlevé analysis of the non-linear Schrödinger family of equations. *J. Phys. A: Math. Gen.* 1987; 20: 2003-2024.