The linearization of the non-Hermitian Lindblad equation and its applicaton

Jian $Xu^{1,*}$ and Youneng $Guo^{2,3,\dagger}$

¹College of Electronics and Information Engineering,

Guangdong Ocean University, Zhanjiang, 524088, China

²Interdisciplinary Center for Quantum Information,

National University of Defense Technology, Changsha 410073, People's Republic of China

³School of Electronic Information and Electrical Engineering,

Changsha University, Changsha, Hunan 410022, People's Republic of China

(Dated: October 25, 2021)

Non-Hermiticity and decoherence are two different effects of the open systems. Each of them has triggered many interesting phenomena. In this work, we rewrite the non-Hermitian Lindblad master equation into a linear one, under which the eigenvalues and eigenmatrices are easy to be obtained. Based on this, we theoretically explore the system with non-Hermiticity and decoherence. As an example, beyond the Fermi arc, we fouce on the cases of just parts of eigenvalues met the conditions(Re[λ]=0 or Im[λ]=0). We find that there are new-style steady states when Im[λ]=0 and exceptionla points(EPs) in the present system and discuss the topological properties of eigenstates when Re[λ]=0, and find that normal EPs is a special case of this conclusion. In short, our work brings new insight to the fields of open systems.

PACS numbers: 03.65.Yz, 42.50.p, 03.65.Vf

I. INTRODUCTION

Non-Hermiticity and decoherence are the two different effects and produce different properties, even though both of them are used to describe the open systems where systems of interest interact with their environments. One of interesting properties of the non-Hermiticity is that the systems in i-Fermi arc have entirely real energy spectra [1, 2], e.g., the Hamiltonians with parity-time (PT) symmetry or pseudo-Hermitian. It allows one to realize the intriguing properties of non-Hermiticity in the steady systems [3–8]. Compared to the rigorous conditions of pseudo-Hermitian or PT-symmetry, a new-type non-Hermitian system with one eigenvalue being real and the other being complex has been proposed [9]. The other interesting properties of the non-Hermiticity is the presence of EPs [10, 11], at which two or more complex eigenvalues coincide and the corresponding eigenstates coalesce. The EPs have many intriguing properties [12– 19] that have no counterparts in Hermitian systems.

Different form the non-Hermiticity induced by the interaction of systems with their environments, another dynamics of an open quantum system is characterized by the presence of decoherence terms, describing the loss of energy, coherence, and information into the environments. As we all known, the time evolution of an open quantum system weakly interacting with a Markovian environment, under general hypotheses, can be described by the Lindblad master equation, composed of a Hermitian Hamiltonian part, and a Lindblad dissipative term [20–22]. Meanwhile, the numerical calculations of the Lindblad master equation have been broad applied in the open systems [23–25]. Recently, some works rewrite some Lindblad terms of the equation as a effect non-Hermitian term and theoretically analysis the propesties of Lindblad master equation [26–28]. An other method [29–31] suggested to rewrite the Lindblad equation to a linear equation (a matrix multiplying a vector, if considering discrete systems), whose eigenvalue and eigenstate are completely theoretically solvable [32–34].

In this work, we focus on a system with non-Hermiticity and decoherence. We address this system via the linearization method, whose the eigenvalues and eigenmatrices are easy to be obtained. Byound the condition of i Fermi arc with $\text{Im}[\lambda]=0$, we theoretically propose a general condition of steady states, which the imaginary part of one of the eigenvalues is zero and the other are negative, and find that there are new-style steady states without PT-symmetry or pseudo-Hermiticity in the present system. In addition, we demonstrate that there are topological properties of the eigenstates when the real part of eigenvalue being zero. Our results can apply to EPs and Fermi arc with $\operatorname{Re}[\lambda]=0$. We further demonstrate the non-Hermitian Liouvillian exceptional points (NLEPs) in this system with different properties compared to normal EPs or Liouvillian exceptional points(LEPs). Moreover, we verify the novel properties of this system by numerically solving the Lindblad master equation.

The contents of this paper are as follows. In section II, we introduce the theoretical framework, mapping the non-Hermitian Lindblad master equation into a linear form, and point out the condition of the steady-state. In section III, we consider a two-level open system with non-Hermiticity and decoherence, and reveal the novel properties of this present system. Discussions and conclusions are given in section VI.

^{*} xujian_328@163.com

[†] guoxuyan2007@163.com

II. LINDBLAD EQUATION FOR A GENERAL OPEN SYSTEM

In this section, we focus on a general open quantum system with gain, losses and decoherence. This quantum system of interest coupling to an environment can be experimentally realized (e.g., active and passive cavities or sink and source systems). Under appropriate approximation, the time evolution of an open quantum system can be expressed by using a non-Hermitian Lindblad master equation

$$\frac{\partial \rho(t)}{\partial t} = -i[\hat{H}\rho(t) - \rho(t)\hat{H}^{\dagger}] + \sum_{\mu} D[\hat{\Gamma}_{\mu}]\rho(t), \quad (1)$$

where \hat{H} is a non-Hermitian Hamiltonian of the system with gain and losses. $\rho(t)$ is the density matrix of the system and the dissipator $D[\Gamma_{\mu}]\rho(t)$ is a linear operator and quadratic in the Lindblad operators $\hat{\Gamma}_{\mu}$ describing directly the dissipative effects of the environment. Without loss of generality, the most general quantum dynamical semigroup form of the dissipator can be written as

$$D[\hat{\Gamma}_{\mu}]\rho(t) = \hat{\Gamma}_{\mu}\rho(t)\hat{\Gamma}^{\dagger}_{\mu} - \frac{1}{2}\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}\rho(t) - \frac{1}{2}\rho(t)\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}.$$
 (2)

The effect of $D[\hat{\Gamma}_{\mu}]$ on the density matrix $\rho(t)$ can be divided into two parts: The first one represents the quantum jump terms $\hat{\Gamma}_{\mu}\rho(t)\hat{\Gamma}^{\dagger}_{\mu}$, and the second one is the continuous non-unitary decoherence terms $\frac{1}{2}[\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}\rho(t) + \rho(t)\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}]$, which describes the continuous loss of energy, information, and coherence of the system into the environment.

Given a non-Hermitian Lindblad master equation as in Eq. (1), one can associate a so-called Liouvillian superoperator to it. We rewrite a $N \times N$ density matrix $\rho(t)$ into a column vector, e.g., $|\hat{\rho}(t)\rangle = [\rho(1,:)^T, \rho(2,:)^T, \cdots, \rho(n,:)^T]$ by using the Hilbert-Schmidt expression of the density matrix. After a straightforward algebra, we can recast the non-Hermitian Hamiltonian Lindblad equation (1) into a linear form

$$i\frac{\partial|\hat{\rho}(t)\rangle}{\partial t} = i(\hat{\mathcal{H}}' + \hat{\mathcal{G}}')|\hat{\rho}(t)\rangle$$

$$= i\hat{L}'|\hat{\rho}(t)\rangle,$$
(3)

where $\hat{\mathcal{H}}' = -i(\hat{H} \otimes I - I \otimes \hat{H}^{\dagger})$, and $\hat{\mathcal{G}}' = \sum_n \hat{\Gamma}_\mu \otimes \hat{\Gamma}_\mu^{\dagger} - \frac{\hat{\Gamma}_\mu \hat{\Gamma}_\mu^{\dagger}}{2} \otimes I - I \otimes \frac{\hat{\Gamma}_\mu \hat{\Gamma}_\mu^{\dagger}}{2}$, while \hat{L}' , acting on a basis of vector in this reduced space, is represented by a $N^2 \times N^2$ matrix. It's worth noting that \hat{L}' is included in two different effects of the open system, one is the non-Hermiticity and the other is the decoherence. Eq.(3) indicates that given a non-Hermitian Hamiltonian, together with the decoherence terms, one can fully describe the quantum dynamics of the system within the Lindblad formalism. This framework has an obvious advantage: the eigenvalues and eigenmatrices of the open system that are defined



FIG. 1. (Color online) Energy level scheme about our approach. There are loss γ_1 , gain γ_2 , coupling Ω and decay Γ .

via the relation

$$\mathcal{L}'\hat{\rho}_i = \lambda_i \hat{\rho}_i,\tag{4}$$

thus it can be obtained exactly. In fact, the condition of the steady state is not only all eigenvalues satisfying $\text{Im}[\lambda]=0$ but also at least one satisfying $\text{Im}[\lambda]=0$. Strictly speaking, the condition of the steady state in open systems is that eigenvalues satisfy

$$Im[\lambda_i] = 0 \quad and \quad Im[\lambda_{j\neq i}] \le 0. \tag{5}$$

The first equation ensures that at least one eigenstate is steady, and the second equation avoids the fact that the populations of eigenstates exponentially grow. In the following section, taking into consideration of the non-Hermiticity and decoherence, we will predict that there are new-style stable states and new-style degenerate points in this type of open system.

III. THE TWO-LEVEL SYSTEM WITH NON-HERMITICITY AND DECOHERENCE

Here, we consider a two-level system consisting of lower level $|0\rangle$ and higher level $|1\rangle$ is coupled to an environment with loss γ_1 , gain γ_2 and the spontaneous emission Γ decay from $|1\rangle$ to $|0\rangle$. The Hamiltonian of system is described as

$$\hat{H} = \begin{pmatrix} -i\gamma_1 & \Omega\\ \Omega & i\gamma_2 \end{pmatrix} /2, \tag{6}$$

combining with the decay channel

$$\hat{\Gamma}_{-} = \Gamma \cdot \hat{\sigma}_{-} = \begin{pmatrix} 0 & \Gamma \\ 0 & 0 \end{pmatrix}.$$
(7)

Therefore, the non-Hermitian Liouvillian master equation

$$i\frac{\partial\rho(t)}{\partial t} = [\hat{H}\rho(t) - \rho(t)\hat{H}^{\dagger}] + iD[\hat{\Gamma}_{-}]\rho(t).$$
(8)

Based on the framework in the last section, one can obtain the linear form superoperator of the non-Hermitian Liouvillian master equation

$$\mathcal{L}' = i \begin{pmatrix} -\gamma_1 & \frac{i\Omega}{2} & -\frac{i\Omega}{2} & \Gamma\\ \frac{i\Omega}{2} & \frac{\gamma_2 - \gamma_1 - \Gamma}{2} & 0 & -\frac{i\Omega}{2}\\ -\frac{i\Omega}{2} & 0 & \frac{\gamma_2 - \gamma_1 - \Gamma}{2} & \frac{i\Omega}{2}\\ 0 & -\frac{i\Omega}{2} & \frac{i\Omega}{2} & \gamma_2 - \Gamma \end{pmatrix}.$$
 (9)

Let us classify and explore the topological properties of eigenstates in this system. According to Eq.(4), the eigenvalues and the eigenstates of Eq.(9) have analytic expressions in principle. However, the explicit expressions are too complicated to present in the text. Get back to the Eq.(4), the second and the third rows of the eigen-equation governed by Eq.(9) become

$$-\Omega\rho_{00}/2 + i(\gamma_2 - \gamma_1 - \Gamma)\rho_{01}/2 - \Omega\rho_{11}/2 = \lambda\rho_{01}$$
(10a)

+
$$\Omega \rho_{00}/2 + i(\gamma_2 - \gamma_1 - \Gamma)\rho_{10}/2 + \Omega \rho_{11}/2 = \lambda \rho_{10}$$
 (10b)

We here focus on the topological properties of eigenstates with the case of $Re[\lambda] = 0$. Note that the elements $\rho_{00,11}$ which donate probabilities of population in $|0\rangle$ and $|1\rangle$ should be the real numbers. Therefore, the Eq.(10) reduces to

$$\rho_{01,10} = \mp \frac{i\Omega(\rho_{00} - \rho_{11})}{2Im[\lambda] + \Gamma + \gamma_1 - \gamma_2}.$$
 (11)

The above formula indicates that $\rho_{01,10}$ are either the zero or the imaginary numbers. $\rho_{01,10} = 0$ represents that there is incoherence between $|0\rangle$ and $|1\rangle$. Otherwise, there is coherence in the system and the phase between $|0\rangle$ and $|1\rangle$ must be $|\pi/2|$. In the following, we will discuss the different cases of the present system in detail.

A. Cases of the systems with $\rho_{01,10} = 0$

Suppose $\rho_{01,10} = 0$, namely without coherence in the system, two obvious conditions are determined from Eq.(11): $\rho_{00} = \rho_{11}$ or $\Omega = 0$. Meanwhile, the eigen-equation governed by Eq.(9) becomes

$$\begin{pmatrix} i(\Gamma\rho_{11} - \gamma_1\rho_{00}) \\ \frac{\Omega}{2}(\rho_{11} - \rho_{00}) \\ \frac{\Omega}{2}(\rho_{00} - \rho_{11}) \\ i(\gamma_2 - \Gamma)\rho_{11} \end{pmatrix} = Im[\lambda] \begin{pmatrix} \rho_{00} \\ 0 \\ 0 \\ \rho_{11} \end{pmatrix}.$$
(12)

From Eq.(12), one can conclude that (i) $Im[\lambda] = 0$ and $\Gamma = \gamma_1 = \gamma_2$ when $\rho_{00} = \rho_{11}$; (ii) $Im[\lambda] = (\gamma_2 - \Gamma)$ and $\rho_{00}/\rho_{11} = \Gamma/(\gamma_1 + \gamma_2 - \Gamma)$ when $\Omega = 0$, respectively. Further, their corresponding eigenmatrices in above cases become $\rho_{s1} = (\frac{1}{2}, 0, 0, \frac{1}{2})^T$ and $\rho_{s2} = (\frac{\Gamma}{\gamma_1 + \gamma_2}, 0, 0, \frac{\gamma_1 + \gamma_2 - \Gamma}{\gamma_1 + \gamma_2})$, respectively.

In order to further verify the above conclusions, we plot the eigenvelues of the Eq.(9) and the steady states for the condition Eq.(5), as shown in Fig.2. Because



FIG. 2. (Color online) Support γ_1 is the energy unit in this work. The imaginary parts of the eigenvalues in Eq.(9) as a function of γ_2/γ_1 . (a) $\Omega = \gamma_1/2$ and $\Gamma = \gamma_1$;(c) $\Omega = 0$ and $\Gamma = 2\gamma_1$. Probabilities ρ_{00} (red line) and ρ_{11} (blue line) as the function of $\gamma_1 * t$ with the initially values $\rho_{00} = 1/4$ and $\rho_{11} = 3/4$. (b) $\Omega = \gamma_1/2$ and $\gamma_2 = \Gamma = \gamma_1$; (d) $\Omega = 0$ and $\gamma_2 = \Gamma = 2\gamma_1$.

the real parts of the eigenvalues are zero, we only plot the imaginary parts of these eigenvalues as function of γ_2/γ_1 . First, we consider the conditions of $\Omega = \gamma_1/2$ and $\Gamma = \gamma_1$, as shown in Figs.2(a) and (b). One can find that there exists a steady state ρ_{s1} when $\gamma_2 = \gamma_1$. Moreover, the system should has PT-symmetric broken in the case of $\gamma_{1,2} > \Omega$. On the other hands, the fact that the populations between $|0\rangle$ and $|1\rangle$ are equal is quite different from the situation of the normal decoherence system where the population of $|0\rangle$ is always larger than case of $|1\rangle$. In other words, this steady state is not existent in normal non-Hermitan or decoherence systems. Further, an extreme case is shown in Figs.2(c) and (d)for $\Omega = 0$ and $\Gamma = 2\gamma_1$. In this case, the eigenstates including steady state ρ_{s2} are totally determined by the non-Hermiticity and the decoherence. Namely, the populations of $|0\rangle$ (or $|1\rangle$) in the steady state ρ_{s2} can be entirely controlled by the parameters $\gamma_{1,2}$ and Γ . This property is completely different from that of normal PT-symmetry non-Hermitan or decoherence systems.

More interestingly, for $\Omega = 0$, there is an exceptional point at $\gamma_1 + \gamma_2 = \Gamma$. In this case, λ_1 and λ_2 coalesce to $-i\gamma_1$, and the corresponding eigenmatrices ρ_1 and ρ_2 coalesce to $(1, 0, 0, 0)^T$. This result means that this is a specific degenerate point where there is no coherence between $|0\rangle$ and $|1\rangle$ because of no coupling. Unlike LEPs only induced by the decay channel, this degenerate point is jointly induced by the effects of non-Hermiticity and decoherence, so this point is referred as NLEPs.



FIG. 3. (Color online) The imaginary part and real part of the eigenvalues λ_1 (magenta solid line), λ_2 (green dashed line), λ_3 (red dash-dot line) and λ_4 (blue dotted line) as a function of γ_2/γ_1 in the system with $\Omega = 2\gamma_1$ for different decay channels (a), (b) $\Gamma = 0$ and (c), (d) $\Gamma = \gamma_1$.

B. Cases of the systems with $\rho_{01,10} \neq 0$

According to Eq.(11), there exist non-zero elements of $\rho_{01,10}$ if and only if $\rho_{00} \neq \rho_{11}$ and $\Omega \neq 0$. Therefore, there is coherence in the system and the phase between $|0\rangle$ and $|1\rangle$ must be $|\pi/2|$. In the other words, this state must be the eigensate of the pauli matrix σ_y . In order to observe the coherence properties of the system clearly, we consider an example of $\Omega = 2\gamma_1$, and this system is a normal non-Hermitan when $\Gamma = 0$. The energy spectra as the functions of γ_2/γ_1 are plotted in Figs.3(a) and (b). It can be clearly seen that this system has PT-symmetry in the condition of $\gamma_2 = \gamma_1$. In this case, all the imaginary parts of the eigenvalues are 0, corresponding to all the eigenstates are steady states. Besides, we also find that there are two EPs, at which four eigenvalues coalesce into only one value.

Interestingly, if there is a decay channel (e.g., $\Gamma \neq 0$), as shown in Figs.3(c) and (d), some imaginary parts of the eigenvalues split. Only ρ_1 corresponding to λ_1 , remains a steady state since $Im[\lambda_1] = 0$ and $Im[\lambda_{2,3,4}] < 0$ in this case. We plot the details of this state as the function of $\gamma_1 t$ in Fig.4. It can be seen that, the populations of this steady state do not need to be equal and there are coherences between $|0\rangle$ and $|1\rangle$ in this steady state. It is noted that the phase between $|0\rangle$ and $|1\rangle$ is still $|\pi/2|$. On the other hands, the normal EPs disappear and the NLEPs arise with the splitting of the imaginary parts of the eigenvalues, which clearly indicates the origin of NLEPs. In this case, ρ^3 and ρ^4 , corresponding to λ_3 and



FIG. 4. (Color online)(a) The probabilities ρ_{00} (blue dotted line) and ρ_{11} (red dash-dot line), (b) the imaginary parts and (c) the real parts of ρ_{12} (green dotted line), ρ_{21} (magenta dash-dot line) as the function of $\gamma_1 t$ with all the initially values $\rho_{ij} = 1/2$. The other parameters are $\Omega = 2\gamma_1$, $\gamma_2 = 1.298\gamma_1$ and $\Gamma = \gamma_1$.

 λ_4 , coalesce to

$$\rho_{NLEP} = (1 + \Xi \frac{(\Xi + \xi)}{2\Omega}, -\frac{i(\Xi + \xi)}{2\Omega}, \frac{i(\Xi + \xi)}{2\Omega}, 1)^T, (13)$$

where $\xi = \gamma_2 - \gamma_1 - \Gamma$ and $\Xi = \frac{1}{3} [54\Gamma\Omega^2 + \sqrt{2916\Gamma^2\Omega^2 - 27(\xi^2 - 4\Omega^2)^3}]^{1/3}$. Form this result above obtained, it indicates that there are two NLEPs in the system and there are coherence between $|0\rangle$ and $|1\rangle$, which is quite different from the case of no coupling. And the phase between $|0\rangle$ and $|1\rangle$ is still $|\pi/2|$, which is the same as the case of the normal EP and the case of the steady state.

IV. DISCUSSION AND CONCLUSION

In summary, we have explored the open system with non-Hermitian and decoherence effect via rewriting the non-Hermitian Lindblad equation to a linear equation. Beyond the Fermi arc with $\text{Im}[\lambda]=0$ in hermitian sysytems, we propose a general condition about the steady state and a two-level system example has been discussed. Further, we demonstrate that there are new topological properties in this system when the real part of the eigenvalue is 0, EPs and Fermi arc with $\text{Re}[\lambda]=0$ are the special cases of this conclusion. Our system can be realized in the current experimental conditions. In short, we open a new way to further explore the open systems with non-Hermiticity and decoherence.

ACKNOWLEDGMENTS

This work is supported by the National Natural Science Foundation of China (Grants No.11747107), the

- M. V. Berry, Physics of Nonhermitian Degeneracies, Czech. J. Phys. 54, 1039 (2004).
- [2] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional Topology of Non-Hermitian Systems, arXiv:1912.10048.
- [3] H. Hodaei, M. Miri, M. Heinrich, D. N. Christodoulides, M. Khajavikhan, Parity-time-symmetric microring lasers, Science **346**, 975 (2014).
- [4] L. Xiao, X. Zhan, Z. H. Bian, K. K. Wang, X. Zhang, X. P. Wang, J. Li, K. Mochizuki, D. Kim, N. Kawakami, W. Yi, H. Obuse, B. C. Sanders, and P. Xue, Observation of topological edge states in paritytime-symmetric quantum walks, Nat. Phys. 13, 1117 (2017).
- [5] P.-Y. Chen, M. Sakhdari, M. Hajizadegan, Q.S. Cui, M. M.-Cheng Cheng, R. E.-Ganainy, and A. Alù, Generalized paritytime symmetry condition for enhanced sensor telemetry, Nat. Electron. 1 297 (2018).
- [6] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Paritytime symmetry and exceptional points in photonics, Nature Materials 18, 783 (2019).
- [7] A Ghatak and T. Das, New topological invariants in non-Hermitian systems, J. Phys.: Condens. Matter **31** 263001 (2019).
- [8] Y.-M. Chu, Y. Liu, H. Liu, and J.-M. Cai, Quantum sensing with a single-qubit pseudo-Hermitian system, Phys. Rev. Lett. **124**, 020501 (2020).
- [9] F. Mostafavi, L. Q. Yuan, and H. Ramezani, Eigentates transition without undergoing an adiabatic process, Phys. Rev. Lett. **122**, 050404 (2019).
- [10] W. D. Heiss, and H.L. Harney, The chirality of exceptional points, Eur. Phys. J. D 17, 149 (2001)
- [11] C. Dembowski, B. Dietz, H.-D. Gräf, H. L. Harney, A. Heine, W. D. Heiss, and A. Richter, Observation of a Chiral State in a Microwave Cavity, Phys. Rev. Lett. 90, 034101 (2003).
- [12] J. Doppler, A. A. Mailybaev, J. Böhm, U. Kuhl, A. Girschik, F. Libisch, t. J. Milburn, P. Rabl, n. Moi-seyev, and S. Rotter, Dynamically encircling an exceptional point for asymmetric mode switching, Nature 537, 76 (2016).
- [13] W. Chen, S. K. Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, Nature (London) 548, 192 (2017).
- [14] J. Zhang, B. Peng, S. K. Özdemir, K. Pichler, D. O. Krimer, G. Zhao, F. Nori, Y. X. Liu, S. Rotter, and L. Yang, A phonon laser operating at an exceptionalpoint, Nature Photonics **12**, 479 (2018).
- [15] B. Dòra, M. Heyl, and R. Moessner, The Kibble-Zurek mechanism at exceptional points, Nat. Common. 10, 2254 (2019).
- [16] A. Cerjan, S. Huang, K. P. Chen, Y.-d. Chong, and M. C. Rechtsman, Experimental realization of a Weyl exceptional ring, Nature Photonics 13 623 (2019)

Natural Science Foundation of Guangdong province (Grants No. 2017A030307023), the Scientific Research Project of Hunan Province Department of Education (Grant Nos.19B060). Y N Guo is supported by the Program of Changsha Excellent Young Talents (kq1905005).

- [17] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and Topology in Non-Hermitian Physics, Phys. Rev. X 9, 041015 (2019).
- [18] T. Yoshida, R. Peters, N. Kawakami, and Y. Hatsugai, Symmetry-protected exceptional rings in twodimensional correlated systems with chiral symmetry, Phys. Rev. B 99, 121101(R) (2019).
- [19] S. Weidemann, M. Kremer, T. Helbig, T. Hofmann, A. Stegmaier, M. Greiter, R. Thomale, and A. Szameit, Topological funneling of light, Science aa29727 (2020).
- [20] G.S. Agarwal, Master-equation approach to spontaneous emission. Phys. Rev. A 2, 2038 (1970).
- [21] D. F. Walls, and G. J. Milburn, Quantum Optics (Springer, Berlin, 1994).
- [22] H. Breuer, and F. Petruccione, The Theory of Open Quantum Systems (Oxford University Press, Oxford, 2007).
- [23] T. Prosen, Spectral theorem for the Lindblad equation for quadratic open fermionic systems, J. Stat. Mech.: Theory Exp. 2010, P07020 (2010).
- [24] V. V. Albert and L. Jiang, Symmetries and conserved quantities in Lindblad master equations, Phys. Rev. A 89, 022118 (2014).
- [25] K. Macieszczak, M. Gută, I. Lesanovsky, and J. P. Garrahan, Towards a Theory of Metastability in Open Quantum Dynamics, Phys. Rev. Lett. **116**, 240404 (2016).
- [26] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, Spectral theory of Liouvillians for dissipative phase transitions, Phys. Rev. A 98, 042118 (2018).
- [27] Chia-Yi Ju, Adam Miranowicz, Guang-Yin Chen, and F. Nori, Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps, Phys. Rev. A 100, 062131 (2019).
- [28] F. Minganti, A. Miranowicz, R. W. Chhajlany, and F. Nori, Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps, Phys. Rev. A **100**, 062131 (2019).
- [29] T. F. Havel, Robust procedures for converting among Lindblad, Kraus and matrix representations of quantum dynamical semigroups, J. Math. Phys. 44, 534 (2003)
- [30] M. Am-Shallem, A. Levy, I. Schaefer, and R. Kosloff, Three approaches for representing Lindblad dynamics by a matrix-vector notation, arXiv: 1510. 08634.
- [31] N. Hatano, Exceptional points of the Lindblad operator of a two-level system, Mol.Phys. 117, 2121 (2019).
- [32] M. van Caspel, S. E. T. Arze, and I. P. Castillo, Dynamical signatures of topological order in the drivendissipative Kitaev chain, SciPost Phys. 6, 26 (2019).
- [33] I. I. Arkhipov, A. Miranowicz, F. Minganti, and F. Nori, Quantum and semiclassical exceptional points of a linear system of coupled cavities with losses and gain within the Scully-Lamb laser theory, Phys. Rev. A 101, 013812 (2020).

[34] F. Minganti, A. Miranowicz, R. W. Chhajlany, I. I. Arkhipov, and F. Nori, Hybrid-Liouvillian formalism connecting exceptional points of non-Hermitian Hamiltonians and Liouvillians via post selection of quantum trajectories, arXiv: 2002. 11620.