# The vectorization of the non-Hermitian Lindblad equation and its applications

Jian  $Xu^{1,*}$  and Youneng  $Guo^{2,3,\dagger}$ 

<sup>1</sup>College of Electronics and Information Engineering,

Guangdong Ocean University, Zhanjiang, 524088, China

<sup>2</sup>Interdisciplinary Center for Quantum Information,

National University of Defense Technology, Changsha 410073, People's Republic of China

<sup>3</sup>School of Electronic Information and Electrical Engineering,

Changsha University, Changsha, Hunan 410022, People's Republic of China

(Dated: January 12, 2021)

Non-Hermiticity and dissipations are two different effects of open systems and the interaction between them has yet to be studied detailedly. In this paper, we theoretically and numerically explore an open system with non-Hermiticity and dissipations. We solve non-Hermitian(NH) Hamiltonians combined with dissipative channels by using the vectorized Lindblad equation, which is an exactly solvable method to address open systems with dissipations. According to this frame and taking a two-level system as an example, we predict a new class of steady states where one eigenvalue of this system is real and the others are complex and a new class of exceptional points(EPs) due to the interplay between non-Hermiticity and dissipations. Further, we reveal the topological properties of eigenstates with  $Re[\lambda] = 0$  regime, e.g NH Fermi arcs. In short, our work brings a new insight into NH systems and provides a way to study open systems with non-Hermiticity and dissipations.

PACS numbers: 03.65.Yz, 42.50.p, 03.65.Vf

#### I. INTRODUCTION

Non-Hermiticity and dissipations are two different effects to describe open systems and the interaction between them, which is an attractive topic, has not been studied detailedly. It is well known that many new features that are inaccessible in Hermitian systems can be observed in non-Hermitian systems. One of the famous examples is the steady states of NH systems. In particular, the notion of pseudo-Hermiticity e.g. paritytime (PT) symmetry provides a general perspective on characterizing a wide class of non-Hermitian Hamiltonians with purely real spectra [1-6]. Recently, a newtype non-Hermitian system with one eigenvalue being purely real and the other being complex has been proposed[7]. However, a general tool to explore non-Hermitian systems with part of eigenvalues being purely real is still lacking. Another famous example of NH systems is EPs. Different from degeneracies in Hermitian Hamiltonians where only the eigenvalues coalesce, EPs refer to not only the eigenvalues coalesce, but also the corresponding eigenstates coalesce. The physics of EPs has been studied extensively, both theoretically and experimentally, because of the observation of novel phenomena[8–13] and widespread applications, incl. unidirectional invisibility[14], enhanced sensors[15], linewidth broadening[16], topological classification[17] and optical funneling [18].

Indeed, all of these interesting phenomena of NH systems are induced by the emergence of NH Fermi (i-Fermi) arcs corresponding to the contours with purely imaginary(real) energy in the parameter space[19]. For example, steady states with pseudo-Hermiticity or PT symmetry are all in NH i-Fermi arcs. And EPs are the endpoints of NH Fermi arcs and NH i-Fermi arcs[20]. Besides, NH Fermi(i-Fermi) arcs are also deeply related to the source of novel bulk state[21, 22], new gapless phases[23], knotted NH nodal structures[24] and so on. Based on these developments, two open questions emerge. Firstly, how to find and classify the steady state condition of non-Hermitian Hamiltonians with parts real spectra. Secondly, what is the interesting property of eigenstate in NH Fermi arcs.

In this work, we address non-Hermitian Hamiltonians through the vectorized Lindblad equation, which are usually implemented in the Hermitian Hamiltonians [25–31]. Unlike previous works use the NH systems viewpoint to study the properties of Lindblad master equation [32–34], our work use the vectorized Lindblad equation to explore NH systems and the interaction between non-Hermiticity and dissipations [35–40]. Based on this, we find that there is a new type of steady states with just parts of eigenvalues being real and new EPs due to the interplay between non-Hermiticity and dissipations. Moreover, we demonstrate that there exists a topological phase  $|\pi/2|$  of the eigenstates in NH Fermi arcs regime, while the topological phase in EPs is a specialized case of our results. These results are easy to generalize a variety of open systems, e.g. NH lattice systems[41], NH quasicrystals[42], and high order NH systems [43].

The contents of this paper are as follows. In section II, we introduce the non-Hermitian vectorized Lindblad equation and propose the condition of steady states. In section III, we consider a two-level open system with non-Hermiticity and dissipations and discuss steady states and EPs in different cases. Further, we explore the new

<sup>\*</sup> xujian\_328@163.com

<sup>&</sup>lt;sup>†</sup> guoxuyan2007@163.com

topological properties of eigenstates in NH Fermi arcs regime. Discussions and conclusions are given in section IV.

# II. THE VECTORIZED LINDBLAD EQUATION AND THE CONDITION OF STEADY STATES FOR A GENERAL OPEN SYSTEM

In this section, we extend the Lindblad, namely Lindblad-Gorini-Kossakowski-Sudarshan, master equation of a Hermitian open system[44, 45] to a non-Hermitian case by transforming density matrices into vectors. We here focus on a general non-Hermitian open system with gains, losses, and dissipations. In the limit of weak Markovian time-independent interaction, the time evolution of an open quantum system can be expressed as[46, 47]

$$\frac{\partial \rho(t)}{\partial t} = -i[\hat{H}\rho(t) - \rho(t)\hat{H}^{\dagger}] + \sum_{\mu} D[\hat{\Gamma}_{\mu}]\rho(t).$$
(1)

Note that  $\hat{H}$  is not limited to Hermiticity. We here consider it essentially is a non-Hermitian Hamiltonian  $(\hat{H} \neq \hat{H}^{\dagger})$  so it is quite different from an effective non-Hermitian Hamiltonian  $H_{eff}$  stemming from the dissipation of environment, e.g.,  $H_{eff} = H_{sys} - i \sum_{\mu} \hat{\Gamma}_{\mu} \hat{\Gamma}^{\dagger}_{\mu}/2$ .  $\rho(t)$  is the density matrices of the system and  $D[\Gamma_{\mu}]\rho(t)$ is a linear operator and the most general quantum dynamical semigroup form about this can be written as

$$D[\hat{\Gamma}_{\mu}]\rho(t) = \hat{\Gamma}_{\mu}\rho(t)\hat{\Gamma}^{\dagger}_{\mu} - \frac{1}{2}\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}\rho(t) - \frac{1}{2}\rho(t)\hat{\Gamma}_{\mu}\hat{\Gamma}^{\dagger}_{\mu}.$$
 (2)

with the Lindblad operators  $\hat{\Gamma}_{\mu}$  describing directly the dissipation effects of environments.

To solve Eq.(1), one can rewrite the  $N \times N$  density matrix  $\rho(t)$  into a column vector, e.g.,  $\hat{\rho}(t) = [\rho(1,:)^T, \rho(2,:)^T, \cdots, \rho(n,:)^T]$  by using the Hilbert-Schmidt expression of the density matrix. After a straightforward algebra, we can rewrite the non-Hermitian Hamiltonian Lindblad equation (1) as a vectorized form

$$i\frac{\partial\hat{\rho}(t)}{\partial t} = i(\hat{\mathcal{H}}' + \hat{\mathcal{G}}')\hat{\rho}(t)$$

$$= i\hat{L}'\hat{\rho}(t),$$
(3)

where  $\hat{\mathcal{H}}' = -i(\hat{H} \otimes I - I \otimes \hat{H}^{\dagger})$  and  $\hat{\mathcal{G}}' = \sum_n \hat{\Gamma}_\mu \otimes \hat{\Gamma}^{\dagger}_\mu - \frac{\hat{\Gamma}_\mu \hat{\Gamma}^{\dagger}_\mu}{2} \otimes I - I \otimes \frac{\hat{\Gamma}_\mu \hat{\Gamma}^{\dagger}_\mu}{2}$ .  $\hat{L}' = \hat{\mathcal{H}}' + \hat{\mathcal{G}}'$  is represented as a  $N^2 \times N^2$  matrix. It's worth mentioning that  $\hat{L}'$  is including two different effects of the open system, one is derived from non-Hermiticity of the system with gains and losses, the other is arising from the dissipation of environments. In this vectorized Lindblad framework, there is an obvious advantage: The analytical expressions of eigenvalues and eigenstates of the open system are easily obtained

$$\hat{L}'\hat{\rho}^{i} = \lambda_{i}\hat{\rho}^{i}, (i = 1, 2, ...n)$$
 (4)



FIG. 1. (Color online) Energy level scheme about our example. There are loss  $\gamma_1$ , gain  $\gamma_2$ , coupling  $\Omega$  and dissipations  $\Gamma$ .

where  $\lambda_i$  and  $\hat{\rho}^i$  are respectively the eigenvalues and eigenstates of the Liouvillian  $\hat{L}'$ . Due to non-Hermiticity of  $\hat{L}'$ , energy spectra become complex in general. For a time-independent Liouvillian  $\hat{L}'$ , based on the discussions about the steady state in Hermitian case[48–50], we declare there exists a steady state, if at least one of the eigenvalues(suppose it is the *j*th eigenvalue) has zero imaginary part while the remanent eigenvalues have nonpositive imaginary parts:

$$Im[\lambda_j] = 0 \quad and \quad Im[\lambda_{i\neq j}] \le 0.$$
<sup>(5)</sup>

The first equation ensures that there is at least one eigenstate being steady, and the second equation avoids the fact that the populations of the corresponding eigenstates exponentially grow. Compared to the steady state condition that appeared in Hermitian Lindblad equation, or Pseudo-Hermitian NH systems, our extension provides a more general constraint to find and classify stable open systems.

# III. STEADY STATE, NH FERMI ARCS AND EPS IN A TWO-LEVEL SYSTEM WITH NON-HERMITICITY AND DISSIPATION

For simplicity, we consider a two-level system consisting of lower level  $|0\rangle$  and higher level  $|1\rangle$  coupling to an environment with the loss  $\gamma_1$ , the gain  $\gamma_2$  and the dissipation  $\Gamma$  that describes decaying from  $|1\rangle$  to  $|0\rangle$  as shown in Fig1. This system is described by the Hamiltonian

$$\hat{H} = \frac{1}{2} \begin{pmatrix} -i\gamma_1 & \Omega\\ \Omega & i\gamma_2 \end{pmatrix}, \tag{6}$$

and a dissipative channel

$$\hat{\Gamma}_{-} = \Gamma \cdot \hat{\sigma}_{-} = \begin{pmatrix} 0 & \Gamma \\ 0 & 0 \end{pmatrix}.$$
(7)

Based on the above framework, one can obtain the vectorized superoperator of the non-Hermitian Liouvillian master equation:

$$\mathcal{L}' = i \begin{pmatrix} -\gamma_1 & \frac{i\Omega}{2} & -\frac{i\Omega}{2} & \Gamma\\ \frac{i\Omega}{2} & \frac{\gamma_2 - \gamma_1 - \Gamma}{2} & 0 & -\frac{i\Omega}{2}\\ -\frac{i\Omega}{2} & 0 & \frac{\gamma_2 - \gamma_1 - \Gamma}{2} & \frac{i\Omega}{2}\\ 0 & -\frac{i\Omega}{2} & \frac{i\Omega}{2} & \gamma_2 - \Gamma \end{pmatrix}, \quad (8)$$

whose one form of eigenvalues is

$$\lambda_{1} = i \frac{1}{2} \eta_{-},$$

$$\lambda_{2} = i \frac{1}{2} [\eta_{-} - 2i\Lambda(0)],$$

$$\lambda_{3} = i \frac{1}{2} [\eta_{-} + i\Lambda(\sqrt{3})],$$

$$\lambda_{4} = i \frac{1}{2} [\eta_{-} + i\Lambda(-\sqrt{3})],$$
(9)

and the corresponding eigenstates are given

$$\hat{\rho}^{1} \propto \begin{pmatrix} 0\\1\\1\\0 \end{pmatrix}, \\
\hat{\rho}^{2} \propto \begin{pmatrix} 1 - \frac{[\eta_{+} + 2i\Lambda(0)](3\eta_{+} + \Theta^{2} - 12\Omega^{2})}{6\Omega^{2}\Theta} \\ -\frac{i[\eta_{+} + 2i\Lambda(0)]}{2\Omega} \\ \frac{i[\eta_{+} + 2i\Lambda(0)]}{2\Omega} \\ 1 \end{pmatrix}, \\
\hat{\rho}^{3} \propto \begin{pmatrix} 1 + \frac{[\eta_{+} - i\Lambda(\sqrt{3})][-i\Lambda(\sqrt{3})]}{2\Omega} \\ -\frac{i[\eta_{+} - i\Lambda(\sqrt{3})]}{2\Omega} \\ \frac{i[\eta_{+} - i\Lambda(\sqrt{3})]}{2\Omega} \\ 1 \end{pmatrix}, \\
\hat{\rho}^{4} \propto \begin{pmatrix} 1 + \frac{[\eta_{+} - i\Lambda(-\sqrt{3})][-i\Lambda(-\sqrt{3})]}{2\Omega} \\ -\frac{i[\eta_{+} - i\Lambda(-\sqrt{3})]}{2\Omega} \\ \frac{i[\eta_{+} - i\Lambda(-\sqrt{3})]}{2\Omega} \\ 1 \end{pmatrix}.$$
(10)

Here we define  $\Lambda(x) = \frac{(i+x)\Theta}{6} + \frac{(i-x)(\eta_+^2 - 4\Omega^2)}{2\Theta}$ , with  $\eta_{\pm} = \pm \gamma_1 + \gamma_2 - \Gamma$  and  $\Theta = \left[54\Gamma\Omega^2 + \sqrt{2916\Gamma^2\Omega^4 - 27(\eta_+^2 - 4\Omega^2)^3}\right]^{1/3}$ . EPs can be found and classified by the above equations and the details will be discussed in the next section.

To further explore and classify the properties of the Liouvillian spectrum of Eq.(4), we focus on the second and third rows of the eigen-equation governed by Eq.(8)

$$-\Omega\rho_{00}^{j}/2 + i(\gamma_{2} - \gamma_{1} - \Gamma)\rho_{01}^{j}/2 + \Omega\rho_{11}^{j}/2 = \lambda_{j}\rho_{01}^{j}$$
(11a)

$$\Omega \rho_{00}^j / 2 + i(\gamma_2 - \gamma_1 - \Gamma) \rho_{10}^j / 2 - \Omega \rho_{11}^j / 2 = \lambda_j \rho_{10}^j,$$
(11b)

where  $\rho_{mn}$  (m, n = 0, 1) are the elements of eigenstates  $\rho^{j}$ . In particular, the diagonal elements  $\rho_{00,11}^{j}$  respectively donate probabilities of  $|0\rangle$  and  $|1\rangle$ , while the offdiagonal elements  $\rho_{01,10}^{j}$  represent the coherence between



FIG. 2. (Color online) The imaginary part of the eigenvalues  $\lambda_1$  (green dashed line),  $\lambda_2$  (magenta solid line),  $\lambda_3$  (red dashdot line) and  $\lambda_4$  (blue dotted line) as a function of  $\gamma_2/\gamma_1$  in the system where (a)  $\Omega = 2\gamma_1$  and  $\Gamma = \gamma_1$ ; (c)  $\Omega = 0$  and  $\Gamma = 2\gamma_1$ . Probabilities  $\rho_{00}$  (red dash-dot line) and  $\rho_{11}$  (blue dotted line) as a function of  $\gamma_1 * t$  with the initial state being  $(1/4, 0, 0, 3/4)^T$  where the paremeters of (b) and (d) are the same as those of (a) and (c). Support  $\gamma_1$  is the energy unit in this work.

 $|0\rangle$  and  $|1\rangle$ . Further, the Eq.(11) reduces to

$$\rho_{01,10}^{j} = \frac{\Omega(\rho_{00}^{j} - \rho_{11}^{j})}{2\lambda_{j} - i\eta_{-}}.$$
(12)

The coherence elements  $\rho_{01,10}^j$  are zero if  $\rho_{00}^j = \rho_{11}^j$  or  $\Omega = 0$  or purely imaginary numbers when  $\rho_{00}^j \neq \rho_{11}^j$ ,  $\Omega \neq 0$  and  $Re[\lambda_j] = 0$ . It implies that there is incoherence between  $|0\rangle$  and  $|1\rangle$  in the first case. In the second case, to ensure that the nondiagonal elements of density matrixes are purely imaginary numbers, the phase between  $|0\rangle$  and  $|1\rangle$  of the corresponding pure states must be  $|\pi/2|$ . For mixed states, each coherence part of mixed states also has the above property. Especially, as in the second case, there is not dynamic phase between  $|0\rangle$  and  $|1\rangle$  when  $Re[\lambda_j] = 0$ . So the phase is geometric, and that is topological because of  $|\pi/2|$  being fixed values. In the following, we will discuss the different cases based on the above classification.

## A. Systems without coherence

Considering the case of incoherence between  $|0\rangle$  and  $|1\rangle$ , in other words,  $\rho_{01,10}^{j} = 0$ , the eigen-equation gov-

erned by Eq.(8) becomes

$$\begin{pmatrix} i(\Gamma\rho_{11}^{j} - \gamma_{1}\rho_{00}^{j}) \\ \frac{\Omega}{2}(\rho_{11}^{j} - \rho_{00}^{j}) \\ \frac{\Omega}{2}(\rho_{00}^{j} - \rho_{11}^{j}) \\ i(\gamma_{2} - \Gamma)\rho_{11}^{j} \end{pmatrix} = \lambda_{j} \begin{pmatrix} \rho_{00}^{j} \\ 0 \\ 0 \\ \rho_{11}^{j} \end{pmatrix}.$$
(13)

When  $\rho_{00}^j = \rho_{11}^j$  is satisfied, Eq.(13) has a steady state solution with  $\Gamma = \frac{1}{2}(\gamma_1 + \gamma_2)$ . The corresponding eigenvalue and eigenstate are  $\lambda_{s1} = 0$  and  $\rho^{(s1)} = (\frac{1}{2}, 0, 0, \frac{1}{2})^T$ , respectively. Due to  $Re[\lambda_j] = 0$ , we only focus on the imaginary part of the eigenvalues  $\lambda_i$  as a function of  $\gamma_2/\gamma_1$  in the case of  $\Gamma = \gamma_1$  in Fig.2(a) and can clearly see that there exists a steady state at  $\gamma_2/\gamma_1 = 1$ . To better understand the consequences of this scenario, we have also plotted the time evolution of the state as shown in Fig.2(b). Obviously, only  $\rho^{(s1)}$  survives and the rest decay in the end. On the other hand, when  $\Omega = 0$  is satisfied, the system has an other steady state solution with  $\Gamma = \gamma_2$ . In this case, the eigenvalue and eigenstate are  $\lambda_{s2} = i(\gamma_2 - \Gamma) = 0 \text{ and } \rho^{(s2)} = (\frac{\Gamma}{\gamma_1 + \gamma_2}, 0, 0, \frac{\gamma_1 + \gamma_2 - \Gamma}{\gamma_1 + \gamma_2}),$ respectively. The spectra about  $\gamma_2$  and the time evolution of the state as whown in Fig.2(c,d). Interestingly, this steady state can be entirely controlled by the parameters of gain  $\gamma_1$ , loss  $\gamma_2$  and dissipation  $\Gamma$ . This property is completely different from that of the conventional PT symmetry non-Hermitan or dissipation systems.

What's more, for  $\Omega = 0$ , the equations  $-2\Lambda(0) = \Lambda(\sqrt{3}) = \Lambda(-\sqrt{3}) = 0$  has a solution when  $\gamma_1 + \gamma_2 = \Gamma$ . In this case, all eigenvalues degenerate and all eigenstates coalesce:

$$\lambda^{NLEPs} = -2i\gamma_1, \tag{14a}$$

$$\hat{\rho}^{NLEPs} \propto (1, 0, 0, 0)^T.$$
 (14b)

Unlike Lindblad exceptional points (LEPs) proposed in Ref.[32] in which LEPs are only induced by the decay channel, this degenerate point is jointly induced by the effects of non-Hermiticity and dissipation, therefore this point is referred as non-Hermitian Lindblad exceptional points(NLEPs).

#### B. Systems with coherence

As we previously discussed, there exists coherence if only if  $\rho_{00} \neq \rho_{11}$  and  $\Omega \neq 0$ . We first consider the case of the absence of dissipative channels( $\Gamma = 0$ ). In this case, this system reduces to a normal non-Hermitian system and the eigenvalues are shown in Figs.3(a) and 3(b). So this system exhibits PT-symmetry at  $\gamma_1 = \gamma_2$  and the corresponding eigenvalues are purely real numbers if  $\Omega \geq \gamma_{1,2}$ . On the other hands, there are two exceptional points at  $\gamma_2/\gamma_1 = \pm \frac{2\Omega}{\gamma_1} - 1$  that is also deduced from  $-2\Lambda(0) = \Lambda(\sqrt{3}) = \Lambda(-\sqrt{3}) = 0$ . And the corresponding density matrices are  $(1, \pm i, \mp i, 1)^T$ . More importantly, for  $\gamma_2/\gamma_1 \geq \frac{2\Omega}{\gamma_1} - 1$  and  $\gamma_2/\gamma_1 \leq -\frac{2\Omega}{\gamma_1} - 1$ , there



FIG. 3. (Color online) (a) The imaginary part and (b) the real part of the eigenvalues  $\lambda_1$  (green dashed line),  $\lambda_2$  (magenta solid line),  $\lambda_3$  (red dash-dot line) and  $\lambda_4$  (blue dotted line) as a function of  $\gamma_2/\gamma_1$  in the system for  $\Omega = 2\gamma_1$  and  $\Gamma = 0$ . (c) The imaginary part and (d) the real part of the  $\rho_{10}^{(2)}$  (magenta solid line),  $\rho_{10}^{(3)}$  (red dash-dot line) and  $\rho_{10}^{(4)}$  (blue dotted line) as a function of  $\gamma_2/\gamma_1$ .

are NH Fermi arcs in the purely imaginary eigenvalues regimes. In this case, based on Eq.(12), the off-diagonal elements of the corresponding density matrices must be purely imaginary numbers. In other words, the phase of eigenstates between  $|0\rangle$  and  $|1\rangle$  must be  $|\pi/2|$ . To prove it, the off-diagonal elements  $\rho_{01(10)}^{j}$  with j = 2, 3, 4 (the eigenstate  $\hat{\rho}_1$  is trivial) are shown in Figs.3(c) and 3(d). This result is clearly verified in EPs but have not been theoretically predicted and experimentally demonstrated in NH Fermi arcs regimes.

However, for  $\Gamma \neq 0$ , several distinct features emerge. Firstly, the imaginary parts of the eigenvalues split so it means that there are not PT symmetry steady state solution. This result illustrates in Figs.4(a) and 4(b). But based on Eq.(5), there is still a steady state when  $Im(E_2) = 0$  and  $Im(E_j) < 0(j = 1, 3, 4)$ . And the offdiagonal elements of  $\hat{\rho}_2$  must be purely imaginary numbers through Eq.(12). The numerical calculations show that, for a initial state  $(1/2, 1/2, 1/2, 1/2)^T$ , a steady state emerges and its off-diagonal elements of density matrix are purely imaginary numbers, as plotted in Figs.4(c) and 4(d). Secondly, allowing for  $\Lambda(\sqrt{3}) = \Lambda(-\sqrt{3})$ , we see that degeneracies occur when

$$\eta_{+} = \pm \sqrt{3(4\Gamma^2 \Omega^4)^{1/3} + 4\Omega^2}.$$
 (15)

So in this case, NLEPs emerge and the corresponding



FIG. 4. (Color online)(a) The imaginary part and (b) the real part of the eigenvalues  $\lambda_1$  (green dashed line),  $\lambda_2$  (magenta solid line),  $\lambda_3$  (red dash-dot line) and  $\lambda_4$  (blue dotted line) as a function of  $\gamma_2/\gamma_1$  for  $\Omega = 2\gamma_1$  and  $\Gamma = 2\gamma_1$ . (c) Probabilities  $\rho_{00}$  (blue line),  $\rho_{11}$  (red line) and (d) the real part (green line), imaginary part (carmine line) of  $\rho_{10}$  as a function of  $\gamma_1 * t$  when the initial state is  $(1/2, 1/2, 1/2, 1/2)^T$  for  $\gamma_2 = 1.298\gamma_1$ . (e) The imaginary part and (f) the real part of  $\rho_{10}^{(2)}$  (magenta solid line),  $\rho_{10}^{(3)}$  (red dash-dot line) and  $\rho_{10}^{(4)}$  (blue dotted line) as a function of  $\gamma_2/\gamma_1$  for  $\Omega = 2\gamma_1$  and  $\Gamma = 2\gamma_1$ .

eigenvalues and eigenstates can be computed to

$$\lambda^{NLEPs} = i \frac{1}{2} [\eta_{-} - (2\Gamma\Omega^2)^{1/3}], \qquad (16a)$$

- H. Hodaei, M. Miri, M. Heinrich, D. N. Christodoulides, M. Khajavikhan, Parity-time-symmetric microring lasers, Science **346**, 975 (2014).
- [2] L. Xiao, X. Zhan, Z. H. Bian, K. K. Wang, X. Zhang, X. P. Wang, J. Li, K. Mochizuki, D. Kim, N. Kawakami, W. Yi, H. Obuse, B. C. Sanders, and P. Xue, Observation of topological edge states in parity time-symmetric quantum walks, Nat. Phys. 13, 1117 (2017).
- [3] P.-Y. Chen, M. Sakhdari, M. Hajizadegan, Q.S. Cui, M. M.-Cheng Cheng, R. E.-Ganainy, and A. Alù, General-

$$\hat{\rho}^{NLEPs} \propto \begin{pmatrix} 1 + \frac{[\eta_{+} + (2\Gamma\Omega^{2})^{1/3}](2\Gamma\Omega^{2})^{1/3}}{2\Omega^{2}} \\ -\frac{i[\eta_{+} + (2\Gamma\Omega^{2})^{1/3}]}{2\Omega} \\ \frac{i[\eta_{+} + (2\Gamma\Omega^{2})^{1/3}]}{2\Omega} \\ 1 \end{pmatrix}.$$
(16b)

Thirdly, NH Fermi arcs also exist, i.e., they are not affected by dissipations. Meanwhile, the off-diagonal elements of the corresponding density matrices must be purely imaginary numbers through Eq.(12).  $\rho_{01(10)}^{j}$  with j = 2, 3, 4 (the eigenstate  $\hat{\rho}_1$  is trivial) as a function of  $\gamma_2/\gamma_1$  is shown in Figs.4(e) and 4(f). Obviously, NLEPs, which is shown in Eq.(16), is still a special case of NH Fermi arcs. In other words, although the eigenstates of NH Fermi arcs are mixed states in this case, the phase between  $|0\rangle$  and  $|1\rangle$  of the pure states parts is still  $|\pi/2|$ .

#### IV. DISCUSSION AND CONCLUSION

In summary, we have explored an open system with non-Hermitian and dissipations via mapping the non-Hermitian Lindblad equation to a vectorized equation. According to this frame, we propose a general condition of the steady state with open systems. As an illustration, we study a two-level system coupling to an environment with gains, losses, and dissipations. We find that there is a new class of steady states and EPs due to the interplay between non-Hermiticity and dissipations. We further demonstrate that the phase between  $|0\rangle$  and  $|1\rangle$  of eigenstates of NH Fermi arcs must be  $|\pi/2|$  while EPs is a special case of this. In short, our work brings a new insight into NH systems and provides a new tool to explore the rich variety of open systems.

## ACKNOWLEDGMENTS

This work is supported by the National Natural Science Foundation of China (Grants No.11747107), the Natural Science Foundation of Guangdong province (Grants No. 2017A030307023), the Scientific Research Project of Hunan Province Department of Education (Grant Nos.19B060). Y N Guo is supported by the Program of Changsha Excellent Young Talents (kq1905005).

ized parity time symmetry condition for enhanced sensor telemetry, Nat. Electron. **1** 297 (2018).

- [4] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Paritytime symmetry and exceptional points in photonics, Nature Materials 18, 783 (2019).
- [5] A Ghatak and T. Das, New topological invariants in non-Hermitian systems, J. Phys.: Condens. Matter **31** 263001 (2019).
- [6] Y.-M. Chu, Y. Liu, H. Liu, and J.-M. Cai, Quantum sensing with a single-qubit pseudo-Hermitian system, Phys.

Rev. Lett. 124, 020501 (2020).

- [7] F. Mostafavi, L. Q. Yuan, and H. Ramezani, Eigentates transition without undergoing an adiabatic process, Phys. Rev. Lett. **122**, 050404 (2019).
- [8] W. D. Heiss, and H.L. Harney, The chirality of exceptional points, Eur. Phys. J. D 17, 149 (2001)
- [9] C. Dembowski, B. Dietz, H.-D. Gräf, H. L. Harney, A. Heine, W. D. Heiss, and A. Richter, Observation of a Chiral State in a Microwave Cavity, Phys. Rev. Lett. 90, 034101 (2003).
- [10] J. Doppler, A. A. Mailybaev, J. Böhm, U. Kuhl, A. Girschik, F. Libisch, t. J. Milburn, P. Rabl, n. Moi-seyev, and S. Rotter, Dynamically encircling an exceptional point for asymmetric mode switching, Nature 537, 76 (2016).
- [11] A. Cerjan, S. Huang, K. P. Chen, Y.-d. Chong, and M. C. Rechtsman, Experimental realization of a Weyl exceptional ring, Nature Photonics 13 623 (2019)
- [12] T. Yoshida, R. Peters, N. Kawakami, and Y. Hatsugai, Symmetry-protected exceptional rings in twodimensional correlated systems with chiral symmetry, Phys. Rev. B 99, 121101(R) (2019).
- [13] B. Dòra, M. Heyl, and R. Moessner, The Kibble-Zurek mechanism at exceptional points, Nat. Common. 10, 2254 (2019).
- [14] A. Regensburger, C. Bersch, M.-A. Miri, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Parity-time synthetic photonic lattices, Nature 488, 167 (2012).
- [15] W. Chen, S. K. Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, Nature (London) 548, 192 (2017).
- [16] J. Zhang, B. Peng, S. K. Özdemir, K. Pichler, D. O. Krimer, G. Zhao, F. Nori, Y. X. Liu, S. Rotter, and L. Yang, A phonon laser operating at an exceptional point, Nature Photonics **12**, 479 (2018).
- [17] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and Topology in Non-Hermitian Physics, Phys. Rev. X 9, 041015 (2019).
- [18] S. Weidemann, M. Kremer, T. Helbig, T. Hofmann, A. Stegmaier, M. Greiter, R. Thomale, and A. Szameit, Topological funneling of light, Science aa29727 (2020).
- [19] M. V. Berry, Physics of Non hermitian Degeneracies, Czech. J. Phys. 54, 1039 (2004).
- [20] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional Topology of Non-Hermitian Systems, arXiv:1912.10048.
- [21] H. Zhou, C.Peng, Y.Yoon, C. W. Hsu, K. A Nelson, L. Fu, J. D. Joannopoulos, M. Solja, B. Zhen, Observation of bulk Fermi arc and polarization half charge from paired exceptional points, Science **359**, 1009 (2018).
- [22] V.Kozii, L. Fu, Non-Hermitian Topological Theory of Finite-Lifetime Quasiparticles: Prediction of Bulk Fermi Arc Due to Exceptional Point, Physica E Low Dimens.Syst. Nanostruct. 99, 63 (2018).
- [23] T. Bessho, K. Kawabata, and M. Sato, Topological Classificaton of Non-Hermitian Gapless Phases: Exceptional Points and Bulk Fermi Arcs, JPS Conf. Proc. **30**, 011098 (2020).
- [24] J. Carlström, M. Stalhammar, J. C. Budich, E. J. Bergholtz, Knotted Non-Hermitian Metals, Phys. Rev. B 99, 161115 (2019).

- [25] G.S. Agarwal, Master-equation approach to spontaneous emission. Phys. Rev. A 2, 2038 (1970).
- [26] T. Prosen, Spectral theorem for the Lindblad equation for quadratic open fermionic systems, J. Stat. Mech.: Theory Exp. 2010, P07020 (2010).
- [27] V. V. Albert and L. Jiang, Symmetries and conserved quantities in Lindblad master equations, Phys. Rev. A 89, 022118 (2014).
- [28] K. Macieszczak, M. Gută, I. Lesanovsky, and J. P. Garrahan, Towards a Theory of Metastability in Open Quantum Dynamics, Phys. Rev. Lett. **116**, 240404 (2016).
- [29] M. van Caspel, S. E. T. Arze, and I. P. Castillo, Dynamical signatures of topological order in the drivendissipative Kitaev chain, SciPost Phys. 6, 26 (2019).
- [30] I. I. Arkhipov, A. Miranowicz, F. Minganti, and F. Nori, Quantum and semiclassical exceptional points of a linear system of coupled cavities with losses and gain within the Scully-Lamb laser theory, Phys. Rev. A 101, 013812 (2020).
- [31] F. Minganti, A. Miranowicz, R. W. Chhajlany, I. I. Arkhipov, and F. Nori, Hybrid-Liouvillian formalism connecting exceptional points of non-Hermitian Hamiltonians and Liouvillians via postselection of quantum trajectories, Phys. Rev. A 101, 062112 (2020).
- [32] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, Spectral theory of Liouvillians for dissipative phase transitions, Phys. Rev. A 98, 042118 (2018).
- [33] Chia-Yi Ju, Adam Miranowicz, Guang-Yin Chen, and F. Nori, Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps, Phys. Rev. A 100, 062131 (2019).
- [34] F. Minganti, A. Miranowicz, R. W. Chhajlany, and F. Nori, Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps, Phys. Rev. A **100**, 062131 (2019).
- [35] T. F. Havel, Robust procedures for converting among Lindblad, Kraus and matrix representations of quantum dynamical semigroups, J. Math. Phys. 44, 534 (2003).
- [36] M. Am-Shallem, A. Levy, I. Schaefer, and R. Kosloff, Three approaches for representing Lindblad dynamics by a matrix-vector notation, arXiv:1510.08634.
- [37] M. Keck, S. Montangero, G. E. Santoro, R. Fazio, and D. Rossini. Dissipation in adiabatic quantum computers: Lessons from an exactly solvable model, New J. Phys. 19 113029 (2017).
- [38] V. Reimer, K. G. L. Pedersen, N. Tanger, M. Pletyukhov, and V.Gritsev. Phys. Rev. A 97, 043851 (2018).
- [39] S. Scopa, G. T. Landi, A. Hammoumi, and D. Karevski, Exact solution of time-dependent Lindblad equations with closed algebras, Phys. Rev. A, **99** 022105 (2019).
- [40] N. Hatano, Exceptional points of the Lindblad operator of a two-level system, Mol.Phys. 117, 2121 (2019).
- [41] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological phases of non-Hermitian systems, Phys. Rev. X 8, 031079 (2018).
- [42] S. Longhi, Topological phase transition in non-Hermitian quasicrystals, Phys. Rev. Lett. 122, 237601 (2019).
- [43] W. Tang, X. Jiang, K. Ding, Y.-X. Xiao, Z.-Q. Zhang, C. T. Chan, and G. Ma, Exceptional nexus with a hybrid topological invariant, Science, **370**, 6250, (2020).
- [44] G. Lindblad, On the Generators of Quantum DynamicalSemigroups, Commun. math. Phys.48,119 (1976).
- [45] V. Gorini, A. Kossakowski, and E.C.G. Sudarshan, Completely positive dynamical semigroups of N-level systems,

J. Math. Phys. 17,821 (1976).

- [46] D. F. Walls, and G. J. Milburn, Quantum Optics (Springer, Berlin, 1994).
- [47] H. Breuer, and F. Petruccione, The Theory of Open Quantum Systems (Oxford University Press, Oxford, 2007).
- [48] V. V. Albert and L. Jiang, Symmetries and conserved quantities in Lindblad master equations, Phys. Rev. A

89, 022118 (2014).

- [49] V. V. Albert, B. Bradlyn, M. Fraas, L. Jiang, Geometry and response of Lindbladians, Phys. Rev. X 6, 041031 (2016).
- [50] D. Nigro, On the uniqueness of the steady-state solution of the Lindblad–Gorini–Kossakowski–Sudarshan equation, J. Stat. Mech. 043202 (2019)