

Dependence of temperatures and kinetic freeze-out volume on centrality in Au-Au and Pb-Pb collisions at high energy

Muhammad Waqas^{1,*}, Fu-Hu Liu^{1,†}, Zafar Wazir^{2,‡}

¹*Institute of Theoretical Physics & State Key Laboratory of Quantum Optics and Quantum Optics Devices, Shanxi University, Taiyuan, Shanxi 030006, China*

²*Department of Physics, International Islamic University, H-10, Islamabad, Pakistan*

Abstract: Centrality-dependent double-differential transverse momentum spectra of negatively charged particles (π^- , K^- and \bar{p}) at mid-(pseudo)rapidity interval in nuclear collisions are analyzed by the standard distribution in terms of multi-component. The experimental data measured in gold-gold (Au-Au) collisions by the PHENIX Collaboration at the Relativistic Heavy Ion Collider (RHIC) and in lead-lead (Pb-Pb) collisions by the ALICE Collaboration at the Large Hadron Collider (LHC) are studied. The effective temperature, initial temperature, kinetic freeze-out temperature, transverse flow velocity and kinetic freeze-out volume are extracted from the fitting to transverse momentum spectra. We observed, that the mentioned five quantities increase with the increase of event centrality due to the fact that the average transverse momentum increases with the increase of event centrality. This renders that larger momentum (energy) transfer and further multiple-scattering had happened in central centrality.

Keywords: Effective temperature, Initial temperature, Kinetic freeze-out temperature, Transverse flow velocity, Kinetic freeze-out volume

PACS: 12.40.Ee, 13.85.Hd, 25.75.Ag, 25.75.Dw, 24.10.Pa

1 Introduction

One of the most important questions in high energy collisions is the identification of various phases of dense matter. It is expected to reach a deconfined state of matter (quarks and gluons) at high energy or density. This state of matter is called Quark-Gluon Plasma (QGP), which was obtained in the early universe shortly after the big-bang prior to the condensation in hadrons. The characterization of phase transition in finite system is a fascinating multi-disciplinary topic which has been studied for decades [1, 2] within different phenomenological applications. The Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC) have been providing excellent tools to determine the phase structure of the strongly interacting Quantum Chromodynamics (QCD) matter [3, 4, 5] and to study the properties of QGP [6, 7, 8].

Within the framework of statistical thermal models, it is assumed that the initial stage of collisions of nuclei at the RHIC and LHC [9, 10, 11] gives a tremendous amount of temperature, where a hot and dense “fireball” over an extended region for a very short period of

time (almost 10^{-22} seconds) is formed. The fireball consists of QGP and it instantly cools which results in the expansion of the system (the change of the volume or density of the system) and the partons recombine into blizzard of hadronic matter [12]. After the hadronization of the fireball, the hadrons continuously interact with each other and the particle number changes. This process results in decrease of temperature and at a certain value where the reaction process stops and the temperature at this point is called the “chemical freeze-out temperature” (T_{ch}). At the stage of chemical freeze-out, the yield ratios of different types of particles remain invariant [13].

However, the rescattering process still take place which continues to build up the collective (hydrodynamical) expansion. Resultantly, the matter becomes dilute and the mean free path of the given hadrons for the elastic reaction processes become comparable with the size of the system. At this stage, the rescattering process stops, which results in the decoupling of hadrons from the rest of the system [14]. This stage is called as the kinetic or thermal freeze-out stage, and the temperature at this stage is called the kinetic or thermal

*E-mail: waqas_phy313@yahoo.com

†Corresponding author. E-mail: fuhuliu@163.com; fuhuliu@sxu.edu.cn

‡E-mail: zafar.wazir@iiu.edu.pk

freeze-out temperature (T_0). After this stage the particle's energy/momentum spectrum is frozen in time and it is the least stage of the system evolution. Meanwhile, at this stage, particle's movement is also affected by the flow effect which should be excluded when one extracts T_0 . To describe the flow effect, one may use the transverse flow velocity β_T .

The above discussed T_0 and β_T can be extracted from transverse momentum (p_T) spectra of particles. Also, from p_T spectra, one can extract the initial temperature T_i according to the color string percolation model [15, 16, 17]. Generally, if the flow effect is not excluded in the temperature parameter, this type of temperature is called the effective temperature (T). At least, three types of temperatures, T , T_i and T_0 , can be extracted from p_T spectra. Although the yield ratios of different types of particles can be also obtained from the normalizations of p_T spectra and then T_{ch} can be also extracted from p_T spectra, we mainly extract other three types of temperatures and β_T from p_T spectra in this paper due to their more pending situations.

In addition, volume is also an important parameter in high energy collisions. The volume occupied by the ejectiles when the mutual interactions become negligible, and the only force they feel is the columbic repulsive force, is known as the kinetic freeze-out volume (V). There are various freeze-out volumes at various freeze-out stages, but we are only focusing on the kinetic freeze-out volume V in the present work. As we know, V gives the information of the co-existence of phase-transition, and is important in the extraction of multiplicity, micro-canonical heat capacity and it's negative branch or shape of the caloric curves under the thermal constraints [18, 19, 20, 21, 22]. In this paper, the fifth quantity extracted from p_T spectra is V . By way of parenthesis, the mean p_T , i.e. $\langle p_T \rangle$, is also obtained.

The study of three types of temperatures, transverse flow velocity and kinetic freeze-out volume is very wide, interesting and of course a huge project. However, in this paper we will only analyze the centrality dependences of the five quantities in gold-gold (Au-Au) collisions at 200 GeV and in lead-lead (Pb-Pb) collisions at 2.76 TeV. Only the p_T spectra of negatively charged pions (π^-) and kaons (K^-) and antiprotons (\bar{p}) are used in the extraction. These representational spectra are enough to extract the necessary centrality dependences.

The remainder of this paper orderly consists of the method and formalism, results and discussion, and summary and conclusions which are presented in section 2, 3 and 4 respectively.

2 The method and formalism

Soft excitation and hard scattering processes are the two generally considered processes for the particle production. Soft excitation process contributes in a narrow p_T range which is less than 2–3 GeV/ c or up to 4–5 GeV/ c and is responsible for the production of most of the light flavored particles. The soft excitation process has various choices of formalisms including but are not limited to the Hagedorn thermal model (Statistical-Bootstrap model) [23], the standard distribution [24], the blast-wave model with Boltzmann-Gibbs statistics [25, 26, 27], the blast-wave model with Tsallis statistics [28, 29, 30], and current thermodynamical related models [31, 32, 33, 34]. The main contributor to the produced particles is the soft excitation process.

If necessary, for the hard excitation process, there is limited choice of formalisms [35, 36, 37] and can be described by the theory of strong interaction. In fact, the contribution of hard scattering process is parameterized to an inverse power law, i.e. the Hagedorn function [23]

$$f_H(p_T) = Ap_T \left(1 + \frac{p_T}{p_0} \right)^{-n}, \quad (1)$$

where p_0 and n are free parameters, and A is the normalized constant related to the free parameters. The inverse power law is obtained from the calculus of QCD [3, 4, 5], and has at least three revisions, which is out of focus of the present work and will not be discussed further.

Different probability density functions can be used to describe the contributions of soft excitation and hard scattering processes. Due to few fraction and being earlier than the kinetic freeze-out stage, the hard scattering process does not contribute largely to T_0 and β_T in general. In fact, the contribution of hard scattering process can be neglected if we study the p_T spectra in a narrow range, say $p_T < 2-3$ GeV/ c or extending to $p_T < 4-5$ GeV/ c , for which only the contribution of soft excitation process is indeed needed. In our opinion, various distributions show similar behaviors in case of fitting the data with acceptable representations, which results in similar $\langle p_T \rangle$ ($\sqrt{\langle p_T^2 \rangle}$) with different parameters.

For the spectra contributed by the soft excitation process, we can choose the standard distribution, as it is very close in concept to the ideal gas model. The standard distribution is the combination of Boltzmann, Fermi-Dirac and Bose-Einstein distributions. The probability density function of the standard distribution in

terms of p_T at mid-rapidity is generally as [24]:

$$f_S(p_T) = C \frac{gV'}{(2\pi)^2} p_T \sqrt{p_T^2 + m_0^2} \times \left[\exp\left(\frac{\sqrt{p_T^2 + m_0^2}}{T}\right) + S \right]^{-1}, \quad (2)$$

where C is the normalization constant, V' is the fitted kinetic freeze-out volume (in terms of interaction volume) of the emission source at the stage of kinetic freeze-out as discussed above, g is the degeneracy factor for pion and kaon (or proton) and has the value of 3 (or 2), m_0 is the rest mass of the considered particle, $S = -1$ (+1) is for bosons (fermions), and T is the effective temperature as discussed above.

By considering the experimental rapidity range $[y_{\min}, y_{\max}]$ around the mid-rapidity, the accurate form of Eq. (2) is [24]

$$f_S(p_T) = C \frac{gV'}{(2\pi)^2} p_T \sqrt{p_T^2 + m_0^2} \int_{y_{\min}}^{y_{\max}} \cosh y \times \left[\exp\left(\frac{\sqrt{p_T^2 + m_0^2} \cosh y}{T}\right) + S \right]^{-1} dy. \quad (3)$$

T and V' in Eqs.(2) and (3) are free parameters. In most cases, the single component standard distribution is not enough for the simultaneous description of low- p_T region. So we have to use a two-component standard distribution.

In particular, in some cases, the multi-component (l -component) standard distribution has to be used, which can be demonstrated as:

$$f_S(p_T) = \sum_{i=1}^l k_i C_i \frac{gV'_i}{(2\pi)^2} p_T \sqrt{p_T^2 + m_0^2} \int_{y_{\min}}^{y_{\max}} \cosh y \times \left[\exp\left(\frac{\sqrt{p_T^2 + m_0^2} \cosh y}{T_i}\right) + S \right]^{-1} dy, \quad (4)$$

where C_i is the normalization constant, k_i represents the fraction contributed by the i th component, and T_i and V'_i are free parameter denoted the kinetic freeze-out temperature and volume respectively corresponding to the i th component.

In case of considering both the contributions of soft excitation and hard scattering processes, we use the superposition in principle

$$f_0(p_T) = k_S f_S(p_T) + (1 - k_S) f_H(p_T), \quad (5)$$

where k_S is the contribution ratio of soft excitation process. The contribution ranges of the soft excitation and hard scattering processes described by Eq. (5) are intersectant in low- p_T region.

Another type of superposition which uses the usual step function $\theta(x)$ based on the Hagedorn model [23] is

$$f_0(p_T) = A_1 \theta(p_1 - p_T) f_S(p_T) + A_2 \theta(p_T - p_1) f_H(p_T), \quad (6)$$

where A_1 and A_2 are the normalized constants which synthesize $A_1 f_S(p_1) = A_2 f_H(p_1)$. The contribution ranges of the soft excitation and hard scattering processes described by Eq. (6) are segregative at $p_T = p_1$.

In the present work, we will study only the first component in Eqs. (5) or (6), because we do not study a very wide p_T range. Meanwhile, we use the two-component standard distribution, i.e. $l = 2$ in Eq. (4) in which the first or second components has no particular priority. As probability density functions, Eqs. (1)–(6) are normalized to 1 naturally. When we compare the probability density functions with the experimental data which appear usually in other forms, a suitable transformation for the probability density function is certainly needed. Considering the treatment of normalization, the real fitted kinetic freeze-out volume should be $V'_i/(C_i k_i)$ which will be simply used in the following section as the symbols, V_i or V ($= \sum_{i=1}^l V_i$).

3 Results and discussion

Figure 1 presents the event centrality dependent double differential p_T spectra, $(1/2\pi p_T) d^2N/dp_T dy$, of the identified particles (π^- , K^- and \bar{p}) produced in Au-Au collisions at $\sqrt{s_{NN}} = 200$ GeV in the mid-pseudorapidity interval of $|\eta| < 0.35$, where N and y stands for the number of particles and rapidity respectively. The symbols are the the experimental data measured by the PHENIX Collaboration [38]. The spectra of centralities 0–5%, 5%–10%, 10%–15%, 15–20%, 20–30%, 30–40%, 40–50%, 50-60%, 60–70%, 70–80% and 80–92% in the three panels are multiplied by 20, 10, 5, 2.5, 1.5, 1, 1, 1, 1, 1 and 1, respectively. In addition, the spectra of centralities 15–20%, 20–30%, 30–40%, 40–50%, 50-60%, 60–70%, 70–80% and 80–92% in panel (a) are re-multiplied by 0.8, 0.7, 0.6, 0.45, 0.3, 0.24, 0.17 and 0.15 respectively. The curves are our fitting results by using the two-component standard distribution, Eq. (4) with $l = 2$. The substantially together event centralities, the values of free parameters (T_1 , T_2 , V_1 and V_2), χ^2 and the number of degree of freedom (ndof) are listed in Table 1. One can see the well approximate description of the model results to the experimental data of the PHENIX Collaboration in special p_T ranges in high energy Au-Au collisions at the RHIC.

Figure 2 is the same as Fig. 1 but it gives the results

Table 1. Values of effective temperatures (T_1 and T_2), volumes (V_1 and V_2), χ^2 , and number of degree of freedom (ndof) corresponding to the curves in Figs. 1–2.

Figure	Particle	Centrality	T_1 (GeV)	T_2 (GeV)	V_1 ((fm) ³)	V_2 ((fm) ³)	χ^2	ndof	
Fig. 1 Au-Au 200 GeV	π^-	0–5%	0.181 ± 0.014	0.268 ± 0.006	745 ± 100	1967 ± 309	45	24	
		5–10%	0.138 ± 0.013	0.270 ± 0.006	564 ± 80	2054 ± 174	14	24	
		10–15%	0.179 ± 0.015	0.250 ± 0.005	430 ± 62	2004 ± 153	233	24	
		15–20%	0.167 ± 0.012	0.247 ± 0.003	354 ± 61	1862 ± 320	63	24	
		20–30%	0.151 ± 0.009	0.245 ± 0.004	360 ± 39	1854 ± 216	87	24	
		30–40%	0.125 ± 0.015	0.237 ± 0.006	195 ± 30	2005 ± 189	276	24	
		40–50%	0.148 ± 0.014	0.227 ± 0.004	106 ± 25	1585 ± 171	388	24	
		50–60%	0.098 ± 0.011	0.227 ± 0.003	59 ± 11	1048 ± 153	146	24	
		60–70%	0.125 ± 0.010	0.274 ± 0.006	37 ± 11	984 ± 109	71	24	
		70–80%	0.112 ± 0.014	0.209 ± 0.007	18 ± 3	296 ± 37	256	24	
		80–92%	0.068 ± 0.015	0.221 ± 0.006	8.0 ± 1.0	213 ± 13	69	24	
		K^-	0–5%	0.201 ± 0.009	0.271 ± 0.008	196 ± 36	1081 ± 178	17	12
	5–10%		0.188 ± 0.007	0.255 ± 0.008	32 ± 6	1116 ± 71	18	12	
	10–15%		0.211 ± 0.013	0.250 ± 0.006	37 ± 6	1030 ± 102	24	12	
	15–20%		0.135 ± 0.014	0.248 ± 0.008	35 ± 5	836 ± 105	14	12	
	20–30%		0.219 ± 0.015	0.241 ± 0.006	10 ± 1	733 ± 66	18	12	
	30–40%		0.156 ± 0.010	0.238 ± 0.007	4.0 ± 0.3	665 ± 82	13	12	
	40–50%		0.130 ± 0.011	0.235 ± 0.008	13 ± 1.2	253 ± 39	24	12	
	50–60%		0.170 ± 0.013	0.230 ± 0.006	1.6 ± 0.2	577 ± 48	9	12	
	60–70%		0.163 ± 0.015	0.225 ± 0.006	0.50 ± 0.02	544 ± 50	35	12	
	70–80%		0.155 ± 0.015	0.220 ± 0.007	0.30 ± 0.01	505 ± 60	215	12	
	80–92%		0.130 ± 0.010	0.218 ± 0.005	0.030 ± 0.001	460 ± 130	66	12	
	\bar{p}		0–5%	0.280 ± 0.017	0.320 ± 0.003	36 ± 7	786 ± 120	17	18
		5–10%	0.272 ± 0.015	0.317 ± 0.002	31 ± 7	775 ± 70	20	18	
10–15%		0.265 ± 0.018	0.313 ± 0.006	16 ± 3	747 ± 86	162	18		
15–20%		0.185 ± 0.016	0.313 ± 0.005	23 ± 4	710 ± 80	35	18		
20–30%		0.211 ± 0.013	0.309 ± 0.006	13 ± 1	678 ± 65	41	18		
30–40%		0.247 ± 0.018	0.317 ± 0.007	10 ± 1	634 ± 50	11	18		
40–50%		0.244 ± 0.015	0.308 ± 0.008	4.0 ± 0.5	597 ± 37	39	18		
50–60%		0.237 ± 0.013	0.300 ± 0.004	1.6 ± 0.3	574 ± 70	43	18		
60–70%		0.210 ± 0.013	0.285 ± 0.006	0.50 ± 0.02	476 ± 45	28	18		
70–80%		0.170 ± 0.014	0.267 ± 0.005	0.30 ± 0.01	455 ± 50	110	18		
80–92%		0.172 ± 0.010	0.218 ± 0.004	0.40 ± 0.01	406 ± 40	5	18		
Fig. 2 Pb-Pb 2.76 TeV		π^-	0–5%	0.179 ± 0.010	0.340 ± 0.004	2517 ± 208	4099 ± 252	206	37
	5–10%		0.172 ± 0.012	0.331 ± 0.004	2226 ± 135	3468 ± 201	101	37	
	10–20%		0.165 ± 0.010	0.312 ± 0.005	1395 ± 70	2813 ± 176	70	37	
	20–30%		0.161 ± 0.014	0.300 ± 0.006	1012 ± 103	2427 ± 231	94	37	
	30–40%		0.147 ± 0.013	0.280 ± 0.005	534 ± 56	3502 ± 165	244	37	
	40–50%		0.129 ± 0.012	0.289 ± 0.005	307 ± 38	3600 ± 198	110	37	
	50–60%		0.108 ± 0.010	0.281 ± 0.005	176 ± 32	3380 ± 213	206	37	
	60–70%		0.114 ± 0.015	0.290 ± 0.005	94 ± 13	3150 ± 147	125	35	
	70–80%		0.130 ± 0.010	0.287 ± 0.007	45 ± 9	2930 ± 206	502	35	
	80–90%		0.118 ± 0.014	0.287 ± 0.006	16 ± 2	1945 ± 102	119	35	
	K^-		0–5%	0.277 ± 0.010	0.361 ± 0.007	170 ± 36	3442 ± 208	3	32
			5–10%	0.270 ± 0.009	0.359 ± 0.009	152 ± 10	3237 ± 240	15	32
		10–20%	0.252 ± 0.009	0.354 ± 0.007	97 ± 11	3190 ± 187	8	32	
		20–30%	0.265 ± 0.009	0.350 ± 0.006	53 ± 7	3136 ± 200	22	32	
		30–40%	0.247 ± 0.008	0.395 ± 0.006	27 ± 4	3014 ± 198	12	32	
		40–50%	0.210 ± 0.008	0.341 ± 0.006	22 ± 4	2828 ± 190	119	32	
		50–60%	0.213 ± 0.011	0.345 ± 0.005	8.1 ± 1.0	2659 ± 210	26	32	
		60–70%	0.226 ± 0.009	0.335 ± 0.005	5.8 ± 1.3	2132 ± 176	46	32	
		70–80%	0.225 ± 0.010	0.332 ± 0.006	2.6 ± 0.3	1736 ± 134	71	32	
		80–90%	0.170 ± 0.009	0.320 ± 0.006	0.50 ± 0.05	1150 ± 120	75	32	
		\bar{p}	0–5%	0.426 ± 0.012	0.421 ± 0.007	55 ± 6	1965 ± 195	49	33
			5–10%	0.300 ± 0.010	0.425 ± 0.005	75 ± 6	1782 ± 164	373	33
	10–20%		0.427 ± 0.010	0.397 ± 0.005	50 ± 5	1748 ± 140	43	33	
	20–30%		0.405 ± 0.013	0.395 ± 0.006	31 ± 4	1690 ± 130	52	33	
	30–40%		0.400 ± 0.012	0.382 ± 0.005	29 ± 4	1534 ± 158	20	33	
	40–50%		0.352 ± 0.011	0.383 ± 0.006	15 ± 2	401 ± 135	109	33	
	50–60%		0.331 ± 0.012	0.380 ± 0.005	6.0 ± 0.5	1357 ± 125	67	33	
	60–70%		0.310 ± 0.012	0.369 ± 0.006	8.7 ± 1.0	1069 ± 90	75	33	
	70–80%		0.321 ± 0.015	0.333 ± 0.004	1.0 ± 0.3	978 ± 50	104	33	
	80–90%		0.310 ± 0.014	0.328 ± 0.006	0.70 ± 0.04	708 ± 60	122	33	




Fig. 1. Centrality dependant $(1/2\pi p_T)d^2N/dp_T dy$ of (a) π^- , (b) K^- and (c) \bar{p} produced in $|\eta| < 0.35$ in Au-Au collisions at 200 GeV. The symbols represent experimental data of the PHENIX Collaboration [38], while the curves are the results of our fits by using the two-component standard distribution, Eq. (4) with $l = 2$. The spectra of centralities 0–5%, 5%–10%, 10%–15%, 15–20%, 20–30%, 30–40%, 40–50%, 50–60%, 60–70%, 70–80% and 80–92% in the three panels are multiplied by 20, 10, 5, 2.5, 1.5, 1, 1, 1, 1 and 1, respectively. In addition, the spectra of centralities 15–20%, 20–30%, 30–40%, 40–50%, 50–60%, 60–70%, 70–80% and 80–92% in panel (a) are re-multiplied by 0.8, 0.7, 0.6, 0.45, 0.3, 0.24, 0.17 and 0.15 respectively.

for π^- , K^- and \bar{p} in different centrality bins in Pb-Pb collisions at 2.76 TeV in the mid-rapidity interval $|y| < 0.5$. The experimental data of the ALICE Collaboration is represented by the symbols [39], where as the spectra of Pb-Pb is scaled by the factor of 2^n and n changes from 9 to 0 with the change of event centrality such as from 0–5% to 80–90%. The related parameters and the existed centralities are listed together in Table 1. One can see the well approximate description of the model results to the experimental data of the ALICE Collaboration in special p_T ranges in high energy Pb-Pb collisions at the LHC.

To study the change trend of parameters with centrality, Fig. 3 shows the dependence of effective temperature on centrality for the productions of π^- , K^- and \bar{p} in different centrality bins in Au-Au and Pb-Pb collisions at 200 GeV and 2.76 TeV respectively. Panel (a) show the result for Au-Au collisions, while panel (b) show the result for Pb-Pb collisions. Different symbols represent different particles. One can see the clear decrease of effective temperature from central to peripheral collisions. The reason behind it is, the more violent collisions in central collisions where it can get a degree of higher excitation and also involve more number of




Fig. 2. Same as Fig. 1, but for the spectra of π^- , K^- and \bar{p} in $|y| < 0.5$ in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. The symbols represent the measured data of the ALICE Collaboration [39], where the spectra are scaled by factors 2^{2n} and n changes from 9 to 0 as the event centrality changes from 0–5% to 80–90%.

participants in interactions, while they decrease from central to peripheral collisions. The effective temperatures in collisions at the LHC in different centrality bins are higher than those at the RHIC due to more energy deposition in collisions at the LHC.

Figure 4 is the same to Fig. 3, however, it shows the result for the dependence of kinetic freeze-out volume V in events with different centralities C , where $V = V_1 + V_2$ due to the additivity of volume. One can see that the kinetic freeze-out volume decreases from central to peripheral collisions, as the number of participant nucleons decreases from central to peripheral collisions depending on the interaction volume. Due to large number of binary collisions by the rescattering of partons, the system with more participants reaches quickly to equilibrium state, but the decreases in centrality, the de-

creases the number of participants and the system goes slowly to equilibrium state. The large volume and more number of participants in the central collisions may indicate the occupation of super-dense hadronic matter, but of course further and more complete information about the local energy density of super-hadronic matter is needed to study the possible phase transition of QGP. The figure shows a volume differential scenario too. The heavier the particle is, the smaller the kinetic freeze-out volume has, which shows the early freeze-out of heavier particles as compared to the lighter particles and suggests different freeze-out surfaces for different particles. Such result can be found in literature [40, 41].

The dependence of mean transverse momentum $\langle p_T \rangle$ in different centrality events is shown in Fig. 5. The symbols represent $\langle p_T \rangle$ for different particles obtained




Fig. 3. Dependence of effective temperature T in different centrality bins C in (a) Au-Au collisions at 200 GeV and (b) Pb-Pb collisions at 2.76 TeV. The obtained values corresponded to identified particles are extracted from the experimental p_T spectra.




Fig. 4. Same as Fig. 3, but for dependence of kinetic freeze-out volume V on centrality C .

from the fitting function Eq. (4) with $l = 2$ over a p_T range from 0 to 5 GeV/ c , where the parameter values are listed in Table 1. One can see that $\langle p_T \rangle$ decreases from central to peripheral Au-Au and Pb-Pb collisions for all particle species, and it may be caused due to decreasing the participant nucleons from central to peripheral collisions and this result is similar to ref. [42]. It is also important to notice that \bar{p} spectra exhibit a concave shape in the peripheral events, which is well described by the power law parametrization as observed in ref. [43], but this curvature decreases with the increasing centrality and it leads to an almost exponential dependence on $\langle p_T \rangle$ for the most central collisions. Furthermore, $\langle p_T \rangle$

for heavier particle is larger than that for lighter ones, and $\langle p_T \rangle$ at LHC energy is slightly larger than that at RHIC energy. The increase of $\langle p_T \rangle$ in central collisions and with the massive mass of the particle may indicate the collective radial flow, and the same behavior is observed at a few GeV [44] and more than 10 GeV [45].

Figure 6 is the same to Fig. 5, however it demonstrates the result for the initial temperature T_i , where T_i is obtained by the root-mean-square p_T divide by $\sqrt{2}$, i.e. $\sqrt{\langle p_T^2 \rangle}/2$ according to the color string percolation model [15, 16, 17]. The symbols are the representation of the results obtained from the fitting function Eq.(4) with $l = 2$ over a p_T range in 0–5 GeV/ c and with the




Fig. 5. Same as Fig. 3, but for dependence of mean transverse momentum $\langle p_T \rangle$ on centrality C .




Fig. 6. Same as Fig. 3, but for dependence of initial temperature T_i on centrality C .

parameter values listed in Table 1. The mass differential temperature scenario is also observed. It is necessary to point out that the initial temperature obtained in this work is larger than the effective temperature which is in agreement with the order of time evolution of interacting system.

Figure 7 shows the dependence of (a)(b) effective temperature T on rest mass m_0 and (c)(d) mean transverse momentum $\langle p_T \rangle$ on mean energy or mean moving mass \bar{m} in the rest frame of emission source for (a)(c) Au-Au collisions at 200 GeV and (b)(d) Pb-Pb collisions at 2.76 TeV in different centrality classes. The symbols represent the derived quantities accord-

ing to the free parameters listed in Table 1, where $T = (T_1 V_1 + T_2 V_2)/(V_1 + V_2)$, $\langle p_T \rangle = \int p_T f_S(p_T) dp_T$, and $\bar{m} = \langle \sqrt{p_T^2 + p_z^2 + m_0^2} \rangle$, where p_z is longitudinal momentum and its distribution can be obtained according to p_T distribution if we assume isotropic emission in the source rest frame [46]. The lines are the results fitted for the values of derived quantities if we assume linear correlations are existent.

The intercepts in Figs. 7(a) and 7(b) are regarded as the kinetic freeze-out temperature T_0 , and the slopes in Figs. 7(c) and 7(d) are regarded as the transverse flow velocity β_T [47, 48, 49, 50]. The dependences of (a) T_0 on C and (b) β_T on C , as well as the correlation




Fig. 7. Dependence of (a)(b) effective temperature T on rest mass m_0 and (c)(d) mean transverse momentum $\langle p_T \rangle$ on mean energy or mean moving mass \bar{m} for (a)(c) Au-Au collisions at 200 GeV and (b)(d) Pb-Pb collisions at 2.76 TeV in different centrality classes. The lines are the results fitted for the values of derived quantities.

between (c) β_T and β_T are presented in Fig. 8. One can see that T_0 and β_T decrease with the increase of C , T_0 increases with the increase of β_T , which renders that central collisions display higher excitation and quicker expansion than peripheral collisions due to more energy deposition in central collisions.

It should be noted that T_0 shown in Fig. 8 is larger than 160–170 MeV which is regarded as the chemical freeze-out temperature T_{ch} of phase transition from hadronic matter to QGP. As expectation, T_0 should be less than or equal to T_{ch} due to time evolution. The difference between our results and T_{ch} is regarded as different “thermometers” (methods) used in the extraction of temperature. In our opinion, an unified “ther-

mometer” (method) should be used in the extraction of temperature. Or, one may find a relation to convert one temperature to another one, as what one did between Celsius Temperature and degree Fahrenheit.

Meanwhile, there is a positive correlation between T_0 and β_T as shown in Fig. 8(c). Some studies show negative correlation between T_0 and β_T when one uses the blast-wave model [25, 26, 27, 28, 29, 30]. At present, one could not decide which correlation is correct. In our opinion, for a given p_T spectrum, T_0 and β_T are negatively correlative if one uses the blast-wave model which gives a large T_0 to correspond to a small β_T . However, for a set of p_T spectra with varying centralities and energies, the situation is dissimilar. The present work uses




Fig. 8. Dependence of (a) kinetic freeze-out temperature T_0 on centrality C and (b) transverse flow velocity β_T on centrality C .

an alternative method to extract T_0 and β_T and obtains a positive correlation.

In addition, one can see some fluctuations in Figs. 3, 4, 5, 6, and 8. These fluctuations have no particular physics meaning, but reflect the statistical and/or systematical fluctuations in the data itself. Although there are fluctuations in the dependences of parameters on centrality, one can see the general decreasing trend of parameters with decreasing the centrality. The considered parameters have similar trend due to their consistent meaning on the energy deposition which is reflected in terms of excitation and expansion degree.

Both the mean transverse momentum $\langle p_T \rangle$ and initial temperature T_i are obtained from p_T spectra. The

relation of $\langle p_T \rangle$ and T_i is certainly positive correlation. Figure 9 shows this correlation. One can see that $\langle p_T \rangle$ increases with the increase of T_i . This correlation is natural due to T_i is defined by $\sqrt{\langle p_T^2 \rangle}/2$. One can also see that $\sqrt{\langle p_T^2 \rangle}/2$ increases with the increase of collision energy and the size of system, but the later has a very small effect, so we can neglect it.

From Figs. 4–9 one can see that the considered quantities in the most peripheral Pb-Pb collisions at 2.76 TeV overlaps the most central Au-Au collisions with lower energy of 200 GeV, which may be the indication of formation of similar systems in the most peripheral collisions at higher energies, and in the most central collisions at lower energies and it may support the hy-




Fig. 9. Dependence of mean transverse momentum $\langle p_T \rangle$ on initial temperature T_i .

pothesis of the effective energy for the particle production [51, 52, 53, 54].

In the considered Au-Au collisions at 200 GeV and Pb-Pb collisions at 2.76 TeV, the decreasing trend of temperatures and kinetic freeze-out volume from central to peripheral collisions renders that more energy deposition and then higher excitation and quicker expansion in central collisions. Some studies [9, 13, 55, 56, 57] which use other methods such as the blast-wave model show that the kinetic freeze-out temperature in central collisions is less than that in peripheral collisions, though this opposite result can be explained as longer freeze-out time in central collisions.

Indeed, the kinetic freeze-out temperature and transverse flow velocity and other quantities are model-dependent. We notice that the current blast-wave model uses a small or almost zero transverse flow velocity in peripheral collisions and obtains a larger kinetic freeze-out temperature in peripheral collisions comparing to that in central collisions. If we use a large transverse flow velocity in peripheral collisions, we can obtain a smaller kinetic freeze-out temperature in peripheral collisions comparing to that in central collisions [58].

In addition, the kinetic freeze-out temperature T_0 and transverse flow velocity β_T are also transverse momentum range dependent. In our opinion, to obtain the parameters as accurately as possible, we should use the transverse momentum range as accurately as possible. The transverse momentum range should not be too narrow or too wide. A too narrow transverse momentum range will exclude the contributions of some particles

which should be included. A too wide transverse momentum range will include the contributions of some particles which should be excluded. In fact, model- and p_T -range-independent T_0 and β_T are ideal.

Generally, the mean transverse momentum $\langle p_T \rangle$ and the root-mean-square transverse momentum $\sqrt{\langle p_T^2 \rangle}$ are model-independent. Obtaining the initial temperature by $T_i = \sqrt{\langle p_T^2 \rangle}/2$ is a suitable treatment which is regardless of model, though it is from the color string percolation model [15, 16, 17]. It is expected that T_0 and β_T are related to $\langle p_T \rangle$, which results in model-independent T_0 and β_T .

As what we did in our recent work [59], let $T_0 \equiv k\langle p_T \rangle/2$ and $\beta_T \equiv (1 - k)\langle p_T \rangle/2\bar{m}$, where k is a parameter which can be approximately taken to be $0.3 - 0.01 \ln(\sqrt{s_{NN}})$ ($\sqrt{s_{NN}}$ is in the units of GeV) [60], $1/2$ is used due to both contributions of projectile and target participants, and \bar{m} denotes the mean energy of the considered particles in the source rest frame. If p_T -range is wide enough, T_0 and β_T are also p_T -range-independent.

If we use T_0 by the above new definition instead of the intercept in the linear relation between T and m_0 , the mean T_0 ($\sim 0.10 - 0.12$ GeV) obtained from Fig. 9 is obviously less than that in the present work which is too large comparing to others. Meanwhile, if we use β_T by the above new definition instead of the slope in the linear relation between $\langle p_T \rangle$ and \bar{m} , the mean β_T ($\sim 0.23 - 0.28$ c) obtained from Fig. 9 is less than those in the present work. Regardless of size, the new definitions of T_0 and β_T are model-independent.

Before summary and conclusions, we would like to point out that this paper fits only the transverse momentum spectra measured from collisions with varying centralities by the two-component standard distribution. Some centrality dependence of related parameters are found. In our recent work [60, 61, 62, 63], we have fitted the spectra measured from collisions with varying energies by the (two-component) standard distribution and/or Tsallis statistics. Some spectra are from small system size and others are from large system size. The related parameters are found to depend also on energy and the larger nucleus in projectile and target nuclei.

In particular, with the increasing energy, the kinetic freeze-out temperature increases quickly from a few GeV to around 10 GeV and then slowly or slightly from around 10 GeV to more than 10 TeV. This implies that around 10 GeV is a special energy at which the interaction mechanism had changed. In fact, the collision system undergone from baryon-dominated to meson-dominated final state [64]. This implies that the critical energy of phase transition from hadronic matter to QGP is possibly existent at around 10 GeV.

The dependence on the larger nucleus in projectile and target nuclei is consistent to the dependence on centrality. This implies possibly that there is a critical size from small to large system, and from peripheral to central collisions. The data measured by the NA61/SHINE Collaboration [65] show that the nucleon number in projectile or target nucleus on the onset of deconfinement is ≈ 10 . Meanwhile, the energy on the onset of deconfinement is ≈ 10 GeV. This double 10 signature is very interesting and should be studied further by various models and methods in future.

We have studied three types of temperatures, namely the effective temperature, initial temperature, and kinetic freeze-out temperature, in this paper. Although the three types of temperatures are extracted from the transverse momentum spectra, they have different physics meanings. The effective temperature is obtained directly from the fit function, which describes together the degree of the thermal motion and flow effect at the stage of kinetic freeze-out. In the case of excluding the contribution of flow effect from the effective temperature, we expect to obtain the kinetic freeze-out temperature which describes only the thermal motion. The initial temperature in this paper is quoted directly from the color string percolation model [15, 16, 17], which is expected to describe the excitation degree of initial state as what we did in our recent work [66].

It is regretful that the chemical freeze-out temperature is not discussed in this paper, though it has wider applications and discussions in literature [67, 68, 69].

The chemical freeze-out temperature describes the excitation degree of collision system at the stage of chemical freeze-out. Generally, the chemical freeze-out temperature can be obtained from the ratio of particle yields, and can be used to map the phase diagram with the chemical potential. In the extensive statistics and/or axiomatic/generic non-extensive statistics [67, 68, 69], one may discuss the chemical and/or kinetic freeze-out parameters systematically.

4 Summary and conclusions

The main observations and conclusions are summarized here.

(a) The transverse momentum spectra of π^- , K^- and \bar{p} at mid-(pseudo)rapidity produced in different centrality events in Au-Au collisions at 200 GeV and Pb-Pb collisions at 2.76 TeV have been analyzed. The experimental data measured by the PHENIX and ALICE Collaborations are fitted by the two-component standard distribution in which the temperature concept is quite close to the ideal gas model.

(b) The effective temperature, initial temperature, kinetic freeze-out temperature, transverse flow velocity and mean transverse momentum increase with the increase of event centrality from peripheral to central collisions, which results in higher excitation degree and quicker expansion velocity in central collisions. The kinetic freeze-out volume increases with the increase of event centrality from peripheral to central collisions due to more number of participant nucleons taking part in central collisions.

(c) The mass dependent differential effective temperature, initial temperature, kinetic freeze-out volume and mean transverse momentum are observed. The kinetic freeze-out temperature and transverse flow velocity extracted in this paper does not show mass dependent differential scenario due to the reason of methodology. Many quantities are model or method dependent.

(d) The formation of similar system is possible in the most peripheral nucleus-nucleus collisions at high energy and in the most central nucleus-nucleus collisions at low energy. This observation confirms the hypothesis of the effective energy for the particle production. There are many similarities in high energy collisions.

Data availability

The data used to support the findings of this study are included within the article and are cited at relevant places within the text as references.

Ethical Approval

The authors declare that they are in compliance with ethical standards regarding the content of this paper.

Disclosure

The funding agencies have no role in the design of the study; in the collection, analysis, or interpretation of the data; in the writing of the manuscript; or in the decision to publish the results.

Conflict of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant Nos. 11575103 and 11947418, the Chinese Government Scholarship (China Scholarship Council), the Scientific and Technological Innovation Programs of Higher Education Institutions in Shanxi (STIP) under Grant No. 201802017, the Shanxi Provincial Natural Science Foundation under Grant No. 201901D111043, and the Fund for Shanxi “1331 Project” Key Subjects Construction.

References

- [1] M. E. Fisher and A. N. Berker, “Scaling for first-order transitions in thermodynamic and finite systems,” *Physics Letters B*, vol. 26, pp. 2507–2513, 1982.
- [2] E. D. Davis and H. G. Miller, ““Phase transitions” at finite temperature in finite systems,” *Physics Letters B*, vol. 196, pp. 277–280, 1987.
- [3] S. Mukherjee and V. Skokov, “Universality driven analytic structure of QCD crossover: Radius of convergence in baryon chemical potential,” arXiv:1909.04639 [hep-ph].
- [4] W.-J. Fu, J. M. Pawłowski, and F. Rennecke, “The QCD phase structure at finite temperature and density,” arXiv:1909.02991 [hep-ph].
- [5] G.-Y. Shao, W.-B. He, and X.-Y. Gao, “Deformed QCD phase structure and entropy oscillation in the presence of a magnetic background,” *Physical Review D*, vol. 100, Article ID 014020, 2019.
- [6] R. Sahoo, “Possible formation of QGP-droplets in proton-proton collisions at the CERN Large Hadron Collider,” *AAPPS Bulletin*, vol. 29, no. 4, pp. 16–21, 2019.
- [7] M. Gyulassy and L. McLerran, “New forms of QCD matter discovered at RHIC,” *Nuclear Physics A*, vol. 750, pp. 30–63, 2005.
- [8] M. T. AlFiky, O. T. ElSherif, and A. M. Hamed, “Onset of the jet quenching phenomenon,” *JETP Letters*, vol. 111, pp. 8–17, 2020.
- [9] STAR Collab. (L. Adamczyk *et al.*), “Bulk properties of the medium produced in relativistic heavy ion collisions from the Beam Energy Scan program,” *Physical Review C*, vol. 96, Article ID 044904, 2017.
- [10] ALICE Collab. (B. B. Abelev *et al.*), “Multiplicity dependence of pion, kaon, proton and Lambda production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Physics Letters B*, vol. 728, pp. 25–38, 2014.
- [11] ALICE Collab. (S. Acharya *et al.*), “Anisotropic flow in Xe-Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV,” *Physics Letters B*, vol. 784, pp. 82–95, 2018.
- [12] L. Van Hove, “Two problems concerning hot hadronic matter and high-energy collisions (Equilibrium formation, plasma deflagration),” *Zeitschrift für Physik C Particles and Fields*, vol. 21, pp. 93–98, 1983.
- [13] S. Chatterjee, S. Das, L. Kumar, D. Mishra, B. Mohanty, R. Sahoo, and N. Sharma, “Freeze-out parameters in heavy-ion collisions at AGS, SPS, RHIC, and LHC energies,” *Advances in High Energy Physics*, vol. 2015, Article ID 349013, 2015.
- [14] S. Uddin, I. Bashir, and R. A. Bhat, “Transverse momentum distributions of hadrons produced in Pb-Pb collisions at LHC energy $\sqrt{s_{NN}} = 2.76$ TeV,” *Advances in High Energy Physics*, vol. 2015, Article ID 154853, 2015.
- [15] L. J. Gutay, A. S. Hirsch, C. Pajares, R. P. Scharenberg, and B. K. Srivastava, “De-confinement in small systems: Clustering of color sources in high multiplicity $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV,” *International Journal of Modern Physics E*, vol. 24, Article ID 1550101, 2015.
- [16] A. S. Hirsch, C. Pajares, R. P. Scharenberg, and B. K. Srivastava, “De-confinement in high multiplicity proton-proton collisions at LHC energies,” *Physical Review D*, vol. 100, Article ID 114040, 2019.
- [17] P. Sahoo, S. De, S. K. Tiwari, and R. Sahoo, “Energy and centrality dependent study of deconfinement phase transition in a color string percolation approach at RHIC energies,” *The European Physical Journal A*, vol. 54, article 136, 2018.
- [18] D. H. E. Gross, “Microcanonical thermodynamics and statistical fragmentation of dissipative systems: The topological structure of the N-body phase space,” *Physics Reports*, vol. 279, pp. 119–201, 1997.
- [19] B. Borderie, “Dynamics and thermodynamics of the liquid-gas phase transition in hot nuclei studied with the INDRA array,” *Journal of Physics G*, vol. 28, p. R217–R247, 2002.
- [20] M. D’Agostino, F. Gulminelli, Ph. Chomaz, M. Bruno, F. Cannata, R. Bougault, N. Colonna, F. Gramegna, I. Iori, N. Le Neindre, G. V. Margagliotti, P. F. Mastinu, P. M. Milazzo, A. Moroni, and G. Vannini, “Negative heat capacity in the critical region of nuclear fragmentation: An experimental evidence of the liquid-gas phase

- transition,” *Physics Letters B*, vol. 473, pp. 219–225, 2000.
- [21] M. D’Agostino, R. Bougault, F. Gulminelli, M. Bruno, F. Cannata, Ph. Chomaz, F. Gramegna, I. Iori, N. Le Neindre, G. V. Margagliotti, A. Moroni, and G. Vanini, “On the reliability of negative heat capacity measurements,” *Nuclear Physics A*, vol. 699, pp. 795–818, 2002.
- [22] P. Chomaz, V. Duflot, and F. Gulminelli, “Caloric curves and energy fluctuations in the microcanonical liquid-gas phase transition,” *Physical Review Letters*, vol. 85, pp. 3587–3590, 2000.
- [23] R. Hagedorn, “Multiplicities, p_T distributions and the expected hadron \rightarrow quark-gluon phase transition,” *La Rivista del Nuovo Cimento*, vol. 6, no. 10, pp. 1–50, 1983.
- [24] J. Cleymans and D. Worku, “Relativistic thermodynamics: Transverse momentum distributions in high-energy physics,” *The European Physical Journal A*, vol. 48, article 160, 2012.
- [25] E. Schnedermann, J. Sollfrank, and U. Heinz, “Thermal phenomenology of hadrons from 200 A GeV S+S collisions,” *Physical Review C*, vol. 48, pp. 2462–2475, 1993.
- [26] STAR Collab. (B. I. Abelev *et al.*), “Systematic measurements of identified particle spectra in pp, d+Au and Au+Au collisions from STAR,” *Physical Review C*, vol. 79, Article ID, 034909, 2009.
- [27] STAR Collab. (B. I. Abelev *et al.*), “Identified particle production, azimuthal anisotropy, and interferometry measurements in Au+Au collisions at $\sqrt{s_{NN}} = 9.2$ GeV,” *Physical Review C*, vol. 81, Article ID 024911, 2010.
- [28] Z. B. Tang, Y. C. Xu, L. J. Ruan, G. van Buren, F. Q. Wang, and Z. B. Xu, “Spectra and radial flow in relativistic heavy ion collisions with Tsallis statistics in a blast-wave description,” *Physical Review C*, vol. 79, Article ID 051901(R), 2009.
- [29] Z. B. Tang, L. J. Ruan, M. Shao, H. F. Chen, C. Li, B. Mohanty, P. Sorensen, A. H. Tang, and Z. B. Xu, “Statistical origin of constituent-quark scaling in the QGP hadronization,” *Chinese Physics Letters*, vol. 30, Article ID 031201, 2013.
- [30] K. Jiang, Y. Y. Zhu, W. T. Liu, H. F. Chen, C. Li, L. J. Ruan, M. Shao, Z. B. Tang, and Z. B. Xu, “Onset of radial flow in p+p collisions,” *Physical Review C*, vol. 91, Article ID 024910, 2015.
- [31] J. Cleymans, H. Oeschler, K. Redlich, and S. Wheaton, “Comparison of chemical freeze-out criteria in heavy-ion collisions,” *Physical Review C*, vol. 73, Article ID 034905, 2006.
- [32] A. Andronic, P. Braun-Munzinger, and J. Stachel, “Hadron production in central nucleus-nucleus collisions at chemical freeze-out,” *Nuclear Physics A*, vol. 772, pp. 167–199, 2006.
- [33] S. Uddin, J. S. Ahmad, W. Bashir, and R. A. Bhat, “A unified approach towards describing rapidity and transverse momentum distributions in thermal freeze-out model,” *Journal of Physics G*, vol. 39, Article ID 015012, 2012.
- [34] R. P. Adak, S. Das, S. K. Ghosh, R. Ray, and S. Samanta, “Centrality dependence of chemical freeze-out parameters from net-proton and net-charge fluctuations using a hadron resonance gas model,” *Physical Review C*, vol. 96, Article ID 014902, 2017.
- [35] R. Odorico, “Does a transverse energy trigger actually trigger on large p_T jets?,” *Physics Letters B*, vol. 118, pp. 151–154, 1982.
- [36] ALICE Collab. (K. Aamodt *et al.*), “Transverse momentum spectra of charged particles in proton-proton collisions at $\sqrt{s} = 900$ GeV with ALICE at the LHC,” *Physics Letters B*, vol. 693, pp. 53–68, 2010.
- [37] M. Biyajima, T. Mizoguchi, and N. Suzuki, “Analyses of whole transverse momentum distributions in $p\bar{p}$ and pp collisions by using a modified version of Hagedorn’s formula,” *International Journal of Modern Physics A*, vol. 32, Article ID 1750057, 2017.
- [38] PHENIX Collab. (S. S. Adler *et al.*), “Identified charged particle spectra and yields in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Physical Review C*, vol. 69, Article ID 034909, 2004.
- [39] ALICE Collab. (B. Abelev *et al.*), “Centrality dependence of π , K , and p production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Physical Review C*, vol. 88, Article ID 044910, 2013.
- [40] D. Thakur, S. Tripathy, P. Garg, R. Sahoo and J. Cleymans, “Indication of differential kinetic freeze-out at RHIC and LHC energies,” *Acta Physica Polonica B Proceedings Supplement*, vol. 9, pp. 329–332, 2016.
- [41] D. Thakur, S. Tripathy, P. Garg, R. Sahoo, and J. Cleymans, “Indication of a differential freeze-out in proton-proton and heavy-ion collisions at RHIC and LHC energies,” *Advances in High Energy Physics*, vol. 2016, Article ID 4149352, 2016.
- [42] PHENIX Collab. (K. Adcox *et al.*), “Centrality dependence of π^\pm , K^\pm , p and \bar{p} production from $\sqrt{s_{NN}} = 13$ GeV Au+Au collisions at RHIC,” *Physical Review Letters*, vol. 88, Article ID 242301, 2002.
- [43] UA1 Collab. (C. Albajar *et al.*), “A study of the general characteristics of $p\bar{p}$ collisions at $\sqrt{s} = 0.2$ to 0.9 TeV,” *Nuclear Physics B*, vol. 335, pp. 261–287, 1990.
- [44] E802 Collab. (Y. Akiba *et al.*), “Particle production in Au+Au collisions from BNL E866,” *Nuclear Physics A*, vol. 610, pp. 139–152, 1996.
- [45] NA44 Collab. (I. G. Bearden *et al.*), “Collective expansion in high-energy heavy ion collisions,” *Physical Review Letters*, vol. 78, pp. 2080–2083, 1997.
- [46] P.-P. Yang, Q. Wang and F.-H. Liu, “Mutual derivation between arbitrary distribution forms of momenta and momentum components,” *International Journal of Theoretical Physics*, vol. 58, pp. 2603–2618, 2019.
- [47] STAR Collab. (B. I. Abelev *et al.*), “Systematic measurements of identified particle spectra in pp, d+Au,

- and Au+Au collisions at the STAR detector,” *Physical Review C*, vol. 79, Article ID 034909, 2009.
- [48] S. Takeuchi, K. Murase, T. Hirano, P. Huovinen and Y. Nara, “Effects of hadronic rescattering on multistrange hadrons in high-energy nuclear collisions,” *Physical Review C*, vol. 92, Article ID 044907, 2015.
- [49] H.-R. Wei, F.-H. Liu, and R. A. Lacey, “Disentangling random thermal motion of particles and collective expansion of source from transverse momentum spectra in high energy collisions,” *Journal of Physical G*, vol. 43, Article ID 125102, 2016.
- [50] H.-R. Wei, F.-H. Liu, and R. A. Lacey, “Kinetic freeze-out temperature and flow velocity extracted from transverse momentum spectra of final-state light flavor particles produced in collisions at RHIC and LHC,” *The European Physical Journal A*, vol. 52, article 102, 2016.
- [51] E. K. G. Sarkisyan and A. S. Sakharov, “Relating multihadron production in hadronic and nuclear collisions,” *The European Physical Journal C*, vol. 70, article 533, 2010.
- [52] A. N. Mishra, R. Sahoo, E. K. G. Sarkisyan and A. S. Sakharov, “Effective-energy budget in multiparticle production in nuclear collisions,” *The European Physical Journal C*, vol. 74, article 3147, 2014 and “Erratum,” *The European Physical Journal C*, vol. 75, article 70, 2015.
- [53] E. K. Sarkisyan-Grinbaum, A. Nath Mishra, R. Sahoo, and A. S. Sakharov, “Effective-energy universality approach describing total multiplicity centrality dependence in heavy-ion collisions,” *EPL*, vol. 127, article 62001, 2019.
- [54] E. K. G. Sarkisyan, A. N. Mishra, R. Sahoo, and A. S. Sakharov, “Multihadron production dynamics exploring the energy balance in hadronic and nuclear collisions,” *Physical Review D*, vol. 93, Article ID 054046, 2016 and “Addendum,” *Physical Review D*, vol. 93, Article ID 079904, 2016.
- [55] D. Prorok, “Centrality dependence of freeze-out temperature fluctuations in Pb-Pb collisions at the LHC,” *The European Physical Journal A*, vol. 55, article 37, 2019.
- [56] B. De, “Non-extensive statistics and understanding particle production and kinetic freeze-out process from p_T -spectra at 2.76 TeV,” *The European Physical Journal A*, vol. 50, article 138, 2014.
- [57] R. Pasechnik and M. Šumbera, “Phenomenological review on quark-gluon plasma: Concepts vs. observations,” *Universe*, vol. 3, article 7, 2017.
- [58] H.-L. Lao, F.-H. Liu, B.-C. Li and M.-Y. Duan, “Kinetic freeze-out temperatures in central and peripheral collisions: Which one is larger?,” *Nuclear Science and Techniques*, vol. 29, article 82, 2018.
- [59] M. Waqas and F.-H. Liu, “Initial, effective, and kinetic freeze-out temperatures from transverse momentum spectra in high-energy proton(deuteron)Cnucleus and nucleusCnucleus collisions,” *The European Physical Journal Plus*, vol. 135, article 147, 2020.
- [60] L.-L. Li, F.-H. Liu, M. Waqas, R. Al-Yusufi and A. Mujear, “Excitation functions of related parameters from transverse momentum (mass) spectra in high energy collisions,” arXiv:1911.07419 [hep-ph] (2019).
- [61] L.-L. Li and F.-H. Liu, “Energy dependent kinetic freeze-out temperature and transverse flow velocity in high energy collisions,” *The European Physical Journal A*, vol. 54, article 169, 2018.
- [62] L.-L. Li and F.-H. Liu, “Excitation functions of kinetic freeze-out temperature and transverse flow velocity in proton-proton collisions,” arXiv:1805.03342 [hep-ph].
- [63] H.-L. Lao, F.-H. Liu, B.-C. Li, M.-Y. Duan, and R. A. Lacey, “Examining the model dependence of the determination of kinetic freeze-out temperature and transverse flow velocity in small collision system,” *Nuclear Science and Techniques*, vol. 29, article 164, 2018.
- [64] J. Cleymans, “The physics case for the $\sqrt{s_{NN}} \approx 10$ GeV energy region,” in Walter Greiner Memorial Volume, edited by P. O. Hess (World Scientific, Singapore, 2018), arXiv:1711.02882 [hep-ph].
- [65] M. P. Lewicki and L. Turko (On behalf of NA61/SHINE Collaboration), “NA61/SHINE shining more light on the onset of deconfinement,” *CERN EP Newsletter*, pp. 67–72, Dec. 2019. arXiv:2002.00631 [hep-ex].
- [66] Q. Wang and F.-H. Liu, “Initial and final state temperatures of antiproton emission sources in high energy collisions,” *International Journal of Theoretical Physics* vol. 58, pp. 4119–4138, 2019.
- [67] A. N. Tawfik, H. Yassin, and E. R. Abo Elyazeed, “Extensive/nonextensive statistics for p_T distributions of various charged particles produced in p+p and A+A collisions in a wide range of energies,” arXiv:1905.12756 [hep-ph].
- [68] A. N. Tawfik, “Axiomatic nonextensive statistics at NICA energies,” *The European Physical Journal A*, vol. 52, article 253, 2016.
- [69] A. N. Tawfik, H. Yassin, and E. R. Abo Elyazeed, “Chemical freezeout parameters within generic nonextensive statistics,” *Indian Journal of Physics*, vol. 92, pp. 1325–1335, 2018.