

PROJECTIVE MODULES AND THE HOMOTOPY CLASSIFICATION OF (G, n) -COMPLEXES

JOHN NICHOLSON

ABSTRACT. A (G, n) -complex is an n -dimensional CW-complex with fundamental group G and whose universal cover is $(n - 1)$ -connected. If G has periodic cohomology then, for appropriate n , we show that there is a one-to-one correspondence between the homotopy types of finite (G, n) -complexes and the orbits of the stable class of a certain projective $\mathbb{Z}G$ -module under the action of $\text{Aut}(G)$. We develop techniques to compute this action explicitly and use this to give an example where the action is non-trivial.

1. INTRODUCTION

For a group G and $n \geq 2$, a (G, n) -complex is a connected n -dimensional CW-complex X for which $\pi_1(X) \cong G$ and \tilde{X} is $(n - 1)$ -connected. For example, a finite $(G, 2)$ -complex is equivalently a finite 2-complex X with $\pi_1(X) \cong G$. An example of a finite $(G, 3)$ -complex is a closed 3-manifold M with $\pi_1(M) \cong G$ finite. Given a group G and $n \geq 2$, a finite (G, n) -complex exists if and only if G is finitely presented and has type FP_n (see [4, p6] and Proposition 5.1).

Let $\text{HT}(G, n)$ be the set of homotopy types of finite (G, n) -complexes, which can be viewed as a graph with edges between each X and $X \vee S^n$. It is well known that $\text{HT}(G, n)$ is a tree [48], i.e. a connected acyclic graph, and has a grading coming from $(-1)^n \chi(X)$ which takes a minimum value $\chi_{\min}(G, n)$. The problem of determining the structure of $\text{HT}(G, n)$ as a tree has a long history which dates back to Cockcroft-Swan [8] and Dyer-Sieradski [15, 16].

In the case of finite abelian groups, the structure of $\text{HT}(G, n)$ has been classified through a series of articles by Metzler [30], Dyer-Sieradski [38], Browning [5] and Linnell [25]. However, much less is known for non-abelian groups and an important class of examples are the groups with k -periodic cohomology, i.e. where the Tate cohomology groups satisfy $\hat{H}^i(G; \mathbb{Z}) \cong \hat{H}^{i+k}(G; \mathbb{Z})$ for all $i \in \mathbb{Z}$. For example, if G is finite and n is even, then it was shown by Browning [6] that $\chi(X) = \chi(Y)$ implies $X \vee S^n \simeq Y \vee S^n$ (see also [19]). However, when n is odd, this is known only when G does not have k -periodic cohomology for $k \mid n + 1$ (see Question 7.4).

The aim of this article is to make new progress towards the classification over groups with periodic cohomology, building upon work of Dyer [13] and Johnson [23].

1.1. Main results. Let $\text{PHT}(G, n)$ be the tree of polarised homotopy types of finite (G, n) -complexes, i.e. the homotopy types of pairs (X, ρ) where $\rho : \pi_1(X) \cong G$.

Let $C(\mathbb{Z}G)$ denote the projective class group, i.e. the equivalence classes of finitely generated projective $\mathbb{Z}G$ -modules where $P \sim Q$ if $P \oplus \mathbb{Z}G^i \cong Q \oplus \mathbb{Z}G^j$ for some i, j . Note that a class $[P] \in C(\mathbb{Z}G)$ can be viewed as the set of (non-zero) projective $\mathbb{Z}G$ -modules P_0 for which $P_0 \sim P$, and this has the structure of a graded

tree with edges between each P_0 and $P_0 \oplus \mathbb{Z}G$. If G has k -periodic cohomology, then the *Swan finiteness obstruction* is an element $\sigma_k(G) \in C(\mathbb{Z}G)/T_G$ which vanishes if and only if there exists a finite CW-complex X with $\pi_1(X) \cong G$ and $\tilde{X} \simeq S^{k-1}$.

Recall that G has the *D2 property* if every cohomologically 2-dimensional finite complex X with $\pi_1(X) \cong G$ is homotopy equivalent to a finite 2-complex.

Theorem A. *Let G have k -periodic cohomology and let $n = ik$ or $ik - 2$ for some $i \geq 1$. Then there is an injective map of graded trees*

$$\Psi : \text{PHT}(G, n) \rightarrow [P_{(G, n)}]$$

for any projective $\mathbb{Z}G$ -module $P_{(G, n)}$ with $\sigma_{ik}(G) = [P_{(G, n)}] \in C(\mathbb{Z}G)/T_G$. Furthermore, Ψ is bijective if and only if $n \geq 3$ or if $n = 2$ and G has the D2 property.

Remark 1.1. (a) If G satisfies the Eichler condition, then $[P_{(G, n)}]$ has *cancellation* in the sense that $P_1 \oplus \mathbb{Z}G \cong P_2 \oplus \mathbb{Z}G$ implies $P_1 \cong P_2$ for all $P_1, P_2 \in [P_{(G, n)}]$ (see [21]). This implies that $\text{PHT}(G, n)$ and $\text{HT}(G, n)$ have cancellation in the sense that $X \vee S^n \simeq Y \vee S^n$ implies $X \simeq Y$, and recovers the main result of Dyer [13]. (b) An equivalent statement appeared in [23] in the case $n = 2$, though the proof contained a small gap which was patched up in [36] using a theorem of Browning [6].

Our proof is based on the work of Hambleton-Kreck [19] and is independent of [6, 23]. After establishing preliminaries in Sections 2 and 3, we will prove general cancellation theorems for chain complexes of projective modules in Section 4. This suffices to prove Theorem A due to the correspondence between $\text{PHT}(G, n)$ and the tree of algebraic n -complexes (see Proposition 5.1). In Theorem 5.3, we give a detailed version of Theorem A which contains an explicit description of the map Ψ .

We then use of this description of Ψ to determine the induced action of $\text{Aut}(G)$ on $[P_{(G, n)}]$ via the bijection $\text{HT}(G, n) \cong \text{PHT}(G, n)/\text{Aut}(G)$. To state the induced action, consider the following two operations for M a (left) projective $\mathbb{Z}G$ -module:

- (1) If $\theta \in \text{Aut}(G)$, then let M_θ be the $\mathbb{Z}G$ -module whose abelian group is that of M but with action $g \cdot x = \theta(g)x$ for $g \in G, x \in M$ (see Lemma 6.1).
- (2) If r represents a class in $(\mathbb{Z}/|G|)^\times$ and $I \subseteq \mathbb{Z}G$ is the augmentation ideal, then (I, r) is a projective $\mathbb{Z}G$ -module. The tensor product $(I, r) \otimes M$ is a projective $\mathbb{Z}G$ -module since (I, r) is a two-sided ideal (see Lemma 3.5).

In Section 6, we will prove the following which is the main result of this article.

Theorem B. *Let G have k -periodic cohomology and let $n = ik$ or $ik - 2$ for some $i \geq 1$. Then Ψ induces an injective map of graded trees*

$$\bar{\Psi} : \text{HT}(G, n) \rightarrow [P_{(G, n)}]/\text{Aut}(G),$$

where the action by $\theta \in \text{Aut}(G)$ is given by $\theta : P \mapsto (I, \psi_k(\theta)^i) \otimes P_\theta$ for some map $\psi_k : \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$ which depends only on G and k . Furthermore, $\bar{\Psi}$ is bijective if and only if $n \geq 3$ or if $n = 2$ and G has the D2 property.

This reduces the problem of determining when cancellation occurs in the homotopy trees to the purely algebraic problem of determining cancellation for $[P]$ and $[P]/\text{Aut}(G)$ which will be dealt with in [33].

1.2. Computing the action of $\text{Aut}(G)$. The rest of this article will be devoted to exploring the action of $\text{Aut}(G)$ on $[P_{(G, n)}]$. This includes establishing some general theory in preparation for the more detailed computations in [33].

Firstly, and perhaps somewhat surprisingly, we could find no example where the $\text{Aut}(G)$ -action described in Theorem B does not take the form of the simpler action $P \mapsto P_\theta$. In all examples computed, we had $(I, \psi_k(\theta)) \cong \mathbb{Z}G$ which implies $(I, \psi_k(\theta)^i) \cong \mathbb{Z}G$ and so $\theta(P) \cong P_\theta$. We therefore ask the following:

Question 7.3. *Does there exist G with k -periodic cohomology and $\theta \in \text{Aut}(G)$ for which $(I, \psi_k(\theta))$ is not free?*

There are two approaches to finding examples where $(I, \psi_k(\theta))$ is not free. The first is to find an example where $(I, \psi_k(\theta))$ is not even stably free. It was shown by Dyer [13, p276] and Davis [12] that $(I, \psi_k(\theta))$ is stably free when $\sigma_k(G) = 0$. Davis asked whether this also holds when $\sigma_k(G) \neq 0$ [12, p488]. The second approach is to find an example where $(I, \psi_k(\theta))$ is stably free but not free. This is likely to be difficult since the general question of whether (I, r) can be stably free but not free is still open and dates back to Wall's 1979 Problems List [45, Problem A5].

In Section 8, we develop a general method to compute the action $P \mapsto P_\theta$. We will then use this to give the following example where the action is non-trivial. Let Q_{4n} denote the quaternion group of order $4n$, which has 4-periodic cohomology. Since $\sigma_4(Q_{4n}) = 0$, we can take $[P_{(Q_{4n}, 2)}] = [\mathbb{Z}Q_{4n}] = \bigcup_{r \geq 1} \text{SF}_r(\mathbb{Z}Q_{4n})$ where $\text{SF}_r(\mathbb{Z}Q_{4n})$ is the set of stably free $\mathbb{Z}Q_{4n}$ -modules of rank $r \geq 1$. As above, let $\theta \in \text{Aut}(Q_{4n})$ act on $[\mathbb{Z}Q_{4n}]$ by $\theta : P \mapsto (I, \psi_4(\theta)^i) \otimes P_\theta$ for some $i \geq 1$. We show:

Theorem 9.1. *$\text{Aut}(Q_{24})$ acts non-trivially on $[\mathbb{Z}Q_{24}]$. More specifically, we have $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{24}) / \text{Aut}(Q_{24})| = 2$.*

This is in contrast to the case Q_{4n} for $2 \leq n \leq 5$ where $|\text{SF}_1(\mathbb{Z}Q_{4n})| = 1$ and the case Q_{28} where $|\text{SF}_1(\mathbb{Z}Q_{28})| = |\text{SF}_1(\mathbb{Z}Q_{28}) / \text{Aut}(Q_{28})| = 2$ (see Figure 2).

1.3. Overview of the wider project. This article is the first of a two-part series (followed by [33]) in which we explore the classification of finite (G, n) -complexes over groups with periodic cohomology. These results are motivated by the following.

Wall's D2 problem for groups with 4-periodic cohomology. In the language above, the D2 problem asks whether every finitely presented group G has the D2 property. This dates back to Wall's 1965 paper on finiteness conditions [44] and is currently open. The case where G has 4-periodic cohomology was proposed to contain a counterexample to the D2 problem in 1977 [9], and has since been studied extensively. In this case, Johnson proved Theorem A when $n = 2$ and, using results of Swan [43], he established the D2 property for many new groups [23]. In [34, 36], we extended these results and determined when $\text{PHT}(G, 2)$ has cancellation.

In the case where $\text{PHT}(G, 2)$ has non-cancellation, the D2 property has only been proven for Q_{28} (see [29, 36]). This motivated Theorem B in the case $n = 2$ since one imagines it might be easier to prove that $\bar{\Psi}$ is bijective rather than Ψ . The question of when $\text{HT}(G, 2)$ has cancellation is answered in [33, Theorem A].

Stable and unstable classification of manifolds. If X is a finite (G, n) -complex, then there exists an embedding $i : X \hookrightarrow \mathbb{R}^{2n+1}$. The boundary of a smooth regular neighbourhood of i gives a smooth closed $2n$ -manifold $M(X)$. If X is determined up to simple homotopy, then $M(X)$ is well-defined up to s -cobordism (which coincides with homeomorphism for reasonable G) and $M(X \vee S^n) \cong M(X) \# (S^n \times S^n)$.

In 1984, Kreck-Schafer used this to construct smooth closed $4n$ -manifolds M_1 and M_2 for every $n \geq 1$ such that $M_1 \# (S^{2n} \times S^{2n}) \cong M_2 \# (S^{2n} \times S^{2n})$ are diffeomorphic but $M_1 \not\cong M_2$ [24]. Their examples have the form $M(X_i)$ where the $X_i \in \text{HT}(G, n)$ are the non-cancellation examples for G abelian found by Metzler, Dyer and Sieradski [30, 37, 38]. Recently, Conway-Crowley-Powell-Sixt constructed examples of both simply-connected M_i [1] and infinitely many M_i [2] for all $n \geq 2$. However, the examples of Kreck-Schafer remain the only known examples in dimension 4. In classifying $\text{HT}(G, n)$ when G has periodic cohomology, we hope to create a second family of examples both in dimension 4 and in higher dimensions.

2. EXTENSIONS OF MODULES

Throughout this section R will be a (unital) ring. From now on, we will assume without further mention that all R -modules are finitely generated left R -modules.

For R -modules A and B , define $\text{Ext}_R^n(A, B)$ to be the set of exact sequences

$$E = (0 \rightarrow B \xrightarrow{i} E_{n-1} \xrightarrow{\partial_{n-1}} E_{n-2} \xrightarrow{\partial_{n-2}} \cdots \xrightarrow{\partial_2} E_1 \xrightarrow{\partial_1} E_0 \xrightarrow{\varepsilon} A \rightarrow 0)$$

for R -modules E_i considered up to congruence, i.e. the equivalence relation generated by *elementary congruences* which are chain maps of the form

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \begin{pmatrix} 0 \rightarrow B \rightarrow E_{n-1} \rightarrow \cdots \rightarrow E_0 \rightarrow A \rightarrow 0 \\ \downarrow \text{id} \quad \downarrow \varphi_{n-1} \quad \quad \downarrow \varphi_0 \quad \downarrow \text{id} \\ 0 \rightarrow B \rightarrow E'_{n-1} \rightarrow \cdots \rightarrow E'_0 \rightarrow A \rightarrow 0 \end{pmatrix}.$$

That is, two extensions E and E' are *congruent* if there exists extensions $E^{(i)}$ for $0 \leq i \leq n$ such that $E = E^{(0)}$, $E' = E^{(n)}$ and, for $i \leq n-1$, there exists an elementary congruence of the form $\varphi : E^{(i)} \rightarrow E^{(i+1)}$ or $\varphi : E^{(i+1)} \rightarrow E^{(i)}$.

We write extensions in $\text{Ext}_R^n(A, B)$ as $E = (E_*, \partial_*)$ where the maps $i : B \rightarrow E_{n-1}$ and $\varepsilon : E_0 \rightarrow A$ are understood. We will often write $\partial_i = \partial_i^E$, $i = i_E$ and $\varepsilon = \varepsilon_E$ when the need arises to distinguish different extensions.

This is an abelian group under Baer sum, and coincides with the usual definition of $\text{Ext}_R^n(A, B)$ [47, Section 3.4]. We will assume familiarity with the basic operations on extensions such as pullback, pushout and Yoneda product [23, Section 24].

Worth emphasising however is the operation of stabilisation. If $E = (E_*, \partial_*) \in \text{Ext}_R^n(A, B)$, then define the *stabilised complex* $E \oplus R \in \text{Ext}_R^n(A, B \oplus R)$ by

$$E \oplus R = (0 \rightarrow B \oplus R \xrightarrow{\cdot \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} E_{n-1} \oplus R \xrightarrow{\cdot \begin{pmatrix} \partial_{n-1} \\ 0 \end{pmatrix}} E_{n-2} \rightarrow \cdots \rightarrow E_0 \rightarrow A \rightarrow 0).$$

This gives a well-defined map of abelian groups

$$- \oplus R : \text{Ext}_R^n(A, B) \rightarrow \text{Ext}_R^n(A, B \oplus R).$$

Let $\text{Proj}_R^n(A, B)$ denote the subset of $\text{Ext}_R^n(A, B)$ consisting of extensions (P_*, ∂_*) with the P_i projective. This is closed under Baer sum, and so is a subgroup, and is also preserved by pullbacks, pushouts, Yoneda product and stabilisation. The following is a consequence of the co-cycle description of Ext [46, Lemma 1.1].

Lemma 2.1 (Shifting). *If A, B, C, D are R -modules, $E \in \text{Proj}_R^k(B, C)$ and $k, n, m \geq 1$, then Yoneda product induces bijections*

$$- \circ E : \text{Ext}_R^n(C, D) \rightarrow \text{Ext}_R^{n+k}(B, D), \quad E \circ - : \text{Ext}_R^m(A, B) \rightarrow \text{Ext}_R^{m+k}(A, C).$$

This can be viewed as a sort of cancellation theorem for extensions up to congruence in the sense that $F \circ E \cong F' \circ E$ or $E \circ F \cong E \circ F'$ implies that $F \cong F'$.

A simple consequence of this is the following lemma. This can be interpreted as a kind of duality theorem for projective extensions.

Lemma 2.2 (Duality). *If A, B, C are R -modules, $F \in \text{Proj}_R^k(A, C)$ and $k > n \geq 1$, then there are bijections*

$$\begin{aligned} \Psi_F : \text{Proj}_R^n(A, B) &\rightarrow \text{Proj}_R^{k-n}(B, C), & \Psi_F^{-1} : \text{Proj}_R^{k-n}(B, C) &\rightarrow \text{Proj}_R^n(A, B). \\ E &\mapsto (- \circ E)^{-1}(F) & E' &\mapsto (E' \circ -)^{-1}(F) \end{aligned}$$

We now turn our attention to an equivalence relation on $\text{Ext}_R^n(A, B)$ which is weaker than congruence. For R -modules A, B and $E, E' \in \text{Ext}_R^n(A, B)$, a chain map $\varphi : E \rightarrow E'$ is said to be a *chain homotopy equivalence* if the restriction to the un-augmented chain complexes $\varphi : (E_*, \partial_*)_{0 \leq * < n} \rightarrow (E'_*, \partial'_*)_{0 \leq * < n}$ is a chain homotopy equivalence.

If $E, E' \in \text{Proj}_R^n(A, B)$ then, since a chain map between projective chain complexes is a chain homotopy equivalence if and only if it is a homology equivalence [23, Theorem 46.6], a chain homotopy equivalence $\varphi : E \rightarrow E'$ can equivalently be defined as a chain map of the form

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \begin{pmatrix} 0 \rightarrow B \rightarrow P_{n-1} \rightarrow \cdots \rightarrow P_0 \rightarrow A \rightarrow 0 \\ \quad \downarrow \varphi_B \quad \downarrow \varphi_{n-1} \quad \quad \downarrow \varphi_0 \quad \downarrow \varphi_A \\ 0 \rightarrow B \rightarrow P'_{n-1} \rightarrow \cdots \rightarrow P'_0 \rightarrow A \rightarrow 0 \end{pmatrix}$$

where φ_A and φ_B are R -module isomorphisms, which we will often abbreviate to $\varphi = (\varphi_B, \varphi_{n-1}, \dots, \varphi_0, \varphi_A)$. It follows easily that a congruence is a chain homotopy equivalence. We define $\text{hProj}_R^n(A, B)$ to be set of equivalence classes in $\text{Proj}_R^n(A, B)$ up to chain homotopy equivalences, which is an abelian group under Baer sum.

For special choices of modules, the shifting lemma and the duality lemma also hold for chain homotopy equivalences. We define \mathbb{Z} to be the R -module with underlying abelian group \mathbb{Z} and trivial R -action, i.e. $r \cdot n = n$ for all $r \in R, n \in \mathbb{Z}$.

Lemma 2.3 (Shifting). *If A, B are R -modules, $F \in \text{Proj}_R^k(\mathbb{Z}, \mathbb{Z})$ and $n, m, k \geq 1$, then Yoneda product induces bijections*

$$- \circ F : \text{hProj}_R^n(\mathbb{Z}, A) \rightarrow \text{hProj}_R^{n+k}(\mathbb{Z}, A), \quad F \circ - : \text{hProj}_R^m(B, \mathbb{Z}) \rightarrow \text{hProj}_R^{m+k}(B, \mathbb{Z}).$$

Proof. Firstly note that $- \circ F$ induces maps on the chain homotopy classes by extending the map to $\pm \text{id}$ on F . This is necessarily surjective. To see that it is injective, suppose that there is a chain homotopy equivalence $\varphi : E_1 \circ F \rightarrow E_2 \circ F$. By considering $-\varphi$ if necessary, we can assume that $\varphi_{\mathbb{Z}} = \text{id}$, so that

$$E_2 \circ F \cong (\varphi_A)_*(E_1 \circ F) = (\varphi_A)_*(E_1) \circ F.$$

By Lemma 2.1, this implies that $E_2 \cong (\varphi_A)_*(E_1)$ and so $E_1 \simeq E_2$ as required. \square

The proof of the duality lemma in this setting is similar as so will be omitted.

Lemma 2.4 (Duality). *If A is an R -module, $F \in \text{Proj}_R^k(\mathbb{Z}, \mathbb{Z})$ and $k > n \geq 1$, then there are bijections*

$$\begin{aligned} \Psi_F : \text{hProj}_R^n(\mathbb{Z}, A) &\rightarrow \text{hProj}_R^{k-n}(A, \mathbb{Z}), & \Psi_F^{-1} : \text{hProj}_R^{k-n}(A, \mathbb{Z}) &\rightarrow \text{hProj}_R^n(\mathbb{Z}, A). \\ E &\mapsto (- \circ E)^{-1}(F) & E' &\mapsto (E' \circ -)^{-1}(F) \end{aligned}$$

We now specialise to the case where R is a ring with involution, i.e. a ring with an anti-automorphism $r \mapsto \bar{r}$ such that $\bar{\bar{r}} = r$ for all $r \in R$. For example, for a group G and a ring R , the group ring RG has involution $\sum_{i=1}^n r_i g_i \mapsto \sum_{i=1}^n r_i g_i^{-1}$ where $r_i \in R, g_i \in G$. Using this involution, any right R -module A can be viewed as a left R -module under the action $r \cdot x = x \cdot \bar{r}$ for $r \in R, x \in A$. If A is a left R -module, then $A^* = \text{Hom}_R(A, R)$ is a right R -module under the action $(\varphi \cdot r)(x) = \varphi(x)r$ for $\varphi \in A^*, r \in R$. We will view A^* as a left R -module using the involution on R .

Note that $(\cdot)^*$ can be viewed as a functor of R -modules: if $f : A_1 \rightarrow A_2$ is a map of R -modules, we can define $f^* : A_2^* \rightarrow A_1^*$ by sending $\varphi \mapsto \varphi \circ f$. For $E = (P_*, \partial_*) \in \text{Proj}_R^n(A, B)$, define the *dual extension* by

$$E^* = (0 \rightarrow A^* \xrightarrow{\varepsilon^*} P_0^* \xrightarrow{\partial_1^*} P_1^* \xrightarrow{\partial_2^*} \cdots \xrightarrow{\partial_{n-2}^*} P_{n-2}^* \xrightarrow{\partial_{n-1}^*} P_{n-1}^* \xrightarrow{i^*} B^* \rightarrow 0).$$

The dual of a projective module is projective since $P \oplus Q \cong R^n$ implies that $P^* \oplus Q^* \cong (R^n)^* \cong R^n$. In particular, the P_i^* are projective R -modules.

Whilst E^* is not exact in general, it is true under mild assumptions on the modules involved. We say that an R -module A is an *R -lattice* if its underlying abelian group is torsion-free. For example, all projective R -modules are R -lattices. Recall that the *evaluation map* is the map $e_A : A \rightarrow A^{**}, x \mapsto (f \mapsto f(x))$.

Lemma 2.5. *If A is a R -lattice, then $e_A : A \rightarrow A^{**}$ is an isomorphism.*

This follows by noting that, if $A \cong_{\text{Ab}} \mathbb{Z}^k$, then the R -module structure is determined by a map $\rho_A : R \rightarrow M_k(\mathbb{Z})$. It can be shown that $\rho_{A^*}(r) = \rho_A(\bar{r})^T$ using the induced identification $A^* \cong_{\text{Ab}} \mathbb{Z}^k$, from which the claim follows.

It follows easily from this that the reflexivity property of R -lattices also holds on the level of extensions.

Lemma 2.6 (Reflexivity). *If A, B are R -lattices and $n \geq 1$, then dualising gives an isomorphism of abelian groups*

$$* : \text{hProj}_R^n(A, B) \rightarrow \text{hProj}_R^n(B^*, A^*).$$

*If $E \in \text{Proj}_R^n(A, B)$, then there is a chain homotopy equivalence $e : E \rightarrow E^{**}$ induced by the evaluation maps.*

This has the following useful consequence which, in the language of [23, Theorem 28.5], says that projective R -modules are *injective relative to the class of R -lattices*.

Lemma 2.7. *Suppose A, B and E are R -lattices such that $(E, -) \in \text{Ext}_R^1(A, B)$ and P is a projective R -module. Then, for any map $f : B \rightarrow P$, there exists $\tilde{f} : E \rightarrow P$ such that $\tilde{f} \circ i = f$, i.e.*

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \xrightarrow{i} & E & \xrightarrow{\varepsilon} & A \longrightarrow 0 \\ & & \downarrow f & \swarrow \tilde{f} & & & \\ & & P & & & & \end{array}$$

We conclude this section by discussing an important invariant of projective extensions. Let $P(R)$ denote the R -module isomorphism classes of (finitely generated) projective R -modules and define the *projective class group* $C(R)$ as the quotient of $P(R)$ by the stable isomorphisms, where $P, Q \in P(R)$ are *stably isomorphic*, written $[P] = [Q]$, if $P \oplus R^i \cong Q \oplus R^j$ for some $i, j \geq 0$. This forms a group under direct sum and coincides with the Grothendieck group of the monoid $P(R)$.

For a projective extension

$$E = (0 \rightarrow B \xrightarrow{i} P_{n-1} \xrightarrow{\partial_{n-1}} P_{n-2} \xrightarrow{\partial_{n-2}} \cdots \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\varepsilon} A \rightarrow 0),$$

we define the *Euler class* $e(E) = \sum_{i=0}^{n-1} (-1)^i [P_i] \in C(R)$. This is known to be a congruence invariant [46, Lemma 1.3]. In fact, more is true:

Lemma 2.8. *If A, B are R -modules, the Euler class defines a map*

$$e : \mathrm{hProj}_R^n(A, B) \rightarrow C(R),$$

i.e. e is a chain homotopy invariant.

Proof. Suppose $E_1, E_2 \in \mathrm{Proj}_R^n(A, B)$ and that $\varphi : E_1 \rightarrow E_2$ is a chain homotopy equivalence. Then $E_2 \cong (\varphi_A)^*((\varphi_B)_*(E_1))$ and, since e is a congruence invariant, we have that $e(E_2) = e((\varphi_A)^*((\varphi_B)_*(E_1)))$. Since pushout and pullback by automorphisms can be made to not affect the isomorphism classes of the modules in the extension, this implies that $e((\varphi_A)^*((\varphi_B)_*(E_1))) = e(E_1)$ and so e is a chain homotopy invariant. \square

For a class $\chi \in C(R)$, we define $\mathrm{Proj}_R^n(A, B; \chi)$ to be the subset of $\mathrm{Proj}_R^n(A, B)$ consisting of those extensions with $e(E) = \chi$, and we can define $\mathrm{hProj}_R^n(A, B; \chi)$ similarly as a subset of $\mathrm{hProj}_R^n(A, B)$.

We have the following nice interpretations for the extensions $E \in \mathrm{Proj}_R^n(A, B)$ with $e(E) = 0$. This follows easily by repeatedly forming the direct sum with length two extensions $P \xrightarrow{\cong} P$ for various $P \in P(R)$.

Lemma 2.9. *If A, B are R -modules and $n \geq 2$, then every congruence class in $\mathrm{Proj}_R^n(A, B; 0)$ has a representative E of the form $E = (F_*, \partial_*)$ with the F_i free.*

This fails in the case $n = 1$, where it is not possible to form the direct sum with length two extensions $R \xrightarrow{\cong} R$ without altering the chain homotopy type. In fact, for a projective extension

$$E = (0 \rightarrow B \rightarrow P \rightarrow A \rightarrow 0),$$

we can define the *unstable Euler class* $\widehat{e}(E) = P \in P(R)$.

Lemma 2.10. *If A, B are R -modules, the unstable Euler class defines a map*

$$\widehat{e} : \mathrm{hProj}_R^1(A, B) \rightarrow P(R).$$

Proof. For $E_1 = (P_1, -), E_2 = (P_2, -) \in \mathrm{Proj}_R^1(A, B)$, recall that a chain map $\varphi : E_1 \rightarrow E_2$ is a chain homotopy equivalence if it induces a chain homotopy equivalence between the length one chain complexes P_1 and P_2 , i.e. if the restriction $\varphi|_{P_1} : P_1 \rightarrow P_2$ is an isomorphism. \square

3. SWAN MODULES AND PROJECTIVE EXTENSIONS

The aim of this section will be to build upon the results of the previous section in the case where $R = \mathbb{Z}G$ is the integral group ring of a finite group G .

We will begin by defining a special type of projective module first introduced in [41, Section 6]. Let $\varepsilon : \mathbb{Z}G \rightarrow \mathbb{Z}$, $g \mapsto 1$ and let $I = \mathrm{Ker}(\varepsilon) \subseteq \mathbb{Z}G$ be the augmentation ideal. For any $r \in \mathbb{Z}$ coprime to $|G|$, the ideal $(I, r) \subseteq \mathbb{Z}G$ is projective and depends only on $r \bmod |G|$ up to $\mathbb{Z}G$ -isomorphism [41].

The modules (I, r) are known as *Swan modules* and the map

$$S : (\mathbb{Z}/|G|)^\times \rightarrow C(\mathbb{Z}G)$$

given by $r \mapsto [(I, r)]$ is known as the *Swan map*. This is a well-defined group homomorphism [41], and we define the *Swan subgroup* to be $T_G = \text{Im}(S) \leq C(\mathbb{Z}G)$.

Remark 3.1. A closely related module is the ideal $(N, r) \subseteq \mathbb{Z}G$ where $N = \sum_{g \in G} g$ denotes the group norm. This is a Swan module since $(N, r) \cong (I, s)$ for any $s \in \mathbb{Z}$ such that $s \equiv r^{-1} \pmod{|G|}$. To see this, note that both modules are pullbacks of the maps $s : \mathbb{Z} \rightarrow \mathbb{Z}/|G|$, $1 \mapsto s$, and $\varepsilon : \mathbb{Z}G/(\Sigma) \rightarrow \mathbb{Z}/|G|$, $x + (\Sigma) \mapsto \varepsilon(x) + |G|$.

We now aim to classify $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ for a fixed $\mathbb{Z}G$ -module A . The following can be found in [23, Proposition 34.2]. Note that this isomorphism depends on the choice of $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ and so only exists when $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ is non-empty.

Theorem 3.2. *Let A be a $\mathbb{Z}G$ -module and $n \geq 1$. Then, for any $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$, there is a bijection*

$$(m_r)^* : (\mathbb{Z}/|G|)^\times \rightarrow \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$$

given by $r \mapsto (m_r)^(E)$, where $m_r : \mathbb{Z} \rightarrow \mathbb{Z}$ denotes multiplication by r .*

Recall that, if M is a left $\mathbb{Z}G$ -module and $r \in (\mathbb{Z}/|G|)^\times$, then the tensor product $(I, r) \otimes M$ can be considered as a left $\mathbb{Z}G$ -module since (I, r) is a two-sided ideal. This allows us to find the following explicit form for pullbacks of extensions.

Lemma 3.3. *Let A be a $\mathbb{Z}G$ -module, let $n \geq 1$ and let $r \in (\mathbb{Z}/|G|)^\times$. If $E = (P_*, \partial_*) \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$, then there are maps $\bar{\partial}_1, \bar{\varepsilon}$ such that*

$$(m_r)^*(E) \cong (0 \rightarrow A \xrightarrow{i} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_2} P_1 \xrightarrow{\bar{\partial}_1} (I, r) \otimes P_0 \xrightarrow{\bar{\varepsilon}} \mathbb{Z} \rightarrow 0).$$

Proof. As before, we can assume $n = 1$ by the shifting lemma. Let $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ and note that we have the following diagrams

$$\begin{array}{ccc} (I, r) & \xrightarrow{\frac{1}{r}\varepsilon} & \mathbb{Z} \\ \downarrow i & & \downarrow r \\ \mathbb{Z}G & \xrightarrow{\varepsilon} & \mathbb{Z} \end{array} \quad \begin{array}{ccc} (I, r) \otimes P & \xrightarrow{\frac{1}{r}\varepsilon \otimes 1} & \mathbb{Z} \otimes P \\ \downarrow i \otimes 1 & & \downarrow r \otimes 1 \\ \mathbb{Z}G \otimes P & \xrightarrow{\varepsilon \otimes 1} & \mathbb{Z} \otimes P \end{array}$$

where $i : (I, r) \hookrightarrow \mathbb{Z}G$ is inclusion. It can be checked directly that the first diagram is a pullback, and this implies that the second diagram is a pullback since P is projective and so flat. We can choose identifications $\mathbb{Z}G \otimes P \cong P$ and $\mathbb{Z} \otimes P \cong \mathbb{Z}$ for which $\varepsilon \otimes 1$ corresponds to ε^E . We now have a map $(\text{id}_A, \varphi, m_r) : E' \rightarrow E$ where $E' = ((I, r) \otimes P, -)$, and so $E' \cong (m_r)^*(E)$ by the uniqueness of pullbacks. \square

Remark 3.4. By combining with Theorem 3.2, this implies $(I, r) \otimes (I, s) \cong (I, rs)$.

Lemma 3.5. *Let $r \in \mathbb{Z}$ be coprime to $|G|$ and let $P \in P(\mathbb{Z}G)$. Then*

$$[(I, r) \otimes P] = [(I, r)] + [P].$$

Proof. Since there exists a surjection $\varphi : P \rightarrow \mathbb{Z}$, we can find $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ for some $\mathbb{Z}G$ -module A . Applying Schanuel's lemma to the map $\mu : (m_r)^*(E) \rightarrow E$ gives an isomorphism $\hat{\mu} : \mathbb{Z} \oplus ((I, r) \otimes P) \rightarrow \mathbb{Z} \oplus P$ which sends $(1, 0) \mapsto (r, x)$ for some $x \in P$. The result now follows from [41, Lemma 6.2]. \square

By combining all of the above, we now get the following.

Corollary 3.6. *Let e, \hat{e} denote the stable and unstable Euler classes. Let $n \geq 1$ and let A be a $\mathbb{Z}G$ -module such that there exists $E \in \text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$.*

(i) If $e(E) = [P]$, then

$$e(\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)) = [P] + T_G \subseteq C(\mathbb{Z}G).$$

(ii) If $n = 1$ and $E = (P, -) \in \text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$, then

$$\widehat{e}(\text{Proj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)) = T_G \cdot P \subseteq P(\mathbb{Z}G).$$

We conclude this section by specialising to the case $A = \mathbb{Z}$. The following can be extracted from [7, Chapter XII].

Proposition 3.7. *A finitely presented group G has k -periodic cohomology if and only if $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is non-empty.*

If G has k -periodic cohomology then, since $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is non-empty, Corollary 3.6 implies that there exists $P \in P(\mathbb{Z}G)$ for which

$$e(\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})) = [P] + T_G \subseteq C(\mathbb{Z}G).$$

We can then quotient by T_G to get a unique class in $C(\mathbb{Z}G)/T_G$ which depends only on G and k . The *Swan finiteness obstruction* is defined as

$$\sigma_k(G) = [P] \in C(\mathbb{Z}G)/T_G.$$

Recall that a group G has *free period k* if there exists $E = (F_*, \partial_*) \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with the F_i free.

Lemma 3.8. *If G has k -periodic cohomology, then $\sigma_k(G) = 0$ if and only if G has free period k .*

By a construction of Milnor, this is equivalent to the existence of a finite CW-complex X with $\pi_1(X) \cong G$ and $\tilde{X} \simeq S^{k-1}$ [41, Proposition 3.1]. Examples of groups with $\sigma_k(G) \neq 0$ were found by Milgram [31].

We would like to find a relation between $[P]$ and $[P^*]$ when $\sigma_k(G) = [P] + T_G$. This is difficult for general projectives since there exists finite groups G and projectives P for which $[P^*] \neq \pm[P]$, even in $C(\mathbb{Z}G)/T_G$. For example, we can take $G = \mathbb{Z}/37^2$ [11, Theorem 50.56]. However, in our situation, we have the following.

Lemma 3.9. *If G has k -periodic cohomology, and $\sigma_k(G) = [P] + T_G$, then*

$$[P] = -[P^*] \in C(\mathbb{Z}G)/T_G.$$

Proof. By Corollary 3.6, there exists $E \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(E) = [P]$ and, by forming the direct sum with length two extensions $\mathbb{Z}G \xrightarrow{\cong} \mathbb{Z}G$, we can assume that

$$E \cong (0 \rightarrow \mathbb{Z} \xrightarrow{i} P \xrightarrow{\partial_{k-1}} F_{k-2} \xrightarrow{\partial_{k-2}} \cdots \xrightarrow{\partial_1} F_0 \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0)$$

for some F_i free. Dualising then gives that

$$E^* \cong (0 \rightarrow \mathbb{Z} \xrightarrow{\varepsilon^*} F_0 \xrightarrow{\partial_1^*} \cdots \xrightarrow{\partial_{k-2}^*} F_{k-2} \xrightarrow{\partial_{k-1}^*} P^* \oplus F_{k-1} \xrightarrow{i^*} \mathbb{Z} \rightarrow 0)$$

and, since k is necessarily even [7, p261], Schanuel's lemma implies that

$$\mathbb{Z} \oplus P \oplus P^* \oplus F \cong \mathbb{Z} \oplus F'$$

for some F, F' free. By [41, Lemma 6.2], we then get that $[P \oplus P^*] \in T_G$. \square

4. CLASSIFICATION OF PROJECTIVE CHAIN COMPLEXES

We would now like to consider more generally the classification of projective extensions over $\mathbb{Z}G$ with only one fixed end. For $n \geq 0$, a *projective n -complex*

$E = (P_*, \partial_*)$ over $\mathbb{Z}G$ is a chain complex consisting of an exact sequence

$$E = (P_n \xrightarrow{\partial_n} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0)$$

where $H_0(P_*) \cong \mathbb{Z}$ and the P_i are (finitely generated) projective $\mathbb{Z}G$ -modules. An *algebraic n -complex* is a projective n -complex such that the P_i are free.

Let $\text{Proj}(G, n)$ denote the set of chain homotopy types of projective n -complexes over $\mathbb{Z}G$, which is a graded graph with edges between each $E = (P_*, \partial_*)$ and

$$E \oplus \mathbb{Z}G = (P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0).$$

Similarly, let $\text{Alg}(G, n)$ denote the set of chain homotopy types of algebraic n -complexes over $\mathbb{Z}G$, which is also a graded graph under stabilisation. By extending the projective n -complex by $\text{Ker}(\partial_n)$, it is easy to see that there is a bijection

$$\text{Proj}(G, n) \cong \coprod_{A \in \text{Mod}(\mathbb{Z}G)} \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A).$$

By abuse of notation, we will assume they are the same, i.e. that an extension $E \in \text{Proj}(G, n)$ lies in $\text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A)$ for some A . For a class $\chi \in C(\mathbb{Z}G)$, let $\text{Proj}(G, n; \chi)$ denote the subset of projective extensions E with $e(E) = \chi$. Note that $\text{Alg}(G, n) \cong \text{Proj}(G, n; 0)$ for $n \geq 2$.

The following is well-known (see [28, Theorem 1.1] or [20, Proof of Lemma 8.12]).

Theorem 4.1. *If $n \geq 0$ and $\chi \in C(\mathbb{Z}G)$, then $\text{Proj}(G, n; \chi)$ is a graded tree, i.e. if $E, E' \in \text{Proj}(G, n)$ have $e(E) = e(E')$, then $E \oplus \mathbb{Z}G^i \simeq E' \oplus \mathbb{Z}G^j$ for some $i, j \geq 0$.*

We will now prove a cancellation theorem for projective n -complexes. Our proof will be modelled on Hambleton-Kreck's proof that, if X and Y are finite 2-complexes with finite fundamental group such that $X \simeq X_0 \vee S^2$ and $X \vee S^2 \simeq Y \vee S^2$, then $X \simeq Y$ [19, Theorem B]. This idea was applied to algebraic 2-complexes in [18].

If A is a $\mathbb{Z}G$ -module, then $x \in A$ is *unimodular* if there exists a map $f : A \rightarrow \mathbb{Z}G$ such that $f(x) = 1$. Let $\text{Um}(A) \subseteq A$ denote the set of unimodular elements in A .

Lemma 4.2. *Let A be a $\mathbb{Z}G$ -module. Then*

- (i) *If $\varphi : A \rightarrow A$ is an automorphism, then $\varphi(\text{Um}(A)) = \text{Um}(A)$*
- (ii) *If $\varphi : A_0 \oplus \mathbb{Z}G \rightarrow A$ is an isomorphism, then $\varphi(0, 1) \in \text{Um}(A)$.*

Suppose a $\mathbb{Z}G$ -module A has a splitting $A = A_1 \oplus A_2 \oplus \cdots \oplus A_n$. Then a map $f : A_i \rightarrow A_j$ can be viewed as an automorphism of A by extending it to vanish everywhere else. Write $\text{GL}(A)$ for the group of automorphisms of A and define

$$E(A_i, A_j) = \langle 1 + f, 1 + g : f : A_i \rightarrow A_j, g : A_j \rightarrow A_i \rangle \leq \text{GL}(A)$$

to be the subgroup of *elementary automorphisms* for $i \neq j$, where $1 : A \rightarrow A$ denotes the identity map. The main result we will use is the following, which can be proven by combining [19, Corollary 1.12] with [19, Lemma 1.16].

Theorem 4.3. *Suppose A is a $\mathbb{Z}G$ -module for which $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module for all but finitely many primes p . If $F_1, F_2 \cong \mathbb{Z}G$, then*

$$\mathcal{G} = \langle E(F_1, A \oplus F_2), E(F_2, A \oplus F_1) \rangle \leq \text{GL}(A \oplus F_1 \oplus F_2)$$

acts transitively on $\text{Um}(A \oplus F_1 \oplus F_2)$.

Here $\mathbb{Z}_{(p)} = \{\frac{a}{b} : a, b \in \mathbb{Z}, p \nmid b\} \leq \mathbb{Q}$ denotes the localisation at a prime p and $A_{(p)} = A \otimes \mathbb{Z}_{(p)}$. The following can be proven by an easy extension of Maschke's theorem on representations.

Lemma 4.4. *If p is a prime not dividing $|G|$ and $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, A)$ is non-empty for n odd, then $\mathbb{Z}_{(p)} \oplus A_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module.*

Note that the fact that $\text{GL}(A \oplus \mathbb{Z}G^2)$ acts transitively on $\text{Um}(A \oplus \mathbb{Z}G^2)$ already implies the following cancellation theorem for modules.

Corollary 4.5. *Suppose A is a $\mathbb{Z}G$ -module, $A \cong A_0 \oplus \mathbb{Z}G$ and $\mathbb{Z}_{(p)} \oplus (A_0)_{(p)}$ is a free $\mathbb{Z}_{(p)}G$ -module for all but finitely many primes p . Then $A \oplus \mathbb{Z}G \cong A' \oplus \mathbb{Z}G$ implies $A \cong A'$.*

Proof. Let $\psi : A \oplus \mathbb{Z}G \rightarrow A' \oplus \mathbb{Z}G$ be an isomorphism and let $x = \psi^{-1}(0, 1) \in \text{Um}(A \oplus \mathbb{Z}G)$. Since $A = A_0 \oplus \mathbb{Z}G$, Theorem 4.3 implies that $\text{GL}(A \oplus \mathbb{Z}G)$ acts transitively on $\text{Um}(A \oplus \mathbb{Z}G)$ and so there is an isomorphism $\varphi : A \oplus \mathbb{Z}G \rightarrow A \oplus \mathbb{Z}G$ such that $\varphi(0, 1) = x$. Hence $\psi \circ \varphi : A \oplus \mathbb{Z}G \rightarrow A' \oplus \mathbb{Z}G$ has $(\psi \circ \varphi)(0, 1) = (0, 1)$ and so induces an isomorphism $(\psi \circ \varphi)|_A : A \rightarrow A' \oplus \mathbb{Z}G / \text{Im}(0 \oplus \mathbb{Z}G) \cong A'$. \square

We will upgrade the above argument from modules to projective n -complexes. The existence of a well-understood subgroup $\mathcal{G} \leq \text{GL}(A \oplus \mathbb{Z}G^2)$ which acts transitively on $\text{Um}(A \oplus \mathbb{Z}G^2)$ is important since we need only show that elements in \mathcal{G} can be extended to chain homotopy equivalences on the short exact sequences.

Theorem 4.6. *Let $n \geq 0$ be even and let $E, E' \in \text{Proj}(G, n)$. If $E \simeq E_0 \oplus \mathbb{Z}G$ and $E \oplus \mathbb{Z}G \simeq E' \oplus \mathbb{Z}G$, then $E \simeq E'$.*

Proof. Let $E_0 \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0)$, $E = (P_*, \partial_*) \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A)$ and $E' = (P'_*, \partial'_*) \in \text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A')$. If $\psi : E \oplus \mathbb{Z}G \rightarrow E' \oplus \mathbb{Z}G$ denotes the given chain homotopy equivalence in $\text{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0 \oplus \mathbb{Z}G^2)$ and $\psi_A : A_0 \oplus \mathbb{Z}G^2 \rightarrow A' \oplus \mathbb{Z}G$ is the induced map on the left, consider $x = \psi_A^{-1}(0, 1) \in \text{Um}(A_0 \oplus \mathbb{Z}G^2)$.

We now claim that there exists a self chain homotopy equivalence $\varphi : E \oplus \mathbb{Z}G \rightarrow E \oplus \mathbb{Z}G$ such that the induced map $\varphi_A : A \oplus \mathbb{Z}G \rightarrow A \oplus \mathbb{Z}G$ has $\varphi_A(0, 1) = x$.

Let $F_1, F_2 \cong \mathbb{Z}G$ be defined so that $A = A_0 \oplus F_1$ and $A \oplus \mathbb{Z}G = A_0 \oplus F_1 \oplus F_2$. Since $\text{Proj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A_0)$ is non-empty and $n+1$ is odd, we can combine Theorem 4.3 and Lemma 4.4 to get that there exists $\varphi_A \in \mathcal{G} = \langle E(F_1, A_0 \oplus F_2), E(F_2, A_0 \oplus F_1) \rangle \leq \text{GL}(A_0 \oplus F_1 \oplus F_2)$ such that $\varphi_A(0, 0, 1) = x$. We claim that φ_A can be extended to a chain homotopy equivalence $\varphi : E \oplus \mathbb{Z}G \rightarrow E \oplus \mathbb{Z}G$.

First recall that $E(F_2, A_0 \oplus F_1) = E(F_1, A) \leq \text{GL}(A \oplus F_1)$ is generated by elements of the form $\begin{pmatrix} 1 & 0 \\ f & 1 \end{pmatrix}$ for $f : A \rightarrow \mathbb{Z}G$ and $\begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix}$ for $g : \mathbb{Z}G \rightarrow A$.

If $i : A \hookrightarrow P$, then there exists $\tilde{f} : P \rightarrow \mathbb{Z}G$ such that $\tilde{f} \circ i = f$ by Lemma 2.7. It is straightforward to verify that the following diagrams commute, and so are chain homotopy equivalences.

$$\begin{array}{c} E \oplus \mathbb{Z}G \\ \downarrow \varphi_1 \\ E \oplus \mathbb{Z}G \end{array} = \begin{pmatrix} 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \\ \downarrow \begin{pmatrix} 1 & 0 \\ f & 1 \end{pmatrix} \quad \downarrow \begin{pmatrix} 1 & 0 \\ \tilde{f} & 1 \end{pmatrix} \quad \downarrow \text{id}_{P_{n-1}} \quad \downarrow \text{id}_{P_0} \\ 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \end{pmatrix}$$

$$\begin{array}{c} E \oplus \mathbb{Z}G \\ \downarrow \varphi_2 \\ E \oplus \mathbb{Z}G \end{array} = \begin{pmatrix} 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \\ \downarrow \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} \quad \downarrow \begin{pmatrix} 1 & i \circ g \\ 0 & 1 \end{pmatrix} \quad \downarrow \text{id}_{P_{n-1}} \quad \downarrow \text{id}_{P_0} \\ 0 \longrightarrow A \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix}} P_n \oplus \mathbb{Z}G \xrightarrow{(\partial_n, 0)} P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \xrightarrow{\partial_1} P_0 \longrightarrow 0 \end{pmatrix}$$

Similarly, we can show that the generators of $E(F_1, A_0 \oplus F_2)$ extend to chain homotopy equivalences. Hence, by writing $\varphi_A \in \mathcal{G}$ as the composition of maps of

this form, we can get a chain homotopy equivalence $\varphi : E \oplus \mathbb{Z}G \rightarrow E' \oplus \mathbb{Z}G$ by taking the composition of equivalences on each of the generators.

Now consider the map $\psi \circ \varphi = (\psi_A \circ \varphi_A, \psi_P \circ \varphi_P, \text{id}, \dots, \text{id}) : E \oplus \mathbb{Z}G \rightarrow E' \oplus \mathbb{Z}G$. Since $(\psi_A \circ \varphi_A)(0, 1) = (0, 1)$, it must have the form $\psi_A \circ \varphi_A = \begin{pmatrix} \phi_A & 0 \\ 0 & 1 \end{pmatrix}$ since it is an isomorphism. By commutativity, $(\psi_P \circ \varphi_P)(0, 1) = (0, 1)$ and so similarly $\psi_P \circ \varphi_P = \begin{pmatrix} \phi_P & 0 \\ 0 & 1 \end{pmatrix}$ for some $\phi_P : P \rightarrow P'$. We are now done by noting that the triple $(\phi_A, \phi_P, \text{id}, \dots, \text{id})$ defines a chain homotopy equivalence $E \simeq E'$. \square

We say that a graded tree is a *fork* if it has a single vertex at each non-minimal grade and a finite set of vertices at the minimal grade.




FIGURE 1. A graded tree which is a fork

Corollary 4.7. *If $n \geq 0$ is even, G is a finite group and $\chi \in C(\mathbb{Z}G)$, then $\text{Proj}(G, n; \chi)$ is a fork. In particular, $\text{Alg}(G, n)$ is a fork for $n \geq 2$ even.*

This recovers the even-dimensional case of a result of Browning [6, Theorem 5.4]. This fails in odd dimensions, i.e. there are examples of finite groups G for which $\text{Alg}(G, n)$ is not a fork for some n odd [14].

We now consider the case $n = 0$. Note that a projective 0-complex has the form

$$E = (0 \rightarrow A \xrightarrow{i} P \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0),$$

and so consists of a projective module $P \in P(\mathbb{Z}G)$ as well as the additional data (A, i, ε) . If \hat{e} is the unstable Euler class, then $\hat{e} : \text{Proj}(G, 0) \rightarrow P(\mathbb{Z}G)$ is a map of graded graphs since $\hat{e}(E \oplus \mathbb{Z}G) = \hat{e}(E) \oplus \mathbb{Z}G$. We will now show the following:

Theorem 4.8. *If G is a finite group, then the unstable Euler class gives an isomorphism of graded graphs*

$$\hat{e} : \text{Proj}(G, 0) \rightarrow P(\mathbb{Z}G).$$

Remark 4.9. Such a statement is implicit in the proof of [23, Theorem IV, Theorem 57.4], though the argument there contains an error and can only be used to recover the statement above in the case of projective modules of rank one. This, however, suffices since one can instead rely on the cancellation theorems of Hambleton-Kreck [19, Theorem B] or Browning [6, Theorem 5.4] at that stage in the proof.

In order to prove this, we will first note that projective $\mathbb{Z}G$ -modules for G a finite group take the following particularly simple form.

Proposition 4.10. *If G is a finite group and P is a projective $\mathbb{Z}G$ -module, then:*

- (i) *There is a projective ideal $P_0 \leq \mathbb{Z}G$ such that $P \cong P_0 \oplus \mathbb{Z}G^i$ for some $i \geq 0$*

- (ii) P is locally free in the sense that $P \otimes \mathbb{Q} \cong \mathbb{Q}G^{i+1}$ for some $i \geq 0$
- (iii) There exists a surjection $\varphi : P \rightarrow \mathbb{Z}$.

The first part was shown by Swan in [40, Theorem A], and it is straightforward to show that (i) \Rightarrow (ii) \Rightarrow (iii).

Proof of Theorem 4.8. To see that e is surjective, let $P \in P(\mathbb{Z}G)$. By Proposition 4.10, there is a surjection $\varphi : P \rightarrow \mathbb{Z}$ and this defines an extension $E = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, \text{Ker}(\varphi))$ which has $e(E) = P$.

We will now show injectivity. First let $E = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A)$ and let $E' = (P, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A')$. We will begin by considering the case where P has rank one. To show that $E \simeq E'$, it suffices to find isomorphisms $\varphi_A : A \rightarrow A'$ and $\varphi_{\mathbb{Z}} : \mathbb{Z} \rightarrow \mathbb{Z}$ such that the following diagram commutes:

$$\begin{array}{c} E \\ \downarrow \varphi \\ E' \end{array} = \begin{pmatrix} 0 & \longrightarrow & A & \xrightarrow{i} & P & \xrightarrow{\varepsilon} & \mathbb{Z} & \longrightarrow & 0 \\ & & \downarrow \varphi_A & & \downarrow \text{id} & & \downarrow \varphi_{\mathbb{Z}} & & \\ 0 & \longrightarrow & A' & \xrightarrow{i'} & P & \xrightarrow{\varepsilon'} & \mathbb{Z} & \longrightarrow & 0 \end{pmatrix}.$$

Consider the maps $\bar{\varepsilon} = \varepsilon \otimes \mathbb{Q}$, $\bar{\varepsilon}' = \varepsilon' \otimes \mathbb{Q} : P \otimes \mathbb{Q} \cong \mathbb{Q}G \rightarrow \mathbb{Q}$. Since \mathbb{Q} has trivial G -action, each map is determined by the fact that $\bar{\varepsilon}(g) = \bar{\varepsilon}'(g) = 0$ for all $g \in G$ and $\bar{\varepsilon}(1) = \bar{\varepsilon}'(1) = x_i$ for some $x_i \in \mathbb{Q}^\times$. Hence $\text{Ker}(\bar{\varepsilon}) = \text{Ker}(\bar{\varepsilon}')$ and so $(\varepsilon' \circ i) \otimes \mathbb{Q} = 0$. Since A is a $\mathbb{Z}G$ lattice, this implies that $\varepsilon' \circ i = 0$ and so we can define maps φ_A and $\varphi_{\mathbb{Z}}$ as above. Now $\varphi_{\mathbb{Z}}$ is necessarily surjective and so an isomorphism. Hence φ_A is an isomorphism by the five lemma, and so $E \simeq E'$.

Now suppose E, E' as above but with $\text{rank}(P) \geq 2$. By Proposition 4.10, this implies that there exists P_0 of rank one such that $P \cong P_0 \oplus \mathbb{Z}G^i$ for some $i \geq 1$. Since e is surjective, there exists $E_0 = (P_0, -) \in \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A_0)$ for some A_0 . By Theorem 4.1, there exists $j \geq 0$ for which $E_0 \oplus \mathbb{Z}G^{i+j} \simeq E \oplus \mathbb{Z}G^j \simeq E' \oplus \mathbb{Z}G^j$. Since $i \geq 1$, Theorem 4.6 then implies that $E_0 \oplus \mathbb{Z}G^i \simeq E \simeq E'$. \square

The following is immediate from the definition of $\text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A; [P])$, and is the key result that we will use from this section.

Corollary 4.11. *If $P \in P(\mathbb{Z}G)$, then there is an isomorphism of graded trees*

$$\hat{e} : \text{Proj}(G, 0; [P]) \rightarrow [P].$$

5. POLARISED HOMOTOPY CLASSIFICATION OF (G, n) -COMPLEXES

Recall that, for a group G , a G -polarised space is a pair (X, ρ_X) where X is a topological space and $\rho_X : \pi_1(X, *) \rightarrow G$ is a given isomorphism. We say that two G -polarised spaces $(X, \rho_X), (Y, \rho_Y)$ are *polarised homotopy equivalent* if there exists a homotopy equivalence $h : X \rightarrow Y$ such that $\rho_X = \rho_Y \circ \pi_1(h)$.

Let $\text{PHT}(G, n)$ denote the set of polarised homotopy types of finite (G, n) -complexes over G . This is a graded graph with edges between each (X, ρ_X) and $(X \vee S^2, (\rho_X)^+)$ where $(\rho_X)^+$ is induced by ρ_X and the collapse map $X \vee S^2 \rightarrow X$.

If X is a finite CW-complex, then the cellular chain complex $C_*(\tilde{X})$ can be viewed as a chain complex of $\mathbb{Z}[\pi_1(X)]$ -modules under the monodromy action. We can use a G -polarisation $\rho : \pi_1(X) \rightarrow G$ to get a chain complex of $\mathbb{Z}G$ -modules $C_*(\tilde{X}, \rho)$ which is the same as $C_*(\tilde{X})$ as a chain complex of abelian groups but with action $g \cdot x = \rho^{-1}(g)x$ for all $g \in G$ and $x \in C_i(\tilde{X})$ for some $i \geq 0$.

The following is a mild generalisation of [36, Theorem 1.1]:

Proposition 5.1. *Let G be a finitely presented group and let $n \geq 2$. Then there is an injective map graded trees*

$$\tilde{C}_* : \text{PHT}(G, n) \rightarrow \text{Alg}(G, n)$$

induced by the map $(X, \rho) \mapsto C_(\tilde{X}, \rho)$. Furthermore:*

- (i) *If $n \geq 3$, then \tilde{C}_* is bijective.*
- (ii) *If $n = 2$, then \tilde{C}_* is bijective if and only if G has the D2 property.*

Remark 5.2. (a) Even if G does not satisfy the D2 property, Proposition 5.1 can be replaced with an isomorphism $\tilde{C}_* : \text{D2}(G) \rightarrow \text{Alg}(G, 2)$ where $\text{D2}(G)$ denotes the polarised homotopy tree of D2-complexes over G [36, Theorems 1.1].

(b) This is often vacuous in the case $n \geq 3$ since $\text{PHT}(G, n)$ and $\text{Alg}(G, n)$ are non-empty. More specifically, $\text{Alg}(G, n)$ is non-empty if and only if G is of type FP_n , and there exists finitely presented groups which are not of type FP_n for $n \geq 3$ [39].

(c) This fails in general for non-finitely presented groups. In particular, for each $n \geq 2$, Bestvina-Brady constructed a non-finitely presented group G of type FP_n [3]. Here $\text{PHT}(G, n)$ is empty and $\text{Alg}(G, n)$ is non-empty and so \tilde{C}_* is not bijective.

We will now use the results from the previous section to study projective n -complexes over groups with periodic cohomology. By Proposition 5.1, this will lead to a proof of the following more detailed version of Theorem A. Note that, if X is a finite (G, n) -complex, then $\pi_n(X) \cong H_n(\tilde{X}) \cong \text{Ker}(\partial_n : C_n(\tilde{X}) \rightarrow C_{n-1}(\tilde{X}))$.

Theorem 5.3. *Let G have k -periodic cohomology, let $n = ik$ or $ik - 2$ for some $i \geq 1$ and let $P_{(G, n)}$ be a projective $\mathbb{Z}G$ -module with $\sigma_{ik}(G) = [P_{(G, n)}] \in C(\mathbb{Z}G)/T_G$. Let $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ be such that $e(F) = [P_{(G, n)}]$. Then there is an injective map of graded trees*

$$\Psi : \text{PHT}(G, n) \rightarrow [P_{(G, n)}]$$

which is defined as follows.

- (i) *If $n = ik - 2$, then $\Psi : X \mapsto P$, where P is the unique projective $\mathbb{Z}G$ -module for which $(0 \rightarrow \mathbb{Z} \xrightarrow{\alpha} P^* \xrightarrow{\beta} \pi_n(X) \rightarrow 0) \circ C_*(\tilde{X}) \simeq F$ for some α, β .*
- (ii) *If $n = ik$, then $\Psi : X \mapsto P$, where P is the unique projective $\mathbb{Z}G$ -module for which $C_*(\tilde{X}) \simeq (0 \rightarrow \pi_n(X) \xrightarrow{\alpha} P \xrightarrow{\beta} \mathbb{Z} \rightarrow 0) \circ F$ for some α, β .*

Furthermore, Ψ is bijective if and only if $n \geq 3$ or $n = 2$ and G has the D2 property.

Remark 5.4. The definition of $P_{(G, n)}$ depends on G , n and k . Note that n and k determine i except when $k = 2$ where $n = ik = (i + 1)k - 2$. However, in this case there is no ambiguity since G is cyclic [42, Lemma 5.2] and so $\sigma_{2i}(G) = 0$ for all i .

First note that, when G has periodic cohomology, we get the following two relations between projective complexes of different dimensions.

Lemma 5.5. *Suppose G has k -periodic cohomology and let $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. If $n \geq 0$, then we have the following isomorphisms of graded graphs*

$$\begin{aligned} - \circ F : \text{Proj}(G, n) &\rightarrow \text{Proj}(G, n + k) \\ * \circ \psi_F : \text{Proj}(G, n) &\rightarrow \text{Proj}(G, k - (n + 2)) \end{aligned}$$

where $n + 2 \leq k$ is the second case.

Proof. The first isomorphism is immediate from the shifting lemma. The second isomorphism consists of the compositions

$$\mathrm{hProj}_{\mathbb{Z}G}^{n+1}(\mathbb{Z}, A) \xrightarrow{\Psi_F} \mathrm{hProj}_{\mathbb{Z}G}^{k-n-1}(A, \mathbb{Z}) \xrightarrow{*} \mathrm{hProj}_{\mathbb{Z}G}^{k-n-1}(\mathbb{Z}, A)$$

for all $\mathbb{Z}G$ -modules A . These are bijections by the duality and reflexivity lemmas, and has image $\mathrm{Proj}(G, k - (n + 2))$. \square

Remark 5.6. Furthermore, if $E \in \mathrm{Proj}(G, n)$ has $\chi = e(E)$, then it is easy to see that $e(E \circ F) = e(F) + \chi$ since k is even and $e((\psi_F(E))^*) = e(F)^* - \chi^*$.

The proof of Theorem 5.3 will now consist of applying Lemma 5.5 in the case $k \mid n$ or $n + 2$ and then composing with the isomorphism from Theorem 4.8.

We will need the following result of Wall [46, Corollary 12.6].

Proposition 5.7. *If G has k -periodic cohomology, then*

$$2\sigma_k(G) = 0 \in C(\mathbb{Z}G)/T_G.$$

By iterating extensions using the Yoneda product, it can be shown that $n\sigma_k(G) = \sigma_{nk}(G)$ and so this theorem is equivalent to showing that $\sigma_{2k}(G) = 0$, i.e. that the obstruction vanishes whenever k is not the minimal period.

Theorem 5.8. *Suppose G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$. Then there exists $F \in \mathrm{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ such that there are isomorphisms of graded trees*

$$\mathrm{Alg}(G, k) \xrightarrow{(-\circ F)^{-1}} \mathrm{Proj}(G, 0; [P_{(G,n)}]) \xrightarrow{\hat{e}} [P_{(G,n)}].$$

Proof. By Proposition 5.7, we have that $\sigma_k(G) = [P_{(G,n)}] + T_G = -[P_{(G,n)}] + T_G$ and so there exists $F \in \mathrm{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(F) = -[P_{(G,n)}]$ by Corollary 3.6. By Lemma 5.5 and Remark 3.9, we get that $(-\circ F)^{-1}$ is the required isomorphism of graded trees. That e is an isomorphism follows from Corollary 4.11. \square

Theorem 5.9. *Suppose G has k -periodic cohomology and $\sigma_k(G) = [P_{(G,n)}] + T_G$ for some $P_{(G,n)} \in P(\mathbb{Z}G)$. Then there exists $F \in \mathrm{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ such that there are isomorphisms of graded trees*

$$\mathrm{Alg}(G, k - 2) \xrightarrow{* \circ \Psi_F} \mathrm{Proj}(G, 0; [P]) \xrightarrow{\hat{e}} [P].$$

Proof. By Lemma 3.9 and Proposition 5.7, we have that $\sigma_k(G) = [P_{(G,n)}] + T_G = [P_{(G,n)}^*] + T_G$ and so there exists $F \in \mathrm{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ with $e(F) = [P_{(G,n)}^*]$ by Corollary 3.6. The result now follows from Lemma 5.5, Remark 3.9 and Corollary 4.11. \square

Proof of Theorem 5.3. If G has k -periodic cohomology, then it also has ik -periodic cohomology for any $i \geq 1$. Hence, by swapping k for ik , we can assume $i = 1$. By combining Theorems 5.8 and 5.9 with Proposition 5.1, we obtain injective maps of graded trees $\Psi : \mathrm{PHT}(G, n) \rightarrow [P_{(G,n)}]$ for $n = k$ or $k - 2$, which are bijective as required. It remains to show that, in each case, Ψ has the form given in (i), (ii).

If $n = k - 2$, then $(*\circ \Psi_F)(C_*(\tilde{X})) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0)$ for some A and some $P \in [P_{(G,n)}]$. By Lemma 2.6, we have $\Psi_F(C_*(\tilde{X})) \simeq (0 \rightarrow \mathbb{Z} \rightarrow P^* \rightarrow A^* \rightarrow 0)$. Hence $A^* \cong \pi_n(X)$ and $(0 \rightarrow \mathbb{Z} \rightarrow P^* \rightarrow A^* \rightarrow 0) \circ C_*(\tilde{X}) \simeq F$.

If $n = k$, then $(-\circ F)^{-1}(C_*(\tilde{X})) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0)$ for some A and some $P \in [P_{(G,n)}]$. Hence $C_*(\tilde{X}) \simeq (0 \rightarrow A \rightarrow P \rightarrow \mathbb{Z} \rightarrow 0) \circ F$ and $A \cong \pi_n(X)$. \square

This completes the proof of Theorem A.

6. HOMOTOPY CLASSIFICATION OF (G, n) -COMPLEXES

For a finitely presented group G , an automorphism $\theta \in \text{Aut}(G)$ acts on $\text{PHT}(G, n)$ by sending $(X, \rho) \mapsto (X, \theta \circ \rho)$. It is straightforward to see that

$$\text{HT}(G, n) \cong \text{PHT}(G, n) / \text{Aut}(G)$$

and the goal of this chapter will be to determine the induced action of $\text{Aut}(G)$ on $[P_{(G, n)}]$ under the isomorphism $\text{PHT}(G, n) \cong [P_{(G, n)}]$ obtained in Theorem 5.3.

6.1. Preliminaries on the action of $\text{Aut}(G)$. We begin by defining natural actions of $\text{Aut}(G)$ on $\mathbb{Z}G$ -modules and chain complexes of $\mathbb{Z}G$ -modules. Firstly, for a $\mathbb{Z}G$ -module A and $\theta \in \text{Aut}(G)$, let A_θ denote the $\mathbb{Z}G$ -module whose underlying abelian group is that of A and whose action is $g \cdot x = \theta(g)x$ where $g \in G$, $x \in A$. This action has the following basic properties:

Lemma 6.1. *Let $\theta \in \text{Aut}(G)$. Then*

(i) *There is a $\mathbb{Z}G$ -module isomorphism*

$$i_\theta : \mathbb{Z}G \rightarrow \mathbb{Z}G_\theta, \quad \sum_{g \in G} a_i g_i \mapsto \sum_{g \in G} a_i \theta(g_i).$$

(ii) *If $A, B \in \text{Mod}(\mathbb{Z}G)$, then $(A \oplus B)_\theta \cong A_\theta \oplus B_\theta$.*

(iii) *If $P \in P(\mathbb{Z}G)$, then $P_\theta \in P(\mathbb{Z}G)$.*

We can extend the action to chain complexes as follows. If A, B are $\mathbb{Z}G$ -modules and $E = (E_*, \partial_*) \in \text{Ext}_{\mathbb{Z}G}^n(A, B)$, then we define $E_\theta \in \text{Ext}_{\mathbb{Z}G}^n(A_\theta, B_\theta)$ by

$$E_\theta = (0 \rightarrow B_\theta \xrightarrow{\partial_n} (E_{n-1})_\theta \xrightarrow{\partial_{n-1}} (E_{n-2})_\theta \rightarrow \cdots \rightarrow (E_1)_\theta \xrightarrow{\partial_1} (E_0)_\theta \xrightarrow{\partial_0} A_\theta \rightarrow 0).$$

It is easy to see that this is well-defined up to chain homotopy and, by the lemma above, it preserves projective extensions and so also induces a map on $\text{hProj}_{\mathbb{Z}G}^n(A, B)$. This following is immediate from the definition of $\tilde{C}_*(X, \rho)$.

Lemma 6.2. *If $E \in \text{Alg}(G, n)$, then the induced action of $\theta \in \text{Aut}(G)$ on E is given by $\theta \cdot E = E_\theta$, i.e. if $E = \tilde{C}_*(X, \rho)$, then $E_\theta = \tilde{C}_*(X, \theta \circ \rho)$.*

We now establish a few basic properties of this action which we will use later in this section. Firstly, we note that the action commutes with dualising.

Lemma 6.3. *If A and B are $\mathbb{Z}G$ -lattices, $E \in \text{Proj}_{\mathbb{Z}G}^n(A, B)$ for $n \geq 1$ and $\theta \in \text{Aut}(G)$, then*

$$(E_\theta)^* \cong (E^*)_\theta.$$

Proof. We begin by proving the corresponding statement for modules, i.e. that, if A is a $\mathbb{Z}G$ -lattice, then $(A_\theta)^* \cong (A^*)_\theta$. Let $A \cong_{\text{Ab}} \mathbb{Z}^k$, so that the $\mathbb{Z}G$ -module structure is determined by an integral representation $\rho_A : G \rightarrow \text{GL}_k(\mathbb{Z})$. As remarked earlier, we have that $\rho_{A^*}(g) = \rho_A(g^{-1})^T$ and it is easy to see that $\rho_{A_\theta} = \rho_A \circ \theta$. Therefore $(A_\theta)^* \cong (A^*)_\theta$ follows by noting that

$$\rho_{(A_\theta)^*}(g) = \rho_{A_\theta}(g^{-1})^T = \rho_A(\theta(g^{-1}))^T = \rho_A(\theta(g)^{-1})^T$$

and

$$\rho_{(A^*)_\theta}(g) = \rho_{A^*}(\theta(g)) = \rho_A(\theta(g)^{-1})^T.$$

The result for extensions now follows immediately since θ only affects the underlying modules and not the maps between them. \square

In light of this, for $\mathbb{Z}G$ -lattices A and B and $E \in \text{Proj}_{\mathbb{Z}G}^n(A, B)$, it now makes sense to write A_θ^* and E_θ^* . Note that the action also commutes with pullbacks.

Lemma 6.4. *If $\theta \in \text{Aut}(G)$, $f : B_1 \rightarrow B_2$ is a $\mathbb{Z}G$ -module homomorphism and $E \in \text{Ext}_{\mathbb{Z}G}^n(A, B_1)$, then*

$$f_*(E_\theta) \cong (f_*(E))_\theta.$$

6.2. Proof of Theorem B. In the case where $A = B = \mathbb{Z}$, we can consider this as an action on $\text{Proj}_{\mathbb{Z}G}^n(\mathbb{Z}, \mathbb{Z})$ by using the identification $\mathbb{Z}_\theta \cong \mathbb{Z}$.

Lemma 6.5. *If G has k -periodic cohomology, then there exists a unique map $\psi_k : \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$ such that, for every $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and $\theta \in \text{Aut}(G)$, we have*

$$F_\theta \cong (m_{\psi_k(\theta)})_*(F).$$

Proof. Fix an extension $F_0 \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. By dualising and then applying Theorem 3.2, it follows that every extension in $\text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is of the form $(m_r)_*(F_0)$ for some $r \in (\mathbb{Z}/|G|)^\times$. For $\theta \in \text{Aut}(G)$, define $\psi_k(\theta) = r \in (\mathbb{Z}/|G|)^\times$ for any $r \in (\mathbb{Z}/|G|)^\times$ such that $(F_0)_\theta \cong (m_r)_*(F_0)$.

If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$, then $F \cong (m_r)_*(F_0)$ for some $r \in (\mathbb{Z}/|G|)^\times$. By Lemma 6.4, we now have that

$$\begin{aligned} F_\theta &\cong ((m_r)_*(F_0))_\theta \cong (m_r)_*((F_0)_\theta) \cong (m_r)_*((m_{\psi_n(\theta)})_*(F_0)) \\ &\cong (m_{\psi_n(\theta)})_*((m_r)_*(F_0)) \cong (m_{\psi_n(\theta)})_*(F). \end{aligned} \quad \square$$

Lemma 6.6. *If $E, E' \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and $r \in \mathbb{Z}$ coprime to $|G|$, then*

$$E \circ (m_r)_*(E') \cong (m_r)_*(E) \circ E'.$$

Proof. Consider the pushout map $\nu : E' \rightarrow (m_r)_*(E')$. Since this induces m_r on the left copy of \mathbb{Z} , we can extend it to a map $\tilde{\nu} : E \circ E' \rightarrow E \circ (m_r)_*(E')$ which induces multiplication by $r \in \mathbb{Z} \subseteq \mathbb{Z}G$ on every module in E , i.e.

$$\begin{array}{c} E \circ E' \\ \downarrow \tilde{\nu} \\ E \circ (m_r)_*(E') \end{array} = \left(\begin{array}{cccccccccccc} 0 & \rightarrow & \mathbb{Z} & \xrightarrow{i} & P_{k-1} & \xrightarrow{\partial_{k-1}^{-1}} & \dots & \xrightarrow{\partial_1} & P_0 & \xrightarrow{i' \circ \varepsilon} & P'_{k-1} & \xrightarrow{\partial'_{k-1}^{-1}} & \dots & \xrightarrow{\partial'_1} & P'_0 & \xrightarrow{\varepsilon'} & \mathbb{Z} & \rightarrow & 0 \\ & & \downarrow r & & \downarrow r & & & & \downarrow r & & \downarrow \nu_{k-1} & & & & \downarrow \nu_0 & & \downarrow 1 & & \\ 0 & \rightarrow & \mathbb{Z} & \xrightarrow{i} & P_{k-1} & \xrightarrow{\partial_{k-1}^{-1}} & \dots & \xrightarrow{\partial_1} & P_0 & \xrightarrow{i' \circ \varepsilon} & P'_{k-1} & \xrightarrow{\partial'_{k-1}^{-1}} & \dots & \xrightarrow{\partial'_1} & P'_0 & \xrightarrow{\varepsilon'} & \mathbb{Z} & \rightarrow & 0 \end{array} \right).$$

By the uniqueness of pushouts, this implies that $E \circ (m_r)_*(E') \cong (m_r)_*(E \circ E') = (m_r)_*(E) \circ E'$ as required. \square

Note that, if G has k -periodic cohomology and $k \mid n$, then it also has n -periodic cohomology and so ψ_n can still be defined using Lemma 6.5. The above lemma now allows us to give the following relation between ψ_k and ψ_n for $k \mid n$.

Lemma 6.7. *If G has k -periodic cohomology, $i \geq 1$ and $\theta \in \text{Aut}(G)$, then*

$$\psi_{ik}(\theta) = \psi_k(\theta)^i.$$

Proof. Let $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ and consider $F^i \in \text{Proj}_{\mathbb{Z}G}^{ik}(\mathbb{Z}, \mathbb{Z})$. Then Lemma 6.5 implies that $F_\theta \cong (m_{\psi_k(\theta)})_*(F)$ and $(F^i)_\theta \cong (m_{\psi_{ik}(\theta)})_*(F^i)$. Since $(F^i)_\theta \cong (F_\theta)^i$, this implies that $(m_{\psi_{ik}(\theta)})_*(F^i) \cong ((m_{\psi_k(\theta)})_*(F))^i$.

By repeated application of Lemma 6.6, we get that

$$(m_{\psi_{ik}(\theta)})_*(F^i) \cong ((m_{\psi_k(\theta)})_*(F))^i \cong (m_{\psi_k(\theta)})_*^i(F^i) \cong (m_{\psi_k(\theta)^i})_*(F^i)$$

and so $\psi_{ik}(\theta) \cong \psi_k(\theta)^i \bmod |G|$ by the extension of Theorem 3.2 to arbitrary extensions via the shifting lemma. \square

In order to prove Theorem B, it suffices to check what the action of $\text{Aut}(G)$ corresponds to under the isomorphisms described in Section 5. Similarly to Section 5, it will suffice to consider the cases where $k = n$ or $n + 2$.

Theorem 6.8. *Suppose G has k -periodic cohomology and $\sigma_k(G) = [P] + T_G$ for some $P \in P(\mathbb{Z}G)$. If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is such that $e(F) = -[P]$, then*

$$\begin{array}{ccc} \text{hProj}_{\mathbb{Z}G}^{k+1}(\mathbb{Z}, A; 0) & \xrightarrow{(- \circ F)^{-1}} & \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A; [P]) \xrightarrow{\widehat{e}} [P] \\ E & \longmapsto & E' \longmapsto P \\ E_\theta & \longmapsto & (m_{\psi_k(\theta)})^*((E')_\theta) \longmapsto (I, \psi_k(\theta)) \otimes P_\theta \end{array}$$

Proof. For the first map, it suffices to check that $(\psi_k(\theta))^*((E')_\theta) \circ F \simeq E_\theta$. Since $E' \circ F = E$, we have $(E')_\theta \circ F_\theta \simeq E_\theta$. By Lemma 6.5, we have $F_\theta \cong (m_{\psi_n(\theta)})_*(F)$ and so

$$E_\theta \simeq (E')_\theta \circ (m_{\psi_n(\theta)})_*(F) = (m_{\psi_n(\theta)})^*((E')_\theta) \circ F.$$

For the second map, we can use Lemma 3.3 to get that

$$\widehat{e}((m_{\psi_k(\theta)})^*((E')_\theta)) \cong (I, \psi_k(\theta)) \otimes \widehat{e}((E')_\theta) \cong (I, \psi_k(\theta)) \otimes P_\theta. \quad \square$$

Theorem 6.9. *Suppose G has k -periodic cohomology and $\sigma_k(G) = [P] + T_G$ for some $P \in P(\mathbb{Z}G)$. If $F \in \text{Proj}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$ is such that $e(F) = -[P^*]$, then*

$$\begin{array}{ccc} \text{hProj}_{\mathbb{Z}G}^{k-1}(\mathbb{Z}, A; 0) & \xrightarrow{\Psi_F} \text{hProj}_{\mathbb{Z}G}^1(A, \mathbb{Z}; [P^*]) \xrightarrow{*} \text{hProj}_{\mathbb{Z}G}^1(\mathbb{Z}, A^*; [P]) \xrightarrow{\widehat{e}} [P] \\ E & \longmapsto & E' \longmapsto (E')^* \longmapsto P \\ E_\theta & \longmapsto & (m_{\psi_k(\theta)^{-1}})^*((E')_\theta) \longmapsto (m_{\psi_k(\theta)})^*((E')_\theta^*) \longmapsto (I, \psi_k(\theta)) \otimes P_\theta \end{array}$$

Proof. For this first map, it suffices to check that $(m_{\psi_k(\theta)^{-1}})^*((E')_\theta) \circ E_\theta \simeq F$. Since $E' \circ E \simeq F$, we have $(E')_\theta \circ E_\theta \simeq F_\theta$. By Lemma 6.5, we have $F_\theta \cong (m_{\psi_n(\theta)})_*(F)$ and so

$$F \simeq (m_{\psi_k(\theta)^{-1}})^*((E')_\theta \circ E_\theta) \simeq (m_{\psi_k(\theta)^{-1}})^*((E')_\theta) \circ E_\theta.$$

For the second map, it is easy to see that pushouts dualise to pullbacks in the other direction, i.e. if $E_0 = (m_{\psi_k(\theta)^{-1}})^*((E')_\theta)$, then $(m_{\psi_k(\theta)^{-1}})^*(E_0^*) \simeq (E')_\theta^*$ and so $E_0^* \simeq (m_{\psi_k(\theta)})^*((E')_\theta^*)$. For the third map, we can use Lemma 3.3 to get that

$$\widehat{e}((m_{\psi_k(\theta)})^*((E')_\theta^*)) = (I, \psi_k(\theta)) \otimes \widehat{e}((E')_\theta^*) \cong (I, \psi_k(\theta)) \otimes P_\theta. \quad \square$$

If G has k -periodic cohomology and $n = ik$ or $ik - 2$ for some $i \geq 1$, then the above shows that the induced action of $\theta \in \text{Aut}(G)$ on $[P]$ is given by $P \mapsto (I, \psi_{ik}(\theta)) \otimes P_\theta$ and $\psi_{ik}(\theta) = \psi_k(\theta)^i$ by Lemma 6.7.

This completes the proof of Theorem B except for a possible discrepancy in the case where $k = 2$ and i is not determined by the fact that $n = ik$ or $ik - 2$ (see Remark 5.4). However, in this case, G is cyclic and so $(I, r) \cong \mathbb{Z}G$ for all $r \in (\mathbb{Z}/|G|)^\times$ by [41, Corollary 6.1]. Hence $(I, \psi_k(\theta)^i) \cong \mathbb{Z}G$ is independent of i .

7. STABLY FREE SWAN MODULES AND (G, n) -COMPLEXES

Before computing the action of $\text{Aut}(G)$ on $[P_{(G, n)}]$, we will pause to consider to consider the role of Swan modules in the classification of (G, n) -complexes. We

begin by considering the map

$$\psi_k : \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$$

where G has k -periodic cohomology.

If $\theta \in \text{Aut}(G)$, then the action $E \mapsto E_\theta$ induces an action of $\text{Aut}(G)$ on $H^k(G; \mathbb{Z}) = \text{Ext}_{\mathbb{Z}G}^k(\mathbb{Z}, \mathbb{Z})$. This agrees with the usual action coming from the alternate definition of $H^k(-; \mathbb{Z})$ as a functor on groups [7, Chapter XII]. This implies that $\text{Im}(\psi_k) = \text{Aut}_k(G)$ which is defined in [13, Section 8]. We will now give several examples of maps $\psi_k : \text{Aut}(G) \rightarrow (\mathbb{Z}/|G|)^\times$. Here we write $\mathbb{Z}_n = \mathbb{Z}/n$.

Cyclic. If $C_n = \langle x \mid x^n = 1 \rangle$ is the cyclic group of order n , then

$$\text{Aut}(C_n) = \{\theta_i : x \mapsto x^i : i \in \mathbb{Z}_n^\times\}$$

and $\psi_2 : \text{Aut}(C_n) \rightarrow \mathbb{Z}_n^\times$ sends $\theta_i \mapsto i$ by [41, Proposition 8.1]. This is surjective and so recovers the classical results $T_{C_n} = 1$.

Dihedral. If $D_{4n+2} = \langle x, y \mid x^{2n+1} = y^2 = 1, yxy^{-1} = x^{-1} \rangle$ is the dihedral group of order $4n+2$, then

$$\text{Aut}(D_{4n+2}) = \{\theta_{i,j} : x \mapsto x^i, y \mapsto x^j y : i \in \mathbb{Z}_{2n+1}^\times, j \in \mathbb{Z}_{2n+1}\}$$

and $\psi_4 : \text{Aut}(D_{4n+2}) \rightarrow \mathbb{Z}_{4n+2}^\times$ sends $\theta_{i,j} \mapsto i^2$ by the discussion in [22, Section 5]. Since $\mathbb{Z}_{4n+2}^\times = \pm(\mathbb{Z}_{4n+2}^\times)^2$, this recovers the result $T_{D_{4n+2}} = 1$.

Quaternionic. Let $Q_{4n} = \langle x, y \mid x^n = y^2, yxy^{-1} = x^{-1} \rangle$ is the quaternion group of order $4n$. For $n = 2$, it is shown in [41, Proposition 8.3] that $\psi_4 : \text{Aut}(Q_8) \rightarrow \mathbb{Z}_8^\times$ sends $\theta \mapsto 1$ for all $\theta \in \text{Aut}(G)$. For $n \geq 3$, we have

$$\text{Aut}(Q_{4n}) = \{\theta_{i,j} : x \mapsto x^i, y \mapsto x^j y : i \in \mathbb{Z}_{2n}^\times, j \in \mathbb{Z}_{2n}\}$$

and $\psi_4 : \text{Aut}(Q_{4n}) \rightarrow \mathbb{Z}_{4n}^\times$ sends $\theta_{i,j} \mapsto i^2$ by, for example, [17, Proposition 1.1].

The following was noted by Davis [12] and Dyer [13, Note (b)]. It would be interesting to know, as was asked by Davis, whether this holds in the case $\sigma_k(G) \neq 0$.

Proposition 7.1. *If G has free period k , then $S \circ \psi_k = 0$, i.e. $(I, \psi_k(\theta))$ is stably free for all $\theta \in \text{Aut}(G)$.*

Proof. Note that Theorems 6.8 and 6.9 each show that $[P] = [(I, \psi_k(\theta)) \otimes P_\theta]$ for all $P \in P(\mathbb{Z}G)$ such that $\sigma_k(G) = [P] + T_G$. By Lemma 3.5, the composition

$$\text{Aut}(G) \xrightarrow{\psi_k} (\mathbb{Z}/|G|)^\times \xrightarrow{S} T_G \leq C(\mathbb{Z}G)$$

is given by $S \circ \psi_k : \theta \mapsto [P] - [P_\theta]$ which is well-defined since θ gives a well-defined action on $C(\mathbb{Z}G)$. By Lemma 6.1, we have that $(\mathbb{Z}G)_\theta \cong \mathbb{Z}G$ and so the composition is trivial in the case where $\sigma_k(G) = 0$. \square

We say that a finite group G has *weak cancellation* if every stably free Swan module is free. The following was asked by Dyer in [13, p266] and later appeared as Problem A4 in Wall's 1979 Problems List [45].

Question 7.2. *Does there exist G with periodic cohomology and $r \in (\mathbb{Z}/|G|)^\times$ such that (I, r) is stably free but not free?*

This is equivalent to asking whether every group with periodic cohomology has weak cancellation and is still open, even for arbitrary finite groups. There are two important consequences that a negative answer to Question 7.2 would have.

Firstly, recall the following question from the introduction. Note that, if $(I, \psi_k(\theta))$ is free, then the action described in Theorem B has the simpler form $P \mapsto P_\theta$.

Question 7.3. *Does there exist G with k -periodic cohomology and $\theta \in \text{Aut}(G)$ for which $(I, \psi_k(\theta))$ is not free?*

It follows from Proposition 7.1 that, if G has free period k and has weak cancellation, then $(I, \psi_k(\theta)) \cong \mathbb{Z}G$ for all $\theta \in \text{Aut}(G)$. In particular, if Question 7.2 has a negative answer, then the only groups for which the action in Theorem B might not have the form $P \mapsto P_\theta$ are the groups with $\sigma_k(G) \neq 0$.

Secondly, consider the following:

Question 7.4. *Let $n \geq 2$, let G be finite and let X, Y be finite (G, n) -complexes with $\chi(X) = \chi(Y)$. Then $X \vee rS^n \simeq Y \vee rS^n$ for some r . Does $r = 1$ always work?*

This is equivalent to asking whether $\text{HT}(G, n)$ is a fork when G is finite. The case where n is even was proven by Browning [6], and also follows by combining Corollary 4.7 and Proposition 5.1. When n is odd, this is known to hold provided G does not have k -periodic cohomology for any $k \mid n + 1$. If G has k -periodic cohomology for $k \mid n + 1$, then this holds provided G has weak cancellation (see [13, p276-277]). In particular, if Question 7.2 has a negative answer, then Question 7.4 has an affirmative answer. Note that the corresponding question for infinite groups is also still open (see [35, Problem 2]).

8. MILNOR SQUARES AND THE CLASSIFICATION OF PROJECTIVE MODULES

Given the observations in the previous section, the primary obstacle to computing sufficiently interesting examples of $\text{HT}(G, n)$ and $\text{PHT}(G, n)$ for our groups is the classification of projective $\mathbb{Z}G$ -modules.

One method to classify projective R -modules over a ring R is to relate this to the classification of projective modules over simpler rings using Milnor squares. In this section, we will present a refinement of the basic theory of Milnor squares which will also allow us to determine how a ring automorphism $\alpha \in \text{Aut}(R)$ acts on the class of projective R -modules. We will then apply these methods in Section 9.

Suppose R and S are rings and $f : R \rightarrow S$ is a ring homomorphism. We can use this to turn S into an (S, R) -bimodule, with right-multiplication by $r \in R$ given by $x \cdot r = xf(r)$ for any $x \in S$. If M is an R -module, we can define the *extension of scalars* of M by f as the tensor product

$$f_{\#}(M) = S \otimes_R M$$

since S as a right R -module and M as a left R -module, and we consider this as a left S -module where left-multiplication by $s \in S$ is given by $s \cdot (x \otimes m) = (sx) \otimes m$ for any $x \in S$ and $m \in M$. This comes equipped with maps of abelian groups

$$f_* : M \rightarrow f_{\#}(M)$$

sending $m \mapsto 1 \otimes m$, and defines a covariant functor from R -modules to S -modules [10, p227]. It has the following basic properties which follow from the standard properties of tensor products such as associativity [26, p145].

Lemma 8.1. *Let $f : R \rightarrow S$ and $g : S \rightarrow T$ be ring homomorphisms and let M and N be R -modules. Then*

$$(i) \quad f_{\#}(M \oplus N) \cong f_{\#}(M) \oplus f_{\#}(N)$$

- (ii) $f_{\#}(R) \cong S$
- (iii) $(g \circ f)_{\#}(M) \cong (g_{\#} \circ f_{\#})(M)$.

If $P(R)$ denotes the set of projective R -modules, then the first two properties show that $f_{\#}$ induces a map $f_{\#} : P(R) \rightarrow P(S)$ which restricts to each stable class.

Recall that, if R, R_1, R_2 and R_0 are rings, then a pullback diagram

$$\mathcal{R} = \begin{array}{ccc} R & \xrightarrow{i_2} & R_2 \\ \downarrow i_1 & & \downarrow j_2 \\ R_1 & \xrightarrow{j_1} & R_0 \end{array}$$

is a Milnor square if either j_1 or j_2 are surjective. If $P_1 \in P(R_1)$, $P_2 \in P(R_2)$ are such that there is a R_0 -module isomorphism $h : (j_1)_{\#}(P_1) \rightarrow (j_2)_{\#}(P_2)$, then define

$$M(P_1, P_2, h) = \{(x, y) \in P_1 \times P_2 : h((j_1)_*(x)) = (j_2)_*(y)\} \leq P_1 \times P_2,$$

which is an R -module where multiplication by $r \in R$ is given by $r \cdot (x, y) = ((i_1)_*(r)x, (i_2)_*(r)y)$. It was shown by Milnor that $M(P_1, P_2, h)$ is projective [32, Theorem 2.1]. Let $\text{Aut}_R(P)$ denote the set of R -module automorphisms of an R -module P . The main result on Milnor squares is as follows.

Theorem 8.2. *Suppose \mathcal{R} is a Milnor square and $P_i \in P(R_i)$ for $i = 0, 1, 2$ are such that $P_0 \cong (j_1)_{\#}(P_1) \cong (j_2)_{\#}(P_2)$ as R_0 -modules. Then there is a one-to-one correspondence*

$$\text{Aut}_{R_1}(P_1) \backslash \text{Aut}_{R_0}(P_0) / \text{Aut}_{R_2}(P_2) \leftrightarrow \{P \in P(R) : (i_1)_{\#}(P) \cong P_1, (i_2)_{\#}(P) \cong P_2\}$$

given by sending a coset $[h]$ to $M(P_1, P_2, h)$ for any representative h .

Now suppose $\alpha \in \text{Aut}(R)$. If M is an R -module, define M_{α} as the R -module whose abelian group is that of M but whose R -action is given by $r \cdot m = \alpha(r)m$ for $r \in R$ and $m \in M$. For example, if $R = \mathbb{Z}G$, then $\theta \in \text{Aut}(G)$ induces a map $\theta \in \text{Aut}(\mathbb{Z}G)$ and M_{θ} coincides with the definition given earlier.

This is a special case of restriction of scalars, but can also be viewed as a part of extension of scalars as follows.

Lemma 8.3. *Let R be a ring and let $\alpha \in \text{Aut}(R)$. If M is an R -module, then there is an isomorphism of R -modules*

$$\psi : M_{\alpha} \rightarrow (\alpha^{-1})_{\#}(M)$$

given by sending $m \mapsto 1 \otimes m$.

From this, it is clear that this action has basic properties which are analogous to Lemma 6.1. The following is then immediate by combining Lemmas 8.1 and 8.3.

Corollary 8.4. *Suppose $f : R \rightarrow S$ is a ring homomorphism and $\alpha \in \text{Aut}(R)$, $\beta \in \text{Aut}(S)$ are such that $f \circ \alpha = \beta \circ f$. If M is an R -module, then*

$$f_{\#}(M_{\alpha}) \cong f_{\#}(M)_{\beta}.$$

We can turn the set of Milnor squares into a category with morphism defined as follows. If $\mathcal{R}, \mathcal{R}'$ are Milnor squares, then a morphism is a quadruple

$$\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) : \mathcal{R} \rightarrow \mathcal{R}'$$

where $\alpha : R \rightarrow R'$ and $\alpha_i : R_i \rightarrow R'_i$ such that there is a commutative diagram as follows

$$\begin{array}{ccccc}
 R & \xrightarrow{\quad} & R_2 & & \\
 \downarrow & \searrow \alpha & \downarrow & \searrow \alpha_2 & \\
 & R & \xrightarrow{\quad} & R_2 & \\
 & \downarrow & \downarrow & \downarrow & \\
 R_1 & \xrightarrow{\quad} & R_0 & & \\
 & \searrow \alpha_1 & \searrow \alpha_0 & & \\
 & R_1 & \xrightarrow{\quad} & R_0 &
 \end{array}$$

Let $\text{Aut}(\mathcal{R})$ denote the set of automorphisms of a Milnor square \mathcal{R} , i.e. the set of isomorphisms $\hat{\alpha} : \mathcal{R} \rightarrow \mathcal{R}$.

Lemma 8.5. *Let \mathcal{R} is a Milnor square, let $P_1 \in P(R_1)$, $P_2 \in P(R_2)$ be such that there is an R_0 -module isomorphism $h : (j_1)_\#(P_1) \rightarrow (j_2)_\#(P_2)$. If $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$, then*

$$M(P_1, P_2, h)_\alpha \cong M((P_1)_{\alpha_1}, (P_2)_{\alpha_2}, h)$$

where, on the right, we view h as a map $h : (j_1)_\#(P_1)_{\alpha_0} \rightarrow (j_2)_\#(P_2)_{\alpha_0}$.

Proof. Let $P = M(P_1, P_2, h)$ so that, by Theorem 8.2, we have that $(i_1)_\#(P) \cong P_1$ and $(i_2)_\#(P) \cong P_2$. It is easy to see directly that the natural map

$$M((i_1)_\#(P), (i_2)_\#(P), h) \rightarrow M((i_1)_\#(P_\alpha), (i_2)_\#(P_\alpha), h)$$

is an isomorphism. We are then done by applying Corollary 8.4. \square

This has the following simplification when P_1 and P_2 are free of rank one. Here we will use the identification $\text{Aut}_{R_0}(R_0) \cong R_0^\times$ which sends $h : R_0 \rightarrow R_0$ to $h(1) \in R_0^\times$.

Lemma 8.6. *Let \mathcal{R} is a Milnor square and let $u \in R_0^\times$. If $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$, then*

$$M(R_1, R_2, u)_\alpha \cong M(R_1, R_2, \alpha_0^{-1}(u)).$$

Proof. Fix identifications $\psi_i : (j_i)_\#(R_i) \rightarrow R_0$ and let $h : (j_1)_\#(R_1) \rightarrow (j_1)_\#(R_1)$ be such that $(\psi_2 \circ h \circ \psi_1^{-1})(1) = u \in R_0^\times$. By Lemma 8.5, we have that

$$M(R_1, R_2, h)_\alpha \cong M((R_1)_{\alpha_1}, (R_2)_{\alpha_2}, h)$$

where $h : ((j_1)_\#(R_1))_{\alpha_0} \rightarrow ((j_1)_\#(R_1))_{\alpha_0}$ coincides with h as a map of abelian groups. For $i = 0, 1, 2$, let $c_i : R_i \rightarrow (R_i)_{\alpha_i}$ be the isomorphism which sends $1 \mapsto 1$. Then it is easy to see that the following diagram commutes for $i = 1, 2$, where $f : (j_i)_\#((R_i)_{\alpha_i}) \rightarrow ((j_i)_\#(R_i))_{\alpha_0}$ is the isomorphism coming from Corollary 8.4

$$\begin{array}{ccccc}
 (j_i)_\#(R_i) & \xrightarrow{1 \otimes c_i} & (j_i)_\#((R_i)_{\alpha_i}) & \xrightarrow{f} & ((j_i)_\#(R_i))_{\alpha_0} \\
 \downarrow \psi_i & & & & \downarrow \psi_i \\
 R_0 & \xrightarrow{c_0} & (R_0)_{\alpha_0} & &
 \end{array}$$

Using the isomorphisms c_i for $i = 1, 2$, we get that

$$M((R_1)_{\alpha_1}, (R_2)_{\alpha_2}, h) \cong M(R_1, R_2, h_0)$$

where $h_0 : (j_1)_\#(R_1) \rightarrow (j_2)_\#(R_2)$ induces $h : ((j_1)_\#(R_1))_{\alpha_0} \rightarrow ((j_1)_\#(R_1))_{\alpha_0}$ via $f \circ (1 \otimes c_i)$. Let $u_0 = (\psi_2 \circ h_0 \circ \psi_1^{-1})(1) \in R_0^\times$. Then, since the above diagram

commutes, we get the following commutative diagram

$$\begin{array}{ccc}
 R_0 & \xrightarrow{\psi_2 \circ h \circ \psi_1^{-1}} & R_0 \\
 \downarrow c_0 & & \downarrow c_0 \\
 (R_0)_{\alpha_0} & \xrightarrow{\psi_2 \circ h_0 \circ \psi_1^{-1}} & (R_0)_{\alpha_0}
 \end{array}
 \quad
 \begin{array}{ccc}
 1 & \longrightarrow & u_0 \\
 \downarrow & & \downarrow \\
 1 & \longrightarrow & \alpha_0(u_0)
 \end{array}$$

This implies that $u = \alpha_0(u_0)$ and so $u_0 = \alpha_0^{-1}(u)$, as required. \square

We conclude this section by combining Theorem 8.2 and Lemma 8.6. If \mathcal{R} is a Milnor square, we say that $\alpha \in \text{Aut}(R)$ *extends across* \mathcal{R} if there exists $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$.

Corollary 8.7. *Let \mathcal{R} is a Milnor square such that every $\alpha \in \text{Aut}(R)$ extends across \mathcal{R} . Then there is a one-to-one correspondence*

$$R_1^\times \backslash (R_0^\times / \text{Aut}(R)) / R_2^\times \leftrightarrow \{P \in P(R) : (i_1)_\#(P) \cong R_1, (i_2)_\#(P) \cong R_2\} / \text{Aut}(R)$$

where $\alpha \in \text{Aut}(R)$ acts on R_0^\times by sending $r \mapsto \alpha_0^{-1}(r)$ for $r \in R_0^\times$ and any $\alpha_0 \in \text{Aut}(R_0)$ such that $\hat{\alpha} = (\alpha, \alpha_1, \alpha_2, \alpha_0) \in \text{Aut}(\mathcal{R})$.

9. EXAMPLE: QUATERNION GROUPS

The aim of this section will be to illustrate how Theorems A and B can be combined with the known techniques to classify projective $\mathbb{Z}G$ -modules to obtain a detailed classification of finite (G, n) -complexes up to homotopy equivalence.

For $k \geq 2$, recall that the quaternion group of order $4k$ has presentation:

$$Q_{4k} = \langle x, y \mid x^k = y^2, yxy^{-1} = x^{-1} \rangle.$$

It is a finite 3-manifold group and so has free period 4. For $n \geq 2$ even, Theorem A and Proposition 5.1 imply that $\text{PHT}(Q_{4k}, n) \cong [\mathbb{Z}Q_{4k}] = \bigcup_{r \geq 1} \text{SF}_r(\mathbb{Z}Q_{4k})$ where $\text{SF}_r(\mathbb{Z}Q_{4k})$ is the set of stably free $\mathbb{Z}Q_{4k}$ -modules of rank $r \geq 1$.

Since stably free $\mathbb{Z}G$ -modules of rank ≥ 2 are free for G finite [40] (or since $\text{PHT}(G, n)$ is a fork by Corollary 4.7), it remains to compute $\text{SF}_1(\mathbb{Z}Q_{4k})$. This was completed by Swan for $k \leq 9$ [43, Theorem III]. For $k \leq 7$, he showed that $|\text{SF}_1(\mathbb{Z}Q_{4k})| = 1$ for $2 \leq k \leq 5$, $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{28})| = 2$. It also follows from his classification that $\mathbb{Z}Q_{4k}$ has weak cancellation in all these cases and so the action of $\theta \in \text{Aut}(Q_{4k})$ on $[\mathbb{Z}Q_{4k}]$ sends $P \mapsto P_\theta$ (see Section 7).

In the case Q_{28} , the action of $\text{Aut}(Q_{28})$ on $[\mathbb{Z}Q_{28}]$ is trivial since $(\mathbb{Z}Q_{28})_\theta \cong \mathbb{Z}Q_{28}$ for all $\theta \in \text{Aut}(Q_{28})$ and so this must also hold for the non-free stably free module also. The main result of this section will be to compute the action in the case Q_{24} .

Theorem 9.1. *$\text{Aut}(Q_{24})$ acts non-trivially on $[\mathbb{Z}Q_{24}]$. More specifically, we have $|\text{SF}_1(\mathbb{Z}Q_{24})| = 3$ and $|\text{SF}_1(\mathbb{Z}Q_{24}) / \text{Aut}(Q_{24})| = 2$.*

All this can be summarised in the following table, which gives the structure of $\text{PHT}(G, n)$ and $\text{HT}(G, n)$ when $n \neq 2$ is even. These graded trees are both forks by Corollary 4.7 and each dot represents a finite (G, n) -complex at the minimal level.

Remark 9.2. This also holds in the case $n = 2$ provided G has the D2 property. This holds trivially in the cases $Q_8, Q_{12}, Q_{16}, Q_{20}$ and is otherwise only known to be true in the case Q_{28} by [36, Theorem 7.7] using the presentation of Mannan-Popiel [29].

G	Q_8	Q_{12}	Q_{16}	Q_{20}	Q_{24}	Q_{28}
$\text{PHT}(G, n)$	•	•	•	•	•••	••
$\text{HT}(G, n)$	•	•	•	•	••	••

FIGURE 2. Minimal complexes for any n even with $n \neq 2$

We will now proceed to the proof of Theorem 9.1. First let x and y be generators for Q_{24} in the presentation given above. Let $\Lambda = \mathbb{Z}Q_{24}/(x^6 + 1)$ and note that the quotient map $f : \mathbb{Z}Q_{24} \rightarrow \Lambda$ induces a map

$$f_{\#} : \text{SF}_1(\mathbb{Z}Q_{24}) \rightarrow \text{SF}_1(\Lambda)$$

by Lemma 8.1. This is a bijection by the proof of [43, Theorem 11.14].

Now note that the factorisation $x^6 + 1 = (x^2 + 1)(x^4 - x^2 + 1)$ implies that the ideals $I = (x^2 + 1)$ and $J = (x^4 - x^2 + 1)$ have $I \cap J = (x^6 + 1)$ and $I + J = (3, x^2 + 1)$. It follows from [11, Example 42.3] that we have a pullback diagram

$$\begin{array}{ccc} \Lambda & \longrightarrow & \mathbb{Z}Q_{24}/(x^4 - x^2 + 1) \\ \downarrow & & \downarrow \\ \mathbb{Z}Q_{24}/(x^2 + 1) & \longrightarrow & \mathbb{F}_3Q_{24}/(x^2 + 1) \end{array}$$

which is a Milnor square since all maps are surjective.

For a field \mathbb{F} , let $\mathbb{H}_{\mathbb{F}} = \mathbb{F}[i, j]$ denote the quaternions over \mathbb{F} and we define $\mathbb{H}_{\mathbb{Z}} = \mathbb{Z}[i, j]$ and $\mathbb{Z}[\zeta_{12}, j]$ to be subrings of $\mathbb{H}_{\mathbb{R}}$, where $\zeta_{12} = e^{\frac{2\pi i}{12}}$ is the 12th root of unity in the i direction. The following is straightforward to check that there are isomorphisms of rings

$$\begin{aligned} \phi_1 : \mathbb{H}_{\mathbb{Z}} &\rightarrow \mathbb{Z}Q_{24}/(x^2 + 1), & \phi_2 : \mathbb{Z}[\zeta_{12}, j] &\rightarrow \mathbb{Z}Q_{24}/(x^4 - x^2 + 1) \\ i &\mapsto x, j \mapsto y & \zeta_{12} &\mapsto x, j \mapsto y. \end{aligned}$$

Using this, we can rewrite the Milnor square above as follows

$$\begin{array}{ccc} \Lambda & \xrightarrow{i_2} & \mathbb{Z}[\zeta_{12}, j] \\ \downarrow i_1 & & \downarrow j_2 \\ \mathbb{H}_{\mathbb{Z}} & \xrightarrow{j_1} & \mathbb{H}_{\mathbb{F}_3} \end{array} \quad \begin{array}{ccc} x, y & \longmapsto & \zeta_{12}, j \\ \downarrow & & \downarrow \\ i, j & \longmapsto & i, j \end{array}$$

Now note that, by [43, Lemma 8.14], the induced map $(i_2)_* : C(\Lambda) \rightarrow C(\mathbb{Z}[\zeta_{12}, j])$ is an isomorphism. It also follows from [43, p84] that the rings $\mathbb{H}_{\mathbb{Z}}$ and $\mathbb{Z}[\zeta_{12}, j]$ both have stably free cancellation, i.e. that every stably free module is free. It follows easily that

$$\text{SF}_1(\Lambda) = \{P \in P(\Lambda) : (i_1)_{\#}(P) \cong \mathbb{H}_{\mathbb{Z}}, (i_2)_{\#}(P) \cong \mathbb{Z}[\zeta_{12}, j]\}.$$

In particular, by combining with Theorem 8.2, we get that there is a bijection

$$\text{SF}_1(\Lambda) \leftrightarrow \mathbb{H}_{\mathbb{Z}}^{\times} \backslash \mathbb{H}_{\mathbb{F}_3}^{\times} / \mathbb{Z}[\zeta_{12}, j]^{\times}.$$

Lemma 9.3. $\mathbb{H}_{\mathbb{Z}}^{\times} \backslash \mathbb{H}_{\mathbb{F}_3}^{\times} / \mathbb{Z}[\zeta_{12}, j]^{\times} = \{[1], [1 + j], [1 + k]\}.$

Proof. If $N : \mathbb{H}_{\mathbb{F}_3} \rightarrow \mathbb{F}_3$ is the norm, then $\mathbb{H}_{\mathbb{F}_3}^{\times} = N^{-1}(\pm 1)$. Now note that $\mathbb{H}_{\mathbb{Z}}^{\times} = \{\pm 1, \pm i, \pm j, \pm k\}$, and it is easy to check that

$$\mathbb{H}_{\mathbb{Z}}^{\times} \backslash \mathbb{H}_{\mathbb{F}_3}^{\times} = \{[1], [1 + i], [1 + j], [1 + k], [1 + i + j + k], [1 - i - j - k]\}.$$

By [27, Lemma 7.5 (b)], we have that $\mathbb{Z}[\zeta_{12}, j]^\times = \mathbb{Z}[\zeta_{12}]^\times \cdot \langle j \rangle$ and so it remains to determine

$$\mathrm{Im}(\mathbb{Z}[\zeta_{12}, j]^\times \rightarrow \mathbb{H}_\mathbb{Z}^\times \setminus \mathbb{H}_{\mathbb{F}_3}^\times) = \mathrm{Im}(\mathbb{Z}[\zeta_{12}]^\times \rightarrow \mathbb{H}_\mathbb{Z}^\times \setminus \mathbb{H}_{\mathbb{F}_3}^\times) \subseteq \{[1], [1+i]\},$$

where the last inclusion follows since $\zeta_{12} \mapsto i$ and $\mathbb{H}_\mathbb{Z}^\times \setminus \langle 1, i \rangle = \{[1], [1+i]\}$.

Consider the n th cyclotomic polynomial

$$\Phi_n(x) = \prod_{k \in \mathbb{Z}_n^\times} (x - \zeta_n^k).$$

It is well-known, and can be shown using Möbius inversion, that $\Phi_n(1) = 1$ if n is not a prime power. In particular, $\Phi_{12}(1) = 1$ and this implies that $1 - \zeta_{12} \in \mathbb{Z}[\zeta_{12}]^\times$. Hence $[1+i] = [1-i] \in \mathrm{Im}(\mathbb{Z}[\zeta_{12}]^\times \rightarrow \mathbb{H}_\mathbb{Z}^\times \setminus \mathbb{H}_{\mathbb{F}_3}^\times)$. The result then follows since

$$j(1+i+j+k)(1+i) = 1+k, \quad -j(1-i-j-k)(1+i) = 1+j$$

implies that $[1+j] = [1-i-j-k]$, $[1+k] = [1+i+j+k]$ in $\mathbb{H}_\mathbb{Z}^\times \setminus \mathbb{H}_{\mathbb{F}_3}^\times / \mathbb{Z}[\zeta_{12}, j]^\times$. \square

This implies that $|\mathrm{SF}_1(\mathbb{Z}Q_{24})| = 3$, which recovers the result of Swan. In order to determine the action of $\mathrm{Aut}(Q_{24})$ on $\mathrm{SF}_1(\mathbb{Z}Q_{24})$, first recall from Section 7 that

$$\mathrm{Aut}(Q_{24}) = \{\theta_{a,b} : x \mapsto x^a, y \mapsto x^b y \mid a \in \mathbb{Z}_{12}^\times, b \in \mathbb{Z}_{12}\}.$$

If \mathcal{R} denote the Milnor square defined above, then the following is easy to check.

Lemma 9.4. *If $a \in \mathbb{Z}_{12}^\times$, $b \in \mathbb{Z}_{12}$, then $\theta_{a,b} \in \mathrm{Aut}(Q_{24})$ extends to a Milnor square automorphism*

$$\hat{\theta}_{a,b} = (\theta_{a,b}, \theta_{a,b}^1, \theta_{a,b}^2, \bar{\theta}_{a,b}) \in \mathrm{Aut}(\mathcal{R}).$$

If $a = 2a_0 + 1$, $\bar{\theta}_{a,b} \in \mathrm{Aut}(\mathbb{H}_{\mathbb{F}_3})$ is defined by

$$\bar{\theta}_{a,b} : i \mapsto i^a = (-1)^{a_0} i, \quad j \mapsto j^b = \begin{cases} (-1)^{b_0} j, & \text{if } b = 2b_0 + 1 \\ (-1)^{b_0} k, & \text{if } b = 2b_0. \end{cases}$$

In particular, by Corollary 8.7 and Lemma 9.3, there is a bijection

$$\mathrm{SF}_1(\mathbb{Z}Q_{24}) / \mathrm{Aut}(Q_{24}) \leftrightarrow \{[1], [1+j], [1+k]\} / \mathrm{Aut}(Q_{24})$$

where $\theta_{a,b} \in \mathrm{Aut}(Q_{24})$ acts on the double cosets via the action described in Lemma 9.4. In particular

$$\bar{\theta}_{a,b}([1+j]) = \begin{cases} [1 + (-1)^{b_0} j] = [1+j], & \text{if } b = 2b_0 + 1 \\ [1 + (-1)^{b_0} k] = [1+k], & \text{if } b = 2b_0 \end{cases}$$

and so $\bar{\theta}_{a,b}$ acts non-trivially when b is even. Hence $|\mathrm{SF}_1(\mathbb{Z}Q_{24}) / \mathrm{Aut}(Q_{24})| = 2$. This completes the proof of Theorem 9.1.

ACKNOWLEDGEMENTS

I would like to thank F. E. A. Johnson for his guidance and many interesting conversations. I would also like to thank Ian Hambleton, Jonathan Hillman and Wajid Mannan for helpful comments and references. This work was supported by the UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/N509577/1 as well as the Heilbronn Institute for Mathematical Research.

REFERENCES

- [1] M. Powell A. Conway, D. Crowley and J. Sixt, *Simply-connected manifolds with large homotopy stable classes*, (2021), arXiv:2109.00654.

- [2] ———, *Stably diffeomorphic manifolds and the realisation of modified surgery obstructions*, (2021), arXiv:2109.05632.
- [3] M. Bestvina and N. Brady, *Morse theory and finiteness properties of groups*, Invent. Math. **129** (1997), no. 3, 445–470.
- [4] R. Bieri, *Homological dimension of discrete groups*, second ed., Queen Mary College Mathematics Notes, Queen Mary College, Department of Pure Mathematics, London, 1981.
- [5] W. J. Browning, *Finite CW-complexes of cohomological dimension 2 with finite abelian π_1* , ETH preprint - unpublished, 1979.
- [6] ———, *Homotopy types of certain finite CW-complexes with finite fundamental group*, Ph.D. thesis, Cornell University, 1979.
- [7] H. Cartan and S. Eilenberg, *Homological algebra*, vol. 19, Princeton Mathematics Series, 1956.
- [8] W. H. Cockroft and R. G. Swan, *On the homotopy types of certain two-dimensional complexes*, Proc. London Math Soc. **11** (1961), no. 3, 194–202.
- [9] J. M. Cohen, *Complexes dominated by a 2-complex*, Topology **16** (1977), 409–415.
- [10] C. W. Curtis and I. Reiner, *Methods of representation theory: with applications to finite groups and orders*, vol. 1, Wiley Classics Library, 1981.
- [11] ———, *Methods of representation theory: With applications to finite groups and orders*, vol. 2, Wiley Classics Library, 1987.
- [12] J. F. Davis, *A remark on “homotopy equivalences and free modules”*, Topology **22** (1983), 487–488.
- [13] M. N. Dyer, *Homotopy classification of (π, m) -complexes*, J. Pure Appl. Algebra **7** (1976), 249–282.
- [14] ———, *Non-minimal roots in homotopy trees*, Pac. J. Math. **80** (1979), 371–380.
- [15] M. N. Dyer and A. J. Sieradski, *A homotopy classification of 2-complexes with finite cyclic fundamental group*, Bull. Amer. Math. Soc. **79** (1973), no. 1, 75–77.
- [16] ———, *Trees of homotopy types of two-dimensional CW complexes II*, Trans. Amer. Math. Soc. **205** (1975), 115–125.
- [17] M. Golasinski and D. L. Gonçalves, *Spherical space forms: homotopy types and self-equivalences*, Algebraic Topology: Categorical Decomposition Techniques, Progress in Mathematics 215, 2003, pp. 153–165.
- [18] I. Hambleton, *Two remarks on Wall’s D2 problem*, Math. Proc. Cambridge Philos. Soc. **167** (2019), no. 2, 361–368.
- [19] I. Hambleton and M. Kreck, *Cancellation of lattices and finite two-complexes*, J. reine angew. Math **442** (1993), 91–109.
- [20] I. Hambleton, S. Pamuk, and E. Yalçin, *Equivariant CW-complexes and the orbit category*, Comment. Math. Helv. **88** (2013), no. 2, 369–425.
- [21] H. Jacobinski, *Genera and decompositions of lattices over orders*, Acta mathematica **121** (1968), 1–29.
- [22] F. E. A. Johnson, *Explicit homotopy equivalences in dimension two*, Math. Proc. Camb. Phil. Soc. **131** (2002), 411–430.
- [23] ———, *Stable modules and the D(2)-problem*, London Math. Soc. Lecture Note Ser., vol. 301, Cambridge University Press, 2003.
- [24] M. Kreck and J. A. Schafer, *Classification and stable classification of manifolds: some examples*, Comment. Math. Helv. **59** (1984), 12–38.
- [25] P. A. Linnell, *Minimal free resolutions and (G, n) -complexes for finite abelian groups*, Proc. London Math. Soc. **66** (1993), no. 3, 303–326.
- [26] S. MacLane, *Homology*, Springer Berlin-Göttingen-Heidelberg, 1963.
- [27] B. Magurn, R. Oliver, and L. Vaserstein, *Units in Whitehead groups of finite groups*, J. Algebra **84** (1983), 324–360.
- [28] W. H. Mannan, *The D(2) property for D8*, Algebr. Geom. Topol. **7** (2007), 517–528.
- [29] W. H. Mannan and T. Popiel, *An exotic presentation of Q_{28}* , Algebr. Geom. Topol. **21** (2021), no. 4, 2065–2084.
- [30] W. Metzler, *Über den homotopietyp zweidimensionaler CW-komplexe und elementartransformationen bei darstellungen von gruppen durch erzeugende und definierende relationen*, J. reine u. angew. Math **285** (1976), 7–23.

- [31] R. J. Milgram, *Evaluating the Swan finiteness obstruction for periodic groups*, Algebraic and geometric topology (New Brunswick, N.J., 1983), Lecture Notes in Math. 1126 (Berlin), Springer, 1985, pp. 127–158.
- [32] J. Milnor, *Introduction to algebraic K-theory*, Ann. of Math. Stud., vol. 72, Princeton University Press, 1971.
- [33] J. Nicholson, *Cancellation for (G, n) -complexes and the Swan finiteness obstruction*, (2020), arXiv:2005.01664.
- [34] ———, *A cancellation theorem for modules over integral group rings*, Math. Proc. Cambridge Philos. Soc. **171** (2021), no. 2, 317–327.
- [35] ———, *The homotopy type of a finite 2-complex with non-minimal Euler characteristic*, (2021), arXiv:2108.02220.
- [36] ———, *On CW-complexes over groups with periodic cohomology*, Trans. Amer. Math. Soc. **374** (2021), no. 9, 6531–6557.
- [37] A. J. Sieradski, *A semigroup of simply homotopy types*, Math. Zeit. **153** (1977), 135–148.
- [38] A. J. Sieradski and M. N. Dyer, *Distinguishing arithmetic for certain stably isomorphic modules*, J. Pure Appl. Alg. **15** (1979), 199–217.
- [39] J. Stallings, *A finitely presented group whose 3-dimensional integral homology is not finitely generated*, Amer. J. Math. **85** (1963), 541–543.
- [40] R. G. Swan, *Induced representations and projective modules*, Ann. of Math **71** (1960), no. 2, 552–578.
- [41] ———, *Periodic resolutions for finite groups*, Ann. of Math **72** (1960), no. 2, 267–291.
- [42] ———, *Minimal resolutions for finite groups*, Topology **4** (1965), 193–220.
- [43] ———, *Projective modules over binary polyhedral groups*, J. Reine Angew. Math **342** (1983), 66–172.
- [44] C. T. C. Wall, *Finiteness conditions for CW complexes*, Ann. of Math **81** (1965), 56–69.
- [45] ———, *List of problems*, Homological group theory (Proc. Sympos., Durham, 1977), London Math. Soc. Lecture Note Ser., vol. 36, Cambridge University Press, 1979.
- [46] ———, *Periodic projective resolutions*, Proc. London Math. Soc. **39** (1979), no. 3, 509–553.
- [47] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, 1994.
- [48] J. H. C. Whitehead, *Simplicial spaces, nuclei and m -groups*, Proc. London Math. Soc. **45** (1939), no. 2, 243–327.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, LONDON, SW7 2AZ, U.K.

Email address: john.nicholson@imperial.ac.uk