

Alternating Euler T -sums and Euler \tilde{S} -sums

Weiping Wang^{a,*} Ce Xu^{b,†}

a. School of Science, Zhejiang Sci-Tech University,
Hangzhou 310018, P.R. China

b. School of Mathematics and Statistics, Anhui Normal University,
Wuhu 241000, P.R. China

Abstract In this paper, we study the alternating Euler T -sums and related sums by using the method of contour integration. We establish the explicit formulas for all linear and quadratic (alternating) Euler T -sums and related sums. Some interesting new consequences and illustrative examples are considered.

Keywords: Multiple zeta values; multiple t -values; multiple T -values; odd harmonic numbers; Euler T -sums.

AMS Subject Classifications (2020): 11M99; 11M06; 11M32.

1 Introduction and Notations

In our previous paper [7], we introduced and studied the following two variants of the classical Euler sums,

$$T_{p_1 p_2 \dots p_k, q} := \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p_1)} h_{n-1}^{(p_2)} \dots h_{n-1}^{(p_k)}}{(n - 1/2)^q} \quad \text{and} \quad \tilde{S}_{p_1 p_2 \dots p_k, q} := \sum_{n=1}^{\infty} \frac{h_n^{(p_1)} h_n^{(p_2)} \dots h_n^{(p_k)}}{n^q},$$

where $q > 1$, $p_1 \leq p_2 \leq \dots \leq p_k$ are positive integers, and the quantity $w := p_1 + \dots + p_r + q$ is called the weight and the quantity r is called the degree (or order). We often refer these sums as the the Euler T -sums and Euler \tilde{S} -sums, respectively. Here $h_n^{(p)}$ stands for odd harmonic number of order p defined by

$$h_n^{(p)} := \sum_{k=1}^n \frac{1}{(k - 1/2)^p}, \quad h_n \equiv h_n^{(1)} \quad \text{and} \quad h_0^{(p)} := 0.$$

The Euler T -sums and Euler \tilde{S} -sums can be seen as variants of classical Euler sums [1]

$$S_{p_1 p_2 \dots p_k, q} := \sum_{n=1}^{\infty} \frac{H_n^{(p_1)} H_n^{(p_2)} \dots H_n^{(p_k)}}{n^q},$$

where $H_n^{(p)}$ stands for the p -th generalized harmonic number, which is defined by

$$H_n^{(p)} := \sum_{k=1}^n \frac{1}{k^p}, \quad H_n \equiv H_n^{(1)} \quad \text{and} \quad H_0^{(p)} := 0.$$

*Email: wpingwang@yahoo.com, wpingwang@zstu.edu.cn

†Email: 19020170155420@stu.xmu.edu.cn

Like classical Euler sums, the Euler T -sums and Euler \tilde{S} -sums can be evaluated by using the method of contour integration developed by Flajolet and Salvy [1]. In [7], we establish many explicit evaluations of Euler T -sums and Euler \tilde{S} -sums via $\log(2)$, multiple zeta and t -values. Here, for positive integers p_1, \dots, p_k with $p_1 > 1$, the multiple zeta value (MZV for short) [2, 8] and multiple t -values (MtVs for short) [3] are defined by

$$\zeta(p_1, p_2, \dots, p_k) := \sum_{n_1 > \dots > n_k \geq 1} \frac{1}{n_1^{p_1} n_2^{p_2} \cdots n_k^{p_k}}$$

and

$$\begin{aligned} t(p_1, p_2, \dots, p_k) &:= \sum_{\substack{n_1 > \dots > n_k \geq 1 \\ n_i \text{ odd}}} \frac{1}{n_1^{p_1} n_2^{p_2} \cdots n_k^{p_k}} \\ &= \sum_{n_1 > \dots > n_k \geq 1} \frac{1}{(2n_1 - 1)^{p_1} (2n_2 - 1)^{p_2} \cdots (2n_k - 1)^{p_k}}. \end{aligned}$$

As its normalized version,

$$\tilde{t}(p_1, p_2, \dots, p_k) := 2^{p_1 + p_2 + \dots + p_k} t(p_1, p_2, \dots, p_k).$$

In above definitions of MZVs and MtVs, we put a bar on top of p_j ($j = 1, \dots, k$) if there is a sign $(-1)^{n_j}$ appearing in the denominator on the right. Which (one or more the p_j barred) are called the alternating MZVs, alternating multiple t -values. For example,

$$\begin{aligned} \zeta(p_1, \bar{p}_2, p_3, \bar{p}_4) &= \sum_{n_1 > n_2 > n_3 > n_4 > 0} \frac{(-1)^{n_2+n_4}}{n_1^{p_1} n_2^{p_2} n_3^{p_3} n_4^{p_4}}, \\ t(\bar{p}_1, \bar{p}_2, p_3, p_4) &= \sum_{n_1 > n_2 > n_3 > n_4 > 0} \frac{(-1)^{n_1+n_2}}{(2n_1 - 1)^{p_1} (2n_2 - 1)^{p_2} (2n_3 - 1)^{p_3} (2n_4 - 1)^{p_4}}. \end{aligned}$$

In particular, we let

$$\bar{\zeta}(p) := \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k^p} \quad \text{and} \quad \bar{t}(p) := \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{(k - 1/2)^p} \quad (p \geq 1).$$

It is clear that the multiple t -values can be regarded as a level 2 multiple zeta value because of the congruence condition in the summation and of the fact that this value can be written as a linear combination of alternating multiple zeta values. Recently, Kaneko and Tsumura [4, 5] also introduced and studied a new kind of multiple zeta values of level two

$$\begin{aligned} T(p_1, p_2, \dots, p_k) &:= 2^k \sum_{\substack{m_1 > m_2 > \dots > m_k > 0 \\ m_i \equiv k-i+1 \pmod{2}}} \frac{1}{m_1^{p_1} m_2^{p_2} \cdots m_k^{p_k}} \\ &= 2^k \sum_{n_1 > n_2 > \dots > n_k > 0} \frac{1}{(2n_1 - k)^{p_1} (2n_2 - k + 1)^{p_2} \cdots (2n_k - 1)^{p_k}}, \end{aligned}$$

which was called multiple T -values (MtVs).

The subject of this paper are alternating Euler T -sums and alternating Euler \tilde{S} -sums. First, we give the definitions of alternating harmonic number and odd harmonic number. Let p and n

be positive integers, the alternating harmonic number $\bar{H}_n^{(p)}$ and odd harmonic number $\bar{h}_n^{(p)}$ are defined by

$$\begin{aligned}\bar{H}_n^{(p)} &:= \sum_{k=1}^n \frac{(-1)^{k-1}}{k^p}, \quad \bar{H}_0^{(p)} := 0, \quad \bar{H}_n := \bar{H}_n^{(1)}, \\ \bar{h}_n^{(p)} &:= \sum_{k=1}^n \frac{(-1)^{k-1}}{(k-1/2)^p}, \quad \bar{h}_0^{(p)} := 0, \quad \bar{h}_n := \bar{h}_n^{(1)}.\end{aligned}$$

In the definitions of Euler T -sums and Euler \tilde{S} -sums, if replace “ $h_n^{(p_j)}$ ” by “ $\bar{h}_n^{(p_j)}$ ” in the numerator of the summand, we put a “bar” on the top of p_j . In particular, we put a bar on top of q if there is a sign $(-1)^{n-1}$ appearing in the denominator on the right. For example,

$$\begin{aligned}T_{p_1\bar{p}_2\bar{p}_3,\bar{q}} &= \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p_1)}\bar{h}_{n-1}^{(p_2)}\bar{h}_{n-1}^{(p_3)}}{(n-1/2)^q}(-1)^{n-1}, \quad T_{p_1\bar{p}_2\bar{p}_3p_4,q} = \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p_1)}\bar{h}_{n-1}^{(p_2)}\bar{h}_{n-1}^{(p_3)}h_{n-1}^{(p_4)}}{(n-1/2)^q}, \\ \tilde{S}_{\bar{p}_1p_2\bar{p}_3,\bar{q}} &= \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p_1)}h_{n-1}^{(p_2)}\bar{h}_{n-1}^{(p_3)}}{n^q}(-1)^{n-1}, \quad \tilde{S}_{p_1\bar{p}_2p_3\bar{p}_4,q} = \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p_1)}\bar{h}_{n-1}^{(p_2)}h_{n-1}^{(p_3)}\bar{h}_{n-1}^{(p_4)}}{(n-1/2)^q}.\end{aligned}$$

The sums of types above (one or more the p_j or q barred) are called the alternating Euler T -sums and alternating Euler \tilde{S} -sums, respectively. In [6], we systematic studied all classical (alternating) Euler sums. In this paper, we study these two above alternating variants of Euler T -sums or Euler \tilde{S} -sums by using the methods of contour integration and residue theorem.

Next, we introduce some basic notations. Let $A := \{a_k\}$ ($-\infty < k < \infty$) be a sequence of complex numbers with $a_k = o(k^\alpha)$ ($\alpha < 1$) if $k \rightarrow \pm\infty$. For convenience, we let A_1 and A_2 to denote the constant sequence $\{1^k\}$ and alternating sequence $\{(-1)^k\}$, respectively.

Definition 1.1 *With A defined above, we define the parametric digamma function $\Psi(-s; A)$ by*

$$\Psi(-s; A) := \frac{a_0}{s} + \sum_{k=1}^{\infty} \left(\frac{a_k}{k} - \frac{a_k}{k-s} \right) \quad (s \in \mathbb{C} \setminus (\mathbb{N} \cup \{0\})). \quad (1.1)$$

Obviously, if $A = A_1$, then the parametric digamma function $\Psi(-s; A)$ becomes the classical digamma function $\psi(-s) + \gamma$.

Definition 1.2 *Define the cotangent function with sequence A by*

$$\begin{aligned}\pi \cot(\pi s; A) &= -\frac{a_0}{s} + \Psi(-s; A) - \Psi(s; A) \\ &= \frac{a_0}{s} - 2s \sum_{k=1}^{\infty} \frac{a_k}{k^2 - s^2}.\end{aligned} \quad (1.2)$$

It is clear that if letting $A = A_1$ or A_2 in (1.2), respectively, then it become

$$\cot(\pi s; A_1) = \cot(\pi s) \quad \text{and} \quad \cot(\pi s; A_2) = \csc(\pi s).$$

The Definitions 1.1 and 1.2 are also introduced in a previous paper [6] of the second named author.

Definition 1.3 For nonnegative integers $j \geq 1$ and n , we define

$$\begin{aligned}
D^{(A)}(j) &:= \sum_{k=1}^{\infty} \frac{a_k}{k^j}, \quad D^{(A)}(1) := 0, \quad E_n^{(A)}(j) := \sum_{k=1}^n \frac{a_{n-k}}{k^j}, \quad E_0^{(A)}(j) := 0, \\
\bar{E}_n^{(A)}(j) &:= \sum_{k=1}^n \frac{a_{k-n-1}}{k^j}, \quad \bar{E}_0^{(A)}(j) := 0, \quad \hat{E}_n^{(A)}(j) := \sum_{k=1}^n \frac{a_{n-k}}{(k-1/2)^j}, \quad \hat{E}_0^{(A)}(j) := 0, \\
\tilde{E}_n^{(A)}(j) &:= \sum_{k=1}^n \frac{a_{k-n-1}}{(k-1/2)^j}, \quad \tilde{E}_0^{(A)}(j) := 0, \\
\tilde{t}^{(A)}(j) &:= \begin{cases} \sum_{k=1}^{\infty} \left(\frac{a_{k-1}}{k-1/2} - \frac{a_k}{k} \right), & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_{k-1}}{(k-1/2)^j}, & j > 1, \end{cases}, \quad \tilde{t}^{(A)}(j) := \begin{cases} \sum_{k=1}^{\infty} \left(\frac{a_k}{k-1/2} - \frac{a_k}{k} \right), & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_k}{(k-1/2)^j}, & j > 1, \end{cases} \\
F_n^{(A)}(j) &:= \begin{cases} \sum_{k=1}^{\infty} \frac{a_{k+n} - a_k}{k}, & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_{k+n}}{k^j}, & j > 1, \end{cases}, \quad \bar{F}_n^{(A)}(j) := \begin{cases} \sum_{k=1}^{\infty} \frac{a_{k-n} - a_k}{k}, & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_{k-n}}{k^j}, & j > 1, \end{cases}, \\
\hat{F}_n^{(A)}(j) &:= \begin{cases} \sum_{k=1}^{\infty} \left(\frac{a_{k+n}}{k-1/2} - \frac{a_k}{k} \right), & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_{k+n}}{(k-1/2)^j}, & j > 1, \end{cases}, \quad \tilde{F}_n^{(A)}(j) := \begin{cases} \sum_{k=1}^{\infty} \left(\frac{a_{k-n}}{k-1/2} - \frac{a_k}{k} \right), & j = 1, \\ \sum_{k=1}^{\infty} \frac{a_{k-n}}{(k-1/2)^j}, & j > 1, \end{cases}, \\
G_n^{(A)}(j) &:= E_n^{(A)}(j) - \bar{E}_{n-1}^{(A)}(j) - \frac{a_0}{n^j}, \quad G_0^{(A)}(j) := 0, \quad L_n^{(A)}(j) := F_n^{(A)}(j) + (-1)^j \bar{F}_n^{(A)}(j), \\
M_n^{(A)}(j) &:= E_n^{(A)}(j) + (-1)^j F_n^{(A)}(j), \quad \bar{M}_n^{(A)}(j) := \bar{F}_n^{(A)}(j) - \bar{E}_{n-1}^{(A)}(j), \quad n \geq 1, \\
R_n^{(A)}(j) &:= G_n^{(A)}(j) + (-1)^j L_n^{(A)}(j), \quad N_n^{(A)}(j) := \hat{E}_n^{(A)}(j) + (-1)^j \hat{F}_{n-1}^{(A)}(j), \\
\bar{N}_n^{(A)}(j) &:= \tilde{F}_n^{(A)}(j) - \tilde{E}_{n-1}^{(A)}(j), \quad S_n^{(A)}(j) := N_n^{(A)}(j) - \bar{N}_n^{(A)}(j) - \frac{a_0}{(n-1/2)^j}.
\end{aligned}$$

In particular, in Definition 1.3, setting $A = A_1$ or A_2 yield

$$\begin{aligned}
M_n^{(A_1)}(j) &= H_n^{(j)} + (-1)^j \zeta(j), \quad \bar{M}_n^{(A_1)}(j) = \zeta(j) - H_{n-1}^{(j)}, \quad R_n^{(A_1)}(j) = (1 + (-1)^j) \zeta(j), \\
M_n^{(A_2)}(j) &= (-1)^{n-1} \bar{H}_n^{(j)} + (-1)^j \begin{cases} (1 - (-1)^n) \log(2), & j = 1, \\ (-1)^{n-1} \bar{\zeta}(j), & j > 1, \end{cases} \\
\bar{M}_n^{(A_2)}(j) &= (-1)^n \bar{H}_{n-1}^{(j)} + \begin{cases} (1 - (-1)^n) \log(2), & j = 1, \\ (-1)^{n-1} \bar{\zeta}(j), & j > 1, \end{cases} \\
R_n^{(A_2)}(j) &= (-1)^{n-1} (1 + (-1)^j) \bar{\zeta}(j), \\
N_n^{(A_1)}(j) &= h_n^{(j)} + (-1)^j \tilde{t}(j), \quad \tilde{t}(1) := 2 \log(2), \\
N_n^{(A_2)}(j) &= (-1)^{n-1} \bar{h}_n^{(j)} + (-1)^j \begin{cases} (-1)^n \bar{t}(1) + \log(2), & j = 1, \\ (-1)^n \bar{t}(j), & j > 1, \end{cases} \\
\bar{N}_n^{(A_1)}(j) &= \tilde{t}(j) - h_{n-1}^{(j)}, \quad \tilde{t}(1) := 2 \log(2), \\
\bar{N}_n^{(A_2)}(j) &= (-1)^n \bar{h}_{n-1}^{(j)} + \begin{cases} (-1)^{n-1} \bar{t}(1) + \log(2), & j = 1, \\ (-1)^{n-1} \bar{t}(j), & j > 1, \end{cases}
\end{aligned}$$

$$S_n^{(A_1)}(j) = (1 + (-1)^j)\tilde{t}(j), \quad S_n^{(A_2)}(j) = (-1)^{n-1}(1 - (-1)^j)\bar{t}(j).$$

2 Lemmas

In this section, we give some power series expansions for parametric digamma function $\Psi(-s; A)$ and $\cot(\pi s; A)$.

Lemma 2.1 ([6, Theorem 2.2]) *Let p and n be positive integers, if $|s + n| < 1$, then*

$$\frac{\Psi^{(p-1)}(-s; A)}{(p-1)!} = (-1)^p \sum_{j=1}^{\infty} \binom{j+p-2}{p-1} M_n^{(A)}(j+p-1)(s+n)^{j-1}. \quad (2.1)$$

Lemma 2.2 ([6, Theorem 2.3]) *With $\cot(\pi s; A)$ defined above, if $|s-n| < 1$ with $s \neq n$ ($n \in \mathbb{Z}$), then*

$$\pi \cot(\pi s; A) = \frac{a_{|n|}}{s-n} - \sum_{j=1}^{\infty} (-\sigma_n)^j R_{|n|}^{(A)}(j)(s-n)^{j-1}, \quad (2.2)$$

where σ_n is defined by the symbol of n , namely,

$$\sigma_n := \begin{cases} 1, & n \geq 0 \\ -1, & n < 0. \end{cases}$$

Lemma 2.3 *Let $p > 0$ and n be a non-negative integer, if $|s-n| < 1$ then*

$$\frac{\Psi^{(p-1)}(1/2-s; A)}{(p-1)!} = \sum_{j=1}^{\infty} (-1)^{j-1} \binom{j+p-2}{p-1} N_n^{(A)}(j+p-1)(s-n)^{j-1}, \quad (2.3)$$

and if $|s+n| < 1$ then

$$\frac{\Psi^{(p-1)}(1/2-s; A)}{(p-1)!} = \sum_{j=1}^{\infty} (-1)^{j-1} \binom{j+p-2}{p-1} \bar{N}_{n+1}^{(A)}(j+p-1)(s+n)^{j-1}. \quad (2.4)$$

Proof. The proofs of this lemma follows the definition of function $\Psi(-s; A)$. \square

Letting $n = 0$ in (2.3) and (2.4) gives

$$\frac{\Psi^{(p-1)}(1/2-s; A)}{(p-1)!} = (-1)^p \sum_{j=1}^{\infty} \binom{j+p-2}{p-1} \hat{t}^{(A)}(j+p-1)s^{j-1} \quad (|s| < 1). \quad (2.5)$$

Lemma 2.4 *Let $m > 0$ and $n > 1$ be non-negative integer, if $|s-n+1/2| < 1$ then*

$$\frac{d^m}{ds^m}(\pi \cot(\pi s; A)) = (-1)^m m! \sum_{j=1}^{\infty} (-1)^{j-1} \binom{j+m-1}{m} S_n^{(A)}(j+m)(s-n+1/2)^{j-1}. \quad (2.6)$$

Proof. Lemma 2.4 follows immediately from Definition 1.2 and Lemma 2.3. \square

From Lemma 2.4, we have

$$\lim_{s \rightarrow 1/2} \frac{d^m}{ds^m}(\pi \cot(\pi s; A)) = m!((-1)^m \hat{t}^{(A)}(m+1) - \tilde{t}^{(A)}(m+1)), \quad (2.7)$$

$$\pi \cot(\pi s; A) = \frac{a_0}{s} - 2 \sum_{j=1}^{\infty} D^{(A)}(2j) s^{2j-1} \quad (|s| < 1). \quad (2.8)$$

Lemma 2.5 ([6]) Let p and n be positive integers, if $|s - n + 1/2| < 1$ and $s \neq n - 1/2$, then

$$\begin{aligned} & \frac{\Psi^{(p-1)}(1/2 - s; A)}{(p-1)!} \\ &= \frac{1}{(s - n + 1/2)^p} \left\{ a_{n-1} - \sum_{j=1}^{\infty} (-1)^j \binom{j+p-2}{p-1} M_{n-1}^{(A)}(j+p-1)(s-n+1/2)^{j+p-1} \right\}. \quad (2.9) \end{aligned}$$

Proof. This lemma can be immediately obtained from [6, Theorem 2.1]. \square

If $n = 1$ then

$$\frac{\Psi^{(p-1)}(1/2 - s; A)}{(p-1)!} = \frac{a_0}{(s - 1/2)^p} + (-1)^p \sum_{j=1}^{\infty} \binom{j+p-2}{p-1} D^{(A)}(j+p-1)(s-1/2)^{j-1}. \quad (2.10)$$

Finally, we give a residue theorem which was given by Flajolet and Salvy.

Lemma 2.6 ([1]) Let $\xi(s)$ be a kernel function and let $r(s)$ be a rational function which is $O(s^{-2})$ at infinity. Then

$$\sum_{\alpha \in O} \text{Res}[r(s)\xi(s), s = \alpha] + \sum_{\beta \in S} \text{Res}[r(s)\xi(s), s = \beta] = 0. \quad (2.11)$$

where S is the set of poles of $r(s)$ and O is the set of poles of $\xi(s)$ that are not poles $r(s)$. Here $\text{Res}[r(s), s = \alpha]$ denotes the residue of $r(s)$ at $s = \alpha$. The kernel function $\xi(s)$ is defined by the two requirements: 1. $\xi(s)$ is meromorphic in the whole complex plane. 2. $\xi(s)$ satisfies $\xi(s) = o(s)$ over an infinite collection of circles $|s| = \rho_k$ with $\rho_k \rightarrow \infty$.

3 Evaluations of Euler T -sums and Euler \tilde{S} -sums

Let $B := \{b_k\}$, $-\infty < k < \infty$ be a sequence of complex numbers with $b_k = o(k^\beta)$ ($\beta < 1$) if $k \rightarrow \pm\infty$. Flajolet and Salvy [1] applied the kernel function

$$\frac{1}{2}\pi \cot(\pi s) \frac{\psi^{(p-1)}(-s)}{(p-1)!}$$

to the base function $r(s) = s^{-q}$ to prove every linear sum $S_{p,q}$ whose weight $p+q$ is odd is expressible as a polynomial in zeta values. Next, we replace $\cot(\pi s)\psi^{(p-1)}(-s)$ by $\cot(\pi s; A)\Psi^{(p-1)}(-s; B)$, and use contour integration to evaluate linear (alternating) Euler T -sums and Euler \tilde{S} -sums.

Theorem 3.1 Let $p > 0$ and $q > 1$ be positive integers. We have

$$\begin{aligned} & (-1)^{p+q} \sum_{n=1}^{\infty} \frac{\bar{N}_n^{(B)}(p)}{(n - 1/2)^q} a_{n-1} + \sum_{n=1}^{\infty} \frac{N_n^{(B)}(p)}{(n - 1/2)^q} a_n \\ & - (-1)^p \sum_{k=0}^{p-1} \binom{p+q-k-2}{q-1} \sum_{n=1}^{\infty} \frac{b_n S_{n+1}^{(A)}(k+1)}{n^{p+q-k-1}} \\ & - b_0 \left((-1)^{p+q} \tilde{t}^{(A)}(p+q) + \tilde{t}^{(A)}(p+q) \right) \\ & + (-1)^p \sum_{k=1}^q \binom{k+p-2}{p-1} D^{(B)}(k+p-1) \left((-1)^{q-k} \tilde{t}^{(A)}(q-k+1) - \tilde{t}^{(A)}(q-k+1) \right) \\ & = 0. \quad (3.1) \end{aligned}$$

Proof. We consider the kernel function

$$\pi \cot(\pi s; A) \frac{\Psi^{(p-1)}(1/2 - s; B)}{(p-1)!}$$

and base function $r(s) = (s - 1/2)^{-q}$. Clearly, the function $F(s) := \xi(s)r(s)$ only have poles at all integer and $n - 1/2$ (n is a positive integer). The only singularities are poles at the integers. At a negative integer $-n$ and positive integer n these two poles are simple and these residues are

$$\begin{aligned} \text{Res}[F(s), s = -n] &= (-1)^{p+q} \frac{\bar{N}_{n+1}^{(B)}(p)}{(n + 1/2)^q} a_n \quad (n \geq 0), \\ \text{Res}[F(s), s = n] &= \frac{N_n^{(B)}(p)}{(n - 1/2)^q} a_n \quad (n \geq 1), \end{aligned}$$

where we used the identities (2.2)-(2.4). From (2.9), the pole $n - 1/2$ ($n \geq 2$) has order p and the residue is

$$\text{Res}[F(s), s = n - 1/2] = -(-1)^p \sum_{k=0}^{p-1} \binom{p+q-k-2}{q-1} \frac{b_{n-1} S_n^{(A)}(k+1)}{(n-1)^{p+q-k-1}}.$$

From (2.10), the pole $1/2$ has order $p+q$ and the residue is

$$\begin{aligned} \text{Res}[F(s), s = 1/2] &= -b_0 \left((-1)^{p+q} \hat{t}^{(A)}(p+q) + \tilde{t}^{(A)}(p+q) \right) \\ &\quad + (-1)^p \sum_{k=1}^q \binom{k+p-2}{p-1} D^{(B)}(k+p-1) \left((-1)^{q-k} \hat{t}^{(A)}(q-k+1) - \tilde{t}^{(A)}(q-k+1) \right). \end{aligned}$$

Summing these four contributions yields the statement of the theorem. \square

Theorem 3.2 *Let $p > 0$ and $q > 1$ be positive integers. We have*

$$\begin{aligned} &(-1)^{p+q} \sum_{n=1}^{\infty} \frac{\bar{N}_{n+1}^{(B)}(p)}{n^q} a_n + \sum_{n=1}^{\infty} \frac{N_n^{(B)}(p)}{n^q} a_n \\ &- (-1)^p \sum_{k=0}^{p-1} \binom{p+q-k-2}{q-1} \sum_{n=1}^{\infty} \frac{b_{n-1} S_n^{(A)}(k+1)}{(n-1/2)^{p+q-k-1}} \\ &+ a_0 (-1)^p \binom{p+q-1}{q} \hat{t}^{(B)}(p+q) \\ &- 2(-1)^p \sum_{j=1}^{[q/2]} \binom{p+q-2j-1}{p-1} D^{(A)}(2j) \hat{t}^{(B)}(p+q-2j) \\ &= 0. \end{aligned} \tag{3.2}$$

Proof. The proof is similar to the previous proof. We consider the kernel function

$$\frac{1}{2} \pi \cot(\pi s; A) \frac{\Psi^{(p-1)}(1/2 - s; B)}{(p-1)!}$$

and base function $r(s) = s^{-q}$. Then, by a similar argument as in the proof of above, we may easily deduce the desired result. \square

In Theorems 3.1 and 3.2, setting $A, B \in \{A_1, A_2\}$, by straightforward calculations, we can get the following corollaries.

Corollary 3.3 *For positive integers p and $q > 1$,*

$$\begin{aligned} & (1 - (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p)}}{(n - 1/2)^q} \\ &= (-1)^{p+q} \tilde{t}(p+q) - (-1)^p (1 + (-1)^q) \tilde{t}(p) \tilde{t}(q) \\ &\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k - 1) \binom{p+q-k-2}{q-1} \tilde{t}(k+1) \zeta(p+q-k-1) \\ &\quad + (-1)^p \sum_{k=1}^q (1 - (-1)^{q-k}) \binom{k+p-2}{p-1} \tilde{t}(q-k+1) \zeta(k+p-1), \end{aligned} \tag{3.3}$$

$$\begin{aligned} & (1 + (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{\bar{h}_{n-1}^{(p)}}{(n - 1/2)^q} \\ &= -(-1)^{p+q} \bar{t}(p+q) + (-1)^p (1 + (-1)^q) \bar{t}(p) \tilde{t}(q) \\ &\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k + 1) \binom{p+q-k-2}{q-1} \bar{t}(k+1) \zeta(p+q-k-1) \\ &\quad - (-1)^p \sum_{k=1}^q (1 + (-1)^{q-k}) \binom{k+p-2}{p-1} \bar{t}(q-k+1) \bar{\zeta}(k+p-1), \end{aligned} \tag{3.4}$$

$$\begin{aligned} & (1 - (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{\bar{h}_{n-1}^{(p)}}{(n - 1/2)^q} (-1)^{n-1} \\ &= (-1)^{p+q} \tilde{t}(p+q) + (-1)^p (1 - (-1)^q) \bar{t}(p) \bar{t}(q) \\ &\quad + (-1)^p \sum_{k=0}^{p-1} ((-1)^k - 1) \binom{p+q-k-2}{q-1} \tilde{t}(k+1) \bar{\zeta}(p+q-k-1) \\ &\quad - (-1)^p \sum_{k=1}^q (1 - (-1)^{q-k}) \binom{k+p-2}{p-1} \tilde{t}(q-k+1) \bar{\zeta}(k+p-1), \end{aligned} \tag{3.5}$$

$$\begin{aligned} & (1 + (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{h_{n-1}^{(p)}}{(n - 1/2)^q} (-1)^{n-1} \\ &= -(-1)^{p+q} \bar{t}(p+q) - (-1)^p (1 - (-1)^q) \tilde{t}(p) \bar{t}(q) \\ &\quad + (-1)^p \sum_{k=0}^{p-1} ((-1)^k + 1) \binom{p+q-k-2}{q-1} \bar{t}(k+1) \zeta(p+q-k-1) \\ &\quad + (-1)^p \sum_{k=1}^q (1 + (-1)^{q-k}) \binom{k+p-2}{p-1} \bar{t}(q-k+1) \zeta(k+p-1), \end{aligned} \tag{3.6}$$

where $\zeta(1) := 0$ and $\tilde{t}(1) := 2 \log(2)$.

Corollary 3.4 For positive integers p and $q > 1$,

$$\begin{aligned}
& (1 - (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{h_n^{(p)}}{n^q} \\
&= -(-1)^p (1 + (-1)^q) \tilde{t}(p) \zeta(q) - (-1)^p \binom{p+q-1}{p-1} \tilde{t}(p+q) \\
&\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k - 1) \binom{p+q-k-2}{q-1} \tilde{t}(k+1) \tilde{t}(p+q-k-1) \\
&\quad + 2(-1)^p \sum_{j=1}^{[q/2]} \binom{p+q-2j-1}{p-1} \zeta(2j) \tilde{t}(p+q-2j), \\
& (1 + (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{\bar{h}_n^{(p)}}{n^q}
\end{aligned} \tag{3.7}$$

$$\begin{aligned}
&= (-1)^p (1 + (-1)^q) \bar{t}(p) \zeta(q) + (-1)^p \binom{p+q-1}{p-1} \bar{t}(p+q) \\
&\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k + 1) \binom{p+q-k-2}{q-1} \bar{t}(k+1) \tilde{t}(p+q-k-1) \\
&\quad + 2(-1)^p \sum_{j=1}^{[q/2]} \binom{p+q-2j-1}{p-1} \bar{\zeta}(2j) \bar{t}(p+q-2j),
\end{aligned} \tag{3.8}$$

$$\begin{aligned}
& (1 + (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{\bar{h}_n^{(p)}}{n^q} (-1)^{n-1} \\
&= (-1)^p (1 + (-1)^q) \bar{t}(p) \bar{\zeta}(q) - (-1)^p \binom{p+q-1}{p-1} \bar{t}(p+q) \\
&\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k - 1) \binom{p+q-k-2}{q-1} \tilde{t}(k+1) \bar{t}(p+q-k-1) \\
&\quad + 2(-1)^p \sum_{j=1}^{[q/2]} \binom{p+q-2j-1}{p-1} \zeta(2j) \bar{t}(p+q-2j),
\end{aligned} \tag{3.9}$$

$$\begin{aligned}
& (1 - (-1)^{p+q}) \sum_{n=1}^{\infty} \frac{h_n^{(p)}}{n^q} (-1)^{n-1} \\
&= -(-1)^p (1 + (-1)^q) \tilde{t}(p) \bar{\zeta}(q) + (-1)^p \binom{p+q-1}{p-1} \tilde{t}(p+q) \\
&\quad - (-1)^p \sum_{k=0}^{p-1} ((-1)^k + 1) \binom{p+q-k-2}{q-1} \bar{t}(k+1) \bar{t}(p+q-k-1) \\
&\quad + 2(-1)^p \sum_{j=1}^{[q/2]} \binom{p+q-2j-1}{p-1} \bar{\zeta}(2j) \tilde{t}(p+q-2j),
\end{aligned} \tag{3.10}$$

where $\zeta(1) := 0$ and $\tilde{t}(1) := 2 \log(2)$.

Remark 3.5 Note that when $q = 1$, the four identities (3.5), (3.6), (3.9) and (3.10) are also hold. We leave the detail to the interested reader.

Next, we evaluate the quadratic (alternating) Euler T -sums and Euler \tilde{S} -sums.

Theorem 3.6 For positive integers p, m and $q > 1$, then

$$\begin{aligned}
& (-1)^{p+q+m} \sum_{n=1}^{\infty} \frac{\bar{N}_n^{(B)}(m)\bar{N}_n^{(C)}(p)}{(n-1/2)^q} a_{n-1} + \sum_{n=1}^{\infty} \frac{N_n^{(B)}(m)N_n^{(C)}(p)}{(n-1/2)^q} a_n \\
& - (-1)^{p+m} \sum_{k=0}^{p+m-1} \binom{p+q+m-k-2}{q-1} \sum_{n=1}^{\infty} \frac{b_n c_n S_{n+1}^{(A)}(k+1)}{n^{p+q+m-k-1}} \\
& - (-1)^m \sum_{j=1}^m \sum_{k=0}^{m-j} \binom{j+p-2}{p-1} \binom{m+q-k-j-1}{q-1} \sum_{n=1}^{\infty} \frac{M_n^{(C)}(j+p-1)S_{n+1}^{(A)}(k+1)b_n}{n^{m+q-j-k}} \\
& - (-1)^p \sum_{j=1}^p \sum_{k=0}^{p-j} \binom{j+m-2}{m-1} \binom{p+q-k-j-1}{q-1} \sum_{n=1}^{\infty} \frac{M_n^{(B)}(j+m-1)S_{n+1}^{(A)}(k+1)c_n}{n^{p+q-j-k}} \\
& + \text{Res}[F(s), s = 1/2] = 0,
\end{aligned} \tag{3.11}$$

where

$$\begin{aligned}
& \text{Res}[F(s), s = 1/2] \\
& = -b_0 c_0 \left((-1)^{p+q+m} \tilde{t}^{(A)}(p+q+m) + \tilde{t}^{(A)}(p+q+m) \right) \\
& + b_0 (-1)^p \sum_{j=1}^{m+q} \binom{j+p-2}{j-1} D^{(C)}(j+p-1) \\
& \quad \times \left((-1)^{m+q-j} \tilde{t}^{(A)}(m+q-j+1) - \tilde{t}^{(A)}(m+q-j+1) \right) \\
& + c_0 (-1)^m \sum_{j=1}^{p+q} \binom{j+m-2}{j-1} D^{(B)}(j+m-1) \\
& \quad \times \left((-1)^{p+q-j} \tilde{t}^{(A)}(p+q-j+1) - \tilde{t}^{(A)}(p+q-j+1) \right) \\
& + (-1)^{p+m} \sum_{\substack{j_1+j_2 \leq q+1, \\ j_1, j_2 \geq 1}} \binom{j_1+m-2}{j_1-1} \binom{j_2+p-2}{j_2-1} D^{(B)}(j_1+m-1) D^{(C)}(j_2+p-1) \\
& \quad \times \left((-1)^{q+1-j_1-j_2} \tilde{t}^{(A)}(q+2-j_1-j_2) - \tilde{t}^{(A)}(q+2-j_1-j_2) \right).
\end{aligned} \tag{3.12}$$

Proof. We consider the kernel function

$$\cot(\pi s; A) \frac{\Psi^{(m-1)}(1/2-s; B)\Psi^{(p-1)}(1/2-s; C)}{(m-1)!(p-1)!}$$

and base function $r(s) = (s-1/2)^{-q}$. It is obvious that the function

$$F(s) := \cot(\pi s; A) \frac{\Psi^{(m-1)}(1/2-s; B)\Psi^{(p-1)}(1/2-s; C)}{(m-1)!(p-1)!(s-1/2)^q}$$

has simple poles at $s = -n$ ($n \geq 0$) with residues

$$\text{Res}[F(s), s = -n] = (-1)^{p+q+m} \frac{\bar{N}_{n+1}^{(B)}(m) \bar{N}_{n+1}^{(C)}(p)}{(n+1/2)^q} a_n,$$

and simple poles at $s = n$ ($n \geq 1$), with residues

$$\text{Res}[F(s), s = n] = \frac{N_n^{(B)}(m) N_n^{(C)}(p)}{(n-1/2)^q} a_n,$$

where we used the identities (2.3) and (2.4). Clearly, $F(s)$ has poles of order $p+m$ at $s = n-1/2$ ($n \geq 2$). Using (2.6) and (2.9) we find that the residues

$$\begin{aligned} & \text{Res}[F(s), s = n-1/2] \\ &= -(-1)^{p+m} \sum_{k=0}^{p+m-1} \binom{p+q+m-k-2}{q-1} \frac{b_{n-1} c_{n-1} S_n^{(A)}(k+1)}{(n-1)^{p+q+m-k-1}} \\ &\quad - (-1)^m \sum_{j=1}^m \sum_{k=0}^{m-j} \binom{j+p-2}{p-1} \binom{m+q-k-j-1}{q-1} \frac{M_{n-1}^{(C)}(j+p-1) S_n^{(A)}(k+1) b_{n-1}}{(n-1)^{m+q-j-k}} \\ &\quad - (-1)^p \sum_{j=1}^p \sum_{k=0}^{p-j} \binom{j+m-2}{m-1} \binom{p+q-k-j-1}{q-1} \frac{M_{n-1}^{(B)}(j+m-1) S_n^{(A)}(k+1) c_{n-1}}{(n-1)^{p+q-j-k}}. \end{aligned}$$

Moreover, $F(s)$ also has a pole of order $p+q+m$ at $s = 1/2$. Using (2.10) we deduce the formula (3.12) by a direct calculation. Hence, combining these four residue results, we can obtain the desired evaluation. \square

Theorem 3.7 *For positive integers p, m and $q > 1$, then*

$$\begin{aligned} & (-1)^{p+q+m} \sum_{n=1}^{\infty} \frac{\bar{N}_{n+1}^{(B)}(m) \bar{N}_{n+1}^{(C)}(p)}{n^q} a_n + \sum_{n=1}^{\infty} \frac{N_n^{(B)}(m) N_n^{(C)}(p)}{n^q} a_n \\ & - (-1)^{p+m} \sum_{k=0}^{p+m-1} \binom{p+q+m-k-2}{q-1} \sum_{n=1}^{\infty} \frac{b_{n-1} c_{n-1} S_n^{(A)}(k+1)}{(n-1/2)^{p+q+m-k-1}} \\ & - (-1)^m \sum_{j=1}^m \sum_{k=0}^{m-j} \binom{j+p-2}{p-1} \binom{m+q-k-j-1}{q-1} \sum_{n=1}^{\infty} \frac{M_{n-1}^{(C)}(j+p-1) S_n^{(A)}(k+1) b_{n-1}}{(n-1/2)^{m+q-j-k}} \\ & - (-1)^p \sum_{j=1}^p \sum_{k=0}^{p-j} \binom{j+m-2}{m-1} \binom{p+q-k-j-1}{q-1} \sum_{n=1}^{\infty} \frac{M_{n-1}^{(B)}(j+m-1) S_n^{(A)}(k+1) c_{n-1}}{(n-1/2)^{p+q-j-k}} \\ & + \text{Res}[G(s), s = 0] = 0, \end{aligned} \tag{3.13}$$

where

$$\begin{aligned} & \text{Res}[G(s), s = 0] \\ &= a_0 (-1)^{p+m} \sum_{\substack{k_1+k_2=q, \\ k_1, k_2 \geq 0}} \binom{m+k_1-1}{k_1} \binom{p+k_2-1}{k_2} \tilde{t}^{(B)}(m+k_1) \tilde{t}^{(C)}(p+k_2) \end{aligned}$$

$$- 2(-1)^{m+p} \sum_{j=1}^{[q/2]} \sum_{\substack{k_1+k_2=q-2j, \\ k_1, k_2 \geq 0}} \binom{m+k_1-1}{k_1} \binom{p+k_2-1}{k_2} D^{(A)}(2j) \tilde{t}^{(B)}(m+k_1) \tilde{t}^{(C)}(p+k_2). \quad (3.14)$$

Proof. We consider the kernel function

$$\pi \cot(\pi s; A) \frac{\Psi^{(m-1)}(1/2-s; B) \Psi^{(p-1)}(1/2-s; C)}{(m-1)!(p-1)!}$$

and base function $r(s) = s^{-q}$. By the same calculation as in the proof of Theorem 3.6, we thus immediately deduce (3.13) and (3.14) to complete the proof. \square

Theorem 3.8 For positive integers p, m and $q > 1$, then

$$\begin{aligned} & (-1)^{p+q+m} \sum_{n=1}^{\infty} \frac{\bar{M}_n^{(B)}(m) \bar{N}_{n+1}^{(C)}(p)}{n^q} a_n + \sum_{n=1}^{\infty} \frac{M_n^{(B)}(m) N_n^{(C)}(p)}{n^q} a_n \\ & + (-1)^m \sum_{k=0}^m \binom{m+q-k-1}{q-1} \binom{p+k-1}{p-1} \sum_{n=1}^{\infty} \frac{N_n^{(C)}(p+k)}{n^{m+q-k}} a_n b_n \\ & - (-1)^m \sum_{\substack{k+j \leq m+1, \\ k, j \geq 1}} \binom{m+q-k-j}{q-1} \binom{p+k-2}{p-1} \sum_{n=1}^{\infty} \frac{R_n^{(A)}(j) N_n^{(C)}(p+k-1)}{n^{m+q+1-k-j}} b_n \\ & - (-1)^p \sum_{\substack{k_1+k_2+k_3=p-1, \\ k_1, k_2, k_3 \geq 0}} \binom{m+k_2-1}{m-1} \binom{q+k_3-1}{q-1} \sum_{n=1}^{\infty} \frac{S_n^{(A)}(k_1+1) N_n^{(B)}(m+k_2)}{(n-1/2)^{k_3+q}} c_{n-1} \\ & + \text{Res}[H(s), s=0] = 0, \end{aligned} \quad (3.15)$$

where

$$\begin{aligned} & \text{Res}[H(s), s=0] \\ & = a_0 b_0 (-1)^p \binom{p+q+m-1}{p-1} \tilde{t}^{(C)}(p+q+m) \\ & + a_0 (-1)^{p+m} \sum_{j=1}^{q+1} \binom{j+m-2}{m-1} \binom{p+q-j}{p-1} D^{(B)}(j+m-1) \tilde{t}^{(C)}(p+q+1-j) \\ & - 2b_0 (-1)^p \sum_{j=1}^{[(m+q)/2]} \binom{p+q+m-2j-1}{p-1} D^{(A)}(2j) \tilde{t}^{(C)}(p+q+m-2j) \\ & - 2(-1)^{p+m} \sum_{\substack{2j_1+j_2 \leq q+1, \\ j_1, j_2 \geq 1}} \binom{j_2+m-2}{m-1} \binom{p+q-2j_1-j_2}{p-1} \\ & \quad \times D^{(A)}(2j_1) D^{(B)}(j_2+m-1) \tilde{t}^{(C)}(p+q+1-2j_1-j_2). \end{aligned} \quad (3.16)$$

Proof. We consider the kernel function

$$\pi \cot(\pi s; A) \frac{\Psi^{(m-1)}(-s; B) \Psi^{(p-1)}(1/2-s; C)}{(m-1)!(p-1)!}$$

and base function $r(s) = s^{-q}$. By direct residue computations, we can obtain the desired evaluation with the help of formulas (2.1)-(2.6). \square

It is clear that the main results in our previous paper [7] are immediate corollaries of this paper. Moreover, it is possible that of some other relations involving alternating Euler T -sums and related sums can be proved by using the techniques of the present paper. For example, let $A^{(l)} := \{a_k^{(l)}\}$ ($-\infty < k < \infty$, l is any positive integer) be any sequences of complex numbers with $a_k^{(l)} = o(k^\alpha)$ ($\alpha < 1$) if $k \rightarrow \pm\infty$, consider these two function

$$\cot(\pi s; A) \frac{\Psi^{(p_1-1)}(1/2-s; A^{(1)})\Psi^{(p_2-1)}(1/2-s; A^{(2)}) \dots \Psi^{(p_r-1)}(1/2-s; A^{(r)})}{(m-1)!(p-1!)(s-1/2)^q}$$

and

$$\cot(\pi s; A) \frac{\Psi^{(p_1-1)}(1/2-s; A^{(1)})\Psi^{(p_2-1)}(1/2-s; A^{(2)}) \dots \Psi^{(p_r-1)}(1/2-s; A^{(r)})}{(m-1)!(p-1!)s^q}$$

we can deduce the following results

$$\begin{aligned} & (-1)^{p_1+\dots+p_r+q} \sum_{n=1}^{\infty} \frac{\bar{N}_n^{(A^{(1)})}(p_1)\bar{N}_n^{(A^{(2)})}(p_2) \dots \bar{N}_n^{(A^{(r)})}(p_r)}{(n-1/2)^q} a_{n-1} \\ & + \sum_{n=1}^{\infty} \frac{N_n^{(A^{(1)})}(p_1)N_n^{(A^{(2)})}(p_2) \dots N_n^{(A^{(r)})}(p_r)}{(n-1/2)^q} a_n \\ & + \sum (\text{sums of degree } \leq r-1) = 0 \end{aligned}$$

and

$$\begin{aligned} & (-1)^{p_1+\dots+p_r+q} \sum_{n=1}^{\infty} \frac{\bar{N}_{n+1}^{(A^{(1)})}(p_1)\bar{N}_{n+1}^{(A^{(2)})}(p_2) \dots \bar{N}_{n+1}^{(A^{(r)})}(p_r)}{n^q} a_n \\ & + \sum_{n=1}^{\infty} \frac{N_n^{(A^{(1)})}(p_1)N_n^{(A^{(2)})}(p_2) \dots N_n^{(A^{(r)})}(p_r)}{n^q} a_n \\ & + \sum (\text{sums of degree } \leq r-1) = 0, \end{aligned}$$

but we can't get the general explicit formulas.

Acknowledgments. The authors express their deep gratitude to Professors Masanobu Kaneko and Jianqiang Zhao for valuable discussions and comments.

References

- [1] P. Flajolet and B. Salvy, Euler sums and contour integral representations, *Exp. Math.*, 1998, **7**(1): 15–35.
- [2] M.E. Hoffman, Multiple harmonic series, *Pacific J. Math.*, 1992, **152**: 275–290.
- [3] M.E. Hoffman, An odd variant of multiple zeta values, *Comm. Number Theory Phys.*, 2019, **13**, 529–567.

- [4] M. Kaneko, H. Tsumura, Zeta functions connecting multiple zeta values and poly-Bernoulli numbers, arXiv: 1811.07736v1.
- [5] M. Kaneko, H. Tsumura, On a variant of multiple zeta values of level two, arXiv: 1903.03747v2.
- [6] C. Xu, Explicit Formulas for general Euler type sums. arXiv:2002.12107v3.
- [7] C. Xu, W. Wang, Two variants of Euler sums, arXiv:1906.07654v3.
- [8] D. Zagier, Values of zeta functions and their applications, First European Congress of Mathematics, Volume II, Birkhauser, Boston, **120**(1994) 497-512.