On Dissipative Symplectic Integration with Applications to Gradient-Based Optimization

Guilherme França, ${}^{\omega\bar{\omega}*}$ Michael I. Jordan, ${}^{\omega\hat{\omega}\dagger}$ and René Vidal ${}^{\bar{\omega}\ddagger}$

^wDepartment of Electrical Engineering and Computer Sciences, University of California, Berkeley, CA 94720, USA

> ^ŵDepartment of Statistics, University of California, Berkeley, CA 94720, USA

^{$\bar{\omega}$} Mathematical Institute for Data Science, Johns Hopkins University, Baltimore, MD 21218, USA

Abstract

Recently, continuous dynamical systems have proved useful in providing conceptual and quantitative insights into gradient-based optimization, widely used in modern machine learning and statistics. An important question that arises in this line of work is how to discretize the system in such a way that its stability and rates of convergence are preserved. In this paper we propose a geometric framework in which such discretizations can be realized systematically, enabling the derivation of "rate-matching" optimization algorithms without the need for a discrete convergence analysis. More specifically, we show that a generalization of symplectic integrators to dissipative Hamiltonian systems is able to preserve continuous rates of convergence up to a controlled error. Moreover, such methods preserve a perturbed Hamiltonian despite the absence of a conservation law, extending key results of symplectic integrators to dissipative cases. Our arguments rely on a combination of backward error analysis with fundamental results from symplectic geometry.

^{*}guifranca@gmail.com

[†]jordan@cs.berkeley.edu

[‡]rvidal@jhu.edu

1 Introduction

A recent line of research at the interface of machine learning and optimization focuses on establishing connections between continuous dynamical systems and gradient-based optimization methods [1–13]. From this perspective, an optimization algorithm is a particular discretization of a differential equation. Moreover, important *accelerated methods* such as Nesterov's method [14] and Polyak's heavy ball method [15] are modeled as second-order differential equations with a dissipative term [1–6]. There are advantages to working in the continuous-time formalism. In particular, the stability and convergence analysis of a continuous system tends to be simpler and more transparent, making use of general tools such as Lyapunov stability theory and variational formulations. The traditional discrete analysis is usually only applicable on a case-by-case basis and often requires painstaking algebra. This is an unsatisfactory state of affairs given that accelerated optimization methods are the workhorses behind many of the empirical success stories in large-scale machine learning.

A particularly useful step has been the development of a Lagrangian and Hamiltonian perspective on acceleration methods [2]. This framework, which involves the definition of a *Bregman Lagrangian* and a *Bregman Hamiltonian*, places momentum-based methods such as Nesterov acceleration into a larger class of dynamical systems and has accordingly helped to demystify the notion of "acceleration" in optimization. Two difficulties arise, however, when one attempts to further exploit and characterize this class of systems. First, there are many different ways to discretize a continuous system. As pointed out [2], a naive discretization may be unable to preserve the continuous rates of convergence—i.e., the rates of decay to lower-energy level sets—and may even lead to an unstable algorithm. Moreover, it is largely unknown if there is an underlying principle from which one can construct such "rate-matching" discretizations. Thus, a fundamental question arises:

Which classes of discretizations are capable of preserving the rates of convergence of the continuous dynamical systems of interest in optimization?

A second difficulty is that the Hamiltonian formalism has traditionally been applied to conservative systems; in particular, systems characterized by oscillations. Such behavior is incommensurate with the desire to converge to an optimum—a system that converges towards a limit cycle may have favorable stability properties, but the presence of limit cycles may preclude convergence to a point. Thus we have a second fundamental question:

Can we map discrete-time algorithms into dissipative continuous-time dynamical systems that provide analytical insight into the behavior of the original algorithm?

Clearly these two questions are related. Indeed, the ability to map between dynamical systems while preserving rates may be seen as a form of "invariance"—though different from the conservation laws arising from an underlying symmetry in Hamiltonian systems.

In this paper we attempt to provide answers to the above questions. Introducing a class of *dissipative Hamiltonian systems*, we combine fundamental results from symplectic geometry [16, 17] and backward error analysis [18–20] to establish a general quantitative guarantee for the convergence of discrete algorithms based on these continuous dynamics. More specifically, we propose a class of discretizations, which we refer to as *presymplectic integrators*, that are designed to preserve a fundamental geometric structure associated with dissipative Hamiltonian systems.¹

Presymplectic integrators consist of a generalization of the well-known family of symplectic integrators [21–25], which have been developed in the setting of *conservative* Hamiltonian systems. The most important property of symplectic integrators is that in addition to preserving the symplectic structure of conservative Hamiltonian systems they exactly conserve a shadow or perturbed Hamiltonian [18], thus ensuring long-term stability. This crucial result relies on the fact that the Hamiltonian is a constant of motion. On the other hand, there are relatively few results on structure-preserving methods for dissipative systems, though this has been the subject of a nascent literature [26–31]. It is thus unknown if the key stability properties of symplectic integrators can be extended to dissipative cases precisely because a conserved quantity is in general no longer available. We will show that presymplectic integrators allow such properties to be extended into a dissipative setting. In particular we show that they preserve a *time-varying* shadow Hamiltonian and accordingly exhibit long-term stability, despite the absence of a conservation law. Our argument relies on a *symplectification* procedure where the dissipative system is embedded in the phase space of a higher-dimensional conservative system.

Although this paper focuses on bringing a dynamical perspective to accelerated optimization, we note in passing that our results may also be of interest in other fields where the simulation of dissipative systems is important, such as out-of-equilibrium statistical mechanics, complex systems and economics.

The paper is organized as follows. In Appendix A we introduce notation and recall the basic concepts from differential geometry which are needed throughout the paper. In Section 2 we overview, at a high level, the main outlines of our analysis. Section 3 introduces ideas from backward error analysis and symplectic integrators, presenting geometric proofs of earlier results [18–20] so as to anticipate our generalizations to dissipative systems. In Section 4, we introduce dissipative Hamiltonian systems from the point of view of symplectic geometry and construct their symplectification. We then define presymplectic integrators and argue that they extend the useful properties of symplectic integrators into a dissipative setting. In Section 5, we consider the implications of this framework for solving optimization problems. In Section 6, we construct explicit presymplectic integrators for the Bregman Hamiltonian in full generality. Section 7 contains our final remarks.

¹This will be made precise later but briefly the idea is that the phase space of a dissipative Hamiltonian system is a *presymplectic manifold*, endowed with a closed *degenerate* symplectic 2-form, which is exactly preserved by a presymplectic integrator.

2 Overview of the Implications to Accelerated Optimization

Given an *n*-dimensional smooth manifold \mathcal{M} and a function $f : \mathcal{M} \to \mathbb{R}$, consider the optimization problem

$$f(q^{\star}) = \min_{q \in \mathcal{M}} f(q). \tag{2.1}$$

Let H = H(t, q, p) be an explicitly *time-dependent* Hamiltonian over the phase space $(q, p) \in T^*\mathcal{M}$ —the cotangent bundle of \mathcal{M} (see Appendix A)—which determines dynamical evolution through Hamilton's equations:

$$\frac{dq^{j}}{dt} = \frac{\partial H}{\partial p_{j}}, \qquad \frac{dp_{j}}{dt} = -\frac{\partial H}{\partial q^{j}} \qquad (j = 1, \dots, n).$$
(2.2)

We will design dissipative systems whose trajectories tend to a low-energy level set that is consistent with a minimum of f^2 . Specifically, we consider systems arising from the following general family of Hamiltonians:

$$H \equiv e^{-\eta_1(t)} T(t, q, p) + e^{\eta_2(t)} f(q), \qquad (2.3)$$

where η_1 and η_2 are positive and monotone increasing functions that are responsible for introducing dissipation. The kinetic energy T is assumed to be Lipschitz continuous. This Hamiltonian includes many dissipative systems that are relevant to optimization, including the Bregman Hamiltonian [2] and conformal Hamiltonian systems [32]—see Appendix B for a generalization thereof. Note also that (2.3) generalizes the Caldirola-Kanai Hamiltonian [33, 34] which can be seen as the classical limit of the seminal Caldeira-Leggett model [35], important in quantum dissipation and decoherence.

It is possible to characterize the convergence rate that a dissipative system tends to a minimum through a Lyapunov analysis [1-3,7]. This leads to upper bounds:

$$f(q(t)) - f(q^*) = \mathcal{O}\left(\mathcal{R}(t)\right), \qquad (2.4)$$

for a decreasing function of time, $\mathcal{R}(t)$, depending on the landscape of f. Our goal is to construct "rate-matching discretizations," namely general numerical integrators able to reproduce (2.4). As previously mentioned, we propose a class of discretizations called presymplectic integrators to this end—this is formalized in Definition 4.1.

Let $x \equiv (q, p)$. A numerical integrator for the Hamiltonian system (2.2) is a map $\phi_h : \mathbb{R}^{2n} \to \mathbb{R}^{2n}$, with step size h > 0, such that iterations

$$x_{\ell} = \phi_h(x_{\ell-1}), \qquad x_0 = x(0),$$
(2.5)

²By way of contrast, a classical conservative system has a *time-independent* Hamiltonian, H = H(q, p), such that $\frac{dH}{dt} = 0$, implying that trajectories oscillate around a minimum instead of converging.

approximate the true state $x(t_{\ell}) \equiv (q(t_{\ell}), p(t_{\ell}))$ at instants $t_{\ell} = h\ell$ ($\ell = 1, 2, ...$). Let φ_t denote the true flow of (2.2). An integrator ϕ_h is said to be of order $r \geq 1$ if

$$\|\phi_h(x) - \varphi_h(x)\| = \mathcal{O}(h^{r+1}), \qquad (2.6)$$

for any $x \in T^*\mathcal{M}$. We also introduce a Lipschitz assumption for the integrator:

$$\|\phi_h(y) - \phi_h(x)\| \le (1 + hL_\phi) \|y - x\|, \tag{2.7}$$

for some constant $L_{\phi} > 0$, and for all $x, y \in T^* \mathcal{M}^3$. We now state our main result, which we will further explicate and establish formally in the remainder of the paper.

Theorem 2.1. Consider a dissipative Hamiltonian system (2.2) obtained from (2.3). Let ϕ_h be a presymplectic integrator of order r, assumed to obey the Lipschitz condition (2.7). Then ϕ_h preserves the continuous rates of convergence up to a small error, namely

$$\underbrace{f(q_{\ell}) - f(q^{\star})}_{\text{discrete rate}} = \underbrace{f(q(t_{\ell})) - f(q^{\star})}_{\text{continuous rate}} + \underbrace{\mathcal{O}\left(h^r e^{-\eta_2(t_{\ell})}\right)}_{\text{small error}},$$
(2.8)

provided $e^{L_{\phi}t_{\ell}-\eta_1(t_{\ell})} < \infty$ for sufficiently large t_{ℓ} . This holds for exponentially large times $t_{\ell} \equiv h\ell = \mathcal{O}(h^r e^r e^{h_0/h})$, where the constant $h_0 > 0$ is independent of h.

This theorem shows that presymplectic integration can provide answers to the questions posed earlier regarding the possibility of rate-preserving discretization of dissipative systems. The assumptions of the theorem are mild, and the restriction on t_{ℓ} may be irrelevant in practice. More importantly, the error in (2.8) is small and improves with r, though it is dominated by η_2 which suggests that, in this context, higher-order integrators are not likely to be needed. Note that choosing a suitable η_2 is essential since it can make the error negligible; e.g., with $\eta_2 \sim t$ the error is exponentially small.

There is another important aspect of presymplectic integrators worth noting. Since they exactly preserve the phase space geometry, they reproduce the qualitative features of the phase portrait and in particular the stability of critical points. This is not the case for typical discretizations, which in general introduce spurious damping or excitation. In short, given any suitable dissipative system obtained from (2.3), presymplectic integrators constitute a general approach for the construction of optimization algorithms that are guaranteed to respect the stability and rates of convergence of the underlying continuous system. We carry out such an approach and consider explicitly the case of the Bregman Hamiltonian in Section 6 and also provide general examples in Appendix C.

³This condition is satisfied by a large class of methods, even including simple ones such as the explicit Euler method which do not preserve any dynamical invariant [21, 22].

3 Hamiltonian Systems and Symplectic Integrators

In the remainder of the paper we provide a complete theoretical derivation justifying Theorem 2.1. To build up to that derivation, we first recall several essential concepts from backward error analysis, symplectic geometry, and dynamical systems. These concepts are necessary for an understanding of how one can draw conclusions about structure-preserving methods without a discrete-time analysis. For further background on the basic differential geometry that we use to develop our ideas, we refer the reader to Appendix A.

3.1 Numerical integrators and modified equations

Let \mathcal{M} be an *n*-dimensional smooth manifold, and let (\mathcal{U}, x) be a chart such that every point $p \in \mathcal{U} \subset \mathcal{M}$ has local coordinates x^1, \ldots, x^n in \mathbb{R}^n . From now on we refer to a point p by its coordinates x. Given a vector field $X \in T\mathcal{M}$, where $T\mathcal{M}$ denotes the tangent bundle, one has the system of differential equations

$$\frac{dx^{j}}{dt} = X^{j}(x), \qquad x^{j}(0) = x_{0}^{j} \qquad (j = 1, \dots, n).$$
(3.1)

This vector field can be represented by the differential operator⁴

$$X(x) = X^j(x)\partial_j, \tag{3.2}$$

where $\partial_1, \ldots, \partial_n$ is the induced coordinate basis in $T\mathcal{M}$. An integral curve of (3.1) defines a flow $\varphi_t : \mathcal{M} \to \mathcal{M}$. Notice that (3.1) has unique solutions, at least locally, since X is locally Lipschitz due to the smoothness of \mathcal{M} . The flow can be represented by the exponential map

$$\varphi_t = e^{tX} = I + tX + \frac{1}{2}t^2X^2 + \cdots$$
 (3.3)

Through its pullback, denoted by φ_t^* , the Lie derivative of a tensor α of rank (0, q) along X is defined by

$$(\mathcal{L}_X \alpha)(x) \equiv \left. \frac{d}{dt} \right|_{t=0} \varphi_t^* \alpha(x) = \lim_{t \to 0} \frac{\varphi_t^* \alpha(\varphi_t(x)) - \alpha(x)}{t}.$$
(3.4)

The dynamical system is said to preserve α if and only if $\mathcal{L}_X \alpha = 0$.

Let $\phi_h : \mathbb{R}^n \to \mathbb{R}^n$ be a numerical integrator of order $r \ge 1$ for the system (3.1)—as defined in (2.5) and (2.6). Since X is locally Lipschitz, we have $||X(y) - X(x)|| \le L_X ||y - x||$ for a constant $L_X > 0$ and for all $x, y \in \mathcal{M}$ in some region of interest. It follows from classical results [21, 22] that there exists $L_{\phi} > L_X$ such that we obtain control on a global error:

$$\|\phi_h^{\ell}(x_0) - \varphi_{t_{\ell}}(x_0)\| \le C(e^{L_{\phi}t_{\ell}} - 1)h^r,$$
(3.5)

⁴We use Einstein summation convention throughout the paper.

for some constant C > 0 and initial state $x_0 \in \mathcal{M}$. Note that we have denoted $\phi_h^{\ell} \equiv \phi_h \circ \cdots \circ \phi_h$. Thus, for a fixed t_{ℓ} the numerical method is accurate up to order $\mathcal{O}(h^r)$. However, if t_{ℓ} is free the error grows exponentially. For this reason, although higher-order methods may provide accurate solutions in a short span of time, they can be inaccurate and unstable for large times.

Formally, every numerical integrator ϕ_h can be seen as the *exact flow* of a modified or perturbed system, [18–22, 36]

$$\frac{dx^{j}}{dt} = \tilde{X}^{j}(x), \qquad \tilde{X} \equiv X + \Delta X_{1}h + \Delta X_{2}h^{2} + \cdots, \qquad (3.6)$$

where the ΔX 's are expressed in terms of X and its derivatives. We refer to X as the perturbed or *shadow vector field*. In general, the series in (3.6) is divergent, and it is necessary to consider a truncation [36]. Following [19], suppose we have found a truncation⁵

$$\tilde{X}_k = X + \Delta X_1 h + \dots + \Delta X_k h^k \qquad (k \ge r),$$
(3.7)

such that ϕ_h is an integrator of order k, namely $\|\phi_h(x) - \varphi_{h,\tilde{X}_k}(x)\| = \mathcal{O}(h^{k+1})$, where φ_{h,\tilde{X}_k} denotes the exact flow of (3.6) with \tilde{X} replaced by \tilde{X}_k . Define

$$\Delta X_{k+1}(x) \equiv \lim_{h \to 0} \frac{\phi_h(x) - \varphi_{h, \tilde{X}_k}(x)}{h^{k+1}}.$$
(3.8)

Then one can show that

$$\tilde{X}_{k+1} \equiv \tilde{X}_k + \Delta X_{k+1} h^{k+1} \tag{3.9}$$

yields a flow for which ϕ_h is an integrator of order k+1. Proceeding inductively, one can find higher-order vector fields \tilde{X}_k , with increasing k, such that φ_{h,\tilde{X}_k} becomes closer and closer to ϕ_h . However, since (3.6) eventually diverges, there exists a truncation point k^* such that $\|\phi_h(x) - \varphi_{h,\tilde{X}_{k^*}}(x)\|$ is as small as possible. Finding k^* and estimating the size of the ΔX 's terms is quite technical, but it has been carried out in seminal work [18], and the approach has been further improved in subsequent literature [19,36].

Theorem 3.1 (see [19]). Assume that X is real analytic and bounded on a compact subset of its domain. Assume that the numerical method ϕ_h is real analytic. Then, there exists a family of shadow vector fields \tilde{X} such that

$$\left\|X(x) - \tilde{X}(x)\right\| = \mathcal{O}(h^r), \qquad \left\|\phi_h(x) - \varphi_{h,\tilde{X}}(x)\right\| \le Che^{-r}e^{-h_0/h}, \tag{3.10}$$

where the constants $C, h_0 > 0$ do not dependent on the step size h.

This result holds in full generality, for any autonomous dynamical system and any numerical integrator. Next we will discuss how it becomes particularly useful in the case of conservative Hamiltonian systems.

⁵Which exists for k = r by assumption.

3.2 Conservative Hamiltonian systems

Before talking about symplectic integrators, we need to introduce conservative Hamiltonian systems. Hamiltonian systems are ubiquitous because they are naturally attached to the geometry of the cotangent bundle $T^*\mathcal{M}$ of any differentiable manifold \mathcal{M} [16, 17]. We provide a concise introduction to key results that will be necessary later.

Definition 3.2. Let \mathcal{M} be an even-dimensional smooth manifold supplied with a closed nondegenerate 2-form ω . More precisely:

- 1. $d\omega = 0;$
- 2. On any tangent space $T_x\mathcal{M}$, if $\omega(X,Y) = 0$ for all $Y \neq 0 \in T_x\mathcal{M}$, then X = 0.

Then ω is called a symplectic structure and (\mathcal{M}, ω) a symplectic manifold.

Theorem 3.3. Let q^1, \ldots, q^n be local coordinates of \mathcal{M} and $q^1, \ldots, q^n, p_1, \ldots, p_n$ the corresponding induced coordinates of the cotangent bundle $T^*\mathcal{M}$. Then $T^*\mathcal{M}$ admits a closed nondegenerate symplectic structure,

$$\omega = dp_j \wedge dq^j, \tag{3.11}$$

and is therefore a symplectic manifold.

Proof. There always exists a globally defined Liouville-Poincaré 1-form, $\lambda \equiv p_j dq^j \in T^*\mathcal{M}$ [16]. Applying the exterior derivative induces the Poincaré 2-form $\omega \equiv d\lambda = dp_j \wedge dq^j$. Since $d^2 = 0$ (see A.9) we trivially have $d\omega = 0$. It is also easy to see that ω is nondegenerate; e.g., in a matrix representation $\omega = \begin{pmatrix} 0 & -I \\ +I & 0 \end{pmatrix}$. Thus $\det(\omega) = 1 \neq 0$.

Theorem 3.4. A dynamical system with the phase space $T^*\mathcal{M}$ preserves the symplectic structure (3.11) if and only if it is (locally) a conservative Hamiltonian system, namely has the form in (2.2) with a time-independent Hamiltonian H = H(q, p).

Proof. Let $\gamma : \mathbb{R} \to \mathcal{M}$ be a curve parametrized by t, so that $q^j(t) \equiv (q^j \circ \gamma)(t)$ and $p_j(t) \equiv (p_j \circ \gamma)(t)$ are time-dependent coordinates over $T^*\mathcal{M}$. Consider the tangent vector $X \equiv \frac{d}{dt}$ to this curve, which in a coordinate basis is given by

$$X = \frac{dq^j}{dt}\frac{\partial}{\partial q^j} + \frac{dp_j}{dt}\frac{\partial}{\partial p_j}.$$
(3.12)

Note that X lives in the tangent bundle of the phase space $T^*\mathcal{M}$. The vector field X preserves (3.11) if and only if $\mathcal{L}_X \omega = 0$. Recalling Cartan's magic formula

$$\mathcal{L}_X = d \circ i_X + i_X \circ d, \tag{3.13}$$

we conclude that $(d \circ i_X)(\omega) = 0$. The Poincaré lemma thus implies the existence of a (local) Hamiltonian function $H: T^*\mathcal{M} \to \mathbb{R}$, such that

$$i_X(\omega) = -dH. \tag{3.14}$$

This is actually Hamilton's equations (2.2) in disguise; indeed, in component form we have $i_X(\omega) = \dot{p}_j dq^j - \dot{q}^j dp_j$, which by comparison with (3.14) yields (2.2). We have just shown that a vector field that preserves the symplectic form (3.11) genererates Hamiltonian dynamics. It is now easy to show the converse, namely that the flow of a Hamiltonian system preserves the 2-form (3.11). Given a Hamiltonian H, we have the equations of motion (2.2) with an associated vector field X_H . These equations can equivalently be written in the form (3.14), as already shown. Using the identities $d^2H = 0$ and $d\omega = 0$ in (3.13) implies $\mathcal{L}_{X_H}\omega = 0$. \Box

Finally, another fundamental property of Hamiltonian systems is the conservation of energy, which follows immediately from the equations of motion (2.2):

$$\frac{dH}{dt} = 0. \tag{3.15}$$

Thus, any conservative Hamiltonian system have two fundamental properties: 1. its flow preserve the symplectic structure and 2. the Hamiltonian is a constant of motion.

3.3 Symplectic integrators

Consider the following class of numerical methods.

Definition 3.5. A numerical integrator ϕ_h for a conservative Hamiltonian system is a symplectic integrator if it preserves (3.11), i.e., $\phi_h^* \circ \omega \circ \phi_h = \omega$ where ϕ_h^* is the pullback.

It is now easy to see that the flow of the perturbed system (3.6) associated to a symplectic integrator also preserves ω exactly. As a consequence, Theorem 3.4 implies that the perturbed system must be Hamiltonian; i.e., there exists a shadow Hamiltonian \tilde{H} which is a perturbed version of H. Indeed, since ϕ_h preserves ω by assumption, and the flow of the perturbed system (3.6) is an exact description of ϕ_h , the shadow vector field \tilde{X} obeys $\mathcal{L}_{\tilde{X}}\omega = 0$. Replacing the expansion (3.6) and using the linearity of the Lie derivative in the vector field implies $\mathcal{L}_{\Delta X_j}\omega = 0$ for each $j = 1, 2, \ldots$. Hence, Theorem 3.4 implies that not only \tilde{X} but all ΔX_j 's are Hamiltonian vector fields. That is, by the same argument leading to (3.14), there exists a function \tilde{H} and functions H_j 's such that

$$i_{\tilde{X}}(\omega) = -dH, \qquad i_{\Delta X_j}(\omega) = -dH_j.$$
(3.16)

Using the series (3.6), combining these two equations, and using (3.14), we have:

$$-d\tilde{H} = i_X(\omega) + \sum_j h^j i_{\Delta X_j}(\omega) = -dH - \sum_j h^j dH_j, \qquad (3.17)$$

or equivalently

$$\tilde{H} = H + hH_1 + h^2H_2 + \cdots$$
 (3.18)

Moreover, the shadow Hamiltonian \tilde{H} is exactly conserved by the perturbed system, or equivalently by the symplectic integrator ϕ_h : $\tilde{H} \circ \phi_h = \tilde{H}$. If we now consider a truncation (3.9), or equivalently if we truncate (3.18), then the flow φ_{h,\tilde{X}_k} exactly conserves the truncated shadow Hamiltonian \tilde{H}_k :

$$\tilde{H}_k \circ \varphi_{h,\tilde{X}_k} = \tilde{H}_k. \tag{3.19}$$

We are now ready to state the most important property of symplectic integrators. Adapting ideas from [18] we can prove the following.

Theorem 3.6. Let ϕ_h be a symplectic integrator of order r. Assume that the Hamiltonian H is Lipschitz. Then ϕ_h conserves H up to

$$H \circ \phi_h^\ell = H \circ \varphi_{\ell h} + \mathcal{O}(h^r), \tag{3.20}$$

for exponentially large times $t_{\ell} = \mathcal{O}(h^r e^r e^{h_0/h})$. Recall that $\varphi_{\ell h}$ is the true flow and $H = H \circ \varphi_{\ell h}$ is conserved, thus $H(q_{\ell}, p_{\ell}) = H(q_0, p_0) + \mathcal{O}(h^r)$.

Proof. Let $\tilde{X} \equiv \tilde{X}_k$ be a truncated shadow vector field associated to ϕ_h —in what follows we omit k for simplicity. We already know that (3.19) holds true, hence

$$\tilde{H} \circ \phi_h^\ell - \tilde{H} = \sum_{i=0}^{\ell-1} \left(\tilde{H} \circ \phi_h^{i+1} - \tilde{H} \circ \phi_h^i \right) = \sum_{i=0}^{\ell-1} \left(\tilde{H} \circ \phi_h - \tilde{H} \circ \varphi_{h,\tilde{X}} \right) \circ \phi_h^i, \tag{3.21}$$

where the first equality is an identity. Let \tilde{L} be the Lipschitz constant of \tilde{H} . Using the second relation of (3.10) we have:

$$\left| \left(\tilde{H} \circ \phi_{h}^{\ell} - \tilde{H} \right)(x_{0}) \right| \leq \tilde{L} \sum_{i=0}^{\ell-1} \left\| \phi_{h}(x_{i}) - \varphi_{h,\tilde{X}}(x_{i}) \right\| \leq \tilde{L} C \ell h \, e^{-r} e^{-h_{0}/h}, \tag{3.22}$$

so ϕ_h preserves \tilde{H} up to an exponentially small error in h^{-1} . In order to approximate H, write (3.18) in the form $\tilde{H} = H + \mathcal{O}(h^r)$ —since ϕ_h is order r—to obtain (3.20). Note that to ensure that the contribution of (3.22) remains smaller than $\mathcal{O}(h^r)$ we have to choose $t_\ell = h\ell$ such that $t_\ell e^{-r} e^{-h_0/h} \sim h^r$.

Theorem 3.6 explains the benefits of symplectic integrators. Besides preserving the symplectic structure of the system exactly, such methods generate solutions that remain within $\mathcal{O}(h^r)$ of the true energy surface of the system, and thus exhibit long-term stability. It is well known that in practice symplectic integrators tend to outperform alternative approaches when simulating conservative Hamiltonian systems [23–25].

The situation is quite different in the case of dissipative, or more generally time-dependent, Hamiltonian systems. Even if one applies a symplectic integrator to the system written in the extended phase space—so as to transform the nonautonomous system into an autonomous one—the Hamiltonian is no longer conserved. The conservation of the Hamiltonian is the most basic assumption underneath Theorem 3.6; if (3.15) is no longer true, then (3.19) is no longer true, and the argument leading to (3.21) and (3.22) breaks down. Therefore, it is not guaranteed that applying a symplectic integrator to a dissipative Hamiltonian system will closely reproduce the Hamiltonian, which is varying over time, nor is it guaranteed that the method will exhibit long-term stability. One of the main contributions of the current paper is to show that Theorem 3.6 can be extended to a dissipative setting despite the nonexistence of a conservation law.

4 Dissipative Hamiltonian Systems

Consider a time-dependent Hamiltonian H = H(t, q, p).⁶ The evolution of the system is still governed by Hamilton's equations (4.3), however, the energy conservation law (3.15) no longer holds and is replaced by

$$\frac{dH}{dt} = \frac{\partial H}{\partial t}.$$
(4.1)

We begin by introducing an (2n+1)-dimensional extended phase space $T^*\overline{\mathcal{M}}$ in which time becomes a new coordinate, so that $q^0 = t, q^1, \ldots, q^n, p_1, \ldots, p_n$ are local coordinates of $T^*\overline{\mathcal{M}}$. The evolution of the system is now generated by the vector field

$$X \equiv X^{\mu}\partial_{\mu} = \frac{\partial}{\partial q^{0}} + \frac{dq^{j}}{ds}\frac{\partial}{\partial q^{j}} + \frac{dp_{j}}{ds}\frac{\partial}{\partial p_{j}}, \qquad (4.2)$$

where $\mu = 0, 1, ..., n, X^0 = 1$, and s denotes the "new time parameter." Thus, the nonautonomous Hamiltonian system (2.2) is equivalent to the autonomous system

$$\frac{dq^0}{ds} = 1, \qquad \frac{dq^j}{ds} = \frac{\partial H}{\partial p_j}, \qquad \frac{dp_j}{ds} = -\frac{\partial H}{\partial q^j}, \tag{4.3}$$

over $T^*\bar{\mathcal{M}}$, where $H = H(q^0, \ldots, q^n, p_1, \ldots, p_n)$ is independent of s. Importantly, $T^*\bar{\mathcal{M}}$ is odd-dimensional so this phase space is no longer a symplectic manifold.

We thus introduce another dimension by adding a conjugate momentum p_0 that pairs with $q^0 = t$. To make the distinction clear, we denote the (n + 1)-dimensional configuration manifold by $\hat{\mathcal{M}} \equiv \mathbb{R} \times \mathcal{M}$, which has local coordinates q^0, \ldots, q^n . The associated cotangent bundle $T^* \hat{\mathcal{M}}$ is now of even dimensionality 2n + 2, with coordinates $q^0, \ldots, q^n, p_0, \ldots, p_n$. Therefore, $T^* \hat{\mathcal{M}}$ has a natural Liouville-Poincaré 1-form $\Lambda = p_{\mu} dq^{\mu}$ which, as in the proof of Theorem 3.3, induces the closed nondegenerate Poincaré 2-form:

$$\Omega \equiv d\Lambda = dp_{\mu} \wedge dq^{\mu}. \tag{4.4}$$

⁶We often refer to "dissipative Hamiltonian systems," though the reader should keep in mind that everything we say actually holds for any explicitly time-dependent Hamiltonian system.

This makes $T^* \hat{\mathcal{M}}$ a proper symplectic manifold. Requiring that the vector field

$$Y \equiv Y^{\mu}\partial_{\mu} = \frac{dq^{\mu}}{ds}\frac{\partial}{\partial q^{\mu}} + \frac{dp_{\mu}}{ds}\frac{\partial}{\partial p_{\mu}}$$
(4.5)

preserves the symplectic structure (4.4) implies, by the argument leading to (3.14), that

$$i_Y \Omega = -d\mathscr{H},\tag{4.6}$$

for some Hamiltonian $\mathscr{H}: T^* \hat{\mathcal{M}} \to \mathbb{R}$. In components, we have Hamilton's equations:

$$\frac{dq^0}{ds} = \frac{\partial \mathscr{H}}{\partial p_0}, \qquad \frac{dp_0}{ds} = -\frac{\partial \mathscr{H}}{\partial q^0}, \qquad \frac{dq^j}{ds} = \frac{\partial \mathscr{H}}{\partial p_j}, \qquad \frac{dp_j}{ds} = -\frac{\partial \mathscr{H}}{\partial q^j}, \tag{4.7}$$

where j = 1, ..., n ranges over the spatial components. Moreover, since now \mathscr{H} does not depend explicitly on time s, this system is actually conservative:

$$\frac{d\mathscr{H}}{ds} = 0. \tag{4.8}$$

At this stage, the higher-dimensional system (4.7) is not equivalent to (4.3). In order to create this equivalence we need to impose constraints, which means fixing a gauge. Note that the symplectic form (4.4) is $\Omega = dp_0 \wedge dq^0 + \omega$, where ω is the symplectic form (3.11) for the original phase space $T^*\mathcal{M}$. Requiring the vector fields (4.5) and (4.2) to be the same, and using (4.6) together with (3.14), yields:

$$0 = i_Y \Omega + d\mathscr{H} = i_X dp_0 \wedge dq^0 + i_X \omega + d\mathscr{H} = d(-p_0 - H + \mathscr{H}).$$

$$(4.9)$$

Thus, up to an irrelevant constant, we must have

$$\mathscr{H}(q^0, \dots, q^n, p_0, \dots, p_n) = p_0(s) + H(q^0, \dots, q^n, p_1, \dots, p_n).$$
(4.10)

Note that we carefully made the variable dependencies of each term explicit. This equation defines an *embedded submanifold* in the symplectic manifold $T^*\hat{\mathcal{M}}$. Hence, the dynamics of the dissipative Hamiltonian system (4.2) lies on a hypersurface of constant energy \mathscr{H} . More precisely, from (4.10) the first equation in (4.7) gives

$$\frac{dq^0}{ds} = 1,\tag{4.11}$$

which together with the two last equations of (4.7) becomes precisely the original dissipative system (4.3) over the extended phase space $T^*\bar{\mathcal{M}}$. The second equation of (4.7) reproduces the dissipation given by (4.1):

$$\frac{dp_0}{ds} = -\frac{\partial H}{\partial q^0} = -\frac{dH}{ds},\tag{4.12}$$



Figure 1: The symplectification consists of the embeddings $T^*\mathcal{M} \hookrightarrow T^*\bar{\mathcal{M}} \hookrightarrow T^*\hat{\mathcal{M}}$, each of dimension 2n, 2n + 1, and 2n + 2, respectively. The original dissipative Hamiltonian system has phase space $T^*\mathcal{M}$. When the system is written in autonomous form it has phase space $T^*\bar{\mathcal{M}}$, which is a presymplectic manifold that can naturally be embedded as a submanifold of constant energy $\mathscr{H} = 0$ in the symplectic manifold $T^*\hat{\mathcal{M}}$, which can be associated to the phase space of a conservative Hamiltonian system.

where the second equality follows from (4.8) together with (4.10). Hence, up to another irrelevant constant, we have

$$p_0(s) = -H(s), (4.13)$$

with H(s) = H(q(s), p(s)). Note that in (4.13) the Hamiltonian is solely a function of time; the actual trajectories have been replaced. Therefore, p_0 is completely fixed.

We remark that (4.11) and (4.13) are specific choices of coordinates on $T^*\mathcal{M}$ which remove the spurious degrees of freedom that are not present in the original system. This procedure of embedding the phase space of the dissipative Hamiltonian system (4.3), which does not admit a symplectic structure, into a higher-dimensional symplectic manifold is called *symplectification* [16, 17]. We provide an illustration in Figure 1.

Let us comment on another point behind the gauge choice (4.11) and (4.13). On the extended phase space $T^*\bar{\mathcal{M}}$ it is still possible to distinbluesh between position q^0 and momentum p_0 . The flow $\varphi_s = e^{sX}$ defines orbits $s \mapsto q^{\mu}(s)$ on $\hat{\mathcal{M}}$. Such orbits are considered equivalent by time reparametrization, $s \mapsto s'(s)$. However, to match the original orbits of the time-dependent Hamiltonian system over \mathcal{M} , one must fix $s = q^0 = t$. This means fixing a reference frame on $T^*\bar{\mathcal{M}}$. For this reason, time-dependent Hamiltonian systems are not covariant and one *is not free to reparametrize the original time t*. In other words, since the Hamiltonian system is explicitly time-dependent, time transformations are not canonical.

4.1 Presymplectic manifolds

We are now in a position to delineate the specific geometry underlying dissipative Hamiltonian systems. First, we introduce a useful generalization of symplectic manifolds [16].

Definition 4.1. Let \mathcal{M} be a differentiable manifold of dimension $(2n + \bar{n})$, $\bar{n} \geq 0$, supplied with a 2-form ω of rank 2n everywhere. Then ω is called a presymplectic form and (\mathcal{M}, ω) a presymplectic manifold.

When $\bar{n} = 0$ this definition reduces to that of a symplectic manifold, and when $\bar{n} = 1$ it reduces to the definition of a weak contact manifold. A weak contact manifold supplied with a 1-form ϑ obeying $\vartheta \wedge (d\vartheta)^n \neq 0$ is a contact manifold [16,17]. There is an alternative theory of contact manifolds that are intimately related to symplectic manifolds. In particular, they correspond to a particular case of presymplectic manifolds and can naturally be embedded as hypersurfaces in their symplectification, which is a symplectic manifold in 2n + 2 dimensions that is associated to \mathcal{M} . Furthermore, in Appendix B we show that a generalization of the so-called conformal Hamiltonian systems [32] also corresponds to a particular case of the time-dependent Hamiltonian formalism discussed above.

In order to project the symplectic form (4.4) into the hypersurface defined by (4.10) we can simply substitute the gauge choice $q^0 = t$ and $p_0 = -H$ to obtain

$$\Omega = -dH \wedge dt + \omega, \tag{4.14}$$

where we recall that ω is the original symplectic form of $T^*\mathcal{M}$. Moreover, the equations of motion (4.6) reduce to

$$i_X \Omega = 0, \qquad i_X dt = 1. \tag{4.15}$$

The vector field X, whose flow $\varphi_s = e^{sX}$ generates the dynamical evolution on $\hat{\mathcal{M}}$, is thus a zero mode of Ω . Note that, from Cartan's formula (3.13), if we require that a vector field X obeys (4.15), then it immediately implies $\mathcal{L}_X \Omega = 0$ so that Ω is preserved. The symplectic form Ω is closed; however, when projected into $T^* \bar{\mathcal{M}}$ it becomes *degenerate*. One can see this through the matrix representation [37]:

$$\Omega = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -I \\ 0 & +I & 0 \end{pmatrix},$$
(4.16)

which has a vanishing determinant. Therefore, the phase space of the dissipative Hamiltonian system written in the autonomous form (4.3) is a presymplectic manifold (see again Figure 1).

4.2 Presymplectic integrators

Since a dissipative system (4.3) admits a symplectification, we can construct structurepreserving discretizations by imposing constraints on a symplectic integrator. More precisely, we can apply any standard symplectic integrator to the higher-dimensional conservative system (4.7). Then by the gauge choice in (4.11) and (4.13) we obtain an integrator for the dissipative system (4.3). Such a method will preserve a presymplectic structure. Accordingly, we refer to such integrators as *presymplectic integrators*.

Definition 4.2. A numerical map ϕ_h is a presymplectic integrator for a time-dependent Hamiltonian system if it is obtained from a symplectic integrator for its symplectification under the gauge fixing (4.13) and (4.11).

Since the Hamiltonian \mathscr{H} of the higher-dimensional system (4.7) is conserved, we can apply standard results for symplectic integrators to derive conclusions about presymplectic integrators and thereby derive properties of systems without an underlying conservation law.

Theorem 4.3. A presymplectic integrator ϕ_h of order r preserves the time-varying Hamiltonian (assumed to be Lipschitz) up to

$$H \circ \phi_h^\ell = H \circ \varphi_{\ell h} + \mathcal{O}(h^r), \tag{4.17}$$

for exponentially large simulation times $s_{\ell} = h\ell = \mathcal{O}(h^r e^r e^{h_0/h}).$

Proof. Let $\Phi_h : \mathbb{R}^{2n+2} \to \mathbb{R}^{2n+2}$ be a symplectic integrator for the high-dimensional Hamiltonian system (4.7). Since \mathscr{H} is conserved, Theorem 3.6 implies

$$\mathscr{H} \circ \Phi_h^{\ell} = \mathscr{H} \circ \Psi_{\ell h} + \mathcal{O}(h^r), \tag{4.18}$$

with $s_{\ell} = \mathcal{O}(h^r e^r e^{h_0/h})$ and where Ψ_s denotes the true flow. With the gauge choice (4.11) and (4.13) the system is projected into the hypersurface (4.10), thus Ψ_s reduces to φ_s , which is the true flow of (4.3), and Φ_h reduces to $\phi_h : \mathbb{R}^{2n+1} \to \mathbb{R}^{2n+1}$, which approximates φ_s . Thus, using these maps together with the substitution of (4.10) into (4.18), we conclude:

$$H \circ \phi_h^\ell + p_0(\ell h) = H \circ \varphi_{\ell h} + p_0(\ell h) + \mathcal{O}(h^r).$$

$$(4.19)$$

It is important to recall that p_0 is a fixed function of time and this is why it can be canceled on both sides to give (4.17).

Theorem 4.3 is an extension of Theorem 3.6 to dissipative systems. The numerical solutions of presymplectic integrators are thus within a small and bounded error $\mathcal{O}(h^r)$ from the hypersurface (4.10); see Figure 2 for an illustration. Hence, whatever convergence or decay properties the dissipative system may have, a presymplectic integrator will closely reproduce its behavior.

In Appendix C we provide a detailed step-by-step procedure for constructing presymplectic integrators. The procedure is straightforward, essentially involving an application of symplectic integrators with a natural choice for incorporating the time variable. We now turn to the consequences of Theorem 4.3 to optimization.



Figure 2: A presymplectic integrator closely preserves the time-varying Hamiltonian; see Theorem 4.3. The numerical trajectories lie on the same presymplectic manifold as the continuous dissipative system and the numerical value $H(t_{\ell}, q_{\ell}, p_{\ell})$ exhibits a small and bounded error with respect to the true value, $H(t_{\ell}, q(t_{\ell}), p(t_{\ell}))$, of size $\mathcal{O}(h^r)$.

5 Preserving Convergence Rates in Optimization

In this section we consider how presymplectic integrators can be used to construct ratematching optimization algorithms. We consider a standard optimization problem (2.1) and we study dissipative systems obtained from the Hamiltonian (2.3), which we repeat here for convenience:

$$H = e^{-\eta_1(t)}T(t,q,p) + e^{\eta_2(t)}f(q),$$
(5.1)

where the functions η_1 and η_2 are positive and nondecreasing. They are responsible for introducing dissipation in the system. Recall that we assume that the kinetic energy T is Lipschitz continuous. Given a specific system, we assume that a convergence rate $f(q(t)) - f(q^*)$ as described in (2.4) is known, where $f(q^*)$ is a minimum of f in a neighborhood of the initial conditions. We are now able to prove Theorem 2.1 stated in the introduction as a consequence of Theorem 4.3.

Proof of Theorem 2.1. Substituting (5.1) into (4.17), where we now fix s = t, yields:

$$f(q_{\ell}) \leq f(q(t_{\ell})) + e^{-\eta_1(t_{\ell}) - \eta_2(t_{\ell})} \left\{ T\left(t_{\ell}, q(t_{\ell}), p(t_{\ell})\right) - T\left(t_{\ell}, q_{\ell}, p_{\ell}\right) \right\} + Kh^r e^{-\eta_2(t_{\ell})}, \quad (5.2)$$

for all $t_{\ell} = \mathcal{O}(h^r e^r e^{h_0/h})$ and for some constant K > 0. Recall that q_{ℓ} and $q(t_{\ell})$ are the discrete and continuous trajectories, respectively. Let L_T be the Lipschitz constant of the kinetic energy T. Substituting the global error (3.5) into the second term yields:

$$|f(q_{\ell}) - f(q(t_{\ell}))| \le h^r e^{-\eta_2(t_{\ell})} \left(K + L_T C(e^{L_{\phi} t_{\ell}} - 1) e^{-\eta_1(t_{\ell})} \right).$$
(5.3)

We thus conclude that

$$f(q_{\ell}) = f(q(t_{\ell})) + \mathcal{O}\left(h^r e^{-\eta_2(t_{\ell})}\right), \qquad (5.4)$$

provided

$$(e^{L_{\phi}t} - 1)e^{-\eta_1(t)} < \infty, \tag{5.5}$$

for all sufficiently large t.

Therefore, under suitable conditions, presymplectic integrators reproduce the continuous rates of convergence of dissipative Hamiltonian systems. We make several remarks:

- As mentioned earlier, the assumptions of Theorem 2.1 are mild; the kinetic energy being Lipschitz is naturally satisfied in a range of applications, and the Lipschitz condition (2.7) is also satisfied for a large class of integrators. The underlying reason is because \mathcal{M} is a smooth manifold.
- We used a conservative global error (3.5) to bound the kinetic term in (5.2). A specialized analysis may yield a better bound, which could allow condition (5.5) to be relaxed. However, this would have to be justified by a dedicated backward error analysis.
- Increasing the order of accuracy r influences (5.4) in a beneficial way. The error is dominated by $e^{-\eta_2}$, however, which suggests that, in this optimization context, using higher-order integrators may not be of significant practical importance. That said, higher-order methods tend to be more stable so there may be other benefits in their use.
- Interestingly, the error in (5.4) is even smaller than the error in approximating the Hamiltonian (4.17), thanks to the function η_2 . If η_2 grows just enough, this error may become completely negligible, even for large step sizes and integrators of low order.
- We recall that the restriction $t_{\ell} = h\ell \sim h^r e^r e^{h_0/h}$ may be irrelevant. To give an idea, with r = 1, $h_0 = 1$, and h = 0.1 we have iterations $\ell \sim 6 \cdot 10^4$, and with h = 0.01 we have $\ell \sim 10^{43}$.

5.1 The choice of damping

Let us comment on some consequences of the condition (5.5), which can impose some limitations. There are two ways to satisfy it. The first is if the method converges quickly so $e^{L_{\phi}t_{\ell}-\eta_1(t_{\ell})}$ remains bounded regardless of η_1 , or equivalently we enforce $t_{\ell} \leq \bar{t}$, where \bar{t} is a solution to

$$L_{\phi}\bar{t} - \eta_1(\bar{t}) \le C,\tag{5.6}$$

for some constant C > 0. The second way to satisfy (5.5) is if η_1 grows fast enough, allowing t_{ℓ} to be arbitrarily large. For instance, setting:

$$\eta_1(t) = \gamma t, \tag{5.7}$$

for some constant $\gamma \geq L_{\phi}$ suffices for all $t \geq 0$. Curiously, this choice implies constant damping and can be related to the heavy-ball method [15]; see also [5].⁷

⁷ With $\eta_1 = \eta_2 = \gamma t$ and $T = \frac{1}{2} ||p||^2$ into (5.1) the equations of motion give $\ddot{q} + \gamma \dot{q} = -\nabla f(q)$, which is a nonlinear generalization of the damped harmonic oscillator. The heavy-ball method [15] is actually a structure-preserving—a conformal symplectic—discretization of this system [5].

As an alternative, from (5.5) we have

$$\left(1 + L_{\phi}t + \frac{1}{2}(L_{\phi}t)^2 + \cdots\right)e^{-\eta_1(t)} < \infty,$$
(5.8)

which is satisfied if $(L_{\phi}t)^m e^{-\eta_1(t)} < \infty$ for all powers m; i.e., $\eta_1(t) \sim m \log(L_{\phi}t)$. Thus, one can choose

$$\eta_1(t) = \gamma \log t,\tag{5.9}$$

with a suitable $\gamma > 0$ to ensure that (5.8) holds up to some high-order term. Also curiously, the choice (5.9) is related to the damping of Nesterov's method [14] which attains the optimal convergence rate for convex functions $f^{.8}$

Note that the choice (5.9) can guarantee (5.8) only up to a finite power m, thus we are on the edge of violating this condition—in fact we are formally violating it but in practice this may yield a viable algorithm. On the other hand, the choice (5.7) may be overkill. It may be more reasonable to consider an intermediate alternative:

$$\eta_1(t) = \gamma_1 \log t + \gamma_2 t^{\delta}, \tag{5.10}$$

for constants $0 < \delta \leq 1$ and $\gamma_1, \gamma_2 > 0$. With this choice the condition on the *m*th term in the series (5.8) becomes $L_{\phi}^m t^{m-\gamma_1} e^{-\gamma_2 t^{\delta}} < \infty$, which can be guaranteed even if $\gamma_1 < m$ with suitable choices of γ_2 and δ .

Summing up, presymplectic integrators are able to match the continuous rates provided the condition (5.5) is satisfied, which involves an appropriate choice of damping. If the system is overdamped this condition is more likely to hold, however the system would tend to be slower. On the other hand, if the damping is weak the system tends to be fast, but at the cost of violating (5.5). We thus see the tradeoff is delicate and is problem-dependent in general, given that L_{ϕ} in (5.5) is related to the Lipschitz constant of the original vector field.

6 Bregman Dynamics

Given that the Bregman Hamiltonian [2] provides a general framework for deriving continuoustime optimization procedures, we consider this case in some detail. The Hamiltonian has the form

$$H = e^{\alpha + \gamma} \left\{ D_{h^{\star}} \left(\nabla h(q) + e^{-\gamma} p, \nabla h(q) \right) + e^{\beta} f(q) \right\},$$
(6.1)

where α , β , and γ are all functions of time t, and are required to satisfy the following scaling conditions:

$$\frac{d\beta}{dt} \le e^{\alpha}, \qquad \frac{d\gamma}{dt} = e^{\alpha}.$$
 (6.2)

⁸ Choosing $\eta_1 = \eta_2 = \gamma \log t$ yields the differential equation $\ddot{q} + \frac{\gamma}{t}\dot{q} = -\nabla f(q)$. Nesterov's method can be seen as a discretization of this system [1].

The kinetic energy in (6.1) is given in terms of the Bregman divergence,

$$D_h(y,x) \equiv h(y) - h(x) - \langle \nabla h(x), y - x \rangle, \qquad (6.3)$$

which is nonnegative for a given convex function $h : \mathcal{M} \to \mathbb{R}$. Recalling the definition of the convex dual, h^* of h, defined by the Legendre-Fenchel transformation:

$$h^{\star}(p) \equiv \sup_{v} \left\{ \langle p, v \rangle - h(v) \right\}, \tag{6.4}$$

one can show that Hamilton's equations are given by:¹

$$\dot{q} = e^{\alpha} \left\{ \nabla h^* \left(\nabla h(q) + e^{-\gamma} p \right) - q \right\}, \tag{6.5a}$$

$$\dot{p} = -e^{\alpha + \gamma} \nabla^2 h(q) \left\{ \nabla h^* \left(\nabla h(q) + e^{-\gamma} p \right) - q \right\} + e^{\alpha} p - e^{\alpha + \beta + \gamma} \nabla f(q).$$
(6.5b)

In the case where f is convex, the convergence rate of this system is given by [2]:

$$f(q(t)) - f(q^*) = \mathcal{O}\left(e^{-\beta(t)}\right).$$
(6.6)

Moreover, for a given α , the optimal rate is obtained with $\dot{\beta} = e^{\alpha}$, thus $\beta = \gamma + C$, for some constant C, and both are determined in terms of α . We thus have

$$f(q(t)) - f(q^*) = \mathcal{O}\left(e^{-\beta(t)}\right), \qquad \beta(t) = \int^t e^{\alpha(t')} dt', \qquad \gamma(t) = \beta(t) + C.$$
(6.7)

A choice considered by [2] is:

$$\alpha = \log c - \log t, \qquad \beta = c \log t + C, \qquad \gamma = c \log t \qquad (c > 0), \tag{6.8}$$

whereby the convergence rate (6.7) becomes the polynomial $\mathcal{O}(t^{-c})$. Another possibility is

$$\alpha = \log c, \qquad \beta = ct, \qquad \gamma = ct + C \qquad (c > 0), \tag{6.9}$$

leading to an exponential rate, $\mathcal{O}(e^{-ct})$.

6.1 Separable case

To consider a case where the Bregman Hamiltonian is separable, let us start with the choice $h(x) = \frac{1}{2}x \cdot Mx$, for a symmetric positive semidefinite matrix M, so that $h^*(x) = \frac{1}{2}x \cdot M^{-1}x$

$$\ddot{q} + \left(e^{\alpha} - \dot{\alpha}\right)\dot{q} + e^{2\alpha + \beta} \left[\nabla^2 h \left(q + e^{-\alpha} \dot{q}\right)\right]^{-1} \nabla f(q) = 0.$$

We see that α basically controls the damping, while $2\alpha + \beta$ increases the strength of the force $-\nabla f$. Both play a major role in the stability of the system, and they are not independent since $\dot{\beta} \leq e^{\alpha}$.

¹Equivalently, one can write the second-order differential equation

and the Bregman divergence is given by $D_{h^*}(y, x) = \frac{1}{2}(y-x) \cdot M^{-1}(y-x)$. In this case the Hamiltonian (6.1) takes the form

$$H = \frac{1}{2}e^{-\eta_1(t)} p \cdot M^{-1}p + e^{\eta_2(t)}f(q), \qquad (6.10)$$

where we defined

$$\eta_1 \equiv \gamma - \alpha, \qquad \eta_2 \equiv \alpha + \beta + \gamma.$$
 (6.11)

The quadratic kinetic energy is standard in classical mechanics and we see that M plays the role of a mass matrix. We thus have the following equations of motion:

$$\dot{q} = e^{-\eta_1(t)} M^{-1} p, \qquad \dot{p} = -e^{\eta_2(t)} \nabla f(q).$$
 (6.12)

One can now use any presymplectic integrator to obtain an optimization algorithm; see Appendix C. Because the Hamiltonian (6.10) is separable, standard approaches yield explicit methods which are convenient in practice. Assuming that we use a presymplectic integrator of order r, Theorem 2.1 tells us that the continuous rate of convergence will be preserved as

$$f(q_{\ell}) - f^{\star} = \mathcal{O}\left(e^{-\beta(t_{\ell})}\right) + \mathcal{O}\left(h^{r}e^{-\alpha(t_{\ell})-2\beta(t_{\ell})}\right)$$

$$\sim e^{-\beta(t_{\ell})}\left(1 + h^{r}e^{-\alpha(t_{\ell})-\beta(t_{\ell})}\right)$$

$$\sim e^{-\beta(t_{\ell})}, \qquad (6.13)$$

provided η_1 as defined in (6.11) obeys the condition (5.5).

We present two explicit methods. Using the presymplectic Euler method given by (C.2), which is of order r = 1, we obtain:

$$p_{\ell+1} = p_{\ell} - h e^{\eta_2(t_{\ell})} \nabla f(q_{\ell}),$$

$$t_{\ell+1} = t_{\ell} + h,$$

$$q_{\ell+1} = q_{\ell} + h e^{-\eta_1(t_{\ell})} M^{-1} p_{\ell+1}.$$

(6.14)

This algorithm requires only one gradient computation per iteration. In a similar way, using the presymplectic leapfrog method (C.9), which is of order r = 2, we obtain:

$$t_{\ell+1/2} = t_{\ell} + h/2,$$

$$q_{\ell+1/2} = q_{\ell} + (h/2)e^{-\eta_1(t_{\ell+1/2})}M^{-1}p_{\ell},$$

$$p_{\ell+1} = p_{\ell} - he^{\eta_2(t_{\ell+1/2})}\nabla f(q_{\ell+1/2}),$$

$$t_{\ell+1} = t_{\ell+1/2} + h/2,$$

$$q_{\ell+1} = q_{\ell+1/2} + (h/2)e^{-\eta_1(t_{\ell+1/2})}M^{-1}p_{\ell+1}.$$
(6.15)

This method again only requires one gradient computation per iteration, and it is more accurate. One can now choose any suitable scaling functions in (6.11) and substitute into either (6.14) or (6.15) to obtain a specific optimization algorithm.

6.2 Nonseparable case

We turn to the case of a nonseparable Hamiltonian, where the integrators considered thus far yield implicit updates that require solving nonlinear equations. This not only increases the computational burden but can affect the numerical stability. In particular, for the Bregman Hamiltonian (6.1), we need a construction suited to nonseparable cases. In order to obtain *explicit* methods, motivated by [38] we introduce extra degrees of freedom. Thus, given a Hamiltonian H(t, q, p), we double its degrees of freedom by introducing an augmented Hamiltonian:

$$\bar{H}(t,q,p,\bar{t},\bar{q},\bar{p}) \equiv H(t,q,\bar{p}) + H(\bar{t},\bar{q},p) + \frac{\xi}{2} \left(\|q-\bar{q}\|^2 + \|p-\bar{p}\|^2 \right).$$
(6.16)

Here $\xi > 0$ is a coupling constant that controls the strength of the last term, which forces $q = \bar{q}$ and $p = \bar{p}$.² The presymplectic structure of this system is now

$$\Omega = -dH \wedge dt + dp_j \wedge dq^j - dH \wedge d\bar{t} + d\bar{p}_j \wedge d\bar{q}^j \qquad (j = 1, \dots, n).$$
(6.17)

Hamilton's equations, obtained from (6.16), will preserve Ω . The equations of motion are equivalent to those of the original system when $q = \bar{q}$, $p = \bar{p}$ and $t = \bar{t}$. We thus propose the following numerical maps:

$$\begin{split} \phi_{h}^{A} \begin{pmatrix} t \\ q \\ p \\ \bar{t} \\ \bar{q} \\ \bar{p} \end{pmatrix} &= \begin{pmatrix} t \\ q \\ p - h \nabla_{q} H(t, q, \bar{p}) \\ \bar{t} + h \\ \bar{q} + h \nabla_{\bar{p}} H(t, q, \bar{p}) \\ \bar{p} \end{pmatrix}, \quad \phi_{h}^{B} \begin{pmatrix} t \\ q \\ p \\ \bar{t} \\ \bar{q} \\ \bar{p} \end{pmatrix} &= \begin{pmatrix} t \\ q + h \nabla_{p} H(\bar{t}, \bar{q}, p) \\ p \\ \bar{t} \\ \bar{q} \\ \bar{p} \end{pmatrix}, \quad \phi_{h}^{C} \begin{pmatrix} t \\ q \\ p \\ \bar{t} \\ \bar{q} \\ \bar{p} \end{pmatrix} &= \frac{1}{2} \begin{pmatrix} 2t \\ q + \bar{q} + \cos(2\xi h)(q - \bar{q}) + \sin(2\xi h)(p - \bar{p}) \\ p + \bar{p} - \sin(2\xi h)(q - \bar{q}) + \cos(2\xi h)(p - \bar{p}) \\ 2\bar{t} \\ q + \bar{q} - \cos(2\xi h)(q - \bar{q}) - \sin(2\xi h)(p - \bar{p}) \\ p + \bar{p} + \sin(2\xi h)(q - \bar{q}) - \cos(2\xi h)(p - \bar{p}) \end{pmatrix}. \end{split}$$
(6.18)

A presymplectic integrator for any (nonseparable) time-dependent Hamiltonian system can then be constructed by composing these maps. For instance, the Strang composition,

$$\phi_{h/2}^A \circ \phi_{h/2}^B \circ \phi_h^C \circ \phi_{h/2}^B \circ \phi_{h/2}^A, \tag{6.19}$$

is known to generate a method of order r = 2—this is the same composition as the leapfrog. Note that the maps (6.18) are completely explicit in all variables and this approach works for any general time-dependent Hamiltonian.

²It is not necessary to introduce the same term for t and \bar{t} since they will be be equal thanks to (4.11). The constant ξ has to be tuned in practice; see [38] for details.

In particular, for the Bregman Hamiltonian (6.1) it suffices to substitute the gradients $\dot{q} = \nabla_p H$ and $\dot{p} = -\nabla_q H$ from (6.5) into (6.18), followed by the composition (6.19). This yields an explicit, second-order integrator for the general Bregman dynamics. Thanks to Theorem 2.1, this generates an optimization algorithm that may closely preserve the continuous rate of convergence (6.6) for suitable functions α , β and γ .

7 Discussion

We have introduced "presymplectic integrators" as a class of discretizations that are suitable for the simulation of explicitly time-dependent, and in particular dissipative, Hamiltonian systems. This framework accommodates a large class of dissipative dynamical systems that are appropriate for applications to optimization and machine learning. We have also shown that, besides preserving the presymplectic structure of the system, such methods nearly preserve the time-varying Hamiltonian and exhibit long-term stability, despite the absence of a conservation law; see Theorem 4.3. This extends, into a dissipative setting, the most important property of symplectic integrators which are restricted to conservative systems. Thus, our approach and theoretical conclusions may be of interest not only to the field of optimization, but also to a variety of other disciplines where the simulation of dissipative systems play an important role, such as nonequilibrium statistical physics, complex systems, and economics.

Focusing on optimization, we showed that, for a general class of dissipative systems arising from a Hamiltonian in the form (2.3), presymplectic integrators are able to preserve the continuous rates of convergence up to a negligible error, as long as certain conditions are satisfied; see Theorem 2.1. This provides a systematic, first-principles approach to deriving "rate-matching" optimization algorithms, thereby obviating the need for a discrete-time convergence analysis.³ As a concrete example, we considered the general Bregman Hamiltonian, providing general methods from which specific optimization algorithms can be derived, in both the separable and the nonseparable case.

Finally, we comment on two problems that might deserve further study. First, we showed that condition (5.5) is essential and in particular leads to the damping strategies (5.7) and (5.9) related to the heavy ball and Nesterov's method, respectively. We also argued that a more elaborate choice (5.10) may be beneficial. Thus, finding the optimal damping for a given class of problems seems an interesting problem. Second, it would be appealing to find numerical schemes with a better global error—see equation (2.7)—or obtain improved bounds for existing methods, since this would automatically relax the condition (5.5).

³Of course, if one wants to study fine details of a specific algorithm's performance, a discrete analysis, together with an appropriate backward-error analysis, may be necessary.

Acknowledgements

We wish to thank Jelena Diakonikolas and Michael Muehlebach for helpful discussions. This work was supported by grant ARO MURI W911NF-17-1-0304.

A Background on Differential Geometry

In this section we provide a brief, informal overview of those elements of differential geometry that we need for our results. For a fuller presentation see any of the excellent textbooks on the subject (e.g., [16, 17, 37, 39, 40]).

Let $\mathcal{M} \equiv \mathcal{M}^n$ be a smooth *n*-dimensional manifold. Let $p \in \mathcal{M}$ and (\mathcal{U}, x) be a *chart* so that *p* can be assigned local coordinates, $x(p) \equiv (x^1(p), \ldots, x^n(p))$ in \mathbb{R}^n . The coordinates *x* and the point *p* are used interchangeably, and we often refer to the former to indicate a point on the manifold. To each $x \in \mathcal{M}$ there is an associated vector space $T_x\mathcal{M}$ called the *tangent space*. The coordinates *x* induces a basis $\partial_1, \ldots, \partial_n$ in $T_x\mathcal{M}$, where $\partial_j \equiv \frac{\partial}{\partial x^j}$. Thus, $V|_x = V^j(x)\partial_j$ is a representation of the vector $V|_x \in T_x\mathcal{M}$.⁴ The collection of all tangent vectors of \mathcal{M} form the *tangent bundle* $T\mathcal{M}$. One can then represent a (contravariant) vector field by the differential operator

$$V(x) \equiv V^{j}(x)\partial_{j},\tag{A.1}$$

which can be seen as a cross section of the tangent bundle $T\mathcal{M}$. For a given x, the vector field V assigns a single vector, $V(x) = V|_x$. Note that a point in $T\mathcal{M}$ has 2n coordinates, $x^1, \ldots, x^n, V^1, \ldots, V^n$.

To each $V|_x$ there is an associated dual vector, $\alpha|_x : T_x M \to \mathbb{R}$; i.e., α is a linear functional. Dual vectors are called covectors or 1-forms and they live on the *cotangent space* denoted by $T_x^* \mathcal{M}$, which is isomorphic to $T_x \mathcal{M}$. The collection of all 1-forms at every point of \mathcal{M} forms the *cotangent bundle* $T^* \mathcal{M}$. The coordinate basis x induces a dual basis dx^1, \ldots, dx^n in $T_x^* \mathcal{M}$, defined by $dx^j(\partial_k) = \partial_k(dx^j) = \delta_k^j$, where δ is the Kronecker delta. Similarly to (A.1) one can now represent a 1-form field $\alpha \in T^* \mathcal{M}$ as

$$\alpha(x) = \alpha_j(x)dx^j,\tag{A.2}$$

which is a cross section of the cotangent bundle $T^*\mathcal{M}$. The action of dual vectors is thus $\alpha(V)|_x = V(\alpha)|_x = \alpha_j(x)V^j(x)$. Note that $T^*\mathcal{M}$ is a 2*n*-dimensional space where a point has coordinates $x^1, \ldots, x^n, \alpha_1, \ldots, \alpha_n$. The cotangent bundle $T^*\mathcal{M}$ is very special since it can be shown that it is itself a symplectic manifold (see Theorem 3.3).

⁴ We use Einstein's summation convention throughout, where a pair of upper and lower indices are summed over; e.g., $X^j \partial_j \equiv \sum_{j=1}^n X^j \partial_j$, $\alpha_{jk} T^{\ell j k} \equiv \sum_{j=1}^n \sum_{k=n}^n \alpha_{jk} T^{\ell j k}$, etc. Upper indices denote components of vectors—also called contravariant vectors—in $T_x \mathcal{M}$, while lower indices denote components of dual vectors—also called covectors—which belong to the cotangent space $T_x^* \mathcal{M}$.

A general tensor T of rank (p,q) is a multilinear map $T : \bigotimes^p T_x^* \mathcal{M} \bigotimes^q T_x \mathcal{M} \to \mathbb{R}$. In a coordinate basis it is written as

$$T = T^{j_1 \dots j_p}_{k_1 \dots k_q} \partial_{j_1} \cdots \partial_{j_p} dx^{k_1} \cdots dx^{k_q}.$$
(A.3)

In calculations, it is useful to focus on the components $T^{j_1...j_p}_{k_1...k_q}$ alone and omit the basis altogether. A contravariant tensor is a tensor of rank (p, 0), for some $p \ge 1$, and a covariant tensor is a tensor of rank (0, q), for some $q \ge 1$. A *q*-form is a (0, q)-tensor which is totally antisymmetric in its indices. In a basis it is denoted as

$$\alpha = \frac{1}{q!} \alpha_{j_1 \dots j_q} dx^{j_1} \wedge \dots \wedge dx^{j_q}, \tag{A.4}$$

where \wedge denotes the *exterior product*.⁵ Given another *p*-form β , one can compose the (p+q)-form $\alpha \wedge \beta$ which obeys

$$\alpha \wedge \beta = (-1)^{pq} \beta \wedge \alpha. \tag{A.5}$$

It is useful to introduce a notation to denote the space of q-forms at x, namely $\bigwedge^q T^*_x \mathcal{M}$, and as before we obtain a bundle $\bigwedge^q T^* \mathcal{M}$ of q-form fields by allowing the coefficient $\omega_{j_1...j_q}(x)$ to depend on x.

Another important operation is the *exterior derivative*:

$$d: \bigwedge^{q} T^* \mathcal{M} \to \bigwedge^{q+1} T^* \mathcal{M}, \tag{A.6}$$

which can be defined componentwise as

$$d\alpha(x) \equiv \frac{1}{q!} \frac{\partial \alpha_{j_1\dots j_q}(x)}{\partial x^k} dx^k \wedge dx^{j_1} \wedge \dots \wedge dx^{j_q}.$$
 (A.7)

This is a linear operation, $d(c_1\alpha + c_2\beta) = c_1d\alpha + c_2d\beta$, for any forms α and β . Its main properties are given by the equation:

$$d(\alpha \wedge \beta) = (d\alpha) \wedge \beta + (-1)^q \alpha \wedge (d\beta), \tag{A.8}$$

and by the identity

$$d^2 = d \circ d = 0. \tag{A.9}$$

A differential form ω is said to be *closed* if $d\omega = 0$. A differential *q*-form ω is said to be *exact* if $\omega = d\lambda$ for some (q - 1)-form λ . Trivially, from (A.9) every exact form is closed. The *Poincaré lemma* ensures the converse, namely that every closed form is also exact.⁶

Given a vector $v \in T_x \mathcal{M}$ and a q-form $\alpha \in \bigwedge^q T_x^* \mathcal{M}$, the *interior product* $i_v \alpha$ is a (q-1)-form defined by

$$(i_v \alpha)(v_2, \dots, v_q) \equiv \alpha(v, v_2, \dots, v_q).$$
(A.10)

⁵For two 1-forms $\alpha, \beta \in T_x^* \mathcal{M}$ we have $(\alpha \wedge \beta)(v, w) \equiv \alpha(v)\beta(w) - \beta(v)\alpha(w)$, for $v, w \in T_x \mathcal{M}$.

⁶This holds for contractible manifolds, which is the case for smooth manifolds as considered in this paper.

In components, this is simply the contraction $(i_v \alpha)_{j_2...j_q} = v^k \alpha_{kj_2...j_q}$. The interior product is also linear and satisfies an analogous relation to (A.8):

$$i_v(\alpha \wedge \beta) = (i_v \alpha) \wedge \beta + (-1)^q \alpha \wedge (i_v \beta).$$
(A.11)

Since one can only operate on elements of the same vector space, it is necessary to introduce a mapping that makes it possible to move geometric objects over the manifold. In particular, given a function $F : \mathcal{M} \to \mathcal{N}$ between two manifolds \mathcal{M} and \mathcal{N} , for any function $g : \mathcal{N} \to \mathbb{R}$ one defines the *pushforward*—also called the differential—of $v \in T_x \mathcal{M}$ to be the vector $F_* v \in T_{F(x)} \mathcal{N}$ defined by the operation

$$(F_*v)(g) \equiv v(g \circ F). \tag{A.12}$$

The *pullback* goes in the opposite direction, i.e., given a q-form $\alpha \in T^*_{F(x)}\mathcal{N}$ we obtain the q-form $F^*\alpha \in T^*_x\mathcal{M}$ through

$$F^*\alpha(v_1,\ldots,v_q) \equiv \alpha(F_*v_1,\ldots,F_*v_q), \tag{A.13}$$

for $v_j \in T_x \mathcal{M}$, $j = 1, \ldots, q$. The pullback allows us to move q-forms over the manifold. By introducing the concept of a flow: $\varphi_t : \mathcal{M} \to \mathcal{M}$ induced by a vector field X of $T\mathcal{M}$, i.e., $\varphi_t = e^{tX}$, one can define the *Lie derivative* of a q-form as in (3.4). A flow φ_t is a diffeomorphism and thus $\varphi_t^* = (\varphi_{-t})_*$. In this case, one can consider the pullback of not only q-forms but arbitrary (p, q)-tensors. Thus, the Lie derivative of a general tensor field T can be defined by

$$\mathcal{L}_X T|_x \equiv \left. \frac{d}{dt} \right|_{t=0} \varphi_t^* T|_{\varphi_t(x)}. \tag{A.14}$$

Given a differentiable form ω and a vector field X, we say that X preserves ω if and only if

$$\mathcal{L}_X \omega = 0. \tag{A.15}$$

From the definition (A.14) this implies

$$\varphi_t^* \omega = \omega. \tag{A.16}$$

In this sense, any map $\phi : \mathcal{M} \to \mathcal{N}$ is said to be *canonical* if it preserves ω ; i.e., $\phi^* \omega = \omega$. In the case of Hamiltonian systems ω is the symplectic 2-form, and Hamiltonian flows generate canonical transformations, which are symplectomorphisms.

A very useful formula is Cartan's magic formula

$$\mathcal{L}_X \alpha = d \circ i_X \alpha + i_X \circ d\alpha, \tag{A.17}$$

for any differentiable form α . Some other useful formulas are

$$\mathcal{L}_X(\alpha \wedge \beta) = (\mathcal{L}_X \alpha) \wedge \beta + \alpha \wedge (\mathcal{L}_X \beta), \tag{A.18}$$

$$\mathcal{L}_{[X,Y]}\alpha = [\mathcal{L}_X, \mathcal{L}_Y]\alpha, \tag{A.19}$$

$$\mathcal{L}_X \circ d = d \circ \mathcal{L}_x, \tag{A.20}$$

$$i_{[X,Y]} = \mathcal{L}_X \circ i_Y - i_Y \circ \mathcal{L}_X. \tag{A.21}$$

Here, $[\mathcal{L}_X, \mathcal{L}_Y] \equiv \mathcal{L}_X \mathcal{L}_Y - \mathcal{L}_Y \mathcal{L}_X$ is the *Lie bracket*. The same definition holds for [X, Y] = XY - YX, and in this case one refers to $[\cdot, \cdot]$ as the *commutator* of two vector fields.

B Generalized Conformal Hamiltonian Systems

Conformal Hamiltonian systems [32] provide an alternative approach to introducing dissipation into Hamiltonian systems. In this approach, one modifies Hamilton's equations directly by adding a linear term in the momentum. It is easy to construct structure-preserving discretizations for these systems since one can split the system into conservative and dissipative parts, then apply a standard symplectic integrator to the former, while integrating the latter exactly. This approach has been recently explored in optimization [5]. The purpose of this section is to show that a generalization of conformal Hamiltonian systems correspond to a particular case of the explicit time-dependent Hamiltonian formalism introduced in Section 4. As a consequence, one can construct (generalized) conformal symplectic integrators from presymplectic integrators (see Definition 4.2).

Consider a time-independent Hamiltonian, H = H(q, p), and assume a modified form of Hamilton's equations given by

$$\frac{dq^j}{dt} = \frac{\partial H}{\partial p_j}, \qquad \frac{dp_j}{dt} = -\frac{\partial H}{\partial q^j} - \gamma(t)p_j \qquad (j = 1, \dots, n).$$
(B.1)

The conformal case [32] assumes that the damping coefficient $\gamma(t) = \gamma$ is constant. In this formulation, the Hamiltonian vector field is

$$X_H = \frac{\partial H}{\partial q^i} \frac{\partial}{\partial q^i} + \frac{\partial H}{\partial p_i} \frac{\partial}{\partial p_i},\tag{B.2}$$

and by the same geometric approach previously discussed one can see that the equations of motion (B.1) are equivalent to

$$i_{X_H}(\omega) = -dH - \gamma(t)\lambda, \tag{B.3}$$

with ω defined in (3.11) and where $\lambda \equiv p_j dq^j$ is the Liouville-Poincaré 1-form. From Cartan's formula (3.13) we conclude that the symplectic structure contracts as

$$\mathcal{L}_{X_H}\omega = -\gamma(t)\omega,\tag{B.4}$$

or equivalently⁷

$$\varphi_t^* \omega = e^{-\eta(t)} \omega, \qquad \eta(t) \equiv \int^t \gamma(t') dt'.$$
 (B.5)

Finally, in this setting H is the energy of the system and it dissipates as

$$\frac{dH}{dt} = -\gamma(t)\frac{\partial H}{\partial p_j}p_j.$$
(B.6)

⁷From this one concludes that the phase-space volumes contract as $\mathcal{L}_{X_H} \operatorname{vol}^{2n} = -n\gamma(t) \operatorname{vol}^{2n}$, so that $\varphi_t^* \operatorname{vol}^{2n} = e^{-nC(t)} \operatorname{vol}^{2n}$. This is a dissipative version of Liouville's theorem; note the dimension dependency.

We now show that generalized conformal Hamiltonian systems corresponds to a particular case of the time-dependent Hamiltonian formalism. Define the time-dependent Hamiltonian K by

$$K(t,Q,P) \equiv e^{\eta(t)} H(Q, e^{-\eta(t)} P), \qquad Q \equiv q, \qquad P \equiv e^{\eta(t)} p, \tag{B.7}$$

where H is the original Hamiltonian of (B.1). The standard Hamiltonian equations (2.2) yield:

$$\frac{dQ^{j}}{dt} = \frac{\partial K}{\partial P_{j}} = \frac{\partial H(q, p)}{\partial p_{j}}, \qquad \frac{dP_{j}}{dt} = -\frac{\partial K}{\partial Q^{j}} = -e^{\eta(t)}\frac{\partial H(q, p)}{\partial q^{j}}, \tag{B.8}$$

which when written in terms of (q, p) are precisely (B.1). Therefore, one can construct discretizations that preserve the contraction of the symplectic form (B.4) through presymplectic integrators (see Definition 4.2) with the explicit time-dependent Hamiltonian K.

C Constructing Presymplectic Integrators

As stated in Definition 4.2, a presymplectic integrator is a reduction of a higher-dimensional symplectic integrator under the specific choice of coordinates (4.11) and (4.13). To follow this prescription one must perform three steps:

- 1. Pick a symplectic integrator and apply it to the Hamiltonian system (4.7). This results in updates for (q^0, p_0) and for the spatial components (q^j, p_j) ;
- 2. Set $q^0 = t$ and ignore p_0 completely— p_0 is just the actual value of the Hamiltonian as a function of time and does not participate in the dynamics;
- 3. Set s = t.

While these formal steps make clear that we are respecting the symplectification prescription, in practice, these three steps can be reduced to the following:

1. Apply any symplectic integrator to the time-dependent Hamiltonian H(t, q, p) in the "natural way." By this we mean to simply include additional updates for the time variable t with the same rule as the coordinates q^{j} .

We will provide some examples below that should make this procedure clear.

Presymplectic Euler For a conservative Hamiltonian system, one has the following version of the first-order symplectic Euler method [22]:

$$p_{\ell+1} = p_{\ell} - h \nabla_q H(q_{\ell}, p_{\ell+1}),$$

$$q_{\ell+1} = q_{\ell} + h \nabla_p H(q_{\ell}, p_{\ell+1}).$$
(C.1)

Considering a time-dependent Hamiltonian H(t, q, p), since t must be updated in the same way as q, we immediately obtain the following *presymplectic Euler* method:

$$p_{\ell+1} = p_{\ell} - h \nabla_q H(t_{\ell}, q_{\ell}, p_{\ell+1}),$$

$$t_{\ell+1} = t_{\ell} + h,$$

$$q_{\ell+1} = q_{\ell} + h \nabla_p H(t_{\ell}, q_{\ell}, p_{\ell+1}).$$

(C.2)

Note how we have simply added an update for t, following the same structure as the update for q. Also, there is a dual version of (C.1) given by [22]:

$$q_{\ell+1} = q_{\ell} + h\nabla_p H(q_{\ell+1}, p_{\ell}),$$

$$p_{\ell+1} = p_{\ell} - h\nabla_q H(q_{\ell+1}, p_{\ell}).$$
(C.3)

From this we obtain the following alternative to (C.2):

$$t_{\ell+1} = t_{\ell} + h,$$

$$q_{\ell+1} = q_{\ell} + h \nabla_p H(t_{\ell+1}, q_{\ell+1}, p_{\ell}),$$

$$p_{\ell+1} = p_{\ell} - h \nabla_q H(t_{\ell+1}, q_{\ell+1}, p_{\ell}).$$

(C.4)

Both of these methods, namely (C.2) and (C.4), are of order r = 1. Note that for an arbitrary Hamiltonian H in general they are implicit; i.e., nonlinear equations have to be solved to obtain either $q_{\ell+1}$ or $p_{\ell+1}$. However, when the Hamiltonian is separable in the form

$$H = T(t, p) + V(t, q), \tag{C.5}$$

these methods become completely explicit in all variables resulting in cheap implementations.

Presymplectic leapfrog One of the versions of the leapfrog method for a conservative Hamiltonian system is the following second-order method [22]:

$$p_{\ell+1/2} = p_{\ell} - (h/2)\nabla_q H(q_{\ell}, p_{\ell+1/2}),$$

$$q_{\ell+1} = q_{\ell} + (h/2)\left(\nabla_p H(q_{\ell}, p_{\ell+1/2}) + \nabla_p H(q_{\ell+1}, p_{\ell+1/2})\right),$$

$$p_{\ell+1} = p_{\ell+1/2} - (h/2)\nabla_q H(q_{\ell+1}, p_{\ell+1/2}).$$
(C.6)

For a time-dependent Hamiltonian H(t, q, p) we include appropriate updates for time and obtain the following *presymplectic leapfrog* method:

$$p_{\ell+1/2} = p_{\ell} - (h/2)\nabla_q H(t_{\ell}, q_{\ell}, p_{\ell+1/2}),$$

$$t_{\ell+1} = t_{\ell} + h,$$

$$q_{\ell+1} = q_{\ell} + (h/2)\left(\nabla_p H(t_{\ell}, q_{\ell}, p_{\ell+1/2}) + \nabla_p H(t_{\ell+1}, q_{\ell+1}, p_{\ell+1/2})\right),$$

$$p_{\ell+1} = p_{\ell+1/2} - (h/2)\nabla_q H(t_{\ell+1}, q_{\ell+1}, p_{\ell+1/2}).$$

(C.7)

There is another dual version of the leapfrog method (C.6), which for conservative systems is given by [22]:

$$q_{\ell+1/2} = q_{\ell} + (h/2)\nabla_p H(q_{\ell+1/2}, p_{\ell}),$$

$$p_{\ell+1} = p_{\ell} - (h/2) \left(\nabla_q H(q_{\ell+1/2}, p_{\ell}) + \nabla_q H(q_{\ell+1/2}, p_{\ell+1}) \right),$$

$$q_{\ell+1} = q_{\ell+1/2} + (h/2)\nabla_p H(q_{\ell+1/2}, p_{\ell+1}).$$

(C.8)

This leads to an alternative version of the presymplectic leapfrog, given by:

$$t_{\ell+1/2} = t_{\ell} + h/2,$$

$$q_{\ell+1/2} = q_{\ell} + (h/2)\nabla_p H(t_{\ell+1/2}, q_{\ell+1/2}, p_{\ell}),$$

$$p_{\ell+1} = p_{\ell} - (h/2) \left(\nabla_q H(t_{\ell+1/2}, q_{\ell+1/2}, p_{\ell}) + \nabla_q H(t_{\ell+1/2}, q_{\ell+1/2}, p_{\ell+1})\right), \quad (C.9)$$

$$t_{\ell+1} = t_{\ell+1/2} + h/2,$$

$$q_{\ell+1} = q_{\ell+1/2} + (h/2)\nabla_p H(t_{\ell+1/2}, q_{\ell+1/2}, p_{\ell+1}).$$

Again, in general these methods are implicit, but for a separable Hamiltonian (C.5) they become completely explicit in all variables. Moreover, only one gradient computation per iteration is necessary, even though these methods are of order r = 2.

Higher-order methods [41] and [42] have presented an elegant general approach to construct arbitrarily higher-order symplectic integrators. It assumes we are given a base method ϕ_h of order 2r ($r \ge 1$). An integrator of order 2r + 2 is then obtained by the composition:

$$\phi_{\tau_0 h} \circ \phi_{\tau_1 h} \circ \phi_{\tau_0 h}, \qquad \tau_0 = \frac{1}{2 - \frac{2r + 1}{\sqrt{2}}}, \qquad \tau_1 = -\frac{\frac{2r + 1}{\sqrt{2}}}{2 - \frac{2r + 1}{\sqrt{2}}}.$$
 (C.10)

One can start with any base method of choice, such as the leapfrog. From this new integrator of order 2r + 2 one may proceeds recursively to construct even higher-order methods. It is easy to adapt this procedure to presymplectic integrators by carefully adding an update for time t. However, the number of gradient computations of the Suzuki-Yoshida approach grows very fast with increasing order, thus this approach quickly becomes unfeasible. Moreover, the truncation error of such methods tend to be rather large, although the fourth-order method obtained from the leapfrog is competitive and interesting. (See [24] and especially [43] for an interesting discussion.)

References

References

 W. Su, S. Boyd, and E. J. Candès, "A differential equation for modeling Nesterov's accelerated gradient method: Theory and insights," *Journal of Machine Learning Research* 17 no. 153, (2016) 1–43.

- [2] A. Wibisono, A. C. Wilson, and M. I. Jordan, "A variational perspective on accelerated methods in optimization," *Proceedings of the National Academy of Sciences* 113 no. 47, (2016) E7351–E7358.
- [3] G. França, D. P. Robinson, and R. Vidal, "A dynamical systems perspective on nonsmooth constrained optimization." arXiv:1808.04048 [math.OC], 2018.
- [4] G. França, D. P. Robinson, and R. Vidal, "ADMM and accelerated ADMM as continuous dynamical systems," in *International Conference on Machine Learning*. 2018. arXiv:1805.06579 [math.OC].
- [5] G. França, J. Sulam, D. P. Robinson, and R. Vidal, "Conformal symplectic and relativistic optimization." arXiv:1903.04100 [math.OC], 2019.
- [6] W. Krichene, A. Bayen, and P. L. Bartlett, "Accelerated mirror descent in continuous and discrete time," Advances in Neural Information Processing Systems 28 (2015) 2845–2853.
- [7] A. C. Wilson, B. Recht, and M. I. Jordan, "A Lyapunov analysis of momentum methods in optimization." arXiv:1611.02635v3 [math.OC], 2016.
- [8] D. Scieur, V. Roulet, F. Bach, and A. d'Aspremont, "Integration methods and optimization algorithms," in Advances in Neural Information Processing Systems 30. 2017.
- [9] M. Betancourt, M. I. Jordan, and A. C. Wilson, "On symplectic optimization." arXiv:1802.03653 [stat.CO], 2018.
- [10] J. Zhang, A. Mokhtari, S. Sra, and A. Jadbabaie, "Direct Runge-Kutta discretization achieves acceleration," in Advances in Neural Information Processing Systems 31. 2018.
- [11] B. Shi, S. S. Du, W. J. Su, and M. I. Jordan, "Acceleration via symplectic discretization of high-resolution differential equations," in Advances in Neural Information Processing Systems. 2019.
- [12] M. Muehlebach and M. I. Jordan, "A dynamical systems perspective on Nesterov acceleration," in *International Conference of Machine Learning*. 2019.
- [13] J. Diakonikolas and M. I. Jordan, "Generalized momentum-based methods: A Hamiltonian perspective." arXiv:1906.00436 [math.OC], 2019.
- [14] Y. Nesterov, "A method of solving a convex programming problem with convergence rate $O(1/k^2)$," Soviet Mathematics Doklady 27 no. 2, (1983) 372–376.
- [15] B. T. Polyak, "Gradient methods for minimizing functionals," Zh. Vychisl. Mat. Mat. Fiz. 3 no. 4, (1963) 643–653.

- [16] R. Berndt, An Introduction to Symplectic Geometry. American Mathematical Society, 2001.
- [17] B. Aebischer, M. Borer, M. Kalin, and C. Leuenberger, Symplectic Geometry. Springer Basel AG, 1994.
- [18] G. Benettin and A. Giorgilli, "On the Hamiltonian interpolation of near-to-the-identity symplectic mappings with application to symplectic integration algorithms," J. Stat. Phys. 74 (1994) 1117–1143.
- [19] S. Reich, "Backward error analysis for numerical integrators," SIAM J. Numer. Anal. 36 no. 5, (1999) 1549–1570.
- [20] E. Hairer, "Backward analysis of numerical integrators and symplectic methods," Ann. Numer. Math. 1 (1994) 107–132.
- [21] B. Leimkuhler and S. Reich, *Simulating Hamiltonian Dynamics*. Cambridge University Press, 2004.
- [22] E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration*. Springer, 2006.
- [23] J. M. Sanz-Serna, "Symplectic integrators for Hamiltonian problems: An overview," Acta Numerica 1 (1992) 243–286.
- [24] R. I. McLachlan and G. R. W. Quispel, "Geometric integrators for ODEs," J. Phys. A: Math. Gen. 39 (2006) 5251–5285.
- [25] A. Iserles and G. R. W. Quispel, "Why geometric numerical integration?," in *Discrete Mechanics, Geometric Integration and Lie-Butcher Series*, pp. 1–28. Springer International Publishing, 2018.
- [26] A. Bhatt, D. Floyd, and B. E. Moore, "Second order conformal symplectic schemes for damped Hamiltonian systems," *Journal of Scientific Computing* 66 (2016) 1234–1259.
- [27] B. E. Moore, "Multi-conformal-symplectic PDEs and discretizations," Journal of Computational and Applied Mathematics 323 (2017) 1–15.
- [28] A. Bhatt and B. E. Moore, "Exponential integrators preserving local conservation laws of PDEs with time-dependent damping/driving forces," *Journal of Computational and Applied Mathematics* 352 (2019) 341–351.
- [29] X. Shang and H. C. Ottinger, "Structure-preserving integrators for dissipative systems based on reversible-irreversible splitting," Proc. R. Soc. A 476 (2020) 20190446.
- [30] T. Nussle, P. Thibaudeau, and S. Nicolis, "Probing magneto-elastic phenomena through an effective spin-bath coupling model," *Eur. Phys. J. B* 92 no. 29, (2019) 1–10.

- [31] T. Nussle, P. Thibaudeau, and S. Nicolis, "Dynamic magnetostriction for antiferromagnets," *Phys. Rev. B* 100 (2019) 214428.
- [32] R. McLachlan and M. Perlmutter, "Conformal Hamiltonian systems," Journal of Geometry and Physics 39 (2001) 276–300.
- [33] P. Caldirola, "Forze non conservative nella meccanica quantistica," Nuovo Cim. 18 (1941) 393–400.
- [34] E. Kanai, "On the quantization of the dissipative systems," Prog. Theor. Phys. 3 (1948) 440–442.
- [35] A. O. Caldeira and A. J. Leggett, "Influence of Dissipation on Quantum Tunneling in Macroscopic Systems," *Phys. Rev. Lett.* 46 no. 4, (1981) 211–214.
- [36] E. Hairer and C. Lubich, "The life-span of backward error analysis for numerical integrators," Numer. Math. 76 (1997) 441–462.
- [37] S. Sternberg, Lectures on Differential Geometry. Prentice Hall, 1964.
- [38] M. Tao, "Explicit symplectic approximation of nonseparable Hamiltonians: algorithm and long time performance," *Phys. Rev. E* **94** (2016) 043303.
- [39] V. I. Arnold, V. V. Kozlov, and A. I. Neishtadt, Mathematical Aspects of Classical and Celestial Mechanics. Springer-Verlag Berlin Heidelberg, 2006.
- [40] R. Abraham and J. E. Marsden, Foundations of Mechanics. Addison-Wesley Publishing Company, Inc., 1978.
- [41] M. Suzuki, "Fractal decomposition of exponential operators with applications to many-body theories and Monte Carlo simulations," *Phys. Lett. A* 146 (1990) 319–323.
- [42] H. Yoshida, "Construction of higher order symplectic integrators," Phys. Lett. A 150 no. 5, (1990) 262–268.
- [43] R. I. McLachlan, "Families of high-order composition methods," Numerical Algorithms 31 (2002) 233–246.