

SEMISIMPLIFICATION FOR SUBGROUPS OF REDUCTIVE ALGEBRAIC GROUPS

MICHAEL BATE, BENJAMIN MARTIN, AND GERHARD RÖHRLE

ABSTRACT. Let G be a reductive algebraic group—possibly non-connected—over a field k and let H be a subgroup of G . If $G = \mathrm{GL}_n$ then there is a degeneration process for obtaining from H a completely reducible subgroup H' of G ; one takes a limit of H along a cocharacter of G in an appropriate sense. We generalise this idea to arbitrary reductive G using the notion of G -complete reducibility and results from geometric invariant theory over non-algebraically closed fields due to the authors and Herpel. Our construction produces a G -completely reducible subgroup H' of G , unique up to $G(k)$ -conjugacy, which we call a k -*semisimplification* of H . This gives a single unifying construction which extends various special cases in the literature (in particular, it agrees with the usual notion for $G = \mathrm{GL}_n$ and with Serre’s “ G -analogue” of semisimplification for subgroups of $G(k)$ from [21]). We also show that under some extra hypotheses, one can pick H' in a more canonical way using the Tits Centre Conjecture for spherical buildings and/or the theory of optimal destabilising cocharacters introduced by Hesselink, Kempf and Rousseau.

1. INTRODUCTION

The aim of this paper is to present a construction of the *semisimplification* of a subgroup H of a reductive linear algebraic group G over an arbitrary field k . This construction unifies and generalizes many concepts already in the literature within a single framework. For example, the semisimplification of a module for a group is a well-known construction in representation theory, corresponding in our case to the situation where $H \subseteq \mathrm{GL}_n(k)$. Building on this idea, for G a connected reductive linear algebraic group over a field k and H a subgroup of $G(k)$, Serre already introduced the concept of a “ G -analogue” of semisimplification from representation theory in [21, §3.2.4]. This notion is also used for representations of various kinds of algebras, e.g., see [14], [8], [18], [24], and [25]. It is also an ingredient in work of Lawrence-Sawin on the Shafarevich Conjecture for abelian varieties, [15] which involves Galois representations taking values in non-connected reductive p -adic groups.

We begin by recalling how the most basic case works. Let $n \in \mathbb{N}$ and let H be a subgroup of $\mathrm{GL}_n(k)$. There is an H -module filtration of k^n such that the successive quotients are irreducible, by the Jordan-Hölder Theorem. In terms of matrices, this implies that, by changing basis if necessary, we may assume that H is in upper block-triangular form, with the action of H on each quotient being represented by the corresponding block on the diagonal. Letting H' be the subgroup of $\mathrm{GL}_n(k)$ consisting of the block diagonal matrices obtained by taking each element of H and replacing the entries above the block diagonal with 0s, we obtain a subgroup which acts semisimply on k^n —that is, H' is completely reducible. Since

2010 *Mathematics Subject Classification.* 20G15 (14L24).

Key words and phrases. Semisimplification, G -complete reducibility, geometric invariant theory, rationality, cocharacter-closed orbits, degeneration of G -orbits.

this construction is independent of the choice of basis up to $\mathrm{GL}_n(k)$ -conjugacy, again by the Jordan-Hölder Theorem, it is therefore reasonable to call H' the *semisimplification* of H .

We now explain some of the ingredients of our construction in the case that k is algebraically closed, which removes some technicalities. Recall [2], [21] that if G is connected and H is a subgroup of G then H is *G -completely reducible* (G -cr for short) if for any parabolic subgroup P of G such that P contains H , there is a Levi subgroup L of P such that L contains H . If $G = \mathrm{GL}_n$ then H is G -cr if and only if k^n is completely reducible as an H -module; this follows from the usual characterisation of parabolic subgroups of GL_n as stabilizers of flags of subspaces. We make the same definition for arbitrary reductive G , replacing parabolic subgroups and Levi subgroups with R-parabolic subgroups and R-Levi subgroups instead (see Section 2 for details).

To perform our construction, we apply a characterisation of G -complete reducibility in terms of geometric invariant theory (GIT). We see this idea already in our original example: we can view H' as a degeneration of H in the following sense. Let the sizes of the blocks down the diagonal be n_1, \dots, n_r , and define a cocharacter $\lambda: \mathbb{G}_m \rightarrow \mathrm{GL}_n$ by

$$\lambda(a) = \mathrm{diag}(a^r, \dots, a^r, \dots, a^1, \dots, a^1), \text{ with } n_i \text{ occurrences of } a^{r-i+1}, 1 \leq i \leq r.$$

For each $a \in k^*$, define $H_a = \lambda(a)H\lambda(a)^{-1}$ for $a \in k^*$. Then $H' = \lim_{a \rightarrow 0} H_a$ in an appropriate sense.

Our definition of k -semisimplification (Definition 4.1) is new, generalizes the one given by Serre in [21, §3.2.4], and is closely related to the definition given in [6] using optimal destabilising cocharacters; the two notions agree whenever the latter makes sense (cf. also [16, Sec. 4] for the algebraically closed case). We prove that the k -semisimplification of a subgroup H of G is unique up to conjugacy (Theorem 4.5), generalizing [21, Prop. 3.3(b)]. We give a short and self-contained exposition, bringing together some results (such as Corollary 3.6) that follow from previous work but are not easily extracted from earlier papers.

2. COCHARACTER-CLOSED ORBITS

Following [7] and our earlier work [6], [1], we regard an affine variety over a field k as a variety X over the algebraic closure \bar{k} together with a choice of k -structure. We denote the separable closure of k by k_s . We write $X(k)$ for the set of k -points of X and $X(\bar{k})$ (or just X) for the set of \bar{k} -points of X . By a subvariety of X we mean a closed \bar{k} -subvariety of X ; a k -subvariety is a subvariety that is defined over k . We denote by M_n the associative algebra of $n \times n$ matrices over k . Below G denotes a possibly non-connected reductive linear algebraic group over k . By a subgroup of G we mean a closed \bar{k} -subgroup and by a k -subgroup we mean a subgroup that is defined over k . We discuss non-closed subgroups in Remark 4.12(a). By G^0 we denote the identity component of G , and likewise for subgroups of G .

We define $Y_k(G)$ to be the set of k -defined cocharacters of G and $Y(G) := Y_{\bar{k}}(G)$ to be the set of all cocharacters of G .

Let H be a subgroup of G . Even if H is k -defined, the (set-theoretic) centralizer $C_G(H)$ need not be k -defined in general. It is useful to have criteria to ensure that $C_G(H)$ is k -defined (see Proposition 3.5 and Section 5). For instance, if k is perfect and H is k -defined then $C_G(H)$ is k -defined. We say that H is *separable* if the scheme-theoretic centralizer $\mathcal{C}_G(H)$ is smooth [2, Def. 3.27]; for instance, any subgroup of GL_n is separable [2, Ex. 3.28]

(see [5] for more examples of separable subgroups). If H is k -defined and separable then $C_G(H)$ is k -defined (see [1, Prop. 7.4]).

Next we recall some basic notation and facts concerning parabolic subgroups in (non-connected) reductive groups G from [2, §6] and [6]. Given $\lambda \in Y(G)$, we define

$$P_\lambda = \{g \in G \mid \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1} \text{ exists}\}$$

and $L_\lambda = C_G(\text{Im}(\lambda))$ (for the definition of a limit, see [22, Sec. 3.2.13]). We call P_λ an *R-parabolic subgroup* of G and L_λ an *R-Levi subgroup* of P_λ ; they are subgroups of G . We have $P_\lambda = L_\lambda = G$ if $\text{Im}(\lambda)$ belongs to the centre of G . For ease of reference, we record without proof some basic facts about these subgroups.

Lemma 2.1. (i) *If P is a k -defined R-parabolic subgroup then $R_u(P)$ is k -defined.*
(ii) *If P is a parabolic subgroup of G^0 then the normalizer $N_G(P)$ is an R-parabolic subgroup of G , and $N_G(P)$ is k -defined if P is.*

If G is connected then every pair (P, L) consisting of a parabolic k -subgroup P of G and a Levi k -subgroup L of P is of the form $(P, L) = (P_\lambda, L_\lambda)$ for some $\lambda \in Y_k(G)$, and vice versa [22, Lem. 15.1.2(ii)]. In general, if $\lambda \in Y_k(G)$ then P_λ and L_λ are k -defined [6, Lem. 2.5], but the converse is not so straightforward. If P is an R-parabolic k -subgroup and L is an R-Levi k -subgroup of P then for any maximal k -torus T of L , there exists $\lambda \in Y_{k_s}(T)$ such that $P = P_\lambda$ and $L = L_\lambda$. However, it is possible that P is a k -defined R-parabolic subgroup and yet there does not exist any $\mu \in Y_k(G)$ such that $P = P_\mu$, and similarly for R-Levi subgroups — see [6, Rem. 2.4]. This complicates some of the arguments below.

Lemma 2.2. *Let P be an R-parabolic subgroup of G and L an R-Levi subgroup of P .*

- (i) *We have $P \cong L \rtimes R_u(P)$, and this is a k -isomorphism if P and L are k -defined.*
- (ii) *Any two R-Levi k -subgroups of an R-parabolic k -subgroup P are $R_u(P)(k)$ -conjugate.*

We denote the canonical projection from P to L by c_L ; this is k -defined if P and L are. If we are given $\lambda \in Y(G)$ such that $P = P_\lambda$ and $L = L_\lambda$ then we often write c_λ instead of c_L . We have $c_\lambda(g) = \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1}$ for $g \in P_\lambda$; the kernel of c_λ is the unipotent radical $R_u(P_\lambda)$ and the set of fixed points of c_λ is L_λ .

Let $n \in \mathbb{N}$. Below we consider the action of G on G^n by simultaneous conjugation: $g \cdot (g_1, \dots, g_n) = (gg_1g^{-1}, \dots, gg_ng^{-1})$. Given $\lambda \in Y(G)$, we have a map $P_\lambda^n \rightarrow L_\lambda^n$ given by $\mathbf{g} \mapsto \lim_{a \rightarrow 0} \lambda(a) \cdot \mathbf{g}$; we abuse notation slightly and also call this map c_λ . For any $\mathbf{g} \in P_\lambda^n$, there exists an R-Levi k -subgroup L of P_λ such that $\mathbf{g} \in L^n$ if and only if $c_\lambda(\mathbf{g}) = u \cdot \mathbf{g}$ for some $u \in R_u(P_\lambda)(k)$.

Our main tool from GIT is the notion of cocharacter-closure, introduced in [6] and [1].

Definition 2.3. Let X be an affine G -variety and let $x \in X$ (we do not require x to be a k -point). We say that the orbit $G(k) \cdot x$ is *cocharacter-closed over k* if for all $\lambda \in Y_k(G)$ such that $x' := \lim_{a \rightarrow 0} \lambda(a) \cdot x$ exists, x' belongs to $G(k) \cdot x$. If $k = \bar{k}$ then it follows from the Hilbert-Mumford Theorem that $G(k) \cdot x$ is cocharacter-closed over k if and only if $G(k) \cdot x$ is closed [13, Thm. 1.4]. If \mathcal{O} is a $G(k)$ -orbit in X then we say that \mathcal{O} is *accessible from x over k* if there exists $\lambda \in Y_k(G)$ such that $x' := \lim_{a \rightarrow 0} \lambda(a) \cdot x$ belongs to \mathcal{O} .

The following result is [1, Thm. 1.3].

Theorem 2.4 (Rational Hilbert-Mumford Theorem). *Let G , X , x be as above. Then there is a unique $G(k)$ -orbit \mathcal{O} such that \mathcal{O} is cocharacter-closed over k and accessible from x over k .*

3. G -COMPLETE REDUCIBILITY

Definition 3.1. Let H be a subgroup of G . We say that H is *G -completely reducible over k* (G -cr over k) if for any R-parabolic k -subgroup P of G such that P contains H , there is an R-Levi k -subgroup L of P such that L contains H . We say that H is *G -irreducible over k* (G -ir over k) if H is not contained in any proper R-parabolic k -subgroup of G at all.

Remark 3.2. We say that H is G -cr if H is G -cr over \bar{k} —cf. Section 1. More generally, if k'/k is an algebraic field extension then we may regard G as a k' -group and it makes sense to ask whether H is G -cr over k' .

For more on G -complete reducibility, see [20], [21], [2].

Note that the definitions make sense even if H is not k -defined. It is immediate that G -irreducibility over k implies G -complete reducibility over k . We have $P_{g\lambda} = gP_\lambda g^{-1}$ and $L_{g\lambda} = gL_\lambda g^{-1}$ for any $\lambda \in Y(G)$ and any $g \in G$ (see, e.g., [2, §6]). It follows that if H is G -cr over k (resp., G -ir over k) then so is any $G(k)$ -conjugate of H . More generally, one can show that if H is G -cr over k (resp., G -ir over k) then so is $\phi(H)$, for any k -defined automorphism ϕ of G . If $k = \bar{k}$ and H is G -cr then H is reductive [21, Prop. 4.1], [2, §2.4, §6.2]. It follows from Proposition 3.5 below that if H is k -defined, k is perfect and H is G -cr over k then H is reductive. We see below (Corollary 3.6) that the converse holds in characteristic 0. On the other hand, the converse is false in general.

Example 3.3. (See [23, Proof of Prop. 1.10].) Suppose k is imperfect and $\text{char}(k) = 2$. Let $G = \text{PGL}_2$. Choose $a \in k \setminus k^2$ and set

$$H = \left\{ \overline{\begin{pmatrix} x & ay \\ y & x \end{pmatrix}} \mid x, y \in \bar{k}, x^2 - ay^2 \neq 0 \right\},$$

where $A \mapsto \bar{A}$ denotes the canonical projection from GL_2 to PGL_2 . Then $H = C_G(H)$ is k -defined, H is a connected unipotent group—in particular, H is not G -cr—and H is G -cr over k . Note that H is a wound unipotent group (cf. [10, Ex. B.2.3]).

We now explain the link between G -complete reducibility and GIT. Fix a k -embedding $\iota : G \rightarrow \text{GL}_n$ for some $n \in \mathbb{N}$. Let H be a subgroup of G . Let $m \in \mathbb{N}$ and let $\mathbf{h} = (h_1, \dots, h_m) \in H^m$. We call \mathbf{h} a *generic tuple for H with respect to ι* if h_1, \dots, h_m generates the subalgebra of M_n generated by H [6, Def. 5.4]. Note that we don't insist that \mathbf{h} is a k -point. Our constructions below do not depend on the choice of ι , so we suppress the words “with respect to ι ”. It is immediate that if $\mathbf{h} \in H^m$ is a generic tuple for H and $g \in G$ then $g \cdot \mathbf{h}$ is a generic tuple for gHg^{-1} .

Theorem 3.4 ([1, Thm. 9.3]). *Let H be a subgroup of G and let $\mathbf{h} \in H^m$ be a generic tuple for H . Then H is G -completely reducible over k if and only if $G(k) \cdot \mathbf{h}$ is cocharacter-closed over k .*

Using this result one can derive many results on G -complete reducibility: for instance, see [2] for the algebraically closed case and [6], [1] for arbitrary k . Note that if $\mathbf{h} \in H^m$ is a generic tuple for H then the centralizer $C_G(H)$ coincides with the stabilizer $G_{\mathbf{h}}$.

Proposition 3.5. *Let H be a k -subgroup of G . Suppose k is perfect. Then H is G -completely reducible over k if and only if H is G -completely reducible.*

Proof. If k is perfect then \bar{k}/k is separable and $C_G(H)$ is k -defined. The result now follows from [1, Cor. 9.7(i)]. \square

Corollary 3.6. *Suppose $\text{char}(k) = 0$. Let H be a k -subgroup of G . Then H is G -completely reducible over k if and only if H is reductive.*

Proof. If $k = \bar{k}$ then this is well known (see [21, Prop. 4.2], [2, §2.2, §6.3], for example). The result for arbitrary k now follows from Proposition 3.5. \square

Proposition 3.7. *Let H be a subgroup of G . If $C_G(H)$ is k -defined and H is G -completely reducible then H is G -completely reducible over k .*

Proof. This follows from Remark 5.2 below. \square

The next result strengthens [1, Cor. 9.7(ii)]: we do not need the hypotheses that H and $C_G(H)$ are k -defined. See also [21, Prop. 3.2].

Proposition 3.8. *Let $S \subseteq G$ be a k -split torus and let $L = C_G(S)$. Let H be a subgroup of L . Then H is G -completely reducible over k if and only if H is L -completely reducible over k .*

Proof. This follows from Theorem 3.4 together with [1, Thm. 5.4(ii)]. \square

Corollary 3.9. *Let L be an R-Levi k -subgroup of G and let H be a subgroup of L . Then H is G -completely reducible over k if and only if H is L -completely reducible over k .*

Proof. We can choose $\lambda \in Y_{k_s}(G)$ such that $L = C_G(\text{Im}(\lambda))$. Let $\lambda = \lambda_1, \lambda_2, \dots, \lambda_r \in Y_{k_s}(G)$ be the $\text{Gal}(k_s/k)$ -conjugates of λ and let S be the subtorus of $Z(L)^0$ generated by the subtori $\text{Im}(\lambda_i)$. Then S is k -defined and $L = C_G(S)$. The product map $\lambda_1 \times \dots \times \lambda_r$ gives an epimorphism from $\bar{k}^* \times \dots \times \bar{k}^*$ onto S . But a quotient of a split k -torus is k -split [7, III.8.4 Cor.], so S is split. The result now follows from Proposition 3.8. \square

We finish the section with some results involving non-connected reductive groups. Note that if Q is an R-parabolic k -subgroup of G and M is an R-Levi k -subgroup of Q then Q^0 is a parabolic k -subgroup of G^0 and M^0 is a Levi k -subgroup of Q^0 ; see [2, Sec. 6].

Lemma 3.10. *Let P be an R-parabolic subgroup of G and let T be a maximal torus of P . Then there is a unique R-Levi subgroup L of P such that $T \subseteq L$. If P and T are k -defined then L is k -defined.*

Proof. The first assertion is [2, Cor. 6.5]. For the second, suppose P and T are k -defined. There exists an R-Levi k -subgroup L' of P . Now L' contains a maximal k -torus T' , and T' is also a maximal k -torus of P . So $gT'g^{-1} = T$ for some $g \in P(k_s)$. Let $L = gL'g^{-1}$, an R-Levi k_s -subgroup of P . Then L is the unique R-Levi subgroup of P containing T . This implies that L is $\text{Gal}(k_s/k)$ -stable, so L is k -defined, as required. \square

Lemma 3.11. (a) *Let Q be an R-parabolic k -subgroup of G and set $P = Q^0$. Then the R-Levi k -subgroups of Q are precisely the subgroups of the form $N_Q(L)$ for L a Levi k -subgroup of P .*

- (b) Let Q, P be as in (a) and let H be a subgroup of P . Then H is contained in an R-Levi k -subgroup of Q if and only if H is contained in a Levi k -subgroup of P . Moreover, if L is a Levi k -subgroup of P then $c_{N_Q(L)}(H)$ is $N_Q(L)$ -completely reducible over k if and only if $c_L(H)$ is L -completely reducible over k .
- (c) Let H be a subgroup of G^0 . Then H is G -completely reducible over k if and only if H is G^0 -completely reducible over k .

Proof. (a) As observed above, if M is an R-Levi subgroup of Q then M^0 is a Levi subgroup of P , and $N_Q(M^0)^0 = N_P(M^0)^0 = M^0$. Let L be a Levi subgroup of P and let T be a maximal torus of L . By Lemma 3.10 there is a unique R-Levi subgroup M of Q such that $T \subseteq M$. The Levi subgroups M^0 and L of P both contain T , so by Lemma 3.10 they are equal; in particular, M normalizes L . Now $N_Q(T)$ normalizes L by Lemma 3.10, so $N_Q(L)$ meets every component of Q . Since $Q = M \ltimes R_u(Q)$, M also meets every component of Q . It follows that $M = N_Q(L)$. Finally, L contains a maximal k -torus of P if and only if $N_Q(L)$ does, so L is k -defined if and only if $N_Q(L)$ is, by Lemma 3.10.

(b) The first assertion follows immediately from (a), and part (c) now follows. For the second assertion of (b), note that the restriction of $c_{N_Q(L)}(H)$ to P is c_L ; the desired result now follows from part (c) applied to the reductive k -group $N_Q(L)$. \square

4. k -SEMISIMPLIFICATION

Now we come to our main definition.

Definition 4.1. Let H be a subgroup of G . We say that a subgroup H' of G is a k -semisimplification of H (for G) if there exist an R-parabolic k -subgroup P of G and an R-Levi k -subgroup L of P such that $H \subseteq P$, $H' = c_L(H)$ and H' is G -completely reducible (or equivalently by Corollary 3.9, L -completely reducible) over k . We say the pair (P, L) yields H' .

- Remarks 4.2.* (a) Let H be a subgroup of G . If H is G -cr over k then clearly H is a k -semisimplification of itself, yielded by the pair (G, G) .
- (b) Suppose (P, L) yields a k -semisimplification H' of H . Let L_1 be another R-Levi k -subgroup of P . Then $L_1 = uLu^{-1}$ for some $u \in R_u(P)(k)$, so $c_{L_1}(H) = uc_L(H)u^{-1}$. Hence (P, L_1) also yields a k -semisimplification of H . We say that P yields a k -semisimplification of H .
- (c) It is straightforward to check that if ϕ is an automorphism of G (as a k -group), H is a subgroup of G and (P, L) yields a k -semisimplification H' of H then $\phi(H')$ is a k -simplification of $\phi(H)$, yielded by $(\phi(P), \phi(L))$.
- (d) For G connected and H a subgroup of $G(k)$, Definition 4.1 generalizes Serre's " G -analogue" of a semisimplification from [21, §3.2.4]. For $k = \bar{k}$, Definition 4.1 generalizes the definition of $\mathcal{D}(H)$ following [16, Lem. 4.1].

Remark 4.3. Let $\mathbf{h} \in H^m$ be a generic tuple for H . Note that c_λ extends in the obvious way to a homomorphism from a parabolic subalgebra \mathcal{P}_λ of M_n onto a Levi subalgebra \mathcal{L}_λ of \mathcal{P}_λ , and \mathcal{P}_λ contains the subalgebra generated by H . It is easily checked that $c_\lambda(\mathbf{h})$ is a generic tuple for $c_\lambda(H)$. Hence by Theorem 3.4, $c_\lambda(H)$ is a k -semisimplification of H if and only if $G(k) \cdot c_\lambda(\mathbf{h})$ is cocharacter-closed over k . It follows from Theorem 2.4 that H admits at least one k -semisimplification: for we can choose $\lambda \in Y_k(G)$ such that $G(k) \cdot c_\lambda(\mathbf{h})$ is cocharacter-closed over k , so $c_\lambda(H)$ is a k -semisimplification of H , yielded by (P_λ, L_λ) .

Remark 4.4. Suppose H' is a k -semisimplification of H , yielded by a pair (P, L) . By the discussion in Section 2, there exists a maximal k -torus T of L and $\lambda \in Y_{k_s}(T)$ such that $P = P_\lambda$ and $L = L_\lambda$. Choose a finite Galois extension k'/k such that T splits over k' . Let $\Gamma = \text{Gal}(k'/k)$ and let $\mu = \sum_{\gamma \in \Gamma} \gamma \cdot \lambda \in Y_k(T)$. One checks easily that $H \subseteq P_\mu$ and $c_\mu|_H = c_\lambda|_H$ (cf. the proof of [6, Lem. 2.5(ii)]). Hence (P_μ, L_μ) also yields H' .

Here is our main result, which was proved in the special case $k = \bar{k}$ in [6, Prop. 5.14(i)], cf. [21, Prop. 3.3(b)].

Theorem 4.5. *Let H be a subgroup of G . Then any two k -semisimplifications of H are $G(k)$ -conjugate.*

Proof. Let H_1, H_2 be k -semisimplifications of H . By Remark 4.4, there exist $\lambda_1, \lambda_2 \in Y_k(G)$ such that $(P_{\lambda_1}, L_{\lambda_1})$ realizes H_1 and $(P_{\lambda_2}, L_{\lambda_2})$ realizes H_2 . Let $\mathbf{h} \in H^m$ be a generic tuple for H . Then $c_{\lambda_i}(\mathbf{h})$ is a generic tuple for H_i for $i = 1, 2$, and each orbit $G(k) \cdot c_{\lambda_i}(\mathbf{h})$ is cocharacter-closed over k . It follows from the uniqueness result in Theorem 3.4 that the closed subset $C_{\mathbf{h}} := \{g \in G \mid g \cdot c_{\lambda_1}(\mathbf{h}) = c_{\lambda_2}(\mathbf{h})\}$ is non-empty.

Pick $g \in C_{\mathbf{h}}$. If $H_2 = gH_1g^{-1}$ then we are done. Otherwise there exists $h \in H$ such that $gc_{\lambda_1}(h)g^{-1} \notin H_2$ or $g^{-1}c_{\lambda_2}(h)g \notin H_1$. Without loss assume the former. We can repeat the above argument, replacing \mathbf{h} with the generic tuple $\mathbf{h}' := (\mathbf{h}, h) \in H^{m+1}$; note that $C_{\mathbf{h}'}$ is properly contained in $C_{\mathbf{h}}$. The result now follows by a descending chain condition argument. \square

The uniqueness asserted in Theorem 4.5 is akin to the theorem of Jordan–Hölder.

Definition 4.6. We define $\mathcal{D}_k(H)$ to be the set of $G(k)$ -conjugates of any k -semisimplification of H (cf. the discussion preceding [16, Thm. 1.4]). This is well-defined by Theorem 4.5.

Example 4.7. Let H be a subgroup of G . As noted in Remark 4.2(a), if H is G -cr over k then H is a k -semisimplification of itself, yielded by the pair (G, G) .

Note that if H is a G -ir subgroup of G then H is the only k -semisimplification of H : this shows that not every element of $\mathcal{D}_k(H)$ need be a k -semisimplification of H . Similarly, let P be a maximal proper R-parabolic k -subgroup of G and let L be an R-Levi k -subgroup of P . Then (P, L) yields a k -semisimplification L of P , and if L_1 is a $G(k)$ -conjugate of L then L_1 is a k -semisimplification of P if and only if $L_1 \subseteq P$. If P and Q are arbitrary R-parabolic k -subgroups of G and $Q \supseteq P$ then it is easily seen that Q yields a k -semisimplification of P if and only if $P^0 = Q^0$.

Example 4.8. Let H be a subgroup of G and let P be minimal among the R-parabolic k -subgroups that contain H . Let L be an R-Levi k -subgroup of P . We claim that $c_L(H)$ is L -ir over k (cf. [21, Prop. 3.3(a)] and [2, Sec. 3]); it then follows from Corollary 3.9 that $c_L(H)$ is a k -semisimplification of H . Suppose $c_L(H)$ is not L -ir: say, $c_L(H) \subseteq Q$, where Q is a proper R-parabolic k -subgroup of L . There exist a maximal k -torus T of Q and cocharacters $\lambda, \mu \in Y_{k_s}(T)$ such that $P = P_\lambda$, $L = L_\lambda$ and $Q = P_\mu$. Now $H \subseteq QR_u(P) \subsetneq P$, and clearly $QR_u(P)$ is k -defined. But it is easily checked that $QR_u(P) = P_{m\lambda+\mu}$ for suitably large $m \in \mathbb{N}$ (cf. [2, Lem. 6.2(i)]), so $QR_u(P)$ is an R-parabolic k -subgroup of G , contradicting the minimality of P . Conversely, if P is an R-parabolic k -subgroup with R-Levi k -subgroup L such that $P \supseteq H$ and $c_L(H)$ is L -ir over k then a similar argument shows that P is minimal among the R-parabolic k -subgroups containing H . This proves the claim.

In particular, let G , H , λ and H' be as in the GL_n example in Section 1. Let $P = P_\lambda$ be the parabolic subgroup of block upper triangular matrices with blocks of size n_1, \dots, n_r down the leading diagonal. Let $L = L_\lambda$ be the subgroup of block diagonal matrices with blocks of size n_1, \dots, n_r down the leading diagonal. Since each $n_i \times n_i$ block yields an irreducible representation of $H' := c_\lambda(H)$, H' is L -ir over k , so P is minimal among the R-parabolic k -subgroups of G containing H ; hence H' is the k -semisimplification of H yielded by (P, L) .

Example 4.9. Suppose $\mathrm{char}(k) = 0$. Let H be a k -subgroup of G and let P be an R-parabolic subgroup of G with R-Levi subgroup L such that $P \supseteq H$. Then Corollary 3.6 implies that $c_L(H)$ is a k -semisimplification of H if and only if $R_u(H) \subseteq R_u(P)$.

Example 4.10. Let H be a subgroup of G^0 and let H' be a k -semisimplification of H for G^0 yielded by a pair (P, L) . Pick any R-parabolic k -subgroup Q of G such that $Q^0 = P$; note there is at least one, namely $Q = N_G(P)$. Then Lemma 3.11 implies that $N_Q(L)$ is an R-Levi k -subgroup of Q and $H' = c_L(H) = c_{N_Q(L)}(H)$ is $N_Q(L)$ -cr over k . Hence H' is a k -semisimplification of H for G , yielded by the pair $(Q, N_Q(L))$.

Example 4.11. Given a reductive k -group G and a subgroup H of G , we may (as in Remark 3.2) regard G as a \bar{k} -group by forgetting the k -structure, so it makes sense to consider the semisimplification (i.e., the \bar{k} -semisimplification) of H . It can happen that H is G -cr over k but not G -cr, or vice versa: see Example 3.3, [2, Ex. 5.11] and [5, Ex. 7.22]. So there is no direct relation between the notions of k -semisimplification and semisimplification. More generally, if k'/k is an algebraic field extension then we may view G as a k' -group, and again there is no direct relation between the notions of k -semisimplification and k' -semisimplification. On the other hand, if H' is a k -semisimplification of H and H' is G -cr over k' then it is clear that H' is also a k' -semisimplification of H .

Remarks 4.12. (a) The notion of G -complete reducibility over k makes sense for a non-closed subgroup H of $G(\bar{k})$. An R-parabolic subgroup (resp., R-Levi subgroup) of G contains H if and only if it contains the Zariski-closure \overline{H} , so we see that H is G -cr over k if and only if \overline{H} is. Hence we can define a k -semisimplification of H to be a subgroup H' of the form $c_L(H)$, for any pair (P, L) that yields a k -semisimplification of \overline{H} . We obtain analogues of the results above with obvious modifications (we leave the details to the reader).

(b) Suppose now that H is a possibly non-smooth subgroup scheme of G . Again the notion of G -complete reducibility over k makes sense for H . The GIT results from [1] and [6] don't extend directly: one cannot capture all the properties of H just by looking at its group of \bar{k} -points $H(\bar{k})$. We plan to develop a more general theory in future work, where we allow G to be a smooth k -group such that G^0 is pseudo-reductive [10].

5. OPTIMALITY AND NORMAL SUBGROUPS

We have seen that in general not every element of $\mathcal{D}_k(H)$ is a k -semisimplification of H . On the other hand, it can happen that H is contained in many different R-parabolic subgroups of G , and there may exist many conjugate, but different, k -semisimplifications. We now recall two constructions that give under some extra hypotheses a more canonical

choice of R -parabolic subgroup yielding a k -semisimplification. They apply in particular when $G = \mathrm{GL}_n$ (see Example 5.6); this does not seem to be well known even when $k = \overline{k}$.

First construction: Suppose G is connected, H is a subgroup of G and H is not G -cr over k . We use the theory of spherical buildings (see [20], [21]) and the argument of [3, Proof of Thm. 1.1]. Recall that the spherical building $\Delta_k(G)$ of G is a simplicial complex whose simplices are the parabolic k -subgroups of G , ordered by reverse inclusion (the proper k -parabolic subgroups correspond to the non-empty simplices). The apartments of $\Delta_k(G)$ are the sets of all k -parabolic subgroups of G that contain a fixed maximal split k -torus S of G . The set Σ of parabolic k -subgroups P of G such that $P \supseteq H$ is a convex subcomplex of $\Delta_k(G)$, and Σ is not completely reducible in the sense of [21, §2.2] because H is not G -cr over k (see [21, §3.2.1]). By the Tits Centre Conjecture—see, e.g., [4, §2.6], and [21, §2.4] and the references therein— Σ has a so-called “centre”: a proper parabolic k -subgroup $P_c \in \Sigma$ such that P_c is fixed by any building automorphism of $\Delta_k(G)$ that stabilizes Σ . In particular, P_c is stabilized by any k -automorphism of G that stabilizes H .

Lemma 5.1. *Let G , H and Σ be as above. Let P_c be a centre for Σ such that P_c is not properly contained in any other centre for Σ . Then P_c yields a k -semisimplification of H .*

Proof. Let Λ be the set of k -parabolic subgroups Q of G such that $Q \subseteq P_c$. Fix a Levi k -subgroup L of P_c . We have an inclusion-preserving bijection ψ from Λ to $\Delta_k(L)$ given by $Q \mapsto Q \cap L$, with inverse given by $R \mapsto RR_u(P_c)$. Let Σ_L be the subset of $\Delta_k(L)$ consisting of all the k -parabolic subgroups of L that contain $c_L(H)$. It is clear that $\psi(\Sigma \cap \Lambda) = \Sigma_L$. If ϕ is a building automorphism of $\Delta_k(G)$ that fixes P_c then ϕ stabilizes Λ , and we get an automorphism ϕ_L of $\Delta_k(L)$ (as a simplicial complex) given by $\phi_L(Q \cap L) = \phi(Q) \cap L$; moreover, if ϕ stabilizes Σ then ϕ_L stabilizes Σ_L .

We claim that ϕ_L is a building automorphism of $\Delta_k(L)$. It is enough to show that ϕ_L maps apartments to apartments. Let S be a maximal split k -torus of L (and hence of G). Since ϕ is a building automorphism, there is a maximal split k -torus S' of G such that for every k -parabolic subgroup Q of G that contains S , $\phi(Q)$ contains S' . In particular, $S' \subseteq P_c$ since $\phi(P_c) = P_c$. By Lemma 3.10 there is a k -Levi subgroup L' of P_c such that $S' \subseteq L'$. By Lemma 2.2(ii) there exists $u \in R_u(P_c)(k)$ such that $uS'u^{-1} \subseteq L$. Let $R \in \Delta_k(L)$ such that $S \subseteq R$: say, $R = Q \cap L$ for $Q \in \Lambda$. Then $S' \subseteq \phi(Q)$. Since $\phi(Q) \subseteq P_c$, $R_u(\phi(Q))$ contains $R_u(P_c)$, so $uS'u^{-1} \subseteq \phi(Q)$. Hence $uS'u^{-1} \subseteq \phi(Q) \cap L = \phi_L(R)$. This proves the claim.

Now suppose P_c does not yield a k -semisimplification of H . Then $c_L(H)$ is not L -cr over k . By the discussion before the lemma, Σ_L has a centre $R \subsetneq L$. We have $R = Q \cap L$ for some $Q \in \Lambda$ with $Q \subsetneq P_c$. But the results in the previous paragraph imply that Q is a centre for Σ , contradicting the minimality of P_c . \square

More generally, allow G to be non-connected and suppose H is a subgroup of G^0 such that H is not G -cr over k . Then H is not G^0 -cr over k by Lemma 3.11, so P_c chosen as in Lemma 5.1 yields a k -semisimplification of H for G^0 . Hence $N_G(P_c)$ yields a k -semisimplification of H for G by Example 4.10, and $N_G(P_c)$ is stabilized by any k -automorphism of G that stabilizes H . In particular, if M is a k -subgroup of G that normalizes H then $M(k)$ normalizes $N_G(P_c)$.

Second construction: We allow G to be non-connected again. Suppose the following property holds for a subgroup H of G :

(*) there exists an R-parabolic k -subgroup P of G such that $H \subseteq P$ but H is not contained in any R-Levi subgroup—i.e., any R-Levi \bar{k} -subgroup—of P .

This hypothesis implies in particular that H is not G -cr over k . The construction in [6, §5] then yields a canonical so-called “optimal destabilising” R-parabolic k -subgroup P_{opt} of G such that $H \subseteq P_{\text{opt}}$ but H is not contained in any R-Levi subgroup of P_{opt} . If L is an R-Levi k -subgroup of P_{opt} then $c_L(H)$ is G -cr by [6, Prop. 4.15], so (P_{opt}, L) yields a \bar{k} -semisimplification H' of H ; if H' is also G -cr over k then H' is a k -semisimplification of H . Moreover, P_{opt} is stabilized by any k -automorphism of G that stabilizes H ; in particular, if M is a k -subgroup of G that normalizes H then $M(k)$ normalizes P_{opt} . See [6, Sec. 5.2] for details.

This construction rests on the notion of an “optimal destabilising cocharacter” due to work of Hesselink [12], Kempf [13] and Rousseau [19]. Roughly speaking, the idea is as follows. Take a generic tuple $\mathbf{h} \in H^m$ for H . There is a unique closed $G(\bar{k})$ -orbit \mathcal{O} that is accessible from \mathbf{h} via a k -defined cocharacter. We define $\lambda_{\text{opt}} \in Y_k(G)$ to be the cocharacter that takes \mathbf{h} into \mathcal{O} as quickly as possible (in an appropriate sense), and we define P_{opt} to be $P_{\lambda_{\text{opt}}}$. (In fact, we need a slight variation—due to Hesselink—on this construction: rather than taking a single generic tuple \mathbf{h} , one considers the action of a cocharacter λ on all elements of H at once.) Note that P_{opt} is not uniquely determined (see [6, Rem. 5.22]).

Remark 5.2. Suppose $C_G(H)$ is k -defined. One can show that if H is G -cr then H is G -cr over k . (The converse is false: see Example 3.3.) In fact, we prove a slightly stronger result: if H is not G -cr over k then hypothesis (*) holds. To see this, choose a generic tuple $\mathbf{h} \in H^m$. We can find $\lambda \in Y_k(G)$ such that (P_λ, L_λ) yields a semisimplification H' of H ; so $G(k) \cdot c_\lambda(\mathbf{h})$ is cocharacter-closed over k but $G(k) \cdot \mathbf{h}$ is not. If H is contained in an R-Levi \bar{k} -subgroup L of P_λ then $c_\lambda(\mathbf{h}) = u \cdot \mathbf{h}$ for some $u \in R_u(P_\lambda)$. But then [1, Thm. 7.1] implies that $c_\lambda(\mathbf{h}) = u_1 \cdot \mathbf{h}$ for some $u_1 \in R_u(P_\lambda)(k)$, so $G(k) \cdot c_\lambda(\mathbf{h}) = G(k) \cdot \mathbf{h}$, a contradiction.

In particular, if k is perfect, H is not G -cr over k and H is k -defined then $C_G(H)$ is k -defined, so we obtain a \bar{k} -semisimplification H' of H from P_{opt} , as described above. But H' is also k -defined, so $C_G(H')$ is k -defined, and we deduce from the preceding paragraph that H' is G -cr over k ; hence H' is also a k -semisimplification of H .

Remark 5.3. Suppose H is k -defined. Let M be a k -subgroup of G such that M normalizes H . Let us now extend our field from k to k_s . Let P be the k_s -parabolic subgroup obtained from one of the two constructions above: so $P = P_c$ or P_{opt} , but for the field k_s . Recall from above that $M(k_s)$ normalizes P . It follows that M normalizes P . But one can show that P is k -defined (cf. [3, Proof of Thm. 1.1] and [13, Sec. 4])—in fact, P is the k -parabolic subgroup obtained from one of the two constructions above for the field k . So M is a subgroup of $N_G(P)$. Without the hypothesis that H is k -defined, we know only that $M(k)$ is a subgroup of $N_G(P)$.

Remark 5.4. Without the hypothesis on $C_G(H)$, it can happen that the \bar{k} -simplification H' yielded by P_{opt} is not G -cr over k . For instance, let $G_1 = \text{SL}_2$ and let H_1 (resp., B_1) be the subgroup of upper unitriangular (resp., upper triangular) matrices in G_1 . Let G_2 be reductive and let H_2 be a subgroup of G_2 which is G -cr but not G -cr over k . Set $G = G_1 \times G_2$ and $H = H_1 \times H_2$. One can show that $P_{\text{opt}} = B_1 \times G_2$, and $H' = \{1\} \times H_2$ is G -cr but not G -cr over k .

To compound the problem, we do not (yet) know how to define a version of P_{opt} for a subgroup H that is not G -cr over k if $(*)$ does not hold; the obstruction is that there is not a theory of optimal destabilising subgroups that holds for arbitrary fields. See [6, Sec. 1 and Ex. 5.21] for further discussion of the latter point.

Combining the two constructions above we obtain the following.

Proposition 5.5. *Let M be a k -subgroup of G and let H be a normal k -subgroup of M . Suppose at least one of the following holds:*

- (i) k is perfect.
- (ii) $H \subseteq G^0$.

Then:

- (a) *If M is G -completely reducible over k then H is G -completely reducible over k .*
- (b) *There is an R -parabolic subgroup P of G such that $M \subseteq P$ and P yields both a k -semisimplification of M and a k -semisimplification of H . In particular, there exist k -semisimplifications M' (resp., H') of M (resp., of H) such that H' is normal in M' .*

Proof. Suppose H is not G -cr over k . Choose $P = P_{\text{opt}}$ in case (i) and $P = P_c$ in case (ii). Then $M \subseteq N_G(P)$ by Remark 5.3. Since H is not contained in any R -Levi k -subgroup of P , H is not contained in any R -Levi k -subgroup of $N_G(P)$ (Lemma 3.11). Hence M is not contained in any R -Levi k -subgroup of $N_G(P)$. It follows that M is not G -cr over k . This proves part (a).

For (b), pick $\lambda \in Y_k(G)$ such that (P_λ, L_λ) yields a semisimplification $M' := c_\lambda(M)$ of M . Then $c_\lambda(M)$ is G -cr over k and $c_\lambda(H)$ is normal in $c_\lambda(M)$. Now $c_\lambda(M)$ and $c_\lambda(H)$ satisfy the hypotheses of the proposition—note that $c_\lambda(H)$ is contained in G^0 if H is—so $c_\lambda(H)$ is G -cr over k by (a). Hence (P_λ, L_λ) yields a semisimplification $H' := c_\lambda(H)$ of H as well, and H' is normal in M' . \square

Example 5.6. Let H be a k -subgroup of $G = \text{GL}_n$ such that H is not completely reducible over k . Since H is separable, $C_G(H)$ is k -defined, so H is not G -completely reducible; we obtain a parabolic k -subgroup P_{opt} as above which yields a subgroup H' that is both a \bar{k} -semisimplification of H and a k -semisimplification of H . The parabolic subgroup P_{opt} is the stabilizer of some flag \mathcal{F} of k -defined subspaces of \bar{k}^n , and \mathcal{F} does not admit a complementary H -stable flag of subspaces of \bar{k}^n . By Remark 5.3, $C_G(H)$ is a subgroup of P_{opt} —that is, $C_G(H)$ stabilizes \mathcal{F} —and likewise the normalizer $N_G(H)$ stabilizes \mathcal{F} if $N_G(H)$ is k -defined. If k is perfect then $N_G(H)$ is automatically k -defined but it need not be k -defined in general; see [11] for further discussion.

Example 5.7. Hesselink gives an example [12, (8.5) Ex.] of a subgroup H of an almost simple group G of type C_2 such that P_{opt} is not a minimal centre for Σ , the subcomplex of the building $\Delta_k(G)$ of G consisting of all parabolic subgroups of G that contain H . This shows that the two constructions above can yield different R -parabolic subgroups. Nevertheless, their k -semisimplifications of H are $G(k)$ -conjugate in general, thanks to Theorem 4.5. In this particular instance both k -semisimplifications are the trivial subgroup in G , as H is connected unipotent.

Here are the details. Let k be an arbitrary field. Fix a Borel subgroup B of G and a maximal torus T of B . Recall that the positive roots of G with respect to B and T are $\alpha, \beta, \alpha + \beta$ and $2\alpha + \beta$, where α is short and β is long. Then H is the product of root groups

$U_{\alpha+\beta}U_{2\alpha+\beta}$. We have $B = N_G(H)$, and Hesselink shows that P_{opt} is a maximal parabolic subgroup of G which contains B .

Now suppose $k = \bar{k}$ and $\text{char}(k) = 2$. Let Σ be as above and let P_c be a centre for Σ ; recall there is at least one centre. We show that $P_c = B$. Since Σ is stable under conjugation by B , B is contained in P_c . So $P_c = B$, $P_c = P_{\text{opt}}$, or $P_c = Q$, where $Q \neq P_{\text{opt}}$ is the other maximal parabolic subgroup of G that contains B . There is an isogeny $\phi: G \rightarrow G$ such that ϕ stabilises T , ϕ interchanges U_α and U_β and ϕ interchanges $U_{\alpha+\beta}$ and $U_{2\alpha+\beta}$, cf. [9, Exp. 22.4]. In particular, ϕ stabilises B and H , and ϕ interchanges P_{opt} and Q . Now ϕ gives rise to a building automorphism of $\Delta_k(G)$ by mapping a parabolic subgroup P of G to $\phi(P)$. Since ϕ stabilises Σ , we must have $\phi(P_c) = P_c$. This forces $P_c = B$, as claimed.

Acknowledgements:

We are grateful to Brian Lawrence for his questions, which motivated this paper, and for his comments on an earlier draft.

REFERENCES

- [1] M. Bate, S. Herpel, B. Martin, G. Röhrle, *Cocharacter-closure and the rational Hilbert-Mumford theorem*, Math. Z. **287** (2017), no. 1–2, 39–72.
- [2] M. Bate, B. Martin, G. Röhrle, *A geometric approach to complete reducibility*, Invent. Math. **161**, no. 1 (2005), 177–218.
- [3] ———, *Complete reducibility and separable field extensions*, C. R. Acad. Sci. Paris Ser. I Math. **348** (2010), 495–497.
- [4] ———, *The strong Centre Conjecture: an invariant theory approach*, J. Algebra **372** (2012), 505–530.
- [5] M. Bate, B. Martin, G. Röhrle, R. Tange, *Complete reducibility and separability*, Trans. Amer. Math. Soc. **362** (2010), no. 8, 4283–4311.
- [6] ———, *Closed orbits and uniform S -instability in geometric invariant theory*, Trans. Amer. Math. Soc. **365** (2013), no. 7, 3643–3673.
- [7] A. Borel, *Linear algebraic groups*, Graduate Texts in Mathematics, **126**, Springer-Verlag 1991.
- [8] T. Brüstle, L. Hille, G. Röhrle, G. Zwara, *The Bruhat-Chevalley order of parabolic group actions in general linear groups and degeneration for Δ -filtered modules*, Adv. Math., **148**, (1999), no. 2, 203–242.
- [9] C. Chevalley, *Classification des groupes algébriques semi-simples*. Collected works. Vol. 3. Springer-Verlag, Berlin, 2005.
- [10] B. Conrad, O. Gabber, G. Prasad, *Pseudo-reductive groups*, second ed., New Mathematical Monographs, vol. 26, Cambridge University Press, Cambridge, 2015.
- [11] S. Herpel, D.I. Stewart, *On the smoothness of normalisers, the subalgebra structure of modular Lie algebras, and the cohomology of small representations*. Doc. Math. **21** (2016), 1–37.
- [12] W.H. Hesselink, *Uniform instability in reductive groups*, J. Reine Angew. Math. **303/304** (1978), 74–96.
- [13] G.R. Kempf, *Instability in invariant theory*, Ann. Math. **108** (1978), 299–316.
- [14] H. Kraft, *Geometric Methods in Representation Theory*. Representations of algebras (Puebla, 1980), pp. 180–258, Lecture Notes in Math. **944**, Springer, Berlin-New York, 1982.
- [15] B. Lawrence, W. Sawin, *The Shafarevich conjecture for hypersurfaces in abelian varieties*, preprint (2020).
- [16] B. Martin, *Generic stabilisers for actions of reductive groups*, Pacific J. Math. **279**, 397–422.
- [17] R.W. Richardson, *On orbits of algebraic groups and Lie groups*, Bull. Austral. Math. Soc. **25** (1982), no. 1, 1–28.
- [18] C. Riedtmann, *Degenerations for representations of quivers with relations*, Ann. Sci. cole Normale Sup. (4) **19**(2) (1986), 275–301.
- [19] G. Rousseau, *Immeubles sphériques et théorie des invariants*, C.R.A.S. **286** (1978), 247–250.

- [20] J.-P. Serre, *La notion de complète réductibilité dans les immeubles sphériques et les groupes réductifs*, Séminaire au Collège de France, résumé (1997).
- [21] ———, *Complète réductibilité*, Séminaire Bourbaki, 56ème année, 2003–2004, n° 932.
- [22] T.A. Springer, *Linear algebraic groups*, Second edition. Progress in Mathematics, 9. Birkhäuser Boston, Inc., Boston, MA, 1998.
- [23] T. Uchiyama, *Complete reducibility of subgroups of reductive algebraic groups over nonperfect fields I*, J. Algebra **463** (2016), 168–187.
- [24] G. Zwara, *Degenerations for Modules over representation-finite Algebras*, Amer. Math. Soc. 127 (1999), 1313–1322.
- [25] ———, *Degenerations of finite-dimensional modules are given by extensions*. Compositio Math. 121 (2000), no. 2, 205–218.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UNITED KINGDOM

E-mail address: michael.bate@york.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ABERDEEN, KING'S COLLEGE, FRASER NOBLE BUILDING, ABERDEEN AB24 3UE, UNITED KINGDOM

E-mail address: b.martin@abdn.ac.uk

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY

E-mail address: gerhard.roehrle@rub.de