BLOCKS OF THE CATEGORY $\mathcal{O}^{\mathfrak{p}}$ IN TYPE E

JIEREN HU, WEI XIAO, AND AILIN ZHANG

ABSTRACT. In this paper, we determine the blocks of $\mathcal{O}^{\mathfrak{p}}$ associated with semisimple Lie algebras of type E.

1. Introduction

Suppose that \mathfrak{g} is a complex semisimple Lie algebra with a Borel subalgebra \mathfrak{b} and a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{b}$. Let $\mathfrak{p} \supset \mathfrak{b}$ be a standard parabolic subalgebra of \mathfrak{g} . If two simple modules of $\mathcal{O}^{\mathfrak{p}}$ [R, H2] extend nontrivially, they belong to the same block of $\mathcal{O}^{\mathfrak{p}}$ [H2]. The equivalence relation generated by this relation partitions the simple modules into blocks. The category $\mathcal{O}^{\mathfrak{p}}$ can be decomposed as follows:

$$\mathcal{O}^{\mathfrak{p}}=\bigoplus_{\chi}\mathcal{O}_{\chi}^{\mathfrak{p}},$$

where $\chi = \chi_{\lambda}$ for some $\lambda \in \mathfrak{h}^*$ is the infinitesimal character of the full subcategory $\mathcal{O}^{\mathfrak{p}}_{\lambda} := \mathcal{O}^{\mathfrak{p}}_{\chi_{\lambda}}$. If $\mathfrak{p} = \mathfrak{b}$ or λ is regular, then $\mathcal{O}^{\mathfrak{p}}_{\lambda}$ is a block [H2]. Brundan shows that this also holds when the root systems of irreducible components of \mathfrak{g} is of type A [Br]. When Φ is not of type A, there are examples such that $\mathcal{O}^{\mathfrak{p}}_{\lambda}$ is a direct sum of more than one block [ES, BN, P1, X2] (In some paper, the category $\mathcal{O}^{\mathfrak{p}}_{\lambda}$ is called a "block" of $\mathcal{O}^{\mathfrak{p}}$ and a block is called a "linkage class" [P1]). This makes the problem of blocks to be quite subtle.

Theoretically, the extensions between simple modules depend on the leading coefficients (the μ -function) of Kazhdan-Lusztig polynomials [KL], which can be calculated by a recursion process. Along this line, lower rank cases F_4 and G_2 was solved using computer programs [P1]. When the rank of $\mathfrak g$ is large, the computation of μ -functions turns out to be very difficult [Lu, Xi, LX] since the recursion formulae quickly become unusable. In [X2], the second authors show that Jantzen coefficients [XZ] also determine the blocks. This Jantzen coefficients (which is relatively easy to calculate) come from the well-known Jantzen filtration [J1, J2] for standard modules. They can be used to obtain the radical filtrations of generalized Verma modules in many cases [HX]. Applying the theory of Jantzen coefficients, we determine the block decomposition of $\mathcal{O}^{\mathfrak p}$ in this paper for semisimple algebras of type E.

The problem of blocks has a deep relation with some other problems about $\mathcal{O}^{\mathfrak{p}}$, including simplicity of generalized Verma modules [J2, He, HKZ, BX] as well as

1

²⁰¹⁰ Mathematics Subject Classification. 17B10, 22E47.

Key words and phrases. category $\mathcal{O}^{\mathfrak{p}}$, block, reduction process, Jantzen coefficient.

The second author is supported by the National Science Foundation of China (Grant No. 11701381) and Guangdong Natural Science Foundation (Grant No. 2017A030310138). The third author is supported by the National Science Foundation of China (Grant No. 11504246).

homomorphism between generalized Verma modules [Bo, BC, BEJ, BN, L1, L2, M1, M2, M3, X1] and representation types of blocks of $\mathcal{O}^{\mathfrak{p}}$ [BN, P2].

This paper is organized as follows. In section 2, we provide necessary notations and definitions. The relation between Jantzen coefficients and blocks is discovered in section 3. In section 4, we give the algorithm to compute blocks. An example is described in section 5 to illustrate our algorithm. The full data about blocks of type E is presented in section 6.

2. Notations and definitions

Denote by $\Phi \subset \mathfrak{h}^*$ the root system of $(\mathfrak{g}, \mathfrak{h})$ with a positive system Φ^+ and a simple system $\Delta \subset \Phi^+$ corresponding to \mathfrak{b} . Let Φ_I be a subsystem of Φ associated with $I \subset \Delta$. Let W (resp. W_I) be the Weyl group of Φ (resp. Φ_I). Denote by $l(\cdot)$ the length function on W. For $\alpha \in \Phi$ and $\lambda \in \mathfrak{h}^*$, write $s_{\alpha}\lambda = \lambda - \langle \lambda, \alpha^{\vee} \rangle \alpha$, where $\langle \cdot, \cdot \rangle$ is the bilinear form on \mathfrak{h}^* induced from the Killing form and $\alpha^{\vee} := 2\alpha/\langle \alpha, \alpha \rangle$ is the coroot of α . The weight $\lambda \in \mathfrak{h}^*$ is called regular if $\langle \lambda, \alpha^{\vee} \rangle \neq 0$ for all roots $\alpha \in \Phi$. We say λ is integral if $\langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}$ for all $\alpha \in \Phi$. An integral weight $\lambda \in \mathfrak{h}^*$ is dominant if $\langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}^{\geq 0}$ (resp. $\langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}^{\leq 0}$) for all $\alpha \in \Delta$. If λ is integral, there exists a unique dominant weight $\overline{\lambda}$ in the orbit $W\lambda$.

Let \mathfrak{p} be the standard parabolic subalgebra corresponding to I. Then $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$, where \mathfrak{l} is a Levi subalgebra and \mathfrak{u} is a nilpotent radical of \mathfrak{g} . Set

$$\Lambda_I^+ := \{ \lambda \in \mathfrak{h}^* \mid \langle \lambda, \alpha^\vee \rangle \in \mathbb{Z}^{>0} \text{ for all } \alpha \in I \}.$$

For $\lambda \in \Lambda_I^+$, let $F(\lambda - \rho)$ be a finite dimensional simple \mathfrak{l} -modules of highest weight $\lambda - \rho$, where $\rho := \frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha$. The generalized Verma module is defined by

$$M_I(\lambda) := U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} F(\lambda - \rho).$$

Here $F(\lambda - \rho)$ is viewed as a \mathfrak{p} -module with trivial \mathfrak{u} -action. The generalized Verma module $M_I(\lambda)$ and its simple quotients $L(\lambda)$ share the same infinitesimal character χ_{λ} , where χ_{λ} is an algebra homomorphism from the center $Z(\mathfrak{g})$ of $U(\mathfrak{g})$ to \mathbb{C} so that $z \cdot v = \chi_{\lambda}(z)v$ for all $z \in Z(\mathfrak{g})$ and all $v \in M_I(\lambda)$. If $I = \emptyset$, then $\mathfrak{p} = \mathfrak{b}$ and $\mathcal{O}^{\mathfrak{p}}$ collapses to the well known BGG category \mathcal{O} [BGG]. Moreover, $M(\lambda) := M_{\emptyset}(\lambda)$ is the Verma module with highest weight $\lambda - \rho$. Denote by ch M the formal character of module $M \in \mathcal{O}$. The module M has a composition series with simple quotients isomorphic to some $L(\lambda)$. Denote by $[M: L(\lambda)]$ the multiplicity of $L(\lambda)$.

For $\lambda \in \mathfrak{h}^*$, set

$$\Phi_{\lambda} := \{ \beta \in \Phi \mid \langle \lambda, \beta \rangle = 0 \}.$$

It is obvious that Φ_{λ} is a subsystem of Φ . In particular, if λ is integral, then $\Phi_{\lambda} = w\Phi_{J}$ for some $w \in W$, where $J = \{\alpha \in \Delta \mid \langle \overline{\lambda}, \alpha \rangle = 0\}$. Define

$$^{I}W^{J} = \{ w \in W \mid \ell(xwy) = \ell(x) + \ell(w) + \ell(y) \text{ for any } x \in W_{I}, y \in W_{J} \},$$

Any simple module of $\mathcal{O}^{\mathfrak{p}}_{\lambda}$ can be written as $L(w\overline{\lambda})$ for $w \in {}^{I}W^{J}$.

3. Jantzen Coefficients and Blocks

In this section, we will discuss the relation between Jantzen coefficients and blocks of category $\mathcal{O}^{\mathfrak{p}}$. Some results here are already obtained in [X2], we write down the full proof for self containment.

3.1. Jantzen filtration and Jantzen coefficients. For $\lambda \in \Lambda_I^+$, we have the following result (see Proposition 9.6 in [H2]):

(3.1)
$$\operatorname{ch} M_I(\lambda) = \sum_{w \in W_I} (-1)^{\ell(w)} \operatorname{ch} M(w\lambda),$$

The right side of (3.1), which we denoted by $\theta(\lambda)$, is a valid character formula for arbitrary $\lambda \in \mathfrak{h}^*$. Denote

$$\Psi_{\lambda}^{+} = \{ \beta \in \Phi^{+} \backslash \Phi_{I} \mid \langle \lambda, \beta^{\vee} \rangle \in \mathbb{Z}^{>0} \},$$

$$\Psi_{\lambda}^{++} = \{ \beta \in \Psi_{\lambda}^{+} \mid \langle s_{\beta} \lambda, \alpha \rangle \neq 0 \text{ for all } \alpha \in \Phi_{I} \}.$$

The Jantzen filtration is a useful tool in the study of Verma modules[J1, H2]. With Lemma 3, Satz 2 and the observation in the Bemerkung before Lemma 4 in [J2], along Jantzen's line for Verma modules (see [J1] or [H2]), one can obtain the following result:

Theorem 3.2 (Jantzen filtration and sum formula for generalized Verma modules). Let $\lambda \in \Lambda_I^+$, then $M_I(\lambda)$ has a filtration by submodules

$$M_I(\lambda) = M_I(\lambda)^0 \supset M_I(\lambda)^1 \supset M_I(\lambda)^2 \supset \dots$$

with $M_I(\lambda)^i = 0$ for large i, such that

- (1) Every nonzero quotient $M_I(\lambda)^i/M_I(\lambda)^{i+1}$ has a nondegenerate contravariant form.
- (2) $M_I(\lambda)^1$ is the unique maximal submodule of $M_I(\lambda)$.
- (3) There is a formula:

(3.3)
$$\sum_{i>0} \operatorname{ch} M_I(\lambda)^i = \sum_{\beta \in \Psi_\lambda^+} \theta(s_\beta \lambda).$$

Now recall the Jantzen coefficients defined in [XZ]. The right side of (3.3) can be written as

(3.4)
$$\sum_{\beta \in \Psi_{\lambda}^{+}} \theta(s_{\beta}\lambda) = \sum_{\beta \in \Psi_{\lambda}^{++}} \theta(s_{\beta}\lambda) = \sum_{\nu \in \Lambda_{I}^{+}} c(\lambda, \nu) \operatorname{ch} M_{I}(\nu),$$

where $c(\lambda, \nu)$ are called the *Jantzen coefficients* associated with (Φ_I, Φ) . Evidently $c(\lambda, \nu)$ are nonzero for only finitely many $\nu \in \Lambda_I^+$. Write $\mu \leq \lambda$ when $\operatorname{Hom}_{\mathcal{O}}(M(\mu), M(\lambda)) \neq 0$ for $\lambda, \mu \in \mathfrak{h}^*$. If $\lambda, \mu \in \Lambda_I^+$ and $\operatorname{Ext}^1_{\mathcal{O}_{\mathcal{P}}}(L(\mu), L(\lambda)) \neq 0$, then either $\mu < \lambda$ or $\lambda < \mu$. If $c(\lambda, \mu) = 0$, then $\lambda > \mu$.

Definition 3.5. Fix $I \subset \Delta$. Let $\lambda, \mu \in \Lambda_I^+$. We write $\lambda \succeq \mu$ if there exist $\lambda^0, \lambda^1, \ldots, \lambda^k \in \Lambda_I^+$ $(k \geq 0)$ so that $\lambda^0 = \lambda$, $\lambda^k = \mu$ and $c(\lambda^{i-1}, \lambda^i) \neq 0$ for $1 \leq i \leq k$. This induces an ordering on Λ_I^+ . In particular, if $\lambda \succ \mu$ and there exists no $\lambda > \nu > \mu$ with $\lambda \succ \nu \succ \mu$, we say λ is adjacent to μ .

Lemma 3.6. Let $\lambda, \mu \in \Lambda_I^+$.

- (1) If $[M_I(\lambda), L(\mu)] > 0$, then $\lambda \succeq \mu$.
- (2) If λ is adjacent to μ , then $[M_I(\lambda), L(\mu)] > 0$.

Proof. (1) The case $\lambda = \mu$ is trivial. Assume that $[M_I(\lambda), L(\mu)] > 0$ and $\mu < \lambda$. We obtain

(3.7)
$$\sum_{i>0} [M_I(\lambda)^i : L(\mu)] = \sum_{\nu \in \Lambda_I^+} c(\lambda, \nu) [M_I(\nu) : L(\mu)],$$

in view of (3.3) and (3.4). The left side of (3.7) is nonzero. If $c(\lambda,\mu) \neq 0$, then $\lambda \succ \mu$. If $c(\lambda,\mu) = 0$, there exists $\lambda > \lambda^1 > \mu$ such that $c(\lambda,\lambda^1) \neq 0$ and $[M_I(\lambda^1):L(\mu)] \neq 0$. Thus $\lambda \succ \lambda^1$. We get (1) when $\lambda^1 \succ \mu$, otherwise $c(\lambda^1,\mu) = 0$. With λ replaced by λ^1 in the previous argument, there exists $\lambda^1 \succ \lambda^2 > \mu$ so that $c(\lambda^1,\lambda^2)[M_I(\lambda^2):L(\mu)] \neq 0$. In this spirit, we can eventually get $\lambda \succ \lambda^1 \succ \ldots \succ \lambda^{k-1} \succ \mu$ for some $k \geq 1$.

(2) Suppose that λ is adjacent to μ . Then $c(\lambda, \mu) \neq 0$ by definition. If $c(\lambda, \nu)[M_I(\nu) : L(\mu)] \neq 0$ for some $\nu \in \Lambda_I^+$, the argument of the first part shows that $\lambda \succ \nu \succeq \mu$. The adjacent condition implies $\nu = \lambda$. So $c(\lambda, \nu)[M_I(\nu) : L(\mu)] = 0$ unless $\nu = \mu$. In view of (3.7), one has

(3.8)
$$\sum_{i>0} [M_I(\lambda)^i : L(\mu)] = c(\lambda, \mu) \neq 0.$$

Hence $[M_I(\lambda):L(\mu)]>0$.

3.2. Blocks and lined roots.

Definition 3.9. Fix $I \subset \Delta$. For $\lambda, \mu \in \Lambda_I^+$, write $\lambda \leftrightarrow \mu$ if $\operatorname{Ext}_{\mathcal{O}^{\mathfrak{p}}}^1(L(\mu), L(\lambda)) \neq 0$. In general, write $\lambda \leftrightarrow \mu$ if $\lambda = \mu$ or there exist $\lambda^0, \lambda^1, \ldots, \lambda^r \in \Lambda_I^+$ such that

$$\lambda = \lambda^0 \leftrightarrow \lambda^1 \leftrightarrow \ldots \leftrightarrow \lambda^r = \mu$$
,

we say $\lambda, \mu \in \Lambda_I^+$ are Ext^1 -connected relative to (Φ_I, Φ) . For convenience, if $\lambda \leftrightarrow \mu$ relative to (Φ_I, Φ) , we write $w_1 \lambda \leftrightarrow w_2 \mu$ for any $w_1, w_2 \in W_I$.

The definition gives an equivalence relation on all the Φ_I -regular weights such that each W_I orbit is contained in one equivalence class. The following result is well known.

Lemma 3.10. Let $\lambda, \mu \in \Lambda_I^+$ with $\mu < \lambda$. If $\operatorname{Ext}_{\mathcal{O}^{\mathfrak{p}}}^{1}(L(\mu), L(\lambda)) \neq 0$, then $[M_I(\lambda), L(\mu)] > 0$. Conversely, if $[M_I(\lambda), L(\mu)] > 0$, then $\lambda \leftrightarrow \mu$.

Lemma 3.11. If $c(\lambda, \mu) \neq 0$ for $\lambda, \mu \in \lambda_I^+$, there exists $\beta \in \Psi_{\lambda}^+$ and $w \in W_I$ so that $\mu = ws_{\beta}\lambda$. Moreover, $\lambda \leftrightarrow \mu$.

Proof. It suffices to prove the first statement. Note that $c(\lambda, \mu) = \sum_{\beta \in \Psi_{\lambda, \mu}^+} (-1)^{\ell(w_\beta)}$ (see [XZ], §4), where

$$\Psi_{\lambda,\mu}^+ := \{ \beta \in \Psi_{\lambda}^+ \mid \mu = w_{\beta} s_{\beta} \lambda \text{ for some } w_{\beta} \in W_I \}.$$

So $\Psi_{\lambda,\mu}^+$ is not empty in view of $c(\lambda,\mu) \neq 0$. Choose $\beta \in \Psi_{\lambda,\mu}^+$ and set $w = w_\beta$. \square

Lemma 3.11 and the dual invariance of Jantzen coefficients (see Lemma 4.17 in [XZ]) yield the following result.

Lemma 3.12. Let $I, J \subset \Delta$. Suppose that λ (resp. μ) is a dominant weight with $\Phi_{\lambda} = \Phi_{J}$ (resp. $\Phi_{\mu} = \Phi_{I}$). Choose $x, w \in {}^{I}W^{J}$. Then $x\lambda \leftrightarrow w\lambda$ (relative to (Φ_{I}, Φ)) if and only if $x^{-1}\mu \leftrightarrow w^{-1}\mu$ (relative to (Φ_{J}, Φ)).

The above lemma shows a kind of parabolic-singular duality for category $\mathcal{O}^{\mathfrak{p}}$ [So, BGS, B]. Lemma 3.11 and the conjugate invariance of Jantzen coefficients (see Lemma 4.18 in [XZ]) implies the following lemma.

Lemma 3.13. Let $I, J \subset \Delta$. Suppose that Φ_I and Φ_J are W-conjugate, choose $w \in W$ so that $\Phi_J^+ = w\Phi_I^+$. Let $\lambda, \mu \in \Lambda_I^+$. Then $\lambda \leftrightarrow \mu$ (relative to (Φ_I, Φ)) if and only if $w\lambda \leftrightarrow w\mu$ (relative to (Φ_J, Φ)).

Remark 3.14. In type E_7 , there are subsets $I, I' \subset \Delta$ such that Φ_I and $\Phi_{I'}$ are of the same type, but they are not W-conjugate (see [CM] or [P2]). This happens when Φ_I is of type A_5 , $A_3 \times A_1$ and A_1^3 . In each of these cases, there are two conjugate classes, which are denoted by Φ_I' and Φ_I'' .

Lemma 3.15. Let $\lambda, \mu \in \Lambda_I^+$. If $\lambda \leftrightarrow \mu$ relative to (Φ_I, Φ) , then $\lambda \leftrightarrow \mu$ relative to $(\Phi_{I'}, \Phi)$ for any $I' \subset I$.

Proof. This follows from the fact that

$$\operatorname{Ext}^1_{\mathcal{O}^{\mathfrak{p}_I}}(L(\mu),L(\lambda)) \simeq \operatorname{Ext}^1_{\mathcal{O}^{\mathfrak{b}}}(L(\mu),L(\lambda)) \simeq \operatorname{Ext}^1_{\mathcal{O}^{\mathfrak{p}_{I'}}}(L(\mu),L(\lambda)).$$

Definition 3.16. Suppose $\lambda, \mu \in \Lambda_I^+$. We say λ, μ are linked (relative to (Φ_I, Φ)) when $c(\lambda, \mu) \neq 0$. If $\mu = ws_{\beta}\lambda$ for some $w \in W_I$ and $\beta \in \Psi_{\lambda}^+$, we say β is a linked root from λ to μ or λ, μ are linked by β . In this situation, we will say $w\beta$ is a linked root from μ to λ (keeping in mind that $\lambda = w^{-1}s_{w\beta}\mu$). In general, if β is linked root of $\lambda, \mu \in \Lambda_I^+$, we also say $w_1\beta$ is a linked root from $w_1\lambda$ to $w_2\mu$ (keeping in mind that $w_2\mu = w_2ww_1^{-1}s_{w_1\beta}w_1\lambda$) for $w_1, w_2 \in W_I$.

Lemma 3.17. For $\lambda, \mu \in \Lambda_I^+$, $\lambda \leftrightarrow \mu$ if and only if we can find $\lambda^0 = \lambda, \lambda^1, \dots, \lambda^r = \mu \in \Lambda_I^+$ with linked roots β_i from λ^{i-1} to λ^i for $1 \le i \le r$.

Proof. (1) Assume that $\lambda \leftrightarrow \mu$. By Definition 3.9, it suffices to consider the case $\operatorname{Ext}^1_{\mathcal{O}^{\mathfrak{p}}}(L(\mu),L(\lambda))\neq 0$ with $\mu<\lambda$. In view of Lemma 3.10, we have $[M_I(\lambda),L(\mu)]>0$. Lemma 3.6 yields a chain $\lambda=\lambda^0\succ\lambda^1\succ\cdots\succ\lambda^k=\mu$ such that $\lambda^i\in\Lambda_I^+$ and $c(\lambda,\mu)\neq 0$. Then Lemma 3.11 gives linked roots β_i from λ^{i-1} to λ^i .

(2) The converse follows from Lemma
$$3.11$$
.

The Jantzen coefficients have several invariant properties under a reduction process (see [XZ], §3). Choose $\beta \in \Phi$. Let $\Phi_{\beta,0}$ be the irreducible component of Φ with $\beta \in \Phi_{\beta,0}$. Set

$$\Phi_{\beta,1} := (\mathbb{Q}\Phi_I + \mathbb{Q}\beta) \cap \Phi;$$

$$\Phi_{\beta,2} := (\mathbb{Q}\Phi_\lambda + \mathbb{Q}\beta) \cap \Phi.$$

For $\beta \in \Psi_{\lambda}^+$, consider the following chain of subsystems

$$\Phi = \Phi_0(\beta) \supset \Phi_1(\beta) \supset \Phi_2(\beta) \supset \ldots \supset \Phi_m(\beta) \supset \ldots$$

such that for $i \in \mathbb{Z}^{>0}$,

$$\Phi_{i+1}(\beta) = \begin{cases}
(\Phi_i(\beta))_{\beta,0} & \text{if } i \equiv 0 \text{ (mod 4)}; \\
(\Phi_i(\beta))_{\beta,1} & \text{if } i \equiv 1 \text{ (mod 4)}; \\
(\Phi_i(\beta))_{\beta,0} & \text{if } i \equiv 2 \text{ (mod 4)}; \\
(\Phi_i(\beta))_{\beta,2} & \text{if } i \equiv 3 \text{ (mod 4)}.
\end{cases}$$

The chain is stationary, that is, there exists $k \in \mathbb{Z}^{>0}$ with $\Phi_k(\beta) = \Phi_{k+1}(\beta) = \dots$ Denote $\Phi(\beta) = \Phi_k(\beta)$. Recall that for any subsystem Φ' of Φ , there exists a unique weight $\lambda|_{\Phi'}$ in the subspace $\mathbb{C}\Phi'$ so that

$$\langle \lambda |_{\Phi'}, \alpha^{\vee} \rangle = \langle \lambda, \alpha^{\vee} \rangle$$

for all $\alpha \in \Phi'$.

Lemma 3.18 ([XZ, Lemma 3.17]). Let $\lambda \in \Lambda_I^+$ and $\beta \in \Psi_{\lambda}^+$. Then $\Phi(\beta)$ is irreducible and $\lambda|_{\Phi(\beta)}$ is an integral weight on $\Phi(\beta)$. Moreover,

$$rank(\Phi_I \cap \Phi(\beta)) = rank(\Phi_\lambda \cap \Phi(\beta)) = rank \ \Phi(\beta) - 1.$$

In [XZ], the system (Φ, Φ_I, Φ_J) is called a basic system if Φ is irreducible and $\operatorname{rank}\Phi_I = \operatorname{rank}\Phi_\lambda = \operatorname{rank}\Phi - 1$. The above lemma shows that $(\Phi(\beta), \Phi_I \cap \Phi(\beta), \Phi_\lambda \cap \Phi(\beta))$ is a basic system.

Proposition 3.19. Let $\lambda \in \Lambda_I^+$ and $\beta \in \Psi_{\lambda}^+$. Then λ and $s_{\beta}\lambda$ are linked by β (relative to (Φ_I, Φ)) if and only if $\lambda|_{\Phi(\beta)}$ and $s_{\beta}\lambda|_{\Phi(\beta)}$ are linked by β (relative to $(\Phi_I \cap \Phi(\beta), \Phi(\beta))$).

Proof. The invariant properties of Jantzen coefficients ([XZ], §4) implies that $s_{\beta}\lambda$ is Φ_{I} -regular if and only if $\lambda' = s_{\beta}(\lambda|_{\Phi(\beta)}) = s_{\beta}\lambda|_{\Phi(\beta)}$ is $\Phi_{I} \cap \Phi(\beta)$ -regular. There exist $\mu \in \Lambda_{I}^{+}$ (resp. $\mu' \in \Lambda^{+}(\lambda|_{\Phi(\beta)}, \Phi_{I} \cap \Phi(\beta), \Phi(\beta))$) and $w \in W_{I}$ (resp. $w' \in W(\Phi_{I} \cap \Phi(\beta))$) with $\mu = ws_{\beta}\lambda$ (resp. $\mu' = w's_{\beta}\lambda'$). Moreover, the Jantzen coefficient $c(\lambda, \mu)$ associated with (Φ_{I}, Φ) is nonzero if and only if the Jantzen coefficients $c(\lambda', \mu')$ associated with $(\Phi_{I} \cap \Phi(\beta), \Phi(\beta))$ is nonzero.

Lemma 3.20. Let $\Phi = D_n$ and $\beta \in \Phi^+ \backslash \Phi_I$. Suppose that $\{e_{n-1} \pm e_n\}$ is not a subset of I. If λ and $s_{\beta}\lambda$ are Φ_I -regular, then β is always a linked root from λ .

Proof. Keeping in mind of Lemma 3.11, this lemma is an easy consequence of Theorem 7.27 in [XZ]. \Box

4. The algorithm

Although we can run a computer program to determine the blocks of type E, it might take too much time when |I| + |J| are too small. For example, if $I = J = \emptyset$, the category $\mathcal{O}^{\mathfrak{p}}_{\lambda}$ has 696729600 simple modules. This is beyond the processing ability of an ordinary PC. So we need to simplify to calculation. For $\Phi = E_n$ (n = 6, 7, 8), let $\alpha_1, \dots, \alpha_n$ be the standard ordering of simple roots ([H1], §11.4).

Lemma 4.1. Let $\Phi = E_n$ for n = 6, 7, 8. Choose $I \subset \Delta \setminus \{\alpha_1, \alpha_2\}$ or $I \subset \Delta \setminus \{\alpha_1, \alpha_3\}$. Suppose that $\lambda, s_{\beta}\lambda$ are Φ_I -regular weights with $\beta = e_i \pm e_j$ for $1 \leq j < i \leq n - 1$. Then $\lambda \leftrightarrow s_{\beta}\lambda$.

Proof. This is obvious when $\beta \in \Phi_I$. If $\beta \notin \Phi_I$, set $\Phi' := \Phi_{\Delta'} \simeq D_{n-1}$ and $\lambda' = \lambda|_{\Phi'}$, where $\Delta' = \Delta \setminus \{\alpha_1\}$. Lemma 3.20 implies that β is linked root from λ . With $\alpha_1 \notin I$, one has $\Phi_{I'} = \Phi_I \cap \Phi' = \Phi_I$. It follows that $\Phi_{\beta,1} = \Phi'_{\beta,1}$ and $\Phi(\beta) = \Phi'(\beta)$. In view of Proposition 3.19, β is also a linked root from λ and thus $\lambda \leftrightarrow s_{\beta}\lambda$.

Lemma 4.2. Let $\Phi = E_8$. Assume that $|I| + |J| \le 7$ and $|I| \le 3$. Suppose that (Φ, Φ_I, Φ_J) contains only one block. If $I' \subset I$, then $(\Phi, \Phi_{I'}, \Phi_J)$ has only one block.

Proof. Since the case $I = \emptyset$ is trivial, we can assume that $1 \le |I| \le 3$. This forces $\Phi_I \simeq A_3, A_2 \times A_1, A_1^3, A_2, A_1^2$ or A_1 . With Lemma 3.13, it suffices to consider the cases $I = \{\alpha_6, \alpha_7, \alpha_8\}, \{\alpha_5, \alpha_7, \alpha_8\}, \{\alpha_4, \alpha_6, \alpha_8\}, \{\alpha_7, \alpha_8\}, \{\alpha_6, \alpha_8\}$ or $\{\alpha_8\}$.

For any $\Phi_{I'}$ -regular weight with $\Phi_{\overline{\lambda}} = \Phi_J$, we claim there exists $\hat{\lambda} \in \Lambda_I^+$ so that $\hat{\lambda} \leftrightarrow \lambda$ (relative to $(\Phi_{I'}, \Phi)$). The lemma is an easy consequence of the claim. In fact, given $\lambda, \mu \in {}^{I'}W^J\overline{\lambda}$, there exist Φ_I -regular weights $\hat{\lambda}, \hat{\mu} \in \Lambda_I^+$ so that $\lambda \leftrightarrow \hat{\lambda}$ and $\mu \leftrightarrow \hat{\mu}$ (relative to $(\Phi_{I'}, \Phi)$). Since (Φ, Φ_I, Φ_J) contains only one block, one

has $\hat{\lambda} \leftrightarrow \hat{\mu}$ (relative to (Φ_I, Φ)). Applying Lemma 3.15, we get $\lambda \leftrightarrow \hat{\lambda} \leftrightarrow \hat{\mu} \leftrightarrow \mu$ (relative to $(\Phi_{I'}, \Phi)$), that is, $(\Phi, \Phi_{I'}, \Phi_J)$ has only one block.

It remains to prove the claim, which can be achieved in a case-by-case fashion. We only discuss the case $I = \{\alpha_6, \alpha_7, \alpha_8\} = \{e_5 - e_4, e_6 - e_5, e_7 - e_6\}$, while the reasoning for the other cases are similar. Set $\Phi' := \Phi_{\Delta'}$ and $\lambda' = \lambda|_{\Phi'}$, where $\Delta' = \Delta \setminus \{\alpha_1\}$. Note that

$$\operatorname{rank}\Phi_{\lambda'}' = \operatorname{rank}\Phi_{\lambda} \cap \Phi' \leq \operatorname{rank}\Phi_{\lambda} = \operatorname{rank}\Phi_{J} \leq 7 - |I| = 4$$

in this case. Let $a_1 > \cdots > a_m$ be all the different numbers in $\{|\lambda_i| \mid 1 \leq i \leq 7\}$. Put $n_i = |\{1 \leq i \leq 7 \mid |\lambda_i| = a_i\}|$. If $a_m > 0$, then $\Phi'_{\lambda'} \simeq A_{n_1-1} \times \cdots \times A_{n_m-1}$, where $A_0 = 1$. We obtain

$$rank\Phi'_{1} = (n_1 - 1) + \dots + (n_m - 1) = 7 - m < 4,$$

that is, $m \geq 3$. If $a_m = 0$ and $n_m = 1$, similar reasoning shows $m \geq 3$. If $a_m = 0$ and $n_m \geq 2$, then $\Phi'_{\lambda'} \simeq A_{n_1-1} \times \cdots \times D_{n_m}$ and $m \geq 4$, where $D_2 \simeq A_1 \times A_1$ and $D_3 \simeq A_3$.

Set $\mu = \lambda$ and $S_{\mu} = \{|\mu_i| \mid 4 \leq i \leq 7\}$. If $a_1 \in S_{\mu}$, there exists $4 \leq i_1 \leq 7$ so that $|\mu_{i_1}| = a_1$. Otherwise we can find $1 \leq i \leq 3$ with $|\mu_i| = a_1$. Thus $s_{e_i - e_7} \mu$ is $\Phi_{I'}$ -regular (since $a_1 \notin S_{\mu}$). Replacing μ by $s_{e_i - e_7} \mu$, we get $\lambda \leftrightarrow \mu$ (relative to $(\Phi_{I'}, \Phi)$) and $|\mu_7| = a_1$ $(i_1 = 7)$. In a similar spirit, one can eventually get $\Phi_{I'}$ -regular weight $\mu \leftrightarrow \lambda$ and $|\mu_{i_j}| = a_j$ for $4 \leq i_j \leq 7$ and $1 \leq j \leq \min\{m, 4\}$. If μ_4, \dots, μ_7 are not distinct, one must have m = 3 and $\mu_{i_4} = \mu_{i_j} = \pm a_j \neq 0$ for some $1 \leq j \leq 3$, where $i_4 \in \{4, 5, 6, 7\} \setminus \{i_1, i_2, i_3\}$. Choose $l \in \{1, 2, 3\} \setminus \{j\}$. Replace μ by $s_{e_{i_4} + e_{i_l}} \mu$. Then μ_4, \dots, μ_7 are distinct and $\lambda \leftrightarrow \mu$ (relative to $(\Phi_{I'}, \Phi)$). Now we can find $w \in W_I$ so that $w \in \Lambda_I^+$. Since μ_4, \dots, μ_7 are distinct, Lemma 4.1 yields $\mu \leftrightarrow w \mu$ (relative to $(\Phi_{I'}, \Phi)$). Set $\hat{\lambda} = w \mu$.

Keeping in mind of Remark 3.14, the proof of the following two results are similar and easier.

Lemma 4.3. Let $\Phi = E_7$. Assume that $|I| + |J| \le 6$ and $|I| \le 3$. Suppose that (Φ, Φ_I, Φ_J) contains only one block. If $I' \subset I$, then $(\Phi, \Phi_{I'}, \Phi_J)$ has only one block.

Lemma 4.4. Let $\Phi = E_6$. Assume that $|I| + |J| \le 5$ and $|I| \le 2$. Suppose that (Φ, Φ_I, Φ_J) contains only one block. If $I' \subset I$, then $(\Phi, \Phi_{I'}, \Phi_J)$ has only one block.

4.1. **The algorithm.** In this subsection, we will provide an algorithm to find data of the system (Φ, Φ_I, Φ_J) .

First consider $\Phi = E_8$. Let $\varpi_J = \sum_{\alpha_j \notin J} \varpi_j$, where $\varpi_1, \ldots, \varpi_8$ are fundament weights. It is a dominant weight with $\Phi_{\varpi_J} = \Phi_J$. We need to find all the integral weights $\lambda = w\varpi_J$ with $w \in {}^IW^J$. Assume that $\lambda = \sum_{i=1}^8 x_i\varpi_k$ for $x_i \in \mathbb{Z}$. In particular, $x_i \in \mathbb{Z}^{>0}$ when $\alpha_i \in I$ (since $\lambda \in \Lambda_I^+$). Changing basis,

$$\lambda = \frac{x_2 - x_3}{2}e_1 + \frac{x_2 + x_3}{2}e_2 + (\frac{x_2 + x_3}{2} + x_4)e_3 + (\frac{x_2 + x_3}{2} + x_4 + x_5)e_4 + (\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6)e_5 + (\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6 + x_7)e_6 + (\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6 + x_7 + x_8)e_7 + (2x_1 + \frac{5x_2 + 7x_3}{2} + 5x_4 + 4x_5 + 3x_6 + 2x_7 + x_8)e_8.$$

With $\langle \lambda, \lambda \rangle = \langle w \varpi_J, w \varpi_J \rangle = \langle \varpi_J, \varpi_J \rangle$, x_i $(1 \le i \le 8)$ are integral solutions (with $x_i > 0$ for $\alpha_i \in I$) of the equation

$$\langle \varpi_J, \varpi_J \rangle = \left(\frac{x_2 - x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6 + x_7\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6 + x_7 + x_8\right)^2 + \left(2x_1 + \frac{5x_2 + 7x_3}{2} + 5x_4 + 4x_5 + 3x_6 + 2x_7 + x_8\right)^2.$$

The algorithm is summarized as follows.

- (1) Choose $I, J \subset \Delta$ with $|I| + |J| \ge 7$.
- (2) Find all the integral solutions (x_1, \dots, x_8) of the above equations with $x_i > 0$ for $\alpha_i \in I$ using enumeration method.
- (3) For each solution (x_1, \dots, x_8) obtained in (2), write $\mu = \lambda = \sum_{i=1}^8 x_i \varpi_i$. If $\langle \lambda, \alpha \rangle < 0$ for some $\alpha \in \Delta$, replace μ by $s_{\alpha}\mu$. We eventually arrive at a dominant weight μ . If $\mu = \varpi_J$, keep the solution; otherwise it is discarded.
- (4) Let $\lambda^1, \dots, \lambda^N$ be all the vector solutions obtained in (3) with i < j whenever $\langle \lambda^i, \varpi_I \rangle > \langle \lambda^j, \varpi_I \rangle$. Set $a(\lambda^i) = i$. First take $\lambda = \lambda^1$. For any $\beta \in \Psi_{\lambda}^{++}$, we can find $\mu \in \Lambda_I^+$ so that $\mu = ws_{\beta}\lambda$ for some $w \in W_I$. Then μ is also a vector solution of (3). If $c(\lambda, \mu) \neq 0$ and $a(\lambda) \neq a(\mu)$, set $a(\lambda^i) = \min\{a(\lambda), a(\mu)\}$ for any $1 \leq i \leq N$ with $a(\lambda^i) = a(\lambda)$ or $a(\mu)$. Next take $\lambda = \lambda^2, \dots, \lambda^N$ sequentially and iterate the above procedure.
- (5) When step (4) is finished, let S be the set of all the different $a(\lambda^i)$. Thus the system contains |S| blocks. If |S| > 1, we output vectors of each block. Otherwise we output the number N.

It turns out that one always has |S| = 1 when |I| + |J| = 7. The following result is an immediate consequence of Lemma 4.2.

Lemma 4.5. Let $\Phi = E_8$. Assume that $|I| + |J| \le 7$. Then (Φ, Φ_I, Φ_J) contains only one block.

The algorithms for type E_7 , E_6 are similar, while the equation for type E_7 is

$$\langle \varpi_J, \varpi_J \rangle = \left(\frac{x_2 - x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6 + x_7\right)^2 + 2\left(\frac{3x_3 + 3x_5 + x_7}{2} + x_1 + x_2 + 2x_4 + x_6\right)^2$$

and the equation for type E_6 is

$$\langle \varpi_J, \varpi_J \rangle = \left(\frac{x_2 - x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2}\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5\right)^2 + \left(\frac{x_2 + x_3}{2} + x_4 + x_5 + x_6\right)^2 + 3\left(\frac{x_2}{2} + \frac{2x_1 + 2x_5 + x_6}{3} + \frac{5x_3}{6} + x_4\right)^2.$$

We also have the following results.

Lemma 4.6. Let $\Phi = E_7$. Assume that $|I| + |J| \le 6$. Then (Φ, Φ_I, Φ_J) contains only one block.

Lemma 4.7. Let $\Phi = E_6$. Assume that $|I| + |J| \le 5$. Then (Φ, Φ_I, Φ_J) contains only one block.

5. An example

In this section, we give a typical example of system with two blocks. The result is obtained by the algorithm in the previous section. Let $\Phi = E_8$, $I = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$ and $J = \{\alpha_3, \alpha_4, \alpha_5, \alpha_7\}$. Thus $\Phi_I \simeq D_6$ and $\Phi_J \simeq A_3 \times A_1$. Evidently, $\varpi_J = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, \frac{5}{2}, \frac{17}{2})$. All the weights of ${}^IW^J\varpi_J$ are given in Table 1.

$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	\(\frac{1}{2}\)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	\(\frac{1}{2}\)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	\(\frac{1}{2}\)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	<u>\</u>)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$)
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
10 $(0,1,2,3,4,6,-3,3)$ 32 $(\frac{1}{2},\frac{3}{2},\frac{5}{2},\frac{7}{2},\frac{9}{2},\frac{11}{2},-\frac{7}{2},\frac{1}{2})$	
)
11 $(\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}, \frac{7}{2}, \frac{1}{2})$ 33 $(0, 1, 2, 3, 4, 7, 1, -2)$	′
(2, 2, 2, 2, 2, 2, 2)	
$ \boxed{ 12 \ \ (\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}, -\frac{5}{2}, \frac{5}{2}) \ \ 34 \ \ (0, 1, 2, 3, 4, 7, -2, -1) } $	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	<u>'</u>)
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$\frac{5}{2}$)
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	<u>'</u>)
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	
$ 22 \left \left(-\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{13}{2}, -\frac{1}{2}, \frac{1}{2} \right) \right 44 \left \left(0, 1, 2, 3, 4, 5, -2, -5 \right) \right $	

Table 1.

Table 1 shows that the system (Φ, Φ_I, Φ_J) contains 44 simple highest weight modules. Now we explore the relation between them. Write $c_{i,j} = c(\lambda^i, \lambda^j)$. It was proved in [XZ] that $|c_{i,j}| \leq 1$ in this case. All the nonzero Jantzen coefficients are given in Table 2.

Suppose that $c_{i,j} \neq 0$. If there exists no sequence $i = i_0 < i_1 < \dots < i_k = j$ such that $c(i_{t-1}, i_t) \neq 0$ for $1 \leq t \leq k$ and k > 1, we say λ^i and λ^j are adjacent. In this

i	$\{j \mid c_{i,j} = 1\}$	$\{j \mid c_{i,j} = -1\}$	i	$\{j \mid c_{i,j} = 1\}$	$\{j \mid c_{i,j} = -1\}$
1	2, 4, 14, 29, 41, 43	8, 20, 32	23	33, 34, 40	37,44
2	7, 9, 16, 24, 44	8, 13, 19	24	41	_
3	6, 10, 17, 30, 35, 42	15, 26, 39	25	26, 27, 35	31
4	7, 8, 20, 32, 38	14, 29, 41	26	30,42	39
5	11, 21, 23, 24, 37, 44	16, 33, 40	27	30, 31	35
6	12, 15, 26, 31, 39	17, 30, 42	28	29, 32, 36, 38	34, 37, 43
7	8, 13, 19, 40	9, 16, 24	29	32, 34, 37, 43	36, 38, 41
8	9, 14, 16, 24, 29, 37, 41	13, 19, 20, 32	30	39	42
9	13, 14, 19, 33	16, 24	31	35, 39	_
10	12, 22, 27, 35, 42	18, 25, 31	32	41	_
11	16, 19, 28, 33, 40	23, 37, 44	33	36, 37, 44	40
12	18, 25, 31, 39	22, 27, 35	34	36, 38	37,43
13	16, 20, 23, 24	19	35	42	_
14	20, 32, 36	29,41	36	37, 43	38
15	17, 18, 27, 30, 42	26,39	37	38,40	43,44
16	19, 23, 29, 37, 44	24, 33, 40	38	40, 43	_
17	22, 25, 26, 39	30,42	39	42	_
18	22, 27, 30, 35	25, 31	40	44	_
19	24, 32		41	_	
20	29, 34, 41	32	42	_	_
21	28, 34, 37, 41, 43	29, 36, 38	43	44	_
22	25, 26, 31	27, 35	44		

Table 2.

case, it was shown in Lemma 3.6 that $L(\lambda^j)$ is a subquotient of $M_I(\lambda^i)$. We connect i and j when λ^i and λ^j are adjacent. This gives us the poset in Figure 1. Two simple modules $L(\lambda^i)$ and $L(\lambda^j)$ belongs to the same block if and only if i and j are connected in the poset. The poset exposes the block decomposition of the system (Φ, Φ_I, Φ_J) . The system decomposes into two blocks. One block contains 28 simple modules and the other contains 16. It can be easily found in Table 2 and Figure 1 that the system has three simple generalized Verma modules $M_I(\lambda^{41})$, $M_I(\lambda^{42})$ and $M_I(\lambda^{44})$. For i = 1, 3, 5, the simple module $L(\lambda^i)$ is not a composition factor of generalized Verma modules other than $M_I(\lambda^i)$.

6. Disconnected systems

In this section, we will present the data of disconnected systems, that is, those systems which contains more than one block.




Figure 1.

There are 418 different pairs of (I,J) such that the corresponding system is disconnected for type E_8 , while there are 294 pairs for type E_7 and 110 pairs for type E_6 . Of course we can not describe the full information associated with each pair as the example in the previous section. In view of Lemma 3.12 and Lemma 3.13, the block decomposition of the system (Φ, Φ_I, Φ_J) has some invariant properties up to conjugate classes of Φ_I and Φ_J . The disconnected systems are listed in Table 3, Table 5 and Table 7 according to these conjugate classes. By duality, we only illustrated data for those pairs with $|I| \geq |J|$. If |I| = |J| and $I \neq J$, we pick either one of the two pairs. The conjugate classes of Φ_I are listed in the first and the forth columns in each table, while the conjugate classes of Φ_J are listed in the second and the fifth columns. If a conjugate class Φ_I contains k subsets I with k > 1, we write $\Phi_I(k)$ in the table. For example $D_4 \times A_1(2)$ in Table 3 means that there are two subsets I (which are $\{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_7\}$ and $\{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_8\}$) such that Φ_I are in the conjugate class represented by $D_4 \times A_1$. The number of blocks and simple modules are given in the third and sixth columns. For example, if Φ_I and Φ_J are

both conjugate to $D_4 \times A_2$, the formula $2 \times 12 + 20$ means that the corresponding systems have three blocks, two of them contain 12 simple modules, while the other one has 20 simple modules.

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
E_7	$A_1^3(21)$	2×1	$A_5(4)$	$D_4 \times A_1(2)$	72 + 201
D_7	$A_2 \times A_1^3(8)$	2×1	$A_5 \times A_1(3)$	D_4	96 + 144
A_7	$A_3 \times A_2 \times A_1(4)$	2×1	$D_4 \times A_2$	D_4	2×192
$E_6 \times A_1$	$A_3 \times A_1(20)$	2×1	$A_5(4)$	D_4	192 + 504
$D_5 \times A_1(3)$	$A_5(4)$	2×1	$D_4 \times A_1(2)$	$D_4 \times A_1(2)$	288 + 450
$D_5 \times A_2$	$A_3^2(2)$	2×2	$D_4 \times A_1(2)$	D_4	576 + 1224
D_6	$A_3 \times A_2(10)$	2×2	D_4	D_4	1152 + 3366
D_6	$A_2^2 \times A_1^2(2)$	2×4	$E_6 \times A_1$	$A_2^2 \times A_1(8)$	3×1
$E_6 \times A_1$	$A_2^2(8)$	2×6	D_6	$A_3 \times A_2^1(10)$	3×4
$A_4 \times A_3$	$D_4 \times A_2$	2×6	$D_4 \times A_2$	$A_5 \times A_1(3)$	3×6
$A_4 \times A_3$	$A_{5}(4)$	2×6	$D_4 \times A_2$	$D_4 \times A_2$	$2 \times 12 + 20$
E_6	$A_2^2(8)$	2×12	$A_5(4)$	$A_5(4)$	$2 \times 12 + 20 + 36$
D_6	$A_3 \times A_1(20)$	16 + 28	$A_5 \times A_1(3)$	$A_5(4)$	5×6
$D_4 \times A_2$	$A_{5}(4)$	12 + 37	$A_4 \times A_3$	$A_5 \times A_1(3)$	6×1
$A_5 \times A_1(3)$	$D_4 \times A_1(2)$	36 + 60	$A_4 \times A_3$	$A_4 \times A_3$	7×1
E_6	$A_3(7)$	65 + 80	$A_5 \times A_1(3)$	$A_5 \times A_1(3)$	12×1
$D_4 \times A_2$	$D_4 \times A_1(2)$	72 + 96			

Table 3. Disconnected systems of E_8

When $\Phi = E_8$, Table 3 shows that a disconnected system could contain 2, 3, 4, 5, 6, 7 or 12 blocks. It has at most 4518 simple modules (see (E_8, D_4, D_4)). We say a system is semisimple if each of its block has exactly one simple module. 9 of these conjugate pairs corresponding to disconnected semisimple systems, while 9 conjugate pairs corresponding to connected semisimple systems. All of them are listed in Table 4.

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
E_8	Ø	1	E_7	$A_1^3(21)$	2
E_7	$A_2(7)$	1	D_7	$A_2 \times A_1^3(8)$	2
D_7	$A_2^2(8)$	1	A_7	$A_3 \times A_2 \times A_1(4)$	2
$A_6 \times A_1$	$A_4 \times A_2 \times A_1$	1	$E_6 \times A_1$	$A_3 \times A_1(20)$	2
$D_5 \times A_2$	$A_4 \times A_2(4)$	1	$D_5 \times A_1(3)$	$A_{5}(4)$	2
E_6	D_4	1	$E_6 \times A_1$	$A_2^2 \times A_1(8)$	3
D_6	$A_4(6)$	1	$A_4 \times A_3$	$A_5 \times A_1(3)$	6
$A_6(3)$	$D_4 \times A_2$	1	$A_4 \times A_3$	$A_4 \times A_3$	7
$D_5(2)$	$D_5(2)$	1	$A_5 \times A_1(3)$	$A_5 \times A_1(3)$	12

Table 4. Semisimple systems of E_8

When $\Phi = E_7$, Table 5 shows that a disconnected system could contain 2 or 3 blocks. It has at most 150 simple modules (see (E_7, D_4, A_3)). 9 of these conjugate pairs correspond to disconnected semisimple systems, while 5 conjugate pairs correspond to connected semisimple systems. All of them are listed in Table 6.

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
D_6	$(A_1^3)'(10)$	2×1	$D_4 \times A_1$	$(A_3 \times A_1)'(9)$	8 + 12
$A_5 \times A_1$	$(A_3 \times A_1)'(9)$	2×1	$A_4(5)$	$A_4(5)$	2×10
$(A_5)'(2)$	$A_3 \times A_1^2(3)$	2×1	D_4	$(A_3 \times A_1)'(9)$	2×24
$A_4 \times A_1(5)$	$A_4 \times A_1(5)$	2×1	$D_4 \times A_1$	$A_{3}(6)$	24 + 33
$(A_5)'(2)$	$A_2^2 \times A_1(3)$	2×2	D_4	$A_3(6)$	48 + 102
$A_3 \times A_2(3)$	$D_4 \times A_1$	2×2	$A_5 \times A_1$	$A_2^2 \times A_1(3)$	3×1
$A_4 \times A_1(5)$	$A_4(5)$	2×3	$(A_5)'(2)$	$(A_3 \times A_1)'(9)$	3×2
$(A_5)'(2)$	$A_{3}(6)$	8 + 12			

Table 5. Disconnected systems of E_7

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
E_7	Ø	1	$D_4 \times A_1$	$A_4(5)$	1
E_6	$(A_1^3)''$	1	$A_4 \times A_2$	$A_3 \times A_2 \times A_1$	1
D_6	$A_2(6)$	1	D_6	$(A_1^3)'(10)$	2
A_6	$A_2 \times A_1^3$	1	$A_5 \times A_1$	$(A_3 \times A_1)'(9)$	2
$D_5 \times A_1$	$A_2^2(4)$	1	$(A_5)'(2)$	$A_3 \times A_1^2(3)$	2
$D_5(2)$	$(A_3 \times A_1)''(2)$	1	$A_4 \times A_1(5)$	$A_4 \times A_1(5)$	2
$(A_5)''$	D_4	1	$A_5 \times A_1$	$A_2^2 \times A_1(3)$	3

Table 6. Semisimple systems of E_7

When $\Phi=E_6$, Table 5 shows that a disconnected system could contain 2 or 3 blocks. It has at most 25 simple modules (see (E_6,A_3,A_3)). 6 of these conjugate pairs correspond to disconnected semisimple systems, while 4 conjugate pairs correspond to connected sesimple systems. All of them are listed in Table 6.

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
A_5	$A_1^3(5)$	2×1	$A_3(5)$	$A_{3}(5)$	8 + 17
$A_2^2 \times A_1$	$A_3 \times A_1(4)$	2×1	$A_2^2 \times A_1$	$A_2^2 \times A_1$	3×1
$A_3 \times A_1(4)$	$A_3(5)$	2×4	$A_3 \times A_1(4)$	$A_3 \times A_1(4)$	3×1
D_4	$A_2(5)$	2×6			

Table 7. Disconnected systems of E_6

Φ_I	Φ_J	blocks	Φ_I	Φ_J	blocks
E_6	Ø	1	$A_4(4)$	$A_3(5)$	1
$D_5(2)$	$A_1^2(10)$	1	A_5	$A_1^3(5)$	2
A_5	$A_2(5)$	1	$A_2^2 \times A_1$	$A_3 \times A_1(4)$	2
$A_4 \times A_1(2)$	$A_2 \times A_1^2(5)$	1	$A_2^2 \times A_1$	$A_2^2 \times A_1$	3
D_4	A_2^2	1	$A_3 \times A_1(4)$	$A_3 \times A_1(4)$	3

Table 8. Semisimple systems of E_6

References

- [B] E. Backlin, Koszul duality for parabolic and singular category \mathcal{O} . Represent. Theory $\mathbf{3}(1999),\ 139-152.$
- [BX] Z. Q. Bai and W. Xiao, Irreducibility of generalized Verma modules for hermitian symmetric pairs, submitted.
- [BGS] A. Beilinson, V. Ginzburg and W. Soergel, Koszul duality patterns in representation theory, J. Amer. Math. Soc. 9 (1996), 473-527.
- [BGG] Bernstein, I. N., Gelfand, I. M., Gelfand, S. I.: Structure of representations generated by vectors of highest weight. Functional Analysis Appl. 5, 1–9 (1971)
- [Bo] B. D. Boe, Homomorphisms between generalized Verma modules, Trans. Amer. Math. Soc. 288(1985), 791-799
- [BC] B. D. Boe and D. Collingwood, A comparison theory for the structure of induced representations, J. Algebra 94(1985), 511-545.
- [BEJ] B. D. Boe, T. J. Enright , and B. SHELTON , Determination of the intertwining operators for holomorphically induced representations of Hermitian symmetric pairs, Pacific J. Math. 131 (1988), 39-50.
- [BN] B. D. Boe and L. Nakano, Representation type of the blocks of category \mathcal{O}_s , Adv. in Math., 196 (2005) 193-256.
- [Br] J. Brundan, Centers of degenerate cyclotomic Hecke algebras and parabolic category \mathcal{O} , Represent. Theory, 12(2008), 236-259.
- [CM] D. H. Collingwood and W. M. McGovern, Nilpotent orbits in semisimple Lie algebrash, Van Nostrand Reinhold Mathematics Series. Van Nostrand Reinhold Co., New York, 1993.
- [ES] T. J. Enright and B. Shelton, Categories of highest weight modules: applications to classical Hermitian symmetric pairs, Mem. Amer. Math. Soc. 67(1987), no. 367
- [He] H. He, On the reducibility of scalar generalized Verma modules of abelian type, Algebr. Represent. Theory 19 (2016), no. 1, 147-170.
- [HKZ] H. He, T. Kubo and R. Zierau, On the reducibility of scalar generalized Verma modules associated to maximal parabolic subalgebras, to appear in Kyoto. J. Math.
- [HX] J. Hu and W. Xiao, Jantzen coefficients and radical filtrations of generalized Verma modules, preprint.
- [H1] J. Humphreys, Introduction to Lie algebras and representation theory, Springer-Verlag, New York, 1972.
- [H2] J. Humphreys, Representations of Semisimple Lie Algebras in the BGG Category O, GSM. 94, Amer. Math. Soc. Providence, 2008.
- [I] R. S. Irving, Singular blocks of the category \mathcal{O} , Math. Z. 204(1990), 209-224.
- [J1] J. C. Jantzen, Moduln mit einem höchsten Gewicht. Lecture Notes in Mathematics, 750. Springer, Berlin, 1979
- [J2] J. C. Jantzen, Kontravariante Formen auf induzierten Darstellungen halbeinfacher Lie-Algebren, Math. Ann. 226(1977), 53-65.
- [KL] D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, Invent. Math. 53 (1979) 165-184.
- [Kr] H. Kraft, Parametrisierung von Konjugationsklassen in \mathfrak{sl}_n , Math. Ann. 234 (1978), no. 3, 209-220.

- [Ku] T. Kubo, Conformally invariant systems of differential operators associative to two-step nilpotent maximal parabolics of non-Heisenberg type, thesis (Ph.D.)-Oklahoma State University. 2012.
- [LX] S. Leonard and N. Xi, Some non-trivial Kazhdan-Lusztig coefficients of an affine Weyl group of type \tilde{A}_n , Sci. China Math. 53(2010), no. 8, 1919-1930.
- [L1] J. Lepowsky, Conical vectors in induced modules, Trans. Amer. Math. Soc. 208(1975), 219-272.
- [L2] J. Lepowsky, Existence of conical vectors in induced modules, Ann. of Math. (2) 102(1975), 17-40.
- [Lu] G. Lusztig, Cells in affine Weyl groups, in: Algebraic Groups and Related Topics, in: Adv. Stud. Pure Math., vol. 6, KinokuniaCNorth-Holland, 1985, 255-287.
- [M1] H. Matumoto, The homomorphisms between scalar generalized Verma modules associated to maximal parabolic subalgebras, Duke Math. J. 131(2006), no. 1, 75-118.
- [M2] H. Matumoto, On the homomorphisms between scalar generalized Verma modules. Compos. Math. 150 (2014), no. 5, 877-892.
- [M3] H. Matumoto, Homomorphisms between scalar generalized Verma modules for gl(n,C). Int. Math. Res. Not. no. 12(2016),3525-3547.
- [P1] K. J. Platt, Representation type of the blocks of category \mathcal{O}_S in types F_4 and G_2 , J. Algebra 322 (2009), no. 11, 3823-3838.
- [P2] K. J. Platt, Nonzero Infinitesimal Blocks of Category \mathcal{O}_S , Algebr Represent Theory 14 (2011), no. 4, 665-689.
- [R] A. Rocha-Caridi, Splitting criteria for g-modules induced from a parabolic and the Bernstein-Gelfand-Gelfand resolution of a finite-dimensional, irreducible g-module, Trans. Amer. Math. Soc. 262 (1980), 335-366.
- [So] W. Soergel, Kategorie O, Perverse Garben Und Moduln Uber Den Koinvariantez Zur Weylgruppe, J. Amer. Math. Soc. 3 (1990) 421-445.
- [Xi] N. Xi, The leading coefficient of certain Kazhdan-Lusztig polynomials of the permutation group, J. Algebra, 285(2005), no. 1, 136-145.
- [X1] W. Xiao, Leading weight vectors and homomorphisms between generalized Verma modules, J. Algebra, 430(2015), 62-93.
- [X2] W. Xiao, Jantzen coefficients and Blocks of the category $\mathcal{O}^{\mathfrak{p}}$, preprint
- [XZ] W. Xiao and A. Zhang, Basic generalized Verma modules and Jantzen's simplicity criterion, preprint
- (Hu) College of Computer Science and Software Engineering, Shenzhen University, Shenzhen, 518060, Guangdong, P. R. China

 $E ext{-}mail\ address: hujieren97@qq.com}$

(Xiao) College of Mathematics and Statistics, Shenzhen Key Laboratory of Advanced Machine Learning and Applications, Shenzhen University, Shenzhen, 518060, Guangdong, P. R. China

E-mail address: xiaow@szu.edu.cn

(Zhang) College of Mathematics and Statistics, Shenzhen Key Laboratory of Advanced Machine Learning and Applications, Shenzhen University, Shenzhen, 518060, Guangdong, P. R. China

E-mail address: az304@szu.edu.cn