GENERALIZATION OF THE CONWAY-GORDON THEOREM AND INTRINSIC LINKING ON COMPLETE GRAPHS

HIROKO MORISHITA AND RYO NIKKUNI

This article is dedicated to Professor Yoshiyuki Ohyama on his 60th birthday.

ABSTRACT. Conway and Gordon proved that for every spatial complete graph on six vertices, the sum of the linking numbers over all of the constituent two-component links is odd, and Kazakov and Korablev proved that for every spatial complete graph with arbitrary number of vertices greater than six, the sum of the linking numbers over all of the constituent two-component Hamiltonian links is even. In this paper, we show that for every spatial complete graph whose number of vertices is greater than six, the sum of the square of the linking numbers over all of the two-component Hamiltonian links is determined explicitly in terms of the one over all of the triangle-triangle constituent links. This gives an integral lift of the Kazakov-Korablev theorem.

1. INTRODUCTION

Throughout this paper we work in the piecewise linear category. Let G be a finite graph. An embedding f of G into \mathbb{R}^3 is called a *spatial embedding* of G, and the image f(G) is called a *spatial graph* of G. Two spatial embeddings f and g of G are said to be *equivalent* if there exists a self homeomorphism Φ on \mathbb{R}^3 such that $\Phi(f(G)) = g(G)$. Let γ be a subgraph of G homeomorphic to the circle. We call γ a *cycle* of G. For a cycle γ (resp. a disjoint union of cycles λ) and a spatial embedding f of G, we call $f(\gamma)$ (resp. $f(\lambda)$) a *constituent knot* (resp. a *constituent link*) of the spatial graph f(G). In particular, we say that a constituent knot (or link) of a spatial graph f(G) is *Hamiltonian* if it contains all vertices of f(G).

Let K_n be the *complete graph* on *n* vertices, that is the graph consisting of *n* vertices $1, 2, \ldots, n$ such that each pair of its distinct vertices *i* and *j* is connected by exactly one edge ij. Then the following fact is well-known as the *Conway-Gordon* theorems.

Theorem 1.1. (Conway-Gordon [3])

- (1) For any spatial embedding f of K_6 , the sum of the linking numbers over all of the constituent 2-component Hamiltonian links of $f(K_6)$ is odd.
- (2) For any spatial embedding f of K_7 , the sum of the second coefficients of the Conway polynomials over all of the constituent Hamiltonian knots of $f(K_7)$ is odd.

²⁰¹⁰ Mathematics Subject Classification. Primary 57M15; Secondary 57M25.

Key words and phrases. Spatial graphs, Conway-Gordon theorems.

The second author was supported by JSPS KAKENHI Grant Numbers JP15K04881 and $19\mathrm{K}03500.$

Theorem 1.1 also implies that every spatial graph of K_6 contains a nonsplittable 2-component link, and every spatial graph of K_7 contains a nontrivial knot. Furthermore, the authors gave in [7] a generalized Conway-Gordon theorem for K_n with arbitrary number of vertices $n \ge 6$ as follows, where the case of n = 6 has been proven by the second author in [8, Theorem 1.3].

Theorem 1.2. (Morishita-Nikkuni [7, Theorem 1.3]) Let $n \ge 6$ be an integer. For any spatial embedding f of K_n , we have

$$\sum_{\gamma \in \Gamma_n(K_n)} a_2(f(\gamma)) - (n-5)! \sum_{\gamma \in \Gamma_5(K_n)} a_2(f(\gamma))$$
$$= \frac{(n-5)!}{2} \left(\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 - \binom{n-1}{5} \right),$$

where lk denotes the linking number and a_2 denotes the second coefficient of the Conway polynomial.

Moreover, it has been also shown in [7] that for a spatial embedding f of K_n with $n \ge 7$, the sum of a_2 over all of the constituent Hamiltonian knots of $f(K_n)$ modulo (n-5)! does not depend on how to choice f and the modulo (n-5)! reduction has been given explicitly as follows.

Corollary 1.3. ([7, Corollary 1.5]) Let $n \ge 7$ be an integer. For any spatial embedding f of K_n , we have the following congruence modulo (n-5)!:

$$\sum_{\gamma \in \Gamma_n(K_n)} a_2(f(\gamma)) \equiv \begin{cases} -\frac{(n-5)!}{2} \binom{n-1}{5} & (n \equiv 0 \pmod{8}) \\ 0 & (n \not\equiv 0, 7 \pmod{8}) \\ \frac{(n-5)!}{2} \binom{n}{6} & (n \equiv 7 \pmod{8}). \end{cases}$$

On the other hand, for linking numbers of constituent 2-component Hamiltonian links of a spatial graph of K_n with $n \ge 7$, the followings are known.

- **Theorem 1.4.** (1) (Vesnin-Litvintseva [11]) Let $n \ge 7$ be an integer. For any spatial embedding f of K_n , there exists a constituent 2-component Hamiltonian link of $f(K_n)$ with odd linking number.
 - (2) (Kazakov-Korablev [6]) Let $n \ge 7$ be an integer. For any spatial embedding f of K_n , the sum of the linking numbers over all of the constituent 2-component Hamiltonian links of $f(K_n)$ is even.

Our purposes in this paper are to refine Theorem 1.4 (2) by giving its integral lift in terms of the square of the linking number and to investigate the behavior of the nonsplittable 2-component Hamiltonian links in a spatial complete graph. In the following, we call a cycle of G containing exactly p edges a p-cycle of G and denote the set of all p-cycles of G by $\Gamma_p(G)$. Moreover, we denote the set of all pairs of two disjoint cycles of G consisting of a p-cycle and a q-cycle by $\Gamma_{p,q}(G)$. For a spatial embedding f of G and an element λ in $\Gamma_{p,q}(G)$, we also call $f(\lambda)$ a constituent 2-component link of type (p,q). Then we have the following, where the case of p = 4, q = 3 has already been obtained in [7, Theorem 2.3] (see also (2.19)). **Theorem 1.5.** Let $n \ge 6$ be an integer and $p, q \ge 3$ two integers satisfying n = p + q. For any spatial embedding f of K_n , we have

(1.1)
$$\sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 = \begin{cases} (n-6)! \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 & (p=q) \\ 2 \cdot (n-6)! \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 & (p \neq q). \end{cases}$$

In particular, we also have

(1.2)
$$\sum_{p+q=n} \sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 = (n-5)! \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Note that Theorem 1.4 (2) can be recovered by taking the modulo two reduction of (1.2), namely (1.2) gives an integral lift of Theorem 1.4 (2). Since K_n contains K_6 if $n \ge 6$, by Theorem 1.1 (1) we have $\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 > 0$. Therefore (1.1) implies that every spatial graph of K_n with $n \ge 6$ contains a constituent 2-component Hamiltonian link of any type (p, q) with nonzero linking number. In that sense, (1.1) is also a refinement of Theorem 1.4 (1).

By Theorem 1.5, we obtain the following congruences as a corollary, which are remarkable generalizations of Theorem 1.4 (2).

Corollary 1.6. Let $n \ge 6$ be an integer and $p, q \ge 3$ two integers satisfying n = p + q. Let f be a spatial embedding of K_n .

(1) If p = q, then we have the following congruence modulo $2 \cdot (n-6)!$:

$$\sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 \equiv \begin{cases} \frac{n!}{6!} & (n \equiv 6, 7 \pmod{8}) \\ 0 & (n \not\equiv 6, 7 \pmod{8}). \end{cases}$$

(2) If $p \neq q$, then we have the following congruence modulo $4 \cdot (n-6)!$:

$$\sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 \equiv \begin{cases} 2 \cdot \frac{n!}{6!} & (n \equiv 6, 7 \pmod{8}) \\ 0 & (n \not\equiv 6, 7 \pmod{8}). \end{cases}$$

(3) We have the following congruence modulo $2 \cdot (n-5)!$:

$$\sum_{p+q=n}\sum_{\lambda\in\Gamma_{p,q}(K_n)}\mathrm{lk}(f(\lambda))^2\equiv\begin{cases} (n-5)\cdot\frac{n!}{6!} & (n\equiv 6,7\pmod{8})\\ 0 & (n\not\equiv 6,7\pmod{8}). \end{cases}$$

Moreover, by Theorem 1.5, we also obtain the following inequalities as a corollary, where the case of p = q = 3 has already been obtained from Theorem 1.1 (1), and the case of p = 3, q = 4 has also been observed in [5, Theorem 1].

Corollary 1.7. Let $n \ge 6$ be an integer and $p, q \ge 3$ two integers satisfying n = p + q. For any spatial embedding f of K_n , we have

(1.3)
$$\sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 \ge \begin{cases} \frac{n!}{6!} & (p=q)\\ 2 \cdot \frac{n!}{6!} & (p \neq q) \end{cases}$$

In particular, we have

(1.4)
$$\sum_{p+q=n} \sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 \ge (n-5) \cdot \frac{n!}{6!}.$$

The lower bound of each of the inequalities in Corollary 1.7 is sharp, see Remark 3.3. We also remark here that the minimum number of nonsplittable constituent 2-component links of a spatial graph of K_n with $n \leq 9$ has been investigated by Fleming-Mellor [5] and Abrams-Mellor-Trott [1]. It can be said that (1.4) gives an algebraic estimation from below of the number of nonsplittable constituent 2-component Hamiltonian links of a spatial graph of K_n with $n \geq 6$ (see also Remark 3.7).

In addition, Corollary 1.7 also gives an information of the maximum value of the absolute value of the linking numbers over all of the 2-component Hamiltonian links of type (p, q) of a spatial graph of K_n with $n \ge 6$.

Corollary 1.8. Let $n \ge 6$ be an integer and $p, q \ge 3$ two integers satisfying n = p + q. For any spatial embedding f of K_n , we have

$$\max_{\lambda \in \Gamma_{p,q}(K_n)} |\mathrm{lk}(f(\lambda))| \ge \frac{\sqrt{pq}}{3\sqrt{10}}$$

Corollary 1.8 says that if n is sufficiently large then every spatial graph of K_n contains a 2-component Hamiltonian link of type (p, q), the absolute value of whose linking number is arbitrary large. Actually we have the following.

Corollary 1.9. Let $p, q \geq 3$ be two integers. For a spatial embedding f of K_{p+q} and a positive integer m, if $pq > 90(m-1)^2$ then there exists a pair of disjoint cycles $\lambda \in \Gamma_{p,q}(K_{p+q})$ such that $|\text{lk}(f(\lambda))| \geq m$.

Shirai-Taniyama showed in [10] that for any spatial embedding f of $K_{6\cdot2^k}$, there exists a pair of disjoint cycles $\lambda \in \Gamma_{3\cdot2^k,3\cdot2^k}(K_{6\cdot2^k})$ such that $|\text{lk}(f(\lambda))| \ge 2^k$. Since for every positive integer m, there exists an integer k such that $2^{k-1} < m \le 2^k$, they also showed that $K_{12m} \supset K_{6\cdot2^k}$ and therefore for any spatial embedding f of K_{12m} , there exists a pair of disjoint cycles $\lambda \in \Gamma_{3\cdot2^k,3\cdot2^k}(K_{12m})$ such that $|\text{lk}(f(\lambda))| \ge 2^k \ge m$. The new knowledge obtained in Corollary 1.9 is the fact that for any positive integer m, there exists a positive integer n such that every spatial graph of K_n contains a 2-component Hamiltonian link of any type, the absolute value of whose linking number is greater than or equal to m.

The paper is organized as follows. In Section 2, we prove Theorem 1.5 in the case of n = 8. In Section 3, we prove Theorem 1.5 in general case and Corollaries 1.6, 1.7, 1.8 and 1.9. At the end of the paper, we add an appendix, in which we give similar results as Corollaries 1.8 and 1.9 for the maximum value of a_2 over all of the Hamiltonian knots of a rectilinear spatial graph of K_n .

2. Proof of Theorem 1.5: n = 8

First we show Theorem 1.5 in the case of n = 8. Namely we prove the following.

Theorem 2.1. For any spatial embedding f of K_8 , we have

(2.1)
$$\sum_{\lambda \in \Gamma_{4,4}(K_8)} \operatorname{lk}(f(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{3,3}(K_8)} \operatorname{lk}(f(\lambda))^2,$$

(2.2)
$$\sum_{\lambda \in \Gamma_{3,5}(K_8)} \operatorname{lk}(f(\lambda))^2 = 4 \sum_{\lambda \in \Gamma_{3,3}(K_8)} \operatorname{lk}(f(\lambda))^2.$$

We show four lemmas which are needed to prove Theorem 2.1. In the following, we introduce some subgraphs of K_n with $n \ge 4$ which are used in these proofs (and

§3). We denote the edge of K_n connecting two distinct vertices i and j by ij, and denote a path of length 2 of K_n consisting of two edges ij and jk by ijk. We denote the subgraph of K_n obtained from K_n by deleting the vertex m and all of the edges incident to m by $K_{n-1}^{(m)}$ (m = 1, 2, ..., n). Actually $K_{n-1}^{(m)}$ is isomorphic to K_{n-1} for any m. For $1 \leq i < j \leq n$ and $i, j \neq m$, let $F_{ij}^{(m)}$ be the subgraph of K_n obtained from K_n by deleting the edges ij and mk for all k with $1 \leq k \leq n, k \neq i, j$. Note that $F_{ij}^{(m)}$ is homeomorphic to K_{n-1} , namely $F_{ij}^{(m)}$ is obtained from $K_{n-1}^{(m)}$ by subdividing the edge ij by the vertex m, see Fig. 2.1.



FIGURE 2.1. $K_{n-1}^{(m)}$ and $F_{ij}^{(m)}$ (n = 8)

Moreover, we denote a path of length 3 that is not a 3-cycle of K_n consisting of three mutually distinct edges ik, kl and lj by iklj. For two distinct vertices k and l of K_n , we denote the subgraph of K_n obtained from K_n by deleting the vertices k, l and all of the edges incident to k, l by $K_{n-2}^{(kl)} (= K_{n-2}^{(lk)})$. Actually $K_{n-2}^{(kl)}$ is isomorphic to K_{n-2} for any k, l. For two distinct vertices i, j of K_n with $i, j \neq k, l$, let $F_{ij}^{(kl)}$ be the subgraph of K_n obtained from K_n by deleting the edges ij, $\overline{km} (1 \leq m \leq n, m \neq i, l)$ and $\overline{lm'} (1 \leq m' \leq n, m' \neq j, k)$, and $F_{ij}^{(lk)}$ the subgraph of K_n obtained from K_n by deleting the edges $ij, \overline{km} (1 \leq m \leq n, m \neq j, l)$ and $\overline{lm'} (1 \leq m' \leq n, m' \neq i, k)$. Note that both $F_{ij}^{(kl)}$ and $F_{ij}^{(lk)}$ are obtained from $K_{n-2}^{(kl)}$ by subdividing the edge \overline{ij} by the vertices k, l, see Fig 2.2.



FIGURE 2.2. $K_{n-2}^{(kl)}, F_{ij}^{(kl)}$ and $F_{ij}^{(lk)}$ (n=8)

Lemma 2.2. For any spatial embedding f of K_8 , we have

$$\sum_{\gamma \in \Gamma_{7}(K_{8})} a_{2}(f(\gamma)) - 6 \sum_{\gamma \in \Gamma_{5}(K_{8})} a_{2}(f(\gamma)) = 2 \sum_{\lambda \in \Gamma_{3,3}(K_{8})} \operatorname{lk}(f(\lambda))^{2} - 48.$$

Proof. Let f be a spatial embedding of K_8 . Then by applying Theorem 1.2 in the case of n = 7 to the embedding f restricted to $K_7^{(m)}$ (m = 1, 2, ..., 8), we have

$$(2.3)\sum_{\gamma\in\Gamma_7(K_7^{(m)})}a_2(f(\gamma))-2\sum_{\gamma\in\Gamma_5(K_7^{(m)})}a_2(f(\gamma))=\sum_{\lambda\in\Gamma_{3,3}(K_7^{(m)})}\mathrm{lk}(f(\lambda))^2-6.$$

Let us take the sum of both sides of (2.3) for all m. Since for each t-cycle γ of K_8 is shared by exactly $(8-t) K_7^{(m)}$'s, we have

(2.4)
$$\sum_{m=1}^{\circ} \sum_{\gamma \in \Gamma_t(K_7^{(m)})} a_2(f(\gamma)) = (8-t) \sum_{\gamma \in \Gamma_t(K_8)} a_2(f(\gamma)).$$

On the other hand, as we will show later in (3.17), we have

(2.5)
$$\sum_{m=1}^{8} \sum_{\lambda \in \Gamma_{3,3}(K_7^{(m)})} \operatorname{lk}(f(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{3,3}(K_8)} \operatorname{lk}(f(\lambda))^2.$$

By combining (2.4) and (2.5) with (2.3), we have the result.

Lemma 2.3. For any spatial embedding f of K_8 , we have

$$16\sum_{\gamma\in\Gamma_{8}(K_{8})}a_{2}(f(\gamma)) - 14\sum_{\gamma\in\Gamma_{7}(K_{8})}a_{2}(f(\gamma)) - 12\sum_{\gamma\in\Gamma_{5}(K_{8})}a_{2}(f(\gamma))$$

= $5\sum_{\lambda\in\Gamma_{3,5}(K_{8})}\operatorname{lk}(f(\lambda))^{2} - 336.$

Proof. Let f be a spatial embedding of K_8 . Then by applying Theorem 1.2 in the case of n = 6 to the embedding f restricted to $F_{ij}^{(kl)}$ $(1 \le i < j \le 8, i, j \ne k, l)$, we have

$$(2.6) \qquad 2\left(\sum_{\substack{\gamma \in \Gamma_{8}(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} a_{2}(f(\gamma)) + \sum_{\substack{\gamma \in \Gamma_{6}(F_{ij}^{(kl)}) \\ iklj \not \not \subset \gamma}} a_{2}(f(\gamma)) \right) \\ - \sum_{\substack{\gamma \in \Gamma_{7}(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} a_{2}(f(\gamma)) - \sum_{\substack{\gamma \in \Gamma_{5}(F_{ij}^{(kl)}) \\ iklj \not \not \subset \gamma}} a_{2}(f(\gamma))\right) \\ = \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{5}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_{3}(F_{ij}^{(kl)})}} \operatorname{lk}(f(\lambda))^{2} + \sum_{\lambda \in \Gamma_{3,3}(F_{ij}^{(kl)}) \\ klj \subset \gamma} \operatorname{lk}(f(\lambda))^{2} - 1.$$

Note that by applying Theorem 1.2 in the case of n = 6 to the embedding f restricted to $F_{ij}^{(lk)}$, we also have a similar formula as (2.6). Let us take the sum of both side of (2.6) over $1 \le i < j \le 8$ and $i, j \ne k, l$. For a $t(\ge 4)$ -cycle γ of K_8 containing \overline{kl} , let i and j be the two vertices of K_8 which are adjacent to \overline{kl} in γ ($1 \le i < j \le 8$). Then γ is a t-cycle of $F_{ij}^{(lk)}$ or $F_{ij}^{(lk)}$. This implies that

$$(2.7) \sum_{\substack{1 \le i < j \le 8\\ i,j \ne k,l}} \left(\sum_{\substack{\gamma \in \Gamma_t(F_{ij}^{(kl)})\\ iklj \subset \gamma}} a_2(f(\gamma)) + \sum_{\substack{\gamma \in \Gamma_t(F_{ij}^{(lk)})\\ ilkj \subset \gamma}} a_2(f(\gamma)) \right) = \sum_{\substack{\gamma \in \Gamma_t(K_8)\\ \overline{kl} \subset \gamma}} a_2(f(\gamma)).$$

For a t-cycle γ of $K_6^{(kl)}$, let \overline{ij} be an edge of $K_6^{(kl)}$ which is not contained in γ . Then γ is a t-cycle of $F_{ij}^{(kl)}$ (resp. $F_{ij}^{(lk)}$) which does not contain \overline{iklj} (resp. \overline{ilkj}). Note that there are (15-t) ways to to choose such a pair of i and j. This implies that

(2.8)
$$\sum_{\substack{1 \le i < j \le 8\\ i, j \ne k, l}} \sum_{\substack{\gamma \in \Gamma_t(F_{ij}^{(kl)})\\ \frac{\gamma + 1}{2kl \neq \alpha < \gamma}}} a_2(f(\gamma)) = (15-t) \sum_{\gamma \in \Gamma_t(K_6^{(kl)})} a_2(f(\gamma)),$$

(2.9)
$$\sum_{\substack{1 \le i < j \le 8\\i,j \ne k,l}} \sum_{\substack{\gamma \in \Gamma_t(F_{ij}^{(lk)})\\ilkj \not \subset \gamma}} a_2(f(\gamma)) = (15-t) \sum_{\gamma \in \Gamma_t(K_6^{(kl)})} a_2(f(\gamma)).$$

On the other hand, as we will show later in (3.33), (3.36) and (3.37), we have

$$(2.10) \sum_{\substack{1 \leq i < j \leq 8\\ i,j \neq k, \overline{l}}} \left(\sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(F_{ij}^{(kl)})\\\gamma \in \Gamma_5(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_3(F_{ij}^{(kl)})\\\overline{iklj} \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(F_{ij}^{(lk)})\\\overline{iklj} \subset \gamma}} \operatorname{lk}(f(\lambda))^2 \right)$$
$$= \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(K_8)\\\gamma \in \Gamma_5(K_8), \ \gamma' \in \Gamma_3(K_8)\\\overline{kl} \subset \gamma}} \operatorname{lk}(f(\lambda))^2,$$

(2.11)
$$\sum_{\substack{1 \le i < j \le 8\\ i, j \ne k, l}} \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(kl)})\\ ikl j \not \subset \gamma}} \operatorname{lk}(f(\lambda))^2 = 9 \sum_{\lambda \in \Gamma_{3,3}(K_6^{(kl)})} \operatorname{lk}(f(\lambda))^2,$$

(2.12)
$$\sum_{\substack{1 \le i < j \le 8\\ i, j \ne k, l}} \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(lk)})\\ \overline{ilkj} \not \subset \gamma}} \operatorname{lk}(f(\lambda))^2 = 9 \sum_{\substack{\lambda \in \Gamma_{3,3}(K_6^{(kl)})}} \operatorname{lk}(f(\lambda))^2.$$

By combining (2.7), (2.8), (2.9), (2.10), (2.11) and (2.12) with (2.6), we have

$$(2.13) \qquad 2 \sum_{\substack{\gamma \in \Gamma_{8}(K_{8}) \\ \overline{kl} \subset \gamma}} a_{2}(f(\gamma)) + 36 \sum_{\gamma \in \Gamma_{6}(K_{6}^{(kl)})} a_{2}(f(\gamma)) \\ -2 \sum_{\substack{\gamma \in \Gamma_{7}(K_{8}) \\ \overline{kl} \subset \gamma}} a_{2}(f(\gamma)) - 40 \sum_{\substack{\gamma \in \Gamma_{5}(K_{6}^{(kl)})}} a_{2}(f(\gamma)) \\ = \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(K_{8}) \\ \gamma \in \Gamma_{5}(K_{8}), \ \gamma' \in \Gamma_{3}(K_{8})}} \operatorname{lk}(f(\lambda))^{2} + 18 \sum_{\substack{\lambda \in \Gamma_{3,3}(K_{6}^{(kl)})}} a_{2}(f(\lambda))^{2} - 30.$$

Then, by applying Theorem 1.2 in the case of n = 6 to the embedding f restricted to $K_6^{(kl)}$, we have

$$(2.14) \quad 36 \sum_{\gamma \in \Gamma_{6}(K_{6}^{(kl)})} a_{2}(f(\gamma)) - 40 \sum_{\gamma \in \Gamma_{5}(K_{6}^{(kl)})} a_{2}(f(\gamma)) \\ = 18 \left(2 \left(\sum_{\gamma \in \Gamma_{6}(K_{6}^{(kl)})} a_{2}(f(\gamma)) - \sum_{\gamma \in \Gamma_{5}(K_{6}^{(kl)})} a_{2}(f(\gamma)) \right) \right) - 4 \sum_{\gamma \in \Gamma_{5}(K_{6}^{(kl)})} a_{2}(f(\gamma)) \\ = 18 \left(\sum_{\lambda \in \Gamma_{3,3}(K_{6}^{(kl)})} a_{2}(f(\lambda))^{2} - 1 \right) - 4 \sum_{\gamma \in \Gamma_{5}(K_{6}^{(kl)})} a_{2}(f(\gamma)).$$

By combining (2.14) with (2.13), we have

$$(2.15) \qquad 2\sum_{\substack{\gamma \in \Gamma_{8}(K_{8})\\\overline{kl} \subset \gamma}} a_{2}(f(\gamma)) - 2\sum_{\substack{\gamma \in \Gamma_{7}(K_{8})\\\overline{kl} \subset \gamma}} a_{2}(f(\gamma)) - 4\sum_{\substack{\gamma \in \Gamma_{5}(K_{6}^{(kl)})\\\overline{kl} \subset \gamma}} a_{2}(f(\gamma))$$
$$= \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(K_{8})\\\gamma \in \Gamma_{5}(K_{8}), \ \gamma' \in \Gamma_{3}(K_{8})\\\overline{kl} \subset \gamma}} \operatorname{lk}(f(\lambda))^{2} - 12.$$

Now we take the sum of both side of (2.15) over $1 \le k < l \le 8$. For a *t*-cycle γ of K_8 , let \overline{kl} be an edge of K_8 which is contained in γ . Note that there are *t* ways to choose such two vertices *k* and *l*. This implies that

(2.16)
$$\sum_{\substack{1 \le k < l \le 8}} \sum_{\substack{\gamma \in \Gamma_t(K_8) \\ \overline{kl} \subset \gamma}} a_2(f(\gamma)) = t \sum_{\gamma \in \Gamma_t(K_8)} a_2(f(\gamma)).$$

For a 5-cycle γ of K_8 , let k and l be two distinct vertices of K_8 which are not contained in γ . Then γ is a 5-cycle of $K_6^{(kl)}$. Note that there are three ways to choose such two vertices k and l. This implies that

(2.17)
$$\sum_{1 \le i < j \le 8} \sum_{\gamma \in \Gamma_5(K_6^{(kl)})} a_2(f(\gamma)) = 3 \sum_{\gamma \in \Gamma_5(K_8)} a_2(f(\gamma)).$$

On the other hand, as we will show later in (3.43), we have

(2.18)
$$\sum_{\substack{1 \le k < l \le 8 \\ \gamma \in \Gamma_5(K_8), \ \gamma' \in \Gamma_3(K_8) \\ \overline{kl} \subset \gamma}} \lim_{k < f(\lambda)^2} |\mathbf{k}(f(\lambda))|^2 = 5 \sum_{\substack{\lambda \in \Gamma_{3,5}(K_8) \\ \lambda \in \Gamma_{3,5}(K_8)}} \lim_{k < f(\lambda)^2} |\mathbf{k}(f(\lambda))|^2.$$

By combining (2.16), (2.17) and (2.18) with (2.15), we have the reslut.

As we mentioned before, (1.1) has already shown if p = 3, q = 4 [7, Theorem 2.3]. Namely for any spatial embedding f of K_7 , we have

(2.19)
$$\sum_{\lambda \in \Gamma_{3,4}(K_7)} \operatorname{lk}(f(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{3,3}(K_7)} \operatorname{lk}(f(\lambda))^2.$$

Moreover, the following has also been shown by the authors in [7, Lemma 2.1 (2)].

Lemma 2.4. For any spatial embedding f of K_n with $n \ge 7$, we have

$$\sum_{\lambda \in \Gamma_{3,4}(K_n)} \operatorname{lk}(f(\lambda))^2 = 2(n-6) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

On the other hand, we also have the following in the case of n = 8.

Lemma 2.5. For any spatial embedding f of K_8 , we have

$$8\sum_{\lambda\in\Gamma_{4,4}(K_8)} \mathrm{lk}(f(\lambda))^2 + 5\sum_{\lambda\in\Gamma_{3,5}(K_8)} \mathrm{lk}(f(\lambda))^2 = 36\sum_{\lambda\in\Gamma_{3,3}(K_8)} \mathrm{lk}(f(\lambda))^2.$$

Proof. For any spatial embedding g of K_7 , by (2.19) we have

$$\sum_{\lambda \in \Gamma_{3,4}(K_7)} \operatorname{lk}(g(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{3,3}(K_7)} \operatorname{lk}(g(\lambda))^2.$$

Then we have the result by setting n = 8, a = 2, t = 3 and s = 1 in Lemma 3.1 which will be shown in §3.

Proof of Theorem 2.1. Let f be a spatial embedding of K_8 . Then by Theorem 1.2 in the case of n = 8, we have

(2.20)
$$\sum_{\gamma \in \Gamma_8(K_8)} a_2(f(\gamma)) - 6 \sum_{\gamma \in \Gamma_5(K_8)} a_2(f(\gamma)) = 3 \sum_{\lambda \in \Gamma_{3,3}(K_8)} \operatorname{lk}(f(\lambda))^2 - 63.$$

On the other hand, by Lemmas 2.2 and 2.3, we have

(2.21)
$$16 \sum_{\gamma \in \Gamma_8(K_8)} a_2(f(\gamma)) - 96 \sum_{\gamma \in \Gamma_5(K_8)} a_2(f(\gamma))$$
$$= 28 \sum_{\lambda \in \Gamma_{3,3}(K_8)} \operatorname{lk}(f(\lambda))^2 + 5 \sum_{\lambda \in \Gamma_{3,5}(K_8)} \operatorname{lk}(f(\lambda))^2 - 1008.$$

Then by combining (2.20) and (2.21), we have (2.2), and by combining Lemma 2.5 and (2.2), we have (2.1). This completes the proof.

In a similar way as Lemma 2.4, we have the following by (2.2).

Lemma 2.6. Let $n \ge 8$ be an integer. For any spatial embedding f of K_n , we have

$$\sum_{\lambda \in \Gamma_{3,5}(K_n)} \operatorname{lk}(f(\lambda))^2 = 2(n-6)(n-7) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

Proof. Note that each pair of two disjoint 3-cycles of K_n is shared by exactly $\binom{n-6}{2}$ subgraphs isomorphic to K_8 if $n \ge 8$. Then by applying (2.2) to the embedding f restricted to each of the subgraphs of K_n isomorphic to K_8 and taking the sum of both sides of this over all of them, we have the result.

3. Proof of Theorem 1.5: General Case

In this section, we show Theorem 1.5 in the case of $n \ge 9$. First we show two lemmas which are needed later.

Lemma 3.1. Let $n \ge 7$ be an integer and t, s two integers satisfying n-1 = 2t+s, where $t \ge 3$ and $s \ge 0$. Assume that there exist a constant a such that

(3.1)
$$\sum_{\lambda \in \Gamma_{t,t+s}(K_{n-1})} \operatorname{lk}(g(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-1})} \operatorname{lk}(g(\lambda))^2$$

for any spatial embedding g of K_{n-1} . Then for any spatial embedding f of K_n , we have

(3.2)
$$\sum_{\lambda \in \Gamma_{t,t+1}(K_n)} \operatorname{lk}(f(\lambda))^2 = 2a(n-6) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if s = 0*,*

(3.3)
$$2(t+1) \sum_{\lambda \in \Gamma_{t+1,t+1}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+2) \sum_{\lambda \in \Gamma_{t,t+2}(K_n)} \operatorname{lk}(f(\lambda))^2$$
$$= a(n-6)(2t+3) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if s = 1 and

(3.4)
$$(t+1)\sum_{\lambda\in\Gamma_{t+1,t+s}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+s+1)\sum_{\lambda\in\Gamma_{t,t+s+1}(K_n)} \operatorname{lk}(f(\lambda))^2 \\ = a(n-6)(2t+s+2)\sum_{\lambda\in\Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if $s \geq 2$.

Proof. Let f be a spatial embedding of K_n . For the embedding f restricted to $F_{ij}^{(m)}$, by the assumption we have

$$(3.5) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t+s}(F_{ij}^{(m)})\\\gamma\in\Gamma_{t+1}(F_{ij}^{(m)}), \ \gamma'\in\Gamma_{t+s}(F_{ij}^{(m)})\\imj\subset\gamma}} \operatorname{lk}(f(\lambda))^{2} + \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t,t+s+1}(F_{ij}^{(m)})\\imj\subset\gamma}} \operatorname{lk}(f(\lambda))^{2} \\ + \sum_{\substack{\lambda\in\Gamma_{t,t+s}(K_{n-1}^{(m)})\\ij\not\in\lambda}} \operatorname{lk}(f(\lambda))^{2} \\ = a\left(\sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,4}(F_{ij}^{(m)})\\\gamma\in\Gamma_{4}(F_{ij}^{(m)}), \ \gamma'\in\Gamma_{3}(F_{ij}^{(m)})\\imj\subset\gamma}} \operatorname{lk}(f(\lambda))^{2} + \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})\\ij\not\in\lambda}} \operatorname{lk}(f(\lambda))^{2}\right).$$

First we assume that $s \ge 1$. Let us take the sum of both side of (3.5) over $1 \le i < j \le n$ and $i, j \ne m$. For a pair of disjoint cycles λ of K_n consisting of a (t+1)-cycle γ which contains the vertex m and a (t+s)-cycle γ' , let i and j be the two vertices of K_n which are adjacent to m in γ . Then λ is a pair of disjoint cycles of $F_{ij}^{(m)}$ consisting of a (t+1)-cycle γ which contains \overline{imj} and a (t+s)-cycle γ' . This implies that

$$(3.6) \sum_{\substack{1 \le i < j \le n \\ i, j \ne m}} \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+1, t+s}(F_{ij}^{(m)}) \\ \gamma \in \Gamma_{t+1}(F_{ij}^{(m)}), \ \gamma' \in \Gamma_{t+s}(F_{ij}^{(m)})}{\frac{1}{imj \subset \gamma}} \operatorname{lk}(f(\lambda))^2 = \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+1, t+s}(K_n) \\ \gamma \in \Gamma_{t+1}(K_n), \ \gamma' \in \Gamma_{t+s}(K_n)}} \operatorname{lk}(f(\lambda))^2.$$

In the same way as (3.6), we also have

$$(3.7) \sum_{\substack{1 \leq i < j \leq n \\ i,j \neq m}} \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t,t+s+1}(F_{ij}^{(m)}) \\ \gamma \in \Gamma_t(F_{ij}^{(m)}), \ \gamma' \in \Gamma_{t+s+1}(F_{ij}^{(m)}) \\ \hline imj \subset \gamma'}} \operatorname{lk}(f(\lambda))^2 = \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t,t+s+1}(K_n) \\ \gamma \in \Gamma_t(K_n), \ \gamma' \in \Gamma_{t+s+1}(K_n) \\ m \subset \gamma'}} \operatorname{lk}(f(\lambda))^2 = \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,4}(K_n) \\ m \subset \gamma'}} \operatorname{lk}(f(\lambda))^2.$$

For a pair of disjoint cycles λ of $K_{n-1}^{(m)}$ consisting of a *t*-cycle and a (t+s)-cycle, let \overline{ij} be an edge of $K_{n-1}^{(m)}$ which is not contained in λ . Note that there are $\binom{n-1}{2} - (2t+s)$ ways to choose such a pair of *i* and *j*. This implies that

$$(3.9)\sum_{\substack{1\leq i< j\leq n\\i,j\neq m}}\sum_{\lambda\in\Gamma_{t,t+s}(K_{n-1}^{(m)})}\operatorname{lk}(f(\lambda))^{2} = \left(\binom{n-1}{2} - (2t+s)\right)\sum_{\lambda\in\Gamma_{t,t+s}(K_{n-1}^{(m)})}\operatorname{lk}(f(\lambda))^{2}.$$

In the same way as (3.9), we also have

$$(3.10)\sum_{\substack{1\leq i< j\leq n\\i,j\neq m}}\sum_{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})\atop ij\not\in\lambda}\operatorname{lk}(f(\lambda))^2 = \left(\binom{n-1}{2} - 6\right)\sum_{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})}\operatorname{lk}(f(\lambda))^2.$$

By combining (3.6), (3.7), (3.8), (3.9) and (3.10) with (3.5), we have

$$(3.11) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t+s}(K_n)\\\gamma\in\Gamma_{t+1}(K_n),\ \gamma'\in\Gamma_{t+s}(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t,t+s}(K_n)\\m\subset\gamma'}} \operatorname{lk}(f(\lambda))^2 \\ + \left(\binom{n-1}{2} - (2t+s)\right) \sum_{\substack{\lambda\in\Gamma_{t,t+s}(K_{n-1}^{(m)})\\\lambda\in\Gamma_{t,t+s}(K_{n-1}^{(m)})}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,4}(K_n)\\\gamma\in\Gamma_4(K_n),\ \gamma'\in\Gamma_3(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + a \left(\binom{n-1}{2} - 6\right) \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 \\ \end{cases}$$

Then for the embedding f restricted to $K_{n-1}^{(m)}$, by the assumption we have

(3.12)
$$\sum_{\lambda \in \Gamma_{t,t+s}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.12) with (3.11), we have

$$(3.13) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t+s}(K_n)\\\gamma\in\Gamma_{t+1}(K_n),\ \gamma'\in\Gamma_{t+s}(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t,t+s+1}(K_n)\\m\subset\gamma'}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,4}(K_n)\\\gamma\in\Gamma_4(K_n),\ \gamma'\in\Gamma_3(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + a(2t+s-6) \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2.$$

Now we assume that $s \geq 2$. Let us take the sum of both sides of (3.13) over m = 1, 2, ..., n. For a pair of disjoint cycles λ of K_n consisting of a (t + 1)-cycle γ and a (t + s)-cycle γ' , let m be a vertex of K_n which is contained in γ . Note that there are t + 1 ways to choose such a vertex m. This implies that

$$(3.14) \sum_{m=1}^{n} \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t+s}(K_n)\\\gamma\in\Gamma_{t+1}(K_n), \ \gamma'\in\Gamma_{t+s}(K_n)\\m\in\gamma}} \operatorname{lk}(f(\lambda))^2 = (t+1) \sum_{\lambda\in\Gamma_{t+1,t+s}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

In the same way as (3.14), we also have

$$(3.15) \quad \sum_{m=1}^{n} \sum_{\substack{\lambda=\gamma \cup \gamma' \in \Gamma_{t,t+s+1}(K_n) \\ \gamma \in \Gamma_t(K_n), \ \gamma' \in \Gamma_{t+s+1}(K_n) \\ m \subset \gamma'}} \operatorname{lk}(f(\lambda))^2 = (t+s+1) \sum_{\lambda \in \Gamma_{t,t+s+1}(K_n)} \operatorname{lk}(f(\lambda))^2,$$

$$(3.16) \quad \sum_{m=1}^{n} \sum_{\substack{\lambda=\gamma \cup \gamma' \in \Gamma_{3,4}(K_n) \\ \gamma \in \Gamma_4(K_n), \ \gamma' \in \Gamma_3(K_n) \\ m \subset \gamma}} \operatorname{lk}(f(\lambda))^2 = 4 \sum_{\lambda \in \Gamma_{3,4}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

For a pair of two disjoint 3-cycles λ of K_n , let m be a vertex of K_n which is not contained in λ . Then λ is a pair of two disjoint 3-cycles of $K_{n-1}^{(m)}$. Note that there are n-6 ways to choose such a vertex m. This implies that

(3.17)
$$\sum_{m=1}^{n} \sum_{\lambda \in \Gamma_{3,3}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2 = (n-6) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

By combining (3.14), (3.15), (3.16) and (3.17) with (3.13), we have

(3.18)
$$(t+1)\sum_{\lambda\in\Gamma_{t+1,t+s}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+s+1)\sum_{\lambda\in\Gamma_{t,t+s+1}(K_n)} \operatorname{lk}(f(\lambda))^2 \\ = 4a\sum_{\lambda\in\Gamma_{3,4}(K_n)} \operatorname{lk}(f(\lambda))^2 + a(2t+s-6)(n-6)\sum_{\lambda\in\Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Then by (3.18) and Lemma 2.4, we have (3.4).

Next, assume that s = 1. Note that we also have

(3.19)
$$\sum_{\substack{m=1 \ \lambda \in \Gamma_{t+1,t+1}(K_n) \\ m \subset \lambda}}^{n} \ln(f(\lambda))^2 = 2(t+1) \sum_{\substack{\lambda \in \Gamma_{t+1,t+1}(K_n) \\ m \subset \lambda}} \ln(f(\lambda))^2$$

in a similar way as (3.14). By combining (3.15), (3.16), (3.17) and (3.19) with (3.13), we have

(3.20)
$$2(t+1) \sum_{\lambda \in \Gamma_{t+1,t+1}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+2) \sum_{\lambda \in \Gamma_{t,t+2}(K_n)} \operatorname{lk}(f(\lambda))^2 \\ = 4a \sum_{\lambda \in \Gamma_{3,4}(K_n)} \operatorname{lk}(f(\lambda))^2 + a(2t-5)(n-6) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Then by (3.20) and Lemma 2.4, we have (3.3).

Finally, assume that s = 0. By (3.5), we have

$$(3.21) \qquad \sum_{\substack{\lambda \in \Gamma_{t+1,t}(F_{ij}^{(m)}) \\ \gamma \in \Gamma_{t+1}(F_{ij}^{(m)}), \ \gamma' \in \Gamma_t(F_{ij}^{(m)}) \\ \overline{imj} \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda \in \Gamma_{1,t}(K_{n-1}^{(m)}) \\ \overline{ij} \not \subset \lambda}} \operatorname{lk}(f(\lambda))^2 \\ = a \Biggl(\sum_{\substack{\lambda \in \Gamma_{3,4}(F_{ij}^{(m)}) \\ \gamma \in \Gamma_4(F_{ij}^{(m)}), \ \gamma' \in \Gamma_3(F_{ij}^{(m)}) \\ \overline{imj} \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda \in \Gamma_{3,3}(K_{n-1}^{(m)}) \\ \overline{ij} \not \subset \lambda}} \operatorname{lk}(f(\lambda))^2 \Biggr).$$

Let us take the sum of both side of (3.21) over $1 \le i < j \le n$ and $i, j \ne m$. Note that (3.6) and (3.9) also hold if s = 0. By combining (3.6), (3.8), (3.9) and (3.10) with (3.21), we have

$$(3.22) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t}(K_n)\\\gamma\in\Gamma_{t+1}(K_n),\ \gamma'\in\Gamma_t(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + \left(\binom{n-1}{2} - 2t\right) \sum_{\substack{\lambda\in\Gamma_{t,t}(K_{n-1}^{(m)})\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,4}(K_n)\\\gamma\in\Gamma_4(K_n),\ \gamma'\in\Gamma_3(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + a \left(\binom{n-1}{2} - 6\right) \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2.$$

Then for the embedding f restricted to $K_{n-1}^{(m)},$ by the assumption we have

(3.23)
$$\sum_{\lambda \in \Gamma_{t,t}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.22) and (3.23), we have

$$(3.24) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+1,t}(K_n)\\\gamma\in\Gamma_{t+1}(K_n),\ \gamma'\in\Gamma_t(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,4}(K_n)\\\gamma\in\Gamma_4(K_n),\ \gamma'\in\Gamma_3(K_n)\\m\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + 2a(t-3) \sum_{\lambda\in\Gamma_{3,3}(K_{n-1}^{(m)})} \operatorname{lk}(f(\lambda))^2.$$

Now we take the sum of both sides of (3.24) over $m = 1, 2, \dots, n$. Note that (3.14) also holds if s = 0. By combining (3.14), (3.16) and (3.17) with (3.24), we have

(3.25)
$$(t+1)\sum_{\lambda\in\Gamma_{t+1,t}(K_n)} \operatorname{lk}(f(\lambda))^2 \\ = 4a\sum_{\lambda\in\Gamma_{3,4}(K_n)} \operatorname{lk}(f(\lambda))^2 + 2a(t-3)(n-6)\sum_{\lambda\in\Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Then by (3.25) and Lemma 2.4, we have (3.2). This completes the proof.

Lemma 3.2. Let $n \ge 8$ be an integer and t, s two integers satisfying n-2 = 2t+s, where $t \ge 3$ and $s \ge 0$. Assume that there exist a constant a such that

(3.26)
$$\sum_{\lambda \in \Gamma_{t,t+s}(K_{n-2})} \operatorname{lk}(g(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-2})} \operatorname{lk}(g(\lambda))^2$$

for any spatial embedding g of K_{n-2} . Then for any spatial embedding f of K_n , we have

(3.27)
$$\sum_{\lambda \in \Gamma_{t,t+2}(K_n)} \operatorname{lk}(f(\lambda))^2 = 2a(n-6)(n-7) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if
$$s = 0$$
,

(3.28)
$$2(t+2) \sum_{\lambda \in \Gamma_{t+2,t+2}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+4) \sum_{\lambda \in \Gamma_{t,t+4}(K_n)} \operatorname{lk}(f(\lambda))^2$$
$$= 2a(n-6)(n-7)(t+3) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if s = 2 and

(3.29)
$$(t+2) \sum_{\lambda \in \Gamma_{t+2,t+s}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+s+2) \sum_{\lambda \in \Gamma_{t,t+s+2}(K_n)} \operatorname{lk}(f(\lambda))^2$$
$$= a(n-6)(n-7)(2t+s+4) \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

if s = 1 or $s \ge 3$.

Proof. Let f be a spatial embedding of K_n . Then for the embedding f restricted to $F_{ij}^{(kl)}$ $(1 \le i < j \le n, i, j \ne k, l)$, by the assumption we have

$$(3.30) \qquad \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+2,t+s}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{t+2}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_{t+s}(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} \operatorname{lk}(f(\lambda))^{2} + \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t,t+s+2}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{t}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_{t+s+2}(F_{ij}^{(kl)}) \\ iklj \subset \gamma'}} \operatorname{lk}(f(\lambda))^{2} \\ = a \left(\sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{5}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_{3}(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} \operatorname{lk}(f(\lambda))^{2} + \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(kl)}) \\ iklj \subset \lambda}} \operatorname{lk}(f(\lambda))^{2} \right).$$

Note that for the embedding f restricted to $F_{ij}^{(lk)}$, we also have a similar formula as (3.30). Let us take the sum of both side of (3.30) over $1 \le i < j \le n$ and $i, j \ne k, l$. For a pair of disjoint cycles λ of K_n consisting of a (t+2)-cycle γ which contains the edge \overline{kl} and a (t+s)-cycle γ' , let i and j be two distinct vertices of K_n which are adjacent to \overline{kl} in γ . Then λ is a pair of disjoint cycles of $F_{ij}^{(kl)}$ or $F_{ij}^{(lk)}$ consisting of a (t+2)-cycle γ' . This implies

that

$$(3.31) \sum_{\substack{1 \le i < j \le n \\ i_j \ne k, l}} \left(\sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+2,t+s}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{t+2}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_{t+s}(F_{ij}^{(kl)})} } \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+2,t+s}(F_{ij}^{(lk)}) \\ iklj \subset \gamma}} \frac{\operatorname{lk}(f(\lambda))^2}{\gamma \in \Gamma_{t+2}(F_{ij}^{(lk)}), \ \gamma' \in \Gamma_{t+s}(F_{ij}^{(lk)})} } \right)$$
$$= \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+2,t+s}(K_n) \\ \gamma \in \Gamma_{t+2}(K_n), \ \gamma' \in \Gamma_{t+s}(K_n) \\ kl \subset \gamma}} \operatorname{lk}(f(\lambda))^2.$$

In the same way as (3.31), we also have

$$(3.32) \sum_{\substack{1 \leq i < j \leq n \\ i, j \neq k, l}} \left(\sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t, t+s+2}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_t(F_{ij}^{(kl)}), \gamma' \in \Gamma_{t+s+2}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_t(F_{ij}^{(kl)}), \gamma' \in \Gamma_{t+s+2}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_t(K_n), \frac{\gamma'}{\gamma' \in \Gamma_{t+s+2}(K_n)} \\ \gamma \in \Gamma_t(K_n), \frac{\gamma'}{kl \subset \gamma'} \in \Gamma_{t+s+2}(K_n) \\ \gamma \in \Gamma_5(F_{ij}^{(kl)}), \gamma' \in \Gamma_{3,5}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_5(F_{ij}^{(kl)}), \gamma' \in \Gamma_3(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_5(F_{ij}^{(kl)}), \gamma' \in \Gamma_3(K_n) \\ \gamma \in \Gamma_5(K_n), \gamma' \in \Gamma_3(K_n) \\ \gamma \in \Gamma_5(K_n), \gamma' \in \Gamma_3(K_n) \\ kl \subset \gamma \end{cases} kl(f(\lambda))^2.$$

For a pair of disjoint cycles λ of K_n consisting of a *t*-cycle and a (t+s)-cycle, let \overline{ij} be an edge of $K_{n-2}^{(kl)}$ which is not contained in λ . Then λ is a pair of disjoint cycles of $F_{ij}^{(kl)}$ (resp. $F_{ij}^{(lk)}$) which does not contain \overline{iklj} (resp. \overline{ilkj}). Note that there are $\binom{n-2}{2} - (2t+s)$ ways to choose such a pair of *i* and *j*. This implies that

$$(3.34)\sum_{\substack{1\leq i< j\leq n\\i,j\neq k,l}}\sum_{\lambda\in\Gamma_{t,t+s}(F_{ij}^{(kl)})}\operatorname{lk}(f(\lambda))^{2} = \left(\binom{n-2}{2} - (2t+s)\right)\sum_{\lambda\in\Gamma_{t,t+s}(K_{n-2}^{(kl)})}\operatorname{lk}(f(\lambda))^{2},$$

$$(3.35)\sum_{\substack{1\leq i< j\leq n\\i,j\neq k,l}}\sum_{\lambda\in\Gamma_{t,t+s}(F_{ij}^{(lk)})}\operatorname{lk}(f(\lambda))^{2} = \left(\binom{n-2}{2} - (2t+s)\right)\sum_{\lambda\in\Gamma_{t,t+s}(K_{n-2}^{(kl)})}\operatorname{lk}(f(\lambda))^{2}.$$

In the same way as (3.34) and (3.35), we also have

$$(3.36) \quad \sum_{\substack{1 \le i < j \le n \\ i, j \ne k, l}} \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(kl)}) \\ \overline{iklj \not\subset \lambda}}} \operatorname{lk}(f(\lambda))^2 = \left(\binom{n-2}{2} - 6\right) \sum_{\substack{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)}) \\ \overline{iklj \not\subset \lambda}}} \operatorname{lk}(f(\lambda))^2,$$

$$(3.37) \quad \sum_{\substack{1 \le i < j \le n \\ i, j \ne k, l}} \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(lk)}) \\ \overline{ilkj \not\subset \lambda}}} \operatorname{lk}(f(\lambda))^2 = \left(\binom{n-2}{2} - 6\right) \sum_{\substack{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)}) \\ \overline{iklj \not\subset \lambda}}} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.31), (3.32), (3.33), (3.34), (3.35), (3.36) and (3.37) with (3.30), we have

$$(3.38) \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+2,t+s}(K_n)\\\gamma\in\Gamma_{t+2}(K_n),\ \gamma'\in\Gamma_{t+s}(K_n)\\\overline{kl}_{C\gamma}}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t,t+s+2}(K_n)\\\gamma\in\Gamma_t(K_n),\ \gamma'\in\Gamma_{t+s+2}(K_n)}} \operatorname{lk}(f(\lambda))^2 + 2\left(\binom{n-2}{2} - (2t+s)\right) \sum_{\substack{\lambda\in\Gamma_{t,t+s}(K_{n-2}^{(kl)})\\\lambda\in\Gamma_{t,t+s}(K_{n-2}^{(kl)})}} \operatorname{lk}(f(\lambda))^2 = a\left(\sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,5}(K_n)\\\gamma\in\Gamma_5(K_n),\ \gamma'\in\Gamma_3(K_n)\\\overline{kl}_{C\gamma}}} \operatorname{lk}(f(\lambda))^2 + 2\left(\binom{n-2}{2} - 6\right) \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-2}^{(kl)})\\k\in\gamma}} \operatorname{lk}(f(\lambda))^2\right).$$

Then for the embedding f restricted to $K_{n-2}^{\left(kl\right)},$ by the assumption we have

(3.39)
$$\sum_{\lambda \in \Gamma_{t,t+s}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.38) and (3.39), we have

$$(3.40) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+2,t+s}(K_n)\\\gamma\in\Gamma_{t+2}(K_n), \ \gamma'\in\Gamma_{t+s}(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t,t+s+2}(K_n)\\\overline{kl}\subset\gamma'}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,5}(K_n)\\\gamma\in\Gamma_{5}(K_n), \ \gamma'\in\Gamma_{3}(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + 2a(2t+s-6) \sum_{\lambda\in\Gamma_{3,3}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2.$$

Now we assume that s = 1 or $s \ge 3$. Let us take the sum of both sides of (3.40) over $1 \le k < l \le n$. For a pair of disjoint cycles λ of K_n consisting of a (t+2)-cycle γ and a (t+s)-cycle γ' , let \overline{kl} be an edge of K_n which is contained in γ . Note that there are t+2 ways to choose such an edge \overline{kl} . This implies that

$$(3.41)\sum_{1\leq k< l\leq n}\sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+2,t+s}(K_n)\\\gamma\in\Gamma_{t+2}(K_n),\ \gamma'\in\Gamma_{t+s}(K_n)\\\overline{kl}\subset\gamma}} \mathrm{lk}(f(\lambda))^2 = (t+2)\sum_{\lambda\in\Gamma_{t+2,t+s}(K_n)} \mathrm{lk}(f(\lambda))^2.$$

In the same way as (3.41), we also have

$$(3.42) \sum_{\substack{1 \le k < l \le n \\ \gamma \in \Gamma_t(K_n), \ \gamma' \in \Gamma_{t,t+s+2}(K_n) \\ kl \subset \gamma'}} \sum_{\substack{k = \gamma \cup \gamma' \in \Gamma_{1,t+s+2}(K_n) \\ kl \subset \gamma'}} \operatorname{lk}(f(\lambda))^2 = (t+s+2) \sum_{\lambda \in \Gamma_{t,t+s+2}(K_n)} \operatorname{lk}(f(\lambda))^2,$$

$$(3.43) \sum_{\substack{1 \le k < l \le n \\ \gamma \in \Gamma_5(K_n), \ \gamma' \in \Gamma_{3,5}(K_n) \\ kl \subset \gamma}} \sum_{\substack{k \in \Gamma_{3,5}(K_n) \\ kl \subset \gamma}} \operatorname{lk}(f(\lambda))^2 = 5 \sum_{\lambda \in \Gamma_{3,5}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

For a pair of two disjoint 3-cycles λ of K_n , let k, l be two distinct vertices of K_n which is not contained in λ . Then λ is a pair of two disjoint 3-cycles of $K_{n-2}^{(kl)}$. Note that there are $\binom{n-6}{2}$ ways to choose such two vertices k, l. This implies that

(3.44)
$$\sum_{1 \le k < l \le n} \sum_{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2 = \binom{n-6}{2} \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.41), (3.42), (3.43) and (3.44) with (3.40), we have

(3.45)
$$(t+2) \sum_{\lambda \in \Gamma_{t+2,t+s}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+s+2) \sum_{\lambda \in \Gamma_{t,t+s+2}(K_n)} \operatorname{lk}(f(\lambda))^2$$
$$= 5a \sum_{\lambda \in \Gamma_{3,5}(K_n)} \operatorname{lk}(f(\lambda))^2 + 2a(2t+s-6) \binom{n-6}{2} \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Then by (3.45) and Lemma 2.6, we have (3.29).

Next, assume that s = 2. Then by (3.40), we have

$$(3.46) \qquad \sum_{\substack{\lambda \in \Gamma_{t+2,t+2}(K_n) \\ \overline{kl} \subset \lambda}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t,t+4}(K_n) \\ \gamma \in \Gamma_t(K_n), \ \gamma' \in \Gamma_{t+4}(K_n)}} \operatorname{lk}(f(\lambda))^2$$
$$= a \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(K_n) \\ \gamma \in \Gamma_5(K_n), \ \gamma' \in \Gamma_3(K_n) \\ \overline{kl} \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + 2a(2t-4) \sum_{\substack{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)}) \\ \lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)})}} \operatorname{lk}(f(\lambda))^2.$$

Let us take the sum of both sides of (3.46) over $1 \le k < l \le n$. For a pair of disjoint (t+2)-cycles λ of K_n , let \overline{kl} be an edge of K_n which is contained in λ . Note that there are 2(t+2) ways to choose such an edge \overline{kl} . This implies that

$$(3.47)\sum_{1\le k< l\le n}\sum_{\lambda\in\Gamma_{t+2,t+2}(K_n)} \mathrm{lk}(f(\lambda))^2 = 2(t+2)\sum_{\lambda\in\Gamma_{t+2,t+2}(K_n)} \mathrm{lk}(f(\lambda))^2.$$

By combining (3.42), (3.43), (3.44) and (3.47) with (3.46), we have

(3.48)
$$2(t+2) \sum_{\lambda \in \Gamma_{t+2,t+2}(K_n)} \operatorname{lk}(f(\lambda))^2 + (t+4) \sum_{\lambda \in \Gamma_{t,t+4}(K_n)} \operatorname{lk}(f(\lambda))^2 \\ = 5a \sum_{\lambda \in \Gamma_{3,5}(K_n)} \operatorname{lk}(f(\lambda))^2 + 4a(t-2)\binom{n-6}{2} \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

Then by (3.48) and Lemma 2.6, we have (3.28).

Finally, assume that s = 0. By (3.30), we have

$$(3.49) \qquad \sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{t+2,t}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_{t+2}(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_t(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda \in \Gamma_{t,t}(F_{ij}^{(kl)}) \\ iklj \not\in \lambda}} \operatorname{lk}(f(\lambda))^2 \\ = a \left(\sum_{\substack{\lambda = \gamma \cup \gamma' \in \Gamma_{3,5}(F_{ij}^{(kl)}) \\ \gamma \in \Gamma_5(F_{ij}^{(kl)}), \ \gamma' \in \Gamma_3(F_{ij}^{(kl)}) \\ iklj \subset \gamma}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda \in \Gamma_{3,3}(F_{ij}^{(kl)}) \\ iklj \not\in \lambda}} \operatorname{lk}(f(\lambda))^2 \right).$$

Note that, for the embedding f restricted to $F_{ij}^{(lk)}$, we have a similar formula as (3.49). Let us take the sum of both side of (3.49) over $1 \le i < j \le n$ and $i, j \ne k, l$. By combining (3.31), (3.33), (3.34), (3.35), (3.36) and (3.37) with (3.49), we have

$$(3.50) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+2,t}(K_n)\\\gamma\in\Gamma_{t+2}(K_n),\ \gamma'\in\Gamma_t(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + 2\left(\binom{n-2}{2} - 2t\right) \sum_{\substack{\lambda\in\Gamma_{t,t}(K_{n-2}^{(kl)})\\kl\subset\gamma}} \operatorname{lk}(f(\lambda))^2$$
$$= a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,5}(K_n)\\\gamma\in\Gamma_5(K_n),\ \gamma'\in\Gamma_3(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + 2a\left(\binom{n-2}{2} - 6\right) \sum_{\substack{\lambda\in\Gamma_{3,3}(K_{n-2}^{(kl)})\\kl\subset\gamma}} \operatorname{lk}(f(\lambda))^2.$$

Then for the embedding f restricted to $K_{n-2}^{\left(kl\right)},$ by the assumption we have

(3.51)
$$\sum_{\lambda \in \Gamma_{t,t}(K_{n-2}^{(kl)})} \operatorname{lk}(\lambda))^2 = a \sum_{\lambda \in \Gamma_{3,3}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2.$$

By combining (3.50) and (3.51), we have

$$(3.52) \qquad \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{t+2,t}(K_n)\\\gamma\in\Gamma_{t+2}(K_n),\ \gamma'\in\Gamma_t(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 \\ = a \sum_{\substack{\lambda=\gamma\cup\gamma'\in\Gamma_{3,5}(K_n)\\\gamma\in\Gamma_5(K_n),\ \gamma'\in\Gamma_3(K_n)\\\overline{kl}\subset\gamma}} \operatorname{lk}(f(\lambda))^2 + 2a(2t-6) \sum_{\lambda\in\Gamma_{3,3}(K_{n-2}^{(kl)})} \operatorname{lk}(f(\lambda))^2.$$

Now we take the sum of both sides of (3.52) over $1 \le k < l \le n$. By combining (3.41), (3.43) and (3.44) with (3.52), we have

(3.53)
$$(t+2) \sum_{\lambda \in \Gamma_{t+2,t}(K_n)} \operatorname{lk}(f(\lambda))^2 = 5a \sum_{\lambda \in \Gamma_{3,5}(K_n)} \operatorname{lk}(f(\lambda))^2 + 4a(t-3)\binom{n-6}{2} \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$

Then by (3.53) and Lemma 2.6, we have (3.27).

Proof of Theorem 1.5. First we show in the case of p = q = n/2 by the induction on p. If p = 4, then by Theorem 2.1, we have the result. Assume that $p \ge 5$ and it holds that

(3.54)
$$\sum_{\lambda \in \Gamma_{p-1,p-1}(K_{2p-2})} \operatorname{lk}(g(\lambda))^2 = (2p-8)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p-2})} \operatorname{lk}(g(\lambda))^2$$

for any spatial embedding g of K_{2p-2} . Then by (3.54) and setting n = 2p - 1, a = (2p - 8)!, t = p - 1 and s = 0 in Lemma 3.1, we have

(3.55)
$$\sum_{\lambda \in \Gamma_{p-1,p}(K_{2p-1})} \operatorname{lk}(h(\lambda))^2 = 2(2p-7)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p-1})} \operatorname{lk}(h(\lambda))^2$$

for any spatial embedding h of K_{2p-1} . Then by (3.55) and setting n = 2p, a = 2(2p-7)!, t = p-1 and s = 1 in Lemma 3.1, we have

(3.56)
$$2p \sum_{\lambda \in \Gamma_{p,p}(K_{2p})} \operatorname{lk}(f(\lambda))^{2} + (p+1) \sum_{\lambda \in \Gamma_{p-1,p+1}(K_{2p})} \operatorname{lk}(f(\lambda))^{2}$$
$$= 2(2p-6)(2p+1) \sum_{\lambda \in \Gamma_{3,3}(K_{2p})} \operatorname{lk}(f(\lambda))^{2}$$

for any spatial embedding f of K_{2p} . On the other hand, by (3.54) and setting n = 2p, a = (2p - 8)!, t = p - 1 and s = 0 in Lemma 3.2, we also have

(3.57)
$$\sum_{\lambda \in \Gamma_{p-1,p+1}(K_{2p})} \operatorname{lk}(f(\lambda))^2 = 2(2p-6)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p})} \operatorname{lk}(f(\lambda))^2$$

Then by (3.56) and (3.57), we have the result.

Next, we show in the case of n = p + q (p < q) by the induction on $r = q - p \ge 1$. In the case of r = 1, as we showed in the first half of this proof, we have

(3.58)
$$\sum_{\lambda \in \Gamma_{p,p}(K_{2p})} \operatorname{lk}(g(\lambda))^2 = (2p-6)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p})} \operatorname{lk}(g(\lambda))^2$$

for any spatial embedding g of K_{2p} . Then by (3.58) and setting n = 2p + 1, a = (2p - 6)!, t = p and s = 0 in Lemma 3.1, we have

$$\sum_{\lambda \in \Gamma_{p,p+1}(K_{2p+1})} \mathrm{lk}(f(\lambda))^2 = 2(2p-5)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p+1})} \mathrm{lk}(f(\lambda))^2$$

for any spatial embedding f of K_{2p+1} . Thus we have the result.

In the case of r = 2, by (3.58) and setting n = 2p + 2, a = (2p - 6)!, t = p and s = 0 in Lemma 3.2, we have

$$\sum_{\lambda \in \Gamma_{p,p+2}(K_{2p+2})} \mathrm{lk}(f(\lambda))^2 = 2(2p-4)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p+2})} \mathrm{lk}(f(\lambda))^2$$

for any spatial embedding f of K_{2p+1} . Thus we have the result.

Assume that $r\geq 3$ and it holds that

$$(3.59)\sum_{\lambda\in\Gamma_{p,p+(r-1)}(K_{2p+r-1})} \operatorname{lk}(g(\lambda))^2 = 2(2p+r-7)!\sum_{\lambda\in\Gamma_{3,3}(K_{2p+r-1})} \operatorname{lk}(g(\lambda))^2$$

for any spatial embedding g of K_{2p+r-1} . Then by (3.59) and setting n = 2p + r, a = 2(2p + r = 7)!, t = p and $s = r - 1 \ge 2$ in Lemma 3.1, we have

(3.60)
$$(p+1) \sum_{\lambda \in \Gamma_{p+1,p+(r-1)}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2 + (p+r) \sum_{\lambda \in \Gamma_{p,p+r}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2$$

= $2(2p+r-6)!(2p+r+1) \sum_{\lambda \in \Gamma_{3,3}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2.$

Here, by the induction hypothesis, we also have

$$(3.61) \sum_{\lambda \in \Gamma_{p+1,p+(r-1)}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2 = \sum_{\lambda \in \Gamma_{p+1,(p+1)+(r-2)}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2 \\ = 2(2p+r-6)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2.$$

By (3.60) and (3.61), we have

$$(3.62) \quad 2(p+1)(2p+r-6)! \sum_{\lambda \in \Gamma_{3,3}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2 + (p+r) \sum_{\lambda \in \Gamma_{p,p+r}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2 = 2(2p+r-6)!(2p+r+1) \sum_{\lambda \in \Gamma_{3,3}(K_{2p+r})} \operatorname{lk}(f(\lambda))^2.$$

By (3.62), we have the result.

Finally we show (1.2). Assume that n is odd. Then there are (n-5)/2 ways to choose a pair of two integers p, q with n = p + q. Then by (1.1), we have

$$\sum_{p+q=n} \sum_{\lambda \in \Gamma_{p,q}(K_n)} \operatorname{lk}(f(\lambda))^2 = \frac{n-5}{2} \cdot 2(n-6)! \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$$
$$= (n-5)! \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2.$$

On the other hand, assume that n is even. Then there are (n-6)/2 ways to choose a pair of two integers p, q with $p \neq q$ and n = p + q, and there is only one way to choose a pair of two integers p, q with p = q and n = p + q (namely p = q = n/2)). Then by (1.1), we have

$$\sum_{p+q=n} \sum_{\substack{\lambda \in \Gamma_{p,q}(K_n) \\ p \neq q}} \operatorname{lk}(f(\lambda))^2$$

$$= \sum_{\substack{p+q=n \\ p \neq q}} \sum_{\substack{\lambda \in \Gamma_{p,q}(K_n) \\ \lambda \in \Gamma_{3,3}(K_n)}} \operatorname{lk}(f(\lambda))^2 + \sum_{\substack{\lambda \in \Gamma_{3,3}(K_n) \\ \lambda \in \Gamma_{3,3}(K_n)}} \operatorname{lk}(f(\lambda))^2 + (n-6)! \sum_{\substack{\lambda \in \Gamma_{3,3}(K_n) \\ \lambda \in \Gamma_{3,3}(K_n)}} \operatorname{lk}(f(\lambda))^2.$$

This completes the proof.

Proof of Corollary 1.6. We show (1). For any two spatial embeddings f and g of K_n , by (1.1), we have

(3.63)
$$\sum_{\lambda \in \Gamma_{p,p}(K_n)} \operatorname{lk}(f(\lambda))^2 - \sum_{\lambda \in \Gamma_{p,p}(K_n)} \operatorname{lk}(g(\lambda))^2$$
$$= (n-6)! \Big(\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 - \sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(g(\lambda))^2\Big).$$

Since $\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2$ and $\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(g(\lambda))^2$ have the same parity, that is also equal to the parity of $\binom{n}{6}$, by (3.63), we have

$$(3.64) \quad \sum_{\lambda \in \Gamma_{p,p}(K_n)} \operatorname{lk}(f(\lambda))^2 \equiv \sum_{\lambda \in \Gamma_{p,p}(K_n)} \operatorname{lk}(g(\lambda))^2 \pmod{2(n-6)!}.$$

Note that there exists a spatial embedding g of K_n such that

(3.65)
$$\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk} (g(\lambda))^2 = \binom{n}{6}.$$

see Remark 3.3. Thus by (3.64) and (3.65), we have

$$\sum_{k \in \Gamma_{3,3}(K_n)} \operatorname{lk} (f(\lambda))^2 \equiv \binom{n}{6} \pmod{2(n-6)!}$$

for any spatial embedding f of K_n . Since $\binom{n}{6}$ is odd if and only if $n \equiv 6, 7 \pmod{8}$, we have the result. (2) and (3) can be shown in the same way as (1).

Proof of Corollary 1.7. Note that no pair of two disjoint 3-cycles λ of K_n is shared by two distinct subgraphs of K_n isomorphic to K_6 . Then for any spatial embedding f of K_n , Theorem 1.1 (1) implies that

(3.66)
$$\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk} (f(\lambda))^2 \ge \binom{n}{6}.$$

Thus by (3.66) and Theorem 1.5, we have the result.

Remark 3.3. Endo-Otsuki introduced in [4] a canonical book presentation of K_n , which is a certain special spatial embedding f_b of K_n , and Otsuki showed in [9] that $f_b(K_n)$ contains exactly $\binom{n}{6}$ Hopf links corresponding to all the pairs of two disjoint 3-cycles of K_n if $n \ge 6$ (see also Example 3.5 in the case of n = 8). Thus the lower bound of each of the inequalities in Corollary 1.7 is sharp.

Example 3.4. Let g and h be two spatial embeddings of K_7 as illustrated in Fig. 3.1. The embedding g was given in [3] as an embedding that the image contains exactly one nontrivial knot, and the embedding h is obtained from g by a single crossing change at the crossing between $g(\overline{24})$ and $g(\overline{35})$. Then we can see that all of the nonsplittable constituent 2-component links of type (3,3) of $g(K_7)$ are exactly 7 Hopf links, and the ones of $h(K_7)$ are exactly 9 Hopf links. Thus by Theorem 1.5 ((2.19)) we have

(3.67)
$$\sum_{\lambda \in \Gamma_{3,4}(K_7)} \operatorname{lk}(g(\lambda))^2 = 14,$$

(3.68)
$$\sum_{\lambda \in \Gamma_{3,4}(K_7)} \operatorname{lk}(h(\lambda))^2 = 18$$

Actually $g(K_7)$ contains exactly 14 Hopf links and $h(K_7)$ contains exactly 18 Hopf links as all of the nonsplittable constituent Hamiltonian 2-component links. Note that for any spatial embedding f of K_7 , by Corollary 1.6 (2), we have

(3.69)
$$\sum_{\lambda \in \Gamma_{3,4}(K_7)} \operatorname{lk}(f(\lambda))^2 \equiv 2 \pmod{4}$$

By (3.67) and (3.68), the congruence (3.69) is the best possible.



FIGURE 3.1. Two spatial graphs of K_7

Example 3.5. Let g and h be two spatial embeddings of K_8 as illustrated in Fig. 3.2. The embedding g is a canonical book presentation of K_8 (see Remark 3.3), and the embedding h was given in [5] as an embedding that the image contains only Hopf links as nonsplittable constituent 2-component links. Then we can see that all of the nonsplittable constituent 2-component links of type (3,3) of $g(K_8)$ are exactly 28 Hopf links, and the ones of $h(K_8)$ are exactly 30 Hopf links. Thus by Theorem 1.5 (Theorem 2.1) we have

(3.70)
$$\sum_{\lambda \in \Gamma_{3,5}(K_8)} \operatorname{lk}(g(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{4,4}(K_8)} \operatorname{lk}(g(\lambda))^2 = 112,$$

(3.71)
$$\sum_{\lambda \in \Gamma_{3,5}(K_8)} \operatorname{lk}(h(\lambda))^2 = 2 \sum_{\lambda \in \Gamma_{4,4}(K_8)} \operatorname{lk}(h(\lambda))^2 = 120$$

Actually $g(K_8)$ contains exactly 112 Hopf links of type (3,5) and exactly 52 Hopf links and one (2, 4)-torus link $g([1357] \cup [2468])$ of type (4, 4) as all of the nonsplittable constituent Hamiltonian 2-component links, where $[i_1i_2 \cdots i_p]$ denotes a p-cycle $\overline{i_1i_2} \cup \overline{i_2i_3} \cup \cdots \cup \overline{i_pi_1}$, and $h(K_8)$ contains exactly 120 Hopf links of type (3,5) and exactly 60 Hopf links of type (4, 4) as all of the nonsplittable constituent Hamiltonian 2-component links [5]. Here we checked the number of nonsplittable links by a computer program *Gordian* [2]. Note that for any spatial embedding fof K_8 , by Corollary 1.6 (1) and (2), we have

(3.72)
$$\sum_{\lambda \in \Gamma_{3,5}(K_8)} \operatorname{lk}(f(\lambda))^2 \equiv 0 \pmod{8},$$

(3.73)
$$\sum_{\lambda \in \Gamma_{4,4}(K_8)} \operatorname{lk}(f(\lambda))^2 \equiv 0 \pmod{4}$$

By (3.70) and (3.71), both the congruences (3.72) and (3.73) are the best possible.



FIGURE 3.2. Two spatial graphs of K_8

We strongly think that all of the congruences in Corollary 1.6 are the best possible for each $n \ge 6$. To show it, it is sufficient to give the affirmative answer to the following question.

Question 3.6. For each integer $n \ge 9$, does there exist a spatial embedding f of K_n satisfying $\sum_{\lambda \in \Gamma_{3,3}(K_n)} \operatorname{lk}(f(\lambda))^2 = \binom{n}{6} + 2?$

Remark 3.7. The spatial graph $g(K_8)$ in Example 3.5 also satisfies

(3.74)
$$\sum_{\lambda \in \Gamma_{3,4}(K_8)} \operatorname{lk}(g(\lambda)) = 112$$

by Lemma 2.4, and actually $g(K_8)$ contains exactly 112 Hopf links as all of the nonsplittable constituent 2-component links of type (3, 4). Therefore the number of nonsplittable constituent 2-component links of $g(K_8)$ is 28 + 112 + 53 + 112 = 305. This gives an upper bound of the minimum number of nonsplittable constituent 2-component links of a spatial graph of K_8 (see [5, Theorem 2]). On the other hand, if a spatial graph $f(K_8)$ contains only 2-component links with $|lk| \leq 1$ as the nonsplittable constituent 2-component links, then it follows from (3.66), Lemma 2.4 and Corollary 1.7 that the number of nonsplittable constituent 2-component links of $f(K_8)$ is greater than or equal to 28 + 112 + 56 + 112 = 308. This implies that if a spatial graph of K_8 realizes the minimum number of the nonsplittable constituent 2-component links then it must contain a 2-component link with $|lk| \geq 2$.

Proof of Corollary 1.8. Note that the number of all elements in $\Gamma_{p,q}(K_n)$ is

$$\begin{cases} \frac{1}{2} \binom{n}{p} \cdot \frac{(p-1)!}{2} \cdot \frac{(p-1)!}{2} = \frac{n!}{8p^2} \quad (p=q) \\ \binom{n}{p} \cdot \frac{(p-1)!}{2} \cdot \frac{(q-1)!}{2} = \frac{n!}{4pq} \quad (p \neq q). \end{cases}$$

Then by Corollary 1.7, if p = q then we have

(3.75)
$$\left(\max_{\lambda\in\Gamma_{p,p}(K_n)}\left|\operatorname{lk}(f(\lambda))\right|\right)^2\cdot\frac{n!}{8p^2}\geq\sum_{\lambda\in\Gamma_{p,p}(K_n)}\operatorname{lk}(f(\lambda))^2\geq\frac{n!}{6!},$$

and if $p \neq q$ then we have

(3.76)
$$\left(\max_{\lambda\in\Gamma_{p,q}(K_n)} |\mathrm{lk}(f(\lambda))|\right)^2 \cdot \frac{n!}{4pq} \ge \sum_{\lambda\in\Gamma_{p,q}(K_n)} \mathrm{lk}(f(\lambda))^2 \ge 2 \cdot \frac{n!}{6!}$$

Therefore by (3.75) and (3.76), we have

$$\left(\max_{\lambda\in\Gamma_{p,q}(K_n)}\left|\operatorname{lk}(f(\lambda))\right|\right)^2 \ge \frac{pq}{90}$$

Thus we have the result.

Proof of Corollary 1.9. If $pq > 90(m-1)^2$, then by Corollary 1.8 we have

$$\max_{\in \Gamma_{p,q}(K_n)} |\mathrm{lk}(f(\lambda))| \ge \frac{\sqrt{pq}}{3\sqrt{10}} > m - 1.$$

This implies the desired conclusion.

 λ

Appendix A. Maximum value of a_2 for Hamiltonian knots in a rectilinear spatial complete graph

A spatial embedding f_r of a simple graph G is said to be *rectilinear* if for any edge e of G, $f_r(e)$ is a straight line segment in \mathbb{R}^3 . We are also interested in rectilinear spatial graphs because they appear in polymer chemistry as a mathematical model for chemical compounds. We refer the reader to [7] for intrinsic knotting on rectilinear spatial graphs of K_n . Note that every constituent knot (resp. link) of a rectilinear spatial graph is a polygonal knot (resp. link) with some sticks, and it is well-known that every polygonal knot with less than or equal to five sticks is trivial. Then by combining this fact and (3.66) with Theorem 1.2, we have the following.

Theorem A.1. (Morishita-Nikkuni [7, Corollary 1.7]) Let $n \ge 6$ be an integer. For any rectilinear spatial embedding f_r of K_n , we have

$$\sum_{\gamma \in \Gamma_n(K_n)} a_2(f_{\mathbf{r}}(\gamma)) \ge \frac{(n-5)(n-6) \cdot (n-1)!}{2 \cdot 6!}.$$

Corollary A.2. Let $n \ge 6$ be an integer. For any rectilinear spatial embedding f_r of K_n , we have

$$\max_{\gamma \in \Gamma_n(K_n)} a_2(f_{\mathbf{r}}(\gamma)) \ge \frac{(n-5)(n-6)}{6!}$$

Proof of Corollary A.2. Note that the number of all elements in $\Gamma_n(K_n)$ is (n-1)!/2. Then by Theorem A.1, we have

$$\max_{\lambda \in \Gamma_n(K_n)} a_2(f_{\mathbf{r}}(\gamma)) \cdot \frac{(n-1)!}{2} \ge \sum_{\lambda \in \Gamma_n(K_n)} a_2(f_{\mathbf{r}}(\gamma)) \ge \frac{(n-5)(n-6) \cdot (n-1)!}{2 \cdot 6!}.$$

Thus we have the result.

Corollary A.2 says that if n is sufficiently large then every rectilinear spatial graph of K_n contains a Hamiltonian knot whose value of a_2 is arbitrary large. Actually we have the following.

Corollary A.3. Let $n \ge 6$ be a positive integer. For a rectilinear spatial embedding f_r of K_n and a positive integer m, if $n > (11 + \sqrt{2880m - 2879})/2$ then there exists a cycle $\gamma \in \Gamma_n(K_n)$ such that $a_2(f_r(\gamma)) \ge m$.

Proof of Corollary A.3. For an integer $n \ge 6$ and a positive integer m, we can see that $n > (11 + \sqrt{2880m - 2879})/2$ if and only if (n - 5)(n - 6)/6! > m - 1. Then by Corollary A.2 we have

$$\max_{\lambda \in \Gamma_n(K_n)} a_2(f_{\mathbf{r}}(\gamma)) \ge \frac{(n-5)(n-6)}{6!} > m-1.$$

This implies the desired conclusion.

Remark A.4. Shirai-Taniyama showed in [10] that for any spatial embedding f of K_n (which does not need to be rectilinear), if $n \ge 96\sqrt{m}$ there exists a cycle γ of K_n such that $a_2(f(\gamma)) \ge m$. Moreover, if $m = 2^{2k}$ for some non-negative integer k, then $n = 48\sqrt{m}$ is sufficient. Note that in their argument, we cannot know whether $f(\gamma)$ is Hamiltonian or not. Corollary A.3 says that if we restrict ourselves to rectilinear spatial embeddings, then $n > (11 + \sqrt{2880m - 2879})/2$ is more sufficient, and rectilinear spatial graph of K_n always contains a Hamiltonian knot with $a_2 \ge m$.

Acknowledgment

The authors are grateful to Professor Jae Choon Cha for his valuable comments.

References

- L. Abrams, B. Mellor and L. Trott, Counting links and knots in complete graphs, *Tokyo J. Math.* 36 (2013), 429–458.
- [2] L. Abrams, B. Mellor and L. Trott, Gordian (Java computer program), available at http://myweb.lmu.edu/bmellor/research/Gordian 510 (2001), 245-267.
- [3] J. H. Conway and C. McA. Gordon, Knots and links in spatial graphs, J. Graph Theory 7 (1983), 445–453.
- [4] T. Endo and T. Otsuki, Notes on spatial representations of graphs, Hokkaido Math. J. 23 (1994), 383–398.
- [5] T. Fleming and B. Mellor, Counting links in complete graphs, Osaka J. Math. 46 (2009), 173–201.
- [6] A. A. Kazakov and Ph. G. Korablev, Triviality of the Conway-Gordon function ω₂ for spatial complete graphs, J. Math. Sci. (N.Y.) 203 (2014), 490–498.
- [7] H. Morishita and R. Nikkuni, Generalizations of the Conway-Gordon theorems and intrinsic knotting on complete graphs, J. Math. Soc. Japan 71 (2019), 1223–1241.
- [8] R. Nikkuni, A refinement of the Conway-Gordon theorems, *Topology Appl.* 156 (2009), 2782–2794.
- [9] T. Otsuki, Knots and links in certain spatial complete graphs, J. Combin. Theory Ser. B 68 (1996), 23–35.
- [10] M. Shirai and K. Taniyama, A large complete graph in a space contains a link with large link invariant, J. Knot Theory Ramifications 12 (2003), 915–919.
- [11] A. Yu. Vesnin and A. V. Litvintseva, On linking of hamiltonian pairs of cycles in spatial graphs (in Russian), Sib. Èlektron. Mat. Izv. 7 (2010), 383–393

Division of Mathematics, Graduate School of Science, Tokyo Woman's Christian University, 2-6-1 Zempukuji, Suginami-ku, Tokyo 167-8585, Japan

 $E\text{-}mail\ address:\ \texttt{d17m104@cis.twcu.ac.jp}$

DEPARTMENT OF MATHEMATICS, SCHOOL OF ARTS AND SCIENCES, TOKYO WOMAN'S CHRISTIAN UNIVERSITY, 2-6-1 ZEMPUKUJI, SUGINAMI-KU, TOKYO 167-8585, JAPAN

E-mail address: nick@lab.twcu.ac.jp