

A STABILITY THEOREM FOR PROJECTIVE CR MANIFOLDS

JUDITH BRINKSCHULTE, C. DENSON HILL, AND MAURO NACINOVICH

ABSTRACT. We consider smooth deformations of the *CR* structure of a smooth 2-pseudoconcave compact *CR* submanifold M of a reduced complex analytic variety X outside the intersection $D \cap M$ with the support D of a Cartier divisor of a positive line bundle F_X . We show that nearby structures still admit projective *CR* embeddings. Special results are obtained under the additional assumptions that X is a projective space or a Fano variety.

CONTENTS

1. Introduction	2
2. Preliminaries and notation	4
2.1. <i>CR</i> manifolds of type (n, k)	4
2.2. Natural projection onto the canonical bundle	6
2.3. <i>CR</i> maps	7
2.4. q -pseudoconcavity	8
2.5. Deformation of <i>CR</i> structures	8
3. $\bar{\partial}_M$ -cohomology	8
3.1. Differential presentation of the tangential <i>CR</i> complex	9
3.2. <i>CR</i> line bundles	10
3.3. Deformations and <i>CR</i> line bundles	12
4. An example	13
5. Vanishing results	16
5.1. Embedded <i>CR</i> manifolds	16
5.2. Some estimates for abstract <i>CR</i> manifolds	17
5.3. Deformation of tangential <i>CR</i> complexes	19
5.4. Estimates for deformations	20
5.5. Vanishing theorems	21
6. Proof of Theorem 1.1	22
6.1. The general case	22
6.2. Generic <i>CR</i> submanifolds of the projective space	23
6.3. Generic <i>CR</i> submanifolds of Fano varieties	24
References	25

1. INTRODUCTION

In this paper we study projective CR manifolds, meaning those that have a CR embedding into some $\mathbb{C}P^N$. We consider a small CR deformation of such an M , and investigate under what conditions the deformed CR manifold still has a projective CR embedding. Our main result is the following CR -embedding theorem.

Theorem 1.1. *Let X be an $(n+k)$ -dimensional reduced complex analytic variety and D the support of a Cartier divisor of a positive line bundle F_X .*

Let M be a smooth compact submanifold of real dimension $2n+k$ of X_{reg} and $\{M_t\}_{t \in \mathbb{R}}$ a family of CR structures of type (n, k) on M , smoothly depending on a real parameter t . We assume that

- (1) M_0 is induced by the embedding $M \hookrightarrow X$;
- (2) M_0 is 2-pseudoconcave;
- (3) the CR structures of all M_t agree to infinite order on $M \cap D$.

Then we can find $\epsilon_0 > 0$ such that, for every $|t| < \epsilon_0$, M_t admits a generic CR embedding into a projective complex variety X_t .

More precise results are obtained under the additional assumptions that X is a projective space or a Fano variety (Theorems 6.1 and 6.2).

The example of §4 shows that an assumption of 1-pseudoconcavity on M_0 is not sufficient to obtain the statement of Theorem 1.1.

Precise definitions and explanation of terms used will be found in the following sections; let us start by explaining how our main theorem relates to other results in the field of CR geometry or analysis on CR manifolds.

The notion of a CR structure of type (n, k) on a differentiable manifold M of real dimension $2n+k$ arises very naturally in two, a priori different, contexts. One is quite geometric in nature, and the other is more analytical, being connected with fundamental questions about partial differential equations.

From the geometric perspective, a CR structure of type (n, k) on a smooth real submanifold M of dimension $2n+k$ of some complex manifold X of complex dimension $n+k$ is induced as the tangential part of the ambient complex structure on X . In particular, one has the tangential Cauchy-Riemann equations on M . In this situation M is said to be generically CR embedded in X . It is also possible that M might be embedded in a complex manifold X of complex dimension greater than $n+k$, with the CR structure on M also being induced as the tangential part of the ambient complex structure in X . In this situation the real codimension of M is greater than the CR dimension k , and M is said to be non generically CR embedded in X . This is an important distinction only globally, because: if M can be locally non generically CR embedded, then M can be locally generically CR embedded (see e.g. [25]).

From the PDE perspective, one begins with (at least locally) a system of n smooth complex vector fields defined on M , which are linearly independent over \mathbb{C} , and satisfy a formal integrability condition. Again n is the *CR* dimension, and k is the *CR* codimension. One then asks if the given complex vector fields might represent (in a local basis) the tangential *CR* equations on M given by some (local) *CR* embedding of M . The criterion for this, is that one must be able to find (locally) a maximal set of functionally independent characteristic coordinates, which means to find $n+k$ independent complex valued functions that are solutions of the homogeneous system associated to the n complex vector fields. When the vector fields can be chosen to have real analytic coefficients, then this is possible ([1]). Hence locally, in the real analytic case, the geometric perspective and the analytical PDE perspective are equivalent. The famous example of L. Nirenberg [33] showed that the two perspectives are not locally equivalent if the coefficients of the vector fields are assumed to be only smooth. Why does this matter? From an analytical PDE viewpoint, it is too restrictive to allow only real analytic structures, one needs cut off functions, partitions of unity; etc., in order to employ calculus and modern analysis. So from this perspective it is more natural to define what is known as an abstract *CR* manifold M of type (n, k) , by reformulating the local vector field basis description given above into a global and invariant version, and to do so within the smooth category. Precise definitions and details are below.

The example of Nirenberg [33] was a single complex vector field in \mathbb{R}^3 , so it endows \mathbb{R}^3 with the structure of an abstract *CR* manifold of type $(1, 1)$. To make matters worse, the complex vector field of Nirenberg's example is a small smooth perturbation, near some point, of the tangential Cauchy Riemann equation to the unit sphere in \mathbb{C}^2 . In other words, the generically *CR* embedded and real analytic *CR* structure of type $(1, 1)$ on the sphere S^3 , inherited from \mathbb{C}^2 , becomes not even locally *CR* embeddable near some point, if it is perturbed in a smooth way to obtain a very nearby abstract *CR* structure of type $(1, 1)$ on the same S^3 . Here the original *CR* structure on S^3 is strictly pseudoconvex, and remains so under the small perturbation.

Rossi [36] constructed small real analytic deformations of the *CR* structure on S^3 in \mathbb{C}^2 , such that the resulting perturbed abstract *CR* structures fail to *CR* embed globally into \mathbb{C}^2 , even though they do locally *CR* embed into \mathbb{C}^2 . On the other hand, sticking with strictly pseudoconvex *CR* structures of hypersurface type, i.e. type $(n, 1)$, Boutet de Monvel proved that when M is compact and $n > 1$, then M has a *CR* embedding into \mathbb{C}^N for some perhaps large N . Catlin and Lempert [15] showed that there are compact strictly pseudoconvex *CR* manifolds of hypersurface type $(n, 1)$, for $n \geq 1$, that are *CR* embedded into some \mathbb{C}^N , which admit small deformations that are also embeddable, but their embeddings cannot be chosen close to the original embedding. Thus these examples exhibit the phenomenon of *instability* of the *CR* embedding. On the other hand, for $n = 1$, Lempert [31] proved that if a compact strictly pseudoconvex M is *CR* embeddable

into \mathbb{C}^2 , hence generically embedded, and if the CR structure gets perturbed by a family of CR embeddable CR structures, then the embedding is *stable*. This means that the embedding of the perturbed structure stays close to the unperturbed one. Additional references to other work on the strictly pseudoconvex hypersurface type case can be found in the above mentioned papers.

In the higher codimensional case ($k > 1$) Polyakov [35] proved a stability theorem for compact 3-pseudoconcave CR manifolds M , generically CR embedded in a complex manifold, under some additional hypotheses, such as the vanishing of a certain cohomology group. The notion of flexible/inflexible, which is closely related to the notion of stability/instability, was introduced in [10], also for the higher codimension case, where the emphasis was on CR manifolds M that are not compact. A CR manifold M of type (n, k) , generically CR embedded in \mathbb{C}^{n+k} , is called flexible if it admits a compactly supported CR deformation whose deformed structure is no longer CR embeddable in \mathbb{C}^{n+k} ; in other words, the deformation causes the CR structure to “flex out” of the space where the original manifold lies. On the other hand, M is inflexible if any compactly supported deformation stays in the class of manifolds globally CR embeddable into \mathbb{C}^{n+k} . In [10], [11] it was proved that if M is a CR submanifold of type (n, k) in \mathbb{C}^{n+k} , and it is only 2-pseudoconcave, then M is inflexible. This relates to the present paper as follows: If a CR structure of a 2-pseudoconcave compact CR manifold of a reduced complex analytic variety gets perturbed by a smooth family of CR structures that “glues to” the unperturbed structure along a Cartier divisor of a positive line bundle, then the perturbed CR structures *do not* “flex out” of the class of globally CR embeddable CR manifolds (Theorem 1.1). Moreover, the embedding is stable if the unperturbed CR manifold sits generically in a projective space (Theorem 6.1).

We emphasize that in this paper we discuss compact CR manifolds and we allow higher codimension and that we are interested in global results, not local ones.

2. PRELIMINARIES AND NOTATION

2.1. CR manifolds of type (n, k) . Let M be a smooth manifold of real dimension m and n, k nonnegative integers with $2n+k=m$. A CR structure of type (n, k) on M is a formally integrable distribution $T_M^{0,1}$ of n -dimensional complex subspaces of the complexified tangent bundle $\mathbb{C}T_M$ of M , transversal to T_M . By this transversality condition, the real parts of vectors in $T_M^{0,1}$ are the elements of the *vector CR distribution* H_M and for each $p \in M$ and $v \in H_{M,p}$ (subscript “ p ” means “the fiber at p ”) there is a unique $v' \in H_{M,p}$ such that $v + iv' \in T_{M,p}^{0,1}$. The correspondence $v \rightarrow v'$ defines a bundle self map $J_M: H_M \rightarrow H_M$ which yields a complex structure on the fibers of H_M , as $J_M^2 = -I_{H_M}$. The formal integrability condition means that

$$(2.1) \quad [\Gamma^\infty(M, T_M^{0,1}), \Gamma^\infty(M, T_M^{0,1})] \subseteq \Gamma^\infty(M, T_M^{0,1}),$$

which is equivalent to the vanishing of the Nijenhuis tensor on H_M , i.e.

$$J_M([X, Y] - [J_M X, J_M Y]) = [J_M X, Y] + [X, J_M Y], \quad \forall X, Y \in \Gamma^\infty(M, H_M).$$

We use the standard notation

$$\begin{aligned} T_M^{1,0} &= \{X - iJ_M X \mid X \in H_M\} \subset \mathbb{C}T_M, \\ T_M^{0,1} &= \{X + iJ_M X \mid X \in H_M\} \subset \mathbb{C}T_M, \\ T_M^{*1,0} &= \{\phi \in \mathbb{C}T_M^* \mid \phi(Z) = 0, \forall Z \in \Gamma^\infty(M, T_M^{0,1})\}, \\ T_M^{*0,1} &= \{\phi \in \mathbb{C}T_M^* \mid \phi(Z) = 0, \forall Z \in \Gamma^\infty(M, T_M^{1,0})\}, \\ H_M^0 &= \{\xi \in T_M^* \mid \xi(X) = 0, \forall X \in \Gamma^\infty(M, H_M)\}. \end{aligned}$$

We have

$$T_M^{1,0} \cap T_M^{0,1} = \{0\}, \quad T_M^{1,0} + T_M^{0,1} = \mathbb{C}H_M, \quad T_M^{*1,0} \cap T_M^{*0,1} = \mathbb{C}H_M^0, \quad T_M^{*1,0} + T_M^{*0,1} = \mathbb{C}T_M^*.$$

Deformations of CR structures were considered especially for the hypersurface-type case $k = 1$ [2, 17]. Since, by a theorem of Darboux (see [16]), 1-codimensional contact structures are locally equivalent, deformations of non degenerate CR hypersurfaces can be described by smooth curves $\{t \rightarrow J_t\}$ of formally integrable complex structures on a fixed contact distribution H_M of a smooth real manifold M . Differently, generalized contact distributions of higher codimension may be not locally equivalent. Therefore, to describe general deformations of abstract CR manifolds of type (n, k) with arbitrary k it will be convenient to focus our attention on their *canonical bundles*, which carry comprehensive information on both their contact and partial complex structures, as explained below.

Denote by \mathcal{A}_M the sheaf of Grassmann algebras of germs of complex valued smooth exterior differential forms on M . The subsheaf \mathcal{A}_M^q of homogeneous elements of degree q consists of the germs of smooth sections of the \mathbb{C} -vector bundle $\Lambda^q(\mathbb{C}T_M^*)$ of complex q -covectors on M .

We can identify the Grassmannian $\mathcal{G}_n^{\mathbb{C}}(M)$ of n -dimensional complex subspaces of $\mathbb{C}T_M$ with the subbundle of the projective bundle $\mathbb{P}(\Lambda^{n+k}(\mathbb{C}T_M^*))$ whose representatives belong to the cone $\mathcal{M}on(\Lambda^{n+k}(\mathbb{C}T^*M))$ of degree $n+k$ monomials¹ of the Grassmann algebra of $\mathbb{C}T^*M$ (see e.g. [14]). Then a distribution of complex n -planes of $\mathbb{C}T_M$ is the datum of a smooth complex line bundle K_M on M , with $K_M \subset \mathcal{M}on(\Lambda^{n+k}(\mathbb{C}T^*M))$.

Definition 2.1. Let $(M, T_M^{0,1})$ be an abstract CR manifold of type (n, k) . The complex line bundle K_M corresponding to the distribution $T_M^{0,1}$ of its *anti-holomorphic tangent vectors* is called its *canonical bundle*.

Notation 2.1. Let us denote by \mathcal{K}_M the subsheaf of ideals in \mathcal{A}_M generated by the sheaf \underline{K}_M of germs of smooth sections of K_M .

¹We recall that, if V is a vector space, the *monomials of degree q* of its exterior algebra Λ^*V are the exterior products $v_1 \wedge \cdots \wedge v_q$ of vectors v_1, \dots, v_q of V .

Proposition 2.1. *A CR structure of type (n, k) on M is completely determined by the datum of a complex line bundle $K_M \subset \text{Mon}(\Lambda^{n+k}(\mathbb{C}T_M^*))$ such that the ideal sheaf \mathcal{K}_M of \mathcal{A}_M generated by the germs its smooth sections satisfies the following conditions:*

- (1) \mathcal{K}_M is closed, i.e. $d\mathcal{K}_M \subseteq \mathcal{K}_M$ (FORMAL INTEGRABILITY);
- (2) $\{X \in \Gamma^\infty(M, T_M) \mid X \lrcorner \mathcal{K}_M \subseteq \mathcal{K}_M\} = \{0\}$ (TRANSVERSALITY TO T_M).

Indeed, when (1) and (2) are valid, K_M is the canonical bundle of the CR manifold $(M, T_M^{0,1})$ with

$$(2.2) \quad \Gamma^\infty(M, T_M^{0,1}) = \{Z \in \Gamma^\infty(M, \mathbb{C}T_M) \mid Z \lrcorner \mathcal{K}_M \subseteq \mathcal{K}_M\}$$

Proof. In fact, (1) is equivalent to the formal integrability (2.1) and (2) to the transversality of $T_M^{0,1}$ to the real tangent distribution T_M . \square

By the Newlander-Nirenberg theorem ([34]) abstract CR manifolds of type $(n, 0)$ are n -dimensional complex manifolds.

2.2. Natural projection onto the canonical bundle. On a CR manifold M of type (n, k) we have an exact sequence of smooth bundle maps

$$(2.3) \quad 0 \longrightarrow \mathbb{C}H_M^0 \longrightarrow T_M^{*1,0} \oplus T_M^{*0,1} \longrightarrow \mathbb{C}T_M^* \longrightarrow 0,$$

where

$$\begin{aligned} \mathbb{C}H_M^0 \ni \xi &\longrightarrow (\xi, \xi) \in T_M^{*1,0} \oplus T_M^{*0,1}, \\ T_M^{*1,0} \oplus T_M^{*0,1} \ni (\xi_1, \xi_2) &\longrightarrow \xi_1 - \xi_2 \in \mathbb{C}T_M^*. \end{aligned}$$

A CR gauge is a bundle map

$$(2.4) \quad \lambda : \mathbb{C}T_M^* \longrightarrow T_M^{*1,0}, \quad \text{such that } \xi - \frac{1}{2}\lambda(\xi) \in T_M^{*0,1}, \quad \forall \xi \in \mathbb{C}T_M^*.$$

Note that, in particular,

$$\lambda(T_M^{*0,1}) \subseteq \mathbb{C}H_M^0 \quad \text{and} \quad \xi - \frac{1}{2}\lambda(\xi) \in \mathbb{C}H_M^0, \quad \forall \xi \in T_M^{*1,0}.$$

Therefore a CR gauge yields a short exact sequence inverting (2.3):

$$0 \longrightarrow \mathbb{C}T_M^* \xrightarrow{(\lambda/2) \oplus (\text{id} - \lambda/2)} T_M^{*1,0} \oplus T_M^{*0,1} \xrightarrow{(\lambda/2 - \text{id}, \lambda/2)} \mathbb{C}H_M^0 \longrightarrow 0,$$

where id denotes the identity on $\mathbb{C}T_M^*$.

There are several possible choices of λ . A CR gauge λ is said to be *of the real type* if

$$(2.5) \quad \lambda(\xi) + \overline{\lambda(\bar{\xi})} = 2 \cdot \xi, \quad \forall \xi \in \mathbb{C}T_M^*$$

and *balanced* if

$$(2.6) \quad \lambda(\xi) = \xi, \quad \forall \xi \in H_M^0.$$

Existence of balanced CR gauges of the real type was proved in [32, §2A]. The following is then easy to deduce from the above properties.

Proposition 2.2. *The bundle map λ of a balanced CR gauge*

- *is a projection of $\mathbb{C}T_M^*$ onto $T_M^{*1,0}$;*

- its restriction to T_M^* is an isomorphism with a real complement of $i \cdot H_M^0$ in $T_M^{*1,0}$.

Let us define a linear map

$$(2.7) \quad \lambda^p : \Lambda^p(\mathbb{C}T_M^*) \rightarrow \Lambda^p(T_M^{*1,0})$$

in such a way that on the monomials we have

$$\lambda^p(\xi_1 \wedge \cdots \wedge \xi_p) = \lambda(\xi_1) \wedge \cdots \wedge \lambda(\xi_p).$$

Proposition 2.3. *The map λ^{n+k} defines a projection of $\Lambda^{n+k}(\mathbb{C}T_M^*)$ onto K_M , which is independent of the particular choice of a balanced CR gauge λ of the real type.*

Proof. Fix a point p of M and a basis

$$(\ddagger) \quad \xi_1, \dots, \xi_n, \tau_1, \dots, \tau_k, \zeta_1, \dots, \zeta_n$$

of $\mathbb{C}T_{M,p}^*$ with $\xi_i \in T_{M,p}^{*1,0}$, for $1 \leq i \leq n$, $\tau_j \in H_M^0$ for $1 \leq j \leq k$ and $\zeta_i \in T_{M,p}^{*0,1}$ for $1 \leq i \leq n$.

If λ is a balanced CR gauge of the real type, then $\lambda(\xi_i) = \xi_i$ for $1 \leq i \leq n$, $\lambda(\tau_j) = \tau_j$ for $1 \leq j \leq k$, $\lambda(\zeta_i) \in \mathbb{C}H_M^0$ for $1 \leq i \leq n$. Hence the image by λ^{n+k} of all monomials of degree $n+k$ in the elements of the basis (\ddagger) which contain a factor ζ_i contain at least $k+1$ factors in $\mathbb{C}H_M^0$ and is therefore 0, while

$$\lambda^{n+k}(\xi_1 \wedge \cdots \wedge \xi_n \wedge \tau_1 \wedge \cdots \wedge \tau_k) = \xi_1 \wedge \cdots \wedge \xi_n \wedge \tau_1 \wedge \cdots \wedge \tau_k.$$

This completes the proof. \square

Definition 2.2. We call λ^{n+k} the *natural projection* onto the canonical bundle.

2.3. CR maps. Let (M_i, K_{M_i}) be abstract CR manifolds of type (n_i, k_i) ($i=1, 2$), $f : M_1 \rightarrow M_2$ a smooth map and $f_* : \mathbb{C}T_{M_1} \rightarrow \mathbb{C}T_{M_2}$ the bundle map obtained by complexifying its differential.

Definition 2.3. The map f is said to be:

- CR if $f_*(T_{M_1}^{0,1}) \subseteq T_{M_2}^{0,1}$,
- a CR immersion (embedding) if it is a smooth immersion (embedding) and moreover $T_{M_1}^{0,1} = \{Z \in \mathbb{C}T_{M_1} \mid f_*(Z) \in T_{M_2}^{0,1}\}$.
- a CR submersion if it is a smooth submersion and $f_*(T^{0,1}M_1) = T^{0,1}M_2$.

A CR immersion (embedding) is called *generic* when $n_1 + k_1 = n_2 + k_2$.

On an abstract CR manifold M of type (n, k) , we consider the Grassmann subalgebra Ω_M^* of $\Lambda^*(\mathbb{C}T_M^*)$ generated by $T_M^{*1,0}$:

$$\Omega_M^* = \sum_{p=0}^{n+k} \Omega_M^p, \quad \text{with} \quad \Omega_M^p = \Omega_M^* \cap \Lambda^p(\mathbb{C}T_M^*).$$

With this notation, $K_M = \Omega_M^{n+k}$. CR maps and immersion can be described by using the bundles Ω_M^* .

Proposition 2.4. *Let M_i be abstract CR manifolds of type (n_i, k_i) , for $i=1, 2$ and $f : M_1 \rightarrow M_2$ a smooth map. Then f is*

- *CR* iff $f^*(\Omega_{M_2}^1) \subseteq \Omega_{M_1}^1$;
- a *CR immersion* iff f is a smooth immersion and $f^*(\Omega_{M_2}^{n_1+k_1}) = \mathcal{K}_{M_1}$.

A generic *CR-immersion* $f : M_1 \rightarrow M_2$ is characterized by

$$(2.8) \quad f^*(\mathcal{K}_{M_2}) = \mathcal{K}_{M_1}. \quad \square$$

2.4. q -pseudoconcavity. Let q be a nonnegative integer. Following [21], we say that a *CR* manifold M , of type (n, k) , is q -pseudoconcave if, for every $p \in M$ and every characteristic covector $\xi \in H_p^0 M \setminus \{0\}$, the scalar Levi form $\mathcal{L}_p(\xi, \cdot)$ has at least q negative and q positive eigenvalues.

We recall that

$$\mathcal{L}_p(\xi, v) = \xi([J_M X, X]) = d\tilde{\xi}(v, J_M(v)), \quad \text{for } v \in H_p M,$$

with $X \in \Gamma^\infty(M, H_M)$, $\tilde{\xi} \in \Gamma^\infty(M, H^0 M)$ and $X_p = v$, $\tilde{\xi}_p = \xi$.

The *CR* dimension n of a q -pseudoconcave *CR* manifold is at least $2q$ (more precise lower bounds on n , depending also on the *CR* codimension k , can be obtained e.g. from [7]).

Various equivalent definitions of the Levi form and more of its aspects are explained in [13].

2.5. Deformation of *CR* structures. With the preparation of §2.1, we can define *CR* deformations of a *CR* manifold M of general type (n, k) in terms of canonical bundles, allowing in this way also a deformation of the underlying contact structure H_M .

Definition 2.4. A smooth one-parameter deformation of a *CR* structure \mathcal{K}_M on M is a smooth map $M \times (-\epsilon_0, \epsilon_0) \ni (p, t) \rightarrow [\mathcal{K}_{M_t, p}] \in \mathbb{P}(\Lambda^{n+k}(\mathbb{C}T^*))$ such that, for each $t \in (-\epsilon_0, \epsilon_0)$, \mathcal{K}_{M_t} is the canonical complex line bundle of a *CR* structure on M and $\mathcal{K}_{M_0} = \mathcal{K}_M$.

Line bundles over a contractible base are trivial (see e.g. [27, Cor.3.4.8]). Hence, for a compact M , we can find a finite open covering $\mathcal{U} = \{U_j\}$ to describe the deformation by the data of smooth maps

$$(2.9) \quad M \times (-\epsilon_0, \epsilon_0) \ni (p, t) \rightarrow \omega_j(p, t) \in \Lambda^{n+k} \mathbb{C}T^* M$$

such that $\omega_{j,t}(\cdot, t) \in \Gamma^\infty(U_j, \mathcal{K}_{M_t})$ for each $t \in (-\epsilon_0, \epsilon_0)$.

3. $\bar{\partial}_M$ -COHOMOLOGY

Let M be an abstract *CR* manifold of type (n, k) .

The ideal sheaf \mathcal{I}_M of \mathcal{A}_M generated by the germs of smooth sections of $T_M^{*1,0}$ is, according to [14], the *characteristic system* of $T_M^{0,1}$. Hence, formal integrability is equivalent to

$$d(\mathcal{I}_M) \subseteq \mathcal{I}_M.$$

For nonnegative integers p , its exterior powers \mathcal{I}_M^p are also d-closed ideals and we can consider the quotients of the de Rham complex:

$$(*) \quad \bar{\partial}_M : \mathcal{I}_M^p / \mathcal{I}_M^{p+1} \longrightarrow \mathcal{I}_M^p / \mathcal{I}_M^{p+1}$$

(we put $\mathcal{I}_M^0 = \mathcal{A}_M$). Since $\mathcal{I}_M^p = 0$ when $p > n+k$, the map $(*)$ is nontrivial if and only if $0 \leq p \leq n+k$.

The canonical \mathbb{Z} -grading of \mathcal{A}_M induces a \mathbb{Z} -grading $\mathcal{I}_M^p = \bigoplus_{q=0}^n \mathcal{I}_M^{p,q}$ of \mathcal{I}_M^p , with $\mathcal{I}_M^{p,q} = \mathcal{I}_M^p \cap \mathcal{A}_M^{p+q}$. Let us set $Q_M^{p,*} = \mathcal{I}_M^p / \mathcal{I}_M^{p+1}$. By passing to the quotients, the \mathbb{Z} -gradings of the exterior powers of \mathcal{I}_M induce the \mathbb{Z} -gradings

$$Q_M^{p,*} = \bigoplus_{q=0}^n Q_M^{p,q}.$$

For each pair of integers (p, q) with $0 \leq p \leq n+k$, $0 \leq q \leq n$ the summand $Q_M^{p,q}$ is the sheaf of germs of smooth sections of a complex vector bundle $Q_M^{p,q}$ on M . Then for each $0 \leq p \leq n+k$, the map $(*)$ is a *differential complex*

$$0 \rightarrow Q_M^{p,0} \xrightarrow{\bar{\partial}_M} Q_M^{p,1} \longrightarrow \dots \xrightarrow{\bar{\partial}_M} Q_M^{p,n-1} \xrightarrow{\bar{\partial}_M} Q_M^{p,n} \rightarrow 0,$$

which is called the *tangential Cauchy-Riemann complex in degree p* (for more details see e.g. [21]).

Since $\mathcal{I}_M^{n+k+1} = 0$, the bundles $Q_M^{n+k,q}$ are subbundles of $\Lambda^{n+k+q}(\mathbb{C}T_M^*)$ and in particular $Q_M^{n+k,0}$ is the same as the *canonical bundle* \mathbb{K}_M . As we will explain below, it is a *CR line bundle* and the tangential Cauchy-Riemann complex in degree $n+k$ is also its $\bar{\partial}_M$ -complex as a *CR line bundle*. This is a special example of a notion that was described e.g. in [22, §7] and that we will quickly recall in the next subsection.

The tangential Cauchy-Riemann complexes can be described in a way that is suitable to deal with smooth deformations of the *CR structure*.

Let us set $T_M^{*p,0} = \Lambda^p(T_M^{*1,0})$ and denote by $\mathcal{T}_M^{*p,0}$ the sheaf of germs of smooth sections of $T_M^{*p,0}$.

3.1. Differential presentation of the tangential CR complex. Let E_M be a complement of $T_M^{*1,0}$ in $\mathbb{C}T_M^*$. We can e.g. fix a smooth Hermitian product on the fibers of $\mathbb{C}T_M^*$ and take $E_M = (T_M^{*1,0})^\perp$. The fiber bundle E_M has rank n and we obtain a bigradation of the Grassmann algebra of the complexified cotangent bundle of M by setting

$$\Lambda^*(\mathbb{C}T_M^*) = \sum_{\substack{0 \leq p \leq n+k \\ 0 \leq q \leq n}} E_M^{p,q}, \quad \text{with } E_M^{p,q} = \Lambda^q(E_M) \otimes_M T_M^{*p,0}.$$

Denote by $\mathcal{E}_M^{p,q}$ the sheaf of germs of smooth sections of $E_M^{p,q}$.

Lemma 3.1. *For every integer p with $0 \leq p \leq n+k$ we have:*

- (1) $\mathcal{E}_M^{p,q} \subseteq \mathcal{I}_M^p$.
- (2) *The restriction of the projection onto the quotient defines isomorphisms*

$$(3.1) \quad E_M^{p,q} \rightarrow Q_M^{p,q}, \quad \mathcal{E}_M^{p,q} \rightarrow Q_M^{p,q}$$

of fiber bundles and of their sheaves of sections. \square

By using the isomorphisms of (3.1) we can rewrite the tangential Cauchy-Riemann complexes in the form

$$0 \longrightarrow \mathcal{E}_M^{p,0} \xrightarrow{d''_M} \mathcal{E}_M^{p,1} \xrightarrow{d''_M} \dots \rightarrow \mathcal{E}_M^{p,n-1} \xrightarrow{d''_M} \mathcal{E}_M^{p,n} \longrightarrow 0.$$

We get explicit representations of d''_M , and hence of $Q_M^{p,q}$ and $\bar{\partial}_M$, by choosing suitable coframes of $\mathbb{C}T_M^*$. Take indeed on an open subset U of M a coframe $\omega_1, \dots, \omega_n, \eta_1, \dots, \eta_{n+k}$ of T_M^*

$$\text{with } \omega_1, \dots, \omega_n \text{ in } \mathcal{E}_M \text{ and } \eta_1, \dots, \eta_{n+k} \in \Gamma^\infty(U, T_M^{*1,0}).$$

Then there are uniquely determined smooth complex vector fields $\bar{L}_1, \dots, \bar{L}_n, P_1, \dots, P_{n+k}$ such that

$$du = \sum_{j=1}^n (\bar{L}_j u) \omega_j + \sum_{i=1}^{n+k} (P_i u) \eta_i$$

The vector fields \bar{L}_h , for $1 \leq h \leq n$, are characterized by

$$\omega_j(\bar{L}_h) = \delta_{j,h} \text{ (Kronecker delta)}, \quad 1 \leq j \leq n, \quad \eta_i(\bar{L}_h) = 0, \quad 1 \leq i \leq n+k,$$

and hence are sections of $T_M^{0,1}$. By our definition,

$$d''_M u = \sum_{j=1}^n (\bar{L}_j u) \omega_j, \quad \forall u \in C^\infty(U).$$

For each j with $1 \leq j \leq n$ we have

$$d\omega_j = \sum_{1 \leq j_1 \leq j_2 \leq n} c_j^{j_1, j_2} \omega_{j_1} \wedge \omega_{j_2} + \zeta_j,$$

with $c_j^{j_1, j_2} \in C^\infty(U)$, $\zeta_j \in \mathcal{I}_M(U)$. This gives in particular

$$d''_M \omega_j = \sum_{1 \leq j_1 \leq j_2 \leq n} c_j^{j_1, j_2} \omega_{j_1} \wedge \omega_{j_2}$$

and hence, computing by recurrence on the number of factors, we get

$$\begin{aligned} d''_M(\omega_{j_0} \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_q}) &= d''_M(\omega_{j_0}) \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_q} + \omega_{j_0} \wedge d''_M(\omega_{j_1} \wedge \dots \wedge \omega_{j_q}), \\ d''_M(\omega_{j_1} \wedge \dots \wedge \omega_{j_q} \wedge \eta_{i_1} \wedge \dots \wedge \eta_{i_p}) &= (d''_M(\omega_{j_1} \wedge \dots \wedge \omega_{j_q})) \wedge \eta_{i_1} \wedge \dots \wedge \eta_{i_p}. \end{aligned}$$

These formulas yield explicit local expressions for d''_M . We note that for $p=0$ they only depend on the choice of the coframe $\omega_1, \dots, \omega_n$ of \mathcal{E}_M .

In particular, if M is compact and $\{M_t\}_{|t| < \epsilon_0}$ a smooth deformation of its CR structure, then \mathcal{E}_M stays transversal to $T_{M_t}^{*1,0}$ for small $|t|$ and therefore the tangential Cauchy-Riemann complex in degree 0 for M_t can be defined on sections of the same vector bundles for all small t .

3.2. CR line bundles. A CR line bundle on M is the datum of a smooth complex line bundle $\pi: F \rightarrow M$, together with a CR structure on F which is compatible with the linear structure of the fibers and for which π is a CR submersion. The compatible CR structures on F are parametrized (modulo CR equivalence) by special cohomology classes of $H^1(M, \mathcal{H}_M^{0,1})$, where $\mathcal{H}_M^{0,1}$ is the sheaf of germs of local cohomology classes of degree $(0, 1)$ on M : In a

local trivialization $\{U_i, s_i\}$ of F , with transition functions $(\gamma_{i,j})$, the structure is described by the datum of $\bar{\partial}_M$ -closed forms ϕ_i in $Q_M^{0,1}(U_i)$ with

$$\bar{\partial}_M^F s_i = \phi_i \otimes s_i \text{ on } U_i \text{ and } \phi_i - \phi_j = \gamma_{i,j}^{-1} \bar{\partial}_M \gamma_{i,j} \text{ on } U_i \cap U_j.$$

We can consistently define a differential operator $\bar{\partial}_M^F$ on smooth sections of F , with values in the smooth sections of $Q_M^{0,1} \otimes F$, in such a way that²

$$\bar{\partial}_M^F(f s_i) = (\bar{\partial}_M f \otimes s_i) + (f \cdot \phi_i) \otimes s_i, \quad \forall f \in C^\infty(U_i).$$

Let us use the notation \mathcal{F}_M to indicate the sheaf of germs of smooth sections of F_M and $\mathcal{F}_M^{p,q}$ for the sheaf of germs of smooth sections on M of the bundle $Q_M^{p,q} \otimes F$. Then we can define, in general, a differential operator $\bar{\partial}_M^F$ mapping sections of $\mathcal{F}_M^{p,q}$ into sections of $\mathcal{F}_M^{p,q+1}$ by setting for U open in M , $\alpha \in Q_M^{p,q}(U)$ and $\sigma \in \mathcal{F}_M(U)$,

$$\bar{\partial}_M^F(\alpha \otimes \sigma) = (\bar{\partial}_M \alpha) \otimes \sigma + (-1)^{p+q} \alpha \wedge \bar{\partial}_M^F \sigma.$$

In this way we obtain the *tangential Cauchy-Riemann complexes with coefficients in F* :

$$\begin{aligned} 0 \longrightarrow \mathcal{F}_M^{p,0}(U) &\xrightarrow{\bar{\partial}_M^F} \mathcal{F}_M^{p,1}(U) \longrightarrow \dots \\ \dots &\longrightarrow \mathcal{F}_M^{p,n-1}(U) \xrightarrow{\bar{\partial}_M^F} \mathcal{F}_M^{p,n}(U) \longrightarrow 0. \end{aligned}$$

We denote its cohomology groups by

$$H^{p,q}(U, F_M) = \frac{\ker(\bar{\partial}_M^F : \mathcal{F}_M^{p,q}(U) \rightarrow \mathcal{F}_M^{p,q+1}(U))}{\text{image}(\bar{\partial}_M^F : \mathcal{F}_M^{p,q-1}(U) \rightarrow \mathcal{F}_M^{p,q}(U))}.$$

When the CR structure on M is induced from a generic embedding into an $(n+k)$ -dimensional complex manifold X , the CR line bundle K_M is the restriction to M of the holomorphic canonical line bundle K_X of X . In general, the canonical bundle of an abstract CR manifold might not even be locally CR trivialisable, as explained e.g. in [9, 22, 28].

For a CR submanifold M of an N -dimensional projective space $\mathbb{C}P^N$, the restrictions $\mathcal{O}_M(-1)$ and $\mathcal{O}_M(1)$ to M of its tautological and hyperplane bundles are CR line bundles on M , dual to each other. By the Euler sequence, the canonical bundle $K_{\mathbb{C}P^N}$ of the projective space is isomorphic to the $(N+1)$ -th power $\mathcal{O}_{\mathbb{C}P^N}(-N-1)$ of the tautological bundle (see e.g. [19, p.146]) and therefore its dual $K_{\mathbb{C}P^N}^*$, being isomorphic to the $(N+1)$ -th power $\mathcal{O}_{\mathbb{C}P^N}(N+1)$ of the hyperplane bundle $\mathcal{O}_{\mathbb{C}P^N}(1)$, which is very ample, is very ample. In particular, if M is a generic CR submanifold of $\mathbb{C}P^N$, then its anti-canonical bundle K_M^* is the restriction to M of a very ample line bundle. If E_X is a very ample line bundle on a complex compact complex manifold X , then its

²This is obtained by identifying F with its vertical bundle and first defining $\bar{\partial}_M^F$ on the smooth local sections s of F by setting

$$(\dagger) \quad \bar{\partial}_M^F s(X + iJ_M X) = ds(X) + J_F ds(J_M X), \quad \forall X \in HM.$$

In fact, when s is a section, the right hand side of (\dagger) is vertical.

first Chern class $c_1(E_X)$ is positive and the Kodaira-Nakano theorem tells us that this property is sufficient to ensure that the cohomology groups $H^q(X, K_X \otimes E_X)$ vanish for $q > 0$ (see e.g. [19, p.154]). On the other hand, it is known that also the anti-canonical bundles of the generalized complex flag manifolds are positive (see e.g. [6, 8]). They are therefore examples of the larger class of *Fano varieties*, which are characterized by the ampleness of their anti-canonical bundle K_X^* .

When X is a flag manifold of a complex semisimple Lie group \mathbf{G} , the minimal orbits in X of its real forms $\mathbf{G}_{\mathbb{R}}$, studied e.g. in [3, 37], are examples of generic *CR* submanifolds of Fano varieties.

3.3. Deformations and *CR* line bundles. A Cartier divisor in X is the effective divisor of a section of a line bundle F_X on X . A positive F_X has a power which is very ample and therefore the support of its Cartier divisor is the support of the pole divisor of a global meromorphic function f on X .

Let X be a complex space, f a global meromorphic function on X . The corresponding complex line bundle F_X can be described in the following way: Let $\{U_i\}$ be an open covering of X by connected open subsets with the property that, for each index i , there are relatively prime $f'_i, f''_i \in \mathcal{O}_X(U_i)$ such that $f'_i/f''_i = f$ on U_i . The quotients $f'_i/f'_j = f''_i/f''_j = g_{i,j}$ define nowhere vanishing holomorphic functions on $U_i \cap U_j$, which are the transition functions of a holomorphic line bundle F_X on the trivialization atlas $\{U_i\}$. Then

$$(3.2) \quad D = \bigcup_i \{p \in U_i \mid f''_i(p) = 0\}$$

is the support of the pole divisor of f .

Proposition 3.2. *Let X be a complex space, D the support of the pole divisor of a global meromorphic function f on X and F_X the corresponding line bundle. Let M be a smooth *CR* submanifold of X and $\{M_t\}$ a deformation on M of the *CR* structure M_0 induced on M by X such that for each t the *CR* structures of M_t and M_0 agree to infinite order on $D \cap M$. Assume that the *CR* embedding $M_0 \hookrightarrow X$ is generic. Then we can define *CR* line bundles F_{M_t} on each M_t in such a way that $\{F_{M_t}\}$ is a smooth deformation of F_M , and their *CR* structures agree to infinite order on the fibers over D .*

Proof. We keep the notation introduced at the beginning of the subsection. The restrictions of the transition functions $g_{i,j}$ define a *CR* line bundle on M_0 , because

$$g_{i,j}^{-1} \bar{\partial}_M g_{i,j} = 0 \quad \text{on } U_i \cap U_j \cap M.$$

Let us consider now an element of the deformation $\{M_t\}$. We note that

$$g_{j,i} \bar{\partial}_{M_t} g_{i,j} = \frac{f''_j f'_j \bar{\partial}_{M_t} f''_i - f''_i \bar{\partial}_{M_t} f'_j}{f''_i f''_j} = \frac{\bar{\partial}_{M_t} f''_i}{f''_i} - \frac{\bar{\partial}_{M_t} f'_j}{f'_j}, \quad \text{on } U_i \cap U_j \cap M$$

is the difference of two smooth $\bar{\partial}_{M_t}$ -closed forms, the first one defined on $U_i \cap M$, the second on $U_j \cap M$. Indeed

$$\phi_i^{(t)} = \begin{cases} f_i''^{-1} \cdot \bar{\partial}_{M_t} f_i'', & \text{on } (U_i \cap M) \setminus D, \\ 0, & \text{on } U_i \cap M \cap D, \end{cases}$$

is a smooth $\bar{\partial}_{M_t}$ -closed $(0, 1)$ form on $U_i \cap M$, because $\bar{\partial}_{M_t} f_i''$ vanishes to infinite order on $D \cap U_i \cap M$ by the assumption that the CR structures of M and M_t agree to infinite order on $D \cap M$, while f_i'' , which extends holomorphically to the neighbourhood U_i of $U_i \cap M$ in X , cannot vanish to infinite order at any point of $D \cap M$.

The complex smooth line bundle F_M is obtained by gluing the trivial line bundles $\{U_i \times \mathbb{C}\}$ by the transition functions $\{g_{i,j}\}$. Let σ_i be the section $U_i \times \{1\}$ on U_i . The CR structure on F_{M_t} is obtained by setting

$$(3.3) \quad \bar{\partial}_{M_t}^F \sigma_i = \phi_i^{(t)} \cdot \sigma_i \quad \text{on } U_i \text{ for all } i.$$

Clearly it agrees to infinite order on D with that of F_{M_0} . \square

4. AN EXAMPLE

Let m be an integer ≥ 2 and M the quadric hypersurface of $\mathbb{C}P^m$ described in homogeneous coordinates by

$$M = \{z_0 \bar{z}_0 + z_1 \bar{z}_1 = z_2 \bar{z}_2 + \cdots + z_m \bar{z}_m\}.$$

The rational function $\zeta_1 = z_1/z_0$ has divisor $(\zeta_1) = D_1 - D_0$ with

$$D_0 = \{z_0 = 0\}, \quad D_1 = \{z_1 = 0\} \quad \text{and} \quad D_1 \cap D_2 \cap M = \emptyset.$$

Since the point $[1, 0, \dots, 0]$ does not belong to M , the projection map

$$\pi : \mathbb{C}P^m \setminus \{[1, 0, \dots, 0]\} \ni [z_0, z_1, \dots, z_m] \longrightarrow [z_1, \dots, z_m] \in \mathbb{C}P^{m-1}$$

is well defined on M . Its image is the complement in $\mathbb{C}P^{m-1}$ of the open ball

$$B = \{z_2 \bar{z}_2 + \cdots + z_m \bar{z}_m < z_1 \bar{z}_1\}.$$

Let F be a smooth function on $\mathbb{C}P^{m-1}$ which is holomorphic on B and q any point of its boundary ∂B . If we can find an open neighbourhood V of q in $\mathbb{C}P^{m-1}$ and a solution v to the Cauchy problem

$$\begin{cases} v \in C^\infty(V \setminus \bar{B}) \cap C^0(V \setminus B), \\ \bar{\partial} v = \bar{\partial} F, & \text{on } V \setminus \bar{B}, \\ v = 0, & \text{on } V \cap \partial B, \end{cases}$$

then

$$F' = \begin{cases} F, & \text{on } V \cap B, \\ F - v, & \text{on } V \setminus B \end{cases}$$

continues analytically $F|_B$ to the point q .

There are holomorphic functions on B which can be continued smoothly, but not analytically, to all points of ∂B . Thus we can find $F \in C^\infty(\mathbb{C}P^{m-1})$, holomorphic on B , such that $F|_B$ has no analytic continuation to any point

of $\partial\mathbf{B}$ and moreover, since $\partial\mathbf{B} \cap \{z_1 = 0\} = \emptyset$, we can require that its support does not intersect the hyperplane $\{z_1 = 0\}$. Then $\alpha = \bar{\partial}F$ is a smooth $(0, 1)$ form on $\mathbb{C}\mathbb{P}^{m-1}$ with the properties:

- $\bar{\partial}\alpha = 0$ and $\text{supp}(\alpha) \subset \mathbb{C}\mathbb{P}^{m-1} \setminus \mathbf{B}$, $\text{supp}(\alpha) \cap \{z_1 = 0\} = \emptyset$;
- If V is any open neighbourhood of a point $q \in \partial\mathbf{B}$ and $v \in C^\infty(V \setminus \bar{\mathbf{B}}) \cap C^0(V \setminus \mathbf{B})$ satisfies $\bar{\partial}v = \alpha$ on $V \setminus \bar{\mathbf{B}}$, then v is not 0 at some point of $V \cap \partial\mathbf{B}$.

Let us fix a $(0, 1)$ form α on $\mathbb{C}\mathbb{P}^{m-1}$ with these properties.

We consider the coordinate charts $(U_0, \zeta), (U_1, \xi)$ in $\mathbb{C}\mathbb{P}^m$ with

$$U_0 = \{z_0 \neq 0\} = \mathbb{C}\mathbb{P}^m \setminus D_0, \quad \zeta_1 = \frac{z_1}{z_0}, \quad \zeta_2 = \frac{z_2}{z_0}, \dots, \quad \zeta_m = \frac{z_m}{z_0},$$

$$U_1 = \{z_1 \neq 0\} = \mathbb{C}\mathbb{P}^m \setminus D_1, \quad \xi_1 = \frac{z_0}{z_1}, \quad \xi_2 = \frac{z_2}{z_1}, \dots, \quad \xi_m = \frac{z_m}{z_1}.$$

The pullback $\pi^*\alpha$ vanishes to infinite order on the points of $M \cap D_0$, which are mapped by π onto the boundary of \mathbf{B} . Then $\zeta_1 \cdot \pi^*\alpha$, defined on $M \setminus D_0$, extends to a smooth $(0, 1)$ form β on M , vanishing to infinite order on $M \cap D_0$, whose support is contained in $U_1 \cap M$.

We can define a smooth CR structure M_t on M by requiring that its canonical bundle has sections

$$\begin{cases} s_{t,0} = d\zeta_1 \wedge d\zeta_2 \wedge \dots \wedge d\zeta_m, & \text{on } U_0 \cap M \setminus \text{supp}(\beta), \\ s_{t,1} = (d\xi_1 + t\beta) \wedge d\xi_2 \wedge \dots \wedge d\xi_m, & \text{on } U_1 \cap M. \end{cases}$$

Note that

$$\begin{aligned} ds_{t,1} &= t \left(-\frac{d\xi_1}{\xi_1^2} \wedge \pi^*(\alpha) + \frac{1}{\xi_1} \pi^*(d\alpha) \right) \wedge d\xi_2 \wedge \dots \wedge d\xi_m \\ &= -t \frac{d\xi_1}{\xi_1} \wedge \beta \wedge d\xi_2 \wedge \dots \wedge d\xi_m = \frac{t}{\xi_1} \beta \wedge s_{t,1} \end{aligned}$$

because

$$(\dagger) \quad \begin{cases} \pi^*(d\alpha) \wedge d\xi_2 \wedge \dots \wedge d\xi_m = \pi^*(d\alpha \wedge d\xi_2 \wedge \dots \wedge d\xi_m) \\ = \pi^*((\bar{\partial}\alpha) \wedge d\xi_2 \wedge \dots \wedge d\xi_m) = 0. \end{cases}$$

Note that the form $\beta/\xi_1 = \pi^*\alpha/\xi_1^2$ extends to a smooth form β' , defined on M and vanishing to infinite order on D_0 , so that

$$(\ddagger) \quad ds_{t,0} = 0 \text{ on } U_0 \cap M \setminus \text{supp}(\beta) \text{ and } ds_{t,1} = t \cdot \beta' \wedge s_{t,1} \text{ on } U_1 \cap M$$

shows that the line bundles K_{M_t} define formally integrable complex valued distributions.

These CR structures agree to infinite order on $M \cap D_0$ with the CR structure M_0 on M which is induced by its embedding into $\mathbb{C}\mathbb{P}^m$

Their canonical bundles are isomorphic complex line bundles, since the transition functions are in all cases $g_{0,1} = s_{t,0}/s_{t,1} = -\zeta_1^{m+1} = -\xi_1^{-m-1}$, $g_{1,0} = g_{0,1}^{-1}$ on $M \setminus (D_0 \cup D_1 \cup \text{supp}(\beta))$.

We note that ξ_2, \dots, ξ_m are CR functions on $M_t \setminus D_0$ for all t . A necessary condition for M_t being locally embeddable at a point p of D_0 is that

one could find an open neighbourhood U' of p in $U_1 \cap M$ and a function $u \in C^\infty(U')$ such that

$$d(e^u s_{t,1}) = 0 \quad \text{on } U'.$$

This equation can be rewritten, by factoring out $e^u \zeta_1$ and taking into account (\dagger), as

$$(*) \quad \left(\xi_1 du + \frac{t}{\xi_1} \pi^* \alpha \right) \wedge d\xi_2 \wedge \cdots \wedge d\xi_m = 0 \quad \text{on } U' \setminus D_0.$$

The fibers of π above points of $\mathbb{C}P^{m-1} \setminus \bar{B}$ are circles and reduce to single points over ∂B . Since the projection π is proper, by shrinking we may assume that $\pi^{-1}(\pi(U')) = U'$. By integrating over the fibers we define

$$v(q) = \frac{-1}{2\pi i} \oint_{\pi^{-1}(q)} u \xi_1 d\xi_1, \quad \text{for } q \in \pi(U') \setminus \partial B.$$

This is a smooth function on $\pi(U') \setminus \partial B$, which extends to a continuous function, that we still denote by v , vanishing on $\pi(U') \cap \partial B$. Then from (*) we obtain that

$$dv \wedge d\xi_2 \wedge \cdots \wedge d\xi_m = t \alpha \wedge d\xi_2 \wedge \cdots \wedge d\xi_m, \quad \text{on } \pi(U') \setminus \bar{B}.$$

This is equivalent to

$$\bar{\partial}v = t \alpha, \quad \text{on } \pi(U') \setminus \bar{B}$$

and hence contradicts the choice of α if $t \neq 0$, proving that M_t cannot be locally embedded at any point $p \in D_0$ if $t \neq 0$.

This example from [24] shows that we can construct a smooth deformation $\{M_t\}$ of the CR structure M_0 of the quadric M such that no M_t can be locally embedded into a complex manifold at any point of D_0 if $t \neq 0$.

The CR structure of the M_t 's are of type $(m-1, 1)$ and M_0 is 1-pseudoconcave. Hence the example shows that Theorem 1.1 cannot be valid if we only require M_0 being 1-pseudoconcave.

Remark 4.1. Equations (\ddagger) is of the type of those we need to solve to construct projective embeddings in §6. It can be rewritten as

$$\begin{cases} \bar{\partial}_{M_t}^K s_{t,0} = 0, \\ \bar{\partial}_{M_t}^K s_{t,1} = t \cdot \beta' s_{t,1}. \end{cases}$$

We also have

$$\bar{\partial}_{M_t} u = \bar{\partial}_{M_0} u - t \cdot \frac{\partial u}{\partial \xi_1} \beta.$$

Then we obtain

$$\bar{\partial}_{M_t}^K \left(\frac{1}{\xi_1} s_{t,1}^{-1} \right) = -t \left(\frac{1}{\xi_1^2} \beta + \frac{1}{\xi_1} \beta' \right) s_{t,1}^{-1} = -2 \frac{t}{\xi_1^3} \pi^* \alpha s_{t,1}^{-1}.$$

An argument similar to the one used above shows that the equation

$$(**) \quad \bar{\partial}(w \cdot s_{t,1}^{-1}) = (\bar{\partial}_{M_0} w - t \frac{\partial w}{\partial \xi_1} \beta - t \cdot w \beta') s_{t,1}^{-1} = -2 \frac{t}{\xi_1^3} \pi^* \alpha s_{t,1}^{-1}$$

has no smooth solution on M . Indeed (***) can be rewritten in the form

$$\left(\mathbf{d}w - \frac{t}{\xi_1^2} \frac{\partial(\xi_1 w)}{\partial \xi_1} \pi^* \alpha + \frac{2t}{\xi_1^3} \pi^* \alpha \right) \mathbf{d}\xi_2 \wedge \cdots \wedge \mathbf{d}\xi_m = 0$$

Multiplying by ξ_1^2 and integrating on the fiber, we see that the term in the middle vanishes, while

$$v(q) = \frac{-1}{4\pi i} \oint_{\pi^{-1}(q)} (\xi_1^2 w) \mathbf{d}\xi_1$$

yields a solution in $C^\infty(\mathbb{C}\mathbb{P}^{m-1} \setminus \bar{\mathbf{B}}) \cap C^0(\mathbb{C}\mathbb{P}^{m-1} \setminus \mathbf{B})$, vanishing on $\partial\mathbf{B}$, of the equation $\bar{\partial}v = \alpha$ on $\mathbb{C}\mathbb{P}^{m-1} \setminus \bar{\mathbf{B}}$, contradicting the choice of α .

This adds a further argument for the need of conditions stronger than 1-pseudoconcavity for the validity of the conclusions of Theorem 1.1.

5. VANISHING RESULTS

We keep the notation introduced in §2, §3.

5.1. Embedded CR manifolds. We have the following vanishing result:

Theorem 5.1. *Assume that M is a 2-pseudoconcave generic smooth CR submanifold of type (n, k) of a complex variety X . Let F_X be a positive line bundle on X and F_M its restriction to M . Then we can find a positive integer ℓ_0 such that*

$$(5.1) \quad H^{p,q}(M, F_M^{-\ell}) = 0, \quad \forall \ell \geq \ell_0, \quad 0 \leq p \leq n+k, \quad q = 0, 1.$$

Proof. We know from [21] that a generic 2-pseudoconcave M has a fundamental system of tubular neighborhoods U in X that are $(n-2)$ -pseudoconvex and $(n+k-2)$ -pseudoconcave in the sense of Andreotti-Grauert³ and for which, for all $0 \leq p \leq n+k$, the natural restriction maps

$$H^q(U, \Omega^p(F_U)) \longrightarrow H^q(M, \Omega^p(F_M)), \quad q = 0, 1,$$

are isomorphisms for any holomorphic line bundle F_U over U .

This was proved in [21, Thm.4.1] for the trivial line bundle. The arguments of the proof given there apply also in general, yielding the isomorphism for arbitrary holomorphic vector bundles. The statement is then a consequence of [5, Prop.27], because U is *strongly $(n+k-2)$ -concave* in the sense of Andreotti-Grauert and hence also $F_U^{-1} := F_X^{-1}|_U$ is strongly $(n+k-2)$ -concave as a complex space, according to [5] and therefore there is a positive integer ℓ_0 such that $H^{p,q}(U, F_X^{-\ell}) = 0$ for $\ell \geq \ell_0$, $0 \leq p \leq n+k$ and $q = 0, 1$. \square

³A real valued smooth function ϕ on an N -dimensional complex manifold X is strongly q -pseudoconvex in the sense of [5] at points where its complex Hessian has at least $(N-q+1)$ positive eigenvalues. Then X is called *strictly q -pseudoconvex* if there is an exhaustion function $\phi \in C^\infty(X, \mathbb{R})$ which is strictly q -pseudoconvex outside a compact subset of X and strictly q -pseudoconcave if there is an exhaustion function $\phi \in C^\infty(X, \mathbb{R})$ such that $(-\phi)$ is strictly q -pseudoconvex outside a compact subset of X .

5.2. Some estimates for abstract CR manifolds. Let M be an abstract compact CR manifold of type (n, k) and F_M a complex CR line bundle on M . To discuss its cohomology groups $H^{p,q}(M, F_M)$ by using functional analytic methods, it is convenient to fix a smooth Riemannian metric on M and a smooth Hermitian norm on the fibers of F_M . In this way we are able, for each real $r \geq 0$, to define in a standard way on the smooth sections of $Q_M^{p,q} \otimes F_M$ the Sobolev W^r -norm, involving the L^2 -norms of their fractional derivatives up to order r . The completions of $\mathcal{F}^{p,q}(M)$ with respect to these norms are Hilbert spaces, that we denote by $W^r(M, Q_M^{p,q} \otimes F_M)$. We use the notation

$$(v_1 | v_2)_r \quad \text{and} \quad \|v\|_r = \sqrt{(v|v)_r}, \quad \text{for } v_1, v_2, v \in W^r(M, Q_M^{p,q} \otimes F_M)$$

for their Hermitian scalar products and norms (see e.g. [20]).

In [21] subelliptic estimates were proved for the $\bar{\partial}_M$ complex under pseudoconcavity conditions. Since the arguments are local, they trivially extend to forms with coefficients in a complex CR line bundle. We have in particular the following statement.

Proposition 5.2. *Let M be a compact smooth abstract CR manifold of type (n, k) and F_M any complex CR line bundle on M . Assume that M is 2-pseudoconcave. Then, for every $0 \leq p \leq n+k$ and every real $r \geq 0$ there is a positive constant $c_{r,p} > 0$ such that*

$$(5.2) \quad \begin{cases} c_{r,p} \cdot \|u\|_{r+(1/2)} \leq \|\bar{\partial}_M^F u\|_r^2, & \forall u \in \mathcal{F}^{p,0}(M), \\ c_{r,p} \cdot \|v\|_{r+(1/2)} \leq \|\bar{\partial}_M^F v\|_r^2 + \|(\bar{\partial}_M^F)^* v\|_r^2 + \|v\|_0^2, & \forall v \in \mathcal{F}^{p,1}(M). \quad \square \end{cases}$$

Let us define the subspace

$$(5.3) \quad \mathcal{N}_F^{p,1}(M) = \{v \in \mathcal{F}^{p,1}(M) \mid \bar{\partial}_M^F v = 0, \quad (\bar{\partial}_M^F)^* v = 0\}.$$

Keeping the assumptions and the notation above, we have

Proposition 5.3. *Under the assumptions of Proposition 5.2:*

- The space $\mathcal{N}_F^{p,1}(M)$ is finite dimensional and equals its weak closure with respect to the L^2 norm;
- For every pair of real numbers r, r' with $-(1/2) \leq r' < r, r \geq 0$, there is a constant $C_{r,r',p} > 0$ such that

$$(5.4) \quad \begin{cases} \|v\|_{r'+(1/2)} \leq C_{r,r',p} \left(\|\bar{\partial}_M^F v\|_r^2 + \|(\bar{\partial}_M^F)^* v\|_r^2 \right), \\ \forall v \in \mathcal{F}^{p,1}(M) \cap (\mathcal{N}_F^{p,1}(M))^\perp. \end{cases}$$

[The orthogonal is taken with respect to the L^2 -scalar product on $\mathcal{F}^{p,1}(M)$.]

- $H^{p,1}(M, F_M) \simeq \mathcal{N}_F^{p,1}(M)$.

Proof. The first statement is a consequence of (5.2): this subelliptic estimate for $r=0$ implies in particular that the L^2 and $W^{1/2}$ -Sobolev norms are equivalent on $\mathcal{N}_F^{p,1}(M)$. Then by Rellich's theorem the unit L^2 ball in $\mathcal{N}_F^{p,1}(M)$ is compact and this implies that $\mathcal{N}_F^{p,1}(M)$ is finite dimensional and thus also equals its weak closure with respect to the L^2 norm.

The second statement can be proved by contradiction: assume that there are real numbers r, r' , with $-(1/2) \leq r' < r, r \geq 0$, for which (5.5) is not valid for any positive constant $C_{r,r',p}$.

Then we can find a sequence $\{v_\nu\} \subset \mathcal{F}^{p,1}(\mathbb{M}) \cap (\mathcal{N}_F^{p,1}(\mathbb{M}))^\perp$ with

$$\|v_\nu\|_{r'+(1/2)} = 1, \quad \|\bar{\partial}_M^F v\|_r^2 + \|(\bar{\partial}_M^F)^* v\|_r^2 < 2^{-\nu}.$$

By (5.2) this sequence is bounded in the $W^{r+(1/2)}$ -norm and therefore has a subsequence $\{v_\nu\}$ which, weakly in $W^{r+(1/2)}$ and hence strongly in $W^{r'+(1/2)}$, converges to a v_∞ . In particular $\|v_\infty\|_{r'+(1/2)} = 1$ shows that $v_\infty \neq 0$. This limit is a weak solution of $\bar{\partial}_M^F v_\infty = 0$ and $(\bar{\partial}_M^F)^* v_\infty = 0$ and therefore, by the hypoellipticity of $\bar{\partial}_M^F \oplus (\bar{\partial}_M^F)^*$ following from the subellipticity estimate (5.2), is a smooth section and thus a nonzero element of $\mathcal{N}_F^{p,1}(\mathbb{M})$.

This completes the proof of the second statement.

It is clear that the elements of $\mathcal{N}_F^{p,1}(\mathbb{M})$ represent cohomology classes in $H^{p,1}(\mathbb{M}, \mathbb{F}_M)$. Indeed, if $v \in \mathcal{N}_F^{p,1}$ and $v = \bar{\partial}_M^F u$ for some $u \in \mathcal{F}^{p,0}(\mathbb{M})$, then

$$(\bar{\partial}_M^F)^* \bar{\partial}_M^F u = 0 \implies \|v\|_0^2 = (v | \bar{\partial}_M^F u)_0 = ((\bar{\partial}_M^F)^* v | u)_0 = 0$$

shows that $v = 0$. On the other hand, if $f \in \mathcal{F}^{p,1}(\mathbb{M})$ and $\bar{\partial}_M^F f = 0$, then we can decompose f into the sum $f = f_0 + f_1$, with $f_0 \in \mathcal{N}_F^{p,1}(\mathbb{M})$ and $f_1 \in (\mathcal{N}_F^{p,1}(\mathbb{M}))^\perp$.

We get

$$\begin{aligned} |(f_1 | v)_0| &\leq \|f_1\|_0 \|v\|_0 \leq C_{-1/2,0,p} (\|\bar{\partial}_M^F v\|_0^2 + \|(\bar{\partial}_M^F)^* v\|_0^2)^{1/2}, \\ &\forall v \in \mathcal{F}^{p,1}(\mathbb{M}) \cap (\mathcal{N}_F^{p,1}(\mathbb{M}))^\perp. \end{aligned}$$

By Riesz' representation theorem there is a unique $w \in \mathcal{F}^{p,1}(\mathbb{M}) \cap (\mathcal{N}_F^{p,1}(\mathbb{M}))^\perp$ (w is smooth because of the subellipticity estimate (5.2)) such that

$$(\bar{\partial}_M^F w | \bar{\partial}_M^F v)_0 + ((\bar{\partial}_M^F)^* w | (\bar{\partial}_M^F)^* v)_0 = (f_1 | v), \quad \forall v \in \mathcal{F}^{p,1}(\mathbb{M}) \cap (\mathcal{N}_F^{p,1}(\mathbb{M}))^\perp.$$

Since f_1 is L^2 -orthogonal to $\mathcal{N}_F^{p,1}(\mathbb{M})$, this equality holds true for all v in $\mathcal{F}^{p,1}(\mathbb{M})$ and integration by parts yields

$$(\bar{\partial}_M^F)^* \bar{\partial}_M^F w = 0, \quad \bar{\partial}_M^F (\bar{\partial}_M^F)^* w = f_1.$$

This shows that the orthogonal projection of $\ker(\bar{\partial}_M^F : \mathcal{F}^{p,1}(\mathbb{M}) \rightarrow \mathcal{F}^{p,2}(\mathbb{M}))$ onto $\mathcal{N}_F^{p,1}(\mathbb{M})$ yields, by passing to the injective quotient, an isomorphism between $H^{p,1}(\mathbb{M}, \mathbb{F})$ and $\mathcal{N}_F^{p,1}(\mathbb{M})$.

The proof is complete. \square

Likewise, we have a similar statement for CR sections of \mathbb{F} on \mathbb{M} . Set

$$\mathcal{N}_F^{p,0}(\mathbb{M}) = \{u \in \mathcal{F}^{p,0}(\mathbb{M}) \mid \bar{\partial}_M^F u = 0\}.$$

Proposition 5.4. *Under the assumptions of Proposition 5.2:*

- The space $\mathcal{N}_F^{p,0}(\mathbb{M})$ is finite dimensional and equals its weak closure with respect to the L^2 norm;

- For every pair of real numbers r, r' with $-(1/2) \leq r' < r, r \geq 0$, there is a constant $C_{r,r',p} > 0$ such that

$$(5.5) \quad \|u\|_{r'+(1/2)} \leq C_{r,r',p} \|\bar{\partial}_M^F u\|_r^2, \quad \forall u \in \mathcal{F}^{p,0}(M) \cap (\mathcal{N}_F^{p,0}(M))^\perp.$$

5.3. Deformation of tangential CR complexes. Let us consider the situation described at the beginning of §5.2. Here we make the additional assumption that M is a generic compact smooth CR submanifold of a complex variety X of dimension $n+k$ and that the complex CR line bundle F_M is the pullback of a holomorphic line bundle F_X on X , associated to a global meromorphic function on X , whose pole divisor has support D . We indicate by M_0 the CR structure, of type (n, k) , induced on M by the embedding $M \hookrightarrow X$ and consider a smooth one-parameter deformation $\{M_t\}$ on M of this CR structure M_0 , with the constraint:

$$(5.6) \quad \text{The CR structures of } M_t \text{ agree to infinite order on } M \cap D.$$

We observed in §3.3 that in this case F_M can be also considered as a CR line bundle on each M_t . We need to consider the different tangential Cauchy-Riemann complexes with coefficients in F_{sfM} as complexes of partial differential operators acting on the same spaces of vector valued functions.

We showed in §3.1 that we can represent the tangential CR complex (on the trivial line bundle) by fixing a complement $E_M = E_M^{0,1}$ of $T_M^{1,0}$. This yields isomorphisms between $E_M^{p,q} = \Lambda^q(E_M) \wedge_M \Lambda^p(T_M^{1,0})$ and $Q_M^{p,q}$ for all integers p, q with $0 \leq p \leq n+k, 0 \leq q \leq n$.

We will be interested in deformations M_t corresponding to *small* values of the parameter t . In particular, after shrinking, we can assume that E_M is transversal to $T_{M_t}^{1,0}$ for all values of the parameter t . Then the tangential CR complexes

$$0 \longrightarrow \mathcal{E}_M^{0,0} \xrightarrow{d''_{M_t}} \mathcal{E}_M^{0,1} \xrightarrow{d''_{M_t}} \dots \rightarrow \mathcal{E}_M^{0,n-1} \xrightarrow{d''_{M_t}} \mathcal{E}_M^{0,n} \longrightarrow 0$$

are defined as complexes of linear partial differential operators, smoothly depending on the parameter t , which act on the germs of sections of the same fiber bundles.

To obtain an analogous presentation for positive values of p , we note that every τ in $T_{M_t}^{1,0}$ uniquely decomposes into the sum $\tau' + \tau''$ of a $\tau' \in T_{M_0}^{1,0}$ and $\tau'' \in E_M$. The correspondence $\tau \leftrightarrow \tau'$ yields smooth isomorphisms $Q_{M_t}^{p,0} \simeq Q_{M_0}^{p,0}$ of complex vector bundles, from which we obtain also for positive p representations

$$0 \longrightarrow \mathcal{E}_M^{p,0} \xrightarrow{d''_{M_t}} \mathcal{E}_M^{p,1} \xrightarrow{d''_{M_t}} \dots \rightarrow \mathcal{E}_M^{0,n-1} \xrightarrow{d''_{M_t}} \mathcal{E}_M^{p,n} \longrightarrow 0$$

of the tangential CR complexes on M_t by linear partial differential operators acting on the same complex vector bundles and smoothly depending on the parameter t .

Finally, for the line bundle F_M , we use the trivialization given by the sections σ_i of §3.3 and (3.3) to compute their $\bar{\partial}_{M_t}^F$. The form $\phi_i^{(t)} \in Q^{0,1}(U_i)$ decomposes into the sum of a $\psi_i^{(t)} \in \mathcal{E}^{0,1}(U_i)$ and a form in $\Gamma^\infty(U_i, T_{M_t}^{1,0})$.

Then we can define for every $0 \leq p \leq n+k$ and $0 \leq q < n$ and all values of the parameter t a linear partial differential operators $d_{M_t}^{\prime\prime F}$ mapping germs of smooth sections of $F_M \otimes E_M^{p,q}$ to germs of smooth sections of $F_M \otimes E_M^{p,q+1}$ in such a way that

$$d_{M_t}^{\prime\prime F}(\alpha \cdot \sigma_i) = (d_{M_t}^{\prime\prime} \alpha + (-1)^{p+q} \psi_i^{(t)} \wedge \alpha) \cdot \sigma_i, \quad \forall i, \forall \alpha \in \mathcal{E}_M^{p,q}(U_i).$$

In this way we represent the different tangential CR complexes for forms with coefficients in F_M as complexes of differential operators, smoothly depending on the parameter t , but acting on the same smooth complex vector bundles:

$$(5.7) \quad \begin{aligned} 0 \rightarrow \mathcal{E}^{p,0}(M, F_M) &\xrightarrow{d_{M_t}^{\prime\prime F}} \mathcal{E}^{p,1}(M, F_M) \rightarrow \dots \\ \dots \rightarrow \mathcal{E}^{p,n-1}(M, F_M) &\xrightarrow{d_{M_t}^{\prime\prime F}} \mathcal{E}^{p,n}(M, F_M) \rightarrow 0. \end{aligned}$$

5.4. Estimates for deformations. We keep the notation and assumptions of §5.3. Our goal is to show that the results of Propositions 5.2, 5.3 remain valid for small deformations.

Proposition 5.5. *Let M be a compact generic CR submanifold of type (n, k) of a complex variety X . Let F_X be the holomorphic line bundle associated to a global meromorphic function on X and D the support of its polar divisor. We assume that M is 2-pseudoconcave. Let $\{M_t\}$ be a deformation of the CR structure induced on $M=M_0$ by the embedding into X , smoothly depending on the parameter t and with CR structures agreeing to infinite order on $D \cap M$. Then we can find $\epsilon_0 > 0$ such that for all $|t| < \epsilon_0$*

- (1) *for every $0 \leq p \leq n+k$ and every real $r \geq 0$ there is a positive constant $c_{r,p} > 0$ such that*

$$(5.8) \quad \begin{cases} c_{r,p} \cdot \|u\|_{r+(1/2)}^2 \leq \|d_{M_t}^{\prime\prime F} u\|_r^2 + \|u\|_0^2, & \forall u \in \mathcal{F}^{p,0}(M) \\ c_{r,p} \cdot \|v\|_{r+(1/2)}^2 \leq \|d_{M_t}^{\prime\prime F} v\|_r^2 + \|(d_{M_t}^{\prime\prime F})^* v\|_r^2 + \|v\|_0^2, & \forall v \in \mathcal{F}^{p,1}(M). \end{cases}$$

- (2) *for every pair r', r of real numbers with $-(1/2) \leq r' < r$, $r \geq 0$ and $0 \leq p \leq n+k$, there is a positive constant $C_{r,r',p} > 0$ such that*

$$(5.9) \quad \begin{cases} \|u\|_{r'+(1/2)}^2 \leq C_{r,r',p} \|d_{M_t}^{\prime\prime F} u\|_r^2 & \forall u \in \mathcal{F}^{p,0}(M) \cap (\mathcal{N}_F^{p,0}(M_t))^\perp \\ \|v\|_{r'+(1/2)}^2 \leq C_{r,r',p} (\|d_{M_t}^{\prime\prime F} v\|_r^2 + \|(d_{M_t}^{\prime\prime F})^* v\|_r^2) & \\ & \forall v \in \mathcal{F}^{p,1}(M) \cap (\mathcal{N}_F^{p,1}(M_t))^\perp. \end{cases}$$

- (3) *$\dim_{\mathbb{C}}(H^{p,0}(M_t, F_M)) \leq \dim_{\mathbb{C}}(H^{p,0}(M_0, F_M)) < +\infty$ and $\dim_{\mathbb{C}}(H^{p,1}(M_t, F_M)) \leq \dim_{\mathbb{C}}(H^{p,1}(M_0, F_M)) < +\infty$ for all $0 \leq p \leq n+k$.*

Proof. The proof of the subelliptic estimates in [21] yield (5.8), the constants being uniform for $|\mathfrak{t}| < \epsilon_0$ because they depend upon the coefficients of the linear differential operators $\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}}$ and their derivatives.

We prove (2) and (3) in the case of forms ($q = 1$), as the case of sections ($q=0$) is analogous and even simpler.

We begin by proving that, for any pair r, r' with $(-1/2) < r' < r$, $r \geq 0$ and for all $0 \leq p \leq n+k$, we can find $\epsilon > 0$ and $C_{r,r',p} > 0$ such that

$$(5.10) \quad \begin{aligned} \|v\|_{r'+(1/2)}^2 &\leq C_{r,r',p} \left(\|\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}} v\|_r^2 + \|(\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}})^* v\|_r^2 \right), \\ \forall |\mathfrak{t}| < \epsilon, \quad \forall v &\in \mathcal{E}^{p,1}(M, \mathbb{F}_M) \cap (\mathcal{N}_{\mathbb{F}}^{p,1}(M_0))^\perp. \end{aligned}$$

We argue by contradiction: if this was false, we could find a sequence $\{v_\nu\}$ in $\mathcal{E}^{p,1}(M, \mathbb{F}_M) \cap (\mathcal{N}_{\mathbb{F}}^{p,1}(M_0))^\perp$ and a sequence $\{\mathfrak{t}_\nu \rightarrow 0\}$ such that

$$\|v_\nu\|_{r'+(1/2)}^2 = 1, \quad \|\mathfrak{d}_{M_{\mathfrak{t}_\nu}}''^{\mathbb{F}} v_\nu\|_r^2 + \|(\mathfrak{d}_{M_{\mathfrak{t}_\nu}}''^{\mathbb{F}})^* v_\nu\|_r^2 < 2^{-\nu}.$$

By (1), the sequence $\{v_\nu\}$ is uniformly bounded in $W^{r+1/2}$ and then, by the compactness of the inclusion $W^{r+(1/2)} \hookrightarrow W^{r'+(1/2)}$, contains a subsequence which strongly converges in $W^{r'+(1/2)}$ to a v_∞ , which on one hand should be different from 0 because $\|v_\infty\|_{r'+(1/2)} = 1$, on the other hand should be zero because it belongs to $\mathcal{N}_{\mathbb{F}}^{p,1}(M_0) \cap (\mathcal{N}_{\mathbb{F}}^{p,1}(M_0))^\perp$.

Estimate (5.10) implies that $\mathcal{N}_{\mathbb{F}}^{1,p}(M_{\mathfrak{t}}) \cap (\mathcal{N}_{\mathbb{F}}^{p,1}(M_0))^\perp = \{0\}$ for $|\mathfrak{t}| < \epsilon$. This yields (3), because the dimension of $H^{1,p}(M_{\mathfrak{t}}, \mathbb{F})$ equals that of $\mathcal{N}_{\mathbb{F}}^{1,p}(M_{\mathfrak{t}})$.

We can prove (2) by an argument similar to the one used for (5.10). \square

5.5. Vanishing theorems. We will use Proposition 5.5 to solve $\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}} u_{\mathfrak{t}} = f$, for forms f in $\mathcal{E}^{0,1}(M, \mathbb{F}_M)$, with $u_{\mathfrak{t}}$ satisfying uniform estimates in the Sobolev norms W^r , for sufficiently large r , yielding by the immersion theorems control upon the derivatives of $u_{\mathfrak{t}}$.

Theorem 5.6. *Let us keep the assumptions of Proposition 5.5. Assume moreover that \mathbb{F}_M is a positive line bundle on X . Then we can find $\epsilon_0 > 0$ such that*

$$(5.11) \quad H^{p,q}(M_{\mathfrak{t}}, \mathbb{F}_M^{-\ell}) = 0, \quad \forall \ell \geq \ell_0, \quad 0 \leq p \leq n+k, \quad q = 0, 1, \quad |\mathfrak{t}| < \epsilon_0.$$

Moreover, for every $f \in \mathcal{E}^{0,1}(M, \mathbb{F}_M^{-\ell})$ satisfying $\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}} f = 0$ there is a unique $u_{\mathfrak{t}}$ in $\Gamma^\infty(M, \mathbb{F}_M^{-\ell})$ such that, for $r' < r$ and $r \geq 1/2$,

$$(5.12) \quad \mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}} u_{\mathfrak{t}} = f \quad \text{and} \quad \|u_{\mathfrak{t}}\|_{r'} \leq C_{r',r} \|f\|_{r-(1/2)}.$$

Proof. The first statement is a consequence of Theorem 5.1 and Proposition 5.5. These results allow us to fix ϵ_0 in such a way that, by (5.9),

$$\|v\|_{\mathfrak{t}} = \left(\|\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}} v\|_0^2 + \|(\mathfrak{d}_{M_{\mathfrak{t}}}''^{\mathbb{F}})^* v\|_0^2 \right)^{1/2}$$

is a pre-Hilbertian norm on $\mathbb{E}^{0,1}(M_{\mathfrak{t}}, \mathbb{F}_M^{-\ell})$. Let $\mathbf{S}_{\mathfrak{t}}$ be its completion with respect to this norm. Given $f \in \mathbb{E}^{0,1}(M, \mathbb{F}_M^{-\ell})$, we can consider $v \rightarrow (v|f)_0$ as a

linear continuous functional on \mathbf{S}_t and hence by Riesz representation Theorem there is a unique $w \in \mathbf{S}_t$ such that

$$(v|f)_0 = (d''_{M_t} v | d''_{M_t} w)_0 + ((d''_{M_t})^* v | (d''_{M_t})^* w)_0, \quad \forall v \in E^{0,1}(M, F_M^{-\ell}).$$

By the subelliptic estimate (5.8), in fact $w \in E^{0,1}(M, F_M^{-\ell})$, i.e. w is smooth, because we assumed that f is smooth. If f satisfies the integrability condition $d''_{M_t} f = 0$, then integration by parts yields

$$\begin{aligned} 0 &= (d''_{M_t} w | d''_{M_t} f)_0 = ((d''_{M_t})^* d''_{M_t} w | f)_0 \\ &= (d''_{M_t} (d''_{M_t})^* d''_{M_t} w | d''_{M_t} w)_0 + ((d''_{M_t})^* (d''_{M_t})^* d''_{M_t} w | (d''_{M_t})^* w)_0 \\ &= ((d''_{M_t})^* d''_{M_t} w | (d''_{M_t})^* d''_{M_t} w)_0 \implies (d''_{M_t})^* d''_{M_t} w = 0 \\ &\implies 0 = ((d''_{M_t})^* d''_{M_t} w | w)_0 = \|d''_{M_t} w\|_0^2 \implies d''_{M_t} w = 0. \end{aligned}$$

By (5.9), for all $0 < r < 1/2$

$$\begin{aligned} (u_t | u_t)_0 &= ((d''_{M_t})^* w, (d''_{M_t})^* w)_0 = (w, d''_{M_t} (d''_{M_t})^* w)_0 = (w, f)_0 \\ &\leq \|f\|_{-r} \|w\|_r \leq C_r'' \|f\|_{-r} \|u_t\|_{r-\frac{1}{2}} \leq C_r'' \|f\|_{-r} \|u_t\|_0 \end{aligned}$$

Hence

$u_t = (d''_{M_t})^* w$ solves $d''_{M_t} u_t = f$ and $\|u_t\|_0 \leq C_r'' \|f\|_{-r}$, $\forall 0 < r < 1/2$, with constants C_r'' independent of $|t| < \epsilon_0$. Since $\mathcal{O}(M_t, F_M) = \{0\}$, the solution is unique and therefore we obtain the estimates in (5.12) by recurrence, using, for any pseudodifferential operator P on M , the identity

$$d''_{M_t} P(u_t) = [d''_{M_t}, P](u_t) + P(f).$$

If P has order r' , with $0 < r' < 1/2$, then

$$\|P(u_t)\|_0 \leq C_{r'}'' \left(\|[d''_{M_t}, P](u_t)\|_{-r'} + \|P(f)\|_{-r'} \right) \leq \text{const}_P C_{r'}'' (\|u_t\|_0 + \|f\|_0),$$

with a constant const_P which only depends on P . This yields

$$\|u_t\|_{r'} \leq C_{r',(1/2)} \|f\|_0.$$

with constants independent of t . Repeating this argument we obtain the estimate in the statement. \square

6. PROOF OF THEOREM 1.1

6.1. The general case. We keep the notation of §3.3. The (f_i'') define a global section σ of F_M , which is CR for the structure on M_0 and is $\neq 0$ on $M \setminus D$.

Fix an integer ℓ such that F_X^ℓ is very ample and $H^{0,1}(M_t, F_M^{-h}) = 0$ for $h \geq \ell$ and $|t| < \epsilon_0$, which is possible by Theorem 5.6.

For every t we set

$$g_t = \begin{cases} d''_{M_t}^{-\ell} \sigma^{-\ell}, & \text{on } M \setminus D, \\ 0, & \text{on } D. \end{cases}$$

Since the CR structure of M_t agrees with that of M_0 to infinite order on D , the g_t 's are smooth sections in $\mathcal{E}^{0,1}(M, F^{-\ell})$ that vanish to infinite order on

D and satisfy the integrability condition $d''_{M_t} g_t = 0$. By Theorem 5.6, for $|t| < \epsilon_0$, we can find $v_t \in \Gamma^\infty(M, F_M^{-\ell})$ to solve

$$d''_{M_t} v_t = g_t, \quad \text{and we have } \|v_t\|_{r'} \leq C_{r',r} \|g_t\|_{r-(1/2)}$$

for $0 \leq r' < r$, with a constant $C_{r,r'}$ independent of t . Then

$$\sigma^{-\ell} - v_t \in \mathcal{O}_{M_t}(M_t \setminus D, F_M^{-\ell}).$$

By Sobolev's embedding theorems, the sup-norms of the first derivatives of the $v_{i,t}$ are bounded by their $W^{r'}$ -norms for $r' > (2n+k+2)/2$. As $\|g_{i,t}\|_{r-(1/2)} \rightarrow 0$ for $t \rightarrow 0$, we obtain that, for $|t| \ll 1$, the sections

$$\eta_t = (\sigma^{-\ell} - v_t)^{-1} \in \mathcal{O}_{M_t}(M_t \setminus D, F_M^\ell).$$

extend to smooth sections of $\mathcal{O}_{M_t}(M_t, F_M^\ell)$, defining the divisor D .

Since F_X^ℓ is very ample, there are sections $\sigma_0, \sigma_1, \dots, \sigma_m \in \mathcal{O}_{M_0}(X, F_X^\ell)$ providing a holomorphic embedding of X into $\mathbb{C}P^m$.

Their restrictions $(\sigma_{0,0}, \sigma_{1,0}, \dots, \sigma_{m,0})$ to M provide a CR embedding of M_0 into $\mathbb{C}P^m$. For every t and $1 \leq i \leq m$ set

$$f_{i,t} = \begin{cases} d''_{M_t} (\eta_t^{-2} \sigma_i), & \text{on } M \setminus D, \\ 0, & \text{on } D. \end{cases}$$

Then, arguing as above and invoking Theorem 5.6 for $|t| < \epsilon_0$, we can find $u_{i,t} \in \Gamma^\infty(M, F_M^{-\ell})$ to solve

$$d''_{M_t} u_{i,t} = f_{i,t}, \quad \text{and we have } \|u_{i,t}\|_{r'} \leq C_{r',r} \|f_{i,t}\|_{r-(1/2)}$$

for $0 \leq r' < r$, with a constant $C_{r,r'}$ independent of t . Then

$$\sigma_{i,t} = \sigma_i - u_{i,t} \eta_t^2 \in \mathcal{O}_{M_t}(M_t, F_M^\ell).$$

As $\|f_{i,t}\|_{r-(1/2)} \rightarrow 0$ for $t \rightarrow 0$, we obtain that, for $|t| \ll 1$, the sections $(\sigma_{0,t}, \sigma_{1,t}, \dots, \sigma_{m,t})$, being a small C^1 -perturbation of $(\sigma_{0,0}, \sigma_{1,0}, \dots, \sigma_{m,0})$, still provide a CR immersion of M_t into $\mathbb{C}P^m$.

The last part of the statement follows from [23], where it is shown that the maximum degree of transcendence of the field of CR meromorphic functions on M_t is $n+k$: this implies that all pseudoconcave CR manifold of type (n, k) having a projective embedding, can be embedded into a projective complex variety of dimension $n+k$.

6.2. Generic CR submanifolds of the projective space. The line bundles on the projective space $\mathbb{C}P^{n+k}$ are parametrized (modulo equivalence) by the integers. They are all holomorphically equivalent to integral powers of the line bundle $\mathcal{O}_{\mathbb{C}P^{n+k}}$, which can be described, in the covering $\{U_i = \{z_i \neq 0\}\}$ by the transition functions $\{g_{i,j} = z_j^{-1} z_i\}$. The support of the pole divisor of a meromorphic function f on $\mathbb{C}P^{n+k}$ is the set of common zeros of a homogeneous polynomial

$$(6.1) \quad D = \{\wp(z_0, z_1, \dots, z_{n+k}) = 0\}.$$

If $\wp(z)$ has degree d , this is the zero locus of a section of $\Theta_{\mathbb{C}\mathbb{P}^{n+k}}^d$. Replacing \wp by one of its powers if necessary, we can as well assume that $d > n+k$. Then we have

$$H^q(\mathbb{C}\mathbb{P}^{n+k}, \Theta_{\mathbb{C}\mathbb{P}^{n+k}}^d) = \begin{cases} \{\psi \in \mathbb{C}[z_0, z_1, \dots, z_{n+k}] \mid \psi(\lambda z) = \lambda^d \psi(z)\}, & \text{if } q=0, \\ \{0\} & \text{for } q>0. \end{cases}$$

The fact that the pullback Θ_M^d of $\Theta_{\mathbb{C}\mathbb{P}^{n+k}}^d$ is a CR line bundle on M_t implies that also all pullbacks Θ_M^p of $\Theta_{\mathbb{C}\mathbb{P}^{n+k}}^p$ to M , for any $p \in \mathbb{Z}$, are CR line bundles on M_t . Indeed the fact that Θ_M^d is a CR line bundle on M_t means that we can find an open covering of M , that we can take of the form $\{U_{i,\mu}\}$ with $U_{i,\mu} \subseteq U_i = \{z_i \neq 0\}$ and d''_{M_t} -closed forms $\phi_{i,\mu} \in \mathcal{E}^{0,1}(U_{i,\mu})$ such that

$$\left(\frac{z_j}{z_i}\right)^d d''_{M_t} \left(\frac{z_i}{z_j}\right)^d = \phi_{i,\mu} - \phi_{j,\nu} \quad \text{on } U_{i,\mu} \cap U_{j,\nu}.$$

This is equivalent to

$$\left(\frac{z_j}{z_i}\right) d''_{M_t} \left(\frac{z_i}{z_j}\right) = \frac{1}{d} \phi_{i,\mu} - \frac{1}{d} \phi_{j,\nu} \quad \text{on } U_{i,\mu} \cap U_{j,\nu},$$

showing that Θ_M , and therefore all Θ_M^p with $p \in \mathbb{Z}$ are CR line bundles on M_t .

By repeating the proof in §6.1 by using the sections $\sigma_i = z_i$ of Θ_M , we obtain

Theorem 6.1. *Let M be a smooth compact generic CR submanifold of type (n, k) of $\mathbb{C}\mathbb{P}^{n+k}$ and D the zero locus in $\mathbb{C}\mathbb{P}^{n+k}$ of a homogeneous polynomial of degree d in $\mathbb{C}[z_0, z_1, \dots, z_{n+k}]$. Let $\{M_t\}$ be a family of CR structures of type (n, k) on M , smoothly depending on a real parameter t . If*

- (1) M_0 is induced by the embedding $M \hookrightarrow \mathbb{C}\mathbb{P}^{n+k}$;
- (2) M_0 is 2-pseudoconcave;
- (3) the CR structures of all M_t agree to infinite order on $M \cap D$.

Then we can find $\epsilon_0 > 0$ such that, for every $|t| < \epsilon_0$, M_t admits a generic CR embedding into $\mathbb{C}\mathbb{P}^{n+k}$. \square

6.3. Generic CR submanifolds of Fano varieties. We recall that a complex variety X is *Fano* if its anticanonical bundle K_X^{-1} is ample. If we add to the assumptions of Theorem 1.1 the requirement that X is Fano, then we can define a projective embedding of M_0 by using CR sections $\sigma_0, \sigma_1, \dots, \sigma_m$ of a power $K_M^{-\ell}$ of the pullback on M of the anticanonical bundle. Indeed, by [29], the tensor product $K_X^{-\ell} \otimes F_X^{-k}$ is the dual of a positive bundle and we can apply the arguments in §6.1 to obtain sections $\sigma_{i,t}$ in $\mathcal{O}_{M_t}(M_t, K_M^{-\ell})$, which, for $|t| \ll 1$ yield projective CR embeddings $\sigma_t = (\sigma_{0,t}, \sigma_{1,t}, \dots, \sigma_{m,t})$ of M_t into $\mathbb{C}\mathbb{P}^m$. The images $M'_t = \sigma_t(M_t)$ are generic submanifolds of complex $(n+k)$ -dimensional subvarieties X_t of $\mathbb{C}\mathbb{P}^m$ which have, by construction, an ample anticanonical bundle. Indeed we know from [23, Thm.5.2] that, since the M_t are pseudoconcave, global CR meromorphic functions on M_t are restrictions of global meromorphic functions on X_t . Thus we have

Theorem 6.2. *Add to the assumptions of Theorem 1.1 the fact that X is Fano. Then there is $\epsilon_0 > 0$ such that for all $|t| < \epsilon_0$ the abstract CR manifold M_t has a generic CR embedding into a complex Fano variety X_t .*

Acknowledgements. The first author was supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, grant BR 3363/2-2).

REFERENCES

1. A. ANDREOTTI, C.D. HILL: *Complex characteristic coordinates and tangential Cauchy-Riemann equations*. Ann. Scuola Norm. Sup. Pisa **26**, 299–324 (1972)
2. T. AKAHORI: *Deformation Theory of CR-Structures and Its Application to Deformations of Isolated Singularities I*, Advanced Studies in Pure Mathematics **25**, CR-Geometry and Overdetermined Systems, pp. 1-40 (1997).
3. A. ALTOMANI, C. MEDORI, AND M. NACINOVICH: *The CR structure of minimal orbits in complex flag manifolds*, J. Lie Theory **16**, no. 3, 483–530, (2006).
4. A. ANDREOTTI AND G.A. FREDRICKS: *Embeddability of real analytic Cauchy-Riemann manifolds*, Ann. Scuola Norm. Sup. Pisa (4) **6**, 285–304, (1979).
5. A. ANDREOTTI AND H. GRAUERT: *Theorèmes de Finitude pour la cohomologie des espaces complexes*, Bulletin de la Societ’e Mathématique de France, **90**, pp. 193–259 (1962).
6. H. AZAD, R. KOBAYASHI, M.N. QURESHI: *Quasi-potentials and Kähler-Einstein metrics on flag manifolds*, J. Algebra **196**, 620–629 (1997).
7. AU-YEUNG, YIK-HOI: *On Matrices Whose Nontrivial Real Linear Combinations Are Nonsingular*. Proc. of the A.M.S. **29**, no. 1, 17–22 (1971).
8. A. BOREL, F. HIRZEBRUCH: *Characteristic classes and homogeneous spaces*, Amer. J. Math **80**, 459–538 (1958).
9. J. BRINKSCHULTE, C.D. HILL: *Non locally trivializable CR line bundles over compact Lorentzian CR manifolds* Ann. Inst. Fourier **68**, 101–108 (2018).
10. J. BRINKSCHULTE, C.D. HILL: *Inflexible CR submanifolds*, Math. Z. **287**, 461–472 (2017).
11. J. BRINKSCHULTE, C.D. HILL: *Flexible and inflexible CR submanifolds*, Ark. Mat. **57**, 23–33 (2019).
12. J. BRINKSCHULTE, C.D. HILL, M. NACINOVICH: *Obstructions to generic embeddings*. Ann. Inst. Fourier **52**, 1785–1792 (2002).
13. J. BRINKSCHULTE, C.D. HILL, J. LEITERER, M. NACINOVICH: *Aspects of the Levi form*. to appear in B.U.M.I.
14. É. CARTAN: *Les systèmes différentiels extérieurs et leurs applications géométriques*, Actualités Scientifiques et Industrielles, n° 994, Hermann, Paris, 1945.
15. D. CATLIN, L. LEMPert: *A note on the instability of embeddings of Cauchy-Riemann manifolds*. J. Geom. Analysis **2**, pp. 99–104 (1992).
16. G. DARBOUX: *Sur le problème de Pfaff* Bull. Sci. Math. **6**, pp. 14-36, 49-68 (1882).
17. P. DE BARTOLOMEIS AND F. MEYLAN: *Intrinsic deformation theory of CR structures*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) **IX**, pp.459–494 (2010).
18. G.B. FOLLAND, J.J. KOHN: *The Neumann problem for the Cauchy-Riemann complex*. Ann. Math. Studies **75**, Princeton University Press, Princeton, N. J. (1972).
19. PH. A. GRIFFITHS, J. HARRIS: *Principles of Algebraic Geometry*, John Wiley & Sons, New York, 1978.
20. E. HEBBEY: *Sobolev spaces on Riemannian manifolds*, Lecture Notes in Mathematics, v.1635, Springer-Verlag, Berlin, (1996).

21. C.D. HILL, M. NACINOVICH: *Pseudoconcave CR manifolds*. Complex analysis and geometry (V. Ancona, E. Ballico, A. Silva, eds), Marcel Dekker, Inc., New York, 275–297 (1996).
22. C.D. HILL, M. NACINOVICH: *A weak pseudoconcavity condition for abstract almost CR manifolds*. Invent. math. **142**, 251–283 (2000).
23. C.D. HILL, M. NACINOVICH: *Fields of CR meromorphic functions*. Rend. Sem. Mat. Univ. Padova **111**, 179–204 (2004).
24. C.D. HILL, M. NACINOVICH: *Non completely solvable systems of complex first order PDE's*. Rend. Sem. Mat. Univ. Padova **129**, 129–169 (2013).
25. C.D. HILL, G. TAIANI: *Real analytic approximation of locally embeddable CR manifolds*. Compositio Math. **44**, 113–131 (1981).
26. L. HÖRMANDER: *An introduction to complex analysis in several complex variables*. North Holland Mathematical Library (1990).
27. D. HUSEMÖLLER: *Fibre Bundles*, Graduate Texts in Mathematics, Springer, New York, (1994).
28. T. KAJISA: *On the existence of local frames for CR vector bundles*. Hokkaido Math. J. **42**, 121–130 (2013).
29. S.L. KLEIMAN: *Towards a numerical theory of ampleness*. Ann. of Math. **84**, 293–344, (1966).
30. K. KODAIRA, D.C. SPENCER: *On deformations of complex analytic structures, I-II, III*. Ann. of Math. **67**, 328–466 (1958); **71**, 43–76 (1960).
31. L. LEMPERT: *Embeddings of three dimensional Cauchy-Riemann manifolds*. Math. Ann. **300**, 1–15 (1994).
32. C. MEDORI, M. NACINOVICH: *Pluriharmonic functions on abstract CR manifolds*. Ann. Mat. Pura e Appl. (IV) **CLXX**, 377–394 (1996).
33. L. NIRENBERG: *On a problem of Hans Lewy*. Uspeki Math. Naut. **292**, pp. 241–251 (1974).
34. A. NEWLANDER AND L. NIRENBERG: *Complex analytic coordinates in almost complex manifolds*. Annals of Mathematics. **66**, pp. 391–404 (1957).
35. P.L. POLYAKOV: *Versal embeddings of compact 3-pseudoconcave CR submanifolds*. Math. Z. **248**, 267–312 (2004).
36. H. ROSSI: *Attaching analytic spaces to an analytic space along a pseudoconcave boundary*, Proc. Conf. Complex Manifolds (Minneapolis), 1964, Springer-Verlag, New York, pp. 242–256 (1965).
37. J. A. Wolf, *The action of a real semisimple group on a complex flag manifold. I. Orbit structure and holomorphic arc components*, Bull. Amer. Math. Soc. **75** (1969), 1121–1237.

J. BRINKSCHULTE: MATHEMATISCHES INSTITUT, UNIVERSITÄT LEIPZIG, AUGUSTUSPLATZ 10/11, 04109 LEIPZIG (GERMANY)

E-mail address: brinkschulte@math.uni-leipzig.de

C.D. HILL: DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, N.Y. 11794 (USA)

E-mail address: dhill@math.stonybrook.edu

M. NACINOVICH: DIPARTIMENTO DI MATEMATICA, II UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA (ITALY)

E-mail address: nacinovi@mat.uniroma2.it