

THE CONE OF MINIMAL WEIGHTS FOR MOD p HILBERT MODULAR FORMS

FRED DIAMOND AND PAYMAN L KASSAEI

ABSTRACT. We prove that all mod p Hilbert modular forms arise via multiplication by generalized partial Hasse invariants from forms whose weight falls within a certain *minimal cone*. This answers a question posed by Andreatta and Goren, and generalizes our previous results which treated the case where p is unramified in the totally real field. Whereas our previous work made use of deep Jacquet–Langlands type results on the Goren–Oort stratification (not yet available when p is ramified), here we instead use properties of the stratification at Iwahori level which are more readily generalizable to other Shimura varieties.

1. INTRODUCTION

In this paper, we consider weights of mod p Hilbert modular forms associated to F , where F is a totally real field of degree $d = [F : \mathbb{Q}] > 1$. The weights of such forms are d -tuples of integers, and unlike the case of characteristic zero Hilbert modular forms or mod- p classical modular forms, there are non-zero forms of partially negative weight. Indeed the partial Hasse invariants studied by Andreatta and Goren in [1] are examples of such forms, and they ask whether all non-zero forms arise from ones of non-negative weight by multiplication by partial Hasse invariants. In our earlier work [6], we answered the question in the affirmative, under the assumption that p is unramified in F . Indeed we proved a stronger result, namely that all non-zero forms arise from ones whose weight lies in a certain minimal cone contained in the set of non-negative weights (and is strictly smaller unless p splits completely in F).

Here we obtain such a result for arbitrary totally real fields, now allowing p to be ramified in F . The overall strategy is similar to the one used in [6], namely to use \mathbb{P}^1 -bundle fibrations on strata to prove vanishing results, and deduce from these that multiplication by certain partial Hasse invariants defines an isomorphism for weights outside the minimal cone. There is however an essential difference which makes the method in this paper more amenable to generalization: rather than obtaining the desired fibrations from geometric Jacquet–Langlands relations between Shimura varieties for different reductive groups, we derive the fibrations from the Hecke action at primes over p .

We now briefly explain the statements of our main results, referring the reader to the body of the paper for more detailed definitions. We fix a sufficiently small level \mathfrak{n} prime to p , and let \overline{X}^{PR} denote the geometric special fibre of the Pappas–Rapoport model (defined in [15]) of the associated Hilbert modular variety. Then $\overline{X}_{\mathbb{F}_p}^{\text{PR}}$ is

2010 *Mathematics Subject Classification.* 11F41 (primary), 11F33, 14G35 (secondary).
F.D. was partially supported by EPSRC Grant EP/L025302/1.

smooth of dimension d over $\overline{\mathbb{F}}_p$, and comes equipped with automorphic line bundles ω_τ indexed by the embeddings $\tau \in \Sigma = \text{Hom}(F, \overline{\mathbb{Q}}_p)$. For $\mathbf{k} = \sum k_\tau \mathbf{e}_\tau \in \mathbb{Z}^\Sigma$, we consider the space of mod p Hilbert modular forms of weight \mathbf{k} , defined as

$$M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p) = H^0(\overline{X}_{\overline{\mathbb{F}}_p}^{\text{PR}}, \bigotimes_{\tau} \omega_{\tau}^{\otimes k_{\tau}}).$$

For each $\tau \in \Sigma$ there is a generalized partial Hasse invariant H_τ (defined by Reduzzi and Xiao in [16] if p is ramified in F) of weight $\mathbf{h}_\tau = n_\tau \mathbf{e}_{\sigma^{-1}\tau} - \mathbf{e}_\tau$, where the integer $n_\tau \in \{1, p\}$ and σ is a certain “shift” permutation on Σ (generalizing the Frobenius if p is unramified in F , in which case $n_\tau = p$ for all Σ).

Our key theorem then states:

Theorem. *If $n_\tau k_\tau < k_{\sigma^{-1}\tau}$ for some $\tau \in \Sigma$, then multiplication by H_τ induces an isomorphism:*

$$M_{\mathbf{k}-\mathbf{h}_\tau}(\mathfrak{n}; \overline{\mathbb{F}}_p) \xrightarrow{\sim} M_{\mathbf{k}}(\mathfrak{n}; \overline{\mathbb{F}}_p).$$

This generalizes the key result of [6] to the case where p is ramified in F . As mentioned above, the strategy for proving it is similar; however rather than applying deep results of [19]¹ (refining the geometric Jacquet–Langlands relation of [13]) we give an argument which uses more straightforward properties (proved in [8]) on the stratification of the special fibre of the Pappas–Rapoport model of the Hilbert modular variety of Iwahori level at a prime over p and its behaviour under the degeneracy maps to \overline{X}^{PR} . As in [6], the case where there are split primes over p requires a separate argument, for which we need to establish certain properties of the Goren–Oort stratification of the Pappas–Rapoport model; these results may be of independent interest.

The theorem allows us to answer the question posed by Andreatta and Goren in [1, 15.8] in the ramified case as well (or more precisely a version which also accounts for the generalized partial Hasse invariants of [16]). For a non-zero form $f \in M_{\mathbf{k}}(\mathfrak{n}; \overline{\mathbb{F}}_p)$, one can define its *filtration* $\Phi(f)$ to be the minimal weight of a form from which it arises by multiplication by partial Hasse invariants. Defining the *minimal cone* $C^{\min} \subset \mathbb{Q}_{\geq 0}^\Sigma$ to be the set of $\sum x_\tau \mathbf{e}_\tau$ such that $n_\tau x_\tau \geq x_{\sigma^{-1}\tau}$ for all τ ; we obtain the corollary:

Corollary. *If $0 \neq f \in M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p)$, then $\Phi(f) \in C^{\min}$.*

Besides the question of [1], motivation for the corollary comes from the problem of formulating Serre weight conjectures in the context of geometrically-defined Hilbert modular forms, as carried out in [7] when p is unramified in F . The corollary also immediately implies the vanishing of $M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p)$ for weights outside the cone spanned by the \mathbf{h}_τ (proved independently in [9] in the unramified case). Finally, it is also used in the proof of the main theorem of [5], proving the unramifiedness at p of Galois representations associated to (torsion) Hilbert modular forms of parallel weight one.

2. NOTATION

Let F be a totally real field of degree d over \mathbb{Q} , and let \mathcal{O}_F denote its ring of integers. Let p be a prime number, and \mathfrak{n} an ideal of \mathcal{O}_F relatively prime to p which

¹As yet unavailable when p is ramified in F .

is contained in $N\mathcal{O}_F$ for some $N > 3$. Let

$$\mathbb{S} := \{\mathfrak{p} \in \text{Spec } \mathcal{O}_F : \mathfrak{p}|p\}.$$

For any $\mathfrak{p} \in \mathbb{S}$, we let $e_{\mathfrak{p}}$ denote the ramification index, and $f_{\mathfrak{p}}$ denote the degree of the residue field $\mathbb{F}_{\mathfrak{p}} := \mathcal{O}_F/\mathfrak{p}$ over \mathbb{F}_p . We will be considering a finite field \mathbb{F} whose degree over \mathbb{F}_p is divisible by $f = \text{lcm}\{f_{\mathfrak{p}} : \mathfrak{p}|p\}$.

For every $\mathfrak{p} \in \mathbb{S}$, we let $F_{\mathfrak{p}}$ be the completion of F at \mathfrak{p} , $\mathcal{O}_{\mathfrak{p}}$ its ring of integers, $F_{\mathfrak{p}}^{ur}/\mathbb{Q}_p$ its maximal unramified subextension, and $\mathcal{O}_{\mathfrak{p}}^{ur}$ the ring of integers in $F_{\mathfrak{p}}^{ur}$. We identify, once and for all, $W(\mathbb{F}_{\mathfrak{p}})$ with the maximal unramified sub-extension of $F_{\mathfrak{p}}/\mathbb{Q}_p$, and fix a choice of a uniformizer $\varpi_{\mathfrak{p}} \in F_{\mathfrak{p}}$ for each $\mathfrak{p} \in \mathbb{S}$. Let

$$\mathbb{B} := \bigsqcup_{\mathfrak{p} \in \mathbb{S}} \mathbb{B}_{\mathfrak{p}},$$

where $\mathbb{B}_{\mathfrak{p}} := \text{Hom}(F_{\mathfrak{p}}^{ur}, \overline{\mathbb{Q}_p}) \cong \text{Hom}(W(\mathbb{F}_{\mathfrak{p}}), W(\mathbb{F})) \cong \text{Hom}(\mathbb{F}_{\mathfrak{p}}, \mathbb{F})$. Let

$$\Sigma := \bigsqcup_{\mathfrak{p} \in \mathbb{S}} \Sigma_{\mathfrak{p}},$$

where $\Sigma_{\mathfrak{p}} := \text{Hom}(F_{\mathfrak{p}}, \overline{\mathbb{Q}_p})$. Then Σ can be identified with $\text{Hom}(F, \overline{\mathbb{Q}_p})$ via restriction to F . There is an obvious restriction map $\text{res} : \Sigma_{\mathfrak{p}} \rightarrow \mathbb{B}_{\mathfrak{p}}$ which is $e_{\mathfrak{p}}$ -to-1. For $\beta \in \mathbb{B}_{\mathfrak{p}}$, we denote

$$\text{res}^{-1}(\beta) = \{\tau_{\beta,1}, \tau_{\beta,2}, \dots, \tau_{\beta,e_{\mathfrak{p}}}\}.$$

There is no canonical way to order these embeddings, and we choose an ordering as above once and for all. In short, every embedding $\tau : F \rightarrow \overline{\mathbb{Q}_p}$ is the restriction to F of a unique embedding of the form $\tau_{\beta,i}$ for some $\beta \in \mathbb{B}$, and some $1 \leq i \leq e_{\mathfrak{p}}$ (where \mathfrak{p} is such that $\beta \in \mathbb{B}_{\mathfrak{p}}$). We often abuse notation and identify τ with $\tau_{\beta,i}$. Let ϕ denote the automorphism of $W(\mathbb{F})$ that lifts the Frobenius on \mathbb{F} . We then define $\sigma : \Sigma \rightarrow \Sigma$ by $\sigma\tau_{\beta,i} = \tau_{\beta,i+1}$ if $i < e_{\mathfrak{p}}$, and $\sigma\tau_{\beta,e_{\mathfrak{p}}} = \tau_{\phi\beta,1}$.

Let $K \subset \overline{\mathbb{Q}_p}$ be a finite extension of \mathbb{Q}_p that contains all the images $\tau_{\beta,i}(F_{\mathfrak{p}})$ for all $\mathfrak{p} \in \mathbb{S}$, $\beta \in \mathbb{B}_{\mathfrak{p}}$, and $1 \leq i \leq e_{\mathfrak{p}}$. Let \mathcal{O} denote the ring of integers of K (fixing an identification of $W(\mathbb{F})$ with a subring of \mathcal{O}). We fix a uniformizer ϖ of \mathcal{O} . Note that we can take \mathbb{F} to be \mathcal{O}/ϖ .

Let R be an \mathcal{O} -algebra, and Λ be a $(\mathcal{O}_F \otimes_{\mathbb{Z}} R)$ -module. The decomposition

$$\mathcal{O}_F \otimes_{\mathbb{Z}} \mathcal{O} = \prod_{\mathfrak{p}|\mathfrak{p}} \mathcal{O}_{\mathfrak{p}} \otimes_{\mathbb{Z}_p} \mathcal{O} = \prod_{\beta \in \mathbb{B}} \mathcal{O}_{\mathfrak{p}} \otimes_{W(\mathbb{F}_{\mathfrak{p}}), \beta} \mathcal{O}$$

induces a decomposition

$$(2.1) \quad \Lambda = \bigoplus_{\mathfrak{p} \in \mathbb{S}} \Lambda_{\mathfrak{p}} = \bigoplus_{\beta \in \mathbb{B}} \Lambda_{\beta},$$

of Λ into $(\mathcal{O}_F \otimes_{\mathbb{Z}} R)$ -modules, where the action of \mathcal{O}_F on $\Lambda_{\mathfrak{p}}$ extends to a continuous action of $\mathcal{O}_{\mathfrak{p}}$, and for $\beta \in \mathbb{B}_{\mathfrak{p}}$, $\Lambda_{\beta} \subset \Lambda_{\mathfrak{p}}$ is the submodule where the action of $W(\mathbb{F}_{\mathfrak{p}}) \subset \mathcal{O}_{\mathfrak{p}}$ is given by scalar multiplication via $\beta : W(\mathbb{F}_{\mathfrak{p}}) \rightarrow \mathcal{O} \rightarrow R$.

3. HILBERT MODULAR VARIETIES: THE PAPPAS-RAPOPORT MODEL

In this section, we consider the Pappas-Rapoport integral model of the Shimura variety corresponding to the reductive group over \mathbb{Q} defined by

$$G(R) = \{g \in GL_2(F \otimes_{\mathbb{Q}} R) \mid \det(g) \in R^*\},$$

for a \mathbb{Q} -algebra R , where the level is hyperspecial at p . When p is unramified in F , this model is the same as the Deligne-Pappas model. When p is ramified in F , the Pappas-Rapoport model is a desingularization of the Deligne-Pappas model.

Let \mathfrak{J} be a fractional ideal of F , and \mathfrak{J}^+ its cone of totally positive elements. Let \mathfrak{n} be an ideal of \mathcal{O}_F prime to p which contains $N\mathcal{O}_F$ for an integer $N > 3$. Let $X_{\mathfrak{J}}^{\text{DP}}/\mathcal{O}$ be the Hilbert modular scheme classifying \mathfrak{J} -polarized Hilbert-Blumenthal abelian schemes (\mathfrak{J} -polarized HBAS's)

$$\underline{A}/S = (A/S, \iota, \lambda, \alpha),$$

where

- S is a locally noetherian \mathcal{O} -scheme;
- A is an abelian scheme of relative dimension d over S , equipped with real multiplication $\iota: \mathcal{O}_F \rightarrow \text{End}_S(A)$;
- λ is a polarization as in [4], namely, an isomorphism $\lambda: (\mathcal{P}_A, \mathcal{P}_A^+) \rightarrow (\mathfrak{J}, \mathfrak{J}^+)$ such that $A \otimes_{\mathcal{O}_F} \mathfrak{J} \cong A^\vee$. Here, $\mathcal{P}_A = \text{Hom}_{\mathcal{O}_F}(A, A^\vee)^{\text{sym}}$, viewed as sheaf of projective \mathcal{O}_F -modules of rank one in the étale topology, and \mathcal{P}_A^+ is the cone of polarizations;
- the map $\alpha: (\mathcal{O}_F/\mathfrak{n})^2 \rightarrow A[\mathfrak{n}]$ is an \mathcal{O}_F -linear isomorphism of group schemes.

We fix $\{\mathfrak{J}_1, \dots, \mathfrak{J}_r\}$ a set of representatives for the strict class group $\text{Cl}^+(F)$. We define

$$X^{\text{DP}} = \bigcup_{i=1}^r X_{\mathfrak{J}_i}^{\text{DP}}.$$

The Hilbert modular scheme X^{DP} is a normal scheme of relative dimension d over $\text{Spec } \mathcal{O}$. Let $\overline{X}^{\text{DP}} = X^{\text{DP}} \otimes_{\mathcal{O}} \mathbb{F}$ be the special fibre, viewed as a closed subscheme of X^{DP} .

Let S be a locally noetherian \mathcal{O} -scheme of connected components $\{S_i\}_{i \in I}$. By a HBAS over S we mean a function $\nu: I \rightarrow \{1, \dots, r\}$, and a collection $\{\underline{A}_i\}_{i \in I}$, where \underline{A}_i is a $\mathfrak{J}_{\nu(i)}$ -polarized HBAS over S_i for each $i \in I$.

Let \underline{A}/S be a HBAS as above, and let $\epsilon_A: A \rightarrow S$ be the structure map. Let $\omega_{A/S} = \epsilon_{A,*} \Omega_{A/S}^1$ and $\mathbb{H}_{A/S} = R^1 \epsilon_{A,*} \Omega_{A/S}^\bullet$; these are locally free sheaves of ranks, respectively, $d, 2d$ on S . These sheaves carry an action of $\mathcal{O}_F \otimes \mathcal{O}_S$, and since S is an \mathcal{O} -scheme, we obtain decompositions as in Equation 2.1 given by

$$\omega_{A/S} = \bigoplus_{\beta \in \mathbb{B}} \omega_{A/S, \beta},$$

$$\mathbb{H}_{A/S} = \bigoplus_{\beta \in \mathbb{B}} \mathbb{H}_{A/S, \beta}.$$

For $\beta \in \mathbb{B}_{\mathfrak{p}}$ the sheaves $\omega_{A/S, \beta} \subset \mathbb{H}_{A/S, \beta}$ are locally free sheaves of rank $e_{\mathfrak{p}}$ and $2e_{\mathfrak{p}}$, respectively. We denote the corresponding sheaves on \overline{X}^{DP} obtained from the universal HBAS by $\omega = \bigoplus_{\beta \in \mathbb{B}} \omega_{\beta}$ and $\mathbb{H} = \bigoplus_{\beta \in \mathbb{B}} \mathbb{H}_{\beta}$.

By a Pappas-Rapoport filtration on $\omega_{A/S}$, we mean a collection of filtrations,

$$\omega_{A/S}^\bullet = (\omega_{A/S, \beta}^\bullet)_{\beta \in \mathbb{B}},$$

where, for each $\mathfrak{p} \in \mathbb{S}$ and $\beta \in \mathbb{B}_{\mathfrak{p}}$, $\omega_{A/S, \beta}^\bullet$ is a filtration of $\omega_{A/S, \beta}$ given by

$$0 = \omega_{A/S, \beta}^{(0)} \subset \dots \subset \omega_{A/S, \beta}^{(i)} \subset \dots \subset \omega_{A/S, \beta}^{(e_{\mathfrak{p}})} = \omega_{A/S, \beta},$$

satisfying the following conditions:

- (1) each $\omega_{A/S, \beta}^{(i)}$ is \mathcal{O}_F -stable;
- (2) the graded piece $\omega_{A/S, \beta}^{(i)} / \omega_{A/S, \beta}^{(i-1)}$ is a locally free sheaf of rank one over S (in particular, each $\omega_{A/S, \beta}^{(i)}$ is a locally free sheaf of rank i over S);

(3) the action of \mathcal{O}_F on $\omega_{A/S,\beta}^{(i)}/\omega_{A/S,\beta}^{(i-1)}$ is the scalar action via $\mathcal{O}_F \rightarrow \mathcal{O}_{\mathfrak{p}} \xrightarrow{\tau_{\beta,i}} \mathcal{O}$.

In other words, the action of $\varpi_{\mathfrak{p}}$ on $\omega_{A/S,\beta}^{(i)}/\omega_{A/S,\beta}^{(i-1)}$ is given by the scalar action of $\tau_{\beta,i}(\varpi_{\mathfrak{p}}) \in \mathcal{O}$.

There is a scheme X^{PR} over $\text{Spec } \mathcal{O}$ classifying *filtered HBAS's*,

$$(A/S, \iota, \lambda, \alpha, \omega_{A/S}^{\bullet}) = (\underline{A}/S, \omega_{A/S}^{\bullet})$$

where,

- S is a locally noetherian \mathcal{O} -scheme;
- \underline{A}/S is a HBAS of relative dimension d as above;
- $\omega_{A/S}^{\bullet}$ is a Pappas-Rapoport filtration on $\omega_{A/S}$.

The Hilbert modular scheme X^{PR} is a scheme smooth of relative dimension d over $\text{Spec } \mathcal{O}$ (see [15],[17]). There is an obvious forgetful morphism

$$\delta : X^{\text{PR}} \rightarrow X^{\text{DP}}.$$

It is easy to see that δ is an isomorphism when restricted to the Rapoport loci of the source and the target, i.e., the loci where the HBAS in question, \underline{A}/S , satisfies the Rapoport condition: $\omega_{A/S}$ is a locally free $(\mathcal{O}_S \otimes_{\mathbb{Z}} \mathcal{O}_F)$ -module of rank one.

We denote by $\overline{X}^{\text{PR}} = X^{\text{PR}} \otimes_{\mathcal{O}} \mathbb{F}$ the special fibre of X^{PR} . Note that a Pappas-Rapoport filtration over an HBAS \underline{A} over an \mathbb{F} -scheme S is a collection of \mathcal{O}_F -stable filtrations indexed by $\mathfrak{p} \in \mathbb{S}$, $\beta \in \mathbb{B}_{\mathfrak{p}}$:

$$0 = \omega_{A/S,\beta}^{(0)} \subset \dots \subset \omega_{A/S,\beta}^{(i)} \subset \dots \subset \omega_{A/S,\beta}^{(e_{\mathfrak{p}})} = \omega_{A/S,\beta},$$

where the graded pieces are locally free of rank one, and such that we have

$$[\varpi_{\mathfrak{p}}](\omega_{A/S,\beta}^{(i)}) \subset \omega_{A/S,\beta}^{(i-1)},$$

for all $1 \leq i \leq e_{\mathfrak{p}}$. We continue to denote by $\omega, \mathbb{H}, \omega_{\beta}, \mathbb{H}_{\beta}$ the pullback of these sheaves from \overline{X}^{DP} to \overline{X}^{PR} . For any (β, i) as above, we define $\omega_{\beta}^{(i)}$ to be $\omega_{A/S,\beta}^{(i)}$ where A/S is taken to be the universal filtered HBAS over \overline{X}^{PR} . This is a locally free sheaf of rank i on \overline{X}^{PR} . Let $\tau \in \Sigma = \text{Hom}(F, \overline{\mathbb{Q}}_p)$. There are unique $\mathfrak{p} \in \mathbb{S}$ and $\beta \in \mathbb{B}_{\mathfrak{p}}$ such that τ is the restriction to F of $\tau_{\beta,i}$ for some $1 \leq i \leq e_{\mathfrak{p}}$. We denote by $\omega_{\tau} = \omega_{\beta,i}$ the quotient sheaf $\omega_{\beta}^{(i)}/\omega_{\beta}^{(i-1)}$. Each ω_{τ} is a locally free sheaf of rank one over \overline{X}^{PR} .

Let $\mathfrak{p} \in \mathbb{S}$ and $\beta \in \mathbb{B}_{\mathfrak{p}}$. Consider $[\varpi_{\mathfrak{p}}] : \mathbb{H}_{\beta} \rightarrow \mathbb{H}_{\beta}$. Define, for $1 \leq i \leq e_{\mathfrak{p}}$

$$(3.1) \quad \mathbb{H}_{\beta,i} = ([\varpi_{\mathfrak{p}}]^{-1} \omega_{\beta}^{(i-1)}) / \omega_{\beta}^{(i-1)}.$$

Clearly, we have $\omega_{\beta,i} \subset \mathbb{H}_{\beta,i}$. If $\tau = \tau_{\beta,i} \in \Sigma$, we set $\mathbb{H}_{\tau} = \mathbb{H}_{\beta,i}$.

Lemma 3.1. *The sheaf $\mathbb{H}_{\beta,i}$ is locally free of rank 2 on \overline{X}^{PR} . Furthermore, we have a non-canonical isomorphism $\wedge^2 \mathbb{H}_{\beta,i} \cong \mathcal{O}_{\overline{X}^{\text{PR}}}$.*

Proof. First we show that at every closed point, the fibre of $\mathbb{H}_{\beta,i}$ has dimension 2. Since \mathbb{H} is locally free of rank 2 over $\mathcal{O}_{\overline{X}^{\text{PR}}} \otimes_{\mathbb{Z}} \mathcal{O}_F$, we know for all $\beta \in \mathbb{B}_{\mathfrak{p}}$, \mathbb{H}_{β} is locally free of rank 2 over $\mathcal{O}_{\overline{X}^{\text{PR}}} \otimes_{W(\mathbb{F}_{\mathfrak{p}}), \beta} \mathcal{O}_{\mathfrak{p}} \cong \mathcal{O}_{\overline{X}^{\text{PR}}}[[u]]/\langle u^{e_{\mathfrak{p}}} \rangle$, where the isomorphism is chosen so that $\varpi_{\mathfrak{p}}$ is sent to u . Let x be a closed point of \overline{X}^{PR} with residue field κ . Hence $(\mathbb{H}_{\beta})_x \cong (\kappa[[u]]/\langle u^{e_{\mathfrak{p}}} \rangle)^{\oplus 2}$, where a basis is chosen such

that the submodule $\omega_\beta^{(i-1)}$ is given by $\frac{u^\alpha \kappa[u]}{\langle u^{e_p} \rangle} \oplus \frac{u^\beta \kappa[u]}{\langle u^{e_p} \rangle}$ with α, β positive integers. It follows that

$$(\mathbb{H}_{\beta,i})_x = [\varpi_p]^{-1}(\omega_\beta^{(i-1)})_x / (\omega_\beta^{(i-1)})_x \cong \frac{u^{\alpha-1} \kappa[u] / \langle u^{e_p} \rangle}{u^\alpha \kappa[u] / \langle u^{e_p} \rangle} \oplus \frac{u^{\beta-1} \kappa[u] / \langle u^{e_p} \rangle}{u^\beta \kappa[u] / \langle u^{e_p} \rangle}$$

which has dimension 2.

In [16, Cor 2.10] a sheaf \mathcal{H} (denoted there $\mathcal{H}_{p,j}^{(l)}$ for appropriate choices of j, l) is considered that is contained in $\mathbb{H}_{\beta,i}$ by definition. It is proven that this sheaf is locally free of rank 2, and that its determinant is a trivial line bundle. So it is enough to show \mathcal{H} equals $\mathbb{H}_{\beta,i}$. But the quotient sheaf $\mathbb{H}_{\beta,i}/\mathcal{H}$ has all fibres zero, and hence it equals zero. \square

If $i > 1$, the action of ϖ_p induces a map

$$[\varpi_p] : \mathbb{H}_{\beta,i} \rightarrow \mathbb{H}_{\beta,i-1}$$

which clearly has image $\omega_{\beta,i-1}$.

Definition 3.2. Let R be an \mathbb{F} -algebra. Let $\mathbf{k} = \sum k_\tau \mathbf{e}_\tau \in \mathbb{Z}^\Sigma$. We define

$$\omega^{\mathbf{k}} := \bigotimes_{\tau} \omega_{\tau}^{\otimes k_{\tau}}$$

The space of mod p Hilbert modular forms of weight \mathbf{k} and level $\Gamma(\mathbf{n})$ over R is defined to be

$$M_{\mathbf{k}}(\mathbf{n}; R) = H^0(\overline{X}^{PR} \otimes_{\mathbb{F}} R, \omega^{\mathbf{k}}).$$

4. THE GENERALIZED PARTIAL HASSE INVARIANTS

We recall the definition of generalized partial Hasse invariants introduced in [16] which factorize the partial Hasse invariants defined by [1]. Let $\mathfrak{p} \in \mathbb{S}$ and $\beta \in \mathbb{B}_{\mathfrak{p}}$. The Verschiebung morphism of the universal filtered HBAS over the Hilbert modular variety \overline{X}^{PR}

$$\text{Ver} : A^{(\phi)} \rightarrow A$$

induces $\text{Ver}_{\beta}^* : \mathbb{H}_{\beta} \rightarrow \omega_{\phi^{-1}\beta}^{(\phi)} \subset \mathbb{H}_{\phi^{-1}\beta}^{(\phi)}$ (where $\cdot^{(\phi)}$ denotes pullback by absolute Frobenius on \overline{X}^{PR}). The restricted map $\text{Ver}_{\beta}^* : \omega_{\beta} \rightarrow \omega_{\phi^{-1}\beta}^{(\phi)}$ preserves the universal Pappas-Rapoport filtration and hence induces

$$\text{Ver}_{\beta}^* : \omega_{\beta, e_p} = \omega_{\beta} / \omega_{\beta}^{(e_p-1)} \rightarrow \omega_{\phi^{-1}\beta, e_p}^{\otimes p} = \omega_{\phi^{-1}\beta}^{(\phi)} / (\omega_{\phi^{-1}\beta}^{(e_p-1)})^{(\phi)}.$$

We let $H_{\beta} \in H^0(\overline{X}^{PR}, \omega_{\phi^{-1}\beta, e_p}^{\otimes p} \otimes \omega_{\beta, e_p}^{-1})$ denote the section defined by the above morphism. It is easy to see that on the Rapoport locus of \overline{X}^{PR} , H_{β} corresponds under δ to a partial Hasse invariant (at the embedding β) defined by [1] on the Rapoport locus of \overline{X}^{DP} . In [16], Ver_{β}^* is factorized as follows. For $i > 1$, let

$$m_{\beta,i} : \omega_{\beta,i} = \omega_{\beta}^{(i)} / \omega_{\beta}^{(i-1)} \rightarrow \omega_{\beta,i-1} = \omega_{\beta}^{(i-1)} / \omega_{\beta}^{(i-2)}$$

be the morphism defined by the action of $\varpi_p \in \mathcal{O}_{\mathfrak{p}}$. Define a morphism

$$V_{\beta,1} : \mathbb{H}_{\beta,1} \rightarrow \omega_{\phi^{-1}\beta, e_p}^{\otimes p}$$

as follows: Let h be a local section of $\mathbb{H}_{\beta,1}$. Since $[\varpi_{\mathfrak{p}}]h = 0$, it follows that $z = [\varpi_{\mathfrak{p}}]^{e_{\mathfrak{p}}-1}h'$ for h' some local section of \mathbb{H}_{β} . Define $V_{\beta,1}(h)$ to be the image of $\text{Ver}_{\beta}^*(h') \in \omega_{\phi^{-1}\beta}^{(\phi)}$ in $\omega_{\phi^{-1}\beta, e_{\mathfrak{p}}}^{\otimes p}$. It follows that we have the factorization

$$\text{Ver}_{\beta}^* = m_{\beta, e_{\mathfrak{p}}} \circ \cdots \circ m_{\beta, 2} \circ V_{\beta, 1}.$$

Let $H_{\beta,1} \in H^0(\overline{X}^{\text{PR}}, \omega_{\phi^{-1}\beta, e_{\mathfrak{p}}}^{\otimes p} \otimes \omega_{\beta,1}^{-1})$ be the section defined by the restriction of $V_{\beta,1}$ to $\omega_{\beta,1}$. Also, for $1 < i \leq e_{\mathfrak{p}}$, let $H_{\beta,i} \in H^0(\overline{X}^{\text{PR}}, \omega_{\beta,i}^{-1} \otimes \omega_{\beta,i-1})$ be the section defined by $m_{\beta,i}$. It follows that the pullback under δ of the partial Hasse invariant (at β) defined in [1] on the Rapoport locus of \overline{X}^{DP} extends from the Rapoport locus of \overline{X}^{PR} to \overline{X}^{PR} , and factorizes as $H_{\beta} = \prod_{i=1}^{e_{\mathfrak{p}}} H_{\beta,i}$.

Let $\tau \in \Sigma = \text{Hom}(F, \overline{\mathbb{Q}}_p)$, so that $\tau = \tau_{\beta,i}$ for a unique $\mathfrak{p} \in \mathbb{S}$, $\beta \in \mathbb{B}_{\mathfrak{p}}$, and $1 \leq i \leq e_{\mathfrak{p}}$. We define $H_{\tau} = H_{\beta,i}$. Then H_{τ} is a mod p Hilbert modular form of weight \mathbf{h}_{τ} which equals $\mathbf{e}_{\sigma^{-1}\tau} - \mathbf{e}_{\tau}$ if $i > 1$, and $p\mathbf{e}_{\sigma^{-1}\tau} - \mathbf{e}_{\tau}$ if $i = 1$.

5. THE GOREN-OORT STRATIFICATION ON \overline{X}^{PR}

The partial Hasse invariants H_{τ} can be used to define an analogue of the Goren-Oort stratification on \overline{X}^{PR} (see [16]). For any $\tau \in \Sigma = \text{Hom}(F, \overline{\mathbb{Q}}_p)$, define Z_{τ} to be the closed subscheme of \overline{X}^{PR} given by the vanishing of H_{τ} . For $T \subset \Sigma$, define a closed subscheme of \overline{X}^{PR}

$$Z_T = \bigcap_{\tau \in T} Z_{\tau},$$

as well as an open subscheme of Z_T given by

$$W_T = Z_T - \bigcup_{\tau \notin T} Z_{\tau}.$$

In [16, Thm 3.10] it is shown that, for $\tau \in \Sigma$, the closed subschemes Z_{τ} are smooth proper divisors on \overline{X}^{PR} intersecting transversally. In particular, for $T \neq \emptyset$, Z_T is proper. In the following, we prove some properties of this stratification that we use in the proof of our main theorem.

Lemma 5.1. *Let $\mathfrak{p} \in \mathbb{S}$, $\beta \in \mathbb{B}_{\mathfrak{p}}$, and $1 \leq i \leq e_{\mathfrak{p}}$. Let $T \subset \Sigma$ and $\tau_{\beta,i} \in T$. Then, if $i > 1$, we have*

$$\omega_{\beta,i}|_{Z_T} \cong \omega_{\beta,i-1}^{-1}|_{Z_T}.$$

For $i = 1$, we have

$$\omega_{\beta,1}|_{Z_T} \cong \omega_{\phi^{-1}\beta, e_{\mathfrak{p}}}^{-p}|_{Z_T}.$$

Let $U = \overline{X}^{\text{PR}} - Z_{\tau_{\beta,i}}$. Then if $i > 1$, we have

$$\omega_{\beta,i}|_U \cong \omega_{\beta,i-1}|_U.$$

For $i = 1$, we have

$$\omega_{\beta,1}|_U \cong \omega_{\phi^{-1}\beta, e_{\mathfrak{p}}}^p|_U.$$

Proof. Assume $i > 1$ and $\tau_{\beta,i} \in T$. Consider the map $[\varpi_{\mathfrak{p}}] : \mathbb{H}_{\beta,i} \rightarrow \mathbb{H}_{\beta,i-1}$ defined in §3. It has image $\omega_{\beta,i}$. By assumption, $H_{\beta,i} = 0$ on Z_T , and hence the map $m_{\beta,i} : \omega_{\beta,i} \rightarrow \omega_{\beta,i-1}$ vanishes on Z_T . By definition, this implies that

$[\varpi_{\mathfrak{p}}]\omega_{\beta}^{(i)} \subset \omega_{\beta}^{(i-2)}$. In particular, $\omega_{\beta,i}$ sits in the kernel of the map $[\varpi_{\mathfrak{p}}]$ above. Rank considerations show that we have a short exact sequence of sheaves on Z_T :

$$0 \rightarrow \omega_{\beta,i} \rightarrow \mathbb{H}_{\beta,i} \rightarrow \omega_{\beta,i-1} \rightarrow 0,$$

from which it follows $\mathcal{O}_{Z_T} \cong \wedge_{\mathcal{O}_{Z_T}}^2 \mathbb{H}_{\beta,i} \cong \omega_{\beta,i}|_{Z_T} \otimes \omega_{\beta,i-1}|_{Z_T}$.

Now assume $\tau_{\beta,1} \in T$. Consider the morphism $V_{\beta,1} : \mathbb{H}_{\beta,1} \rightarrow \omega_{\phi^{-1}\beta,e_{\mathfrak{p}}}^{\otimes p}$ defined in §4. Since by assumption $H_{\beta,1} = 0$ on Z_T , we have $V_{\beta,1}(\omega_{\beta,1}) = 0$. Rank considerations imply we have an exact sequence

$$0 \rightarrow \omega_{\beta,1} \rightarrow \mathbb{H}_{\beta,1} \rightarrow \omega_{\phi^{-1}\beta,e_{\mathfrak{p}}}^{\otimes p} \rightarrow 0.$$

It follows that $\mathcal{O}_{Z_T} \cong \wedge_{\mathcal{O}_{Z_T}}^2 \mathbb{H}_{\beta,1} \cong \omega_{\beta,1}|_{Z_T} \otimes \omega_{\phi^{-1}\beta,e_{\mathfrak{p}}}^{\otimes p}|_{Z_T}$.

The other two isomorphisms follow immediately from the definition of Hasse invariants. □

Corollary 5.2. *Let $T \subset \Sigma$. For any $\tau \in \Sigma$, we have*

$$\omega_{\tau}^{\otimes(p^{2f_{\mathfrak{p}}}-1)}|_{W_T} \cong \mathcal{O}_{W_T}.$$

Corollary 5.3. *Let $\mathfrak{p} \in \mathbb{S}$. If $\Sigma_{\mathfrak{p}} \subset T \subset \Sigma$, then for all $\tau \in \Sigma_{\mathfrak{p}}$ we have*

$$\omega_{\tau}^{\otimes(p^{2f_{\mathfrak{p}}}-1)}|_{Z_T} \cong \mathcal{O}_{Z_T}.$$

Lemma 5.4. *There are integers m_{τ} for $\tau \in \Sigma$ such that $\bigotimes_{\tau} \omega_{\tau}^{\otimes m_{\tau}}$ is an ample line bundle on \overline{X}^{PR} .*

Proof. Let $\delta : \overline{X}^{\text{PR}} \rightarrow \overline{X}^{\text{DP}}$ be the forgetful morphism. Let $\omega \subset \mathbb{H}$ be the Hodge bundle on \overline{X}^{DP} . Then the determinant $\wedge^d \omega$ is an ample line bundle on \overline{X}^{DP} (see, for example, [3, Theorem 4.3.(ix)]) and we have

$$\delta^*(\wedge^d \omega) = \bigotimes_{\mathfrak{p} \in \mathbb{S}} \bigotimes_{\beta \in \mathbb{B}_{\mathfrak{p}}} \bigotimes_{i=1}^{e_{\mathfrak{p}}} \omega_{\beta,i}.$$

It is easy to see that $\mathcal{M} = \bigotimes_{\mathfrak{p} \in \mathbb{S}} \bigotimes_{\beta \in \mathbb{B}_{\mathfrak{p}}} \bigotimes_{i=1}^{e_{\mathfrak{p}}} \omega_{\beta,i}^{\otimes(i-e_{\mathfrak{p}}-1)}$ is relatively ample with respect to δ (See [16, Lemma 2.7]). By [18, Lem.29.35.7], it follows that for a large enough, $\mathcal{M} \otimes \delta^*(\wedge^d \omega)^{\otimes a} = \bigotimes_{\mathfrak{p} \in \mathbb{S}} \bigotimes_{\beta \in \mathbb{B}_{\mathfrak{p}}} \bigotimes_{i=1}^{e_{\mathfrak{p}}} \omega_{\beta,i}^{\otimes(a+i-e_{\mathfrak{p}}-1)}$ is ample on \overline{X}^{PR} . □

We can now prove the main result of this section.

Proposition 5.5. *Let $T \subset \Sigma$. Then W_T is a quasi-affine scheme.*

Proof. Since W_T is quasi-compact, it is enough to show that there is a line bundle on W_T that is both ample and torsion. Let m_{τ} be integers as in Lemma 5.4. Then $\mathcal{L} = \bigotimes_{\tau} \omega_{\tau}^{\otimes m_{\tau}}$ is ample on W_T . On the other hand, by Corollary 5.2, each line bundle ω_{τ} , and hence \mathcal{L} , is a torsion line bundle on W_T . □

Remark 5.6. In various works, Oort proves quasi-affineness of strata on Shimura varieties via the existence of a torsion ample line bundle on them. In [11] this approach is used to prove that open Goren-Oort strata on a Hilbert modular variety are quasi-affine in the case p is unramified in F ; to construct a torsion ample line bundle on an open stratum, the authors use the relationship between the Ekedahl-Oort stratification and the type stratification. In [2] it is shown that the usual definition of the Ekedahl-Oort stratification is inadequate by giving an example

where there are infinitely many isomorphism classes for the p -torsion over \overline{X}^{DP} . The proof of Proposition 5.5 works directly over the generalized Goren-Oort strata, giving a new proof in the case p is unramified.

Corollary 5.7. *Let $T \subset \Sigma$. Let C be an irreducible component of Z_T . Then for all $\tau \notin T$, we have $C \cap Z_\tau \neq \emptyset$.*

Proof. If not, let $T \subset \Sigma$ be of maximal cardinality where Z_T has a component C such that $C \cap Z_{\tau_0} = \emptyset$ for some $\tau_0 \notin T$ (in particular, $|T| < d$). It follows that $C \cap Z_\tau = \emptyset$ for all $\tau \notin T$. Hence $C \subset W_T$, and being quasi-compact, it follows that C is quasi-affine. We cannot have $T = \emptyset$, since otherwise the assumptions would imply that $Z_\tau = \emptyset$ for all $\tau \in \Sigma$ which is false (see, for example, [1, Cor 8.18]). Since $T \neq \emptyset$, C is proper, and, being quasi-affine, it must be finite. On the other hand, since Z_T is the intersection of $|T|$ divisors, its irreducible components have dimension at least $d - |T| > 0$. This is a contradiction. \square

To close this section we summarize some properties of the stratification on \overline{X}^{PR} .

Proposition 5.8. *The following statements hold.*

- (1) *The collection $\{W_T\}_T$ is a stratification of \overline{X}^{PR} , i.e., for every subset $T \subset \Sigma$, we have $Z_T = \overline{W}_T = \bigcup_{T' \supseteq T} W_{T'}$.*
- (2) *Each stratum Z_T is non-empty and equidimensional of dimension $d - |T|$, and proper if $T \neq \emptyset$.*
- (3) *Each open stratum W_T is non-empty, quasi-affine, and equidimensional of dimension $d - |T|$.*

Proof. We first prove by induction that each Z_T is nonempty and equidimensional of dimension $d - |T|$. This is true for $T = \emptyset$. Assume the statement is true for Z_T , and let $T' = T \cup \{\tau_0\}$ where $\tau_0 \notin T$. Every component C' of $Z_{T'}$ is a component of $C \cap Z_{\tau_0}$ for some component C of Z_T . By Corollary 5.7, $C \cap Z_{\tau_0} \neq \emptyset$, which proves that $Z_{T'} \neq \emptyset$. Furthermore, it follows that for every component $C' \subset Z_{T'}$, we have $\dim(C') \leq d - |T|$. On the other hand, we have $\dim(C') \geq d - |T'|$ (as in the proof of Corollary 5.7), so if equality fails for a component C' , then we have $\dim(C') = d - |T|$. Hence $C' \cap \bigcap_{\tau \notin T'} Z_\tau$, which is contained in Z_Σ and non-empty by Corollary 5.7, will have dimension at least $d - |T| - (d - |T'|) = 1$. On the other hand, $Z_\Sigma = W_\Sigma$ is zero-dimensional as it is both proper and quasi-affine (by Proposition 5.5). This ends the proof by induction. To show that W_T is nonempty for all T , note that $W_T = Z_T - \bigcup_{T' \supsetneq T} Z_{T'}$ and that $\dim(Z_{T'}) < \dim(Z_T)$ if $T' \supsetneq T$. This also shows that W_T is equi-dimensional of dimension $\dim(Z_T) = d - |T|$.

We now prove that $\{W_T\}_T$ form a stratification of \overline{X}^{PR} . Since Z_T is closed, we have $\overline{W}_T \subset Z_T$. By dimension considerations, it follows that \overline{W}_T is a union of components of Z_T . If $\overline{W}_T \neq Z_T$, then a non-empty union of components of Z_T will have to lie inside $Z_T - W_T = \bigcup_{T' \supsetneq T} Z_{T'}$, which is impossible since every $Z_{T'}$ appearing here has dimension smaller than that of $Z_T - W_T$. The quasi-affineness of W_T is Proposition 5.5. \square

6. FILTRATION OF MOD p HILBERT MODULAR FORMS.

In [1, §8.19] a notion of filtration is defined for a nonzero mod p Hilbert modular form f as the minimal weight of a modular form from which f arises by multiplication by the partial Hasse invariants H_β as β runs in \mathbb{B} . As we have seen, over \overline{X}^{PR} , the partial Hasse invariants H_β can be further factorized into a product of generalized partial Hasse invariants. In the following, we define a notion of filtration by considering all generalized partial Hasse invariants; see also [5, §1.4]. The following lemma is clear.

Lemma 6.1. *Let X be a noetherian scheme. Let \mathcal{L}, \mathcal{M} be line bundles on X . Let $h \in H^0(X, \mathcal{L})$ and $f \in H^0(X, \mathcal{M})$ be nonzero sections. Assume that the divisor of h intersects every irreducible component of X . Then there is a maximal $m \geq 0$ such that $f = h^m g$ for some $g \in H^0(X, \mathcal{M} \otimes \mathcal{L}^{-m})$.*

Let $0 \neq f \in M_{\mathbf{k}}(\mathbf{n}, \overline{\mathbb{F}}_p)$. For each $\tau \in \Sigma$, we let a_τ be the maximal power of H_τ that divides f . We define the filtration of f to be

$$\Phi(f) = \mathbf{k} - \sum_{\tau} a_\tau \mathbf{h}_\tau.$$

Since \overline{X}^{PR} is smooth, and since for $\tau_1 \neq \tau_2$ the divisors of H_{τ_1} and H_{τ_2} have no common components (by part (2) of Proposition 5.8), we find that f is divisible by $\prod_{\tau} (H_\tau)^{a_\tau}$. It follows that $\sum_{\tau} a_\tau \mathbf{e}_\tau$ is the unique maximal element of the set

$$\left\{ \sum_{\tau} x_\tau \mathbf{e}_\tau \in \mathbb{Z}_{\geq 0}^\Sigma \mid f = g \prod_{\tau} (H_\tau)^{x_\tau} \text{ for some } g \in M_{\mathbf{k} - \sum_{\tau} x_\tau \mathbf{h}_\tau}(\mathbf{n}, \overline{\mathbb{F}}_p) \right\}$$

under the usual partial ordering. In particular, $\Phi(f)$ is the weight of g corresponding to the above maximal element.

7. PAPPAS-RAPOPORT MODELS IN IWAHORI LEVEL

Fix \mathfrak{p} a prime of \mathcal{O}_F above p . Recall that $\{\mathfrak{J}_1, \dots, \mathfrak{J}_r\}$ is a set of representatives for $\text{Cl}^+(F)$. For each $1 \leq i \leq r$, there is a unique $1 \leq \gamma(i) \leq r$ such that $\mathfrak{p}\mathfrak{J}_i$ and $\mathfrak{J}_{\gamma(i)}$ represent the same class in $\text{Cl}^+(F)$. We fix once and for all a choice of a totally positive isomorphism $\epsilon_i : \mathfrak{J}_{\gamma(i)} \cong \mathfrak{p}\mathfrak{J}_i$.

Let S be a locally noetherian \mathcal{O} -scheme and $\underline{A}/S \equiv (A/S, \iota_A, \lambda_A, \alpha_A, \omega_{A/S}^\bullet)$ an S -point of \mathbb{X}^{PR} . Then $A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1}$ can be naturally equipped with a structure of a HBAS $\underline{A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1}}$. We make this into a filtered HBAS by adding a Pappas-Rapoport filtration on $\omega_{(A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1})/S} \cong \omega_{A/S} \otimes_{\mathcal{O}_F} \mathfrak{p}$ by applying $-\otimes_{\mathcal{O}_F} \mathfrak{p}$ to the Pappas-Rapoport filtration on $\omega_{A/S}$.

Let $\underline{A}/S = (A/S, \iota_A, \lambda_A, \alpha_A, \omega_{A/S}^\bullet)$ and $\underline{B}/S = (B/S, \iota_B, \lambda_B, \alpha_B, \omega_{B/S}^\bullet)$ correspond, respectively, to S -points of $X_{\mathfrak{J}_i}^{\text{PR}}$ and $X_{\mathfrak{J}_{\gamma(i)}}^{\text{PR}}$, for some $1 \leq i \leq r$. An isogeny $f : A \rightarrow B$ is called a cyclic $(\mathcal{O}_F/\mathfrak{p})$ -isogeny of filtered HBAS's if

- (1) f is an \mathcal{O}_F -isogeny of degree $p^{f_{\mathfrak{p}}}$;

- (2) f is compatible with the polarizations, i.e., the following diagram is commutative

$$\begin{array}{ccc} \mathcal{P}_B & \xrightarrow{\lambda_B} & \mathfrak{J}_{\gamma(i)} \\ f^* \downarrow & & \downarrow \bar{\epsilon}_i \\ \mathcal{P}_A & \xrightarrow{\lambda_A} & \mathfrak{J}_i \end{array}$$

where $\bar{\epsilon}_i$ is the composition $\mathfrak{J}_{\gamma(i)} \cong_{\epsilon_i} \mathfrak{p}\mathfrak{J}_i \subset \mathfrak{J}_i$, and $f^*(\eta) = f^\vee \eta f$;

- (3) f is compatible with the level structures, i.e., $\alpha_B = f \circ \alpha_A$;
 (4) f preserves the Pappas-Rapoport filtrations, i.e., for all $\beta \in \mathbb{B}$, the induced map $f_\beta^* : \omega_{B/S, \beta} \rightarrow \omega_{A/S, \beta}$ preserves the filtrations $\omega_{A/S, \beta}^\bullet$ and $\omega_{B/S, \beta}^\bullet$.

Let $Y_{\mathfrak{J}_i}^{\text{PR}}/\mathcal{O}$ be the Hilbert modular scheme classifying the isomorphism classes of tuples $(\underline{A}/S, \underline{B}/S, f : A \rightarrow B, g : B \rightarrow A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1})$, where

- S is a locally Noetherian \mathcal{O} -scheme;
- $\underline{A}/S = (A/S, \iota_A, \lambda_A, \alpha_A, \omega_{A/S}^\bullet)$ and $\underline{B}/S = (B/S, \iota_B, \lambda_B, \alpha_B, \omega_{B/S}^\bullet)$ correspond to S -points of $X_{\mathfrak{J}_i}^{\text{PR}}$ and $X_{\mathfrak{J}_{\gamma(i)}}^{\text{PR}}$, respectively;
- f and g are cyclic $(\mathcal{O}_F/\mathfrak{p})$ -isogenies of filtered HBAS's;
- The isogeny $g \circ f : A \rightarrow A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1}$ is the natural map induced by the inclusion of \mathcal{O}_F in \mathfrak{p}^{-1} . (Equivalently, the isogeny $f \circ (g \otimes_{\mathcal{O}_F} \mathfrak{p}) : B \otimes_{\mathcal{O}_F} \mathfrak{p} \rightarrow B$ is the natural map induced by the inclusion of \mathfrak{p} in \mathcal{O}_F .)

The Hilbert modular variety $Y_{\mathfrak{J}_i}^{\text{PR}}$ is independent of the choice of ϵ_i up to isomorphism. We define

$$Y^{\text{PR}} = \bigcup_{i=1}^r Y_{\mathfrak{J}_i}^{\text{PR}}.$$

Let $\bar{Y}^{\text{PR}} = Y^{\text{PR}} \otimes_{\mathcal{O}} \mathbb{F}$ and denote by π_1 and π_2 the forgetful maps

$$\pi_1 : \bar{Y}^{\text{PR}} \rightarrow \bar{X}^{\text{PR}}, \quad \pi_2 : \bar{Y}^{\text{PR}} \rightarrow \bar{X}^{\text{PR}}$$

where $\pi_1(\underline{A}, \underline{B}, f, g) = \underline{A}$, and $\pi_2(\underline{A}, \underline{B}, f, g) = \underline{B}$.

We recall a stratification defined on \bar{Y}^{PR} in [10] when p is unramified in F , and in [8] in the general case. (See also [12] for a similar construction in a related unitary case). Let $(\underline{A}, \underline{B}, f, g)$ be the universal tuple defined over \bar{Y}^{PR} . Since for all $\beta \in \mathbb{B}$, the pullback morphisms $f_\beta^* : \omega_{B/S, \beta} \rightarrow \omega_{A/S, \beta}$ and $g^* : \omega_{A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1}/S, \beta} \rightarrow \omega_{A/S, \beta}$ preserve Pappas-Rapoport filtrations, we have induced morphisms

$$f_{\beta, i}^* : \omega_{B/S, \beta, i} \rightarrow \omega_{A/S, \beta, i},$$

$$g_{\beta, i}^* : \omega_{(A \otimes_{\mathcal{O}_F} \mathfrak{p}^{-1})/S, \beta, i} \rightarrow \omega_{B/S, \beta, i},$$

where for each $\beta \in \mathbb{B}_{\mathfrak{p}}$, i runs between 1 and $e_{\mathfrak{p}}$. For $\tau = \tau_{\beta, i} \in \Sigma_{\mathfrak{p}}$, let U_τ be the vanishing locus of $f_{\beta, i}^*$ on \bar{Y}^{PR} , and V_τ the vanishing locus of $g_{\beta, i}^*$ on \bar{Y}^{PR} . For φ, η subsets of $\Sigma_{\mathfrak{p}}$ satisfying $\sigma^{-1}\varphi \cup \eta = \Sigma_{\mathfrak{p}}$, we define

$$Z_{\varphi, \eta} = \bigcap_{\tau \in \varphi} U_{\sigma^{-1}\tau} \cap \bigcap_{\tau \in \eta} V_\tau.$$

Notice the shift in φ in our notation versus that in [8], which is consistent with the notation in [10]. One can show that the collection of $Z_{\varphi, \eta}$ defined above form a stratification of \bar{Y}^{PR} . The local and global properties of the strata are studied in

detail in [10] and [8]. In the following, we recall some facts about this stratification that we will use in our argument.

Let r be a non-negative integer. A morphism of \mathbb{F} -schemes $\lambda: Z_1 \rightarrow Z_2$ is called an r -Frobenius factor if there is a morphism $\lambda': Z_2 \rightarrow Z_1$ such that $\lambda\lambda': Z_2 \rightarrow Z_2$ is the r -th power of the absolute Frobenius morphism of Z_2 . By [12, Proposition 4.8], if Z_1, Z_2 are schemes of finite type over \mathbb{F} such that Z_2 is normal, then any proper morphism $\lambda: Z_1 \rightarrow Z_2$ which is a bijection on points is an r -Frobenius factor for some $r \geq 0$.

Fix $\tau_0 \in \Sigma_{\mathbf{p}}$. Recall the definition of \mathbb{H}_{τ_0} in (3.1). We let

$$\text{pr}: \mathbb{P} \rightarrow Z_{\sigma\tau_0}$$

be the \mathbb{P}^1 bundle given on $Z_{\sigma\tau_0} \subset X^{\text{PR}}$ by $\mathbb{P}(\mathbb{H}_{\tau_0})$. Let $\{\tau_0\}^c = \Sigma_{\mathbf{p}} - \{\tau_0\}$.

Proposition 7.1. *If $|\Sigma_{\mathbf{p}}| > 1$, then $\pi_1(Z_{\{\sigma\tau_0\}, \{\tau_0\}^c}) = Z_{\sigma\tau_0}$, and $\pi_2(Z_{\{\sigma\tau_0\}, \{\tau_0\}^c}) = Z_{\tau_0}$. Furthermore, there is an integer $r \geq 0$, and an r -Frobenius factor λ which makes the following diagram commutative:*

$$(7.1) \quad \begin{array}{ccc} Z_{\{\sigma\tau_0\}, \{\tau_0\}^c} & \xrightarrow{\lambda} & \mathbb{P} \\ & \searrow \pi_1 & \downarrow \text{pr} \\ & & Z_{\sigma\tau_0}. \end{array}$$

Proof. See [10, Thm 2.6.4. (2)] and [8, Prop. 4.5.], noting the difference in notation between ours and [8]: the stratum $Z_{\{\sigma\tau_0\}, \{\tau_0\}^c}$ is denoted $Y_{\{\tau_0\}, \{\tau_0\}^c}$ in *loc. cit.* For the final statement, note that from definitions we have $\pi_2(Z_{\{\sigma\tau_0\}, \{\tau_0\}^c}) = \pi_1(Z_{\{\sigma\tau_0\}^c, \{\tau_0\}}) = Z_{\tau_0}$. \square

For $\tau = \tau_{\beta, i} \in \Sigma_{\mathbf{p}}$, we define $n_{\tau} = p$ if $i = 1$, and $n_{\tau} = 1$ otherwise. By the proposition, there exists $\lambda': \mathbb{P} \rightarrow Z_{\{\sigma\tau_0\}, \{\tau_0\}^c}$ such that $\lambda\lambda': \mathbb{P} \rightarrow \mathbb{P}$ is the r -th power of the absolute Frobenius morphism. We will be considering the following composite morphism in the proof of our main theorem:

$$\mathbb{P} \xrightarrow{\lambda'} Z_{\{\sigma\tau_0\}, \{\tau_0\}^c} \xrightarrow{\pi_2} Z_{\tau_0}.$$

Lemma 7.2. *Let $\tau_0 \in \Sigma_{\mathbf{p}}$ and assume that $f_{\mathbf{p}} > 1$. Let x be a geometric point of $Z_{\sigma\tau_0}$, and denote by \mathbb{P}_x^1 the fibre at x of the \mathbb{P}^1 -bundle \mathbb{P} . Then $n_{\tau_0} | p^r$ and for $\tau \in \Sigma_{\mathbf{p}}$ we have:*

$$(\pi_2\lambda')^*\omega_{\tau}|_{\mathbb{P}_x^1} \cong \begin{cases} \mathcal{O}_{\mathbb{P}_x^1}(p^r) & \text{if } \tau = \tau_0, \\ \mathcal{O}_{\mathbb{P}_x^1}(-p^r/n_{\tau_0}) & \text{if } \tau = \sigma^{-1}\tau_0, \\ \mathcal{O}_{\mathbb{P}_x^1} & \text{otherwise.} \end{cases}$$

Proof. Consider the following diagram

$$(7.2) \quad \begin{array}{ccccc} \mathbb{P} & \xrightarrow{\lambda'} & Z_{\{\sigma\tau_0\}, \{\tau_0\}^c} & \xrightarrow{\lambda} & \mathbb{P} \\ \pi_2\lambda' \downarrow & & \swarrow \pi_2 & & \downarrow \text{pr} \\ & & & \searrow \pi_1 & \\ & & & & Z_{\sigma\tau_0}. \end{array}$$

The following statements are consequences of [8, Prop. 4.7] (note that $Z_{\{\sigma\tau_0\},\{\tau_0\}^c}$ is denoted $Y_{\{\tau_0\},\{\tau_0\}^c}$ in *loc. cit.*).

- (1) $\pi_2^* \omega_{\tau_0} = \lambda^* \mathcal{O}_{\mathbb{P}}(1)$;
- (2) $\pi_2^* \omega_{\sigma^{-1}\tau_0}^{\otimes n_{\tau_0}} = \lambda^* \mathcal{O}_{\mathbb{P}}(-1)$;
- (3) $\pi_2^* \omega_{\tau}^{\otimes n_{\sigma\tau}} = \pi_1^* \omega_{\sigma\tau}$, if $\tau \notin \{\tau_0, \sigma^{-1}\tau_0\}$.

From (1) it follows that $(\pi_2 \lambda')^* \omega_{\tau_0} = \lambda'^* \lambda^* \mathcal{O}_{\mathbb{P}}(1) = \mathcal{O}_{\mathbb{P}}(p^r)$. From (2) it follows that $(\pi_2 \lambda')^* \omega_{\sigma^{-1}\tau_0}^{\otimes n_{\tau_0}} = \mathcal{O}_{\mathbb{P}}(-p^r)$. In particular, this implies that $n_{\tau_0} | p^r$, and hence we find $(\pi_2 \lambda')^* \omega_{\sigma^{-1}\tau_0} |_{\mathbb{P}_x^1} = \mathcal{O}_{\mathbb{P}_x^1}(-p^r/n_{\tau_0})$. Finally, (3) implies that for $\tau \neq \tau_0, \sigma^{-1}\tau_0$ we have $(\pi_2 \lambda')^* \omega_{\tau}^{\otimes n_{\sigma\tau}} = (\lambda')^* \pi_1^* \omega_{\sigma\tau} = (\lambda \lambda')^* \text{pr}^* \omega_{\sigma\tau}$. Since $\text{pr}^* \omega_{\sigma\tau}$ restricted to every fibre of pr is the trivial line bundle, it follows that $(\pi_2 \lambda')^* \omega_{\tau}^{\otimes n_{\sigma\tau}}$ restricted to \mathbb{P}_x^1 is trivial, and hence so is the restriction of $(\pi_2 \lambda')^* \omega_{\sigma\tau}$ to each fibre \mathbb{P}_x^1 . \square

8. THE MAIN THEOREM

In this section, we prove our main theorem and discuss some of its consequences. Given what we have proven in the previous sections, the rest of the proof is in essence similar to the case where p is unramified in \mathcal{O}_F , treated in [6]. We would like to emphasize again that in this more general proof we bypass the use of deep geometric results on global properties of the Goren-Oort strata on \overline{X}^{PR} (currently unavailable in the case p is ramified in F), and instead appeal to properties of a stratification on \overline{Y}^{PR} that are more readily generalizable to other Shimura varieties.

Recall that for $\tau \in \Sigma$, we have $H_{\tau} \in M_{\mathbf{h}_{\tau}}(\mathbf{n}; \mathbb{F})$, with $\mathbf{h}_{\tau} = n_{\tau} \mathbf{e}_{\sigma^{-1}\tau} - \mathbf{e}_{\tau}$, where $n_{\tau} = p$ if $i = 1$, and $n_{\tau} = 1$ otherwise.

Theorem 8.1. *Let $\mathfrak{p}|p$ be a prime of \mathcal{O}_F . Let $\mathbf{k} = \sum k_{\tau} \mathbf{e}_{\tau} \in \mathbb{Z}^{\Sigma}$ be a weight. Suppose that we have*

$$n_{\tau_0} k_{\tau_0} < k_{\sigma^{-1}\tau_0},$$

for some $\tau_0 \in \Sigma_{\mathfrak{p}} \subset \Sigma$. Then, multiplication by the generalized partial Hasse invariant H_{τ_0} induces an isomorphism:

$$M_{\mathbf{k}-\mathbf{h}_{\tau_0}}(\mathbf{n}; \overline{\mathbb{F}}_p) \xrightarrow{\sim} M_{\mathbf{k}}(\mathbf{n}; \overline{\mathbb{F}}_p).$$

When $\Sigma_{\mathfrak{p}} = \{\tau_0\}$, a stronger statement holds: if $k_{\tau_0} < 0$, then $M_{\mathbf{k}}(\mathbf{n}; \overline{\mathbb{F}}_p) = 0$.

Proof. First we assume $|\Sigma_{\mathfrak{p}}| > 1$. We have a short exact sequence

$$0 \rightarrow \omega^{\mathbf{k}-\mathbf{h}_{\tau_0}} \rightarrow \omega^{\mathbf{k}} \rightarrow \mathcal{F} \rightarrow 0$$

of sheaves on \overline{X}^{PR} , where the first map is multiplication by H_{τ_0} . The quotient sheaf \mathcal{F} is supported on Z_{τ_0} , and we have $H^0(\overline{X}^{\text{PR}}, \mathcal{F}) = H^0(Z_{\tau_0}, \omega^{\mathbf{k}})$. It is hence enough to show that $H^0(Z_{\tau_0}, \omega^{\mathbf{k}}) = 0$ under the assumptions. Consider the morphism

$$\pi_2 \lambda': \mathbb{P} \rightarrow Z_{\tau_0}$$

(see Proposition 7.1). Since Z_{τ_0} is reduced and $\pi_2 \lambda'$ is surjective, we have an injection

$$(\pi_2 \lambda')^*: H^0(Z_{\tau_0}, \omega^{\mathbf{k}}) \hookrightarrow H^0(\mathbb{P}, (\pi_2 \lambda')^* \omega^{\mathbf{k}}),$$

It is therefore enough to show that $H^0(\mathbb{P}, (\pi_2 \lambda')^* \omega^{\mathbf{k}}) = 0$. We prove that for any $s \in H^0(\mathbb{P}, (\pi_2 \lambda')^* \omega^{\mathbf{k}})$, the restriction of s to every geometric fibre of the \mathbb{P}^1 -bundle

$\text{pr} : \mathbb{P} \rightarrow Z_{\sigma\tau_0}$ is zero. Let x be a geometric point of $Z_{\sigma\tau_0}$, and \mathbb{P}_x^1 the fibre over x . By Lemma 7.2, we have

$$(\pi_2\lambda')^*\omega^{\mathbf{k}}|_{\mathbb{P}_x^1} \cong \mathcal{O}_{\mathbb{P}_x^1}(p^rk_{\tau_0} - (p^r/n_{\tau_0})k_{\sigma^{-1}\tau_0}).$$

By assumption $p^rk_{\tau_0} - (p^r/n_{\tau_0})k_{\sigma^{-1}\tau_0} < 0$, which implies that $s|_{\mathbb{P}_x^1} = 0$ as desired.

Now, assume $\Sigma_{\mathbf{p}} = \{\tau_0\}$. We prove that if $k_{\tau_0} < 0$, then $H^0(\overline{X}^{\text{PR}}, \omega^{\mathbf{k}}) = 0$. Let $\Sigma = \{\tau_0, \tau_1, \dots, \tau_{d-1}\}$. Let $T_i := \{\tau_{i+1}, \dots, \tau_{d-1}\}$ for $0 \leq i \leq d-1$ (where $T_{d-1} = \emptyset$). We prove by induction that $H^0(Z_{T_i}, \omega^{\mathbf{k}}) = 0$ for all $0 \leq i \leq d-1$. Since $Z_{T_{d-1}} = \overline{X}^{\text{PR}}$, this proves the claim.

We first prove the result for $i = 0$. Let C be a component of Z_{T_0} . Part (2) of Proposition 5.8 shows that C is a projective curve. By Corollary 5.7, $H_{\tau_0} \in H^0(\overline{X}^{\text{PR}}, \omega_{\tau_0}^{p-1})$ vanishes at some point on C . It follows that ω_{τ_0} has positive degree on C . By Corollary 5.3, every ω_{τ} for $\tau \neq \tau_0$ restricts to a torsion line bundle on C (since $f_{\mathbf{p}}=1$, and consequently $\Sigma_{\mathbf{q}} \subset T_0$ for all $\mathbf{q} \neq \mathbf{p}$). It follows that every ω_{τ} for $\tau \neq \tau_0$ has degree zero on C . This shows that $\deg(\omega^{\mathbf{k}}|_C) = k_{\tau_0} \deg(\omega_{\tau_0}|_C) < 0$, from which it follows that $H^0(C, \omega^{\mathbf{k}}) = 0$.²

Now assume that $1 \leq i \leq d-1$, and we have $H^0(Z_{T_{i-1}}, \omega^{\mathbf{k}}) = 0$ for all \mathbf{k} with $k_{\tau_0} < 0$. We prove that $H^0(Z_{T_i}, \omega^{\mathbf{k}}) = 0$ for all such \mathbf{k} as well. Consider the short exact sequence of sheaves on Z_{T_i}

$$0 \rightarrow \omega^{\mathbf{k}-\mathbf{h}_{\tau_i}}|_{Z_{T_i}} \rightarrow \omega^{\mathbf{k}}|_{Z_{T_i}} \rightarrow \mathcal{F}_i \rightarrow 0,$$

where the first map is multiplication by H_{τ_i} . The sheaf \mathcal{F}_i is supported on the $Z_{T_{i-1}}$, and we have $H^0(Z_{T_i}, \mathcal{F}_i) = H^0(Z_{T_{i-1}}, \omega^{\mathbf{k}}) = 0$. This shows that multiplication by H_{τ_i} induces an isomorphism

$$H^0(Z_{T_i}, \omega^{\mathbf{k}-\mathbf{h}_{\tau_i}}) \xrightarrow{\sim} H^0(Z_{T_i}, \omega^{\mathbf{k}}).$$

The same result holds after replacing \mathbf{k} with $\mathbf{k}' = \mathbf{k} - M\mathbf{h}_{\tau_i}$ for any $M > 0$, as $k'_{\tau_0} = k_{\tau_0} < 0$. It follows that for all $M > 0$, multiplication by $(H_{\tau_i})^M|_{Z_{T_i}}$ induces an isomorphism

$$H^0(Z_{T_i}, \omega^{\mathbf{k}-M\mathbf{h}_{\tau_i}}) \xrightarrow{\sim} H^0(Z_{T_i}, \omega^{\mathbf{k}}).$$

By Corollary 5.7, $H_{\tau_i}^{\mathfrak{J}}$ has a nonempty vanishing locus on every component of Z_{T_i} . Lemma 6.1 implies that any nonzero section of $\omega^{\mathbf{k}}$ on Z_{T_i} is divisible by at most a finite power of $H_{\tau_i}^{\mathfrak{J}}|_{Z_{T_i}}$. It follows that $H^0(Z_{T_i}, \omega^{\mathbf{k}}) = 0$. This proves the claim by induction. In particular, for $i = d-1$, we get

$$H^0(\overline{X}^{\text{PR}}, \omega^{\mathbf{k}}) = H^0(Z_{T_{d-1}}, \omega^{\mathbf{k}}) = 0.$$

□

In the following, we state two corollaries of our main theorem which answer questions Andreatta and Goren asked in [1]. We have previously answered these questions in the case p is unramified in \mathcal{O}_F in [6]. We first define the *minimal*, *standard*, and *Hasse* cones.

$$C^{\min} = \left\{ \sum_{\tau} x_{\tau} \mathbf{e}_{\tau} \in \mathbb{Q}^{\Sigma} \mid n_{\tau} x_{\tau} \geq x_{\sigma^{-1}\tau} \text{ for all } \tau \in \Sigma \right\},$$

²In fact, by [14, Lecture 11, (VI.)], for this argument we only need that C is Cohen-Macaulay, which is already immediate from Part (2) of Proposition 5.8.

$$C^{\text{st}} = \left\{ \sum_{\tau} x_{\tau} \mathbf{e}_{\tau} \in \mathbb{Q}^{\Sigma} \mid x_{\tau} \geq 0 \text{ for all } \tau \in \Sigma \right\},$$

$$C^{\text{Hasse}} = \left\{ \sum_{\tau} y_{\tau} \mathbf{h}_{\tau} \in \mathbb{Q}^{\Sigma} \mid y_{\tau} \geq 0 \text{ for all } \tau \in \Sigma \right\}.$$

It is easy to see that $C^{\min} \subset C^{\text{st}} \subset C^{\text{Hasse}}$, and equality holds if and only if p is totally split in F .

Corollary 8.2. *If $0 \neq f \in M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p)$ for some weight $\mathbf{k} \in \mathbb{Z}^{\Sigma}$, then $\Phi(f) \in C^{\min}$.*

Proof. Let $\Phi(f) = \sum_{\tau} w_{\tau} \mathbf{e}_{\tau}$. Write $f = g \prod_{\tau} (H_{\tau})^{a_{\tau}}$, where the integers a_{τ} are non-negative, and $g \in M(\mathfrak{n}, \overline{\mathbb{F}}_p)$ is not divisible by any generalized partial Hasse invariant. If $\Phi(f) \notin C^{\min}$, then for some $\tau \in \Sigma$ we have $n_{\tau} w_{\tau} < w_{\sigma^{-1}\tau}$. It then follows by Theorem 8.1 that g is divisible by H_{τ} , which is a contradiction. \square

Corollary 8.3. *If $M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p) \neq 0$, then $\mathbf{k} \in C^{\text{Hasse}}$.*

Proof. Assume $0 \neq f \in M_{\mathbf{k}}(\mathfrak{n}, \overline{\mathbb{F}}_p)$. By Corollary 8.2, $\Phi(f) \in C^{\min} \subset C^{\text{Hasse}}$. By definition of filtration, there are integers $a_{\tau} \geq 0$ such that

$$\mathbf{k} = \Phi(f) + \sum_{\tau} a_{\tau} \mathbf{h}_{\tau} \in C^{\text{Hasse}}.$$

\square

REFERENCES

- [1] Fabrizio Andreatta, Eyal Z. Goren, *Hilbert modular forms: mod p and p -adic aspects*, Mem. Amer. Math. Soc. **173** (2005), no. 819.
- [2] Fabrizio Andreatta, Eyal Z. Goren *Geometry of Hilbert modular varieties over totally ramified primes*, IMRN, Volume 2003, Issue 33, (2003), 1785–1835.
- [3] Ching-Li Chai, *Arithmetic minimal compactification of the Hilbert-Blumenthal moduli spaces*, Appendix to The Iwasawa conjecture for totally real fields by A. Wiles. Ann. of Math. (2) **131** (1990), no. 3, 541–554.
- [4] Pierre Deligne, Georgios Pappas, *Singularités des espaces de modules de Hilbert en caractéristiques divisant le discriminant*, Compos. Math. **90** (1994), no. 1, 59–79.
- [5] Shaunak V. Deo, Mladen Dimitrov, Gabor Wiese, *Unramifiedness of weight one Hilbert Hecke algebras*, preprint, available at <https://arxiv.org/abs/1911.11196>.
- [6] Fred Diamond, Payman L Kassaei, *Minimal weights of Hilbert modular forms in characteristic p* , Compos. Math. **153** (2017), no. 9, 1769–1778.
- [7] Fred Diamond, Shu Sasaki, *A Serre weight conjecture for geometric Hilbert modular forms in characteristic p* , preprint, available at <https://arxiv.org/abs/1712.03775>, 2017.
- [8] Matthew Emerton, Davide Reduzzi, Liang Xiao, *Unramifiedness of Galois representations arising from Hilbert modular surfaces*, Forum Math. Sigma **5** (2017), e29, 70.
- [9] Wushi Goldring, Jean-Stefan Koskivirta, *Automorphic vector bundles on G Zip-schemes*, Compositio Math., **154** (2018), pp. 2586–2605.
- [10] Eyal Goren, Payman L Kassaei, *Canonical subgroups over Hilbert modular varieties*, J. Reine Angew. Math. **670** (2012), 1–63.
- [11] Eyal Z. Goren, Frans Oort, *Stratifications of Hilbert modular varieties*, J. Algebraic Geom. **9** (2000), no. 1, 111–154.
- [12] David Helm, *A geometric Jacquet-Langlands correspondence for $U(2)$ Shimura varieties*, Israel Journal of Mathematics, volume 187, 3780 (2012).
- [13] David Helm, *Towards a geometric Jacquet-Langlands correspondence for unitary Shimura varieties*, Duke Math. J. **155** (2010), no. 3, 483–518.
- [14] David Mumford, *Lectures on curves on an algebraic surface*, Annals of Math. Studies **59**, Princeton Univ. Press, 1966.

- [15] Giorgios Pappas, Michael Rapoport, *Local models in the ramified case. II. Splitting models*, Duke Math. J. **127** (2005), no. 2, 193–250.
- [16] Davide Reduzzi, Liang Xiao, *Partial Hasse invariants on splitting models of Hilbert modular varieties*, Annales Scientifiques de l'ENS **50** (2017), 579–607.
- [17] Shu Sasaki, *Integral models of Hilbert modular varieties in the ramified case, deformations of modular Galois representations, and weight one forms*, Invent. Math. **215** (2018), 171–264.
- [18] The Stacks Project Contributors, *The Stacks Project*, open source textbook, available at <http://stacks.math.columbia.edu>.
- [19] Yichao Tian, Liang Xiao, *On Goren–Oort stratification for quaternionic Shimura varieties*, Comp. Math. **152** (2016), no. 10, 2134–2220.

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, WC2R 2LS, UK
E-mail address: `fred.diamond@kcl.ac.uk`

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, WC2R 2LS, UK
E-mail address: `payman.kassaei@kcl.ac.uk`