

ON THE COMPACTNESS OF BERGMAN-TYPE INTEGRAL OPERATORS

LJIA DING AND JUNMEI FAN

ABSTRACT. Bergman-type integral operators are classical operators in complex analysis and operator theory. Recently, the first author and his collaborator [9] completely characterized the L^p - L^q boundedness of Bergman-type integral operators K_α, K_α^+ and the L^p - L^q compactness of K_α on the unit ball. In this paper, we will use a substantially new method to completely characterize the L^p - L^q compactness of K_α^+ , but also prove that the L^p - L^q compactness of operators K_α, K_α^+ is in fact equivalent. Moreover, we completely characterize Schatten class and Macaev class Bergman-type integral operator K_α on L^2 space and Bergman space via inequalities related to the dimension of the unit ball, and we also give an intrinsic characterization by introducing the concept of Hausdorff dimension of compact operators. The Dixmier trace of K_α are also calculated in this paper.

1. INTRODUCTION

This paper is a systematic research on the L^p - L^q compactness of Bergman-type integral operators on the unit ball. While there exist abundant works on the boundedness of Bergman-type integral operators, the investigation of compactness aspect was started only a few years.

Let us start with recalling some notations and terminologies. Let \mathbb{B}^d be the open unit ball in the usual complex Euclidian norm on the d -dimensional space \mathbb{C}^d with the normalized Lebesgue measure dv , which means the measure of \mathbb{B}^d is one. For any $\alpha \in \mathbb{R}$, the Bergman-type integral operator K_α on $L^1(\mathbb{B}^d, dv)$ is defined by

$$K_\alpha f(z) = \int_{\mathbb{B}^d} \frac{f(w)}{(1 - \langle z, w \rangle)^\alpha} dv(w),$$

where $\langle z, w \rangle = z_1 \bar{w}_1 + \cdots + z_d \bar{w}_d$ is the standard Hermitian inner product on \mathbb{C}^d . In particular, when $\alpha = d + 1$, K_{d+1} is the standard Bergman projection on \mathbb{B}^d , since the function $K(z, w) = \frac{1}{(1 - \langle z, w \rangle)^{d+1}}$ is the Bergman kernel of \mathbb{B}^d with the measure dv . However, the Bergman-type integral operator K_α^+ on $L^1(\mathbb{B}^d, dv)$ is given by

$$K_\alpha^+ f(z) = \int_{\mathbb{B}^d} \frac{f(w)}{|1 - \langle z, w \rangle|^\alpha} dv(w).$$

Bergman-type integral operators play an important role in complex analysis and operator theory [19, 25, 26, 27, 28]. In particular, for any $\alpha > 0$, if restrict K_α to Bergman spaces, then every K_α is a special fractional radial differential operator [25, 26, 27]

2010 *Mathematics Subject Classification.* 32A25; 47B07; 47B10; 42B20.

Key words and phrases. Bergman kernel; Compact operator; Dixmier trace; Hausdorff dimension; Schatten class; Macaev class.

The first author was partially supported by PCPSF (2020T130016).

which is a kind of useful operators in the Bergman space theory on the unit ball, see Lemma 3.6 below. On the other hand, the operator K_α^+ in the complex analysis category, to some extent, is analogous to the Riesz potential operators in the real analysis. The classical Riesz potential operator R_α is defined on the real Euclidian space \mathbb{R}^d , whose basic result concerning mapping properties is the Hardy-Littlewood-Sobolev theorem, which characterizes the boundedness of Riesz potential operator $R_\alpha : L^p(\mathbb{R}^d) \rightarrow L^q(\mathbb{R}^d)$ and has been applied to the PDE theory for a long time; we refer the reader to [8, 15, 21].

For convenience, we replace $L^p(\mathbb{B}^d, dv)$ by $L^p(\mathbb{B}^d)$ or L^p for any $1 \leq p \leq \infty$ without confusion arises. Bergman-type operators K_α, K_α^+ are called L^p - L^q bounded (or compact) if $K_\alpha, K_\alpha^+ : L^p(\mathbb{B}^d) \rightarrow L^q(\mathbb{B}^d)$ are bounded (or compact), where $1 \leq p, q \leq \infty$. Researches on L^p - L^q problems of Bergman operators K_α, K_α^+ go back to the boundedness of Bergman projection on bounded domains. F. Forelli and W. Rudin [11] proved that the Bergman projection on the unit ball is L^p - L^p bounded if and only if $1 < p < \infty$, in their method an (Forelli-Rudin) asymptotic estimate of integral for Bergman kernel plays an important role; indeed, following the same method, one can characterize the L^p - L^p boundedness of more general Bergman-type operators, see [26, 27]. Around the same time, H. Phong and E. Stein [20] proved more general results on a class of bounded strongly pseudoconvex domains; see [14, 17, 18] for more results along this line.

Recently, G. Cheng and X. Fang et al [4] completely characterized the L^p - L^q boundedness of K_α on the unit disk \mathbb{D} , i.e. the case $d = 1$, and they conjectured that there should exist similar results in the high dimensional case. Their proofs depend on techniques of harmonic analysis and coefficient multiplier theory of Bergman space on the unit disk. Unfortunately, this method can not be directly applied to the case of unit ball. But, for the spacial case $\alpha = 1$, G. Cheng, C. Liu et al [5] solved the L^p - L^q boundedness problem of K_1, K_1^+ on \mathbb{B}^d . The author and his collaborator [9] completely characterized the L^p - L^q boundedness of K_α, K_α^+ and established some Hardy-Littlewood-Sobolev type inequalities on \mathbb{B}^d , motivated by an observation in [27] we further investigated the L^p - L^q compactness of K_α . Techniques of complex and harmonic analysis are synthetically utilized in [9], such as the Carleson measure theory is applied to the L^p - L^q compactness problem. The key method used for compactness relies on the fact that K_α maps L^p space into Bergman space or Bloch space; however, K_α^+ has no such analytic property, thus the compactness characterization of K_α^+ remains open.

The first purpose of the present paper is to continue to characterize the L^p - L^q compactness of K_α^+ . The following two theorems are our main results on the L^p - L^q compactness.

Theorem 1. Suppose $d+1 < \alpha < d+2$, then the following statements are equivalent.

- (1) $K_\alpha : L^p \rightarrow L^q$ is bounded.
- (2) $K_\alpha^+ : L^p \rightarrow L^q$ is bounded.
- (3) $K_\alpha : L^p \rightarrow L^q$ is compact.
- (4) $K_\alpha^+ : L^p \rightarrow L^q$ is compact.
- (5) p, q satisfy one of the following inequalities:
 - (a) $\frac{1}{d+2-\alpha} < p < \infty, \frac{1}{q} > \frac{1}{p} + \alpha - (d+1)$;

$$(b) \ p = \infty, q < \frac{1}{\alpha - (d+1)}.$$

The equivalences of (1),(2),(3) and (5) in Theorem 1 have been proved in [9], from Theorem 1 we know the L^p - L^q boundedness and compactness of operators K_α, K_α^+ are equivalent when $d+1 < \alpha < d+2$.

Theorem 2. Suppose $0 < \alpha \leq d+1$, then the following statements are equivalent:

- (1) $K_\alpha : L^p \rightarrow L^q$ is compact.
- (2) $K_\alpha^+ : L^p \rightarrow L^q$ is compact.
- (3) p, q satisfy one of the following inequalities:
 - (a) $p = 1, q < \frac{d+1}{\alpha}$;
 - (b) $1 < p < \frac{d+1}{d+1-\alpha}, \frac{1}{q} > \frac{1}{p} + \frac{\alpha}{d+1} - 1$;
 - (c) $p = \frac{d+1}{d+1-\alpha}, q < \infty$;
 - (d) $\frac{d+1}{d+1-\alpha} < p \leq \infty$.

The equivalence of (1) and (3) in Theorem 2 has been proved in [9], combing Theorem 2 with [9, Theorem 2,3], we know the L^p - L^q boundedness and compactness of operators K_α, K_α^+ are different when $0 < \alpha \leq d+1$. When $\alpha \leq 0$, Proposition 3.11 below shows that K_α, K_α^+ are all L^p - L^q compact for any $1 \leq p, q \leq \infty$; moreover, K_α is a finite rank operator if α is a nonpositive integer and K_α^+ is a finite rank operator if α is a nonpositive even integer. In contrast, when $\alpha \geq d+2$, there exist no $1 \leq p, q \leq \infty$ such that $K_\alpha^+ : L^p \rightarrow L^q$ is compact, see Corollary 3.13 below.

Theorem 2 and Proposition 3.11 imply Bergman-type operator K_α are compact on the Hilbert space L^2 only when $\alpha < d+1$. As we all known, Schatten classes and Macaev classes are more refined classification of compact operators on Hilbert spaces, but also involve global estimates of the spectrum of compact operators. The next two theorems deal with the problem of Schatten class and Macaev class Bergman-type operators. Let H be a separable Hilbert space, denote by $\mathcal{L}^p(H)$ and $\mathcal{L}^{p,\infty}(H)$ the Schatten p -class (or ideal) and Macaev p -class (or ideal) on H respectively, where $0 < p < \infty$. More precisely, let T be a compact operator on H and

$$\{\mu_n(|T|) : \mu_0(|T|) \geq \mu_1(|T|) \geq \mu_2(|T|) \geq \dots\} \quad (1.1)$$

be the set of eigenvalues (counting multiplicities) of $|T| := (T^*T)^{\frac{1}{2}}$, then $T \in \mathcal{L}^p(H)$ if and only if $\{\mu_n(|T|)\} \in \ell^p$, and $T \in \mathcal{L}^{p,\infty}(H)$ if and only if $\mu_n(|T|) = O(n^{-\frac{1}{p}})$. We also use $\mathcal{L}^\infty(H) = \mathcal{L}^{\infty,\infty}(H)$ to stand for the compact operator ideal on H . Moreover, $\mathcal{L}^p(H)$ and $\mathcal{L}^{p,\infty}(H)$ will become Banach spaces when provided suitable norms for $1 \leq p < \infty$. We refer the reader to [7, 23, 27, 28] for more details about Schatten class and Macaev class operators. Let us denote the set $\mathcal{S}_d \subset \mathbb{R}$ by

$$\mathcal{S}_d = \{\alpha < d+1 : -\alpha \notin \mathbb{N}\},$$

where \mathbb{N} is the set of all nonnegative integers. For $1 \leq p \leq \infty$, let $A^p = \text{Hol}(\mathbb{B}^d) \cap L^p$ be the p -integrable Bergman space on \mathbb{B}^d . In particular, A^2 is a separable Hilbert space.

Theorem 3. Suppose $\alpha \in \mathcal{S}_d$ and $0 < p < \infty$, then the following statements are equivalent.

- (1) $K_\alpha \in \mathcal{L}^p(L^2)$.
- (2) $K_\alpha \in \mathcal{L}^p(A^2)$.
- (3) $\widetilde{K}_\alpha \in L^p(d\lambda)$.

$$(4) \quad p > \frac{d}{d+1-\alpha}.$$

Where \widetilde{K}_α is the Berezin transform of K_α on the Bergman space A^2 and $d\lambda$ is the Möbius invariant measure on \mathbb{B}^d , the exact definition will be given in the following Section 4.

Theorem 4. Suppose $\alpha \in \mathcal{S}_d$ and $0 < p < \infty$, then the following statements are equivalent.

- (1) $K_\alpha \in \mathcal{L}^{p,\infty}(L^2)$.
- (2) $K_\alpha \in \mathcal{L}^{p,\infty}(A^2)$.
- (3) $p \geq \frac{d}{d+1-\alpha}$.

In particular, $K_\alpha \notin \mathcal{L}^{\frac{d}{d+1-\alpha}}$ but $K_\alpha \in \mathcal{L}^{\frac{d}{d+1-\alpha},\infty}$ when $\alpha \in \mathcal{S}_d$; however, $K_\alpha \in \mathcal{L}^{\frac{d}{d+1-\alpha}}$ and $K_\alpha \in \mathcal{L}^{\frac{d}{d+1-\alpha},\infty}$ if α is a nonpositive integer. As an application of Theorem 3,4 we calculate the Dixmier trace of K_α whenever $K_\alpha \in \mathcal{L}^{1,\infty}$, see Proposition 4.10 below. Moreover, we remark that the condition $\alpha \in \mathcal{S}_d$ in Theorem 3,4 is necessary and sharp, namely \mathcal{S}_d is the maximal set ensures the theorems holds. Note that Theorem 3,4 are two results related to the underlying domain \mathbb{B}^d . We also give a more intrinsic characterization of Schatten class and Macave class Bergman-type operator K_α by introducing the concept of Hausdorff dimension of compact operators, see Definition 4.8 and Corollary 4.12 below. Hausdorff dimension of compact operators is similar to the Hausdorff dimension of subsets in metric spaces [16].

It is worth mentioning that our results are totally new even in the one-dimensional case and the results of this paper can be generalized to the weighted Lebesgue integrable spaces and more general kernel operators on the unit ball. Moreover, we in fact provide two approaches to solve the Schatten class Bergman-type operators on the unit ball, one of which essentially depend on the spectrum estimate without Forelli-Rudin estimate. On the other hand, we know that the unit ball is a spacial bounded strongly pseudoconvex domain and every bounded strongly pseudoconvex domain with noncompact automorphism group is biholomorphic to the unit ball [24], so we guess that there exist some similar results about boundedness and compactness on bounded strongly pseudoconvex domains.

The paper is organized as follows. In Section 2, we give the proof of Theorem 1 by using a criteria of precompactness in L^p space and the interpolation of compact operators. Section 3 is devoted to complete the proof of Theorem 2, which is based on the hypergeometric function theory and the fractional radial differential operator theory. We also describe the phenomenon of Bergman-type operators for $\alpha \leq 0$. In Section 4, we will characterize Schatten class and Macaev class Bergman-type operator K_α by estimates of the spectrum and techniques of operator theory.

2. THE CASE OF $d+1 < \alpha < d+2$

In this section, we will prove the main Theorem 1. We need some lemmas. The following lemma gives the regularity of the image of K_α, K_α^+ .

Lemma 2.1. For any $\alpha \in \mathbb{R}$ and $f \in L^p(\mathbb{B}^d)$, $1 \leq p \leq \infty$, then the followings hold.

- (1) $K_\alpha f$ is holomorphic on \mathbb{B}^d .
- (2) $K_\alpha^+ f$ is smooth on \mathbb{B}^d .

Proof. (1) It is sufficient to show that $K_\alpha f, f \in L^1(\mathbb{B}^d)$ is holomorphic on every point of \mathbb{B}^d . Suppose z is an arbitrary point in \mathbb{B}^d . Denote $B_r = \{w \in \mathbb{C}^d : |w| < r\}, r > 0$. Choose $|z| < r_0 < r'_0 < 1$, and then choose $1 < r_1 < \frac{1}{r'_0}$. It is easy to see that $\{z\} \times \mathbb{B}^d \subset B_{r_0} \times \mathbb{B}^d \subset B_{r'_0} \times B_{r_1}$ and $0 < r_0 < r_0 r_1 < 1$. It follows that the function $K_\alpha(z, w) = \frac{1}{(1-\langle z, \bar{w} \rangle)^\alpha}$ is holomorphic on the domain $B_{r'_0} \times B_{r_1}$. Then [22, Proposition 1.2.6] implies that the binary function $K_\alpha(z, w) = \frac{1}{(1-\langle z, \bar{w} \rangle)^\alpha}$ has a global power series expansion on $B_{r'_0} \times B_{r_1}$. Therefor, we can suppose that

$$K_\alpha(z, w) = \sum_{n, m \geq 0} a_{n, m} z^n \bar{w}^m, \quad (2.1)$$

on $B_{r'_0} \times B_{r_1}$. Here n, m are multi-indexes, $n = (n_1, \dots, n_d) \geq 0$ means $n_j \geq 0$ for any j , and $z^n = \prod_{j=1}^d z_j^{n_j}$. Note that $\overline{B_{r_0}} \times \overline{\mathbb{B}^d} \subset B_{r'_0} \times B_{r_1}$, we get that the power series (2.1) is uniformly converge on compact set $\overline{B_{r_0}} \times \overline{\mathbb{B}^d}$. Then the dominated convergence theorem and the maximum modulus principle of analytic function imply that

$$K_\alpha f(z) = \sum_{n, m \geq 0} a_{n, m} \left(\int_{\mathbb{B}^d} f \bar{w}^m dv \right) z^n,$$

on $B_{r_0} \ni z$. This implies that $K_\alpha f$ is holomorphic at $z \in \mathbb{B}^d$.

(2) Similarly, it suffices to consider the case $f \in L^1$. Denote the holomorphic function $G_\alpha(z, w, u, v)$ on $B_{r'_0} \times B_{r_1} \times B_{r_1} \times B_{r'_0}$ by

$$G_\alpha(z, w, u, v) = \frac{1}{(1 - \langle z, \bar{w} \rangle)^{\frac{\alpha}{2}}} \frac{1}{(1 - \langle u, \bar{v} \rangle)^{\frac{\alpha}{2}}}.$$

By [22, Proposition 1.2.6] again, we know that $G_\alpha(z, w, u, v)$ has a global power series expansion, thus we can suppose that

$$G_\alpha(z, w, u, v) = \sum_{n, m, k, l \geq 0} a_{n, m, k, l} z^n w^m u^k v^l$$

on $B_{r'_0} \times B_{r_1} \times B_{r_1} \times B_{r'_0}$. Since

$$\overline{B_{r_0}} \times \overline{\mathbb{B}^d} \times \overline{\mathbb{B}^d} \times \overline{B_{r_0}} \subset B_{r'_0} \times B_{r_1} \times B_{r_1} \times B_{r'_0},$$

it follows from the dominated convergence theorem and the maximum modulus principle of analytic function that

$$\begin{aligned} F_\alpha(u, v) &\stackrel{\text{def}}{=} \int_{\mathbb{B}^d} f(w) G_\alpha(u, \bar{w}, w, v) dv(w) \\ &= \sum_{n, m, k, l \geq 0} a_{n, m, k, l} \left(\int_{\mathbb{B}^d} f w^k \bar{w}^m dv \right) u^n v^l, \end{aligned}$$

is holomorphic on $B_{r_0} \times B_{r_0}$. Then

$$K_\alpha^+ f(z) = \int_{\mathbb{B}^d} \frac{f(w) dv(w)}{|1 - \langle z, w \rangle|^\alpha} = \int_{\mathbb{B}^d} f(w) G_\alpha(z, \bar{w}, w, \bar{z}) dv(w) = F_\alpha(z, \bar{z}) \quad (2.2)$$

on $B_{r_0} \ni z$. It shows that $K_\alpha^+ f$ is smooth at $z \in \mathbb{B}^d$. \square

Remark 2.2. The formula (2.2) in fact shows that, for any $f \in L^1$, the function $K_\alpha^+ f$ admits a global power series expansion in the form

$$K_\alpha^+ f(z) = \sum_{n,m \geq 0} a_{n,m} z^n \bar{z}^m,$$

and $K_\alpha^+ f$ uniquely determines a holomorphic function on $\mathbb{B}^d \times \mathbb{B}^d$.

From Lemma 2.1, we know that the image of K_α^+ is continuous. The next lemma provides a result on the equicontinuity.

Lemma 2.3. Suppose $\alpha \in \mathbb{R}$, $1 \leq p \leq \infty$ and $K \subset \mathbb{B}^d$ is a compact subset, then for any $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$|K_\alpha^+ f(z_1) - K_\alpha^+ f(z_2)| \leq \|f\|_p \cdot \varepsilon, \quad \forall f \in L^p(\mathbb{B}^d),$$

whenever $z_1, z_2 \in K$ and $|z_1 - z_2| < \delta$.

Proof. Let B_r be the ball with radius r defined in Lemma 2.1. Since $K \subset \mathbb{B}^d$ is a compact subset, there exists a $0 < r < 1$ such that $K \subset B_r \subset \overline{B_r} \subset \mathbb{B}^d$. Since the function $\frac{1}{|1 - \langle z, w \rangle|^\alpha}$ is uniformly continuous on the compact set $\overline{B_r} \times \overline{B_r}$, it yields that, for any $\varepsilon > 0$, there exists a $\delta > 0$, such that

$$\left| \frac{1}{|1 - \langle z_1, w \rangle|^\alpha} - \frac{1}{|1 - \langle z_2, w \rangle|^\alpha} \right| \leq \varepsilon,$$

whenever $z_1, z_2 \in K$ and $|z_1 - z_2| < \delta$. Thus, for any $f \in L^p(\mathbb{B}^d)$, we have

$$|K_\alpha^+ f(z_1) - K_\alpha^+ f(z_2)| \leq \int_{\mathbb{B}^d} |f| \left| \frac{1}{|1 - \langle z_1, w \rangle|^\alpha} - \frac{1}{|1 - \langle z_2, w \rangle|^\alpha} \right| dv(w) \leq \|f\|_p \cdot \varepsilon$$

whenever $z_1, z_2 \in K$ and $|z_1 - z_2| < \delta$. It completes the proof. \square

A subset in a Banach space is called precompact if the closure of the subset in the norm topology is compact. Obviously, an operator between two Banach spaces is compact if and only if the operator maps every bounded set to precompact one. Suppose $\Omega \subset \mathbb{C}^d$ is an arbitrary bounded domain. The following lemma provides a criteria of precompactness in $L^p(\Omega)$ with $1 \leq p < \infty$, see [1, Theorem 2.33] for more details.

Lemma 2.4. [1] Let $1 \leq p < \infty$ and $K \subset L^p(\Omega)$. Suppose there exists a sequence $\{\Omega_j\}$ of subdomains of Ω having the following properties:

- (1) $\Omega_j \subset \Omega_{j+1}$.
- (2) The set of restrictions to Ω_j of the functions in K is precompact in $L^p(\Omega_j)$ for each j .
- (3) For every $\varepsilon > 0$ there exists a j such that

$$\int_{\Omega - \Omega_j} |f|^p dv < \varepsilon, \quad \forall f \in K.$$

Then K is precompact in $L^p(\Omega)$.

The Forelli-Rudin asymptotic estimate of integral for Bergman kernel on the unit ball \mathbb{B}^d is a fundamental result in function spaces and operator theory, see [22, Propostion 1.4.10] or [27, Theorem 1.12] for more details. In the rest of this article, we will frequently use this estimate, for the sake of convenience, we state as follows.

Lemma 2.5. (Forelli-Rudin) Suppose c is real and $t > -1$. Then the integral

$$J_{c,t}(z) = \int_{\mathbb{B}^d} \frac{(1 - |w|^2)^t dv(w)}{|1 - \langle z, w \rangle|^{c+t}}, \quad z \in \mathbb{B}^d,$$

has the following asymptotic properties.

- (1) If $c < d + 1$, then $J_{c,t}$ is bounded on \mathbb{B}^d .
- (2) If $c = d + 1$, then

$$J_{c,t}(z) \sim -\log(1 - |z|^2), \quad |z| \rightarrow 1^-.$$

- (3) If $c > d + 1$, then

$$J_{c,t}(z) \sim (1 - |z|^2)^{d+1-c}, \quad |z| \rightarrow 1^-.$$

The notation $A(z) \sim B(z)$ means that the ratio $\frac{A(z)}{B(z)}$ has a positive finite limit as $|z| \rightarrow 1^-$.

To complete the proof of Theorem 1, we also need a result on the interpolation of compact operators. The following lemma is first proved by A. Krasnoselskii [12], see also [6] or [13, Theorem 3.10].

Lemma 2.6. [6, 12, 13] Suppose that $1 \leq p_1, p_2, q_1, q_2 \leq \infty$ and $q_2 \neq \infty$. If a linear operator T such that $T : L^{p_1} \rightarrow L^{q_1}$ is bounded and $T : L^{p_2} \rightarrow L^{q_2}$ is compact, then $T : L^p \rightarrow L^q$ is compact, if there exists a $\theta \in (0, 1)$ such that

$$\frac{1}{p} = \frac{\theta}{p_1} + \frac{1-\theta}{p_2}, \quad \frac{1}{q} = \frac{\theta}{q_1} + \frac{1-\theta}{q_2}.$$

Proof of Theorem 1. The equivalence of (1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (5) is the main result of [9, Theorem 1], so it is enough to prove (4) \Leftrightarrow (5). Note that the equivalence of (3) and (5), it implies that (4) \Rightarrow (5), since compact operators are all bounded between two Banach spaces. Now we prove (5) \Rightarrow (4), which means we need to show K_α^+ is L^p - L^q compact if K_α^+ is L^p - L^q bounded. It is easy to see that $1 \leq q < p \leq \infty$ under the assumption in (5). By Lemma 2.6, the proof will be completed once we prove the following two conclusions:

- (a) If $p = \infty, q < \frac{1}{\alpha - (d+1)}$, then $K_\alpha^+ : L^p \rightarrow L^q$ is compact.
- (b) If $p > \frac{1}{(d+2) - \alpha}, q = 1$, then $K_\alpha^+ : L^p \rightarrow L^q$ is compact.

Observe the operator K_α^+ is adjoint by Fubini's theorem, combing this with the well known fact that an operator between two Banach spaces is compact if and only if its adjoint is compact, we conclude that conclusions (a) and (b) are in fact equivalent. Consequently, the proof is completed if conclusion (a) is proved. Now we turn to prove the conclusion (a). Suppose $\{f_j\}$ is an arbitrary bounded sequence in L^∞ , without loss of generality, we can suppose that

$$\|f_j\|_\infty \leq 1, \quad j = 1, 2, \dots \quad (2.3)$$

Denote the bounded domain B'_j by $B'_j = \{z \in \mathbb{C}^d : |z| < 1 - \frac{1}{j}\}, j = 1, 2, \dots$. Clearly, B'_j is compactly contained in \mathbb{B}^d and $B'_j \subset B'_{j+1} \subset \mathbb{B}^d$, for every j . We first prove that the set of restrictions to B'_j of the functions in $\{K_\alpha^+ f_n\}$ is precompact in $L^p(B'_j)$ for each j . In view of Lemma 2.1, we know the functions in $\{K_\alpha^+ f_n\}$ are all continuous on \mathbb{B}^d and uniformly continuous on every $\overline{B'_j}$. Combing with the fact that the embedding $C(\overline{B'_j}) \subset L^p(B'_j)$ is continuous for every j , it is enough to prove $\{(K_\alpha^+ f_n)|_{B'_j}\}$ is precompact in $C(\overline{B'_j})$ for every j . Note that (2.3) and the definition of B'_j , we have that

$$\begin{aligned} \|(K_\alpha^+ f_n)|_{B'_j}\|_\infty &= \sup_{z \in \overline{B'_j}} |K_\alpha^+ f_n| = \sup_{z \in \overline{B'_j}} \left| \int_{\mathbb{B}^d} \frac{f_n(w) dv(w)}{|1 - \langle z, w \rangle|^\alpha} \right| \\ &\leq \|f_n\|_\infty \sup_{z \in \overline{B'_j}} \int_{\mathbb{B}^d} \frac{dv(w)}{|1 - \langle z, w \rangle|^\alpha} \\ &\leq j^\alpha \|f_n\|_\infty \\ &\leq j^\alpha, \end{aligned} \quad (2.4)$$

for every j . The estimate (2.4) implies that $\{(K_\alpha^+ f_n)|_{B'_j}\}$ are bounded in $C(\overline{B'_j})$ for every j . It follows from Lemma 2.3 that $\{(K_\alpha^+ f_n)|_{B'_j}\}$ is equicontinuous on $\overline{B'_j}$ for every j . Then Arzelà-Ascoli theorem implies that $\{(K_\alpha^+ f_n)|_{B'_j}\}$ is precompact in $C(\overline{B'_j})$ for every j . By the well known fact that, for $t \in \mathbb{R}$, $(1 - |z|^2)^t \in L^1(\mathbb{B}^d)$ if and only if $t > -1$, we obtain that, for any fixed $t > -1$ and for any $\varepsilon > 0$, there exists a $J > 0$ satisfying

$$\int_{\mathbb{B}^d - B'_j} (1 - |z|^2)^t dv < \varepsilon, \quad \forall j > J, \quad (2.5)$$

since the absolute continuity of the integral and $\lim_{j \rightarrow \infty} v(\mathbb{B}^d - B'_j) = 0$. From the assumption in (a) and $d + 1 < \alpha < d + 2$, we get that $0 < q(\alpha - (d + 1)) < 1$, then by (2.5) we obtain that for any $\varepsilon > 0$, there exists a $J > 0$ such that

$$\int_{\mathbb{B}^d - B'_j} (1 - |z|^2)^{-q(\alpha - (d + 1))} dv < \varepsilon, \quad \forall j > J. \quad (2.6)$$

Combing (2.6) with Lemma 2.5, there exists a positive constant C such that, for any $\varepsilon > 0$, there exists a $J > 0$ satisfying

$$\begin{aligned} \int_{\mathbb{B}^d - B'_j} |K_\alpha^+ f_n(z)|^q dv(z) &= \int_{\mathbb{B}^d - B'_j} \left| \int_{\mathbb{B}^d} \frac{f_n(w) dv(w)}{|1 - \langle z, w \rangle|^\alpha} \right|^q dv(z) \\ &\leq \|f_n\|_\infty^q \int_{\mathbb{B}^d - B'_j} \left| \int_{\mathbb{B}^d} \frac{dv(w)}{|1 - \langle z, w \rangle|^\alpha} \right|^q dv(z) \\ &\leq C \|f_n\|_\infty^q \int_{\mathbb{B}^d - B'_j} (1 - |z|^2)^{-q(\alpha - (d + 1))} dv \\ &\leq C\varepsilon, \end{aligned}$$

for any $j > J$ and $n = 1, 2, \dots$. Thus $\{K_\alpha^+ f_n\}$ is precompact in L^q by Lemma 2.4, it completes the proof. \square

Remark 2.7. Theorem 1 indicates that Bergman operators K_α, K_α^+ from L^p into L^q are compact if and only if they are bounded when $1 \leq q < p \leq \infty$. In fact, this is quite expected, which can be seen from Ando's theorem for a bounded integral operator from L^p into L^q with $1 \leq q < p \leq \infty$, see [3] or [13, Theorem 5.5, 5.13]. Nevertheless, we give a direct proof here.

3. THE CASE OF $\alpha \leq d + 1$

This section is mainly devoted to prove Theorem 2, which characterizes the L^p - L^q compactness of K_α^+ for $0 < \alpha \leq d + 1$. We first recall some results on hypergeometric function theory. For complex numbers α, β, γ and complex variable z , we use the classical notation ${}_2F_1(\alpha, \beta; \gamma; z)$ to denote

$${}_2F_1(\alpha, \beta; \gamma; z) = \sum_{j=0}^{\infty} \frac{(\alpha)_j (\beta)_j}{j! (\gamma)_j} z^j,$$

with $\gamma \neq 0, -1, -2, \dots$, where $(\alpha)_j = \prod_{k=0}^{j-1} (\alpha + k)$ is the Pochhammer for any complex number α . The following lemma calculates the exact value of the hypergeometric function at the point $z = 1$.

Lemma 3.1. [10, Section 2.8] If $Re(\gamma - \alpha - \beta) > 0$, then

$${}_2F_1(\alpha, \beta; \gamma; 1) = \frac{\Gamma(\gamma)\Gamma(\gamma - \alpha - \beta)}{\Gamma(\gamma - \alpha)\Gamma(\gamma - \beta)},$$

where Γ is the usual Euler Gamma function.

Recall the definition of the function $J_{\beta, \gamma}$ in Lemma 2.5. The following lemma is not only a restatement of Lemma 2.5, but also establishes a connection between the integration induced by Bergman kernel and the hypergeometric function. A function is called finite if it has finite value at every point of its domain.

Lemma 3.2. (1)[22] Suppose $\beta \in \mathbb{R}$ and $\gamma > -1$, then

$$J_{\beta-\gamma, \gamma}(z) = \frac{\Gamma(1+d)\Gamma(1+\gamma)}{\Gamma(1+d+\gamma)} {}_2F_1\left(\frac{\beta}{2}, \frac{\beta}{2}; 1+d+\gamma; |z|^2\right), \quad (3.1)$$

for $z \in \mathbb{B}^d$.

(2) $J_{\beta-\gamma, \gamma}$ is finite on the closed ball $\overline{\mathbb{B}^d}$ if and only if $\beta < d + 1 + \gamma$. Moreover, in this case, the identity (3.1) actually holds on the closed ball $\overline{\mathbb{B}^d}$.

Proof. (1) In this case, the identity (3.1) is a restatement of Lemma 2.5, we refer the reader to [22, Proposition 1.4.10] for more details.

(2) It suffices to prove $J_{\beta-\gamma, \gamma}$ is finite on the unit sphere $\mathbb{S}^d = \{z \in \mathbb{C}^d : |z| = 1\}$ if and only if $\beta < d + 1 + \gamma$. Due to the unitary invariance of the Lebesgue measure, we know that

$$J_{\beta-\gamma, \gamma}(\eta) = \int_{\mathbb{B}^d} \frac{(1 - |w|^2)^\gamma dv(w)}{|1 - \langle \eta, w \rangle|^\beta} = \int_{\mathbb{B}^d} \frac{(1 - |w|^2)^\gamma dv(w)}{|1 - w_1|^\beta},$$

for any $\eta \in \mathbb{S}^d$. Then [26, Lemma 1.8,1.9] imply that

$$\begin{aligned}
J_{\beta-\gamma,\gamma}(\eta) &= 2d \int_0^1 r^{2d-1} (1-r^2)^\gamma dr \int_{\mathbb{S}^d} \frac{1}{|1-r\xi_1|^\beta} d\sigma(\xi) \\
&= 2d \int_0^1 r^{2d-1} (1-r^2)^\gamma (d-1) \int_{\mathbb{B}^1} \frac{(1-|w|^2)^{d-2}}{|1-rw|^\beta} dv_1(w) dr \\
&= 2d \int_0^1 r^{2d-1} (1-r^2)^\gamma {}_2F_1\left(\frac{\beta}{2}, \frac{\beta}{2}; d; r^2\right) dr \\
&= 2d \sum_{j=0}^{\infty} \frac{(\frac{\beta}{2})_j (\frac{\beta}{2})_j}{j!(d)_j} \int_0^1 r^{2j+2d-1} (1-r^2)^\gamma dr \\
&= \frac{\Gamma(d+1)\Gamma(\gamma+1)}{\Gamma(d+\gamma+1)} \sum_{j=0}^{\infty} \frac{(\frac{\beta}{2})_j (\frac{\beta}{2})_j}{j!(d+\gamma+1)_j}
\end{aligned} \tag{3.2}$$

for any $\eta \in \mathbb{S}^d$, where $d\sigma$ and dv_1 are normalized Lebesgue measures on \mathbb{S}^d and \mathbb{B}^1 , respectively. Note that $\Gamma(s+j) = (s)_j \Gamma(s)$ for all $s \in \mathbb{C}$ except the nonpositive integers. Then Stirling's formula and (3.2) yield that $J_{\beta-\gamma,\gamma}$ is finite on \mathbb{S}^d if and only if $\beta < d+1+\gamma$. From Lemma 3.1 and (3.2), we know that, if $\beta < d+1+\gamma$, then

$$\begin{aligned}
J_{\beta-\gamma,\gamma}(\eta) &= \frac{\Gamma(d+1)\Gamma(\gamma+1)}{\Gamma(d+\gamma+1)} \sum_{j=0}^{\infty} \frac{(\frac{\beta}{2})_j (\frac{\beta}{2})_j}{j!(d+\gamma+1)_j} \\
&= \frac{\Gamma(d+1)\Gamma(\gamma+1)}{\Gamma(d+\gamma+1)} {}_2F_1\left(\frac{\beta}{2}, \frac{\beta}{2}; d+\gamma+1; 1\right) \\
&= \frac{\Gamma(d+1)\Gamma(\gamma+1)\Gamma(d+\gamma+1-\beta)}{\Gamma^2(d+\gamma+1-\frac{\beta}{2})}
\end{aligned} \tag{3.3}$$

for any $\eta \in \mathbb{S}^d$, which means that the identity (3.1) also holds on the closed ball \mathbb{S}^d . It completes the proof. \square

From Lemma 3.2, we know that $J_{\beta-\gamma,\gamma}$ is a radial function on \mathbb{B}^d . Moreover, when $\beta < d+1+\gamma$, $J_{\beta-\gamma,\gamma}$ is increasing on $\overline{\mathbb{B}^d}$ in the following sense,

$$J_{\beta-\gamma,\gamma}(z_1) < J_{\beta-\gamma,\gamma}(z_2), \tag{3.4}$$

whenever $|z_1| < |z_2| \leq 1$, since all Taylor coefficients of the hypergeometric function in (3.1) are positive. Now, we introduce the following auxiliary function I_α for every $\alpha < d+1$. The function $I_\alpha(r, z)$ on $[0, 1) \times \overline{\mathbb{B}^d}$ is denoted by

$$I_\alpha(r, z) = \int_{r \leq |w| < 1} \frac{1}{|1-\langle z, w \rangle|^\alpha} dv(w).$$

Since $\alpha < d + 1$, it follows by (3.3) and (3.4) that

$$\begin{aligned} I_\alpha(r, z) &\leq J_{\alpha,0}(z) \\ &\leq \sum_{j=0}^{\infty} \left(\frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(\frac{\alpha}{2})} \right)^2 \frac{\Gamma(d+1)}{\Gamma(j+1)\Gamma(j+d+1)} \\ &= \frac{\Gamma(d+1)\Gamma(d+1-\alpha)}{\Gamma^2(d+1-\frac{\alpha}{2})}, \end{aligned} \quad (3.5)$$

for any $(r, z) \in [0, 1) \times \overline{\mathbb{B}^d}$, which means that I_α is finite on $[0, 1) \times \overline{\mathbb{B}^d}$. Moreover, I_α is increasing on $[0, 1) \times \overline{\mathbb{B}^d}$ in the following sense.

Lemma 3.3. Suppose $r \in [0, 1)$, then

$$I_\alpha(r, z_1) \leq I_\alpha(r, z_2),$$

whenever $z_1, z_2 \in \overline{\mathbb{B}^d}$ and $|z_1| \leq |z_2|$.

Proof. From (3.5), we know that I_α is finite on $[0, 1) \times \overline{\mathbb{B}^d}$. Now we calculate its exact value. It follows from [26, Lemma 1.8, 1.11] and the unitary invariance of the Lebesgue measure that

$$\begin{aligned} I_\alpha(r, z) &= \int_{r \leq |w| < 1} \frac{1}{|1 - |z||w_1|^\alpha} dv(w) \\ &= \int_{r \leq |w| < 1} \sum_{j=0}^{\infty} \left(\frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(\frac{\alpha}{2})\Gamma(j+1)} \right)^2 |z|^{2j} |w_1^j|^2 dv(w) \\ &= \sum_{j=0}^{\infty} \left(\frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(\frac{\alpha}{2})\Gamma(j+1)} \right)^2 |z|^{2j} \cdot 2d \int_r^1 t^{2d+2j-1} dt \int_{\mathbb{S}^d} |\xi_1^j|^2 d\sigma(\xi) \\ &= \sum_{j=0}^{\infty} \left(\frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(\frac{\alpha}{2})} \right)^2 \frac{\Gamma(d+1)(1-r^{2(j+d)})}{\Gamma(j+1)\Gamma(j+d+1)} |z|^{2j} \end{aligned} \quad (3.6)$$

for any $z \in \overline{\mathbb{B}^d}$. It leads to the desired result since all coefficients of the power series expansion about $|z|$ in (3.6) are positive. \square

Proposition 3.4. If $0 < \alpha < d+1$, then $K_\alpha^+ : L^\infty \rightarrow L^q$ is compact for any $1 \leq q \leq \infty$.

Proof. It suffices to prove that $K_\alpha : L^\infty \rightarrow L^\infty$ is compact. It is clear that $K_\alpha^+ f$ is continuous on the open ball \mathbb{B}^d for any $f \in L^\infty$ by Lemma 2.1. From Lemma 3.2, (3.3) and (3.4), we obtain that $K_\alpha^+ f(\eta)$ exists and

$$|K_\alpha^+ f(\eta)| \leq \|f\|_\infty \frac{\Gamma(d+1)\Gamma(d+1-\alpha)}{\Gamma^2(d+1-\frac{\alpha}{2})}$$

for any $\eta \in \mathbb{S}^d$. We now turn to prove that $K_\alpha^+ f$ is continuous on $\overline{\mathbb{B}^d}$. It suffices to prove that $K_\alpha^+ f$ on \mathbb{S}^d . Thus we need only to prove that, for any $\eta \in \mathbb{S}^d$ and for any point sequence $\{z_n\}$ in \mathbb{B}^d satisfying $z_n \rightarrow \eta$, then $K_\alpha^+ f(z_n) \rightarrow K_\alpha^+ f(\eta)$ as $n \rightarrow \infty$. By Lemma 3.2 and Lemma 3.4, we have

$$|K_\alpha^+ f(z)| \leq \|f\|_\infty J_{\alpha,0}(z) \leq \|f\|_\infty J_{\alpha,0}(\eta) = \|f\|_\infty \frac{\Gamma(d+1)\Gamma(d+1-\alpha)}{\Gamma^2(d+1-\frac{\alpha}{2})}, \quad (3.7)$$

for any $z \in \mathbb{B}^d$. The absolute continuity of the integral implies that, for any $\varepsilon > 0$, there exists a $0 < \delta < 1$, satisfying

$$\int_F \frac{dv(w)}{|1 - \langle \eta, w \rangle|^\alpha} \leq \frac{\varepsilon}{4}, \quad (3.8)$$

whenever $v(F) < \delta$. Denote $F_\delta = \{z \in \mathbb{B}^d : \sqrt[d]{1 - \frac{\delta}{2}} < |z| < 1\}$. Note that $v(F_\delta) = \frac{\delta}{2} < \delta$ and

$$\frac{1}{|1 - \langle z_n, w \rangle|^\alpha} \rightarrow \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \text{ uniformly on } \mathbb{B}^d \setminus F_\delta, \text{ as } n \rightarrow \infty.$$

Then there exists a $N > 0$ such that, for any $n > N$,

$$\int_{\mathbb{B}^d \setminus F_\delta} \left| \frac{1}{|1 - \langle z_n, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) \leq \frac{\varepsilon}{2}.$$

Combing this with Lemma 3.3 and (3.8), we conclude that, for any $n > N$,

$$\begin{aligned} |K_\alpha^+ f(z_n) - K_\alpha^+ f(\eta)| &\leq \|f\|_\infty \int_{\mathbb{B}^d \setminus F_\delta} \left| \frac{1}{|1 - \langle z_n, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) \\ &\quad + \|f\|_\infty \int_{F_\delta} \left| \frac{1}{|1 - \langle z_n, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) \\ &\leq \|f\|_\infty \int_{\mathbb{B}^d \setminus F_\delta} \left| \frac{1}{|1 - \langle z_n, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) \quad (3.9) \\ &\quad + 2\|f\|_\infty \int_{F_\delta} \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} dv(w) \\ &\leq \|f\|_\infty \frac{\varepsilon}{2} + 2\|f\|_\infty \frac{\varepsilon}{4} \\ &= \varepsilon \|f\|_\infty. \end{aligned}$$

It completes the proof that $K_\alpha^+ f$ is continuous on the closed ball $\overline{\mathbb{B}^d}$ for any $f \in L^\infty$. Now we prove that, for any bounded sequence in L^∞ , there exists a subsequence satisfying that its image under K_α^+ is convergent in L^∞ . Suppose that $\{f_n\}$ is a bounded sequence in L^∞ , then we have that $\{K_\alpha^+ f_n\}$ is in $C(\overline{\mathbb{B}^d})$ and $\{K_\alpha^+ f_n\}$ is uniformly bounded by (3.7). Now we prove that $\{K_\alpha^+ f_n\}$ is also equicontinuous. From (3.9), we know that

$$\lim_{\mathbb{B}^d \ni z \rightarrow \eta} \int_{\mathbb{B}^d} \left| \frac{1}{|1 - \langle z, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) = 0, \quad (3.10)$$

for arbitrary fixed $\eta \in \mathbb{S}^d$. Combing (3.10) with the unitary invariance of Lebesgue measure and the symmetry of the unit ball, we have that, for any $\varepsilon > 0$, there exists a $0 < \delta' < 1$, satisfying that

$$\int_{\mathbb{B}^d} \left| \frac{1}{|1 - \langle z, w \rangle|^\alpha} - \frac{1}{|1 - \langle \eta, w \rangle|^\alpha} \right| dv(w) \leq \frac{\varepsilon}{2} \quad (3.11)$$

whenever $z \in \mathbb{B}^d, \eta \in \mathbb{S}^d$ and $|z - \eta| < \delta'$. Denote $B''_{1-\frac{\delta'}{2}} = \{z \in \mathbb{C}^d : |z| \leq 1 - \frac{\delta'}{2}\}$ and $C_{\frac{\delta'}{2}} = \{z \in \mathbb{C}^d : 1 - \frac{\delta'}{2} < |z| \leq 1\}$. Then the closed ball $\overline{\mathbb{B}^d}$ has the following decomposition,

$$\overline{\mathbb{B}^d} = B''_{1-\frac{\delta'}{2}} \cup C_{\frac{\delta'}{2}} \text{ and } B''_{1-\frac{\delta'}{2}} \cap C_{\frac{\delta'}{2}} = \emptyset.$$

Since the function $\frac{1}{|1-\langle z, w \rangle|^\alpha}$ is uniformly continuous on the compact set $B''_{1-\frac{\delta'}{2}} \times \overline{\mathbb{B}^d}$, there exists a $0 < \delta'' < 1$ such that

$$\left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| \leq \varepsilon, \quad (3.12)$$

whenever $(z_1, w), (z_2, w) \in B''_{1-\frac{\delta'}{2}} \times \overline{\mathbb{B}^d}$ and $|z_1 - z_2| < \delta''$. Take $\delta''' = \min\{\frac{\delta'}{2}, \delta''\}$. Now we prove that, for any $z_1, z_2 \in \overline{\mathbb{B}^d}$ such that $|z_1 - z_2| < \delta'''$, then

$$\int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| dv(w) \leq \varepsilon. \quad (3.13)$$

In fact, there are two cases need to be considered. The first case is $z_1 \in C_{\frac{\delta'}{2}}$ or $z_2 \in C_{\frac{\delta'}{2}}$. Without loss of generality, we can assume that $z_1 \in C_{\frac{\delta'}{2}}$, then there exists an $\eta \in \mathbb{S}^d$ satisfying that $|z_1 - \eta| < \delta''' \leq \frac{\delta'}{2}$. Thus, obviously, the triangle inequality implies that $|z_2 - \eta| \leq |z_2 - z_1| + |z_1 - \eta| < \delta'$. This together with (3.11) implies that

$$\begin{aligned} & \int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| dv(w) \\ & \leq \int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle \eta, w \rangle|^\alpha} \right| dv(w) \\ & \quad + \int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle \eta, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| dv(w) \\ & \leq \varepsilon \end{aligned}$$

The second case is $z_1, z_2 \in B''_{1-\frac{\delta'}{2}}$. From (3.12), it implies that

$$\int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| dv(w) \leq \varepsilon \int_{\mathbb{B}^d} dv = \varepsilon.$$

It proves (3.13). Combing (3.13) with

$$|K_\alpha^+ f_n(z_1) - K_\alpha^+ f_n(z_2)| \leq \|f_n\|_\infty \int_{\mathbb{B}^d} \left| \frac{1}{|1-\langle z_1, w \rangle|^\alpha} - \frac{1}{|1-\langle z_2, w \rangle|^\alpha} \right| dv(w)$$

follows that $\{K_\alpha^+ f_n\}$ is equicontinuous. Then Arzelà-Ascoli theorem implies that $\{K_\alpha^+ f_n\}$ has a convergency subsequence in the supremum norm (or is precompact). That finishes the proof. \square

Corollary 3.5. If $0 < \alpha < d + 1$, then $K_\alpha^+ : L^p \rightarrow L^1$ is compact for any $1 \leq p \leq \infty$.

Proof. It follows from Lemma 3.4 and the fact that K_α^+ is adjoint. \square

Now, recall the definition of the fractional radial differential operator $R^{s,t}$ on $\text{Hol}(\mathbb{B}^d)$. For any two real parameters s and t with the property that neither $d+s$ nor $d+s+t$ is a negative integer, the invertible operator $R^{s,t}$ is given by

$$R^{s,t}f(z) = \sum_{n=0}^{\infty} \frac{\Gamma(d+1+s)\Gamma(d+1+n+s+t)}{\Gamma(d+1+s+t)\Gamma(d+1+n+s)} f_n(z),$$

for any $f = \sum_{n=0}^{\infty} f_n \in \text{Hol}(\mathbb{B}^d)$ with homogeneous expansion. From [22, Proposition 1.2.6], we know every holomorphic function $f \in \text{Hol}(\mathbb{B}^d)$ has a global power series expansion, thus the definition is well-defined on $\text{Hol}(\mathbb{B}^d)$. In fact, it can be checked by the direct calculation that the invertible operator of $R^{s,t}$ is just $R^{s+t,-t}$. Be careful that the invertible operator here is unnecessarily continuous. Recall that $A^p = \text{Hol}(\mathbb{B}^d) \cap L^p$ is the p -integrable Bergman space on \mathbb{B}^d for $1 \leq p \leq \infty$.

Lemma 3.6. Suppose $\alpha \in \mathbb{R}$ satisfying α is not a nonpositive integer and $1 \leq p \leq \infty$, then the following holds on A^p

$$K_\alpha = R^{0,\alpha-d-1}.$$

Proof. Since every holomorphic function $f \in \text{Hol}(\mathbb{B}^d)$ has a global power series expansion, we can suppose $f = \sum_{n=0}^{\infty} f_n \in A^p$ with the homogeneous expansion. Then the dominated convergence theorem and formula (1.21) in [26] imply that

$$\begin{aligned} K_\alpha f(z) &= \int_{\mathbb{B}^d} \frac{f(w)}{(1-\langle z, w \rangle)^\alpha} dv(w) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)}{\Gamma(n+1)\Gamma(\alpha)} \int_{\mathbb{B}^d} f(w) \langle z, w \rangle^n dv(w) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)}{\Gamma(n+1)\Gamma(\alpha)} 2d \int_0^1 r^{2d-1} dr \sum_{k=0}^{\infty} r^{k+n} \int_{\mathbb{S}^d} f_k(\xi) \langle z, \xi \rangle^n d\sigma(\xi) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)}{\Gamma(n+1)\Gamma(\alpha)} 2d \int_0^1 r^{2d-1} dr \int_{\mathbb{S}^d} f_n(r\xi) \langle z, r\xi \rangle^n d\sigma(\xi) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(n+\alpha)}{\Gamma(n+1)\Gamma(\alpha)} \int f_n(w) \langle z, w \rangle^n dv(w) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)} f_n(z), \end{aligned} \tag{3.14}$$

for any $z \in \mathbb{B}^d$. It leads to the desired result. \square

Lemma 3.7. [25, Proposition 5] Suppose s, t are real parameters such that neither $n+s$ nor $n+s+t$ is a negative integer. Then, for any nonnegative integer N ,

$$R^{s,t} \frac{1}{(1-\langle z, w \rangle)^{d+1+s+N}} = \frac{h(\langle z, w \rangle)}{(1-\langle z, w \rangle)^{d+1+s+N+t}},$$

where h is a certain one-variable polynomial of degree N . In particular, $h \equiv 1$ if $N = 0$.

Proposition 3.8. Suppose $0 < \alpha \leq d+1$ and $1 < p, q < \infty$, if $K_\alpha^+ : L^p \rightarrow L^q$ is compact, then $\frac{1}{q} > \frac{1}{p} + \frac{\alpha}{d+1} - 1$.

Proof. For every $z \in \mathbb{B}^d$, denote the holomorphic function H_z on \mathbb{B}^d by

$$H_z(w) = \frac{(1 - |z|^2)^{\frac{s}{q}}}{(1 - \langle w, z \rangle)^{\frac{s}{q} + \frac{d+1}{p} + \alpha - d - 1}}, \quad w \in \mathbb{B}^d, \quad (3.15)$$

where s is a positive parameter satisfying $\frac{s}{q} - (1 - \frac{1}{p})(d+1)$ is a large enough positive integer. Clearly H_z is bounded holomorphic on \mathbb{B}^d for each $z \in \mathbb{B}^d$. We first prove $|R^{\alpha-d-1, d+1-\alpha} H_z| \rightarrow 0$ weakly in L^p , as $|z| \rightarrow 1^-$. By [2, Section 8.3.3, Theorem 2], it suffices to prove that

- (1) $\sup_{z \in \mathbb{B}^d} \|R^{\alpha-d-1, d+1-\alpha} H_z\|_p < \infty$;
- (2) $\lim_{|z| \rightarrow 1^-} \int_{\mathbb{B}^d} |R^{\alpha-d-1, d+1-\alpha} H_z| dv = 0$.

It is immediate from Lemma 3.7 that

$$R^{\alpha-d-1, d+1-\alpha} H_z(w) = \frac{h_s(\langle w, z \rangle)(1 - |z|^2)^{\frac{s}{q}}}{(1 - \langle w, z \rangle)^{\frac{s}{q} + \frac{d+1}{p}}}, \quad w \in \mathbb{B}^d, \quad (3.16)$$

where h_s is a certain one-variable polynomial of degree $\frac{s}{q} - (1 - \frac{1}{p})(d+1)$. From Lemma 2.5, there exists a positive constant C such that

$$\int_{\mathbb{B}^d} \frac{|h_s(\langle w, z \rangle)| dv(w)}{|1 - \langle w, z \rangle|^{\frac{s}{q} + \frac{d+1}{p}}} \leq C(1 - |z|^2)^{(1-\frac{1}{p})(d+1) - \frac{s}{q}}, \quad z \in \mathbb{B}^d.$$

Then by (3.16), the following estimate holds,

$$\begin{aligned} \int_{\mathbb{B}^d} |R^{\alpha-d-1, d+1-\alpha} H_z| dv &\leq \int_{\mathbb{B}^d} \frac{|h_s(\langle w, z \rangle)|(1 - |z|^2)^{\frac{s}{q}} dv(w)}{|1 - \langle w, z \rangle|^{\frac{s}{q} + \frac{d+1}{p}}} \\ &\leq C(1 - |z|^2)^{(1-\frac{1}{p})(d+1)}. \end{aligned}$$

Thus, $\lim_{|z| \rightarrow 1^-} \int_{\mathbb{B}^d} |R^{\alpha-d-1, d+1-\alpha} H_z| dv = 0$, namely, condition (2) holds. Similarly, we can verify the condition (1) by Lemma 2.5. Thus we prove that $|R^{\alpha-d-1, d+1-\alpha} H_z| \rightarrow 0$ weakly in L^p , as $|z| \rightarrow 1^-$. Now, combing with the well-known fact that a compact operator maps a weakly convergent sequence into a strongly convergent one, we obtain

$$\lim_{|z| \rightarrow 1^-} \|K_\alpha^+(|R^{\alpha-d-1, d+1-\alpha} H_z|)\|_q = 0.$$

Note that $K_\alpha^+(|f|) \geq |K_\alpha(f)|$ for any $f \in L^1$, then

$$\lim_{|z| \rightarrow 1^-} \|K_\alpha(R^{\alpha-d-1, d+1-\alpha} H_z)\|_q = 0.$$

Since $R^{0, \alpha-d-1} R^{\alpha-d-1, d+1-\alpha} f = f$ for any bounded holomorphic function f on \mathbb{B}^d , it follows from Lemma 3.6 that

$$\lim_{|z| \rightarrow 1^-} \|H_z\|_q = \lim_{|z| \rightarrow 1^-} \|R^{0, \alpha-d-1} R^{\alpha-d-1, d+1-\alpha} H_z\|_q = \lim_{|z| \rightarrow 1^-} \|K_\alpha R^{\alpha-d-1, d+1-\alpha} H_z\|_q = 0,$$

and then

$$\lim_{|z| \rightarrow 1^-} \int_{\mathbb{B}^d} \frac{(1 - |z|^2)^s dv(w)}{|1 - \langle w, z \rangle|^{s + \frac{q(d+1)}{p} + q(\alpha-d-1)}} = \lim_{|z| \rightarrow 1^-} \|H_z\|_q^q = 0.$$

Since $s > 0$ is chosen large enough, by Lemma 2.5 again, we obtain that $\frac{1}{q} > \frac{1}{p} + \frac{\alpha}{d+1} - 1$. \square

Remark 3.9. (1) We have proved $|R^{\alpha-d-1, d+1-\alpha} H_z| \rightarrow 0$ weakly in L^p , as $|z| \rightarrow 1^-$. Note that bounded operators map a weakly convergent sequence into a weakly convergent one. Analogously, we can prove that, if $K_\alpha^+ : L^p \rightarrow L^q$ is bonded as the same condition in Proposition 3.8, then

$$\sup_{z \in \mathbb{B}^d} \int_{\mathbb{B}^d} \frac{(1 - |z|^2)^s dv(w)}{|1 - \langle w, z \rangle|^{s + \frac{q(d+1)}{p} + q(\alpha-d-1)}} < \infty.$$

Then Lemman 2.5 implies that $\frac{1}{q} \geq \frac{1}{p} + \frac{\alpha}{d+1} - 1$. In fact, this gives a new proof of necessity part of [9, Lemma 4.3], which is a key step to characterize the L^p - L^q boundedness of K_α, K_α^+ .

(2) The construction of H_z is inspired by the characterization of vanishing Carleson measure for Bergman space on the unit ball, we refer the reader to [25, 26].

Now we turn to the proof of Theorem 2, we first describe the L^p - L^q compactness of K_{d+1}^+ . It is as analogous to the proof of Theorem 1. Thus we give a sketchy proof here.

Proposition 3.10. $K_{d+1}^+ : L^p \rightarrow L^q$ is compact if and only if $1 \leq q < p \leq \infty$.

Proof. By Lemma 2.6 and the adjointness of K_α^+ , it suffices to prove $K_{d+1}^+ : L^p \rightarrow L^1$ is compact for any $1 < p \leq \infty$. Note that [9, Theorem 3] shows $K_{d+1}^+ : L^\infty \rightarrow L^1$ is bounded, combing with Lemma 2.6 again, it suffices to prove $K_{d+1}^+ : L^p \rightarrow L^1$ is compact for any $d+1 < p < \infty$. Let $\{B'_j\}$ be as in Proof of Theorem 1, and $\{f_j\}$ be a bounded sequence in L^p . Then we can prove $\|(K_\alpha^+ f_n)|_{B'_j}\|_\infty \leq j^\alpha \|f_n\|_p$, combing with Lemma 2.1 and Arzelà-Ascoli theorem implies that $\{(K_\alpha^+ f_n)|_{B'_j}\}$ is precompact in $L^p(B'_j)$. By Lemma 2.5 and Hölder's inequality, it implies, there exists a constant $C > 0$ such that

$$\begin{aligned} \int_{\mathbb{B}^d - B'_j} |K_{d+1}^+ f_n(z)| dv(z) &\leq \|f_n\|_p \int_{\mathbb{B}^d - B'_j} \left| \int_{\mathbb{B}^d} \frac{dv(w)}{|1 - \langle z, w \rangle|^{\frac{p(d+1)}{p-1}}} \right|^{1-\frac{1}{p}} dv(z) \\ &\leq C \|f_n\|_p \int_{\mathbb{B}^d - B'_j} (1 - |z|^2)^{-\frac{d+1}{p}} dv(z). \end{aligned}$$

Thus we obtain that, if $d+1 < p < \infty$, then for any $\varepsilon > 0$, there exists a $J > 0$, such that

$$\int_{\mathbb{B}^d - B'_j} |K_{d+1}^+ f_n| dv \leq C\varepsilon,$$

whenever $j > J$. Therefore, $\{K_\alpha^+ f_n\}$ is precompact in $L^1(\mathbb{B}^d)$ by Lemma 2.4. \square

Proof of Theorem 2. Since the equivalence of (1) \Leftrightarrow (3) is the main result of [9, Theorem 3], it suffices to prove (2) \Leftrightarrow (3). There are two cases $\alpha = d+1$ and $0 < \alpha < d+1$ to be considered. Indeed, the case $\alpha = d+1$ has been proved in Proposition 3.10. Thus, it remains to deal with the case $0 < \alpha < d+1$. By Lemma 2.6 and Corollary 3.5, we conclude that $K_\alpha^+ : L^1 \rightarrow L^q$ is compact if and only if $q < \frac{d+1}{\alpha}$, and then $K_\alpha^+ : L^p \rightarrow L^\infty$ is compact if and only if $p > \frac{d+1}{d+1-\alpha}$. Then apply Lemma 2.6, Proposition 3.4 and Proposition 3.8 to obtain the equivalence of (2) and (3) when $0 < \alpha < d+1$. It completes the proof. \square

We have completely characterized that L^p - L^q compactness of the Bergman-type operators K_α, K_α^+ when $0 < \alpha < d + 2$ so far. However, when $\alpha \geq d + 2$, [9, Theorem 4] shows that there exist no $1 \leq p, q \leq \infty$ such that $K_\alpha^+ : L^p \rightarrow L^q$ is bounded.

To end this section, we describe the phenomenon for $\alpha \leq 0$. Recall that a bounded operator between two Banach spaces is called a finite rank operator if the range of the operator has finite dimension. Obviously, finite rank operators must be compact and the finite rank operators on a Hilbert space belong to every Schatten p -class with $0 < p < \infty$.

Proposition 3.11. Suppose $\alpha \leq 0$, then the followings hold.

- (1) $K_\alpha, K_\alpha^+ : L^p \rightarrow L^q$ are compact for any $1 \leq p, q \leq \infty$.
- (2) If α is a nonpositive integer, then $K_\alpha : L^p \rightarrow L^q$ is a finite rank operator for any $1 \leq p, q \leq \infty$.
- (3) If α is a nonpositive even integer, then $K_\alpha^+ : L^p \rightarrow L^q$ is a finite rank operator for any $1 \leq p, q \leq \infty$.

Proof. (1) It suffices to prove that $K_\alpha, K_\alpha^+ : L^1 \rightarrow L^\infty$ are compact when $\alpha \leq 0$. Suppose $\{f_j\}$ is an arbitrary bounded sequence in L^∞ , without loss of generality, we can suppose that

$$\|f_j\|_\infty \leq 1, \quad j = 1, 2, \dots$$

Then Lemma 2.1 implies that $\{K_\alpha f_j\}$ is continuous function sequences on \mathbb{B}^d . Note that the kernel function $\frac{1}{(1-\langle z, w \rangle)^\alpha}$ is uniformly continuous on the compact set $\overline{\mathbb{B}^d} \times \overline{\mathbb{B}^d}$ when $\alpha \leq 0$, then we obtain that $\{K_\alpha f_j\}$ is in fact continuous function sequence on $\overline{\mathbb{B}^d}$. Since $\alpha \leq 0$, it follows that

$$\begin{aligned} \|K_\alpha f_n\|_\infty &= \sup_{z \in \overline{\mathbb{B}^d}} \left| \int_{\mathbb{B}^d} \frac{f_n(w) dv(w)}{(1 - \langle z, w \rangle)^\alpha} \right| \\ &\leq 2^{-\alpha} \|f_n\|_1 \\ &\leq 2^{-\alpha}, \end{aligned}$$

Thus $\{K_\alpha f_j\}$ is a bounded subset in $C(\overline{\mathbb{B}^d})$. The uniform continuity of the function $\frac{1}{(1-\langle z, w \rangle)^\alpha}$ also yields that, for any $\varepsilon > 0$, there exists a $\delta > 0$, such that

$$\left| \frac{1}{(1 - \langle z_1, w \rangle)^\alpha} - \frac{1}{(1 - \langle z_2, w \rangle)^\alpha} \right| \leq \varepsilon,$$

whenever $z_1, z_2 \in \overline{\mathbb{B}^d}$ and $|z_1 - z_2| < \delta$. Then we have

$$\begin{aligned} |K_\alpha f_j(z_1) - K_\alpha f_j(z_2)| &\leq \int_{\mathbb{B}^d} |f_j| \left| \frac{1}{(1 - \langle z_1, w \rangle)^\alpha} - \frac{1}{(1 - \langle z_2, w \rangle)^\alpha} \right| dv(w) \\ &\leq \|f_j\|_1 \cdot \varepsilon \\ &\leq \varepsilon \end{aligned}$$

whenever $z_1, z_2 \in \overline{\mathbb{B}^d}$ and $|z_1 - z_2| < \delta$. That means $\{K_\alpha f_j\}$ is equicontinuous. Then Arzelà-Ascoli theorem implies that $\{K_\alpha f_j\}$ has a convergency subsequence in the supremum norm (or is precompact). That proves $K_\alpha : L^1 \rightarrow L^\infty$ is compact when $\alpha \leq 0$. Similarly, $K_\alpha^+ : L^1 \rightarrow L^\infty$ is compact when $\alpha \leq 0$.

(2) It suffices to prove that $K_\alpha : L^1 \rightarrow L^\infty$ is a finite rank operator when α a nonpositive integer. Suppose that $\alpha = -N$, where N is a nonnegative integer. Then the binomial theorem implies that

$$(1 - \langle z, w \rangle)^N = \sum_{|s| \leq N} a_{N,s} z^s \bar{w}^s,$$

where $s \geq 0$ is multi-index and $a_{N,s}$ is nonzero constant. Thus, for any $f \in L^1$, we have

$$K_{-N}f(z) = \int_{\mathbb{B}^d} f(w)(1 - \langle z, w \rangle)^N dv = \sum_{|s| \leq N} \left(a_{N,s} \int_{\mathbb{B}^d} f \bar{w}^s dv \right) z^s.$$

It implies that K_{-N} is a finite rank operator.

(3) It suffices to prove that $K_\alpha^+ : L^1 \rightarrow L^\infty$ is a finite rank operator when α a nonpositive even integer. Suppose that $\alpha = -2N$, where N is a nonnegative integer. Then

$$|1 - \langle z, w \rangle|^{2N} = (1 - \langle z, w \rangle)^N (1 - \langle w, z \rangle)^N = \sum_{|s|, |l| \leq N} a_{N,s} a_{N,l} z^s \bar{z}^l w^l \bar{w}^s.$$

Thus, for any $f \in L^1$, we have

$$K_{-2N}f(z) = \int_{\mathbb{B}^d} f |1 - \langle z, w \rangle|^{2N} dv = \sum_{|s|, |l| \leq N} \left(a_{N,s} a_{N,l} \int_{\mathbb{B}^d} f w^l \bar{w}^s dv \right) z^s \bar{z}^l.$$

It implies that K_{-2N} is a finite rank operator. \square

Remark 3.12. K_α^+ will not be a finite rank operator when α is a nonpositive odd integer.

Thus we have the following corollary.

Corollary 3.13. For $\alpha \in \mathbb{R}$, then the following statements are equivalent:

- (1) $\alpha < d + 2$;
- (2) there exist $1 \leq p, q \leq \infty$ such that $K_\alpha : L^p \rightarrow L^q$ is bounded;
- (3) there exist $1 \leq p, q \leq \infty$ such that $K_\alpha^+ : L^p \rightarrow L^q$ is bounded;
- (4) there exist $1 \leq p, q \leq \infty$ such that $K_\alpha : L^p \rightarrow L^q$ is compact;
- (5) there exist $1 \leq p, q \leq \infty$ such that $K_\alpha^+ : L^p \rightarrow L^q$ is compact.

4. SCHATTEN CLASS AND MACAEV CLASS MEMBERSHIP

In the present section, we complete proofs of Theorem 3,4, which give the necessary and sufficient conditions that ensure the Bergman-type operator K_α belongs to Schatten classes and Macaev classes. The proof of Theorem 3 will be respectively given in cases $0 < \alpha < d + 1$ and $\alpha \leq 0$. Although Theorem 3 can be uniformly proved by the method of estimates of the spectrum, we prefer to prove the theorem in the case $0 < \alpha < d + 1$ by using the method of operator theory which is inspired by the characterization of Schatten class Toeplitz operators and Hankel operators in [19, 27, 28].

Denote the point spectrum (the collections of eigenvalues) of K_α on the Bergman space A^2 by $\sigma_{pt}(K_\alpha, A^2)$.

Lemma 4.1. Suppose $\alpha < d + 1$, then the followings hold.

- (1) If $\alpha \in \mathcal{S}_d$, then $\sigma_{pt}(K_\alpha, A^2) = \left\{ \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)} : n \in \mathbb{N} \right\}$.
- (2) If $-\alpha \in \mathbb{N}$, then $\sigma_{pt}(K_\alpha, A^2) = \{0\} \cup \left\{ \frac{(-1)^n \Gamma(1-\alpha)\Gamma(d+1)}{\Gamma(1-\alpha-n)\Gamma(n+d+1)} : 0 \leq n \leq -\alpha \right\}$.

Proof. (1) Denote $\mu_n = \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)}$, $n \in \mathbb{N}$. Due to Lemma 3.6, it suffices to show that $\sigma_{pt}(K_\alpha, A^2) \subset \{\mu_n : n \in \mathbb{N}\}$. Suppose $\mu \in \sigma_{pt}(K_\alpha, A^2)$, then there exists a nonzero $f \in A^2$ such that

$$K_\alpha f = \mu f. \quad (4.1)$$

It is easy to see that $\cup_{n \in \mathbb{N}} P_n$ is a orthonormal basis of A^2 , where P_n is the set of homogeneous polynomials defined by

$$P_n = \left\{ c_k z^k : \sum_{j=1}^d k_j = n, k_j \in \mathbb{N} \right\}, \quad n \in \mathbb{N},$$

and c_k is the normalized positive constant such that $\|c_k z^k\|_2 = 1$ for each k . Thus f has the following representation

$$f = \sum_{n=0}^{\infty} \sum_{e_{n,k} \in P_n} \langle f, e_{n,k} \rangle e_{n,k},$$

where $\langle \cdot, \cdot \rangle$ is the standard Hermitian inner product on A^2 . Combing this with Lemma 3.6 and (4.1), we conclude that

$$\sum_{n=0}^{\infty} (\mu - \mu_n) \sum_{e_{n,k} \in P_n} \langle f, e_{n,k} \rangle e_{n,k} = 0. \quad (4.2)$$

Since f is nonzero, there exists a e_{n_0, k_0} such that $\langle f, e_{n_0, k_0} \rangle \neq 0$. Then (4.2) implies that f has the form

$$f = \sum_{e_{n,k} \in P_{n_0}} \langle f, e_{n,k} \rangle e_{n,k}, \quad (4.3)$$

and $\mu = \mu_{n_0}$. It completes the proof of $\sigma_{pt}(K_\alpha, A^2) \subset \{\mu_n : n \in \mathbb{N}\}$.

(2) Suppose $f = \sum f_n \in A^2$ with the homogeneous expansion. Since $-\alpha$ is a nature number, it follows that

$$\begin{aligned} K_\alpha f(z) &= \int_{\mathbb{B}^d} \sum f_n(w) (1 - \langle z, w \rangle)^{-\alpha} dv(w) \\ &= \int_{\mathbb{B}^d} \sum_{n=0}^{\infty} f_n \sum_{k=0}^{-\alpha} \frac{(-\alpha)! (-1)^k}{(-\alpha - k)! k!} \langle z, w \rangle^k dv(w) \\ &= \sum_{n=0}^{-\alpha} \frac{(-\alpha)! (-1)^n}{(-\alpha - n)! n!} \int f_n(w) \langle z, w \rangle^n dv(w) \\ &= \sum_{n=0}^{-\alpha} \frac{(-1)^n \Gamma(1-\alpha) \Gamma(d+1)}{\Gamma(1-\alpha-n) \Gamma(n+d+1)} f_n(z) \end{aligned}$$

for any $z \in \mathbb{B}^d$. Then, the same argument as above shows that

$$\sigma_{pt}(K_\alpha, A^2) = \{0\} \cup \left\{ \frac{(-1)^n \Gamma(1-\alpha) \Gamma(d+1)}{\Gamma(1-\alpha-n) \Gamma(n+d+1)} : 0 \leq n \leq -\alpha \right\}.$$

□

Theorem 2, Lemma 2.1, Lemma 3.6 and the boundedness of embedding $A^2 \rightarrow L^2$, imply that K_α is a normal compact operator on the Bergman space A^2 when $\alpha < d+1$. In particular, K_α is normal compact on A^2 when $0 < \alpha < d+1$, thus we can apply the functional calculation to K_α . Set function $F_p(x) = x^p$ on $\mathbb{R}_{\geq 0}$ for any $p > 0$. Note that $F_p : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is bijective. We define $K_\alpha^p = F_p(K_\alpha)$, which is the functional calculation of $K_\alpha \in B(A^2)$ with respect to the function $F_p(x) = x^p$, where $B(A^2)$ is the collections of bounded operators on the Bergman space A^2 .

Lemma 4.2. Suppose $0 < \alpha < d+1$. Then for any $0 < p < \frac{d+1}{d+1-\alpha}$, there exists a positive constant C_p such that the following operator inequality

$$\frac{1}{C_p} K_{p\alpha-(p-1)(d+1)} \leq K_\alpha^p \leq C_p K_{p\alpha-(p-1)(d+1)}$$

holds on the Bergman space A^2 .

Proof. Since K_α is positive and compact on A^2 , it follows from [23, Theorem 1.9.2] that K_α admits the following canonical decomposition

$$K_\alpha f = \sum_{n=0}^{\infty} \lambda_n \langle f, e_n \rangle e_n,$$

where $\{\lambda_n\}$ is the sequence of nonzero eigenvalues (counting multiplicities) with decreasing order, $\{e_n\}$ is the corresponding orthonormal sequence of eigenvectors and $\langle \cdot, \cdot \rangle$ is the standard Hermitian inner product on A^2 . Combing with Lemma 4.1, we can further suppose

$$K_\alpha f = \sum_{n=0}^{\infty} \frac{\Gamma(d+1) \Gamma(\alpha+n)}{\Gamma(\alpha) \Gamma(d+1+n)} \sum_{e_{n,k} \in P_n} \langle f, e_{n,k} \rangle e_{n,k} \quad (4.4)$$

is the canonical decomposition of K_α . It implies from the functional calculation and (4.4) that

$$K_\alpha^p f = \sum_{n=0}^{\infty} \left(\frac{\Gamma(d+1) \Gamma(\alpha+n)}{\Gamma(\alpha) \Gamma(d+1+n)} \right)^p \sum_{e_{n,k} \in P_n} \langle f, e_{n,k} \rangle e_{n,k},$$

is the canonical decomposition of K_α^p . Note that the condition $0 < p < \frac{d+1}{d+1-\alpha}$ ensures $p\alpha - (p-1)(d+1) > 0$. By Stirling's formula, we conclude that

$$\left(\frac{\Gamma(d+1) \Gamma(\alpha+n)}{\Gamma(\alpha) \Gamma(d+1+n)} \right)^p \sim n^{p(\alpha-(d+1))} \sim \frac{\Gamma(d+1) \Gamma(p\alpha - (p-1)(d+1) + n)}{\Gamma(p\alpha - (p-1)(d+1)) \Gamma(d+1+n)}, n \rightarrow \infty.$$

Together this with Lemma 3.6 shows that there exists a positive constant C_p satisfying

$$\left\langle \frac{1}{C_p} K_{p\alpha-(p-1)(d+1)} f, f \right\rangle \leq \langle K_\alpha^p f, f \rangle \leq \langle C_p K_{p\alpha-(p-1)(d+1)} f, f \rangle,$$

for any $f \in A^2$. This finishes the proof. □

Now we recall the Berezin transform on the unit ball \mathbb{B}^d . The Bergman kernel of \mathbb{B}^d is given by

$$K_w(z) = K(z, w) = \frac{1}{(1 - \langle z, w \rangle)^{d+1}}, \quad z, w \in \mathbb{B}^d,$$

which is also called the reproducing kernel of A^2 , since

$$f(z) = \langle f, K_z \rangle, \quad z \in \mathbb{B}^d \quad (4.5)$$

for any $f \in A^2$. The normalized reproducing kernel of A^2 is

$$k_w(z) = \frac{K(z, w)}{\sqrt{K(w, w)}} = \frac{(1 - |w|^2)^{\frac{d+1}{2}}}{(1 - \langle z, w \rangle)^{d+1}}, \quad z, w \in \mathbb{B}^d. \quad (4.6)$$

For a bounded operator $T \in B(A^2)$, the Berezin transform \widetilde{T} of T is given by

$$\widetilde{T}(z) = \langle T k_z, k_z \rangle, \quad z \in \mathbb{B}^d.$$

The Möbius invariant measure $d\lambda$ on \mathbb{B}^d is defined by

$$d\lambda(z) = \frac{dv(z)}{(1 - |z|^2)^{d+1}}.$$

The Berezin transform is an important tool in the operator theory on the holomorphic function space, see [26, 27, 28] for more details. In what follows, we calculate the Berezin transform of K_α .

Lemma 4.3. $\widetilde{K}_\alpha(z) = (1 - |z|^2)^{d+1-\alpha}$.

Proof. For every $w \in \mathbb{B}^d$, denote the holomorphic function $K_{\alpha,w}(z)$ on \mathbb{B}^d by

$$K_{\alpha,w}(z) = \frac{1}{(1 - \langle z, w \rangle)^\alpha}.$$

Obviously, $K_{\alpha,w} \in A^2$ for every $w \in \mathbb{B}^d$. By (4.5), we have

$$K_\alpha K_z(w) = \overline{\langle K_{\alpha,w}, K_z \rangle} = \frac{1}{(1 - \langle w, z \rangle)^\alpha}.$$

Combing this with (4.5) and (4.6), we get that

$$\begin{aligned} \widetilde{K}_\alpha(z) &= \langle K_\alpha k_z, k_z \rangle \\ &= (1 - |z|^2)^{d+1} \langle K_\alpha K_z, K_z \rangle \\ &= (1 - |z|^2)^{d+1} \langle K_{\alpha,z}, K_z \rangle \\ &= (1 - |z|^2)^{d+1-\alpha}. \end{aligned}$$

This completes the proof. \square

The following lemma establishes a connection between the Berezin transform and Schatten p -class on the Bergman space A^2 , see [19, Lemma C] and [27, Lemma 7.10] for more details.

Lemma 4.4. [19, 27] If $T \in B(A^2)$ is a positive operator, then the followings hold.

(1) $T \in S_1(A^2)$ if and only if $\tilde{T} \in L^1(d\lambda)$. Moreover, the following trace formula holds,

$$\mathrm{Tr}(T) = \int_{\mathbb{B}^d} \tilde{T} d\lambda. \quad (4.7)$$

(2) For $1 < p < \infty$, then $\tilde{T} \in L^p(d\lambda)$ if $T \in \mathcal{L}^p(A^2)$.

Now we can prove Theorem 3. The proof will be given in cases $0 < \alpha < d + 1$ and $\alpha \leq 0$, respectively.

Proof of Theorem 3 for the case $0 < \alpha < d + 1$. Note that the compact operator K_α is adjoint on L^2 by Fubini's theorem, it follows from [23, Theorem 1.9.2] that K_α on L^2 admits the canonical decomposition

$$K_\alpha f = \sum_{n=0}^{\infty} \lambda_n \langle f, e_n \rangle e_n \quad (4.8)$$

whenever $\alpha < d + 1$, where $\{\lambda_n\}$ is the sequence of nonzero eigenvalues (counting multiplicities) and $\{e_n\}$ is the corresponding orthonormal sequence of eigenvectors. It follows from (4.8) that

$$K_\alpha e_n = \lambda_n e_n, n = 0, 1, \dots.$$

Since $K_\alpha e_n$ is holomorphic by Lemma 2.1, we obtain that e_n is holomorphic for any integer $n \geq 0$. Thus K_α on A^2 and L^2 own the same canonical decomposition (4.8). This completes the proof that (2) is always equivalent to (1).

Now we turn to prove that (2) implies (3). Namely, we need to prove $\tilde{K}_\alpha \in L^p(d\lambda)$ if $K_\alpha \in \mathcal{L}^p(A^2)$. Suppose that $K_\alpha \in \mathcal{L}^p(A^2)$. Note that $\frac{d+1}{d+1-\alpha} > 1$, then Lemma 4.4 shows that $\tilde{K}_\alpha \in L^p(d\lambda)$ if $p \geq \frac{d+1}{d+1-\alpha}$. Thus, it suffices to consider the case $0 < p < \frac{d+1}{d+1-\alpha}$. Observe that $0 < p < \frac{d+1}{d+1-\alpha}$ ensures $p\alpha - (p-1)(d+1) > 0$. Then, by Lemma 4.2, there exists a positive constant C'_p such that

$$\frac{1}{C'_p} K_{p\alpha - (p-1)(d+1)} \leq K_\alpha^p. \quad (4.9)$$

Note that $K_\alpha^p \in \mathcal{L}^1(A^2)$, together with (4.9) shows that $K_{p\alpha - (p-1)(d+1)} \in \mathcal{L}^1(A^2)$. Then Lemma 4.3 and Lemma 4.4 imply

$$\begin{aligned} \int_{\mathbb{B}^d} |\tilde{K}_\alpha|^p d\lambda &= \int_{\mathbb{B}^d} (1 - |z|^2)^{p(d+1-\alpha)} d\lambda \\ &= \int_{\mathbb{B}^d} (1 - |z|^2)^{d+1 - (p\alpha - (p-1)(d+1))} d\lambda \\ &= \int_{\mathbb{B}^d} \tilde{K}_{p\alpha - (p-1)(d+1)} d\lambda \\ &= \mathrm{Tr}(K_{p\alpha - (p-1)(d+1)}) \\ &< \infty. \end{aligned}$$

Then $\tilde{K}_\alpha \in L^p(d\lambda)$. This shows that (2) implies (3).

Suppose $\widetilde{K}_\alpha \in L^p(d\lambda)$, we go to prove that $p > \frac{d}{d+1-\alpha}$. By Lemma 4.3, we have

$$\begin{aligned} \int_{\mathbb{B}^d} |\widetilde{K}_\alpha|^p d\lambda &= \int_{\mathbb{B}^d} |(1 - |z|^2)^{p(d+1-\alpha)}| d\lambda \\ &= \int_{\mathbb{B}^d} (1 - |z|^2)^{(p-1)(d+1)-p\alpha} dv \end{aligned}$$

Together this with the condition $\widetilde{K}_\alpha \in L^p(d\lambda)$ shows that $(1 - |z|^2)^{(p-1)(d+1)-p\alpha} \in L^1(\mathbb{B}^d)$. By the fact that, for $t \in \mathbb{R}$, $(1 - |z|^2)^t \in L^1(\mathbb{B}^d)$ if and only if $t > -1$, we conclude that $(p-1)(d+1) - p\alpha > -1$. Then $p > \frac{d}{d+1-\alpha}$. This shows that (3) implies (4).

Now we turn to prove that (4) implies (2), that means that we need to prove $K_\alpha \in \mathcal{L}^p(A^2)$ if $p > \frac{d}{d+1-\alpha}$. As mentioned above, we know that K_α is compact on the Bergman space A^2 , i.e. $K_\alpha \in \mathcal{L}^\infty(A^2)$. Since $\frac{d+1}{d+1-\alpha} > 1$, it suffices to prove that $K_\alpha \in \mathcal{L}^p(A^2)$ when $\frac{d}{d+1-\alpha} < p < \frac{d+1}{d+1-\alpha}$ by the interpolation theorem of Schatten classes, see [27, Theorem 2.6]. Now Lemma 4.2 shows that, there exists a positive constant C_p'' such that

$$K_\alpha^p \leq C_p'' K_{p\alpha-(p-1)(d+1)} \quad (4.10)$$

The condition $\frac{d}{d+1-\alpha} < p < \frac{d+1}{d+1-\alpha}$ means that

$$0 < p\alpha - (p-1)(d+1) < 1.$$

Then Lemma 4.3 shows that

$$\begin{aligned} \int_{\mathbb{B}^d} |\widetilde{K}_{p\alpha-(p-1)(d+1)}| d\lambda &= \int_{\mathbb{B}^d} (1 - |z|^2)^{p(d+1-\alpha)} d\lambda \\ &= \int_{\mathbb{B}^d} (1 - |z|^2)^{(p-1)(d+1)-p\alpha} dv \\ &< \infty. \end{aligned}$$

Combing with Lemma 4.4 follows that $\widetilde{K}_{p\alpha-(p-1)(d+1)} \in \mathcal{L}^1(A^2)$. Therefore, we obtain that $K_\alpha \in \mathcal{L}^p(A^2)$ by (4.10). This shows that (4) implies (2). That completes the proof. \square

Although K_α on A^2 and L^2 own the same canonical decomposition (4.8) when $0 < \alpha < d+1$, the point spectrum of K_α on A^2 and L^2 differ by the element 0. Indeed, the point spectrum of K_α on L^2 is $\{0\} \cup \{\frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)} : n \in \mathbb{N}\}$.

Proof of Theorem 3 for the case $\alpha \leq 0$. We have proven that (1) and (2) is always equivalent when $\alpha < d+1$. Thus, it suffice to prove (2), (3) and (4) are equivalent if $\alpha \leq 0$ and α is not a integer.

Suppose $\alpha \leq 0$ and α is not a integer. From Lemma 4.1 we know that $\{\mu_n : n \in \mathbb{N}\}$ is exactly the point spectrum of K_α on A^2 , where $\mu_n = \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)}$, $n \in \mathbb{N}$. Let E_n be the eigenspace corresponding to μ_n . It follows from (4.3) that

$$\dim E_n = \#P_n = \frac{(n+1)_{d-1}}{(d-1)!}, \quad n \in \mathbb{N}. \quad (4.11)$$

Combing this with the definition of $\mathcal{L}^p(A^2)$, we know that $K_\alpha \in \mathcal{L}^p(A^2)$ if and only if

$$\sum_{n=0}^{\infty} |\mu_n|^p \dim E_n = \sum_{n=0}^{\infty} \left| \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)} \right|^p \frac{(n+1)_{d-1}}{(d-1)!} < \infty. \quad (4.12)$$

Then Stirling's formula implies that (4.12) is equivalent to

$$\sum_{n=1}^{\infty} \frac{1}{n^{p(d+1-\alpha)-(d-1)}} < \infty.$$

It follows that $K_\alpha \in \mathcal{L}^p(A^2)$ if and only if $p > \frac{d}{d+1-\alpha}$. This implies that (2) and (4) are equivalent if $\alpha \leq 0$ and α is not a integer.

Suppose $\alpha \leq 0$ and α is not a integer. From Lemma 4.3, we obtain that $\tilde{K}_\alpha \in L^p(d\lambda)$ if and only if

$$\int_{\mathbb{B}^d} |\tilde{K}_\alpha|^p d\lambda = \int_{\mathbb{B}^d} (1-|z|^2)^{p(d+1-\alpha)} d\lambda = \int_{\mathbb{B}^d} (1-|z|^2)^{p(d+1-\alpha)-(d+1)} dv < \infty. \quad (4.13)$$

By the well-known fact that $(1-|z|^2)^t \in L^1(\mathbb{B}^d)$ if and only if $t > -1$, we conclude that (4.13) is equivalent to $p > \frac{d}{d+1-\alpha}$. This implies that (3) and (4) are equivalent if $\alpha \leq 0$ and α is not a integer. \square

Now we turn to prove Theorem 4.

Proof of Theorem 4. It follows from Lemma 2.1 and the canonical decomposition (4.8) when $\alpha < d+1$ that the normal compact operator K_α on L^2 and A^2 has the same nonzero eigenvalues (counting multiplicities). We thus obtain that $K_\alpha \in \mathcal{L}^{p,\infty}(L^2)$ if and only if $K_\alpha \in \mathcal{L}^{p,\infty}(A^2)$ when $\alpha \in \mathcal{S}_d$. This implies that (1) and (2) are equivalent. Now we turn to prove that (2) and (3) are equivalent. From Lemma 4.1 we know that $\{\mu_n : n \in \mathbb{N}\}$ is exactly the point spectrum of K_α on A^2 , where $\mu_n = \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)}$, $n \in \mathbb{N}$. By the functional calculus in C^* -algebra and the canonical decomposition (4.4), it implies that

$$|K_\alpha|f = \sum_{n=0}^{\infty} \left| \frac{\Gamma(d+1)\Gamma(\alpha+n)}{\Gamma(\alpha)\Gamma(d+1+n)} \right| \sum_{e_{n,k} \in P_n} \langle f, e_{n,k} \rangle e_{n,k}. \quad (4.14)$$

By the direct calculation, we have

$$\frac{|\mu_{n+1}|}{|\mu_n|} = \frac{\Gamma(\alpha+n+1)\Gamma(d+n+1)}{\Gamma(\alpha+n)\Gamma(d+n+2)} = \frac{\alpha+n}{d+1+n} < 1$$

for each $n > |\alpha|$, which means that

$$|\mu_{n_0}| > |\mu_{n_0+1}| > |\mu_{n_0+2}| > \cdots \quad (4.15)$$

where n_0 is the minimal integer belongs to $\{n \in \mathbb{N} : n > |\alpha|\}$. On the other hand, it follows from (4.14) that $\{|\mu_n|\}$ is the set of nonzero eigenvalues (counting multiplicities) of $|K_\alpha|$ and the multiplicity m_n of the eigenvalue $|\mu_n|$ is equal to $\dim E_n$, combing with (4.11) it follows that

$$m_n = \frac{(n+1)_{d-1}}{(d-1)!}, \quad n \in \mathbb{N}.$$

Denote M_n by

$$M_n = \sum_{j=0}^n m_j,$$

for $n \geq 0$. This along the well-known fact that $\sum_{j=1}^k j^n \sim k^{n+1}$ for any $n \in \mathbb{N}$ implies that

$$M_n = \sum_{k=0}^n m_k \sim n^d. \quad (4.16)$$

Then $K_\alpha \in \mathcal{L}^{p,\infty}$ if and only if $\mu_n(|K_\alpha|) = O(n^{-\frac{1}{p}})$ if and only if

$$\mu_{M_n}(|K_\alpha|) = O(M_n^{-\frac{1}{p}})$$

if and only if

$$n^{\alpha-(d+1)} = O(n^{-\frac{d}{p}})$$

if and only if $p \geq \frac{d}{d+1-\alpha}$. It completes the proof. \square

Remark 4.5. Proposition 3.11 follows that K_α is a finite operator whenever α is a nonpositive integer. In this case, $K_\alpha \in \mathcal{L}^p(L^2)$ for any $p > 0$ rather than $p > \frac{d}{d+1-\alpha}$ and $K_\alpha \in \mathcal{L}^{p,\infty}(L^2)$ for any $p > 0$ rather than $p \geq \frac{d}{d+1-\alpha}$. This shows that the condition $\alpha \in \mathcal{S}_d$ in Theorem 3,4 is necessary and sharp.

Corollary 4.6. For $0 \leq \alpha < d + 1$, K_α is bounded on L^p and A^p for any $1 < p < \infty$. Moreover, the following operator norm identity holds.

$$\|K_\alpha\|_{L^p \rightarrow L^p} = \|K_\alpha\|_{A^p \rightarrow A^p} = 1.$$

Proof. It comes from Theorem 2, Lemma 2.1, formula (3.14) and (4.15). \square

Now we recall the the Definition of Dixmier trace on $\mathcal{L}^{1,\infty}(H)$. Let $0 \leq T \in \mathcal{L}^{1,\infty}(H)$ and $\{\mu_n(T)\}$ be the sequence of its point spectrum (counting multiplicities) arranged in decreasing order, namely

$$\mu_0(T) \geq \mu_1(T) \geq \mu_2(T) \geq \cdots$$

Since $0 \leq T \in \mathcal{L}^{1,\infty}$, it follows that the sequence $\{\frac{1}{\log n} \sigma_n\}_{n>1} \in \ell^\infty$, where $\sigma_n = \sum_{j=0}^n \mu_j(T) = \sum_{j=0}^n \mu_j(|T|)$. Taking an arbitrary continuous linear functional Lim_ω on ℓ^∞ satisfying the following three conditions:

- (1) $\text{Lim}_\omega\{a_n\} \geq 0$ if $a_n \geq 0$;
- (2) $\text{Lim}_\omega\{a_n\} = \lim\{a_n\}$ if $\{a_n\}$ is convergent;
- (3) $\text{Lim}_\omega\{a_1, a_1, a_2, a_2, a_3, a_3, \cdots\} = \text{Lim}_\omega\{a_1, a_2, a_3, \cdots\}$.

Then the Dixmier trace $\text{Tr}_\omega(T)$ is defined by

$$\text{Tr}_\omega(T) = \text{Lim}_\omega \frac{1}{\log n} \sigma_n.$$

In this way we define the Dixmier trace for all positive operators in $\mathcal{L}^{1,\infty}(H)$, and it can be uniquely extended by linearity to the whole $\mathcal{L}^{1,\infty}(H)$. In general, the Dixmier trace Tr_ω depends on the linear functional Lim_ω . However, in some special case, Tr_ω will be independent of Lim_ω , for example $\text{Tr}_\omega(T) = 0$ for any $T \in \mathcal{L}^1$. We refer the reader to [7, Chapter 4] for more details. For positive compact operator $T \notin \mathcal{L}^{1,\infty}(H)$,

we define formally $\text{Tr}_\omega(T) = \infty$. Thus $T \in \mathcal{L}^{p,\infty}(H)$ if and only if $|T|^p \in \mathcal{L}^{1,\infty}(H)$ if and only if $\text{Tr}_\omega(|T|^p) < \infty$. Then the following lemma is trivial.

Lemma 4.7. Suppose that $T \in \mathcal{L}^{p,\infty}(H)$ for some $0 < p < \infty$ and $0 < s < t < \infty$.

- (1) If $\text{Tr}_\omega(|T|^s) < \infty$, then $\text{Tr}_\omega(|T|^t) = 0$.
- (2) If $\text{Tr}_\omega(|T|^t) = \infty$, then $\text{Tr}_\omega(|T|^s) = \infty$.

Lemma 4.7 permits us to introduce the Hausdorff dimension of compact operators as below. Compare to the Hausdorff dimension for subsets in metric space [16, Definition 1.1.13].

Definition 4.8. The Hausdorff dimension of a compact operator T is defined to be

$$\dim_{H_\omega}(T) := \inf\{0 < s < \infty : \text{Tr}_\omega(|T|^s) = 0\},$$

if there exists $0 < p < \infty$ such that $T \in \mathcal{L}^{p,\infty}(H)$. Otherwise, the Hausdorff dimension of the compact operator T is

$$\dim_{H_\omega}(T) := \infty.$$

It is immediate that $\dim_{H_\omega}(T) \leq p$ if $T \in \mathcal{L}^{p,\infty}(H)$. In fact, there is no difficulty in proving that

$$\dim_{H_\omega}(T) = \inf\{0 < p < \infty : T \in \mathcal{L}^{p,\infty}(H)\} \quad (4.17)$$

if $T \in \mathcal{L}^{p,\infty}(H)$ for some $0 < p < \infty$. Which means the definition of Hausdorff dimension of operator is independent of the choice of the linear functional Lim_ω . Thus, we will omit the functional ω in \dim_{H_ω} in the following. In particular, $\dim_H(T) = 0$ for any finite rank operator T . The following example shows that the Hausdorff dimension can take every value in the interval $[0, \infty]$.

Example 4.9. Let T be a positive compact operator and $\{\mu_n(T)\}$ be the sequence of its point spectrum (counting multiplicities) arranged in decreasing order.

- (1) If $\mu_n(T) = 0$ for any $n > n_0$, then $\dim_H(T) = 0$.
- (2) If $\mu_n(T) = n^{-\frac{1}{\beta}}$ for $0 < \beta < \infty$, then $\dim_H(T) = \beta$.
- (3) If $\mu_n(T) = \frac{1}{\log n}$, then $\dim_H(T) = \infty$.

In the rest of the paper, we will use \mathcal{L}^p instead of $\mathcal{L}^p(L^2)$ or $\mathcal{L}^p(A^2)$ and $\mathcal{L}^{p,\infty}$ instead of $\mathcal{L}^{p,\infty}(L^2)$ or $\mathcal{L}^{p,\infty}(A^2)$ for $0 < p < \infty$.

Proposition 4.10. Bergman-type operator $K_\alpha \in \mathcal{L}^{1,\infty}$ if and only if $\alpha \leq 1$. Moreover, the Dixmier trace of K_α is

$$\text{Tr}_\omega(K_\alpha) = \begin{cases} 0, & \alpha < 1; \\ 1, & \alpha = 1. \end{cases} \quad (4.18)$$

Proof. By Theorem 4, it suffices to prove the Dixmier trace formula (4.18). It follows from Theorem 3 that $K_\alpha \in \mathcal{L}^1$ whenever $\alpha < 1$, thus $\text{Tr}_\omega(K_\alpha) = 0$ whenever $\alpha < 1$. In what follows, we calculate the value of $\text{Tr}_\omega(K_1)$. Note that K_1 is positive and its point spectrum is $\{\mu_n = \frac{\Gamma(d+1)\Gamma(1+n)}{\Gamma(d+1+n)} : n \in \mathbb{N}\}$ from Lemma 4.1. By the direct calculation, we have

$$\frac{\mu_{n+1}}{\mu_n} = \frac{\Gamma(1+n+1)\Gamma(d+n+1)}{\Gamma(1+n)\Gamma(d+n+2)} = \frac{1+n}{d+1+n} < 1,$$

for any $n \geq 0$, namely

$$\mu_0 > \mu_1 > \mu_2 > \cdots > 0. \quad (4.19)$$

It follows from (4.3) and (4.11) that the multiplicity m_n of μ_n is

$$m_n = \frac{(n+1)_{d-1}}{(d-1)!} > 0$$

for any $n \geq 0$. Note that for any integer $k \geq 0$, then there exists a unique integer $n \geq 0$ satisfying

$$M_n \leq k < M_{n+1},$$

where

$$M_n = \sum_{j=0}^n m_j,$$

for $n \geq 0$. Combing this with (4.19), it follows that

$$\frac{\sigma_{M_n}}{\log M_{n+1}} \leq \frac{1}{\log k} \sigma_k \leq \frac{\sigma_{M_{n+1}}}{\log M_n}, \quad (4.20)$$

where $\sigma_k = \sum_{j=0}^k \mu_j(K_1) = \sum_{j=0}^k \mu_j(|K_1|)$. It implies from (4.16) that

$$\log M_n = \log \sum_{j=0}^n \frac{(j+1)_{d-1}}{(d-1)!} \approx d \log n,$$

where $a(n) \approx b(n)$ means that $a(n), b(n) \rightarrow +\infty$ and $\frac{a(n)}{b(n)} \rightarrow 1$. The relationship between $\{\mu_n(K_1)\}$ and $\{\mu_n\}$ implies that

$$\sigma_{M_n} = \sum_{j=0}^n \mu_j m_j = \sum_{j=0}^n \frac{\Gamma(d+1)\Gamma(1+j)}{\Gamma(d+1+j)} \frac{(j+1)_{d-1}}{(d-1)!} = \sum_{j=0}^n \frac{d}{j+d} \approx d \log n.$$

Combing with (4.20) follows that

$$\lim_{n \rightarrow \infty} \frac{1}{\log n} \sigma_n = 1.$$

Thus

$$\text{Tr}_\omega(K_1) = \text{Lim}_\omega \frac{1}{\log n} \sigma_n = \lim_{n \rightarrow \infty} \frac{1}{\log n} \sigma_n = 1,$$

by the condition (2) in the definition of Dixmier trace. It completes the proof. \square

Corollary 4.11. The followings hold.

- (1) If $\alpha \in \mathcal{S}_d$, then $\dim_{\mathbb{H}}(K_\alpha) = \frac{d}{d+1-\alpha}$.
- (2) If α is a nonpositive integer, then $\dim_{\mathbb{H}}(K_\alpha) = 0$.

Proof. (1) It comes from Theorem 3 and (4.17).

(2) It comes from Lemma 3.11 that K_α is finite rank if α is nonpositive integer. \square

Then Theorem 3,4, Proposition 3.11 and Corollary 4.11 imply the following intrinsic characterization for Schatten class and Macaev class Bergman-type operator K_α .

Corollary 4.12. For $0 < p < \infty$, the followings hold.

- (1) If K_α is compact, then $K_\alpha \in \mathcal{L}^p$ if and only if $p > \dim_{\mathbb{H}}(K_\alpha)$.
- (2) If K_α is compact, then $K_\alpha \in \mathcal{L}^{p,\infty}$ if and only if $p \geq \dim_{\mathbb{H}}(K_\alpha)$.

The compact condition in Corollary 4.12 can be removed, for example, since both $K_\alpha \in \mathcal{L}^p$ and $p > \dim_{\text{H}}(K_\alpha)$ can derive the compactness of K_α . Corollary 4.12 provides an example that the Schatten membership of a compact operator can be characterized by its Hausdorff dimension. However, compare to [7, Proposition 4.3.14], which means that the Schatten membership of some compact operators can be characterized by the Hausdorff dimension of some subsets in metric spaces.

Acknowledgements. The authors would like to thank Qi'an Guan and Kai Wang for their helpful discussions.

REFERENCES

- [1] A. Adama, F. Fournier, *Sobolev Spaces*, Pure and Applied Mathematics (Amsterdam), Elsevier/Academic Press, Amsterdam (2003)
- [2] P. Akilov, V. Kantorovich, *Functional analysis*, Pergamon, Oxford (1982)
- [3] T. Ando, *On the compactness of integral operators*, Indag. Math., **24**, 235-239 (1962)
- [4] G. Cheng, X. Fang, Z. Wang, J. Yu, *The hyper-singular cousin of the Bergman projection*, Trans. Amer. Math. Soc. **369**, 8643-8662 (2017)
- [5] G. Cheng, X. Hou, C. Liu, *The singular integral operator induced by Drury-Arveson kernel*, Complex Anal. Oper. Theory, **12**, 917-929 (2018)
- [6] F. Cobos, J. Peetre, *Interpolation of compactness using Aronszajn-Gagliardo functors*, Israel J. Math. **68**, 220-240 (1989)
- [7] A. Connes, *Noncommutative Geometry*, Academic Press Inc., San Diego, CA (1994)
- [8] S. Cui, *Introduction to modern theory of partial differential equations* (Chinese), Modern Mathematics Foundation Series, Higher Education Press, Beijing (2016)
- [9] L. Ding, K. Wang, *The L^p - L^q problems of Bergman-type operators*, arXiv:2003.00479
- [10] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Higher Transcendental Functions, Vol. I*, McGraw-Hill, New York (1953)
- [11] F. Forelli, W. Rudin, *Projections on spaces of holomorphic functions in balls*, Indiana Univ. Math. J. **24**, 593-602 (1974)
- [12] A. Krasnoselskii, *On a theorem of M. Riesz*, Soviet Math. Dokl. **1** 229-231 (1960)
- [13] A. Krasnoselskii et al., *Integral operators in spaces of summable functions*, Nordhoff, Groningen (1976)
- [14] L. Lanzani, E. Stein, *The Bergman projection in L^p for domains with minimal smoothness*, Illinois J. Math. **56**, 127-154 (2013)
- [15] H. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. Math. **118**, 349-374 (1983)
- [16] F. Lin, X. Yang, *Geometric measure theory—an introduction*, Advanced Mathematics, Science Press Beijing, Beijing (2002)
- [17] D. Mcneal, *The Bergman projection as a singular integral operator*, J. Geom. Anal. **1**, 91-103 (1994)
- [18] D. Mcneal, E. Stein, *Mapping properties of the Bergman projection on convex domains of finite type*, Duke Math. J. **73**, 177-199 (1994)
- [19] J. Pau, *Characterization of Schatten-class Hankel operators on weighted Bergman spaces*, Duke Math. J., **165**, 2771-2791 (2016)
- [20] H. Phong, E. Stein, *Estimates for the Bergman and Szegő projections on strongly pseudoconvex domains*, Duke Math. J. **44**, 695-704 (1977)
- [21] N. Plessis, *Some theorems about the Riesz fractional integral*, Trans. Amer. Math. Soc. **80**, 124-134 (1955)

- [22] W. Rudin, *Function Theory in the Unit Ball of \mathbb{C}^n* , Grundlehren der Math. Springer, New York (1980)
- [23] J. Ringrose, *Compact non-self-adjoint operators*, Van Nostrand Reinhold Co., London (1971)
- [24] B. Wong, *Characterization of the unit ball in \mathbb{C}^n by its automorphism group*, Invent. Math. **41**, 253-257 (1977)
- [25] R. Zhao, K. Zhu, *Theory of Bergman Spaces in the Unit Ball of \mathbb{C}^n* , Mém. Soc. Math. Fr. **115** (2009)
- [26] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, Graduate Texts in Mathematics, **226**, Springer-Verlag, New York (2005)
- [27] K. Zhu, *Operator Theory in Function Spaces. Second Edition*, Mathematical Surveys and Monographs, 138, American Mathematical Society, Providence (2007)
- [28] K. Zhu, *Schatten class Toeplitz operators on weighted Bergman spaces of the unit ball*, New York J. Math., **13**, 299-316 (2007)

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING, 100086, P. R. CHINA
E-mail address: ljding@pku.edu.cn

DEPARTMENT OF MATHEMATICS SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN, LIAONING, 116024, P. R. CHINA
E-mail address: junmeifan666@mail.dlut.edu.cn