

PREPARED FOR SUBMISSION TO JHEP

Dimension-8 Operators in the Standard Model Effective Field Theory

Christopher W. Murphy

ABSTRACT: We present a complete basis of dimension-8 operators in the Standard Model Effective Field Theory. Attention is paid to operators that vanish in the absence of flavor structure. The 44,807 operators are encoded in 1,029 Lagrangian terms. We also briefly discuss a few aspects of phenomenology involving dimension-8 operators, including light-by-light scattering and electroweak precision data.

Contents

1	Introduction	1
2	Notation and Conventions	2
3	Operator Classification	4
3.1	Bosonic Operators	6
3.2	Two-Fermion Operators	6
3.3	Four-Fermion Operators	10
4	The Complete Set of Dimension-8 Operators	16
4.1	Results for Bosonic Operators	18
4.2	Results for Two-Fermion Operators	20
4.3	Results for Four-Fermion Operators	26
5	Phenomenology	35
5.1	Light-by-Light Scattering	35
5.2	Electroweak Precision Data	36
5.3	Scalar $SU(2)_w$ Quartets	38
6	Conclusions	39
	Acknowledgements	40
A	Dimension-6 and -7 Operators	40

1 Introduction

The Standard Model (SM) is an extremely successful theory that has been rigorously tested at the Large Hadron Collider (LHC) and elsewhere. Nevertheless it is widely expected that the SM is only an effective field theory (EFT), valid up to some cutoff scale Λ . The Standard Model Effective Field Theory (SMEFT) generalizes the SM by adding a complete, but not over-complete basis of operators at every mass-dimension d rather than stopping at $d = 4$.¹

The counting and classification of operators in the SMEFT has a long history. Starting with dimension-5 there is a single type operator [1], $N_{\text{type}} = 1$, and it violates lepton number. At dimension-6, Ref. [2] classified the 76 baryon number preserving (B) Lagrangian terms; see [3] for earlier work. The eight baryon number violating (\not{B}) terms were previously

¹The SMEFT assumes there are no light hidden states such as sterile neutrinos or an axion, and that the Higgs boson form part of an $SU(2)_w$ doublet with hypercharge $y = 1/2$. Other types of EFTs are possible where these assumptions are relaxed, but we do not consider them here.

known [4], yielding a total of $N_{\text{term}} = 84$. In terms of actual operators rather than terms in the Lagrangian, the counts explode when flavor structure is allowed. For three generations of fermions, $n_g = 3$, there are $N_{\text{op}} = 2499$ independent B operators [5] and 546 \not{B} operators [6]. Hilbert series methods were applied to the SMEFT in Refs. [7–10], providing an elegant way to count the number of operators for arbitrary dimension d . Computing tools **Sim2Int** [11], **DEFT** [12], **BasisGen** [13], and **ECO** [14] were subsequently developed, allowing for automated counting of operators.

Beyond counting operators, work has been done on their explicit forms as well. Refs. [15, 16] classified the 18 dimension-7 operators. So far only partial sets of dimension-8 operators exist in the literature. This includes, however, all of the bosonic operators (in a basis where the number of derivatives is minimized) [17–19]. Our goal in this work is to find a complete set of dimension-8 operators. A subtlety in constructing the dimension-8 operator basis is that some operators vanish in the absence of flavor structure. Our basis contains 231 types of operators corresponding to 1029 Lagrangian terms. For comparison, with $n_g = 1$ there are 993 operators, while for $n_g = 3$ there are instead 44807 operators [10]. We find there are 36 terms that vanish when $n_g = 1$, consistent with the counting of Ref. [10].

Although the counting and classification of operators is certainly interesting in its own right, there is also a wide range of phenomenological implications of dimension-8 operators as well. For some phenomena dimension-8 is the lowest dimension where the interactions become possible. Most famous among these processes is light-by-light scattering. Another area where dimension-8 effects have been studied is electroweak precision data (EWPD) where contributions to the U parameter first arise at dimension-8 [20]. Formally the dimension-6 operators are the leading terms in the EFT expansion. However there are various scenarios in which this is not the case practically speaking. Perhaps the most obvious among these is when the interference between the dimension-6 amplitude and the SM amplitude is suppressed or even vanishes. Additionally there could be a difference in the experimental precision of the measurements being considered [21].

The rest of the paper is organized as follows. Section 2 lays down the notation and conventions we use, including the semantics of number of operators versus number of types of operators. We then discuss how we performed the operator classification in Section 3 with the results given in Sec. 4. We briefly explore light-by-light scattering, EWPD, as well as models involving scalar $SU(2)_w$ quartets where there is interesting interplay between dimension-6 and dimension-8 effects in Section 5 before concluding in Sec. 6. For convenience we provide tables of dimension-6 and -7 operators in Appendix A.

2 Notation and Conventions

We start by considering the various uses of the word operator. See Ref. [22] for further discussion. We define operator to be a gauge and Lorentz invariant contraction of fields and derivatives with specific flavor indices. A Lagrangian term, or just term or short, collects all of operators with the same gauge and Lorentz structure into a single unit, *i.e.* a term collapses the flavor indices of otherwise identical operators. By construction, a

Lagrangian term of mass dimension $d \leq 8$ may contain no more than n_g^4 operators.² Lastly, Ref. [22] defines a type of field as the collection of terms with the combination of fields (and derivatives) where conjugates are counted separately. In this work we use a broader definition of a type of operator where the conjugate fields are counted in unison with the un-conjugated fields. Our types of operators are therefore supersets of those in [22], of which there are 541 to our 231. This definition of a type of operator allows us systematically label the operators in a phenomenologically friendly way.

Moving onto physics conventions, the SM Lagrangian is given by

$$\begin{aligned} \mathcal{L}_{\text{SM}} = & -\frac{1}{4} \sum_X X_{\mu\nu} X^{\mu\nu} + (D_\mu H^\dagger)(D^\mu H) + \sum_\psi \bar{\psi} i \not{D} \psi \\ & - \lambda \left(H^\dagger H - \frac{v^2}{2} \right)^2 - \left[H_j^\dagger \bar{d} Y_d q^j + \tilde{H}_j^\dagger \bar{u} Y_u q^j + H_j^\dagger \bar{e} Y_e l^j + \text{h.c.} \right]. \end{aligned} \quad (2.1)$$

In Eq (2.1), and throughout this work, we generically refer to field strengths as $X = \{G^A, W^I, B\}$, and to fermions as $\psi = \{l, e, q, u, d\}$.

The gauge covariant derivative is

$$(D_\mu q)^{j\alpha} = ((\partial_\mu + i g_1 \mathbf{y} B_\mu) \delta_\beta^\alpha \delta_k^j + i g_2 (t^I)^j_k W_\mu^I \delta_\beta^\alpha + i g_3 (T^A)^\alpha_\beta A_\mu^A \delta_k^j) q^{k\beta}, \quad (2.2)$$

where the generators of $SU(3)_c$ and $SU(2)_w$ are T^a and $t^I = \tau^I/2$, respectively. The $U(1)_y$ hypercharge is given by \mathbf{y} with $Q = \tau^3 + \mathbf{y}$. For $SU(3)_c$ fundamental and adjoint indices are denoted α, β, γ and A, B, C , respectively, while for $SU(2)_w$ the fundamental and adjoint indices are respectively labeled j, k, m and I, J, K .

Anti-symmetrization of indices is denoted by a pair of square brackets, $[\mu\nu]$, and symmetrization is denoted by a pair of round brackets, $(\mu\nu)$. The definition of \tilde{H} is

$$\tilde{H}_j = \epsilon_{jk} H^{\dagger k} \quad (2.3)$$

where $\epsilon_{jk} = \epsilon_{[jk]}$ is the $SU(2)$ invariant tensor with $\epsilon_{12} = 1$. The dual field strength is defined as

$$\tilde{X}_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} X^{\rho\sigma} \quad (2.4)$$

with $\epsilon_{0123} = 1$.

We will sometimes refer to the following combinations of field strength as they are typically what are used when counting coefficients,

$$X_{L,R}^{\mu\nu} = \frac{1}{2} (X^{\mu\nu} \mp i \tilde{X}^{\mu\nu}). \quad (2.5)$$

These field strengths have simple Lorentz transformation properties, $X_L \sim (1, 0)$, $X_R \sim (0, 1)$ under $SU(2)_L \otimes SU(2)_R$. Similarly l and q are left-handed fermion fields, whereas e , u , and d are right-handed fields. When necessary Lorentz indices in the fundamental representations are indicated by a, b, \dot{a}, \dot{b} , *e.g.* $q_L \sim (q_L)_a$, $B_R \sim (B_R)_{(\dot{a}\dot{b})}$.

²Starting at dimension-9 n_g^6 is possible.

The SMEFT extends the SM by adding all of the higher-dimensional operator that are gauge invariant under the SM with the caveat that redundant operators should not be included. Schematically the operators of mass-dimension d are given by

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_{d>4} \mathcal{L}^{(d)}, \quad \mathcal{L}^{(d)} = \sum_i C_i Q_i. \quad (2.6)$$

For the dimension-6 operators we keep notation that has been well-established in the literature, see *e.g.* [5]. On the other hand, we use a systematic, if at times cumbersome, notation for labelling the operators of mass-dimension 7 and above. The subscript label gives the type of operator, denoted as the fields and the derivatives in the operator raised to the power of the number of times that type of object appears in the operator. If a type of operator has multiple terms, not counting Hermitian conjugates, there is also a superscript label that iterates over the terms. The notation is best illustrated with an example,

$$\mathcal{L}^{(d=8)} \supset C_{u^2 GH^2 D, pr}^{(1)} Q_{u^2 GH^2 D, pr}^{(1)} + C_{u^2 GH^2 D, pr}^{(2)} Q_{u^2 GH^2 D, pr}^{(2)} + \left[C_{leBH^3, pr} Q_{leBH^3, pr} + \text{h.c.} \right], \quad (2.7)$$

with

$$\begin{aligned} Q_{u^2 GH^2 D, pr}^{(1)} &= (\bar{u}_p \gamma^\nu T^A u_r) D^\mu (H^\dagger H) G_{\mu\nu}^A, \\ Q_{u^2 GH^2 D, pr}^{(2)} &= (\bar{u}_p \gamma^\nu T^A u_r) D^\mu (H^\dagger H) \tilde{G}_{\mu\nu}^A, \\ Q_{leBH^3, pr} &= (\bar{l}_p \sigma^{\mu\nu} e_r) H (H^\dagger H) B_{\mu\nu}. \end{aligned} \quad (2.8)$$

Flavor indices explicitly appear in Eq. (2.7). Fermion fields have a flavor index p, r, s, t that runs over 1, 2, 3 for three generations. The fermion fields themselves are in the weak eigenstate basis. The Yukawa matrices, $Y_{e,u,d}$, in Eq. (2.1) are matrices in flavor space.

Note that we do not explicitly label the transpose of a spinor in fermion bilinears involving a charge conjugation operator, *e.g.* $\psi_1 C \psi_2 \equiv \psi_1^T C \psi_2$, $\psi_1 C \sigma_{\mu\nu} \psi_2 \equiv \psi_1^T C \sigma_{\mu\nu} \psi_2$. Finally, it is convenient to define Hermitian derivatives *e.g.*

$$\begin{aligned} iH^\dagger \overleftrightarrow{D}_\mu H &= iH^\dagger (D_\mu H) - i(D_\mu H^\dagger) H, \\ iH^\dagger \overleftrightarrow{D}_\mu^I H &= iH^\dagger \tau^I (D_\mu H) - i(D_\mu H^\dagger) \tau^I H. \end{aligned} \quad (2.9)$$

3 Operator Classification

Having defined our notation in Sec. 2 we can more precisely state our goal. We are trying to find the minimum number of Lagrangian terms needed to give all of the operators at dimension-8 subject to the constraint that no term may contain more than n_g^4 operators. For bosonic and two-fermion operators this constraint is trivially satisfied as those terms always contain one and n_g^2 operators, respectively. Four-fermion operators where two or more of the fields are identical constitute the interesting cases.

We use existing results from the literature when they are available. All of the bosonic operators have been classified previously [17–19]. Ref. [18] also gave partial results for three

of the two-fermion classes that were sufficient to allow us to deduce the remaining operators in those classes. As this work was being finalized Refs. [23, 24] appeared, which classified a subset of four-fermion operators with two derivatives. However there are still non-trivial results for us to work out in that class.

When classifying the dimension-8 operators we exploit the fact that not only are the types of operators known, but the number of operators is also known, first being in given Ref. [10]. In particular, we leverage the Python package **BasisGen** [13], which we use to get the number of operators for each type of operator. Additionally, we use the Mathematica program **Sym2Int** [11], which not only gives the number of operators per type, but also the flavor representations when there are identical particles in the operator. Furthermore, **Sym2Int** gives the number of Lagrangian terms per type of operator except when the operator contains both derivatives and identical particles. In that case a range is given because the permutation symmetry of operators with derivatives is ambiguous due to integration by parts (IBP) redundancies.

Ref. [22] used **Sym2Int** to derive lower and upper limits on the number of terms of dimension-8 operators, $1025 \leq N_{\text{term}} \leq 1102$. Our basis contains 1029 terms, close to the lower limit. The counting is clear from [10] for that one generation of fermions there are 993 terms. Of the remaining 36 terms we identified, all of which vanish in the absence of flavor structure, four of them involve derivatives. For each of these four terms there is another term of the same type that does not vanish in the absence of flavor structure, which is consistent with the ambiguity originating from terms with derivatives. In principle this could allow us to reduce the number of terms in our basis by four and hit the lower limit. However, the total number of operators in each case, $n_g^4 + \frac{1}{2}n_g^3(n_g - 1)$, exceeds the maximum of operators that can be placed in a single term, n_g^4 . Therefore the four potentially redundant terms do need to be retained as independent terms in the Lagrangian, suggesting we have indeed found the minimum number of terms.

Beyond getting the number of terms correct we need to ensure that the operators in our basis are independent. Operators with derivatives can be related through integration by parts. When there are multiple derivatives care must be taken to select operators for the basis that span the entire space of possible operators for that class. See the discussion of class 16 below for an example of this. Operators can also be related to each other through the equations of motion (EOM). We use the EOM to remove redundant terms, trading them for basis operators in the same class, operators with fewer derivatives, and sometimes operators of lower mass dimension. See the discussion of class 17 below for an example of this, and see *e.g.* [25] for the SM equations of motion. Our basis does not explicitly contain an EOM. Operators with derivatives can be IBP, and some of the resulting terms contain an EOM. However it is never the case that all of the resulting terms have an EOM. Lastly, there are various tensor and spinor identities that may cause operators to be related to each other. There are the Fierz identities, for example for $SU(2)$

$$(\tau^I)_j^k (\tau^I)_m^n = 2\delta_j^n \delta_m^k - \delta_j^k \delta_m^n. \quad (3.1)$$

There are identities involving the Levi-Civita symbol, *e.g.* in two-dimensions

$$\epsilon_{jk}\epsilon_{mn} + \epsilon_{jm}\epsilon_{nk} + \epsilon_{jn}\epsilon_{km} = 0 \quad (3.2)$$

There are identities for products of Dirac matrices, *e.g.* the anti-symmetric Dirac tensor is self-dual

$$\epsilon^{\rho\tau\mu\nu}\sigma_{\mu\nu}P_R = 2i\sigma^{\rho\tau}P_R. \quad (3.3)$$

3.1 Bosonic Operators

1. X^4

The X^4 operators for a single Yang-Mills field were classified in Ref. [17]. Ref. [19] generalized this result to the SM field content. Note that dimension-8 is the lowest dimension where X_L and X_R appear in the same operator together.

2. H^8

$(H^\dagger H)^4$ is the only possibility.

3. $H^6 D^2$

The $H^6 D^2$ operators were classified in Ref. [18].

4. $H^4 D^4$

Both Refs. [18] and [19] classified the $H^4 D^4$ operators.

5. $X^3 H^2$

The $X^3 H^2$ operators were classified in Ref. [18].

6. $X^2 H^4$

The $X^2 H^4$ operators were classified in Ref. [18].

7. $X^2 H^2 D^2$

Both Refs. [18] and [19] classified the $X^2 H^2 D^2$ operators.

8. $X H^4 D^2$

The $X H^4 D^2$ operators were classified in Ref. [18].

3.2 Two-Fermion Operators

9. $\psi^2 X^2 H$

For the dimension-8 class $\psi^2 X^2 H$, 24 terms arise from joining a field strength to a dimension-6 operator of the form $\psi^2 X H$, whereas 48 terms come from the product of two field strengths and a Yukawa interaction, $(X^2)(\psi^2 H)$. See Table 20 for the dimension-6 operators.

10. $\psi^2 X H^3$

In the class $\psi^2 X H^3$, 16 of the 22 terms are identical to the dimension-6 terms $\psi^2 X H$ up to an extra factor of $(H^\dagger H)$. The remaining six terms, all involving $W_{\mu\nu}^I$, instead have the dimension-2 covariant $(H^\dagger \tau^I H)$.

11. $\psi^2 H^2 D^3$

Ref. [18] classified the four terms involving $q^2 H^2 D^3$. The remaining 12 terms in the class can be deduced from the results of Ref. [18].

12. $\psi^2 H^5$

The class $\psi^2 H^5$ is identical to the dimension-6 class $\psi^2 H^3$ up to an extra factor of $(H^\dagger H)$.

13. $\psi^2 H^4 D$

Ref. [18] classified the four operators involving $q^2 H^4 D$. The remaining nine terms in the class can be deduced from the results of Ref. [18].

14. $\psi^2 X^2 D$

We use integration by parts to place the derivative on a fermion field. Then in order for the operator to not be “reduced” to a class with fewer derivatives through the use of the equations of motion the fermionic component of the operator must not be Lorentz invariant. Class 7 also contains two field strengths, see above, and a subset of the operators in class 7 have covariants formed from Higgs fields and derivatives that transform as $(1, 1)$ under $SU(2)_L \otimes SU(2)_R$. We take the field strength components of that subset of class 7 operators and use them for the class 14 operators, contracting them with fermionic covariants of the form $\bar{\psi} \gamma^\mu D^\nu \psi$. In particular, we use $Q_{G^2 H^2 D^2}^{(1)}$ as the template for when a fermion is not charged under a gauge group, $Q_{W^2 H^2 D^2}^{(1, 4-6)}$ when it is charged under a gauge group, and $Q_{WBH^2 D^2}^{(1, 4-6)}$ when it is charged under two gauge groups. Three terms are not covered by this procedure. They involve quarks and two gluon field strength where the $SU(3)_c$ adjoint indices are contracted with the symmetric d^{ABC} symbol.

15. $\psi^2 X H^2 D$

Ref. [18] classified the 12 terms involving $q^2 W H^2 D$. Of the remaining 74 terms, 68 of them have a form analogous to those classified by Ref. [18]. The final six terms in the class are instead analogous to the dimension-6 operator Q_{Hud} with the addition of a field strength.

16. $\psi^2 X H D^2$

Things become more complicated when there are two or more derivatives in the operator. As such it is useful to introduce some additional machinery to classify the operators. We use the procedure given in Ref. [7] for removing terms that are reducible through the use of the equations of motion. In a nutshell, the procedure says Lorentz indices should be symmetrized for representations that are triplets or higher under either $SU(2)_L$ or $SU(2)_R$. In terms of Lorentz indices we have for example

$$D\psi_L \sim (D\psi_L)_{(ab),\dot{a}}, \quad DX_R \sim (DX_R)_{a,(\dot{a}\dot{b}\dot{c})}, \quad D^2 H \sim (D^2 H)_{(ab),(\dot{a}\dot{b})}. \quad (3.4)$$

We now work through a representative example with field content \bar{l}, e, H, B_L following the procedure laid out in Ref. [18]. Using relations like those in (3.4) we see that

operators with a derivative acting on the field strength or two derivatives acting on a fermion can be reduced using the EOM. Ignoring for the time being constraints from IBP this leaves us with four possibilities

$$\begin{aligned}
x_1 &= (D\bar{l})_{a,(\dot{a}\dot{c})}e_{\dot{d}}(DH)_{b,\dot{b}}B_{(cd)}\epsilon^{ac}\epsilon^{bd}\epsilon^{\dot{a}\dot{d}}\epsilon^{\dot{c}\dot{b}}, \\
x_2 &= \bar{l}_{\dot{c}}(De)_{a,(\dot{a}\dot{d})}(DH)_{b,\dot{b}}B_{(cd)}\epsilon^{ac}\epsilon^{bd}\epsilon^{\dot{a}\dot{c}}\epsilon^{\dot{d}\dot{b}}, \\
x_3 &= (D\bar{l})_{a,(\dot{a}\dot{c})}(De)_{b,(\dot{b}\dot{d})}HB_{(cd)}\epsilon^{ac}\epsilon^{bd}\epsilon^{\dot{a}\dot{b}}\epsilon^{\dot{c}\dot{d}}, \\
x_4 &= \bar{l}_{\dot{c}}e_{\dot{d}}(D^2H)_{(ab),(\dot{a}\dot{b})}B_{(cd)}\epsilon^{ac}\epsilon^{bd}\epsilon^{\dot{a}\dot{c}}\epsilon^{\dot{b}\dot{d}},
\end{aligned} \tag{3.5}$$

where we have not shown the $SU(2)_w$ contraction as it is trivial.

To determine redundancies coming from integration by parts we need operators transforming as $(\frac{1}{2}, \frac{1}{2})$ under the Lorentz group with one fewer derivative than the operators of interest. There are three possibilities in this example

$$\begin{aligned}
y_1 &= (D\bar{l})_{a,(\dot{a}\dot{c})}e_{\dot{d}}HB_{(cd)}\epsilon^{ac}\epsilon^{\dot{a}\dot{d}}, \\
y_2 &= \bar{l}_{\dot{c}}(De)_{a,(\dot{a}\dot{d})}HB_{(cd)}\epsilon^{ac}\epsilon^{\dot{a}\dot{c}}, \\
y_3 &= \bar{l}_{\dot{c}}e_{\dot{d}}(DH)_{a,\dot{a}}B_{(cd)}\frac{1}{2}\epsilon^{ac}(\epsilon^{\dot{a}\dot{c}} + \epsilon^{\dot{a}\dot{d}}).
\end{aligned} \tag{3.6}$$

The derivatives of the y_i show which of the x_i are related by IBP

$$\begin{aligned}
D_{b,\dot{b}}y_1 &= x_1 + x_3 = 0, \\
D_{b,\dot{b}}y_2 &= x_2 + x_3 = 0, \\
D_{b,\dot{b}}y_3 &= \frac{1}{2}x_1 + \frac{1}{2}x_2 + x_4 = 0,
\end{aligned} \tag{3.7}$$

where the appropriate contraction of the remaining Lorentz indices in the leftmost terms is understood. By inspection of (3.7) we see that any of the four candidate operator can be transformed into any of the remaining three through the use of IBP.

We repeat the same procedure, omitting the details here, to find the class 17 operators with field content \bar{l}, e, H, B_R . In this case there are eight candidate operators, x_i , and six operators in the Lorentz four-vector representation, y_i . Given the larger number of operators in this case we use Mathematica to solve the system of constraints, yielding two operators.

We are left with a total of three Q_{leHBD^2} terms (+h.c.)

$$\begin{aligned}
&\epsilon^{ac}\epsilon^{bd}\epsilon^{\dot{a}\dot{b}}\epsilon^{\dot{c}\dot{d}}\bar{l}_{\dot{a}}(De)_{a,(\dot{b}\dot{c})}(DH)_{b,\dot{d}}(B_L)_{cd}, \\
&\epsilon^{ab}\epsilon^{\dot{a}\dot{b}}\epsilon^{\dot{c}\dot{d}}\epsilon^{\dot{e}\dot{f}}\bar{l}_{\dot{a}}(De)_{a,(\dot{b}\dot{c})}(DH)_{b,\dot{d}}(B_R)_{(\dot{e}\dot{f})}, \\
&\epsilon^{ab}\epsilon^{\dot{a}\dot{d}}\epsilon^{\dot{b}\dot{e}}\epsilon^{\dot{c}\dot{f}}\bar{l}_{\dot{a}}e_{\dot{b}}(DH)_{a,\dot{c}}(DB_R)_{b,(\dot{d}\dot{e}\dot{f})},
\end{aligned} \tag{3.8}$$

which matches the counting we found using **BasisGen**. Translating the Lorentz contractions from $SU(2)_L \otimes SU(2)_R$ to $SO(3,1)$ and translating B_L and B_R to B and

\tilde{B} we find

$$\begin{aligned} Q_{leBHD^2}^{(1)} &= (\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) (D_\nu H) B_{\rho\nu} \\ Q_{leBHD^2}^{(2)} &= (\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) (D_\nu H) \tilde{B}_{\rho\nu} \\ Q_{leBHD^2}^{(3)} &= (\bar{l}_p \sigma^{\mu\nu} e_r) (D^\rho H) (D_\rho B_{\mu\nu}). \end{aligned} \quad (3.9)$$

The dual field strength makes an appearance because it is not fully contracted with $\sigma^{\mu\nu}$, otherwise Eq. (3.3) would convert back to an ordinary field strength. The remaining 42 terms in class 17 can be deduced from the Q_{leBHD^2} operators in (3.9) +h.c..

17. $\psi^2 H^3 D^2$

Starting with relations like those in (3.4) we see that any operator in this class with two derivatives acting on the same field can be reduced using the EOM. Therefore an operator with one derivative on each of the two fermion fields can be traded for an operator with one derivative on a Higgs field, one derivative on a fermion plus operators with fewer derivatives. From here we extend the results of Ref. [2] to move any remaining derivatives acting on fermions onto Higgs fields, again plus operators with fewer derivatives. The first relation is

$$\begin{aligned} H^\dagger H (D_\mu H) \bar{\psi} \sigma^{\mu\nu} D_\nu \psi &= \frac{i}{2} H^\dagger H (D_\mu H) \bar{\psi} (\gamma^\mu \not{D} - \not{D} \gamma^\mu) \psi \\ &= i H^\dagger H (D_\mu H) \bar{\psi} \gamma^\mu \not{D} \psi - i H^\dagger H (D_\mu H) \bar{\psi} D^\mu \psi \\ &= -i H^\dagger H (D_\mu H) \bar{\psi} D^\mu \psi + \boxed{\psi^2 H^4 D} + \boxed{E}, \end{aligned} \quad (3.10)$$

where \boxed{E} represents operators that vanish via the EOM. The other relation we need is

$$\begin{aligned} 2H^\dagger H (D_\mu H) \bar{\psi} D^\mu \psi &= H^\dagger H (D_\mu H) \bar{\psi} (\gamma^\mu \not{D} + \not{D} \gamma^\mu) \psi \\ &= \left(H^\dagger H (D_\mu H) \bar{\psi} \gamma^\mu \not{D} \psi - H^\dagger H (D_\mu H) (\not{D} \bar{\psi}) \gamma^\mu \psi \right. \\ &\quad \left. - D_\nu [H^\dagger H (D_\mu H)] \bar{\psi} \gamma^\nu \gamma^\mu \psi + \boxed{T} \right) \\ &= -D_\nu [H^\dagger H (D_\mu H)] \bar{\psi} \gamma^\nu \gamma^\mu \psi + \boxed{E} + \boxed{T} \end{aligned} \quad (3.11)$$

where \boxed{T} stands for a total derivative.

After all this we are left with six terms +h.c. where the derivatives act only on Higgs fields. In particular, having already established that operators in this class cannot have two derivatives acting on the same field, the derivatives can either act on H and H^\dagger or there can be one derivative on each of the H fields. For each of these cases the fermion pair can either be in the (0,0) or (0,1) representation of the Lorentz group. (The classification for the Hermitian conjugate operators proceeds in an identical fashion.) Finally when the derivatives act on H and H^\dagger the covariants can either be $SU(2)_w$ singlets or adjoints. The same logic applies for all three choices for the pair of fermions.

3.3 Four-Fermion Operators

18. $\psi^4 H^2$

All 38 of the ψ^4 dimension-6 terms can be multiplied by $(H^\dagger H)$. Focusing on B preserving operators, an additional 23 terms are formed by inserting a τ^I into a ψ^4 operator (with at least two left-handed fermions) and joining it to the dimension-2 covariant $(H^\dagger \tau^I H)$. There are some redundant operators involving identical left-handed fermions. For example, the terms

$$\begin{aligned} Q_{l^4 H^2}^{(3)} &= (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{l}_s \gamma_\mu \tau^I l_t) (H^\dagger H), \\ Q_{l^4 H^2}^{(4)} &= (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{l}_s \gamma_\mu l_t) (H^\dagger \tau^I H), \\ Q_{l^4 H^2}^{(5)} &= \epsilon^{IJK} (\bar{l}_p \gamma^\mu \tau^I l_r) (\bar{l}_s \gamma_\mu \tau^J \tau^K l_t) (H^\dagger \tau^K H), \end{aligned} \quad (3.12)$$

are related to the operators in our basis

$$\begin{aligned} Q_{l^4 H^2}^{(3)} &= 2Q_{l^4 H^2}^{(1)} - Q_{l^4 H^2}^{(1)}_{ptsr}, \\ Q_{l^4 H^2}^{(4)} &= Q_{l^4 H^2}^{(2)}_{stpr}, \\ iQ_{l^4 H^2}^{(5)} &= Q_{l^4 H^2}^{(2)}_{prst} - Q_{l^4 H^2}^{(2)}_{stpr}. \end{aligned} \quad (3.13)$$

This can be seen using Eq. (3.1) and a variation of (3.1)

$$\delta_j^k (\tau^I)_m^n - (\tau^I)_j^k \delta_m^n = i\epsilon^{IJK} (\tau^J)_j^n (\tau^K)_m^k. \quad (3.14)$$

Operators of the type $Q_{l^2 q^2 H^2}$ provide three of these terms, whereas all other types of operators provide one term (+h.c.). There are also 22 $\psi^4 H^2$ terms that, schematically, are products of Yukawa interactions, $(\bar{L}RH)(\bar{L}RH)$ or $(\bar{L}RH)(H^\dagger \bar{R}L)$. Among these there is one new type of operator, $Q_{l^2 ud H^2}$. Bi-Yukawa terms with four fields that are fundamentals under either $SU(2)_w$ or $SU(3)_c$ where two of those fields are identical can be contracted either as singlets or adjoints.

The terms for baryon number violating operators with dimension-6 analogs follow the logic laid out for the B operators. However there are two interesting flavor structures. The dimension-6 operator Q_{qqqe} is symmetric in its q flavor indices [4]. This constraint is broken by the sum over $SU(2)_w$ adjoint indices and the Pauli matrices in $Q_{eq^2 u H^2}$, giving it full flavor rank, n_g^4 . The dimension-6 operator Q_{qqql} also has a flavor constraint [4]

$$Q_{qqql} + Q_{qqql} = Q_{qqql} + Q_{qqql}, \quad (3.15)$$

which can be derived using Eq. (3.2). The operators $Q_{lq^3 H^2}^{(1,2)}$ respect this constraint, leading to each of their Lagrangian terms (+h.c.) containing $\frac{2}{3}n_g^2(2n_g^2 + 1)$ operators. On the other hand, $Q_{lq^3 H^2}^{(3)}$ has mixed symmetry. It is symmetric in p and r and anti-symmetric in r and s , which causes it to vanish when there is only one generation of

fermions. As a result of the six fundamental $SU(2)_w$ indices there are eight redundant operators

$$\begin{aligned}
Q_{lq^3H^2}^{(1a)} &= \epsilon_{\alpha\beta\gamma}\epsilon_{mj}\epsilon_{kn}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger H), \\
Q_{lq^3H^2}^{(1b)} &= \epsilon_{\alpha\beta\gamma}(\tau^I\epsilon)_{mj}(\tau^I\epsilon)_{kn}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger H), \\
Q_{lq^3H^2}^{(2a)} &= \epsilon_{\alpha\beta\gamma}\epsilon_{mj}(\tau^I\epsilon)_{kn}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^I H), \\
Q_{lq^3H^2}^{(2b)} &= \epsilon_{\alpha\beta\gamma}(\tau^I\epsilon)_{jn}\epsilon_{km}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^I H), \\
Q_{lq^3H^2}^{(3a)} &= \epsilon_{\alpha\beta\gamma}(\tau^I\epsilon)_{mj}\epsilon_{kn}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^I H), \\
Q_{lq^3H^2}^{(3b)} &= \epsilon_{\alpha\beta\gamma}\epsilon_{jn}(\tau^I\epsilon)_{km}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^I H), \\
Q_{lq^3H^2}^{(4a)} &= \epsilon_{\alpha\beta\gamma}\epsilon^{IJK}(\tau^I\epsilon)_{mn}(\tau^J\epsilon)_{jk}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^K H), \\
Q_{lq^3H^2}^{(4b)} &= \epsilon_{\alpha\beta\gamma}\epsilon^{IJK}(\tau^I\epsilon)_{mj}(\tau^J\epsilon)_{kn}(q_p^{m\alpha}Cq_r^{j\beta})(q_s^{k\gamma}Cl_t^n)(H^\dagger\tau^K H).
\end{aligned} \tag{3.16}$$

The operators in (3.16) can be written in terms of the operators in our basis using Eq. (3.1), Eq. (3.14), and the following relations obtained from Eq. (3.1)

$$\begin{aligned}
&\epsilon_{mj}(\tau^I\epsilon)_{kn} + (\tau^I\epsilon)_{mj}\epsilon_{kn} = \epsilon_{mn}(\tau^I\epsilon)_{jk} - (\tau^I\epsilon)_{mn}\epsilon_{jk}, \\
&i\epsilon^{IJK}[(\tau^J\epsilon)_{mn}(\tau^K\epsilon)_{jk} - \frac{1}{2}(\tau^J\epsilon)_{mj}(\tau^K\epsilon)_{kn}] = \epsilon_{mn}(\tau^I\epsilon)_{jk} + (\tau^I\epsilon)_{mn}\epsilon_{jk}.
\end{aligned} \tag{3.17}$$

In particular, the relations are

$$\begin{aligned}
-Q_{lq^3H^2}^{(1a)} &= Q_{lq^3H^2}^{(1)}_{prst} + Q_{lq^3H^2}^{(1)}_{rpst}, \\
-Q_{lq^3H^2}^{(1b)} &= Q_{lq^3H^2}^{(1)}_{prst} - Q_{lq^3H^2}^{(1)}_{rpst}, \\
Q_{lq^3H^2}^{(2b)} &= Q_{lq^3H^2}^{(2)}_{rpst}, \\
Q_{lq^3H^2}^{(3b)} &= -Q_{lq^3H^2}^{(3)}_{rpst}, \\
2Q_{lq^3H^2}^{(2a)} &= (Q_{lq^3H^2}^{(2)}_{prst} + Q_{lq^3H^2}^{(2)}_{rpst}) - (Q_{lq^3H^2}^{(3)}_{prst} + Q_{lq^3H^2}^{(3)}_{rpst}), \\
2Q_{lq^3H^2}^{(3a)} &= (Q_{lq^3H^2}^{(2)}_{prst} - Q_{lq^3H^2}^{(2)}_{rpst}) - (Q_{lq^3H^2}^{(3)}_{prst} - Q_{lq^3H^2}^{(3)}_{rpst}), \\
Q_{lq^3H^2}^{(4a)} &= Q_{lq^3H^2}^{(2)}_{rpst} - Q_{lq^3H^2}^{(3)}_{rpst}, \\
\frac{1}{2}Q_{lq^3H^2}^{(4b)} &= -(Q_{lq^3H^2}^{(2)}_{prst} - Q_{lq^3H^2}^{(2)}_{rpst}) - (Q_{lq^3H^2}^{(3)}_{prst} + Q_{lq^3H^2}^{(3)}_{rpst}).
\end{aligned} \tag{3.18}$$

In addition to \mathcal{B} operators with dimension-6 analogs, three new types of operators appear. These were three of the types of operators identified by Ref. [10] as types that vanish in the absence of flavor structure. All three contain a Lorentz singlet pair of quarks in the antisymmetric $\bar{3}$ representation of $SU(3)_c$, yielding $n_g^3(n_g - 1)$ independent operators. The operators of type $Q_{lq^3H^2}$ are different from these

three (and others identified by [10]) in that there is at least one Lagrangian term in the absence of flavor. However not all of the terms are present in the absence of flavor structure. Dimension-8 is the lowest mass dimension where this happens. The vanishing of operators in the absence of flavor structure first occurs at dimension-7.

19. $\psi^4 X$

For a pair of currents there are 114 terms formed an operator by contracting the currents with a field strength, and inserting $SU(2)_w$ and $SU(3)_c$ generators and invariants as necessary. There are at least two terms per JJX operator type, one from X_L and one from X_R . The largest number of terms is eight, which occurs for operator types $Q_{u^2 d^2 G}$, $Q_{q^2 u^2 G}$, and $Q_{q^2 d^2 G}$, where the $SU(3)_c$ combinations are $(8 \otimes 1 \otimes 8)$, $(1 \otimes 8 \otimes 8)$, $(8 \otimes 8 \otimes 8)_A$, and $(8 \otimes 8 \otimes 8)_S$ for the, say, u current, d current and G , respectively.

There are five types of operators that were identified in [10] involving identical currents contracted with the hypercharge field strength, *e.g.* $(\bar{l}\gamma^\mu l)(\bar{l}\gamma^\nu l)B_{\mu\nu}$, which vanish in the absence of flavor structure as the contraction forces the fermions into an anti-symmetric flavor representation. For each Lagrangian term the number of operators is

$$\left(\overline{\square} \otimes \square \otimes \overline{\square} \otimes \square\right)_A = adj \oplus adj \oplus \bar{a}s \oplus \bar{s}a = \frac{1}{2}n_g^2(n_g^2 - 1) \quad (3.19)$$

The relevant group theory results can be found in *e.g.* [26]. The electron is a special case as it does not have $SU(2)_w$ or $SU(3)_c$ indices. As such only half of the operators of this class 19 type are independent with the rest being related through a Fierz identity.

For operators with fermion chirality $(\bar{L}R)(\bar{R}L)$ there are two possibilities per field strength, one with the left-handed field strength and the other with its right-handed counterpart. For operators with fermion chirality $(\bar{L}R)(\bar{L}R)$ there are instead three choices for the Lorentz contractions, all with X_R . Two of these terms involve a tensor bilinear and a scalar bilinear while the third has two tensor bilinears. Additionally when all the fermions are quarks there are two choices for the $SU(3)_c$ contractions.

For the baryon number violating operators with two left-handed and two right-handed fermions there are two possible Lorentz contractions, one with X_L and one with X_R . When these operators involve a gluon field strength there are also two possible arrangements of the $SU(3)_c$ indices, $8 \otimes 3 \otimes 3 \otimes 3 = (3 \oplus \bar{6} \dots) \otimes 3 \otimes 3 = 1 \oplus 1 \dots$. The operators of type $Q_{eq^2 uX}$ with $X = W_R$ or B_L are in antisymmetric flavor representations for the q pair, and as such vanish in the absence of flavor structure. Instead $Q_{eq^2 uX}$ with $X = W_L$ or B_R are symmetric in p and r . The types involving gluons have full flavor rank as the q fields can either be a color $\bar{3}$ or 6, compensating for other (anti)symmetries of the operator.

On the other hand, when the fermions all have the same chirality three Lorentz contractions are possible. For operators of the type $Q_{eu^2 dB}$ the gauge contractions are fixed and it is the Lorentz contractions that dictate the flavor representation of

the u pair is, leading to one symmetric and one full rank term +h.c.. Instead for Q_{eu^2dG} type operators, the additional freedom coming from the color indices of the gluon, allowing for full flavor rank, n_g^4 , in all the Lagrangian terms. In the case of Q_{lq^3X} only two of the three Lorentz contractions are independent due to the identical q fields. There are two terms of Q_{lq^3G} operators (+h.c.) that have the same flavor representations as the their dimension-6 analog, Q_{qqql} , along with one term of Q_{lq^3W} and one term of Q_{lq^3B} , again +h.c.. For each of these types of operators there are an equal number of operator types that have mixed symmetry, $\frac{1}{3}n_g^2(n_g^2 - 1)$, that vanish in the absence of flavor structure similar to $Q_{lq^3H^2}^{(3)}$. Finally there is a third term +h.c. for the operators involving $W_{\mu\nu}^I$, $Q_{lq^3W}^{(2)}$, that is in a symmetric plus mixed flavor representation.

Operators of the type Q_{lq^3W} are another complicated case that deserve further discussion. Here the quarks can be in either the 2 or the 4 representation of both $SU(2)_w$ and the $SU(2)_L$ of the Lorentz group. Naively there are four terms (+h.c.) to consider with redundant operators handled in a similar fashion as the $Q_{lq^3H^2}$ case. However we can unambiguously combine the two terms that are Lorentz quartets into a single term, reducing the number of terms to three +h.c.. To see this consider the quark flavor symmetries of the four cases. Following Ref. [22], specifically its Table 2, we decompose the product of the gauge and Lorentz representations into irreducible representations of the permutation group of three objects, S_3 , which gives us the flavor representations of the three quarks. As we have been seen before, when the quarks are in the 2 of both $SU(2)_w$ and $SU(2)_L$ they have symmetric, mixed, and antisymmetric flavor representations. When one of the two representations is a 2 and the other is a 4 there is only a mixed flavor representation. Finally, when both representations are the 4 there is only the symmetric representation. We can combine the two terms that are in the 4 of $SU(2)_L$ into $Q_{lq^3W}^{(2)}$ as they contain distinct flavor representations. This is not unambiguously possible for $Q_{lq^3W}^{(1)}$ and $Q_{lq^3W}^{(3)}$, or $Q_{lq^3H^2}^{(2)}$ and $Q_{lq^3H^2}^{(3)}$ as each of those terms contain a mixed representation. In equations, the naive terms that are in the 4 of $SU(2)_L$ are

$$\begin{aligned} Q_{lq^3W}^{(2a)} &= \epsilon_{\alpha\beta\gamma}(\tau^I \epsilon)_{mn} \epsilon_{jk} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) W_{\mu\nu}^I, \\ Q_{lq^3W}^{(2b)} &= \epsilon_{\alpha\beta\gamma} \epsilon_{mn} (\tau^I \epsilon)_{jk} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) W_{\mu\nu}^I. \end{aligned} \quad (3.20)$$

They can be combined as

$$2Q_{lq^3W}^{(2)} = (Q_{lq^3W}^{(2a)} - Q_{lq^3W}^{(2a)}) - (Q_{lq^3W}^{(2b)} - Q_{lq^3W}^{(2b)}). \quad (3.21)$$

Other combinations are possible of course, but this combination makes it clear there is a symmetric and a mixed flavor representation.

20. $\psi^4 HD$

In class 20 the operators either have one fermion transforming as $(\frac{1}{2}, 0)$ and three transforming as $(0, \frac{1}{2})$ under the Lorentz group, or vice versa. When describing the classification of this class we assume the former case. Then, from (3.4) the derivative cannot act on the left-handed fermion. Otherwise the operator would be reduced by the EOM.

We start with the baryon number conserving operators. Consider the case when the four fermion fields are distinguishable, *e.g.* \bar{d}, d, \bar{l}, e (with conjugate fields are counted separately). Here there are two independent Lorentz contractions when the derivative acts on the Higgs field. A third Lorentz structure is related to the first two by Eq. (3.2). When the derivative acts on a fermion there is instead only one possible Lorentz contraction. We use type Q_{led^2HD} as an example. It has the five aforementioned candidate terms

$$\begin{aligned}
x_1 &= \bar{d}_a d_{\dot{a}} \bar{l}_{\dot{b}} e_{\dot{c}} (DH)_{b\dot{d}} \epsilon^{ab} \epsilon^{\dot{a}\dot{b}} \epsilon^{\dot{c}\dot{d}}, \\
x_2 &= \bar{d}_a d_{\dot{a}} \bar{l}_{\dot{b}} e_{\dot{c}} (DH)_{b\dot{d}} \epsilon^{ab} \epsilon^{\dot{a}\dot{d}} \epsilon^{\dot{c}\dot{b}}, \\
x_3 &= \bar{d}_a (Dd)_{b(\dot{a}\dot{d})} \bar{l}_{\dot{b}} e_{\dot{c}} H \frac{1}{2} \epsilon^{ab} (2\epsilon^{\dot{a}\dot{b}} \epsilon^{\dot{c}\dot{d}} - \epsilon^{\dot{a}\dot{d}} \epsilon^{\dot{c}\dot{b}}), \\
x_4 &= \bar{d}_a d_{\dot{a}} (D\bar{l})_{a(\dot{b}\dot{d})} e_{\dot{c}} H \frac{1}{2} \epsilon^{ab} (\epsilon^{\dot{a}\dot{b}} \epsilon^{\dot{c}\dot{d}} + \epsilon^{\dot{a}\dot{d}} \epsilon^{\dot{c}\dot{b}}), \\
x_5 &= \bar{d}_a d_{\dot{a}} l_{\dot{b}} (De)_{a(\dot{c}\dot{d})} H \frac{1}{2} \epsilon^{ab} \epsilon^{\dot{a}\dot{b}} \epsilon^{\dot{c}\dot{d}},
\end{aligned} \tag{3.22}$$

where Eq. (3.2) is used to remove redundant Lorentz structures. There are two constraint equations

$$\begin{aligned}
Dy_1 &= x_1 + x_3 + \frac{1}{2}x_4 + \frac{1}{2}x_5 = 0, \\
Dy_2 &= x_2 - x_3 + \frac{1}{2}x_4 = 0,
\end{aligned} \tag{3.23}$$

and we choose to keep x_1 , x_2 , and x_4 in our basis. For other types of operators where all four fermions are distinguishable we keep the analogs of x_1 , x_2 , and x_4 as well. In addition, if there are four fields in the operator that are fundamentals under $SU(2)_w$ or $SU(3)_c$, including the Higgs, then there are two possible contractions of those gauge indices for each possible Lorentz contraction.

The other possibility is that only three of the fermion fields are unique, *e.g.* \bar{d}, d, \bar{q}, d . In this case there is only one way to contract the Lorentz indices when the derivative acts on the Higgs field, and only two possible ways to assign the derivative to fermions. When the derivative acts on the Higgs field or the repeated fermion there are four, two, and one possible gauge contractions when the repeated fermion is q , one of $\{l, u, d\}$, or e , respectively. Instead when the derivative acts on the fermion is not repeated, *e.g.* $\bar{d}, d, (D\bar{q}), d$, there are two possible gauge contractions if the repeated fermion is q and only one otherwise. Here the repeated fermion is in a symmetric Lorentz representation and so the gauge contractions must be antisymmetric, eliminating half the possibilities.

The baryon number violating operators follow the same rules. There are a couple of non-trivial flavor cases. When there are duplicate right-handed fermions they form

a symmetric flavor representation if the derivative acts on the Higgs field, and have full flavor rank if the derivative acts on the one of the duplicate fermions. For the term Q_{eq^3HD} , the derivative acts on one of the q fields, breaking the flavor constrain, Eq. (3.15), giving it full flavor rank.

21. $\psi^4 D^2$

We start with operators with equal numbers of left- and right-handed fermions, both B and \bar{B} . There are two ways to assign the derivatives to the fields that are not related by IBP. The first is to assign the derivatives to fermions that have the same Lorentz representation, and second is to assign the derivatives to fermions with conjugate Lorentz representations. The only exception to this is when both currents are electron currents in which case there is only one independent term. As previously mentioned this is due to the fact that the electron is a singlet both $SU(2)_w$ and $SU(3)_c$. As this work was being completed Ref. [23] appeared, which classified nine terms from class 19. Some of their operators subsume both derivatives into the d'Alembertian operator. Our logic is consistent with the results of Ref. [23]. The difference is we use relations like (3.4) to reduce operators with two derivatives acting on the same fermion to classes with fewer derivatives. As such the $\psi^4 D^2$ operators in our basis where derivatives act symmetrically take the form $D_\mu(\psi_1 \Gamma \psi_2) D^\mu(\psi_3 \Gamma \psi_4)$ where Γ is some, possibly scalar, combination of gamma matrices. The remaining 30 current-current terms and the four terms with chirality $(\bar{L}R)(\bar{R}L)$ in class 19 have a form analogous to the operators classified by Ref. [23]. Also, the derivatives in $Q_{eq^2uD^2}$ break the flavor constraint present in its dimension-6 analog Q_{qqe} .

For operators with either all left-handed or all right-handed fermions there are two possible Lorentz contractions, one where the fermions without derivatives form a scalar and a second where they form a tensor. However through IBP and using (3.10) only one is independent. There are six ways to assign the derivatives to the operators with chirality $(\bar{L}R)(\bar{L}R)$, three of which are independent after IBP. For operators of type Q_{lequD^2} we choose the derivatives assignments to correspond to the exchange of heavy scalars with representations $(1, 2)_{1/2}$, $(3, 2)_{7/6}$, and $(3, 1)_{-1/3}$, respectively. An example of an IBP constraint equation for these types of operators is

$$Q_{lequD^2}^{(1)} + 2(D_\mu \bar{l}_p^j D^\mu e_r) \epsilon_{jk} (\bar{q}_s^k u_t) = \boxed{E}. \quad (3.24)$$

Similarly, for operators of type $Q_{q^2udD^2}$ we choose to assign derivatives to correspond to the exchange of heavy scalars with gauge representations $(1, 2)_{1/2}$, $(8, 2)_{1/2}$, and $(3, 1)_{-1/3}$ (or $(6, 1)_{1/3}$), respectively. There is a single term (+h.c.) for the type $Q_{lq^3D^2}$. As was the case with $Q_{eq^2uD^2}$, the derivatives in $Q_{lq^3D^2}$ break the flavor constraint present in some other operators of type $Q_{lq^3...}$, allowing the term to have full flavor rank. On the other hand, for the first time we encounter a term of the type $Q_{eu^2d...}$ that vanishes in the absence of flavor structure. There is also a second type of $Q_{eu^2dD^2}$ operator (+h.c.) that has full flavor rank.

4 The Complete Set of Dimension-8 Operators

Having gone through our classification of the dimension-8 operators in the previous Section we are now ready to tabulate the results. Table 1 summarizes the results tables that follow it. The links in the rightmost column point to the table(s) of results for a given class. The number of types of operators and the number of Lagrangian terms in the class are given in the third and fourth columns from the left, respectively. For comparison the number of operators from Ref. [10] is given in the second column from the right. Lines separate the classes based on the number of fermions in the class. Four-fermion operators are further divided into subclasses either preserving or violating baryon number. Additionally the number to the right of the + sign in $N_{\text{type,term}}$ for the four-fermion operators is the number of types or terms that vanish in the absence of flavor structure. Tables 2 and 3 contain bosonic operators. Tables 4, 5, 6, 7, 8, and 9 contain two-fermion operators. Tables 10, 11, 12, 13, 14, 15, 16, 17, and 18 contain four-fermions operators.

#	Class	N_{type}	N_{term}	N_{op} [10]	Table(s)
1	X^4	7	43	43	2
2	H^8	1	1	1	2
3	$H^6 D^2$	1	2	2	2
4	$H^4 D^4$	1	3	3	2
5	$X^3 H^2$	3	6	6	3
6	$X^2 H^4$	5	10	10	3
7	$X^2 H^2 D^2$	4	18	18	3
8	$X H^4 D^2$	2	6	6	3
9	$\psi^2 X^2 H$	16	96	$96n_g^2$	4
10	$\psi^2 X H^3$	8	22	$22n_g^2$	5
11	$\psi^2 H^2 D^3$	6	16	$16n_g^2$	5
12	$\psi^2 H^5$	3	6	$6n_g^2$	5
13	$\psi^2 H^4 D$	6	13	$13n_g^2$	5
14	$\psi^2 X^2 D$	21	57	$57n_g^2$	6, 7
15	$\psi^2 X H^2 D$	16	92	$92n_g^2$	7, 8
16	$\psi^2 X H D^2$	8	48	$48n_g^2$	9
17	$\psi^2 H^3 D^2$	3	36	$36n_g^2$	9
18(B)	$\psi^4 H^2$	19	75	$n_g^2(67n_g^2 + n_g + 7)$	10, 11
18(\cancel{B})		4 + 3	12 + 8	$\frac{1}{3}n_g^2(43n_g^2 - 9n_g + 2)$	10
19(B)	$\psi^4 X$	40 + 5	156 + 12	$4n_g^2(40n_g^2 - 1)$	12, 13, 14
19(\cancel{B})		4	44 + 12	$2n_g^3(21n_g + 1)$	15
20(B)	$\psi^4 H D$	16	134 + 2	$n_g^3(135n_g - 1)$	16, 17
20(\cancel{B})		7	32	$n_g^3(29n_g + 3)$	17
21(B)	$\psi^4 D^2$	18	55	$\frac{11}{2}n_g^2(9n_g^2 + 1)$	10, 18
21(\cancel{B})		4	10 + 2	$n_g^3(11n_g - 1)$	10
	B	204 + 5	895 + 14	895(36971), $n_g = 1(3)$	
	\cancel{B}	19 + 3	98 + 22	98(7836), $n_g = 1(3)$	
	Total	223 + 8	993 + 36	993(44807), $n_g = 1(3)$	

Table 1. Summary of the contents of the tables to follow. The links in the rightmost column point to the table(s) of results for a given class. The number of types of operators and the number of Lagrangian terms in the class are given in the third and fourth columns from the left, respectively. For comparison the number of operators from Ref. [10] is given in the second column from the right. Lines separate the classes based on the number of fermions in the class. Four-fermion operators are further divided into subclasses either preserving or violating baryon number. Additionally the number to the right of the + sign in $N_{\text{type(term)}}$ for the four-fermion operators is the number of types(terms) that vanish in the absence of flavor structure.

4.1 Results for Bosonic Operators

$1 : X^4, X^3 X'$		$1 : X^2 X'^2$	
$Q_{G^4}^{(1)}$	$(G_{\mu\nu}^A G^{A\mu\nu})(G_{\rho\sigma}^B G^{B\rho\sigma})$	$Q_{G^2 W^2}^{(1)}$	$(W_{\mu\nu}^I W^{I\mu\nu})(G_{\rho\sigma}^A G^{A\rho\sigma})$
$Q_{G^4}^{(2)}$	$(G_{\mu\nu}^A \tilde{G}^{A\mu\nu})(G_{\rho\sigma}^B \tilde{G}^{B\rho\sigma})$	$Q_{G^2 W^2}^{(2)}$	$(W_{\mu\nu}^I \tilde{W}^{I\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{A\rho\sigma})$
$Q_{G^4}^{(3)}$	$(G_{\mu\nu}^A G^{B\mu\nu})(G_{\rho\sigma}^A G^{B\rho\sigma})$	$Q_{G^2 W^2}^{(3)}$	$(W_{\mu\nu}^I G^{A\mu\nu})(W_{\rho\sigma}^I G^{A\rho\sigma})$
$Q_{G^4}^{(4)}$	$(G_{\mu\nu}^A \tilde{G}^{B\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{B\rho\sigma})$	$Q_{G^2 W^2}^{(4)}$	$(W_{\mu\nu}^I \tilde{G}^{A\mu\nu})(W_{\rho\sigma}^I \tilde{G}^{A\rho\sigma})$
$Q_{G^4}^{(5)}$	$(G_{\mu\nu}^A G^{A\mu\nu})(G_{\rho\sigma}^B \tilde{G}^{B\rho\sigma})$	$Q_{G^2 W^2}^{(5)}$	$(W_{\mu\nu}^I \tilde{W}^{I\mu\nu})(G_{\rho\sigma}^A G^{A\rho\sigma})$
$Q_{G^4}^{(6)}$	$(G_{\mu\nu}^A G^{B\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{B\rho\sigma})$	$Q_{G^2 W^2}^{(6)}$	$(W_{\mu\nu}^I W^{I\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{A\rho\sigma})$
$Q_{G^4}^{(7)}$	$d^{ABE} d^{CDE} (G_{\mu\nu}^A G^{B\mu\nu})(G_{\rho\sigma}^C G^{D\rho\sigma})$	$Q_{G^2 W^2}^{(7)}$	$(W_{\mu\nu}^I G^{A\mu\nu})(W_{\rho\sigma}^I \tilde{G}^{A\rho\sigma})$
$Q_{G^4}^{(8)}$	$d^{ABE} d^{CDE} (G_{\mu\nu}^A \tilde{G}^{B\mu\nu})(G_{\rho\sigma}^C \tilde{G}^{D\rho\sigma})$	$Q_{G^2 B^2}^{(1)}$	$(B_{\mu\nu} B^{\mu\nu})(G_{\rho\sigma}^A G^{A\rho\sigma})$
$Q_{G^4}^{(9)}$	$d^{ABE} d^{CDE} (G_{\mu\nu}^A G^{B\mu\nu})(G_{\rho\sigma}^C \tilde{G}^{D\rho\sigma})$	$Q_{G^2 B^2}^{(2)}$	$(B_{\mu\nu} \tilde{B}^{\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{A\rho\sigma})$
$Q_{W^4}^{(1)}$	$(W_{\mu\nu}^I W^{I\mu\nu})(W_{\rho\sigma}^J W^{J\rho\sigma})$	$Q_{G^2 B^2}^{(3)}$	$(B_{\mu\nu} G^{A\mu\nu})(B_{\rho\sigma} G^{A\rho\sigma})$
$Q_{W^4}^{(2)}$	$(W_{\mu\nu}^I \tilde{W}^{I\mu\nu})(W_{\rho\sigma}^J \tilde{W}^{J\rho\sigma})$	$Q_{G^2 B^2}^{(4)}$	$(B_{\mu\nu} \tilde{G}^{A\mu\nu})(B_{\rho\sigma} \tilde{G}^{A\rho\sigma})$
$Q_{W^4}^{(3)}$	$(W_{\mu\nu}^I W^{J\mu\nu})(W_{\rho\sigma}^I W^{J\rho\sigma})$	$Q_{G^2 B^2}^{(5)}$	$(B_{\mu\nu} \tilde{B}^{\mu\nu})(G_{\rho\sigma}^A G^{A\rho\sigma})$
$Q_{W^4}^{(4)}$	$(W_{\mu\nu}^I \tilde{W}^{J\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{J\rho\sigma})$	$Q_{G^2 B^2}^{(6)}$	$(B_{\mu\nu} B^{\mu\nu})(G_{\rho\sigma}^A \tilde{G}^{A\rho\sigma})$
$Q_{W^4}^{(5)}$	$(W_{\mu\nu}^I W^{I\mu\nu})(W_{\rho\sigma}^J \tilde{W}^{J\rho\sigma})$	$Q_{G^2 B^2}^{(7)}$	$(B_{\mu\nu} G^{A\mu\nu})(B_{\rho\sigma} \tilde{G}^{A\rho\sigma})$
$Q_{W^4}^{(6)}$	$(W_{\mu\nu}^I W^{J\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{J\rho\sigma})$	$Q_{W^2 B^2}^{(1)}$	$(B_{\mu\nu} B^{\mu\nu})(W_{\rho\sigma}^I W^{I\rho\sigma})$
$Q_{B^4}^{(1)}$	$(B_{\mu\nu} B^{\mu\nu})(B_{\rho\sigma} B^{\rho\sigma})$	$Q_{W^2 B^2}^{(2)}$	$(B_{\mu\nu} \tilde{B}^{\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{I\rho\sigma})$
$Q_{B^4}^{(2)}$	$(B_{\mu\nu} \tilde{B}^{\mu\nu})(B_{\rho\sigma} \tilde{B}^{\rho\sigma})$	$Q_{W^2 B^2}^{(3)}$	$(B_{\mu\nu} W^{I\mu\nu})(B_{\rho\sigma} W^{I\rho\sigma})$
$Q_{B^4}^{(3)}$	$(B_{\mu\nu} B^{\mu\nu})(B_{\rho\sigma} \tilde{B}^{\rho\sigma})$	$Q_{W^2 B^2}^{(4)}$	$(B_{\mu\nu} \tilde{W}^{I\mu\nu})(B_{\rho\sigma} \tilde{W}^{I\rho\sigma})$
$Q_{G^3 B}^{(1)}$	$d^{ABC} (B_{\mu\nu} G^{A\mu\nu})(G_{\rho\sigma}^B G^{C\rho\sigma})$	$Q_{W^2 B^2}^{(5)}$	$(B_{\mu\nu} \tilde{B}^{\mu\nu})(W_{\rho\sigma}^I W^{I\rho\sigma})$
$Q_{G^3 B}^{(2)}$	$d^{ABC} (B_{\mu\nu} \tilde{G}^{A\mu\nu})(G_{\rho\sigma}^B \tilde{G}^{C\rho\sigma})$	$Q_{W^2 B^2}^{(6)}$	$(B_{\mu\nu} B^{\mu\nu})(W_{\rho\sigma}^I \tilde{W}^{I\rho\sigma})$
$Q_{G^3 B}^{(3)}$	$d^{ABC} (B_{\mu\nu} \tilde{G}^{A\mu\nu})(G_{\rho\sigma}^B G^{C\rho\sigma})$	$Q_{W^2 B^2}^{(7)}$	$(B_{\mu\nu} W^{I\mu\nu})(B_{\rho\sigma} \tilde{W}^{I\rho\sigma})$
$Q_{G^3 B}^{(4)}$	$d^{ABC} (B_{\mu\nu} G^{A\mu\nu})(G_{\rho\sigma}^B \tilde{G}^{C\rho\sigma})$		
$2 : H^8$		$3 : H^6 D^2$	
Q_{H^8}	$(H^\dagger H)^4$	$Q_{H^6}^{(1)}$	$(H^\dagger H)^2 (D_\mu H^\dagger D^\mu H)$
		$Q_{H^6}^{(2)}$	$(H^\dagger H)(H^\dagger \tau^I H)(D_\mu H^\dagger \tau^I D^\mu H)$
		$4 : H^4 D^4$	
		$Q_{H^4}^{(1)}$	$(D_\mu H^\dagger D_\nu H)(D^\nu H^\dagger D^\mu H)$
		$Q_{H^4}^{(2)}$	$(D_\mu H^\dagger D_\nu H)(D^\mu H^\dagger D^\nu H)$
		$Q_{H^4}^{(3)}$	$(D^\mu H^\dagger D_\mu H)(D^\nu H^\dagger D_\nu H)$

Table 2. The dimension-eight operators in the SMEFT whose field content is either entirely gauge field strengths or Higgs boson fields.

5 : $X^3 H^2$			6 : $X^2 H^4$		
$Q_{G^3 H^2}^{(1)}$	$f^{ABC}(H^\dagger H)G_\mu^{A\nu}G_\nu^{B\rho}G_\rho^{C\mu}$	$Q_{G^2 H^4}^{(1)}$	$(H^\dagger H)^2 G_{\mu\nu}^A G^{A\mu\nu}$		
$Q_{G^3 H^2}^{(2)}$	$f^{ABC}(H^\dagger H)G_\mu^{A\nu}G_\nu^{B\rho}\tilde{G}_\rho^{C\mu}$	$Q_{G^2 H^4}^{(2)}$	$(H^\dagger H)^2 \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$		
$Q_{W^3 H^2}^{(1)}$	$\epsilon^{IJK}(H^\dagger H)W_\mu^{I\nu}W_\nu^{J\rho}W_\rho^{K\mu}$	$Q_{W^2 H^4}^{(1)}$	$(H^\dagger H)^2 W_{\mu\nu}^I W^{I\mu\nu}$		
$Q_{W^3 H^2}^{(2)}$	$\epsilon^{IJK}(H^\dagger H)W_\mu^{I\nu}W_\nu^{J\rho}\tilde{W}_\rho^{K\mu}$	$Q_{W^2 H^4}^{(2)}$	$(H^\dagger H)^2 \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$		
$Q_{W^2 B H^2}^{(1)}$	$\epsilon^{IJK}(H^\dagger \tau^I H)B_\mu^\nu W_\nu^{J\rho}W_\rho^{K\mu}$	$Q_{W^2 H^4}^{(3)}$	$(H^\dagger \tau^I H)(H^\dagger \tau^J H)W_{\mu\nu}^I W^{J\mu\nu}$		
$Q_{W^2 B H^2}^{(2)}$	$\epsilon^{IJK}(H^\dagger \tau^I H)(\tilde{B}^{\mu\nu}W_{\nu\rho}^J W_\mu^{K\rho} + B^{\mu\nu}W_{\nu\rho}^J \tilde{W}_\mu^{K\rho})$	$Q_{W^2 H^4}^{(4)}$	$(H^\dagger \tau^I H)(H^\dagger \tau^J H)\tilde{W}_{\mu\nu}^I W^{J\mu\nu}$		
		$Q_{WBH^4}^{(1)}$	$(H^\dagger H)(H^\dagger \tau^I H)W_{\mu\nu}^I B^{\mu\nu}$		
		$Q_{WBH^4}^{(2)}$	$(H^\dagger H)(H^\dagger \tau^I H)\tilde{W}_{\mu\nu}^I B^{\mu\nu}$		
		$Q_{B^2 H^4}^{(1)}$	$(H^\dagger H)^2 B_{\mu\nu} B^{\mu\nu}$		
		$Q_{B^2 H^4}^{(2)}$	$(H^\dagger H)^2 \tilde{B}_{\mu\nu} B^{\mu\nu}$		
7 : $X^2 H^2 D^2$			8 : $X H^4 D^2$		
$Q_{G^2 H^2 D^2}^{(1)}$	$(D^\mu H^\dagger D^\nu H)G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{WH^4 D^2}^{(1)}$	$(H^\dagger H)(D^\mu H^\dagger \tau^I D^\nu H)W_{\mu\nu}^I$		
$Q_{G^2 H^2 D^2}^{(2)}$	$(D^\mu H^\dagger D_\mu H)G_{\nu\rho}^A G^{\nu\rho}$	$Q_{WH^4 D^2}^{(2)}$	$(H^\dagger H)(D^\mu H^\dagger \tau^I D^\nu H)\tilde{W}_{\mu\nu}^I$		
$Q_{G^2 H^2 D^2}^{(3)}$	$(D^\mu H^\dagger D_\mu H)G_{\nu\rho}^A \tilde{G}^{A\nu\rho}$	$Q_{WH^4 D^2}^{(3)}$	$\epsilon^{IJK}(H^\dagger \tau^I H)(D^\mu H^\dagger \tau^J D^\nu H)W_{\mu\nu}^K$		
$Q_{W^2 H^2 D^2}^{(1)}$	$(D^\mu H^\dagger D^\nu H)W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{WH^4 D^2}^{(4)}$	$\epsilon^{IJK}(H^\dagger \tau^I H)(D^\mu H^\dagger \tau^J D^\nu H)\tilde{W}_{\mu\nu}^K$		
$Q_{W^2 H^2 D^2}^{(2)}$	$(D^\mu H^\dagger D_\mu H)W_{\nu\rho}^I W^{\nu\rho}$	$Q_{BH^4 D^2}^{(1)}$	$(H^\dagger H)(D^\mu H^\dagger D^\nu H)B_{\mu\nu}$		
$Q_{W^2 H^2 D^2}^{(3)}$	$(D^\mu H^\dagger D_\mu H)W_{\nu\rho}^I \tilde{W}^{I\nu\rho}$	$Q_{BH^4 D^2}^{(2)}$	$(H^\dagger H)(D^\mu H^\dagger D^\nu H)\tilde{B}_{\mu\nu}$		
$Q_{W^2 H^2 D^2}^{(4)}$	$\epsilon^{IJK}(D^\mu H^\dagger \tau^I D^\nu H)W_{\mu\rho}^J W_\nu^{K\rho}$				
$Q_{W^2 H^2 D^2}^{(5)}$	$\epsilon^{IJK}(D^\mu H^\dagger \tau^I D^\nu H)(W_{\mu\rho}^J \tilde{W}_\nu^{K\rho} - \tilde{W}_{\mu\rho}^J W_\nu^{K\rho})$				
$Q_{W^2 H^2 D^2}^{(6)}$	$\epsilon^{IJK}(D^\mu H^\dagger \tau^I D^\nu H)(W_{\mu\rho}^J \tilde{W}_\nu^{K\rho} + \tilde{W}_{\mu\rho}^J W_\nu^{K\rho})$				
$Q_{WBH^2 D^2}^{(1)}$	$(D^\mu H^\dagger \tau^I D_\mu H)B_{\nu\rho} W^{\nu\rho}$				
$Q_{WBH^2 D^2}^{(2)}$	$(D^\mu H^\dagger \tau^I D_\mu H)B_{\nu\rho} \tilde{W}^{I\nu\rho}$				
$Q_{WBH^2 D^2}^{(3)}$	$(D^\mu H^\dagger \tau^I D^\nu H)(B_{\mu\rho} W_\nu^{I\rho} - B_{\mu\rho} W_\nu^{I\rho})$				
$Q_{WBH^2 D^2}^{(4)}$	$(D^\mu H^\dagger \tau^I D^\nu H)(B_{\mu\rho} W_\nu^{I\rho} + B_{\mu\rho} W_\nu^{I\rho})$				
$Q_{WBH^2 D^2}^{(5)}$	$(D^\mu H^\dagger \tau^I D^\nu H)(B_{[\mu}^\rho \tilde{W}_{\nu]\rho}^I - \tilde{B}_{[\mu}^\rho W_{\nu]\rho}^I)$				
$Q_{WBH^2 D^2}^{(6)}$	$(D^\mu H^\dagger \tau^I D^\nu H)(B_{(\mu}^\rho \tilde{W}_{\nu)\rho}^I + \tilde{B}_{(\mu}^\rho W_{\nu)\rho}^I)$				
$Q_{B^2 H^2 D^2}^{(1)}$	$(D^\mu H^\dagger D^\nu H)B_{\mu\rho} B_\nu^\rho$				
$Q_{B^2 H^2 D^2}^{(2)}$	$(D^\mu H^\dagger D_\mu H)B_{\nu\rho} B^{\nu\rho}$				
$Q_{B^2 H^2 D^2}^{(3)}$	$(D^\mu H^\dagger D_\mu H)B_{\nu\rho} \tilde{B}^{\nu\rho}$				

Table 3. Bosonic dimension-eight operators in the SMEFT containing both gauge field strengths and Higgs boson fields.

4.2 Results for Two-Fermion Operators

$9 : \psi^2 X^2 H + \text{h.c.}$		$9 : \psi^2 X^2 H + \text{h.c.}$	
$Q_{leG^2H}^{(1)}$	$(\bar{l}_p e_r) H G_{\mu\nu}^A G^{A\mu\nu}$	$Q_{leWBH}^{(1)}$	$(\bar{l}_p e_r) \tau^I H W_{\mu\nu}^I B^{\mu\nu}$
$Q_{leG^2H}^{(2)}$	$(\bar{l}_p e_r) H \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	$Q_{leWBH}^{(2)}$	$(\bar{l}_p e_r) \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$
$Q_{leW^2H}^{(1)}$	$(\bar{l}_p e_r) H W_{\mu\nu}^I W^{I\mu\nu}$	$Q_{leWBH}^{(3)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\rho}^I B_{\nu}^{\rho}$
$Q_{leW^2H}^{(2)}$	$(\bar{l}_p e_r) H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	$Q_{leB^2H}^{(1)}$	$(\bar{l}_p e_r) H B_{\mu\nu} B^{\mu\nu}$
$Q_{leW^2H}^{(3)}$	$\epsilon^{IJK} (\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\rho}^J W_{\nu}^{K\rho}$	$Q_{leB^2H}^{(2)}$	$(\bar{l}_p e_r) H \tilde{B}_{\mu\nu} B^{\mu\nu}$
$Q_{quG^2H}^{(1)}$	$(\bar{q}_p u_r) \tilde{H} G_{\mu\nu}^A G^{A\mu\nu}$	$Q_{qdG^2H}^{(1)}$	$(\bar{q}_p d_r) H G_{\mu\nu}^A G^{A\mu\nu}$
$Q_{quG^2H}^{(2)}$	$(\bar{q}_p u_r) \tilde{H} \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	$Q_{qdG^2H}^{(2)}$	$(\bar{q}_p d_r) H \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$
$Q_{quG^2H}^{(3)}$	$d^{ABC} (\bar{q}_p T^A u_r) \tilde{H} G_{\mu\nu}^B G^{C\mu\nu}$	$Q_{qdG^2H}^{(3)}$	$d^{ABC} (\bar{q}_p T^A d_r) H G_{\mu\nu}^B G^{C\mu\nu}$
$Q_{quG^2H}^{(4)}$	$d^{ABC} (\bar{q}_p T^A u_r) \tilde{H} \tilde{G}_{\mu\nu}^B G^{C\mu\nu}$	$Q_{qdG^2H}^{(4)}$	$d^{ABC} (\bar{q}_p T^A d_r) H \tilde{G}_{\mu\nu}^B G^{C\mu\nu}$
$Q_{quG^2H}^{(5)}$	$f^{ABC} (\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\rho}^B G_{\nu}^{C\rho}$	$Q_{qdG^2H}^{(5)}$	$f^{ABC} (\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\rho}^B G_{\nu}^{C\rho}$
$Q_{quGWH}^{(1)}$	$(\bar{q}_p T^A u_r) \tau^I \tilde{H} G_{\mu\nu}^A W^{I\mu\nu}$	$Q_{qdGWH}^{(1)}$	$(\bar{q}_p T^A d_r) \tau^I H G_{\mu\nu}^A W^{I\mu\nu}$
$Q_{quGWH}^{(2)}$	$(\bar{q}_p T^A u_r) \tau^I \tilde{H} \tilde{G}_{\mu\nu}^A W^{I\mu\nu}$	$Q_{qdGWH}^{(2)}$	$(\bar{q}_p T^A d_r) \tau^I H \tilde{G}_{\mu\nu}^A W^{I\mu\nu}$
$Q_{quGWH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tau^I \tilde{H} G_{\mu\rho}^A W_{\nu}^{I\rho}$	$Q_{qdGWH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \tau^I H G_{\mu\rho}^A W_{\nu}^{I\rho}$
$Q_{quGBH}^{(1)}$	$(\bar{q}_p T^A u_r) \tilde{H} G_{\mu\nu}^A B^{\mu\nu}$	$Q_{qdGBH}^{(1)}$	$(\bar{q}_p T^A d_r) H G_{\mu\nu}^A B^{\mu\nu}$
$Q_{quGBH}^{(2)}$	$(\bar{q}_p T^A u_r) \tilde{H} \tilde{G}_{\mu\nu}^A B^{\mu\nu}$	$Q_{qdGBH}^{(2)}$	$(\bar{q}_p T^A d_r) H \tilde{G}_{\mu\nu}^A B^{\mu\nu}$
$Q_{quGBH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\rho}^A B_{\nu}^{\rho}$	$Q_{qdGBH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\rho}^A B_{\nu}^{\rho}$
$Q_{quW^2H}^{(1)}$	$(\bar{q}_p u_r) \tilde{H} W_{\mu\nu}^I W^{I\mu\nu}$	$Q_{qdW^2H}^{(1)}$	$(\bar{q}_p d_r) H W_{\mu\nu}^I W^{I\mu\nu}$
$Q_{quW^2H}^{(2)}$	$(\bar{q}_p u_r) \tilde{H} \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	$Q_{qdW^2H}^{(2)}$	$(\bar{q}_p d_r) H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$
$Q_{quW^2H}^{(3)}$	$\epsilon^{IJK} (\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\rho}^J W_{\nu}^{K\rho}$	$Q_{qdW^2H}^{(3)}$	$\epsilon^{IJK} (\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\rho}^J W_{\nu}^{K\rho}$
$Q_{quWBH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\rho}^I B_{\nu}^{\rho}$	$Q_{qdWBH}^{(1)}$	$(\bar{q}_p d_r) \tau^I H W_{\mu\nu}^I B^{\mu\nu}$
$Q_{quWBH}^{(1)}$	$(\bar{q}_p u_r) \tau^I \tilde{H} W_{\mu\nu}^I B^{\mu\nu}$	$Q_{qdWBH}^{(2)}$	$(\bar{q}_p d_r) \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$
$Q_{quWBH}^{(2)}$	$(\bar{q}_p u_r) \tau^I \tilde{H} \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	$Q_{qdWBH}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\rho}^I B_{\nu}^{\rho}$
$Q_{quB^2H}^{(1)}$	$(\bar{q}_p u_r) \tilde{H} B_{\mu\nu} B^{\mu\nu}$	$Q_{qdB^2H}^{(1)}$	$(\bar{q}_p d_r) H B_{\mu\nu} B^{\mu\nu}$
$Q_{quB^2H}^{(2)}$	$(\bar{q}_p u_r) \tilde{H} \tilde{B}_{\mu\nu} B^{\mu\nu}$	$Q_{qdB^2H}^{(2)}$	$(\bar{q}_p d_r) H \tilde{B}_{\mu\nu} B^{\mu\nu}$

Table 4. The dimension-eight operators in the SMEFT of class-9 with field content $\psi^2 X^2 H$. All of the operators have Hermitian conjugates. The subscripts p, r are weak-eigenstate indices.

10 : $\psi^2 X H^3 + \text{h.c.}$			11 : $\psi^2 H^2 D^3$		
$Q_{leWH^3}^{(1)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H (H^\dagger H) W_{\mu\nu}^I$		$Q_{l^2 H^2 D^3}^{(1)}$	$i(\bar{l}_p \gamma^\mu D^\nu l_r) (D_{(\mu} D_{\nu)} H^\dagger H)$	
$Q_{leWH^3}^{(2)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) H (H^\dagger \tau^I H) W_{\mu\nu}^I$		$Q_{l^2 H^2 D^3}^{(2)}$	$i(\bar{l}_p \gamma^\mu D^\nu l_r) (H^\dagger D_{(\mu} D_{\nu)} H)$	
Q_{leBH^3}	$(\bar{l}_p \sigma^{\mu\nu} e_r) H (H^\dagger H) B_{\mu\nu}$		$Q_{l^2 H^2 D^3}^{(3)}$	$i(\bar{l}_p \gamma^\mu \tau^I D^\nu l_r) (D_{(\mu} D_{\nu)} H^\dagger \tau^I H)$	
Q_{quGH^3}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} (H^\dagger H) G_{\mu\nu}^A$		$Q_{l^2 H^2 D^3}^{(4)}$	$i(\bar{l}_p \gamma^\mu \tau^I D^\nu l_r) (H^\dagger \tau^I D_{(\mu} D_{\nu)} H)$	
$Q_{quWH^3}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} (H^\dagger H) W_{\mu\nu}^I$		$Q_{e^2 H^2 D^3}^{(1)}$	$i(\bar{e}_p \gamma^\mu D^\nu e_r) (D_{(\mu} D_{\nu)} H^\dagger H)$	
$Q_{quWH^3}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} (H^\dagger \tau^I H) W_{\mu\nu}^I$		$Q_{e^2 H^2 D^3}^{(2)}$	$i(\bar{e}_p \gamma^\mu D^\nu e_r) (H^\dagger D_{(\mu} D_{\nu)} H)$	
Q_{quBH^3}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} (H^\dagger H) B_{\mu\nu}$		$Q_{q^2 H^2 D^3}^{(1)}$	$i(\bar{q}_p \gamma^\mu D^\nu q_r) (D_{(\mu} D_{\nu)} H^\dagger H)$	
Q_{qdGH^3}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H (H^\dagger H) G_{\mu\nu}^A$		$Q_{q^2 H^2 D^3}^{(2)}$	$i(\bar{q}_p \gamma^\mu D^\nu q_r) (H^\dagger D_{(\mu} D_{\nu)} H)$	
$Q_{qdWH^3}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H (H^\dagger H) W_{\mu\nu}^I$		$Q_{q^2 H^2 D^3}^{(3)}$	$i(\bar{q}_p \gamma^\mu \tau^I D^\nu q_r) (D_{(\mu} D_{\nu)} H^\dagger \tau^I H)$	
$Q_{qdWH^3}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) H (H^\dagger \tau^I H) W_{\mu\nu}^I$		$Q_{q^2 H^2 D^2}^{(4)}$	$i(\bar{q}_p \gamma^\mu \tau^I D^\nu q_r) (H^\dagger \tau^I D_{(\mu} D_{\nu)} H)$	
Q_{qdBH^3}	$(\bar{q}_p \sigma^{\mu\nu} d_r) H (H^\dagger H) B_{\mu\nu}$		$Q_{u^2 H^2 D^3}^{(1)}$	$i(\bar{u}_p \gamma^\mu D^\nu u_r) (D_{(\mu} D_{\nu)} H^\dagger H)$	
			$Q_{u^2 H^2 D^3}^{(2)}$	$i(\bar{u}_p \gamma^\mu D^\nu u_r) (H^\dagger D_{(\mu} D_{\nu)} H)$	
			$Q_{d^2 H^2 D^3}^{(1)}$	$i(\bar{d}_p \gamma^\mu D^\nu d_r) (D_{(\mu} D_{\nu)} H^\dagger H)$	
			$Q_{d^2 H^2 D^3}^{(2)}$	$i(\bar{d}_p \gamma^\mu D^\nu d_r) (H^\dagger D_{(\mu} D_{\nu)} H)$	
			$Q_{udH^2 D^3} + \text{h.c.}$	$i(\bar{u}_p \gamma^\mu D^\nu d_r) (\tilde{H}^\dagger D_{(\mu} D_{\nu)} H)$	
12 : $\psi^2 H^5 + \text{h.c.}$			13 : $\psi^2 H^4 D$		
Q_{leH^5}	$(H^\dagger H)^2 (\bar{l}_p e_r H)$	$Q_{l^2 H^4 D}^{(1)}$		$i(l_p \gamma^\mu l_r) (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	
Q_{quH^5}	$(H^\dagger H)^2 (\bar{q}_p u_r \tilde{H})$	$Q_{l^2 H^4 D}^{(2)}$		$i(l_p \gamma^\mu \tau^I l_r) [(H^\dagger \overleftrightarrow{D}_\mu^I H) (H^\dagger H) + (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger \tau^I H)]$	
Q_{qdH^5}	$(H^\dagger H)^2 (\bar{q}_p d_r H)$	$Q_{l^2 H^4 D}^{(3)}$		$i\epsilon^{IJK} (l_p \gamma^\mu \tau^I l_r) (H^\dagger \overleftrightarrow{D}_\mu^J H) (H^\dagger \tau^K H)$	
		$Q_{l^2 H^4 D}^{(4)}$		$\epsilon^{IJK} (l_p \gamma^\mu \tau^I l_r) (H^\dagger \tau^J H) D_\mu (H^\dagger \tau^K H)$	
		$Q_{e^2 H^4 D}$		$i(e_p \gamma^\mu e_r) (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	
		$Q_{q^2 H^4 D}^{(1)}$		$i(q_p \gamma^\mu q_r) (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	
		$Q_{q^2 H^4 D}^{(2)}$		$i(q_p \gamma^\mu \tau^I q_r) [(H^\dagger \overleftrightarrow{D}_\mu^I H) (H^\dagger H) + (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger \tau^I H)]$	
		$Q_{q^2 H^4 D}^{(3)}$		$i\epsilon^{IJK} (q_p \gamma^\mu \tau^I q_r) (H^\dagger \overleftrightarrow{D}_\mu^J H) (H^\dagger \tau^K H)$	
		$Q_{q^2 H^4 D}^{(4)}$		$\epsilon^{IJK} (q_p \gamma^\mu \tau^I q_r) (H^\dagger \tau^J H) D_\mu (H^\dagger \tau^K H)$	
		$Q_{u^2 H^4 D}$		$i(u_p \gamma^\mu u_r) (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	
		$Q_{d^2 H^4 D}$		$i(d_p \gamma^\mu d_r) (H^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	
		$Q_{udH^4 D} + \text{h.c.}$		$i(u_p \gamma^\mu d_r) (\tilde{H}^\dagger \overleftrightarrow{D}_\mu H) (H^\dagger H)$	

Table 5. The dimension-eight operators in the SMEFT of classes-10, through -13, all of which have two fermions. The operators $Q_{udH^4 D}$ and $Q_{udH^2 D^3}$ have Hermitian conjugates. The subscripts p, r are weak-eigenstate indices.

14 : $\psi^2 X^2 D$		14 : $\psi^2 X^2 D$	
$Q_{q^2 G^2 D}^{(1)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r) G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{q^2 G^2 D}^{(4)}$	$if^{ABC}(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} - \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
$Q_{q^2 G^2 D}^{(2)}$	$if^{ABC}(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 G^2 D}^{(5)}$	$if^{ABC}(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} + \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
$Q_{q^2 G^2 D}^{(3)}$	$id^{ABC}(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 GWD}^{(1)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I \overleftrightarrow{D}^\nu q_r)(G_{\mu\rho}^A W_\nu^{I\rho} - G_{\mu\rho}^A W_\nu^{I\rho})$
$Q_{q^2 W^2 D}^{(1)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r) W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{q^2 GWD}^{(2)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I \overleftrightarrow{D}^\nu q_r)(G_{\mu\rho}^A W_\nu^{I\rho} + G_{\mu\rho}^A W_\nu^{I\rho})$
$Q_{q^2 W^2 D}^{(2)}$	$i\epsilon^{IJK}(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r) W_{\mu\rho}^J W_\nu^{K\rho}$	$Q_{q^2 GWD}^{(3)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I \overleftrightarrow{D}^\nu q_r)(G_{[\mu}^{A\rho} \tilde{W}_{\nu]}^I - \tilde{G}_{[\mu}^{A\rho} W_{\nu]}^I)$
$Q_{q^2 B^2 D}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r) B_{\mu\rho} B_\nu^\rho$	$Q_{q^2 GWD}^{(4)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I \overleftrightarrow{D}^\nu q_r)(G_{(\mu}^{A\rho} \tilde{W}_{\nu)}^I + \tilde{G}_{(\mu}^{A\rho} W_{\nu)}^I)$
$Q_{u^2 G^2 D}^{(1)}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r) G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{q^2 GBD}^{(1)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(B_{\mu\rho} G_\nu^{A\rho} - B_{\mu\rho} G_\nu^{A\rho})$
$Q_{u^2 G^2 D}^{(2)}$	$if^{ABC}(\bar{u}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu u_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 GBD}^{(2)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(B_{\mu\rho} G_\nu^{A\rho} + B_{\mu\rho} G_\nu^{A\rho})$
$Q_{u^2 G^2 D}^{(3)}$	$id^{ABC}(\bar{u}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu u_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 GBD}^{(3)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(B_{[\mu}^\rho \tilde{G}_{\nu]\rho}^A - \tilde{B}_{[\mu}^\rho G_{\nu]\rho}^A)$
$Q_{u^2 W^2 D}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r) W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{q^2 GBD}^{(4)}$	$i(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(B_{(\mu}^\rho \tilde{G}_{\nu)\rho}^A + \tilde{B}_{(\mu}^\rho G_{\nu)\rho}^A)$
$Q_{u^2 B^2 D}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r) B_{\mu\rho} B_\nu^\rho$	$Q_{q^2 W^2 D}^{(3)}$	$i\epsilon^{IJK}(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(W_{\mu\rho}^J \tilde{W}_\nu^{K\rho} - \tilde{W}_{\mu\rho}^J W_\nu^{K\rho})$
$Q_{d^2 G^2 D}^{(1)}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r) G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{q^2 W^2 D}^{(4)}$	$i\epsilon^{IJK}(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(W_{\mu\rho}^J \tilde{W}_\nu^{K\rho} + \tilde{W}_{\mu\rho}^J W_\nu^{K\rho})$
$Q_{d^2 G^2 D}^{(2)}$	$if^{ABC}(\bar{d}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu d_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 WBD}^{(1)}$	$i(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(B_{\mu\rho} W_\nu^{I\rho} - B_{\mu\rho} W_\nu^{I\rho})$
$Q_{d^2 G^2 D}^{(3)}$	$id^{ABC}(\bar{d}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu d_r) G_{\mu\rho}^B G_\nu^{C\rho}$	$Q_{q^2 WBD}^{(2)}$	$i(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(B_{\mu\rho} W_\nu^{I\rho} + B_{\mu\rho} W_\nu^{I\rho})$
$Q_{d^2 W^2 D}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r) W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{q^2 WBD}^{(3)}$	$i(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(B_{[\mu}^\rho \tilde{W}_{\nu]\rho}^I - \tilde{B}_{[\mu}^\rho W_{\nu]\rho}^I)$
$Q_{d^2 B^2 D}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r) B_{\mu\rho} B_\nu^\rho$	$Q_{q^2 WBD}^{(4)}$	$i(\bar{q}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu q_r)(B_{(\mu}^\rho \tilde{W}_{\nu)\rho}^I + \tilde{B}_{(\mu}^\rho W_{\nu)\rho}^I)$
		$Q_{u^2 G^2 D}^{(4)}$	$if^{ABC}(\bar{u}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu u_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} - \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
		$Q_{u^2 G^2 D}^{(5)}$	$if^{ABC}(\bar{u}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu u_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} + \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
		$Q_{u^2 GBD}^{(1)}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(B_{\mu\rho} G_\nu^{A\rho} - B_{\mu\rho} G_\nu^{A\rho})$
		$Q_{u^2 GBD}^{(2)}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(B_{\mu\rho} G_\nu^{A\rho} + B_{\mu\rho} G_\nu^{A\rho})$
		$Q_{u^2 GBD}^{(3)}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(B_{[\mu}^\rho \tilde{G}_{\nu]\rho}^A - \tilde{B}_{[\mu}^\rho G_{\nu]\rho}^A)$
		$Q_{u^2 GBD}^{(4)}$	$i(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(B_{(\mu}^\rho \tilde{G}_{\nu)\rho}^A + \tilde{B}_{(\mu}^\rho G_{\nu)\rho}^A)$
		$Q_{d^2 G^2 D}^{(4)}$	$if^{ABC}(\bar{d}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu d_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} - \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
		$Q_{d^2 G^2 D}^{(5)}$	$if^{ABC}(\bar{d}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu d_r)(G_{\mu\rho}^B \tilde{G}_\nu^{C\rho} + \tilde{G}_{\mu\rho}^B G_\nu^{C\rho})$
		$Q_{d^2 GBD}^{(1)}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r)(B_{\mu\rho} G_\nu^{A\rho} - B_{\mu\rho} G_\nu^{A\rho})$
		$Q_{d^2 GBD}^{(2)}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r)(B_{\mu\rho} G_\nu^{A\rho} + B_{\mu\rho} G_\nu^{A\rho})$
		$Q_{d^2 GBD}^{(3)}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r)(B_{[\mu}^\rho \tilde{G}_{\nu]\rho}^A - \tilde{B}_{[\mu}^\rho G_{\nu]\rho}^A)$
		$Q_{d^2 GBD}^{(4)}$	$i(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r)(B_{(\mu}^\rho \tilde{G}_{\nu)\rho}^A + \tilde{B}_{(\mu}^\rho G_{\nu)\rho}^A)$

Table 6. The hadronic dimension-eight operators in the SMEFT of class-14. The subscripts p, r are weak-eigenstate indices.

14 : $\psi^2 X^2 D$		14 : $\psi^2 X^2 D$	
$Q_{l^2 G^2 D}$	$i(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r) G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{l^2 W^2 D}^{(3)}$	$i\epsilon^{IJK}(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(W_{\mu\rho}^J \widetilde{W}_\nu^{K\rho} - \widetilde{W}_{\mu\rho}^J W_\nu^{K\rho})$
$Q_{l^2 W^2 D}^{(1)}$	$i(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r) W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{l^2 W^2 D}^{(4)}$	$i\epsilon^{IJK}(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(W_{\mu\rho}^J \widetilde{W}_\nu^{K\rho} + \widetilde{W}_{\mu\rho}^J W_\nu^{K\rho})$
$Q_{l^2 W^2 D}^{(2)}$	$i\epsilon^{IJK}(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r) W_{\mu\rho}^J W_\nu^{K\rho}$	$Q_{l^2 WBD}^{(1)}$	$i(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(B_{\mu\rho} W_\nu^{I\rho} - B_{\mu\rho} W_\nu^{I\rho})$
$Q_{l^2 B^2 D}$	$i(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r) B_{\mu\rho} B_\nu^\rho$	$Q_{l^2 WBD}^{(2)}$	$i(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(B_{\mu\rho} W_\nu^{I\rho} + B_{\mu\rho} W_\nu^{I\rho})$
$Q_{e^2 G^2 D}$	$i(\bar{e}_p \gamma^\mu \overleftrightarrow{D}^\nu e_r) G_{\mu\rho}^A G_\nu^{A\rho}$	$Q_{l^2 WBD}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(B_{[\mu}^\rho \widetilde{W}_{\nu]\rho}^I - \widetilde{B}_{[\mu}^\rho W_{\nu]\rho}^I)$
$Q_{e^2 W^2 D}$	$i(\bar{e}_p \gamma^\mu \overleftrightarrow{D}^\nu e_r) W_{\mu\rho}^I W_\nu^{I\rho}$	$Q_{l^2 WBD}^{(4)}$	$i(\bar{l}_p \gamma^\mu \tau^I \overleftrightarrow{D}^\nu l_r)(B_{(\mu}^\rho \widetilde{W}_{\nu)\rho}^I + \widetilde{B}_{(\mu}^\rho W_{\nu)\rho}^I)$
$Q_{e^2 B^2 D}$	$i(\bar{e}_p \gamma^\mu \overleftrightarrow{D}^\nu e_r) B_{\mu\rho} B_\nu^\rho$		
15 : $(\bar{R}R)XH^2D$		15 : $(\bar{R}R)XH^2D$	
$Q_{e^2 WH^2 D}^{(1)}$	$(\bar{e}_p \gamma^\nu e_r) D^\mu (H^\dagger \tau^I H) W_{\mu\nu}^I$	$Q_{d^2 GH^2 D}^{(1)}$	$(\bar{d}_p \gamma^\nu T^A d_r) D^\mu (H^\dagger H) G_{\mu\nu}^A$
$Q_{e^2 WH^2 D}^{(2)}$	$(\bar{e}_p \gamma^\nu e_r) D^\mu (H^\dagger \tau^I H) \widetilde{W}_{\mu\nu}^I$	$Q_{d^2 GH^2 D}^{(2)}$	$(\bar{d}_p \gamma^\nu T^A d_r) D^\mu (H^\dagger H) \widetilde{G}_{\mu\nu}^A$
$Q_{e^2 WH^2 D}^{(3)}$	$(\bar{e}_p \gamma^\nu e_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) W_{\mu\nu}^I$	$Q_{d^2 GH^2 D}^{(3)}$	$(\bar{d}_p \gamma^\nu T^A d_r) (H^\dagger \overleftrightarrow{D}^\mu H) G_{\mu\nu}^A$
$Q_{e^2 WH^2 D}^{(4)}$	$(\bar{e}_p \gamma^\nu e_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) \widetilde{W}_{\mu\nu}^I$	$Q_{d^2 GH^2 D}^{(4)}$	$(\bar{d}_p \gamma^\nu T^A d_r) (H^\dagger \overleftrightarrow{D}^\mu H) \widetilde{G}_{\mu\nu}^A$
$Q_{e^2 BH^2 D}^{(1)}$	$(\bar{e}_p \gamma^\nu e_r) D^\mu (H^\dagger H) B_{\mu\nu}$	$Q_{d^2 WH^2 D}^{(1)}$	$(\bar{d}_p \gamma^\nu d_r) D^\mu (H^\dagger \tau^I H) W_{\mu\nu}^I$
$Q_{e^2 BH^2 D}^{(2)}$	$(\bar{e}_p \gamma^\nu e_r) D^\mu (H^\dagger H) \widetilde{B}_{\mu\nu}$	$Q_{d^2 WH^2 D}^{(2)}$	$(\bar{d}_p \gamma^\nu d_r) D^\mu (H^\dagger \tau^I H) \widetilde{W}_{\mu\nu}^I$
$Q_{e^2 BH^2 D}^{(3)}$	$(\bar{e}_p \gamma^\nu e_r) (H^\dagger \overleftrightarrow{D}^\mu H) B_{\mu\nu}$	$Q_{d^2 WH^2 D}^{(3)}$	$(\bar{d}_p \gamma^\nu d_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) W_{\mu\nu}^I$
$Q_{e^2 BH^2 D}^{(4)}$	$(\bar{e}_p \gamma^\nu e_r) (H^\dagger \overleftrightarrow{D}^\mu H) \widetilde{B}_{\mu\nu}$	$Q_{d^2 WH^2 D}^{(4)}$	$(\bar{d}_p \gamma^\nu d_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) \widetilde{W}_{\mu\nu}^I$
$Q_{u^2 GH^2 D}^{(1)}$	$(\bar{u}_p \gamma^\nu T^A u_r) D^\mu (H^\dagger H) G_{\mu\nu}^A$	$Q_{d^2 BH^2 D}^{(1)}$	$(\bar{d}_p \gamma^\nu d_r) D^\mu (H^\dagger H) B_{\mu\nu}$
$Q_{u^2 GH^2 D}^{(2)}$	$(\bar{u}_p \gamma^\nu T^A u_r) D^\mu (H^\dagger H) \widetilde{G}_{\mu\nu}^A$	$Q_{d^2 BH^2 D}^{(2)}$	$(\bar{d}_p \gamma^\nu d_r) D^\mu (H^\dagger H) \widetilde{B}_{\mu\nu}$
$Q_{u^2 GH^2 D}^{(3)}$	$(\bar{u}_p \gamma^\nu T^A u_r) (H^\dagger \overleftrightarrow{D}^\mu H) G_{\mu\nu}^A$	$Q_{d^2 BH^2 D}^{(3)}$	$(\bar{d}_p \gamma^\nu d_r) (H^\dagger \overleftrightarrow{D}^\mu H) B_{\mu\nu}$
$Q_{u^2 GH^2 D}^{(4)}$	$(\bar{u}_p \gamma^\nu T^A u_r) (H^\dagger \overleftrightarrow{D}^\mu H) \widetilde{G}_{\mu\nu}^A$	$Q_{d^2 BH^2 D}^{(4)}$	$(\bar{d}_p \gamma^\nu d_r) (H^\dagger \overleftrightarrow{D}^\mu H) \widetilde{B}_{\mu\nu}$
$Q_{u^2 WH^2 D}^{(1)}$	$(\bar{u}_p \gamma^\nu u_r) D^\mu (H^\dagger \tau^I H) W_{\mu\nu}^I$	$Q_{udGH^2+\text{h.c.}}^{(1)}$	$(\bar{u}_p \gamma^\nu T^A d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^\mu H) G_{\mu\nu}^A$
$Q_{u^2 WH^2 D}^{(2)}$	$(\bar{u}_p \gamma^\nu u_r) D^\mu (H^\dagger \tau^I H) \widetilde{W}_{\mu\nu}^I$	$Q_{udGH^2+\text{h.c.}}^{(2)}$	$(\bar{u}_p \gamma^\nu T^A d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^\mu H) \widetilde{G}_{\mu\nu}^A$
$Q_{u^2 WH^2 D}^{(3)}$	$(\bar{u}_p \gamma^\nu u_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) W_{\mu\nu}^I$	$Q_{udWH^2+\text{h.c.}}^{(1)}$	$(\bar{u}_p \gamma^\nu d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^{I\mu} H) W_{\mu\nu}^I$
$Q_{u^2 WH^2 D}^{(4)}$	$(\bar{u}_p \gamma^\nu u_r) (H^\dagger \overleftrightarrow{D}^{I\mu} H) \widetilde{W}_{\mu\nu}^I$	$Q_{udWH^2+\text{h.c.}}^{(2)}$	$(\bar{u}_p \gamma^\nu d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^{I\mu} H) \widetilde{W}_{\mu\nu}^I$
$Q_{u^2 BH^2 D}^{(1)}$	$(\bar{u}_p \gamma^\nu u_r) D^\mu (H^\dagger H) B_{\mu\nu}$	$Q_{udBH^2+\text{h.c.}}^{(1)}$	$(\bar{u}_p \gamma^\nu d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^\mu H) B_{\mu\nu}$
$Q_{u^2 BH^2 D}^{(2)}$	$(\bar{u}_p \gamma^\nu u_r) D^\mu (H^\dagger H) \widetilde{B}_{\mu\nu}$	$Q_{udBH^2+\text{h.c.}}^{(2)}$	$(\bar{u}_p \gamma^\nu d_r) (\widetilde{H}^\dagger \overleftrightarrow{D}^\mu H) \widetilde{B}_{\mu\nu}$
$Q_{u^2 BH^2 D}^{(3)}$	$(\bar{u}_p \gamma^\nu u_r) (H^\dagger \overleftrightarrow{D}^\mu H) B_{\mu\nu}$		
$Q_{u^2 BH^2 D}^{(4)}$	$(\bar{u}_p \gamma^\nu u_r) (H^\dagger \overleftrightarrow{D}^\mu H) \widetilde{B}_{\mu\nu}$		

Table 7. The leptonic dimension-eight operators in the SMEFT of class-14, and the dimension-eight operators of class-15 with field content $(\bar{R}R)X^2H$. The subscripts p, r are weak-eigenstate indices.

15 : $(\bar{L}L)XH^2D$		15 : $(\bar{L}L)XH^2D$	
$Q_{l^2WH^2D}^{(1)}$	$(\bar{l}_p\gamma^\nu l_r)D^\mu(H^\dagger\tau^I H)W_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(1)}$	$(\bar{q}_p\gamma^\nu T^A\tau^I q_r)D^\mu(H^\dagger\tau^I H)G_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(2)}$	$(\bar{l}_p\gamma^\nu l_r)D^\mu(H^\dagger\tau^I H)\widetilde{W}_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(2)}$	$(\bar{q}_p\gamma^\nu T^A\tau^I q_r)D^\mu(H^\dagger\tau^I H)\widetilde{G}_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(3)}$	$(\bar{l}_p\gamma^\nu l_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)W_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(3)}$	$(\bar{q}_p\gamma^\nu T^A\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)G_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(4)}$	$(\bar{l}_p\gamma^\nu l_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)\widetilde{W}_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(4)}$	$(\bar{q}_p\gamma^\nu T^A\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)\widetilde{G}_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(5)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger H)W_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(5)}$	$(\bar{q}_p\gamma^\nu T^A q_r)D^\mu(H^\dagger H)G_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(6)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger H)\widetilde{W}_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(6)}$	$(\bar{q}_p\gamma^\nu T^A q_r)D^\mu(H^\dagger H)\widetilde{G}_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(7)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^\mu H)W_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(7)}$	$(\bar{q}_p\gamma^\nu T^A q_r)(H^\dagger\overleftrightarrow{D}^\mu H)G_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(8)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^\mu H)\widetilde{W}_{\mu\nu}^I$	$Q_{q^2GH^2D}^{(8)}$	$(\bar{q}_p\gamma^\nu T^A q_r)(H^\dagger\overleftrightarrow{D}^\mu H)\widetilde{G}_{\mu\nu}^A$
$Q_{l^2WH^2D}^{(9)}$	$\epsilon^{IJK}(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger\tau^J H)W_{\mu\nu}^K$	$Q_{q^2WH^2D}^{(1)}$	$(\bar{q}_p\gamma^\nu q_r)D^\mu(H^\dagger\tau^I H)W_{\mu\nu}^I$
$Q_{l^2WH^2D}^{(10)}$	$\epsilon^{IJK}(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger\tau^J H)\widetilde{W}_{\mu\nu}^K$	$Q_{q^2WH^2D}^{(2)}$	$(\bar{q}_p\gamma^\nu q_r)D^\mu(H^\dagger\tau^I H)\widetilde{W}_{\mu\nu}^I$
$Q_{l^2WH^2D}^{(11)}$	$\epsilon^{IJK}(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^{J\mu}H)W_{\mu\nu}^K$	$Q_{q^2WH^2D}^{(3)}$	$(\bar{q}_p\gamma^\nu q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)W_{\mu\nu}^I$
$Q_{l^2WH^2D}^{(12)}$	$\epsilon^{IJK}(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^{J\mu}H)\widetilde{W}_{\mu\nu}^K$	$Q_{q^2WH^2D}^{(4)}$	$(\bar{q}_p\gamma^\nu q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)\widetilde{W}_{\mu\nu}^I$
$Q_{l^2BH^2D}^{(1)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger\tau^I H)B_{\mu\nu}$	$Q_{q^2WH^2D}^{(5)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger H)W_{\mu\nu}^I$
$Q_{l^2BH^2D}^{(2)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)D^\mu(H^\dagger\tau^I H)\widetilde{B}_{\mu\nu}$	$Q_{q^2WH^2D}^{(6)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger H)\widetilde{W}_{\mu\nu}^I$
$Q_{l^2BH^2D}^{(3)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)B_{\mu\nu}$	$Q_{q^2WH^2D}^{(7)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^\mu H)W_{\mu\nu}^I$
$Q_{l^2BH^2D}^{(4)}$	$(\bar{l}_p\gamma^\nu\tau^I l_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)\widetilde{B}_{\mu\nu}$	$Q_{q^2WH^2D}^{(8)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^\mu H)\widetilde{W}_{\mu\nu}^I$
$Q_{l^2BH^2D}^{(5)}$	$(\bar{l}_p\gamma^\nu l_r)D^\mu(H^\dagger H)B_{\mu\nu}$	$Q_{q^2WH^2D}^{(9)}$	$\epsilon^{IJK}(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger\tau^J H)W_{\mu\nu}^K$
$Q_{l^2BH^2D}^{(6)}$	$(\bar{l}_p\gamma^\nu l_r)D^\mu(H^\dagger H)\widetilde{B}_{\mu\nu}$	$Q_{q^2WH^2D}^{(10)}$	$\epsilon^{IJK}(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger\tau^J H)\widetilde{W}_{\mu\nu}^K$
$Q_{l^2BH^2D}^{(7)}$	$(\bar{l}_p\gamma^\nu l_r)(H^\dagger\overleftrightarrow{D}^\mu H)B_{\mu\nu}$	$Q_{q^2WH^2D}^{(11)}$	$\epsilon^{IJK}(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{J\mu}H)W_{\mu\nu}^K$
$Q_{l^2BH^2D}^{(8)}$	$(\bar{l}_p\gamma^\nu l_r)(H^\dagger\overleftrightarrow{D}^\mu H)\widetilde{B}_{\mu\nu}$	$Q_{q^2WH^2D}^{(12)}$	$\epsilon^{IJK}(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{J\mu}H)\widetilde{W}_{\mu\nu}^K$
		$Q_{q^2BH^2D}^{(1)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger\tau^I H)B_{\mu\nu}$
		$Q_{q^2BH^2D}^{(2)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)D^\mu(H^\dagger\tau^I H)\widetilde{B}_{\mu\nu}$
		$Q_{q^2BH^2D}^{(3)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)B_{\mu\nu}$
		$Q_{q^2BH^2D}^{(4)}$	$(\bar{q}_p\gamma^\nu\tau^I q_r)(H^\dagger\overleftrightarrow{D}^{I\mu}H)\widetilde{B}_{\mu\nu}$
		$Q_{q^2BH^2D}^{(5)}$	$(\bar{q}_p\gamma^\nu q_r)D^\mu(H^\dagger H)B_{\mu\nu}$
		$Q_{q^2BH^2D}^{(6)}$	$(\bar{q}_p\gamma^\nu q_r)D^\mu(H^\dagger H)\widetilde{B}_{\mu\nu}$
		$Q_{q^2BH^2D}^{(7)}$	$(\bar{q}_p\gamma^\nu q_r)(H^\dagger\overleftrightarrow{D}^\mu H)B_{\mu\nu}$
		$Q_{q^2BH^2D}^{(8)}$	$(\bar{q}_p\gamma^\nu q_r)(H^\dagger\overleftrightarrow{D}^\mu H)\widetilde{B}_{\mu\nu}$

Table 8. The dimension-eight operators in the SMEFT of class-15 with field content $(\bar{L}L)X^2H$. The subscripts p, r are weak-eigenstate indices.

16 : $\psi^2 X H D^2 + \text{h.c.}$		17 : $\psi^2 H^3 D^2 + \text{h.c.}$	
$Q_{leWH D^2}^{(1)}$	$(\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) \tau^I (D_\nu H) W_{\rho\nu}^I$	$Q_{leH^3 D^2}^{(1)}$	$(D_\mu H^\dagger D^\mu H) (\bar{l}_p e_r H)$
$Q_{leWH D^2}^{(2)}$	$(\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) \tau^I (D_\nu H) \widetilde{W}_{\rho\nu}^I$	$Q_{leH^3 D^2}^{(2)}$	$(D_\mu H^\dagger \tau^I D^\mu H) (\bar{l}_p e_r \tau^I H)$
$Q_{leWH D^2}^{(3)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I (D^\rho H) (D_\rho W_{\mu\nu}^I)$	$Q_{leH^3 D^2}^{(3)}$	$(D_\mu H^\dagger D_\nu H) (\bar{l}_p \sigma^{\mu\nu} e_r H)$
$Q_{leB H D^2}^{(1)}$	$(\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) (D_\nu H) B_{\rho\nu}$	$Q_{leH^3 D^2}^{(4)}$	$(D_\mu H^\dagger \tau^I D_\nu H) (\bar{l}_p \sigma^{\mu\nu} e_r \tau^I H)$
$Q_{leB H D^2}^{(2)}$	$(\bar{l}_p \sigma^{\mu\nu} D^\rho e_r) (D_\nu H) \widetilde{B}_{\rho\nu}$	$Q_{leH^3 D^2}^{(5)}$	$(H^\dagger D_\mu H) (\bar{l}_p e_r D^\mu H)$
$Q_{leB H D^2}^{(3)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) (D^\rho H) (D_\rho B_{\mu\nu})$	$Q_{leH^3 D^2}^{(6)}$	$(H^\dagger D_\mu H) (\bar{l}_p \sigma^{\mu\nu} e_r D_\nu H)$
$Q_{quGH D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A D^\rho u_r) (D_\nu \widetilde{H}) G_{\rho\nu}^A$	$Q_{quH^3 D^2}^{(1)}$	$(D_\mu H^\dagger D^\mu H) (\bar{q}_p u_r \widetilde{H})$
$Q_{quGH D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A D^\rho u_r) (D_\nu \widetilde{H}) \widetilde{G}_{\rho\nu}^A$	$Q_{quH^3 D^2}^{(2)}$	$(D_\mu H^\dagger \tau^I D^\mu H) (\bar{q}_p u_r \tau^I \widetilde{H})$
$Q_{quGH D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) (D^\rho \widetilde{H}) (D_\rho G_{\mu\nu}^A)$	$Q_{quH^3 D^2}^{(3)}$	$(D_\mu H^\dagger D_\nu H) (\bar{q}_p \sigma^{\mu\nu} u_r \widetilde{H})$
$Q_{quWH D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho u_r) \tau^I (D_\nu \widetilde{H}) W_{\rho\nu}^I$	$Q_{quH^3 D^2}^{(4)}$	$(D_\mu H^\dagger \tau^I D_\nu H) (\bar{q}_p \sigma^{\mu\nu} u_r \tau^I \widetilde{H})$
$Q_{quWH D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho u_r) \tau^I (D_\nu \widetilde{H}) \widetilde{W}_{\rho\nu}^I$	$Q_{quH^3 D^2}^{(5)}$	$(D_\mu H^\dagger H) (\bar{q}_p u_r D^\mu \widetilde{H})$
$Q_{quWH D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I (D^\rho \widetilde{H}) (D_\rho W_{\mu\nu}^I)$	$Q_{quH^3 D^2}^{(6)}$	$(D_\mu H^\dagger H) (\bar{q}_p \sigma^{\mu\nu} u_r D_\nu \widetilde{H})$
$Q_{quB H D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho u_r) (D_\nu \widetilde{H}) B_{\rho\nu}$	$Q_{qdH^3 D^2}^{(1)}$	$(D_\mu H^\dagger D^\mu H) (\bar{q}_p d_r H)$
$Q_{quB H D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho u_r) (D_\nu \widetilde{H}) \widetilde{B}_{\rho\nu}$	$Q_{qdH^3 D^2}^{(2)}$	$(D_\mu H^\dagger \tau^I D^\mu H) (\bar{q}_p d_r \tau^I H)$
$Q_{quB H D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) (D^\rho \widetilde{H}) (D_\rho B_{\mu\nu})$	$Q_{qdH^3 D^2}^{(3)}$	$(D_\mu H^\dagger D_\nu H) (\bar{q}_p \sigma^{\mu\nu} d_r H)$
$Q_{qdGH D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A D^\rho d_r) (D_\nu H) G_{\rho\nu}^A$	$Q_{qdH^3 D^2}^{(4)}$	$(D_\mu H^\dagger \tau^I D_\nu H) (\bar{q}_p \sigma^{\mu\nu} d_r \tau^I H)$
$Q_{qdGH D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A D^\rho d_r) (D_\nu H) \widetilde{G}_{\rho\nu}^A$	$Q_{qdH^3 D^2}^{(5)}$	$(H^\dagger D_\mu H) (\bar{q}_p d_r D^\mu H)$
$Q_{qdGH D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) (D^\rho H) (D_\rho G_{\mu\nu}^A)$	$Q_{qdH^3 D^2}^{(6)}$	$(H^\dagger D_\mu H) (\bar{q}_p \sigma^{\mu\nu} d_r D_\nu H)$
$Q_{qdWH D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho d_r) \tau^I (D_\nu H) W_{\rho\nu}^I$		
$Q_{qdWH D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho d_r) \tau^I (D_\nu H) \widetilde{W}_{\rho\nu}^I$		
$Q_{qdWH D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I (D^\rho H) (D_\rho W_{\mu\nu}^I)$		
$Q_{qdB H D^2}^{(1)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho d_r) (D_\nu H) B_{\rho\nu}$		
$Q_{qdB H D^2}^{(2)}$	$(\bar{q}_p \sigma^{\mu\nu} D^\rho d_r) (D_\nu H) \widetilde{B}_{\rho\nu}$		
$Q_{qdB H D^2}^{(3)}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) (D^\rho H) (D_\rho B_{\mu\nu})$		

Table 9. The dimension-eight operators in the SMEFT of classes-16, and -17, which have two fermions and two derivatives. All of the operators have Hermitian conjugates. The subscripts p, r are weak-eigenstate indices.

4.3 Results for Four-Fermion Operators

18 : $(\bar{L}R)(\bar{R}L)H^2 + \text{h.c.}$		18(\mathcal{B}) : $\psi^4 H^2 + \text{h.c.}$	
$Q_{leqdH^2}^{(1)}$	$(\bar{l}_p^j e_r)(\bar{d}_s q_{tj})(H^\dagger H)$	$Q_{lqudH^2}^{(1)}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}(d_p^\alpha C u_r^\beta)(q_s^{j\gamma} C l_t^k)(H^\dagger H)$
$Q_{leqdH^2}^{(2)}$	$(\bar{l}_p e_r)\tau^I(\bar{d}_s q_t)(H^\dagger \tau^I H)$	$Q_{lqudH^2}^{(2)}$	$\epsilon_{\alpha\beta\gamma}(\tau^I \epsilon)_{jk}(d_p^\alpha C u_r^\beta)(q_s^{j\gamma} C l_t^k)(H^\dagger \tau^I H)$
$Q_{l^2udH^2}$	$(\bar{l}_p d_r H)(\tilde{H}^\dagger \tilde{u}_s l_t)$	$Q_{eq^2uH^2}$	$\epsilon_{\alpha\beta\gamma}(\tau^I \epsilon)_{jk}(q_p^{j\alpha} C q_r^{k\beta})(u_s^\gamma C e_t)(H^\dagger H)$
$Q_{lequH^2}^{(5)}$	$(\bar{l}_p e_r H)(\tilde{H}^\dagger \tilde{u}_s q_t)$	$Q_{lq^3H^2}^{(1)}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{mn}\epsilon_{jk}(q_p^{m\alpha} C q_r^{j\beta})(q_s^{k\gamma} C l_t^n)(H^\dagger H)$
$Q_{q^2udH^2}^{(5)}$	$(\bar{q}_p d_r H)(\tilde{H}^\dagger \tilde{u}_s q_t)$	$Q_{lq^3H^2}^{(2)}$	$\epsilon_{\alpha\beta\gamma}(\tau^I \epsilon)_{mn}\epsilon_{jk}(q_p^{m\alpha} C q_r^{j\beta})(q_s^{k\gamma} C l_t^n)(H^\dagger \tau^I H)$
$Q_{q^2udH^2}^{(6)}$	$(\bar{q}_p T^A d_r H)(\tilde{H}^\dagger \tilde{u}_s T^A q_t)$	$Q_{eu^2dH^2}$	$\epsilon_{\alpha\beta\gamma}(d_p^\alpha C u_r^\beta)(u_s^\gamma C e_t)(H^\dagger H)$
		<hr/>	
		$Q_{lq^3H^2}^{(3)}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{mn}(\tau^I \epsilon)_{jk}(q_p^{m\alpha} C q_r^{j\beta})(q_s^{k\gamma} C l_t^n)(H^\dagger \tau^I H)$
		$Q_{lqu^2H^2}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}\epsilon_{mn}(l_p^j C q_r^{m\alpha})(u_s^\beta C u_t^\gamma)\tilde{H}^k \tilde{H}^n$
		$Q_{lqd^2H^2}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}\epsilon_{mn}(l_p^j q_r^{m\alpha})(d_s^\beta C d_t^\gamma)H^k H^n$
		$Q_{eq^2dH^2}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}\epsilon_{mn}(e_p d_r^\alpha)(q_s^{j\beta} C q_t^{m\gamma})\tilde{H}^k \tilde{H}^n$
21 : $(\bar{L}R)(\bar{R}L)D^2 + \text{h.c.}$		21(\mathcal{B}) : $\psi^4 D^2 + \text{h.c.}$	
$Q_{leqdD^2}^{(1)}$	$D_\mu(\bar{l}_p^j e_r)D^\mu(\bar{d}_s q_{tj})$	$Q_{lqudD^2}^{(1)}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}D_\mu(d_p^\alpha C u_r^\beta)D^\mu(q_s^{j\gamma} C l_t^k)$
$Q_{leqdD^2}^{(2)}$	$(\bar{l}_p^j \overleftrightarrow{D}_\mu e_r)(\bar{d}_s \overleftrightarrow{D}^\mu q_{tj})$	$Q_{lqudD^2}^{(2)}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}D_\mu(d_p^\alpha C q_r^{j\beta})D^\mu(u_s^\gamma C l_t^k)$
		$Q_{eq^2uD^2}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{jk}(q_p^{j\alpha} C D_\mu q_r^{k\beta})D^\mu(u_s^\gamma C e_t)$
		$Q_{lq^3D^2}$	$\epsilon_{\alpha\beta\gamma}\epsilon_{mn}\epsilon_{jk}(q_p^{m\alpha} C D_\mu q_r^{j\beta})D^\mu(q_s^{k\gamma} C l_t^n)$
		$Q_{eu^2dD^2}^{(1)}$	$\epsilon_{\alpha\beta\gamma}(u_p^\alpha C D_\mu u_r^\beta)D^\mu(d_s^\gamma C e_t)$
		$Q_{eu^2dD^2}^{(2)}$	$\epsilon_{\alpha\beta\gamma}(u_p^\alpha C u_r^\beta)(D_\mu d_s^\gamma C D^\mu e_t)$

Table 10. The dimension-8 operators of classes-18 and -21 whose fermionic content either has the chiral structure $(\bar{L}R)(\bar{R}L)$ or is baryon number violating. The subscripts p, r, s, t are weak-eigenstate indices. Operators below the dashed lines vanish when there is only one generation of fermions.

18 : $(\bar{L}L)(\bar{L}L)H^2$		18 : $(\bar{R}R)(\bar{R}R)H^2$	
$Q_{l^4 H^2}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma_\mu l_t)(H^\dagger H)$	$Q_{e^4 H^2}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{e}_s \gamma_\mu e_t)(H^\dagger H)$
$Q_{l^4 H^2}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma_\mu \tau^I l_t)(H^\dagger \tau^I H)$	$Q_{u^4 H^2}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger H)$
$Q_{q^4 H^2}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma_\mu q_t)(H^\dagger H)$	$Q_{d^4 H^2}$	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger H)$
$Q_{q^4 H^2}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma_\mu \tau^I q_t)(H^\dagger \tau^I H)$	$Q_{e^2 u^2 H^2}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger H)$
$Q_{q^4 H^2}^{(3)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma_\mu \tau^I q_t)(H^\dagger H)$	$Q_{e^2 d^2 H^2}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger H)$
$Q_{l^2 q^2 H^2}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma_\mu q_t)(H^\dagger H)$	$Q_{u^2 d^2 H^2}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger H)$
$Q_{l^2 q^2 H^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma_\mu q_t)(H^\dagger \tau^I H)$	$Q_{u^2 d^2 H^2}^{(2)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma_\mu T^A d_t)(H^\dagger H)$
$Q_{l^2 q^2 H^2}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma_\mu \tau^I q_t)(H^\dagger H)$		
$Q_{l^2 q^2 H^2}^{(4)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma_\mu \tau^I q_t)(H^\dagger \tau^I H)$		
$Q_{l^2 q^2 H^2}^{(5)}$	$\epsilon^{IJK}(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma_\mu \tau^J q_t)(H^\dagger \tau^K H)$		
18 : $(\bar{L}L)(\bar{R}R)H^2$		18 : $(\bar{L}R)(\bar{L}R)H^2 + \text{h.c.}$	
$Q_{l^2 e^2 H^2}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{e}_s \gamma_\mu e_t)(H^\dagger H)$	$Q_{q^2 u d H^2}^{(1)}$	$(\bar{q}_p^j u_r) \epsilon_{jk} (\bar{q}_s^k d_t)(H^\dagger H)$
$Q_{l^2 e^2 H^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{e}_s \gamma_\mu e_t)(H^\dagger \tau^I H)$	$Q_{q^2 u d H^2}^{(2)}$	$(\bar{q}_p^j u_r)(\tau^I \epsilon)_{jk} (\bar{q}_s^k d_t)(H^\dagger \tau^I H)$
$Q_{l^2 u^2 H^2}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger H)$	$Q_{q^2 u d H^2}^{(3)}$	$(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k T^A d_t)(H^\dagger H)$
$Q_{l^2 u^2 H^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger \tau^I H)$	$Q_{q^2 u d H^2}^{(4)}$	$(\bar{q}_p^j T^A u_r)(\tau^I \epsilon)_{jk} (\bar{q}_s^k T^A d_t)(H^\dagger \tau^I H)$
$Q_{l^2 d^2 H^2}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger H)$	$Q_{lequ H^2}^{(1)}$	$(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)(H^\dagger H)$
$Q_{l^2 d^2 H^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger \tau^I H)$	$Q_{lequ H^2}^{(2)}$	$(\bar{l}_p^j e_r)(\tau^I \epsilon)_{jk} (\bar{q}_s^k u_t)(H^\dagger \tau^I H)$
$Q_{q^2 e^2 H^2}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{e}_s \gamma_\mu e_t)(H^\dagger H)$	$Q_{lequ H^2}^{(3)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)(H^\dagger H)$
$Q_{q^2 e^2 H^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{e}_s \gamma_\mu e_t)(H^\dagger \tau^I H)$	$Q_{lequ H^2}^{(4)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r)(\tau^I \epsilon)_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)(H^\dagger \tau^I H)$
$Q_{q^2 u^2 H^2}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger H)$	$Q_{l^2 e^2 H^2}^{(3)}$	$(\bar{l}_p e_r H)(\bar{l}_s e_t H)$
$Q_{q^2 u^2 H^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{u}_s \gamma_\mu u_t)(H^\dagger \tau^I H)$	$Q_{leqd H^2}^{(3)}$	$(\bar{l}_p e_r H)(\bar{q}_s d_t H)$
$Q_{q^2 u^2 H^2}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma_\mu T^A u_t)(H^\dagger H)$	$Q_{leqd H^2}^{(4)}$	$(\bar{l}_p e_r \tau^I H)(\bar{q}_s d_t \tau^I H)$
$Q_{q^2 u^2 H^2}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{u}_s \gamma_\mu T^A u_t)(H^\dagger \tau^I H)$	$Q_{q^2 u^2 H^2}^{(5)}$	$(\bar{q}_p u_r \tilde{H})(\bar{q}_s u_t \tilde{H})$
$Q_{q^2 d^2 H^2}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger H)$	$Q_{q^2 u^2 H^2}^{(6)}$	$(\bar{q}_p T^A u_r \tilde{H})(\bar{q}_s T^A u_t \tilde{H})$
$Q_{q^2 d^2 H^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{d}_s \gamma_\mu d_t)(H^\dagger \tau^I H)$	$Q_{q^2 d^2 H^2}^{(5)}$	$(\bar{q}_p d_r H)(\bar{q}_s d_t H)$
$Q_{q^2 d^2 H^2}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma_\mu T^A d_t)(H^\dagger H)$	$Q_{q^2 d^2 H^2}^{(6)}$	$(\bar{q}_p T^A d_r H)(\bar{q}_s T^A d_t H)$
$Q_{q^2 d^2 H^2}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{d}_s \gamma_\mu T^A d_t)(H^\dagger \tau^I H)$		

Table 11. Most of the dimension-eight operators in the SMEFT of class-9, which are further divided into subclasses according to their chiral properties. See Table 10 for the remaining class-9 operators. Operators with + h.c. have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices.

19 : $(\bar{L}L)(\bar{L}L)X$		19 : $(\bar{R}R)(\bar{R}R)X$	
$Q_{l^4 W}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma^\nu \tau^I l_t) W_{\mu\nu}^I$	$Q_{u^4 G}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma^\nu T^A u_t) G_{\mu\nu}^A$
$Q_{l^4 W}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma^\nu \tau^I l_t) \widetilde{W}_{\mu\nu}^I$	$Q_{u^4 G}^{(2)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{G}_{\mu\nu}^A$
$Q_{q^4 G}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu T^A q_t) G_{\mu\nu}^A$	$Q_{d^4 G}^{(1)}$	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma^\nu T^A d_t) G_{\mu\nu}^A$
$Q_{q^4 G}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu T^A q_t) \widetilde{G}_{\mu\nu}^A$	$Q_{d^4 G}^{(2)}$	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{q^4 G}^{(3)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) G_{\mu\nu}^A$	$Q_{e^2 u^2 G}^{(1)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma^\nu T^A u_t) G_{\mu\nu}^A$
$Q_{q^4 G}^{(4)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) \widetilde{G}_{\mu\nu}^A$	$Q_{e^2 u^2 G}^{(2)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{G}_{\mu\nu}^A$
$Q_{q^4 W}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu \tau^I q_t) W_{\mu\nu}^I$	$Q_{e^2 u^2 B}^{(1)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma^\nu u_t) B_{\mu\nu}$
$Q_{q^4 W}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu \tau^I q_t) \widetilde{W}_{\mu\nu}^I$	$Q_{e^2 u^2 B}^{(2)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{B}_{\mu\nu}$
$Q_{q^4 W}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) W_{\mu\nu}^I$	$Q_{e^2 d^2 G}^{(1)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma^\nu T^A d_t) G_{\mu\nu}^A$
$Q_{q^4 W}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) \widetilde{W}_{\mu\nu}^I$	$Q_{e^2 d^2 G}^{(2)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{l^2 q^2 G}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu T^A q_t) G_{\mu\nu}^A$	$Q_{e^2 d^2 B}^{(1)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma^\nu d_t) B_{\mu\nu}$
$Q_{l^2 q^2 G}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu T^A q_t) \widetilde{G}_{\mu\nu}^A$	$Q_{e^2 d^2 B}^{(2)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{B}_{\mu\nu}$
$Q_{l^2 q^2 G}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) G_{\mu\nu}^A$	$Q_{u^2 d^2 G}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma^\nu T^A d_t) G_{\mu\nu}^A$
$Q_{l^2 q^2 G}^{(4)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu T^A \tau^I q_t) \widetilde{G}_{\mu\nu}^A$	$Q_{u^2 d^2 G}^{(2)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{l^2 q^2 W}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu \tau^I q_t) W_{\mu\nu}^I$	$Q_{u^2 d^2 G}^{(3)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu d_t) G_{\mu\nu}^A$
$Q_{l^2 q^2 W}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu \tau^I q_t) \widetilde{W}_{\mu\nu}^I$	$Q_{u^2 d^2 G}^{(4)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{l^2 q^2 W}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu q_t) W_{\mu\nu}^I$	$Q_{u^2 d^2 G}^{(5)}$	$f^{ABC} (\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^B d_t) G_{\mu\nu}^C$
$Q_{l^2 q^2 W}^{(4)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu q_t) \widetilde{W}_{\mu\nu}^I$	$Q_{u^2 d^2 G}^{(6)}$	$f^{ABC} (\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^B d_t) \widetilde{G}_{\mu\nu}^C$
$Q_{l^2 q^2 W}^{(5)}$	$\epsilon^{IJK} (\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu \tau^J q_t) W_{\mu\nu}^K$	$Q_{u^2 d^2 G}^{(7)}$	$d^{ABC} (\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^B d_t) G_{\mu\nu}^C$
$Q_{l^2 q^2 W}^{(6)}$	$\epsilon^{IJK} (\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu \tau^J q_t) \widetilde{W}_{\mu\nu}^K$	$Q_{u^2 d^2 G}^{(8)}$	$d^{ABC} (\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^B d_t) \widetilde{G}_{\mu\nu}^C$
$Q_{l^2 q^2 B}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu q_t) B_{\mu\nu}$	$Q_{u^2 d^2 B}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma^\nu d_t) B_{\mu\nu}$
$Q_{l^2 q^2 B}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma^\nu q_t) \widetilde{B}_{\mu\nu}$	$Q_{u^2 d^2 B}^{(2)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{B}_{\mu\nu}$
$Q_{l^2 q^2 B}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu \tau^I q_t) B_{\mu\nu}$	$Q_{u^2 d^2 B}^{(3)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^A d_t) B_{\mu\nu}$
$Q_{l^2 q^2 B}^{(4)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma^\nu \tau^I q_t) \widetilde{B}_{\mu\nu}$	$Q_{u^2 d^2 B}^{(4)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{B}_{\mu\nu}$
<hr/>		<hr/>	
$Q_{l^4 B}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma^\nu l_t) B_{\mu\nu}$	$Q_{e^4 B}^{(1)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{e}_s \gamma^\nu e_t) B_{\mu\nu}$
$Q_{l^4 B}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma^\nu l_t) \widetilde{B}_{\mu\nu}$	$Q_{e^4 B}^{(2)}$	$(\bar{e}_p \gamma^\mu e_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{B}_{\mu\nu}$
$Q_{q^4 B}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu q_t) B_{\mu\nu}$	$Q_{u^4 B}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma^\nu u_t) B_{\mu\nu}$
$Q_{q^4 B}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma^\nu q_t) \widetilde{B}_{\mu\nu}$	$Q_{u^4 B}^{(2)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{B}_{\mu\nu}$
$Q_{q^4 B}^{(3)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma^\nu \tau^I q_t) B_{\mu\nu}$	$Q_{d^4 B}^{(1)}$	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma^\nu d_t) B_{\mu\nu}$
$Q_{q^4 B}^{(4)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma^\nu \tau^I q_t) \widetilde{B}_{\mu\nu}$	$Q_{d^4 B}^{(2)}$	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{B}_{\mu\nu}$

Table 12. The dimension-eight operators in the SMEFT of class-19 with field content JJX with $J = (\bar{L}L)$ or $(\bar{R}R)$. The subscripts p, r, s, t are weak-eigenstate indices. Operators below the dashed lines vanish when there is only one generation of fermions.

19 : $(\bar{L}L)(\bar{R}R)X$		19 : $(\bar{L}L)(\bar{R}R)X$	
$Q_{l^2 e^2 W}^{(1)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{e}_s \gamma^\nu e_t) W_{\mu\nu}^I$	$Q_{q^2 u^2 G}^{(5)}$	$f^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^B u_t) G_{\mu\nu}^C$
$Q_{l^2 e^2 W}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{W}_{\mu\nu}^I$	$Q_{q^2 u^2 G}^{(6)}$	$f^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^B u_t) \widetilde{G}_{\mu\nu}^C$
$Q_{l^2 e^2 B}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{e}_s \gamma^\nu e_t) B_{\mu\nu}$	$Q_{q^2 u^2 G}^{(7)}$	$d^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^B u_t) G_{\mu\nu}^C$
$Q_{l^2 e^2 B}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{B}_{\mu\nu}$	$Q_{q^2 u^2 G}^{(8)}$	$d^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^B u_t) \widetilde{G}_{\mu\nu}^C$
$Q_{l^2 u^2 G}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma^\nu T^A u_t) G_{\mu\nu}^A$	$Q_{q^2 u^2 W}^{(1)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{u}_s \gamma^\nu u_t) W_{\mu\nu}^I$
$Q_{l^2 u^2 G}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{G}_{\mu\nu}^A$	$Q_{q^2 u^2 W}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{W}_{\mu\nu}^I$
$Q_{l^2 u^2 W}^{(1)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{u}_s \gamma^\nu u_t) W_{\mu\nu}^I$	$Q_{q^2 u^2 W}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{u}_s \gamma^\nu T^A u_t) W_{\mu\nu}^I$
$Q_{l^2 u^2 W}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{W}_{\mu\nu}^I$	$Q_{q^2 u^2 W}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{W}_{\mu\nu}^I$
$Q_{l^2 u^2 B}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma^\nu u_t) B_{\mu\nu}$	$Q_{q^2 u^2 B}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma^\nu u_t) B_{\mu\nu}$
$Q_{l^2 u^2 B}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{B}_{\mu\nu}$	$Q_{q^2 u^2 B}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{B}_{\mu\nu}$
$Q_{l^2 d^2 G}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma^\nu T^A d_t) G_{\mu\nu}^A$	$Q_{q^2 u^2 B}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^A u_t) B_{\mu\nu}$
$Q_{l^2 d^2 G}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{G}_{\mu\nu}^A$	$Q_{q^2 u^2 B}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{B}_{\mu\nu}$
$Q_{l^2 d^2 W}^{(1)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{d}_s \gamma^\nu d_t) W_{\mu\nu}^I$	$Q_{q^2 d^2 G}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma^\nu T^A d_t) G_{\mu\nu}^A$
$Q_{l^2 d^2 W}^{(2)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{W}_{\mu\nu}^I$	$Q_{q^2 d^2 G}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{l^2 d^2 B}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma^\nu d_t) B_{\mu\nu}$	$Q_{q^2 d^2 G}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu d_t) G_{\mu\nu}^A$
$Q_{l^2 d^2 B}^{(2)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{B}_{\mu\nu}$	$Q_{q^2 d^2 G}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{G}_{\mu\nu}^A$
$Q_{q^2 e^2 G}^{(1)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{e}_s \gamma^\nu e_t) G_{\mu\nu}^A$	$Q_{q^2 d^2 G}^{(5)}$	$f^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^B d_t) G_{\mu\nu}^C$
$Q_{q^2 e^2 G}^{(2)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{G}_{\mu\nu}^A$	$Q_{q^2 d^2 G}^{(6)}$	$f^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^B d_t) \widetilde{G}_{\mu\nu}^C$
$Q_{q^2 e^2 W}^{(1)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{e}_s \gamma^\nu e_t) W_{\mu\nu}^I$	$Q_{q^2 d^2 G}^{(7)}$	$d^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^B d_t) G_{\mu\nu}^C$
$Q_{q^2 e^2 W}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{W}_{\mu\nu}^I$	$Q_{q^2 d^2 G}^{(8)}$	$d^{ABC}(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^B d_t) \widetilde{G}_{\mu\nu}^C$
$Q_{q^2 e^2 B}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{e}_s \gamma^\nu e_t) B_{\mu\nu}$	$Q_{q^2 d^2 W}^{(1)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{d}_s \gamma^\nu d_t) W_{\mu\nu}^I$
$Q_{q^2 e^2 B}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{e}_s \gamma^\nu e_t) \widetilde{B}_{\mu\nu}$	$Q_{q^2 d^2 W}^{(2)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{W}_{\mu\nu}^I$
$Q_{q^2 u^2 G}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma^\nu T^A u_t) G_{\mu\nu}^A$	$Q_{q^2 d^2 W}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{d}_s \gamma^\nu T^A d_t) W_{\mu\nu}^I$
$Q_{q^2 u^2 G}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma^\nu T^A u_t) \widetilde{G}_{\mu\nu}^A$	$Q_{q^2 d^2 W}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \tau^I q_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{W}_{\mu\nu}^I$
$Q_{q^2 u^2 G}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu u_t) G_{\mu\nu}^A$	$Q_{q^2 d^2 B}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma^\nu d_t) B_{\mu\nu}$
$Q_{q^2 u^2 G}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma^\nu u_t) \widetilde{G}_{\mu\nu}^A$	$Q_{q^2 d^2 B}^{(2)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma^\nu d_t) \widetilde{B}_{\mu\nu}$
		$Q_{q^2 d^2 B}^{(3)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^A d_t) B_{\mu\nu}$
		$Q_{q^2 d^2 B}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma^\nu T^A d_t) \widetilde{B}_{\mu\nu}$

Table 13. The dimension-eight operators in the SMEFT of class-19 with field content $(\bar{L}L)(\bar{R}R)X$. The subscripts p, r, s, t are weak-eigenstate indices.

19 : $(\bar{L}R)(\bar{R}L)X + \text{h.c.}$		19 : $(\bar{L}R)(\bar{L}R)X + \text{h.c.}$	
$Q_{ledqG}^{(1)}$	$(\bar{l}_p^j \sigma^{\mu\nu} e_r)(\bar{d}_s T^A q_{tj}) G_{\mu\nu}^A$	$Q_{q^2udG}^{(1)}$	$(\bar{q}_p^j \sigma^{\mu\nu} T^A u_r) \epsilon_{jk} (\bar{q}_s^k d_t) G_{\mu\nu}^A$
$Q_{ledqG}^{(2)}$	$(\bar{l}_p^j e_r)(\bar{d}_s \sigma^{\mu\nu} T^A q_{tj}) G_{\mu\nu}^A$	$Q_{q^2udG}^{(2)}$	$(\bar{q}_p^j \sigma^{\mu\nu} u_r) \epsilon_{jk} (\bar{q}_s^k T^A d_t) G_{\mu\nu}^A$
$Q_{ledqW}^{(1)}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I (\bar{d}_s q_t) W_{\mu\nu}^I$	$Q_{q^2udG}^{(3)}$	$(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} d_t) G_{\mu\nu}^A$
$Q_{ledqW}^{(2)}$	$(\bar{l}_p e_r) \tau^I (\bar{d}_s \sigma^{\mu\nu} q_t) W_{\mu\nu}^I$	$Q_{q^2udG}^{(4)}$	$(\bar{q}_p^j u_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} T^A d_t) G_{\mu\nu}^A$
$Q_{ledqB}^{(1)}$	$(\bar{l}_p^j \sigma^{\mu\nu} e_r)(\bar{d}_s q_{tj}) B_{\mu\nu}$	$Q_{q^2udG}^{(5)}$	$(\bar{q}_p^j \sigma^{\mu\rho} T^A u_r) \epsilon_{jk} (\bar{q}_s^k \sigma_{\rho\nu} d_t) G_{\mu\nu}^{A\nu}$
$Q_{ledqB}^{(2)}$	$(\bar{l}_p^j e_r)(\bar{d}_s \sigma^{\mu\nu} q_{tj}) B_{\mu\nu}$	$Q_{q^2udG}^{(6)}$	$(\bar{q}_p^j \sigma^{\mu\rho} u_r) \epsilon_{jk} (\bar{q}_s^k \sigma_{\rho\nu} T^A d_t) G_{\mu\nu}^{A\nu}$
		$Q_{q^2udW}^{(1)}$	$(\bar{q}_p^j \sigma^{\mu\nu} u_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k d_t) W_{\mu\nu}^I$
		$Q_{q^2udW}^{(2)}$	$(\bar{q}_p^j u_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k \sigma^{\mu\nu} d_t) W_{\mu\nu}^I$
		$Q_{q^2udW}^{(3)}$	$(\bar{q}_p^j \sigma^{\mu\rho} u_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k \sigma_{\rho\nu} d_t) W_{\mu\nu}^{I\nu}$
		$Q_{q^2udB}^{(1)}$	$(\bar{q}_p^j \sigma^{\mu\nu} u_r) \epsilon_{jk} (\bar{q}_s^k d_t) B_{\mu\nu}$
		$Q_{q^2udB}^{(2)}$	$(\bar{q}_p^j u_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} d_t) B_{\mu\nu}$
		$Q_{q^2udB}^{(3)}$	$(\bar{q}_p^j \sigma^{\mu\rho} u_r) \epsilon_{jk} (\bar{q}_s^k \sigma_{\rho\nu} d_t) B_{\mu\nu}^\nu$
		$Q_{lequG}^{(1)}$	$(\bar{l}_p^j \sigma^{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k T^A u_t) G_{\mu\nu}^A$
		$Q_{lequG}^{(2)}$	$(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} T^A u_t) G_{\mu\nu}^A$
		$Q_{lequG}^{(3)}$	$(\bar{l}_p^j \sigma^{\mu\rho} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma_{\rho\nu} T^A u_t) G_{\mu\nu}^{A\nu}$
		$Q_{lequW}^{(1)}$	$(\bar{l}_p^j \sigma^{\mu\nu} e_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k u_t) W_{\mu\nu}^I$
		$Q_{lequW}^{(2)}$	$(\bar{l}_p^j e_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t) W_{\mu\nu}^I$
		$Q_{lequW}^{(3)}$	$(\bar{l}_p^j \sigma^{\mu\rho} e_r) (\tau^I \epsilon)_{jk} (\bar{q}_s^k \sigma_{\rho\nu} u_t) W_{\mu\nu}^{I\nu}$
		$Q_{lequB}^{(1)}$	$(\bar{l}_p^j \sigma^{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k u_t) B_{\mu\nu}$
		$Q_{lequB}^{(2)}$	$(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t) B_{\mu\nu}$
		$Q_{lequB}^{(3)}$	$(\bar{l}_p^j \sigma^{\mu\rho} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma_{\rho\nu} u_t) B_{\mu\nu}^\nu$

Table 14. The dimension-eight operators in the SMEFT of class-19 with field content $(\bar{L}R)(\bar{R}L)X$ or $(\bar{L}R)(\bar{L}R)X$. All of the operators have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices.

19(\mathcal{B}) : $\psi^4 X + \text{h.c.}$

$Q_{lqu dG}^{(1)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} \epsilon_{jk} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (q_s^{j\gamma} C l_t^k) G_{\mu\nu}^A$
$Q_{lqu dG}^{(2)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} \epsilon_{jk} (d_p^\alpha C u_r^\beta) (q_s^{j\gamma} C \sigma^{\mu\nu} l_t^k) G_{\mu\nu}^A$
$Q_{lqu dG}^{(3)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta \epsilon_{jk} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (q_s^{j\gamma} C l_t^k) G_{\mu\nu}^A$
$Q_{lqu dG}^{(4)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta \epsilon_{jk} (d_p^\alpha C u_r^\beta) (q_s^{j\gamma} C \sigma^{\mu\nu} l_t^k) G_{\mu\nu}^A$
$Q_{lqu dW}^{(1)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{jk} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (q_s^{j\gamma} C l_t^k) W_{\mu\nu}^I$
$Q_{lqu dW}^{(2)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{jk} (d_p^\alpha C u_r^\beta) (q_s^{j\gamma} C \sigma^{\mu\nu} l_t^k) W_{\mu\nu}^I$
$Q_{lqu dB}^{(1)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (q_s^{j\gamma} C l_t^k) B_{\mu\nu}$
$Q_{lqu dB}^{(2)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (d_p^\alpha C u_r^\beta) (q_s^{j\gamma} C \sigma^{\mu\nu} l_t^k) B_{\mu\nu}$
$Q_{eq^2 uG}^{(1)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} \epsilon_{jk} (q_p^{j\alpha} C \sigma^{\mu\nu} q_r^{k\beta}) (u_s^\gamma C e_t) G_{\mu\nu}^A$
$Q_{eq^2 uG}^{(2)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta \epsilon_{jk} (q_p^{j\alpha} C q_r^{k\beta}) (u_s^\gamma C \sigma^{\mu\nu} e_t) G_{\mu\nu}^A$
$Q_{eq^2 uW}^{(1)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{jk} (q_p^{j\alpha} C \sigma^{\mu\nu} q_r^{k\beta}) (u_s^\gamma C e_t) W_{\mu\nu}^I$
$Q_{eq^2 uB}^{(1)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C q_r^{k\beta}) (u_s^\gamma C \sigma^{\mu\nu} e_t) B_{\mu\nu}$
$Q_{lq^3 G}^{(1)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) G_{\mu\nu}^A$
$Q_{lq^3 G}^{(2)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C \sigma^{\mu\nu} l_t^n) G_{\mu\nu}^A$
$Q_{lq^3 W}^{(1)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{mn} \epsilon_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C \sigma^{\mu\nu} l_t^n) W_{\mu\nu}^I$
$Q_{lq^3 W}^{(2)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{mj} \epsilon_{kn} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) W_{\mu\nu}^I$
$Q_{lq^3 B}^{(1)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C \sigma^{\mu\nu} l_t^n) B_{\mu\nu}$
$Q_{eu^2 dG}^{(1)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (u_s^\gamma C e_t) G_{\mu\nu}^A$
$Q_{eu^2 dG}^{(2)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} (u_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (d_s^\gamma C e_t) G_{\mu\nu}^A$
$Q_{eu^2 dG}^{(3)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta (u_p^\alpha C u_r^\beta) (d_s^\gamma C \sigma^{\mu\nu} e_t) G_{\mu\nu}^A$
$Q_{eu^2 dB}^{(1)}$	$\epsilon_{\alpha\beta\gamma} (d_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (u_s^\gamma C e_t) B_{\mu\nu}$
$Q_{eu^2 dB}^{(2)}$	$\epsilon_{\alpha\beta\gamma} (u_p^\alpha C \sigma^{\mu\nu} u_r^\beta) (d_s^\gamma C e_t) B_{\mu\nu}$
$Q_{eq^2 uW}^{(2)}$	$\epsilon_{\alpha\beta\gamma} (\tau^I \epsilon)_{jk} (q_p^{j\alpha} C q_r^{k\beta}) (u_s^\gamma C \sigma^{\mu\nu} e_t) W_{\mu\nu}^I$
$Q_{eq^2 uB}^{(2)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C \sigma^{\mu\nu} q_r^{k\beta}) (u_s^\gamma C e_t) B_{\mu\nu}$
$Q_{lq^3 G}^{(3)}$	$(T^A)_{(\alpha}^\delta \epsilon_{\beta)\gamma} \delta \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) G_{\mu\nu}^A$
$Q_{lq^3 G}^{(4)}$	$(T^A)_\gamma^\delta \epsilon_{\delta\alpha\beta} \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C \sigma^{\mu\nu} l_t^n) G_{\mu\nu}^A$
$Q_{lq^3 W}^{(3)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{mn} (\tau^I \epsilon)_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C \sigma^{\mu\nu} l_t^n) W_{\mu\nu}^I$
$Q_{lq^3 B}^{(2)}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C \sigma^{\mu\nu} q_r^{j\beta}) (q_s^{k\gamma} C l_t^n) B_{\mu\nu}$

Table 15. The baryon number violating dimension-eight operators of class-19. All of the operators have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices. Operators below the dashed line vanish when there is only one generation of fermions.

20 : $\psi^4 HD + \text{h.c.}$		20 : $\psi^4 HD + \text{h.c.}$	
$Q_{l^3 e HD}^{(1)}$	$i(\bar{l}_p \gamma^\mu l_r)[(\bar{l}_s e_t) D_\mu H]$	$Q_{l^2 qu HD}^{(1)}$	$i(\bar{l}_p \gamma^\mu l_r)[(\bar{q}_s u_t) D_\mu \tilde{H}]$
$Q_{l^3 e HD}^{(2)}$	$i(\bar{l}_p \gamma^\mu \tau^I l_r)[(\bar{l}_s e_t) \tau^I D_\mu H]$	$Q_{l^2 qu HD}^{(2)}$	$i(\bar{q}_{p\alpha} \gamma^\mu l_r)[(\bar{l}_s u_t^\alpha) D_\mu \tilde{H}]$
$Q_{l^3 e HD}^{(3)}$	$i(\bar{l}_p \gamma^\mu l_r)[(D_\mu \bar{l}_s e_t) H]$	$Q_{l^2 qu HD}^{(3)}$	$i(\bar{l}_p \gamma^\mu \tau^I l_r)[(\bar{q}_s u_t) \tau^I D_\mu \tilde{H}]$
$Q_{le^3 HD}^{(1)}$	$i(\bar{e}_p \gamma^\mu e_r)[(\bar{l}_s D_\mu e_t) H]$	$Q_{l^2 qu HD}^{(4)}$	$i(\bar{q}_{p\alpha} \gamma^\mu \tau^I l_r)[(\bar{l}_s u_t^\alpha) \tau^I D_\mu \tilde{H}]$
$Q_{leq^2 HD}^{(1)}$	$i(\bar{q}_p \gamma^\mu q_r)[(\bar{l}_s e_t) D_\mu H]$	$Q_{l^2 qu HD}^{(5)}$	$i(\bar{l}_p \gamma^\mu l_r)[(\bar{q}_s D_\mu u_t) \tilde{H}]$
$Q_{leq^2 HD}^{(2)}$	$i(\bar{l}_p \gamma^\mu q_r^\alpha)[(\bar{q}_{s\alpha} e_t) D_\mu H]$	$Q_{l^2 qu HD}^{(6)}$	$i(\bar{l}_p \gamma^\mu \tau^I l_r)[(\bar{q}_s D_\mu u_t) \tau^I \tilde{H}]$
$Q_{leq^2 HD}^{(3)}$	$i(\bar{q}_p \gamma^\mu \tau^I q_r)[(\bar{l}_s e_t) \tau^I D_\mu H]$	$Q_{e^2 qu HD}^{(1)}$	$i(\bar{e}_p \gamma^\mu e_r)[(\bar{q}_s u_t) D_\mu \tilde{H}]$
$Q_{leq^2 HD}^{(4)}$	$i(\bar{l}_p \gamma^\mu \tau^I q_r^\alpha)[(\bar{q}_{s\alpha} e_t) \tau^I D_\mu H]$	$Q_{e^2 qu HD}^{(2)}$	$i(\bar{e}_p \gamma^\mu u_r)[(\bar{q}_{s\alpha} e_t) D_\mu \tilde{H}]$
$Q_{leq^2 HD}^{(5)}$	$i(\bar{q}_p \gamma^\mu q_r)[(\bar{l}_s D_\mu e_t) H]$	$Q_{e^2 qu HD}^{(3)}$	$i(\bar{e}_p \gamma^\mu e_r)[(D_\mu \bar{q}_s u_t) \tilde{H}]$
$Q_{leq^2 HD}^{(6)}$	$i(\bar{q}_p \gamma^\nu \tau^I q_r)[(\bar{l}_s D_\mu e_t) \tau^I H]$	$Q_{q^3 u HD}^{(1)}$	$i(\bar{q}_p \gamma^\mu q_r)[(\bar{q}_s u_t) D_\mu \tilde{H}]$
$Q_{leu^2 HD}^{(1)}$	$i(\bar{u}_p \gamma^\mu u_r)[(\bar{l}_s e_t) D_\mu H]$	$Q_{q^3 u HD}^{(2)}$	$i(\bar{q}_p \gamma^\mu \tau^I q_r)[(\bar{q}_s u_t) \tau^I D_\mu \tilde{H}]$
$Q_{leu^2 HD}^{(2)}$	$i(\bar{u}_{p\alpha} \gamma^\mu e_r)[(\bar{l}_s u_t^\alpha) D_\mu H]$	$Q_{q^3 u HD}^{(3)}$	$i(\bar{q}_p \gamma^\mu T^A q_r)[(\bar{q}_s T^A u_t) D_\mu \tilde{H}]$
$Q_{leu^2 HD}^{(3)}$	$i(\bar{u}_p \gamma^\mu u_r)[(D_\mu \bar{l}_s e_t) H]$	$Q_{q^3 u HD}^{(4)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I q_r)[(\bar{q}_s T^A u_t) \tau^I D_\mu \tilde{H}]$
$Q_{led^2 HD}^{(1)}$	$i(\bar{d}_p \gamma^\mu d_r)[(\bar{l}_s e_t) D_\mu H]$	$Q_{q^3 u HD}^{(5)}$	$i(\bar{q}_p \gamma^\mu q_r)[(D_\mu \bar{q}_s u_t) \tilde{H}]$
$Q_{led^2 HD}^{(2)}$	$i(\bar{d}_{p\alpha} \gamma^\mu e_r)[(\bar{l}_s d_t^\alpha) D_\mu H]$	$Q_{q^3 u HD}^{(6)}$	$i(\bar{q}_p \gamma^\mu \tau^I q_r)[(D_\mu \bar{q}_s u_t) \tau^I \tilde{H}]$
$Q_{led^2 HD}^{(3)}$	$i(\bar{d}_p \gamma^\mu d_r)[(D_\mu \bar{l}_s e_t) H]$	$Q_{qu^3 HD}^{(1)}$	$i(\bar{u}_p \gamma^\mu u_r)[(\bar{q}_s u_t) D_\mu \tilde{H}]$
$Q_{leud HD}^{(1)}$	$i\epsilon_{jk}(\bar{u}_p \gamma^\mu d_r)(\bar{e}_s l_t^j) D_\mu H^k$	$Q_{qu^3 HD}^{(2)}$	$i(\bar{u}_p \gamma^\mu T^A u_r)[(\bar{q}_s T^A u_t) D_\mu \tilde{H}]$
$Q_{leud HD}^{(2)}$	$i\epsilon_{jk}(\bar{e}_p \gamma^\mu d_r^\alpha)(\bar{u}_{s\alpha} l_t^j) D_\mu H^k$	$Q_{qu^3 HD}^{(3)}$	$i(\bar{u}_p \gamma^\mu u_r)[(\bar{q}_s D_\mu u_t) \tilde{H}]$
$Q_{leud^2 HD}^{(3)}$	$i\epsilon_{jk}(\bar{u}_p \gamma^\mu d_r)(\bar{e}_s D_\mu l_t^j) H^k$	$Q_{qud^2 HD}^{(1)}$	$i(\bar{d}_p \gamma^\mu d_r)[(\bar{q}_s u_t) D_\mu \tilde{H}]$
$Q_{le^3 HD}^{(2)}$	$i(\bar{e}_p \gamma^\mu e_r)[(\bar{l}_s e_t) D_\mu H]$	$Q_{qud^2 HD}^{(2)}$	$i(\bar{d}_p \gamma^\mu u_r)[(\bar{q}_s d_t) D_\mu \tilde{H}]$
		$Q_{qud^2 HD}^{(3)}$	$i(\bar{d}_p \gamma^\mu T^A d_r)[(\bar{q}_s T^A u_t) D_\mu \tilde{H}]$
		$Q_{qud^2 HD}^{(4)}$	$i(\bar{d}_p \gamma^\mu T^A u_r)[(\bar{q}_s T^A d_t) D_\mu \tilde{H}]$
		$Q_{qud^2 HD}^{(5)}$	$i(\bar{d}_p \gamma^\mu d_r)[(D_\mu \bar{q}_s u_t) \tilde{H}]$
		$Q_{qud^2 HD}^{(6)}$	$i(\bar{d}_p \gamma^\mu T^A d_r)[(D_\mu \bar{q}_s T^A u_t) \tilde{H}]$

Table 16. The dimension-eight operators in the SMEFT of class-20 whose field content superficially includes either an electron-type or up-quark-type Yukawa interaction. All of the operators have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices. The operator below the dashed line is redundant when there is only one generation of fermions.

20 : $\psi^4 HD + \text{h.c.}$		20(\cancel{B}) : $\psi^4 HD + \text{h.c.}$	
$Q_{l^2 q d HD}^{(1)}$	$i(\bar{l}_p \gamma^\mu l_r)[(\bar{q}_s d_t) D_\mu H]$	$Q_{lu^2 d HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma}[D_\mu H^\dagger(u_p^\alpha C \gamma^\mu l_r)](u_s^\beta C d_t^\gamma)$
$Q_{l^2 q d HD}^{(2)}$	$i(\bar{q}_{p\alpha} \gamma^\mu l_r)[(\bar{l}_s d_t^\alpha) D_\mu H]$	$Q_{lu^2 d HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma}[H^\dagger(u_p^\alpha C \gamma^\mu l_r)](D_\mu u_s^\beta C d_t^\gamma)$
$Q_{l^2 q d HD}^{(3)}$	$i(\bar{l}_p \gamma^\mu \tau^I l_r)[(\bar{q}_s d_t) \tau^I D_\mu H]$	$Q_{lu^2 d HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jk}(d_p^\alpha C \gamma^\mu l_r^j)(d_s^\beta C u_t^\gamma) D_\mu H^k$
$Q_{l^2 q d HD}^{(4)}$	$i(\bar{q}_{p\alpha} \gamma^\mu \tau^I l_r)[(\bar{l}_s d_t^\alpha) \tau^I D_\mu H]$	$Q_{lu^2 d HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jk}(d_p^\alpha C \gamma^\mu l_r^j)(D_\mu d_s^\beta C u_t^\gamma) H^k$
$Q_{l^2 q d HD}^{(5)}$	$i(\bar{l}_p \gamma^\mu l_r)[(\bar{q}_s D_\mu d_t) H]$	$Q_{lq^2 u HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jn} \epsilon_{km} D_\mu H^{n\dagger}(q_p^{m\alpha} C \gamma^\mu u_r^\beta)(q_s^{j\gamma} C l_t^k)$
$Q_{l^2 q d HD}^{(6)}$	$i(\bar{l}_p \gamma^\mu \tau^I l_r)[(\bar{q}_s D_\mu d_t) \tau^I H]$	$Q_{lq^2 u HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{kn} \epsilon_{jm} D_\mu H^{n\dagger}(q_p^{m\alpha} C \gamma^\mu u_r^\beta)(q_s^{j\gamma} C l_t^k)$
$Q_{e^2 q d HD}^{(1)}$	$i(\bar{e}_p \gamma^\mu e_r)[(\bar{q}_s d_t) D_\mu H]$	$Q_{lq^2 u HD}^{(3)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jn} \epsilon_{km} H^{n\dagger}(q_p^{m\alpha} C \gamma^\mu u_r^\beta)(D_\mu q_s^{j\gamma} C l_t^k)$
$Q_{e^2 q d HD}^{(2)}$	$i(\bar{e}_p \gamma^\mu d_r^\alpha)[(\bar{q}_s e_t) D_\mu H]$	$Q_{lq^2 d HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jn} \epsilon_{km} (q_p^{m\alpha} C \gamma^\mu d_r^\beta)(q_s^{j\gamma} C l_t^k) D_\mu H^n$
$Q_{e^2 q d HD}^{(3)}$	$i(\bar{e}_p \gamma^\mu e_r)[(D_\mu \bar{q}_s d_t) H]$	$Q_{lq^2 d HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{kn} \epsilon_{jm} (q_p^{m\alpha} C \gamma^\mu d_r^\beta)(q_s^{j\gamma} C l_t^k) D_\mu H^n$
$Q_{q^3 d HD}^{(1)}$	$i(\bar{q}_p \gamma^\mu q_r)[(\bar{q}_s d_t) D_\mu H]$	$Q_{lq^2 d HD}^{(3)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jn} \epsilon_{km} (q_p^{m\alpha} C \gamma^\mu d_r^\beta)(D_\mu q_s^{j\gamma} C l_t^k) H^n$
$Q_{q^3 d HD}^{(2)}$	$i(\bar{q}_p \gamma^\mu \tau^I q_r)[(\bar{q}_s d_t) \tau^I D_\mu H]$	$Q_{eq^3 HD}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{mn} \epsilon_{jk} (q_p^{j\alpha} C \gamma^\mu e_r)(D_\mu q_s^{k\beta} C q_t^{m\gamma}) H^n$
$Q_{q^3 d HD}^{(3)}$	$i(\bar{q}_p \gamma^\mu T^A q_r)[(\bar{q}_s T^A d_t) D_\mu H]$	$Q_{eq^2 HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma}[D_\mu H^\dagger(u_p^\alpha C \gamma^\mu q_r^\beta)](u_s^\gamma C e_t)$
$Q_{q^3 d HD}^{(4)}$	$i(\bar{q}_p \gamma^\mu T^A \tau^I q_r)[(\bar{q}_s T^A d_t) \tau^I D_\mu H]$	$Q_{eq^2 HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma}[H^\dagger(u_p^\alpha C \gamma^\mu q_r^\beta)](D_\mu u_s^\gamma C e_t)$
$Q_{q^3 d HD}^{(5)}$	$i(\bar{q}_p \gamma^\mu q_r)[(D_\mu \bar{q}_s d_t) H]$	$Q_{equd HD}^{(1)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C \gamma^\mu u_r^\beta)(d_s^\gamma C e_t) D_\mu H^k$
$Q_{q^3 d HD}^{(6)}$	$i(\bar{q}_p \gamma^\mu \tau^I q_r)[(D_\mu \bar{q}_s d_t) \tau^I H]$	$Q_{equd HD}^{(2)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C \gamma^\mu d_r^\beta)(u_s^\gamma C e_t) D_\mu H^k$
$Q_{qu^2 d HD}^{(1)}$	$i(\bar{u}_p \gamma^\mu u_r)[(\bar{q}_s d_t) D_\mu H]$	$Q_{equd HD}^{(3)}$	$i\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C \gamma^\mu u_r^\beta)(d_s^\gamma C D_\mu e_t) H^k$
$Q_{qu^2 d HD}^{(2)}$	$i(\bar{u}_p \gamma^\mu d_r)[(\bar{q}_s u_t) D_\mu H]$		
$Q_{qu^2 d HD}^{(3)}$	$i(\bar{u}_p \gamma^\mu T^A u_r)[(\bar{q}_s T^A d_t) D_\mu H]$		
$Q_{qu^2 d HD}^{(4)}$	$i(\bar{u}_p \gamma^\mu T^A d_r)[(\bar{q}_s T^A u_t) D_\mu H]$		
$Q_{qu^2 d HD}^{(5)}$	$i(\bar{u}_p \gamma^\mu u_r)[(D_\mu \bar{q}_s d_t) H]$		
$Q_{qu^2 d HD}^{(6)}$	$i(\bar{u}_p \gamma^\mu T^A u_r)[(D_\mu \bar{q}_s T^A d_t) H]$		
$Q_{qd^3 HD}^{(1)}$	$i(\bar{d}_p \gamma^\mu d_r)[(\bar{q}_s d_t) D_\mu H]$		
$Q_{qd^3 HD}^{(2)}$	$i(\bar{d}_p \gamma^\mu T^A d_r)[(\bar{q}_s T^A d_t) D_\mu H]$		
$Q_{qd^3 HD}^{(3)}$	$i(\bar{d}_p \gamma^\mu d_r)[(\bar{q}_s D_\mu d_t) H]$		

Table 17. The dimension-eight operators in the SMEFT of class-20 whose field content superficially includes a down-quark-type Yukawa interaction or is baryon number violating. All of the operators have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices.

21 : $(\bar{L}L)(\bar{L}L)D^2$		21 : $(\bar{R}R)(\bar{R}R)D^2$	
$Q_{l^4 D^2}^{(1)}$	$D^\nu(\bar{l}_p \gamma^\mu l_r) D_\nu(\bar{l}_s \gamma_\mu l_t)$	$Q_{e^4 D^2}$	$D^\nu(\bar{e}_p \gamma^\mu e_r) D_\nu(\bar{e}_s \gamma_\mu e_t)$
$Q_{l^4 D^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r)(\bar{l}_s \gamma_\mu \overleftrightarrow{D}_\nu l_t)$	$Q_{u^4 D^2}^{(1)}$	$D^\nu(\bar{u}_p \gamma^\mu u_r) D_\nu(\bar{u}_s \gamma_\mu u_t)$
$Q_{q^4 D^2}^{(1)}$	$D^\nu(\bar{q}_p \gamma^\mu q_r) D_\nu(\bar{q}_s \gamma_\mu q_t)$	$Q_{u^4 D^2}^{(2)}$	$(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(\bar{u}_s \gamma_\mu \overleftrightarrow{D}_\nu u_t)$
$Q_{q^4 D^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(\bar{q}_s \gamma_\mu \overleftrightarrow{D}_\nu q_t)$	$Q_{d^4 D^2}^{(1)}$	$D^\nu(\bar{d}_p \gamma^\mu d_r) D_\nu(\bar{d}_s \gamma_\mu d_t)$
$Q_{q^4 D^2}^{(3)}$	$D^\nu(\bar{q}_p \gamma^\mu \tau^I q_r) D_\nu(\bar{q}_s \gamma_\mu \tau^I q_t)$	$Q_{d^4 D^2}^{(2)}$	$(\bar{d}_p \gamma^\mu \overleftrightarrow{D}^\nu d_r)(\bar{d}_s \gamma_\mu \overleftrightarrow{D}_\nu d_t)$
$Q_{q^4 D^2}^{(4)}$	$(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^{I\nu} q_r)(\bar{q}_s \gamma_\mu \overleftrightarrow{D}_\nu^I q_t)$	$Q_{e^2 u^2 D^2}^{(1)}$	$D^\nu(\bar{e}_p \gamma^\mu e_r) D_\nu(\bar{u}_s \gamma_\mu u_t)$
$Q_{l^2 q^2 D^2}^{(1)}$	$D^\nu(\bar{l}_p \gamma^\mu l_r) D_\nu(\bar{q}_s \gamma_\mu q_t)$	$Q_{e^2 u^2 D^2}^{(2)}$	$(\bar{e}_p \gamma^\mu \overleftrightarrow{D}^\nu e_r)(\bar{u}_s \gamma_\mu \overleftrightarrow{D}_\nu u_t)$
$Q_{l^2 q^2 D^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r)(\bar{q}_s \gamma_\mu \overleftrightarrow{D}_\nu q_t)$	$Q_{e^2 d^2 D^2}^{(1)}$	$D^\nu(\bar{e}_p \gamma^\mu e_r) D_\nu(\bar{d}_s \gamma_\mu d_t)$
$Q_{l^2 q^2 D^2}^{(3)}$	$D^\nu(\bar{l}_p \gamma^\mu \tau^I l_r) D_\nu(\bar{q}_s \gamma_\mu \tau^I q_t)$	$Q_{e^2 d^2 D^2}^{(2)}$	$(\bar{e}_p \gamma^\mu \overleftrightarrow{D}^\nu e_r)(\bar{d}_s \gamma_\mu \overleftrightarrow{D}_\nu d_t)$
$Q_{l^2 q^2 D^2}^{(4)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^{I\nu} l_r)(\bar{q}_s \gamma_\mu \overleftrightarrow{D}_\nu^I q_t)$	$Q_{u^2 d^2 D^2}^{(1)}$	$D^\nu(\bar{u}_p \gamma^\mu u_r) D_\nu(\bar{d}_s \gamma_\mu d_t)$
		$Q_{u^2 d^2 D^2}^{(2)}$	$(\bar{u}_p \gamma^\mu \overleftrightarrow{D}^\nu u_r)(\bar{d}_s \gamma_\mu \overleftrightarrow{D}_\nu d_t)$
		$Q_{u^2 d^2 D^2}^{(3)}$	$D^\nu(\bar{u}_p \gamma^\mu T^A u_r) D_\nu(\bar{d}_s \gamma_\mu T^A d_t)$
		$Q_{u^2 d^2 D^2}^{(4)}$	$(\bar{u}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu u_r)(\bar{d}_s \gamma_\mu T^A \overleftrightarrow{D}_\nu d_t)$
21 : $(\bar{L}L)(\bar{R}R)D^2$		21 : $(\bar{L}R)(\bar{L}R)D^2 + \text{h.c.}$	
$Q_{l^2 e^2 D^2}^{(1)}$	$D^\nu(\bar{l}_p \gamma^\mu l_r) D_\nu(\bar{e}_s \gamma_\mu e_t)$	$Q_{q^2 u d D^2}^{(1)}$	$D_\mu(\bar{q}_p^j u_r) \epsilon_{jk} D^\mu(\bar{q}_s^k d_t)$
$Q_{l^2 e^2 D^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r)(\bar{e}_s \gamma_\mu \overleftrightarrow{D}_\nu e_t)$	$Q_{q^2 u d D^2}^{(2)}$	$D_\mu(\bar{q}_p^j T^A u_r) \epsilon_{jk} D^\mu(\bar{q}_s^k T^A d_t)$
$Q_{l^2 u^2 D^2}^{(1)}$	$D^\nu(\bar{l}_p \gamma^\mu l_r) D_\nu(\bar{u}_s \gamma_\mu u_t)$	$Q_{q^2 u d D^2}^{(3)}$	$D_\mu(\bar{q}_p^j C \bar{q}_{r\beta}^k) \epsilon_{jk} D^\mu(u_s^\alpha C d_t^\beta)$
$Q_{l^2 u^2 D^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r)(\bar{u}_s \gamma_\mu \overleftrightarrow{D}_\nu u_t)$	$Q_{lequ D^2}^{(1)}$	$D_\mu(\bar{l}_p^j e_r) \epsilon_{jk} D^\mu(\bar{q}_s^k u_t)$
$Q_{l^2 d^2 D^2}^{(1)}$	$D^\nu(\bar{l}_p \gamma^\mu l_r) D_\nu(\bar{d}_s \gamma_\mu d_t)$	$Q_{lequ D^2}^{(2)}$	$D_\mu(\bar{l}_p^j u_r^\alpha) \epsilon_{jk} D^\mu(\bar{q}_s^k e_t)$
$Q_{l^2 d^2 D^2}^{(2)}$	$(\bar{l}_p \gamma^\mu \overleftrightarrow{D}^\nu l_r)(\bar{d}_s \gamma_\mu \overleftrightarrow{D}_\nu d_t)$	$Q_{lequ D^2}^{(3)}$	$D_\mu(\bar{l}_p^j C \bar{q}_{r\alpha}^k) \epsilon_{jk} D^\mu(e_s C u_t^\alpha)$
$Q_{q^2 e^2 D^2}^{(1)}$	$D^\nu(\bar{q}_p \gamma^\mu q_r) D_\nu(\bar{e}_s \gamma_\mu e_t)$		
$Q_{q^2 e^2 D^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(\bar{e}_s \gamma_\mu \overleftrightarrow{D}_\nu e_t)$		
$Q_{q^2 u^2 D^2}^{(1)}$	$D^\nu(\bar{q}_p \gamma^\mu q_r) D_\nu(\bar{u}_s \gamma_\mu u_t)$		
$Q_{q^2 u^2 D^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(\bar{u}_s \gamma_\mu \overleftrightarrow{D}_\nu u_t)$		
$Q_{q^2 u^2 D^2}^{(3)}$	$D^\nu(\bar{q}_p \gamma^\mu T^A q_r) D_\nu(\bar{u}_s \gamma_\mu T^A u_t)$		
$Q_{q^2 u^2 D^2}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r)(\bar{u}_s \gamma_\mu T^A \overleftrightarrow{D}_\nu u_t)$		
$Q_{q^2 d^2 D^2}^{(1)}$	$D^\nu(\bar{q}_p \gamma^\mu q_r) D_\nu(\bar{d}_s \gamma_\mu d_t)$		
$Q_{q^2 d^2 D^2}^{(2)}$	$(\bar{q}_p \gamma^\mu \overleftrightarrow{D}^\nu q_r)(\bar{d}_s \gamma_\mu \overleftrightarrow{D}_\nu d_t)$		
$Q_{q^2 d^2 D^2}^{(3)}$	$D^\nu(\bar{q}_p \gamma^\mu T^A q_r) D_\nu(\bar{d}_s \gamma_\mu T^A d_t)$		
$Q_{q^2 d^2 D^2}^{(4)}$	$(\bar{q}_p \gamma^\mu T^A \overleftrightarrow{D}^\nu q_r)(\bar{d}_s \gamma_\mu T^A \overleftrightarrow{D}_\nu d_t)$		

Table 18. Most of the dimension-eight operators in the SMEFT of class-21, which are further divided into subclasses according to their chiral properties. See Table 10 for the remaining class-21 operators. Operators with + h.c. have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices.

5 Phenomenology

In this Section we briefly discuss a few aspects of phenomenology involving dimension-8 operators, focusing on processes that first start at dimension-8 and/or involve interplay between dimension-6 and -8 effects. In particular, we discuss light-by-light scattering and electroweak precision data. We also present a model where dimension-8 effects are arguably more important than dimension-6 effects.

Note that sometimes we will explicitly write factors of the cutoff scale of the effective theory, $C_i \rightarrow c_i/\Lambda^{d-4}$, to better highlight the different orders in the EFT expansion in our phenomenological studies.

5.1 Light-by-Light Scattering

The possibility of non-linear processes involving solely photons had been discussed back in the 1930s [27–30]. Later in the 1950s the cross section for elastic light-by-light scattering was computed in QED [31, 32]. Almost 70 years would pass before this process was finally observed in vacuum in 2019 by the ATLAS collaboration at the LHC [33]. Additionally, the CMS collaboration reports evidence for elastic light-by-light scattering in vacuum [34].

Interactions in the SMEFT involving four photons first start at dimension-8

$$\mathcal{L}_{LbL} = \frac{\alpha^2}{90M_e^4} \left[\mathcal{C}_{LbL}^{(1)} (F_{\mu\nu}F^{\mu\nu})^2 + \mathcal{C}_{LbL}^{(2)} (F_{\mu\nu}\tilde{F}^{\mu\nu})^2 + \tilde{\mathcal{C}}_{LbL} (F_{\mu\nu}F^{\mu\nu}) (F_{\rho\sigma}\tilde{F}^{\rho\sigma}) \right], \quad (5.1)$$

where $F_{\mu\nu}$ is the photon field strength, α is the fine structure constant, and M_e is the mass of the electron. The coefficients appearing in Eq. (5.1) are

$$\begin{aligned} \mathcal{C}_{LbL}^{(1)} &= 1 + \frac{90M_e^4}{\Lambda^4} 16\pi^2 c_{LbL}^{(1)}, \\ \mathcal{C}_{LbL}^{(2)} &= \frac{7}{4} + \frac{90M_e^4}{\Lambda^4} 16\pi^2 c_{LbL}^{(2)}, \\ \tilde{\mathcal{C}}_{LbL} &= \frac{90M_e^4}{\Lambda^4} 16\pi^2 \tilde{c}_{LbL}, \end{aligned} \quad (5.2)$$

with 1 and 7/4 being the leading order SM contributions [35]. The coefficient $\tilde{\mathcal{C}}_{LbL}$ is not generated at leading order in the SM. The combinations of Wilson coefficients from Table 2 appearing in Eq. (5.2) are

$$\begin{aligned} c_{LbL}^{(1)} &= \frac{1}{g_2^4} (c_{W^4}^{(1)} + c_{W^4}^{(3)}) + \frac{1}{g_1^4} c_{B^4}^{(1)} + \frac{1}{g_2^2 g_1^2} (c_{W^2 B^2}^{(1)} + c_{W^2 B^2}^{(3)}), \\ c_{LbL}^{(2)} &= \frac{1}{g_2^4} (c_{W^4}^{(2)} + c_{W^4}^{(4)}) + \frac{1}{g_1^4} c_{B^4}^{(2)} + \frac{1}{g_2^2 g_1^2} (c_{W^2 B^2}^{(2)} + c_{W^2 B^2}^{(4)}), \\ \tilde{c}_{LbL} &= \frac{1}{g_2^4} (c_{W^4}^{(5)} + c_{W^4}^{(6)}) + \frac{1}{g_1^4} c_{B^4}^{(3)} + \frac{1}{g_2^2 g_1^2} (c_{W^2 B^2}^{(5)} + c_{W^2 B^2}^{(6)} + c_{W^2 B^2}^{(7)}), \end{aligned} \quad (5.3)$$

where we have dropped terms of $\mathcal{O}(\Lambda^{-4})$ that are not enhanced by a factor of $16\pi^2$. If the physics beyond the SM (BSM) that generates (5.2) is loop suppressed then these additional terms must be included.

5.2 Electroweak Precision Data

The constraints on BSM physics from electroweak precision data are frequently summarized in terms of the parameters S , T , and U [36–41]. The leading contributions to S and T come from dimension-6 operators. However U first arises from a dimension-8 operator [20]. Here we work out the contributions to S , T , and U up to dimension-8, and point out a couple of subtleties that arise along the way.

Higher-dimensional operators change the definitions of SM parameters in a variety of ways. Field redefinitions are needed to related these combinations of inputs to measured quantities. The relationships are well known in the SM and have been worked out in the SMEFT as well [5, 20, 42, 43] including at dimension-8 [18, 44]. We now review the parts of this procedure that are necessary for defining S , T , and U .

The electroweak (EW) sector of the SMEFT is

$$\begin{aligned}\mathcal{L}_{EW}^{(d\leq 4)} &= -\frac{1}{4}W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} + (D_\mu H^\dagger)(D^\mu H) - \lambda \left(H^\dagger H - \frac{v^2}{2}\right)^2, \\ \mathcal{L}_{EW}^{(d=6)} &= \frac{c_H}{\Lambda^2}(H^\dagger H)^3 + \frac{c_{H\Box}}{\Lambda^2}(H^\dagger H)\Box(H^\dagger H) + \frac{c_{HD}}{\Lambda^2}[(D_\mu H^\dagger)H][H^\dagger(D^\mu H)] \\ &\quad + \frac{c_{HW}}{\Lambda^2}(H^\dagger H)W_{\mu\nu}^I W^{I\mu\nu} + \frac{c_{HB}}{\Lambda^2}(H^\dagger H)B_{\mu\nu}B^{\mu\nu} + \frac{c_{HWB}}{\Lambda^2}(H^\dagger \tau^I H)W_{\mu\nu}^I B^{\mu\nu}, \\ \mathcal{L}_{EW}^{(d=8)} &= \frac{c_{H^8}}{\Lambda^4}(H^\dagger H)^4 + \frac{c_{B^2H^4}^{(1)}}{\Lambda^4}(H^\dagger H)^2 B_{\mu\nu}B^{\mu\nu} + \frac{c_{WBH^4}^{(1)}}{\Lambda^4}(H^\dagger H)(H^\dagger \tau^I H)W_{\mu\nu}^I B^{\mu\nu} \\ &\quad + \frac{c_{W^2H^4}^{(1)}}{\Lambda^4}(H^\dagger H)^2 W_{\mu\nu}^I W^{I\mu\nu} + \frac{c_{W^2H^4}^{(3)}}{\Lambda^4}(H^\dagger \tau^I H)(H^\dagger \tau^J H)W_{\mu\nu}^I W^{J\mu\nu} \\ &\quad + \frac{c_{H^6D^2}^{(1)}}{\Lambda^4}(H^\dagger H)^2(D_\mu H^\dagger D^\mu H) + \frac{c_{H^6D^2}^{(2)}}{\Lambda^4}(H^\dagger H)(H^\dagger \tau^I H)(D_\mu H^\dagger \tau^I D^\mu H).\end{aligned}\tag{5.4}$$

The dimension-8 operators in (5.4) are given in Tables 2 and 3. There are several consequences of the higher-dimensional operators in (5.4).

The minimum of Higgs boson potential is shifted

$$\langle H^\dagger H \rangle \equiv \frac{v_T^2}{2} = \frac{v^2}{2} \left(1 + \frac{v^2}{\Lambda^2} \frac{3c_H}{4\lambda} + \frac{v^4}{\Lambda^4} \frac{9c_H^2 + 4\lambda c_{H^4}}{8\lambda^2} \right)\tag{5.5}$$

with $v_T \approx 246$ GeV.

The gauge fields need to be canonically normalized

$$B_\mu \rightarrow (1 + c_{B,kin})\mathcal{B}_\mu, \quad W_\mu^\pm \rightarrow (1 + c_{W^\pm,kin})\mathcal{W}_\mu^\pm, \quad W_\mu^3 \rightarrow (1 + c_{W^3,kin})\mathcal{W}_\mu^3,\tag{5.6}$$

where \mathcal{W}_μ^I and \mathcal{B}_μ are canonically normalized bosons. Foreshadowing what is to come $c_{W^\pm,kin} \neq c_{W^3,kin}$ due to terms arising at dimension-8

$$\begin{aligned}c_{B,kin} &= \frac{v_T^2}{\Lambda^2}c_{HB} + \frac{v_T^4}{2\Lambda^4}(3c_{HB}^2 + c_{B^2H^4}^{(1)}), \\ c_{W^\pm,kin} &= \frac{v_T^2}{\Lambda^2}c_{HW} + \frac{v_T^4}{2\Lambda^4}(3c_{HW}^2 + c_{W^2H^4}^{(1)}), \\ c_{W^3,kin} &= \frac{v_T^2}{\Lambda^2}c_{HW} + \frac{v_T^4}{2\Lambda^4}(3c_{HW}^2 + c_{W^2H^4}^{(1)} + c_{W^2H^4}^{(3)}).\end{aligned}\tag{5.7}$$

Canonically normalized fields in hand, we can define the S parameter

$$\Delta\mathcal{L} = -\frac{1}{2}\mathcal{W}_{\mu\nu}^+\mathcal{W}^{-\mu\nu} - \frac{1}{4}\mathcal{W}_{\mu\nu}^3\mathcal{W}^{3\mu\nu} - \frac{1}{4}\mathcal{B}_{\mu\nu}\mathcal{B}^{\mu\nu} - \frac{1}{2}\frac{S}{16\pi}\mathcal{W}_{\mu\nu}^3\mathcal{B}^{\mu\nu}. \quad (5.8)$$

The next step is to redefine the gauge couplings to try to compensate for the redefinitions of the gauge bosons we just made. For the hypercharge boson (and similarly for the gluon) by making the change $g_1 \rightarrow \bar{g}_1/(1 + c_{B,kin})$, the product of gauge boson and gauge couple is unchanged, $g_1 B_\mu = \bar{g}_1 \mathcal{B}_\mu$. However, starting at dimension-8, $c_{W^\pm,kin} \neq c_{W^3,kin}$, and so we must choose how to redefine g_2 . Following Ref. [18] we choose $g_2 \rightarrow \bar{g}_2/(1 + c_{W^\pm,kin})$. Given the covariant derivative

$$D_\mu = \partial_\mu + i\frac{\bar{g}_{2,\pm}}{\sqrt{2}}(\tau^+\mathcal{W}^+ + \tau^-\mathcal{W}^-) + i\frac{\bar{g}_{2,3}}{2}\tau^3\mathcal{W}^3 + \dots, \quad (5.9)$$

we can define the U parameter through the couplings that appear in it

$$\frac{U}{16\pi} = \frac{\bar{g}_{2,\pm}^2 - \bar{g}_{2,3}^2}{g_2^2}. \quad (5.10)$$

With our convention $\bar{g}_{2,\pm} = \bar{g}_2$. From this we see the physical impact of U is to break the universality in the coupling strength of the charged- and neutral-currents [20].

Lastly, consider the mass terms for the canonically normalized gauge bosons

$$\Delta\mathcal{L} = M_W^2\mathcal{W}_\mu^+\mathcal{W}^{-\mu} + \frac{1}{2}M_3^2\mathcal{W}_\mu^3\mathcal{W}^{3\mu} + \frac{1}{2}M_B^2\mathcal{B}_\mu\mathcal{B}^\mu + M_3M_B\mathcal{W}_\mu^3\mathcal{B}^\mu, \quad (5.11)$$

where the form of the last term on the right-hand side is fixed by electromagnetic gauge invariance. With this final definition in place we can define T as

$$\rho = 1 + \bar{\alpha} T = \frac{\bar{g}_{2,3}^2 M_W^2}{\bar{g}_{2,\pm}^2 M_3^2}. \quad (5.12)$$

where the shifted fine-structure constant is

$$\frac{1}{4\pi\bar{\alpha}} = \frac{1}{\bar{g}_2^2} + \frac{1}{\bar{g}_1^2} \quad (5.13)$$

Note this shift does not affect our analysis of light-by-light scattering; since all of those effects start at dimension-8 we can freely trade α for $\bar{\alpha}$.

With these definitions it is straightforward to work out the contributions to S , T , and U

$$\begin{aligned} \frac{1}{16\pi}S &= \frac{v_T^2}{\Lambda^2}c_{HWB} + \frac{v_T^4}{2\Lambda^4} \left[2c_{HWB}(c_{HW} + c_{HB}) + c_{WBH^4}^{(1)} \right], \\ \bar{\alpha} T &= -\frac{v_T^2}{2\Lambda^2}c_{HD} + \frac{v_T^4}{4\Lambda^4} \left(c_{HD}^2 - 2c_{H^6D^2}^{(2)} \right), \\ \frac{1}{16\pi}U &= -\frac{v_T^4}{\Lambda^4}c_{W^2H^4}^{(3)}. \end{aligned} \quad (5.14)$$

As we are working beyond leading order in the EFT expansion, changes in the definition of S and T can change the contributions arising from the square of dimension-6 terms, *e.g.* if $\rho \equiv 1 + \bar{\alpha} T$ then in general $\rho^{-1} \neq 1 - \bar{\alpha} T$. This subtlety does not affect contributions to S and T that are linear in Wilson coefficients, whether they are from dimension-6 or dimension-8 operators.

5.3 Scalar $SU(2)_w$ Quartets

Although the dimension-6 operators are formally the leading terms in the EFT expansion, there are various reasons why it may be necessary to consider dimension-8 effects. Here we explore a scenario where the difference in experimental precision in different classes of measurements causes dimension-8 effects to be important. The measurements are double Higgs boson production for which only upper limits exist on the cross section exist [45, 46], and EWPD, which as the name implies, is precisely measured. For example, one of the more precisely measured observables in this class is the width of the Z boson, which has a relative precision of $9 \cdot 10^{-4}$ [47].

The models under consideration add to the SM field content a new scalar field, Θ , that is an $SU(2)_w$ quartet with either $y = 3/2$ or $1/2$. The Lagrangian for the $y = 3/2$ case is

$$\begin{aligned} \mathcal{L}_\Theta = & D_\mu \Theta^\dagger D^\mu \Theta - M^2 \Theta^\dagger \Theta + [\lambda_1 \Theta_{jkm}^\dagger H^j H^k H^m + \text{h.c.}] \\ & - \lambda_{\alpha 1} (H^\dagger H) (\Theta^\dagger \Theta) - \lambda_{\alpha 2} (H_n^\dagger H^m) (\Theta_{jkm}^\dagger \Theta^{jkn}) + \mathcal{O}(\Theta^4). \end{aligned} \quad (5.15)$$

For the $y = 1/2$ case the term linear in Θ is instead $[\lambda_1 \Theta_{jkm}^\dagger H^j H^k \tilde{H}^m + \text{h.c.}]$. Note that Θ is completely symmetric in its $SU(2)_w$ indices. Assuming $M \gg v_T$ we integrate out Θ and match to the SMEFT. At dimension-6 only one operator, $(H^\dagger H)^3$, is generated (in both cases) [21, 48, 49]

$$C_H = \frac{|\lambda_1|^2}{M^2}. \quad (5.16)$$

At this stage the scalar quartets look like great candidates to enhance the double Higgs boson production rate.

However things are not so simple. When electroweak symmetry is broken the term in linear in Θ in Eq. (5.15) will force Θ to have a non-zero vacuum expectation value (vev). If Θ gets a vev, v_Θ , its quantum numbers dictate that it contributes to the T parameter [21]

$$1 + \bar{\alpha} T = \frac{v_T^2}{v_T^2 - 2v_\Theta^2 [\frac{3}{2}(\frac{3}{2} + 1) - 3y^2]} \quad (5.17)$$

With this in mind we extend the matching to dimension-8 starting with the $y = 3/2$ case. We find

$$\begin{aligned} \mathcal{L}_\Theta^{(d=8)} = & -\frac{\lambda_\alpha |\lambda_1|^2}{M^4} (H^\dagger H)^4 + \frac{3|\lambda_1|^2}{M^4} (H^\dagger H)^2 (D_\mu H^\dagger) (D^\mu H) \\ & + \frac{6|\lambda_1|^2}{M^4} (H^\dagger H) H_j^\dagger H^k (D_\mu H_k^\dagger) (D^\mu H^j) \end{aligned} \quad (5.18)$$

with $\lambda_\alpha = \lambda_{\alpha 1} + \lambda_{\alpha 2}$. The last term on the right-hand side of Eq. (5.18) is not in our operator basis. We use the Fierz identity for Pauli matrices, Eq. (3.1), to convert that operator into our basis, yielding for the $y = 3/2$ case

$$C_{H^8} = -\frac{\lambda_\alpha |\lambda_1|^2}{M^4}, \quad C_{H^6 D^2}^{(1)} = \frac{6|\lambda_1|^2}{M^4}, \quad C_{H^6 D^2}^{(2)} = \frac{3|\lambda_1|^2}{M^4}. \quad (5.19)$$

The $y = 1/2$ case is slightly more complicated

$$\begin{aligned}\mathcal{L}_\Theta^{(d=8)} = & -\frac{\lambda_\alpha |\lambda_1|^2}{M^4} (H^\dagger H)^4 + \frac{3|\lambda_1|^2}{M^4} (H^\dagger H)^2 (D_\mu H^\dagger)(D^\mu H) \\ & + \frac{2|\lambda_1|^2}{M^4} (H^\dagger H) \left[H_j^\dagger H^k (D_\mu H_k^\dagger)(D^\mu H^j) \right. \\ & \left. + H_j^\dagger H_k^\dagger (D_\mu H^k)(D^\mu H^j) + H^j H^k (D_\mu H_k^\dagger)(D^\mu H_j^\dagger) \right]\end{aligned}\quad (5.20)$$

Integrating the last two terms by parts, applying the Higgs boson EOM to fields with two derivatives acting on them, and using Eq. (3.1) we find

$$\begin{aligned}C_H = & \left(1 - \frac{2\lambda v^2}{M^2}\right) \frac{|\lambda_1|^2}{M^2}, \quad C_{H^8} = -(\lambda_\alpha + 4\lambda) \frac{|\lambda_1|^2}{M^4}, \quad C_{H^6 D^2}^{(2)} = -\frac{|\lambda_1|^2}{M^4} \\ C_{leH^5} = & -Y_e^\dagger \frac{|\lambda_1|^2}{M^4}, \quad C_{quH^5} = -Y_u^\dagger \frac{|\lambda_1|^2}{M^4}, \quad C_{qdH^5} = -Y_d^\dagger \frac{|\lambda_1|^2}{M^4}.\end{aligned}\quad (5.21)$$

We included the dimension-6 coefficient C_H here as it receives another contribution from mass term in the Higgs EOM, which was used to reduce a dimension-8 operator into operators in our basis. This completes the matching of the scalar quartet models to the SMEFT at dimension-8.

Using (5.14) we see that $T_{y=3/2} = -3T_{y=1/2} = -\frac{3}{2\alpha} |\lambda_1|^2 (\frac{v_T}{M})^4$. Our result for the contributions of these models to the T parameter agrees with what was found in Ref. [21]. The implication of our matching results is that these model cannot in fact provide a large enhancement to double Higgs boson production. This was unclear, from an EFT perspective at least, until the matching was done at dimension-8.

6 Conclusions

In this work we presented a complete basis of dimension-8 operators in the Standard Model Effective Field Theory. There are 1029 Lagrangian terms, 36 of which vanish in the absence of flavor structure. Multiple checks have been passed including that no operator in the basis can be removed completely using the equations of motion. We also presented a counting based argument that 1029 is the minimum number of Lagrangian terms needed to represent all of the dimension-8 SMEFT operators.

As a sample of what can be done with the a complete basis of dimension-8 operators, we briefly considered the phenomenology of light-by-light scattering and electroweak precision data, deriving S , T , and U to dimension-8. Additionally, we matched theories of $SU(2)_w$ quartets onto the SMEFT up to dimension-8 allowing to us showcase the interplay between dimension-6 effects and dimension-8 effects, the latter of which cannot be neglected in those models.

Note added: Ref. [50] is set to appear on the arXiv simultaneously with this work. It also presents a complete basis of dimension-8 SMEFT operators. The basis of Ref. [50] contains more Lagrangian terms than our basis as it limits its terms to have only one flavor representation, whereas we follow the convention of Ref. [2], combining flavor representations together when unambiguously possible to obtain the minimum number of terms.

Acknowledgements

We thank Renato Fonseca, Adam Martin, Verónica Sanz, Michael Trott, and especially Jing Shu for useful discussions. The author is grateful for the privilege of being able to stay home and work on a project like this.

A Dimension-6 and -7 Operators

For the sake of convenience we reproduce the tables of dimension-6 and -7 operators here. Table 19 contains the dimension-7 operators, and is adapted from Ref. [16]. The classification scheme we use for the dimension-7 comes from labelling the classes in Table 2 of Ref. [10] in descending order. Table 20 contains the dimension-6 operators, and is adapted from Ref. [5]. In contrast with [5] we also include in Table 20 the baryon number violating operators as listed in Ref. [6].

1 : $\psi^2 X H^2 + \text{h.c.}$		2 : $\psi^2 H^4 + \text{h.c.}$	
$Q_{l^2 W H^2}$	$\epsilon_{mn}(\tau^I \epsilon)_{jk}(l_p^m C i \sigma^{\mu\nu} l_r^j) H^n H^k W_{\mu\nu}^I$	$Q_{l^2 H^4}$	$\epsilon_{mn} \epsilon_{jk} (l_p^m C l_r^j) H^n H^k (H^\dagger H)$
$Q_{l^2 B H^2}$	$\epsilon_{mn} \epsilon_{jk} (l_p^m C i \sigma^{\mu\nu} l_r^j) H^n H^k B_{\mu\nu}$		
3(B) : $\psi^4 H + \text{h.c.}$		3(B) : $\psi^4 H + \text{h.c.}$	
$Q_{l^3 e H}$	$\epsilon_{jk} \epsilon_{mn} (\bar{e}_p l_r^j) (l_s^k C l_t^m) H^n$	$Q_{l u d^2 H}$	$\epsilon_{\alpha\beta\gamma} (\bar{l}_p d_r^\alpha) (u_s^\beta C d_t^\gamma) \tilde{H}$
$Q_{l e u d H}$	$\epsilon_{jk} (\bar{d}_p l_r^j) (u_s C e_t) H^k$	$Q_{l q^2 d H}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (\bar{l}_p d_r^\alpha) (q_{sm}^\beta C q_t^{j\gamma}) \tilde{H}^k$
$Q_{l^2 q d H}^{(1)}$	$\epsilon_{jk} \epsilon_{mn} (\bar{d}_p l_r^j) (q_s^k C l_t^m) H^n$	$Q_{l d^3 H}$	$\epsilon_{\alpha\beta\gamma} (\bar{l}_p d_r^\alpha) (d_s^\beta C d_t^\gamma) H$
$Q_{l^2 q d H}^{(2)}$	$\epsilon_{jm} \epsilon_{kn} (\bar{d}_p l_r^j) (q_s^k C l_t^m) H^n$	$Q_{e q d^2 H}$	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (\bar{e}_p q_r^{j\alpha}) (d_s^\beta C d_t^\gamma) \tilde{H}^k$
$Q_{l^2 q u H}$	$\epsilon_{jk} (\bar{q}_p^m u_r) (l_{sm} C l_t^j) H^k$		
4 : $\psi^2 H^3 D + \text{h.c.}$		5(B) : $\psi^4 D + \text{h.c.}$	
$Q_{l e H^3 D}$	$\epsilon_{mn} \epsilon_{jk} (l_p^m C \gamma^\mu e_r) H^n H^j i D_\mu H^k$	$Q_{l^2 u d D}$	$\epsilon_{jk} (\bar{d}_p \gamma^\mu u_r) (l_s^j C i D_\mu l_t^k)$
6 : $\psi^2 H^2 D^2 + \text{h.c.}$		5(B) : $\psi^4 D + \text{h.c.}$	
$Q_{l^2 H^2 D^2}^{(1)}$	$\epsilon_{jk} \epsilon_{mn} (l_p^j C D^\mu l_r^k) H^m (D_\mu H^n)$	$Q_{l q d^2 D}$	$\epsilon_{\alpha\beta\gamma} (\bar{l}_p \gamma^\mu q_r^\alpha) (d_s^\beta C i D_\mu d_t^\gamma)$
$Q_{l^2 H^2 D^2}^{(2)}$	$\epsilon_{jm} \epsilon_{kn} (l_p^j C D^\mu l_r^k) H^m (D_\mu H^n)$	$Q_{e d^3 D}$	$\epsilon_{\alpha\beta\gamma} (\bar{e}_p \gamma^\mu d_r^\alpha) (d_s^\beta C i D_\mu d_t^\gamma)$

Table 19. The dimension-seven operators in the SMEFT. The operators are divided into six classes according to their field content. The classes-3 and -5 are further divided into subclasses according to their baryon number. All of the operators have Hermitian conjugates. The subscripts p, r, s, t are weak-eigenstate indices. Operators below the dashed lines in classes-1 and -3 vanish when there is only one generation of fermions.

1 : X^3		2 : H^6		3 : $H^4 D^2$		5 : $\psi^2 H^3 + \text{h.c.}$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H	$(H^\dagger H)^3$	$Q_{H\Box}$	$(H^\dagger H)\Box(H^\dagger H)$	Q_{eH}	$(H^\dagger H)(\bar{l}_p e_r H)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$			Q_{HD}	$(H^\dagger D^\mu H)^* (H^\dagger D_\mu H)$	Q_{uH}	$(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W	$\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$					Q_{dH}	$(H^\dagger H)(\bar{q}_p d_r H)$
$Q_{\tilde{W}}$	$\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$						
4 : $X^2 H^2$		6 : $\psi^2 XH + \text{h.c.}$		7 : $\psi^2 H^2 D$			
Q_{HG}	$H^\dagger H G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{Hl}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$		
$Q_{H\tilde{G}}$	$H^\dagger H \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{Hl}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$		
Q_{HW}	$H^\dagger H W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{He}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$		
$Q_{H\tilde{W}}$	$H^\dagger H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$		
Q_{HB}	$H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$	$(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$		
$Q_{H\tilde{B}}$	$H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$		
Q_{HWB}	$H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$		
$Q_{H\tilde{W}B}$	$H^\dagger \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$	$i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$		
8 : $(\bar{L}L)(\bar{L}L)$		8 : $(\bar{R}R)(\bar{R}R)$		8 : $(\bar{L}L)(\bar{R}R)$			
Q_{ll}	$(\bar{l}_p \gamma^\mu l_r)(\bar{l}_s \gamma_\mu l_t)$	Q_{ee}	$(\bar{e}_p \gamma^\mu e_r)(\bar{e}_s \gamma_\mu e_t)$	Q_{le}	$(\bar{l}_p \gamma^\mu l_r)(\bar{e}_s \gamma_\mu e_t)$		
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{q}_s \gamma_\mu q_t)$	Q_{uu}	$(\bar{u}_p \gamma^\mu u_r)(\bar{u}_s \gamma_\mu u_t)$	Q_{lu}	$(\bar{l}_p \gamma^\mu l_r)(\bar{u}_s \gamma_\mu u_t)$		
$Q_{qq}^{(3)}$	$(\bar{q}_p \gamma^\mu \tau^I q_r)(\bar{q}_s \gamma_\mu \tau^I q_t)$	Q_{dd}	$(\bar{d}_p \gamma^\mu d_r)(\bar{d}_s \gamma_\mu d_t)$	Q_{ld}	$(\bar{l}_p \gamma^\mu l_r)(\bar{d}_s \gamma_\mu d_t)$		
$Q_{lq}^{(1)}$	$(\bar{l}_p \gamma^\mu l_r)(\bar{q}_s \gamma_\mu q_t)$	Q_{eu}	$(\bar{e}_p \gamma^\mu e_r)(\bar{u}_s \gamma_\mu u_t)$	Q_{qe}	$(\bar{q}_p \gamma^\mu q_r)(\bar{e}_s \gamma_\mu e_t)$		
$Q_{lq}^{(3)}$	$(\bar{l}_p \gamma^\mu \tau^I l_r)(\bar{q}_s \gamma_\mu \tau^I q_t)$	Q_{ed}	$(\bar{e}_p \gamma^\mu e_r)(\bar{d}_s \gamma_\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{u}_s \gamma_\mu u_t)$		
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma^\mu u_r)(\bar{d}_s \gamma_\mu d_t)$	$Q_{qu}^{(8)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{u}_s \gamma_\mu T^A u_t)$		
		$Q_{ud}^{(8)}$	$(\bar{u}_p \gamma^\mu T^A u_r)(\bar{d}_s \gamma_\mu T^A d_t)$	$Q_{qd}^{(1)}$	$(\bar{q}_p \gamma^\mu q_r)(\bar{d}_s \gamma_\mu d_t)$		
				$Q_{qd}^{(8)}$	$(\bar{q}_p \gamma^\mu T^A q_r)(\bar{d}_s \gamma_\mu T^A d_t)$		
8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$		8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$		8 : $(\bar{B}) + \text{h.c.}$			
Q_{ledq}	$(\bar{l}_p^j e_r)(\bar{d}_s q_{tj})$	$Q_{quqd}^{(1)}$	$(\bar{q}_p^j u_r) \epsilon_{jk} (\bar{q}_s^k d_t)$	Q_{duql}	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (d_p^\alpha C u_r^\beta) (q_s^{j\gamma} C l_t^k)$		
		$Q_{quqd}^{(8)}$	$(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k T^A d_t)$	Q_{qqqe}	$\epsilon_{\alpha\beta\gamma} \epsilon_{jk} (q_p^{j\alpha} C q_r^{k\beta}) (u_s^\gamma C e_t)$		
		$Q_{lequ}^{(1)}$	$(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$	Q_{qqql}	$\epsilon_{\alpha\beta\gamma} \epsilon_{mn} \epsilon_{jk} (q_p^{m\alpha} C q_r^{j\beta}) (q_s^{k\gamma} C l_t^n)$		
		$Q_{lequ}^{(3)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$	Q_{duue}	$\epsilon_{\alpha\beta\gamma} (d_p^\alpha C u_r^\beta) (u_s^\gamma C e_t)$		

Table 20. The dimension-six operators in the SMEFT. The operators are divided into eight classes according to their field content. The class-8 ψ^4 four-fermion operators are further divided into subclasses according to their chiral and baryonic properties. Operators with + h.c. have Hermitian conjugates, as does the $\psi^2 H^2 D$ operator Q_{Hud} . The subscripts p, r, s, t are weak-eigenstate indices.

References

- [1] S. Weinberg, *Baryon and Lepton Nonconserving Processes*, *Phys. Rev. Lett.* **43** (1979) 1566–1570.
- [2] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085, [[arXiv:1008.4884](#)].
- [3] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys.* **B268** (1986) 621–653.
- [4] L. F. Abbott and M. B. Wise, *The Effective Hamiltonian for Nucleon Decay*, *Phys. Rev.* **D22** (1980) 2208.
- [5] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **04** (2014) 159, [[arXiv:1312.2014](#)].
- [6] R. Alonso, H.-M. Chang, E. E. Jenkins, A. V. Manohar, and B. Shotwell, *Renormalization group evolution of dimension-six baryon number violating operators*, *Phys. Lett.* **B734** (2014) 302–307, [[arXiv:1405.0486](#)].
- [7] L. Lehman and A. Martin, *Hilbert Series for Constructing Lagrangians: expanding the phenomenologist’s toolbox*, *Phys. Rev.* **D91** (2015) 105014, [[arXiv:1503.07537](#)].
- [8] B. Henning, X. Lu, T. Melia, and H. Murayama, *Hilbert series and operator bases with derivatives in effective field theories*, *Commun. Math. Phys.* **347** (2016), no. 2 363–388, [[arXiv:1507.07240](#)].
- [9] L. Lehman and A. Martin, *Low-derivative operators of the Standard Model effective field theory via Hilbert series methods*, *JHEP* **02** (2016) 081, [[arXiv:1510.00372](#)].
- [10] B. Henning, X. Lu, T. Melia, and H. Murayama, *2, 84, 30, 993, 560, 15456, 11962, 261485, ...: Higher dimension operators in the SM EFT*, *JHEP* **08** (2017) 016, [[arXiv:1512.03433](#)]. [Erratum: *JHEP*09,019(2019)].
- [11] R. M. Fonseca, *The Sym2Int program: going from symmetries to interactions*, *J. Phys. Conf. Ser.* **873** (2017), no. 1 012045, [[arXiv:1703.05221](#)].
- [12] B. Gripaios and D. Sutherland, *DEFT: A program for operators in EFT*, *JHEP* **01** (2019) 128, [[arXiv:1807.07546](#)].
- [13] J. C. Criado, *BasisGen: automatic generation of operator bases*, *Eur. Phys. J.* **C79** (2019), no. 3 256, [[arXiv:1901.03501](#)].
- [14] C. B. Marinissen, R. Rahn, and W. J. Waalewijn, ..., *83106786, 114382724, 1509048322, 2343463290, 27410087742, ... Efficient Hilbert Series for Effective Theories*, [[arXiv:2004.09521](#)].
- [15] L. Lehman, *Extending the Standard Model Effective Field Theory with the Complete Set of Dimension-7 Operators*, *Phys. Rev.* **D90** (2014), no. 12 125023, [[arXiv:1410.4193](#)].
- [16] Y. Liao and X.-D. Ma, *Renormalization Group Evolution of Dimension-seven Baryon- and Lepton-number-violating Operators*, *JHEP* **11** (2016) 043, [[arXiv:1607.07309](#)].
- [17] A. Yu. Morozov, *Matrix of Mixing of Scalar and Vector Mesons of Dimension $D \leq 8$ in QCD. (In Russian)*, *Sov. J. Nucl. Phys.* **40** (1984) 505. [*Yad. Fiz.*40,788(1984)].

- [18] C. Hays, A. Martin, V. Sanz, and J. Setford, *On the impact of dimension-eight SMEFT operators on Higgs measurements*, *JHEP* **02** (2019) 123, [[arXiv:1808.00442](#)].
- [19] G. N. Remmen and N. L. Rodd, *Consistency of the Standard Model Effective Field Theory*, *JHEP* **12** (2019) 032, [[arXiv:1908.09845](#)].
- [20] B. Grinstein and M. B. Wise, *Operator analysis for precision electroweak physics*, *Phys. Lett.* **B265** (1991) 326–334.
- [21] S. Dawson and C. W. Murphy, *Standard Model EFT and Extended Scalar Sectors*, *Phys. Rev.* **D96** (2017), no. 1 015041, [[arXiv:1704.07851](#)].
- [22] R. M. Fonseca, *Enumerating the operators of an effective field theory*, *Phys. Rev. D* **101** (2020), no. 3 035040, [[arXiv:1907.12584](#)].
- [23] S. Alioli, R. Boughezal, E. Mereghetti, and F. Petriello, *Novel angular dependence in Drell-Yan lepton production via dimension-8 operators*, [arXiv:2003.11615](#).
- [24] G. N. Remmen and N. L. Rodd, *Flavor Constraints from Unitarity and Analyticity*, [arXiv:2004.02885](#).
- [25] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators I: Formalism and λ Dependence*, *JHEP* **10** (2013) 087, [[arXiv:1308.2627](#)].
- [26] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, *Spin flavor structure of large $N(c)$ baryons*, *Phys. Rev. D* **51** (1995) 3697–3727, [[hep-ph/9411234](#)].
- [27] O. Halpern, *Scattering processes produced by electrons in negative energy states*, *Phys. Rev.* **44** (1933), no. 10 855.2–856.
- [28] H. Euler and B. Kockel, *The scattering of light by light in Dirac’s theory*, *Naturwiss.* **23** (1935), no. 15 246–247.
- [29] W. Heisenberg and H. Euler, *Consequences of Dirac’s theory of positrons*, *Z. Phys.* **98** (1936), no. 11-12 714–732, [[physics/0605038](#)].
- [30] H. Euler, *On the scattering of light by light according to Dirac’s theory*, *Annalen Phys.* **26** (1936), no. 5 398–448. [*Annalen Phys.* 418, no. 5, 398 (1936)].
- [31] R. Karplus and M. Neuman, *Non-Linear Interactions between Electromagnetic Fields*, *Phys. Rev.* **80** (1950) 380–385.
- [32] R. Karplus and M. Neuman, *The scattering of light by light*, *Phys. Rev.* **83** (1951) 776–784.
- [33] **ATLAS** Collaboration, G. Aad et al., *Observation of light-by-light scattering in ultraperipheral Pb+Pb collisions with the ATLAS detector*, *Phys. Rev. Lett.* **123** (2019), no. 5 052001, [[arXiv:1904.03536](#)].
- [34] **CMS** Collaboration, A. M. Sirunyan et al., *Evidence for light-by-light scattering and searches for axion-like particles in ultraperipheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV*, *Phys. Lett.* **B797** (2019) 134826, [[arXiv:1810.04602](#)].
- [35] J. S. Schwinger, *On gauge invariance and vacuum polarization*, *Phys. Rev.* **82** (1951) 664–679. [*116*(1951)].
- [36] M. E. Peskin and T. Takeuchi, *A New constraint on a strongly interacting Higgs sector*, *Phys. Rev. Lett.* **65** (1990) 964–967.

- [37] W. J. Marciano and J. L. Rosner, *Atomic parity violation as a probe of new physics*, *Phys. Rev. Lett.* **65** (1990) 2963–2966. [Erratum: *Phys.Rev.Lett.* 68, 898 (1992)].
- [38] D. Kennedy and P. Langacker, *Precision electroweak experiments and heavy physics: A Global analysis*, *Phys. Rev. Lett.* **65** (1990) 2967–2970. [Erratum: *Phys.Rev.Lett.* 66, 395 (1991)].
- [39] B. Holdom and J. Terning, *Large corrections to electroweak parameters in technicolor theories*, *Phys. Lett. B* **247** (1990) 88–92.
- [40] M. Golden and L. Randall, *Radiative Corrections to Electroweak Parameters in Technicolor Theories*, *Nucl. Phys. B* **361** (1991) 3–23.
- [41] G. Altarelli and R. Barbieri, *Vacuum polarization effects of new physics on electroweak processes*, *Phys. Lett. B* **253** (1991) 161–167.
- [42] L. Berthier and M. Trott, *Towards consistent Electroweak Precision Data constraints in the SMEFT*, *JHEP* **05** (2015) 024, [[arXiv:1502.02570](#)].
- [43] I. Brivio, Y. Jiang, and M. Trott, *The SMEFTsim package, theory and tools*, *JHEP* **12** (2017) 070, [[arXiv:1709.06492](#)].
- [44] A. Helset, A. Martin, and M. Trott, *The Geometric Standard Model Effective Field Theory*, *JHEP* **03** (2020) 163, [[arXiv:2001.01453](#)].
- [45] **CMS Collaboration**, A. M. Sirunyan et al., *Combination of searches for Higgs boson pair production in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **122** (2019), no. 12 121803, [[arXiv:1811.09689](#)].
- [46] **ATLAS Collaboration**, G. Aad et al., *Search for non-resonant Higgs boson pair production in the $b\bar{b}\nu\bar{\nu}$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **801** (2020) 135145, [[arXiv:1908.06765](#)].
- [47] **ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group Collaboration**, S. Schael et al., *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257–454, [[hep-ex/0509008](#)].
- [48] B. Henning, X. Lu, and H. Murayama, *How to use the Standard Model effective field theory*, *JHEP* **01** (2016) 023, [[arXiv:1412.1837](#)].
- [49] J. de Blas, J. Criado, M. Perez-Victoria, and J. Santiago, *Effective description of general extensions of the Standard Model: the complete tree-level dictionary*, *JHEP* **03** (2018) 109, [[arXiv:1711.10391](#)].
- [50] H.-L. Li, Z. Ren, J. Shu, M.-L. Xiao, J.-H. Yu, and Y.-H. Zheng, *Complete Set of Dimension-8 Operators in the Standard Model Effective Field Theory*, to appear (2020).