

SUMS OF k -POTENT MATRICES

IVAN GARGATE AND MICHAEL GARGATE

ABSTRACT. We study sums of k -potent matrices. We show the conditions by which a complex matrix A can be expressed as a sums of k -potent matrices. Also we obtain conditions by which a complex matrix A can be expressed as a sum of finite order elements. This generalize some results obtain by Wu in [2]. Also we study the sum of k -potent matrices in $\mathcal{M}_{cf}(F)$ with F be a field and proof that any matrix in this space can be expressed as a sum of 14 $(k+1)$ -potent matrices preserving the result obtain by Slowik in [3].

1. INTRODUCTION

It is a classical question whether the elements of a ring or a group can be expressed as sums or products of elements of some particular set.

In the case of expressed matrices as a product, there are several authors who have works in this respect, for example see [6, 7, 10, 11, 12].

Recently the authors in [5] gave a way to express matrices with product of commutators of finite order.

In the case of expressed matrices as product of idempotents we can see [2]. In this context, there are various generalizations of concept of matrices idempotents, for instance see [8]. In this paper, the authors explored the k -potent matrices using the definition in [9] or the case $(1, k)$ in [8] and generalize the work done by Wu and Slowik in [2] and [3] respectively and show how to express matrices as sum of k -potent matrices.

Wu in [2] shows a necessary and sufficient condition for a complex square matrix be the sum of finitely many idempotent matrices. In the section 2 the authors generalize this work proving the following result:

Theorem 1.1. *A complex square matrix A is a sum of finitely many $(k+1)$ -potent if and only if there is an polynomial $F(x)$ with coefficient integers non-negatives and degree $\leq k-1$ such that $\text{tr} A = F(\omega)$ and $F(1) \geq \text{rank}(A)$.*

And in the gernal case, we show the following:

Theorem 1.2. *Let A be a complex square matrix and suppose that*

- 1.) *There are $\omega_1, \omega_2, \dots, \omega_n$ and $\beta_1, \beta_2, \dots, \beta_n$ integer numbers with $\omega_i \neq \omega_j$ and $\beta_i \neq \beta_j$ for $i \neq j$ such that ω_i is a β_i th root of unity for $i = 1, 2, \dots, n$.*

Date: November 27, 2024.

Key words and phrases. k -potent matrices; upper triangular matrices; finite order; column-finite matrices .

- 2.) There are polynomials $F_1(x), F_2(x), \dots, F_n(x)$ with positive coefficients integers and $1 \leq \deg(F_i) \leq \beta_i - 1$, such that

$$\text{tr}(A) = a_0 + F_1(\omega_1) + F_2(\omega_2) + \dots + F_n(\omega_n)$$

where a_0 is a positive integer number.

Then, A is a sum of finitely many $\beta_1 + 1, \beta_2 + 1, \dots, \beta_n + 1$ -potent matrices if and only if

$$a_0 + F_1(1) + F_2(1) + \dots + F_n(1) \geq \text{rank}(A).$$

And we enunciated some remarks about linear combinations of these special matrices.

In the section 3, the authors generalize the results of Slowik in [3] for the case of k -potent matrices on $\mathcal{M}_{cf}(F)$, proving the following result:

Theorem 1.3. *Let F any field. Any matrix $A \in \mathcal{M}_{cf}(F)$, can be expressed as a sum of at most 14 , $(k + 1)$ -potent matrices from $\mathcal{M}_{cf}(F)$.*

2. SUMS OF k -POTENT COMPLEX SQUARE MATRICES

In this section we consider matrices with complex entries or complex matrices. We start introducing the notation. Let $k \geq 2$ be an integer. If a square matrix A satisfies $A^k = A$, then A is said to be k -potent (see definition in [4]). For $k = 2$, A is said idempotent. A matrix A is called of order k if $A^k = I$, in the case that $k = 2$, A is called an involution.

In the following, $\text{tr}(A)$ denotes the trace of a matrix A , $\text{ran}(A)$ denotes its range, $\text{rank}(A)$ the dimension of $\text{ran}(A)$, and $\text{ker}(A)$ the kernel of A . Two matrices A and B are similar if $XA = BX$ for some nonsingular matrix X , they are unitarily equivalent if the above X can be chosen to be unitary. If A and B act on spaces H and K , respectively, then

$$A \oplus B = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$$

acts on $H \oplus K$, the orthogonal direct sum of H and K .

Let $\omega \in F$ be a k th root of unity with $\omega \neq 1$, then we have the following lemma:

Lemma 2.1. *Let A be an matrix $(k + 1)$ -potent and ω the k -root of unity. There is a unique polynomial $F(x)$ with non-negative integer coefficients and degree $\leq k - 1$ such that $\text{tr}(A) = F(\omega)$ and $\text{rank}(A) = F(1)$.*

Proof. Consider a $(k + 1)$ -potent matrix A , then their possible eigenvalues satisfies the equation $x^{k+1} - x = 0$, so they can assume the values $0, 1, \omega, \omega^2, \dots, \omega^{k-1}$ where ω is the k -root of unity. Then A is triangularizable and by similarity, the $\text{tr}(A)$ is a polynomial in ω with degree $\leq k - 1$ and define this polynomial as $F(x)$. The coefficient of the variable x^j in this polynomial represented the number of times that appears ω^j in the diagonal of the triangularization of A . So $F(1)$ represented the number of non zero

elements that appears in the diagonal of the triangularization of A and this number is the $\text{rank}(A)$. \square

In the following Lemma we show basic $(k+1)$ -potent matrices.

Lemma 2.2. *Let $\omega \in F$ be a k th root of unity. The matrices of the form*

$$A = \begin{bmatrix} 0 & & & & \\ & \ddots & & & \\ & & 0 & x & \\ & & & \omega & \\ & & & y & 0 \\ & & & & \ddots & \\ & & & & & 0 \end{bmatrix}$$

satisfies $X^k \neq X$ and $X^{k+1} = X$. So A is a $(k+1)$ -potent matrix.

Proof. For any j positive integer we have that

$$A^j = \begin{bmatrix} 0 & & & & \\ & \ddots & & & \\ & & 0 & x\omega^{j-1} & \\ & & & \omega^j & \\ & & & y\omega^{j-1} & 0 \\ & & & & \ddots & \\ & & & & & 0 \end{bmatrix}.$$

\square

In some cases it's not difficult to show that some matrices are sums of $(k+1)$ -potent matrices, see for example the following Remark.

Remark 1. If k is even, then we can expressed the identity matrix I_n as sum of $(k+1)$ -potent matrices, its follows from the property $1 + \omega + \omega^2 + \dots + \omega^{k-1} = 0$, in this case

$$\begin{aligned} I_n &= \sum_{i=1}^{k-1} (-\omega^i) \cdot I_n \\ &= \sum_{i=1}^{k-1} \sum_{j=1}^n \begin{bmatrix} 0 & & & & \\ & \ddots & & & \\ & & 0 & & \\ & & & -\omega^i & \\ & & & 0 & 0 \\ & & & & \ddots & \\ & & & & & j - \text{column} & \\ & & & & & & 0 \end{bmatrix} \end{aligned}$$

Then, I_n can be expressed as sum of $n(k-1)$, $(k+1)$ -potent matrices. Also, consider $A = \alpha I_n$ with α an positive integer number, then A can be expressed as sum of $\alpha n(k-1)$, $(k+1)$ -potent matrices.

Now we prove our first main result.

Proof of Theorem 1.1. For the case $k = 1$ it is proved by Wu in [2], so we consider the case that ≥ 2 .

[\Rightarrow] If $A = \sum_{j=1}^m E_j$, where each E_j is $(k+1)$ -potent, then by the Lemma 2.1 denote by $F_j(x)$ their respective polynomial. Then

$$\text{tr}(A) = \text{tr}\left(\sum_{j=1}^m E_j\right) = \sum_{j=1}^m \text{tr}(E_j) = \sum_{j=1}^m F_j(\omega).$$

Denote by $F(x) = \sum_{j=1}^m F_j(x)$ and observe that

$$F(1) = \sum_{j=1}^m F_j(1) = \sum_{j=1}^m \text{rank}(E_j) \geq \text{rank}\left(\sum_{j=1}^m E_j\right) = \text{rank}(A).$$

[\Leftarrow] Let $r = \text{rank}(A)$, then A is unitary equivalent to a matrix of the form

$$\begin{bmatrix} A_1 & 0 \\ A_2 & 0 \end{bmatrix}$$

with respect to the decomposition $H = \text{rank}(A^*) \oplus \ker(A)$, where A_1 is of size $r \times r$. We have two cases to consider

- (1.) A_1 is not a scalar matrix. By hipotesse, there is a polynomial $F(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{k-1}x^{k-1}$ with integer coefficients and $a_0, a_1, \dots, a_{k-1} > 0$. By the condition $F(1) = a_1 + a_2 + \cdots + a_{k-1} \geq r = \text{rank}(A)$ we can choose integer numbers r_0, r_1, \dots, r_{k-1} such that $a_i \geq r_i > 0$ and $r_0 + r_1 + \cdots + r_{k-1} = r$. Then for each $j = 0, 1, \dots, k-1$, since $a_j \geq r_j$ then we choose positive integers $a_j^{(1)}, a_j^{(2)}, \dots, a_j^{(r_j)}$ such that $a_j = a_j^{(1)} + a_j^{(2)} + \cdots + a_j^{(r_j)}$. Applying this decomposition, we have that

$$\text{tr}(A) = \sum_{i=1}^{r_0} a_0^{(i)} + \left(\sum_{i=1}^{r_1} a_1^{(i)}\right) \omega + \cdots + \left(\sum_{i=1}^{r_{k-1}} a_{k-1}^{(i)}\right) \omega^{k-1}.$$

By the Theorem 2 of Fillmore in [1], A_1 is similar to a matrix B of the form

$$\begin{bmatrix} \begin{bmatrix} a_0^{(1)} & & \\ & \ddots & \\ & & a_0^{(r_0)} \end{bmatrix} & & & * \\ & \begin{bmatrix} a_1^{(1)}\omega & & \\ & \ddots & \\ & & a_1^{(r_1)}\omega \end{bmatrix} & & \\ & & \ddots & \\ * & & & \begin{bmatrix} a_{k-1}^{(1)}\omega^{k-1} & & \\ & \ddots & \\ & & a_{k-1}^{(r_{k-1})}\omega^{k-1} \end{bmatrix} \end{bmatrix}$$

Then, A is similar to

$$C = \left[\begin{array}{c|c} B & 0 \\ \hline * & 0 \end{array} \right]$$

Denote by $X_0^{(1)}, X_0^{(2)}, \dots, X_0^{(r_0)}, X_1^{(1)}, X_1^{(2)}, \dots, X_1^{(r_1)}, \dots, X_{k-1}^{(1)}, X_{k-1}^{(2)}, \dots, X_{k-1}^{(r_{k-1})}$ the first $r = r_0 + r_1 + \dots + r_{k-1}$ column vectors of C , then C can be expressed as

$$\begin{aligned} C = & \sum_{i=1}^{r_0} \begin{bmatrix} 0 & \dots & 0 & X_0^{(i)} & 0 & \dots & 0 \end{bmatrix} \\ & + \sum_{i=1}^{r_1} \begin{bmatrix} 0 & \dots & 0 & X_1^{(i)} & 0 & \dots & 0 \end{bmatrix} \\ & + \dots + \sum_{i=1}^{r_{k-1}} \begin{bmatrix} 0 & \dots & 0 & X_{k-1}^{(i)} & 0 & \dots & 0 \end{bmatrix} \end{aligned}$$

where, the column $X_j^{(i)}$ appears in the position $r_0 + r_1 + \dots + r_{j-1} + i$, or also

$$\begin{aligned} C = & \sum_{i=1}^{r_0} a_0^{(i)} \begin{bmatrix} 0 & \dots & 0 & \frac{1}{a_0^{(i)}} X_0^{(i)} & 0 & \dots & 0 \end{bmatrix} \\ & + \sum_{i=1}^{r_1} a_1^{(i)} \begin{bmatrix} 0 & \dots & 0 & \frac{1}{a_1^{(i)}} X_1^{(i)} & 0 & \dots & 0 \end{bmatrix} \\ & + \dots + \sum_{i=1}^{r_{k-1}} a_{k-1}^{(i)} \begin{bmatrix} 0 & \dots & 0 & \frac{1}{a_{k-1}^{(i)}} X_{k-1}^{(i)} & 0 & \dots & 0 \end{bmatrix} \end{aligned}$$

Observe that, for $j > 0$, the matrices

$$\begin{bmatrix} 0 & \cdots & 0 & \frac{1}{a_j^{(i)}} X_j^{(i)} & 0 & \cdots & 0 \end{bmatrix}$$

are in the form of Lemma 2.2 and from here we have that these matrices are $(k+1)$ -potent and for $j = 0$, follows from Wu in [2] that these respective matrices are idempotents, so $(k+1)$ -potent matrices.

Then we have expressed C as a sum of $\sum_{i=1}^{r_0} a_0^{(i)} + \sum_{i=1}^{r_1} a_1^{(i)} + \cdots + \sum_{i=1}^{r_{k-1}} a_{k-1}^{(i)} = F(1)$ matrices $(k+1)$ -potent and by similarity we concluded that they also hold for A .

- (2.) If A_1 is a scalar matrix, then $\text{tr}(A) = F(1) = r$ and by Wu in [2] we have that A is sum of r idempotent matrices and so $(k+1)$ -potent matrices.

□

Now we prove our second main result.

Proof of Theorem 1.2. The necessity is similar to the proof of Theorem 1.1. For the converse, let $r = \text{rank}(A)$ then A is unitary equivalent to matrix of the form

$$\begin{bmatrix} A_1 & 0 \\ A_2 & 0 \end{bmatrix}$$

with A_1 an matrix of size $r \times r$. Choose positive integer numbers r_0, r_1, \dots, r_n such that $a_0 \geq r_0$, $F_i(1) \geq r_i$ for $i = 1, 2, \dots, n$ and $r_0 + r_1 + \cdots + r_n = \text{rank}(A)$.

Proceeding similar to the proof of Theorem 1.1 we have that A_1 is similar to an matrix

$$B = \begin{bmatrix} B_0 & & & * \\ & B_1 & & \\ & & \ddots & \\ * & & & B_n \end{bmatrix}$$

such that each B_i is a $r_i \times r_i$ matrix with $\text{tr}(B_0) = a_0$ and $\text{tr}(B_i) = F_i(\omega_i)$ for $i = 1, 2, \dots, n$. Observe that

$$B = \begin{bmatrix} B_0 & 0 & & 0 \\ * & 0 & & 0 \\ & & \ddots & \\ * & 0 & & 0 \end{bmatrix} + \begin{bmatrix} 0 & * & & 0 \\ 0 & B_1 & & 0 \\ & & \ddots & \\ 0 & * & & 0 \end{bmatrix} + \cdots + \begin{bmatrix} 0 & 0 & & * \\ 0 & 0 & & * \\ & & \ddots & \\ 0 & 0 & & B_n \end{bmatrix}$$

and applying the Theorem 1.1 in each to the matrices B_i we obtain the result.

□

Wu in [2] gave a result that A is idempotent if and only if $2A - I$ is an involution. Unfortunately, if $k > 2$ this equivalence is not known but in the following Lemmas we show various forms of construct these special matrices from the other.

Lemma 2.3. *If A is a matrix of order k then the matrix*

$$B = \frac{1}{k}(I + A + A^2 + \cdots + A^{k-1}),$$

is k -potent.

Proof. Its can proof using simple calculations. □

In the following Lemma we show how its possible construct new matrices of order k using k -potent matrices.

Lemma 2.4. *Let A be a matrix k -potent and ω a k th root of unity, then:*

1.) $\omega - 1$ is an solution of the equation

$$(1) \quad x^{k-1} + \binom{k}{1}x^{k-2} + \binom{k}{2}x^{k-3} + \cdots + \binom{k}{k-2}x + \binom{k}{k-1} = 0$$

and the matrix $B = (\omega - 1)A^{k-1} + I$ is a matrix of order k .

2.) *If k is even, then $1 + \omega$ is solution of the equation*

$$(2) \quad x^{k-1} - \binom{k}{1}x^{k-2} + \binom{k}{2}x^{k-3} - \cdots + \binom{k}{k-2}x - \binom{k}{k-1} = 0$$

and the matrix $C = (1 + \omega)A^{k-1} + I$ is a matrix of order k .

Proof. Since that $\omega^k = 1$ and using the Newton binomial formula, then

$$\begin{aligned} 1 &= \omega^k \\ 1 &= (\omega - 1 + 1)^k \\ 1 &= \sum_{i=0}^k \binom{k}{i} (\omega - 1)^i \\ 1 &= \sum_{i=1}^k \binom{k}{i} (\omega - 1)^i + 1 \\ 0 &= (\omega - 1) \left[\sum_{i=1}^k \binom{k}{i} (\omega - 1)^{i-1} \right], \end{aligned}$$

So, $\omega - 1$ satisfies the equation (1). Therefore, we observe that, for $i = 2, 3, \dots, k-1$ $(A^{k-1})^i = A^{(k-1)i} = A^{(ki-i)} = A^{(i-1)k+k-i} = (A^k)^{(i-1)} \cdot A^{k-i} =$

$A^{i-1} \cdot A^{k-i} = A^{k-1}$, and using the Newton binomial formula

$$\begin{aligned}
 B^k &= (\alpha A^{k-1} + I)^k \\
 &= \sum_{i=0}^k \binom{k}{i} (\alpha A^{k-1})^i \\
 &= \sum_{i=1}^k \binom{k}{i} \alpha^i A^{k-1} + I \\
 &= \alpha \left(\sum_{i=1}^k \binom{k}{i} \alpha^{i-1} \right) + I \\
 &= I,
 \end{aligned}$$

with $\alpha = \omega - 1$. The proof of item (2) is similar. \square

Using the simple matrices of Lemma 2.2 its possible construct $(k+1)$ -potent matrices.

Lemma 2.5. *Let ω be a k th root of unity and k an even number. With the matrices of the form*

$$A = \begin{bmatrix} 0 & & & & & \\ & \ddots & & & & \\ & & 0 & x & & \\ & & & \omega & & \\ & & & y & 0 & \\ & & & & & \ddots \\ & & & & & & 0 \end{bmatrix}$$

we can define $B = 2A - \omega I$, then B is a matrix of order k .

Proof. The matrix B is in the form

$$B = \begin{bmatrix} -\omega & & & & & \\ & \ddots & & & & \\ & & -\omega & 2x & & \\ & & & \omega & & \\ & & & 2y & -\omega & \\ & & & & & \ddots \\ & & & & & & -\omega \end{bmatrix}$$

and by simple verification we proof the result. \square

Applying the proof of the Theorem 1.1 and the Lemma 2.5 we can enunciated the following

Corollary 2.6. *Let A an $n \times n$ matrix and k an even number. Then A is the sum of finitely many matrices of order k if and only if there is an polynomial $F(x)$ of degree $< k-1$ and positive integer coefficients such that $\text{tr}(A) = F(\omega)$ with ω a k th root of unity.*

Proof. [\Leftarrow] The trace of an matrix of order k is an polynomial with integer coefficients and the same hold for sums of these elements.

[\Rightarrow] Consider $n \geq 2$ and choose an integer m such that $m \geq 2 - \frac{1}{n}F(1)$ with m and $F(1)$ are even or odd at the same time. Define $S = \frac{1}{2}(A + m\omega I)$ and we have that $tr(S) = \frac{1}{2}F(\omega) + \frac{1}{2}mn\omega$. Here define $G(x) = \frac{1}{2}F(x) + \frac{1}{2}mnx$ then $tr(S) = G(\omega)$ and $G(1) \geq n > rank(A)$, by the Theorem 1.1 we have that S is a sum of $(k+1)$ -potent matrices, $S = \sum_{i=1}^r E_i$ where the matrix E_i have the form of Lemma 2.5. Define $B_i = 2E_i - \omega I$ for $i = 1, 2, \dots, r$ then by the same Lemma, B_1, B_2, \dots, B_r are matrices of order k . So

$$\begin{aligned} T &= 2S - m\omega I \\ &= 2\left(\sum_{i=1}^r E_i\right) - m\omega I \\ &= \sum_{i=1}^r (2E_i - \omega I) + (r - m)\omega I \\ &= \sum_{i=1}^r B_i + (r - m)\omega I, \end{aligned}$$

is a sum of matrices of order k . \square

And, using the Theorem 1.2 and the last Corollary we get the following corollary for finite order matrices:

Corollary 2.7. *Let A be a complex square matrix. Then A is the sum of finitely many matrices of finite (different) order if and only if*

- 1.) *There are $\omega_1, \omega_2, \dots, \omega_n$ and $\beta_1, \beta_2, \dots, \beta_n$ all even integer numbers with $\omega_i \neq \omega_j$ and $\beta_i \neq \beta_j$ for $i \neq j$ such that ω_i is a β_i th root of unity for $i = 1, 2, \dots, n$.*
- 2.) *There are polynomials $F_1(x), F_2(x), \dots, F_n(x)$ with positive coefficients integers and $1 \leq \deg(F_i) \leq \beta_i - 1$, such that*

$$tr(A) = a_0 + F_1(\omega_1) + F_2(\omega_2) + \dots + F_n(\omega_n)$$

where a_0 is a positive integer number.

Proof. The proof follows immediately from the proofs of Theorem 1.2 and Corollary 2.6. \square

In the following Lemmas we show some results about k -potent matrices.

Lemma 2.8. *Consider ω a k th root of unity in F . Then the matrices*

$$B = \begin{bmatrix} a & -a \\ a - \omega & \omega - a \end{bmatrix} \quad \wedge \quad C = \begin{bmatrix} a & a \\ \omega - a & \omega - a \end{bmatrix},$$

satisfy the equations $X^k \neq X$ and $X^{k+1} = X$. Then both are $(k+1)$ -potent matrices.

Proof. Its no difficult proof that, for all $i < k$:

$$B^i = \begin{bmatrix} a\omega^{i-1} & -a\omega^{i-1} \\ a\omega^{i-1} - \omega^i & \omega^i - a\omega^{i-1} \end{bmatrix}.$$

And for C we have an identity similar. □

Lemma 2.9. *If $x \in F$, where F is a field, then the matrix*

$$A = \begin{bmatrix} x & 0 \\ 0 & 2\omega - x \end{bmatrix}$$

is the sum of two $(k+1)$ -potent matrices.

Proof. By the Lemma 2.8 is no difficult see that the matrix A is the sum of matrices B and C with $a = \frac{x}{2}$. □

Lemma 2.10. *Let A an matrix and ω a non-trivial k th root of unity, then $(\omega I + A) \oplus (\omega I - A)$ is a sum of two $(k+1)$ -potent matrices.*

Proof. Consider $X = \omega I + A$ then

$$(\omega I + A) \oplus (\omega I - A) = \begin{bmatrix} X & 0 \\ 0 & 2\omega I - X \end{bmatrix}$$

and the result follows from Lemma 2.9. □

With the above lemmas we obtain the following proposition.

Proposition 2.11. *Consider A an matrix $n \times n$, $n \geq 2$ with $\text{tr}(A) = \alpha\omega^j$ for some $\alpha > 3$ an integer number, ω an non-trivial k th root of unity and $1 < j < k$. Suppose that $\text{rank}(A) = 1$, then*

- 1.) *If α is an even number, then A is a sum of $\frac{\alpha-2}{2} + 2$, $(k+1)$ -potent matrices.*
- 2.) *If α is an odd number, then A is a sum of $\frac{\alpha-3}{2} + 3$, $(k+1)$ -potent matrices.*

Proof. In the case that α is an even number, then A is similar to

$$\begin{bmatrix} \alpha\omega^j & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix}$$

Then, we have

$$\begin{bmatrix} \alpha\omega^j & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} = \begin{bmatrix} \beta\omega^j & & 0 \\ & \beta\omega^j & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} + \begin{bmatrix} (\alpha-\beta)\omega^j & & 0 \\ & -\beta\omega^j & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix},$$

where $\beta = \frac{\alpha - 2}{2}$. The first matrix is a sum of β , $(k + 1)$ -potent matrices, and by the Lemma 2.10, the second matrix is a sum of two $(k + 1)$ -potent matrices. In the case that α is an odd number we can observe that

$$\begin{bmatrix} \alpha\omega^j & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} = \begin{bmatrix} \omega^j & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} + \begin{bmatrix} (\alpha - 1)\omega^j & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix},$$

where the first matrix is $(k + 1)$ -potent and, by (1), the second matrix is a sum of $\frac{\alpha - 3}{2} + 2$, $(k + 1)$ -potent matrices. □

In the geral case, we can enunciated the following proposition

Proposition 2.12. *Consider A an matrix $n \times n$, $n \geq 2$, ω an non-trivial k th root of unity and $tr(A) = \alpha_0 + \alpha_1\omega + \alpha_2\omega^2 + \cdots + \alpha_{k-1}\omega^{k-1}$ with $\alpha_i > 3$ for all $i = 0, 1, \dots, k-1$. If $rank(A) = k$ then A is a sum of $(\sum_{i=0}^{k-1} \gamma_i + 2)k$, $(k + 1)$ -potent matrices where*

$$\gamma_i = \begin{cases} \frac{\alpha_i - 2}{2} & , \text{ if } \alpha_i \text{ is an even number,} \\ \frac{\alpha_i - 3}{2} + 3 & , \text{ if } \alpha_i \text{ is an odd number.} \end{cases}$$

Proof. Consider $tr(A) = F(\omega)$ then $F(1) = \sum_{i=0}^{k-1} \alpha_i \geq rank(A) = k$, then by Fillmore in [1], A is similar to

$$\begin{bmatrix} \begin{bmatrix} \alpha_0 & & & 0 \\ & \alpha_1\omega & & \\ & & \alpha_2\omega^2 & \\ & & & \ddots \\ & & & & \alpha_{k-1}\omega^{k-1} \end{bmatrix}_{k \times k} & & \\ & \ddots & \\ & & 0 \end{bmatrix}$$

And, using the Proposition 2.11 in each element in the diagonal we concluded the proof. □

Remark 2. Follows of the proof of Proposition 2.12 that if $tr(A) = \alpha_0 + \alpha_1\omega + \cdots + \alpha_{k-1}\omega^{k-1}$ with $0 < j < k - 1$ of α'_i 's are zero and $k - j$ are > 3 .

If $rank(A) = k - j$, then A is a sum of $\sum_{i=0 \wedge \alpha_i \neq 0}^{k-1} \gamma_i(k - j)$, $(k + 1)$ -potent matrices.

Using the above proposition we can enunciated the more geral case.

Proposition 2.13. *Consider A an matrix $n \times n$, $n \geq 2$, and suppose that*

- 1.) *There are $\omega_1, \omega_2, \dots, \omega_n$ and $\beta_1, \beta_2, \dots, \beta_n$ integer numbers with $\omega_i \neq \omega_j$ and $\beta_i \neq \beta_j$ for $i \neq j$ such that ω_i is a β_i th root of unity for $i = 1, 2, \dots, n$.*
- 2.) *There are polynomials $F_1(x), F_2(x), \dots, F_n(x)$ with $1 \leq \text{degree}(F_i) \leq \beta_i - 1$ such that $F_i(x) = \alpha_{i1}x + \alpha_{i2}x^2 + \dots + \alpha_{i(\beta_i-1)}x^{\beta_i-1}$ with all α_{ij} positive coefficients integers and*

$$\text{tr}(A) = a_0 + F_1(\omega_1) + F_2(\omega_2) + \dots + F_n(\omega_n)$$

where a_0 is a positive integer number.

If $\text{rank}(A) = 1 + \sum_{i=1}^n (\beta_i - 1)$, then A is a sum of $\left\{ \gamma_0 + \sum_{i=1}^n \sum_{j=1}^{\beta_i-1} (\gamma_{ij} + 2) \right\} \times \text{rank}(A)$ many $\beta_1, \beta_2, \dots, \beta_n$ -potent matrices where

$$\gamma_0 = \begin{cases} \frac{\alpha_0 - 2}{2} & , \text{ if } \alpha_0 \text{ is an even number,} \\ \frac{\alpha_0 - 3}{2} + 3 & , \text{ if } \alpha_0 \text{ is an odd number.} \end{cases}$$

and, for $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, \beta_i$,

$$\gamma_{ij} = \begin{cases} \frac{\alpha_{ij} - 2}{2} & , \text{ if } \alpha_{ij} \text{ is an even number,} \\ \frac{\alpha_{ij} - 3}{2} + 3 & , \text{ if } \alpha_{ij} \text{ is an odd number.} \end{cases}$$

Proof. Follows immediately from Theorem 1.2 and the Proposition 2.12. \square

Now, we can observe the following situations.

Example 2.14. *Consider A a complex matrix 3×3 such that $\text{Trace}(A) = 4 - 6\omega + 10\omega^2$ with ω a 3th root of unity. Suppose that $\text{rank}(A) = 3$ then we can expressed A as a sums and differences of 4-potent matrices. This is possible because A is similar to*

$$B = \begin{bmatrix} 4 & & * \\ & -6\omega & \\ * & & 10\omega \end{bmatrix}$$

and

$$B = 4 \begin{bmatrix} 1 & 0 & 0 \\ * & 0 & 0 \\ * & 0 & 0 \end{bmatrix} - 6 \begin{bmatrix} 0 & * & 0 \\ 0 & \omega & 0 \\ 0 & * & 0 \end{bmatrix} + 10 \begin{bmatrix} 0 & 0 & * \\ 0 & 0 & * \\ 0 & 0 & \omega \end{bmatrix}.$$

Then we can conclude the our affirmation since the first matrix is idempotent and the second and third matrices are 4-potent

Example 2.15. *If in the Example 2.14 we suppose that $\text{Trace}(A) = 4 - \frac{9}{2}\omega + \frac{25}{4}\omega^2$ then we obtain that B is a combination linear of one idempotent and two 4-potent matrices.*

Using the above example we can easily extend all results for matrices with all coefficients that appears in their respective trace no necessarily positives. First, consider $F(x) = \alpha_0 + \alpha_1 x + \cdots + \alpha_n x^n$ an polynomial with non-zero coefficients, define the module of polynomial F as $|F| = |\alpha_0| + |\alpha_1| + \cdots + |\alpha_n| = \sum_{i=0}^n |\alpha_i|$ and the number $Min(F) = \min \{|\alpha_j|\}_{j=0}^n$

Corollary 2.16. *Let A be a complex square matrix and suppose that*

- 1.) *There are $\omega_1, \omega_2, \dots, \omega_n$ and $\beta_1, \beta_2, \dots, \beta_n$ integer numbers with $\omega_i \neq \omega_j$ and $\beta_i \neq \beta_j$ for $i \neq j$ such that ω_i is a β_i th root of unity for $i = 1, 2, \dots, n$.*
- 2.) *There are polynomials $F_1(x), F_2(x), \dots, F_n(x)$ with non-zero coefficients, $Min(F_i) \geq 1$ and $1 \leq \text{degree}(F_i) \leq \beta_i - 1$ for $i = 1, 2, \dots, n$, such that*

$$\text{tr}(A) = a_0 + F_1(\omega_1) + F_2(\omega_2) + \cdots + F_n(\omega_n)$$

where a_0 satisfies $|a_0| \geq 1$.

Then, A is a linear combination of finitely many $\beta_1 + 1, \beta_2 + 1, \dots, \beta_n + 1$ -potent matrices if and only if

$$|a_0| + |F_1| + |F_2| + \cdots + |F_n| \geq \text{rank}(A)$$

Proof. Follow immediately from Theorem 1.2 and the above examples. \square

And so is possible enunciate similar results for the case of linear combination of matrices of finite order (see Corollary 2.7) and count the quantity necessary of k -potent matrices for obtain a linear combination in special cases for the rank of A (see Proposition 2.13). We leave this work for the reader.

3. SUMS OF k -POTENT MATRICES IN $\mathcal{M}_{cf}(F)$

In this section we present some results that generalized the paper of Slowik in [3]. We write e_∞ and e_k for $\mathbb{N} \times \mathbb{N}$ and $k \times k$ identity matrices, respectively, and e_{nm} for the $\mathbb{N} \times \mathbb{N}$ matrix with 1 in the position (n, m) and 0 in every other position.

Let $\mathcal{M}_{cf}(F)$ be the set of all column-finite $\mathbb{N} \times \mathbb{N}$ matrices over a field F . Denote $\mathcal{T}_\infty(F)$ the subring of $\mathcal{M}_{cf}(F)$ consisting of all upper triangular matrices and $\mathcal{LT}_{cf}(F)$ the subring of all column-finite lower triangular matrices. Here $\mathcal{T}_n(F)$ and $\mathcal{LT}_n(F)$ will be used for the rings of all $n \times n$ upper or lower triangular matrices respectively, whereas $\mathcal{D}_n(F)$ will denote the ring of all $n \times n$ diagonal matrices. The full $n \times n$ matrix ring will be denoted by $\mathcal{M}_{n \times n}(F)$.

3.1. Triangular matrices. Consider $\omega \in F$ a k th root of unity, $\omega \neq 1$. Here we have the following lemma

Lemma 3.1. *Let F a field and let $t \in \mathcal{T}_\infty(F)$. If*

$$t = \omega e_\infty + \sum_{m-n \geq 1} t_{nm} e_{nm}$$

and $t_{n,n+1} \neq 0$ for all $n \in \mathbb{N}$, then t is similar to

$$\omega e_\infty + \sum_{n=1}^{\infty} t_{n,n+1} e_{n,n+1}.$$

Proof. Let $t = \omega e_\infty + \sum_{m-n \geq 1} t_{nm} e_{nm}$. Define $u = \frac{1}{\omega} \cdot t$, in this case u are in the conditions of Lemma 2.1 in [3], so u is similar to

$$e_\infty + \sum_{n=1}^{\infty} \frac{1}{\omega} \cdot t_{n,n+1} e_{n,n+1}.$$

Therefore, t is similar to

$$\omega e_\infty + \sum_{n=1}^{\infty} t_{n,n+1} e_{n,n+1},$$

what demonstrates the Lemma. \square

Next, proof the following proposition

Proposition 3.2. *Let F be a field. If $t \in \mathcal{T}_\infty(F)$ satisfies the condition $t_{nn} = 2\omega$ for all $n \in \mathbb{N}$, then t is a sum of at most four $(k+1)$ -potent matrices.*

Proof. Following the Slowik demonstration of Lemma 2.1 in [3]. First we define two matrices t_1 and t_2 as follows:

$$(t_1)_{nm} = \begin{cases} \omega & \text{if } m = n, \\ t_{nm} & \text{if } m - n > 1, \\ t_{nm} & \text{if } m - n = 1, \text{ and } t_{nm} \neq 0, \\ \omega & \text{if } m - n = 1, \text{ and } t_{nm} = 0, \end{cases} \quad t_2 = t - t_1$$

One can see that t_1 fulfills the assumptions of Lemma 3.1. Hence, t_1 is similar to the matrix

$$\omega e_\infty + \sum_{n=1}^{\infty} (t_1)_{n,n+1} e_{n,n+1},$$

and we write this matrix as the sum $u + u'$, where

$$u = \sum_1^{\infty} (\omega e_{2n-1,2n-1} + (t_1)_{2n-1,2n} e_{2n-1,2n}) \text{ and } u' = \sum_1^{\infty} (\omega e_{2n,2n} + (t_1)_{2n,2n+1} e_{2n,2n+1}),$$

it is easy to check that u and u' are $(k+1)$ -potent matrices.

Now we will write $t_2 = v' + v''$ where v' is defined by the following inductive rule:

$$(1) \ v'_{11} = (t_2)_{11}, \ v'_{12} = (t_2)_{12};$$

- (2) if $v'_{n,n+1} = 0$, then put we $v'_{n+1,n+1} = \omega$, $v'_{n+1,n+2} = (t_2)_{n+1,n+2}$;
- (3) if $v'_{n,n+1} \neq 0$, then we put $v'_{n+1,n+1} = 0$, $v'_{n+1,n+2} = 0$.

This it, if $v'_{n,n+1} \neq 0$, then the entries from the next row 'go' to v'' . If $v'_{n,n+1} = 0$, then the next row can 'stay' in v' .

From the construction of v' and v'' it follows that they are $(k+1)$ -potent matrices. □

Corollary 3.3. *Let F be a field and $k \in \mathbb{N}$. If t is either from $\mathcal{T}_\infty(F)$ or $\mathcal{LT}_k(F)$ and $t_{nn} = 2\omega$ for all n , $1 \leq n \leq k$, then t is a sum of at most four $(k+1)$ -potent matrices.*

The above corollary will be useful in the proof of the below result.

Proposition 3.4. *If F is a field and $t \in \mathcal{LT}_{Cf}(F)$ is such that $t_{nn} = 3\omega$ for all $n \in \mathbb{N}$, then t can be written as a sum of at most six $(k+1)$ -potent matrices.*

Proof. We apply the method used in the proof of Proposition 2.2 from [3]. First we construct the sequence $(l_m)_{m \in \mathbb{N}}$ as follows:

$$l_m = \max\{i : t_{im} \neq 0\}.$$

Now we define another sequence $(l'_m)_{m \in \mathbb{N}}$ as follows:

$$l'_m = \begin{cases} l_m & \text{if for all } j < m \text{ we have } l_j \leq l_m, \\ \max\{l_j : j < m \wedge l_j > l_m\} & \text{otherwise} \end{cases}$$

The sequence $(l'_m)_{m \in \mathbb{N}}$ is nondecreasing. Observed that for all $m \in \mathbb{N}$ and all $i > l'_m$ we have $t_{im} = 0$. Thus, we can say that t has staircase structure and that the stairs are determined by the sequence (l'_m) .

Now we will write $t = u + v$ where the matrix u is constructed as follows. For all $i \in \mathbb{N}$ we put $u_{ii} = 2\omega$. Now let m be equal to 1 and let n be equal to l'_1 . For all i, j such that $m \leq j < i \leq n$ we put $u_{ij} = t_{ij}$.

Let now m be equal to the preceding n increased by 1 and let the new n be equal to l'_m . Then for all $i, j, m \leq i, j \leq n$ we put $u_{ij} = t_{ij}$.

Proceeding the same way we obtain a block diagonal matrix u and consider $v = t - u$. By Corollary 3.3 each block of u is a sum of at most four $(k+1)$ -potent matrices. From the construction of v it follows that

$$v = \begin{bmatrix} \omega & & & & \\ v_1 & \omega & & & \\ 0 & v_2 & \omega & & \\ 0 & 0 & v_3 & \omega & \\ 0 & 0 & 0 & v_4 & \omega \\ \vdots & \vdots & & \ddots & \ddots \end{bmatrix}$$

and

$$v' = \begin{bmatrix} \omega & & & & \\ v_1 & 0 & & & \\ 0 & 0 & \omega & & \\ 0 & 0 & v_3 & 0 & \\ 0 & 0 & 0 & 0 & \omega \\ \vdots & \vdots & & \ddots & \ddots \end{bmatrix} \quad v'' = v - v' = \begin{bmatrix} 0 & & & & \\ 0 & \omega & & & \\ 0 & v_2 & 0 & & \\ 0 & 0 & 0 & \omega & \\ 0 & 0 & 0 & v_4 & 0 \\ \vdots & \vdots & & \ddots & \ddots \end{bmatrix}.$$

It is easy to check that v', v'' are $(k+1)$ -potent matrices. \square

3.2. Diagonal Matrices. For $\mathcal{D}_\infty(F)$ we have the following proposition:

Proposition 3.5. *Let F any field. Any $D \in \mathcal{D}_\infty(F)$ is the sum of four $(k+1)$ -potent matrices from $\mathcal{M}_{cf}(F)$.*

Proof. The proof follows from Slowik in [3]. For $D = \text{diag}(d_{11}, d_{22}, d_{33}, \dots)$ define the matrices $X, Y \in \mathcal{D}_\infty(F)$ inductively as

$$\begin{aligned} x_{11} &= d_{11}, & y_{11} &= 0, \\ x_{22} &= 2\omega - x_{11}, & y_{22} &= d_{22} - x_{22}, \\ y_{33} &= 2\omega - y_{22}, & x_{33} &= d_{33} - y_{33}, \\ x_{44} &= 2\omega - x_{33}, & y_{44} &= d_{44} - x_{44}, \\ \vdots & & \vdots & \end{aligned}$$

Then, X and Y are in the form

$$\begin{aligned} X &= \begin{bmatrix} x_{11} & 0 \\ 0 & 2\omega - x_{11} \end{bmatrix} \oplus \begin{bmatrix} x_{33} & 0 \\ 0 & 2\omega - x_{33} \end{bmatrix} \oplus \dots \\ Y &= [0] \oplus \begin{bmatrix} y_{22} & 0 \\ 0 & 2\omega - y_{22} \end{bmatrix} \oplus \begin{bmatrix} y_{44} & 0 \\ 0 & 2\omega - y_{44} \end{bmatrix} \oplus \dots \end{aligned}$$

and, by the Lemma 2.9, both X and Y are the sums of $(k+1)$ -potent matrices. \square

Now, we can prove our following result.

Proof of Theorem 1.3. Define t_1, t_2 and d as follows:

$$(t_1)_{ij} = \begin{cases} 2\omega & \text{if } j = i \\ a_{ij} & \text{if } i < j \\ 0 & \text{if } i > j \end{cases} \quad (t_2)_{ij} = \begin{cases} 3\omega & \text{if } j = i \\ a_{ij} & \text{if } i > j \\ 0 & \text{if } i < j \end{cases} \quad d_{ij} = \begin{cases} a_{ij} - 5\omega & \text{if } j = i \\ 0 & \text{if } i \neq j \end{cases}$$

Clearly $A = t_1 + t_2 + d$. From Propositions 3.2 and 3.4 we know that t_1 and t_2 are sums of at most four and six $(k+1)$ -potent matrices, respectively. By Proposition 3.5 the matrix d is a sum of four $(k+1)$ -potent matrices. This proves the claim. \square

REFERENCES

- [1] P. A. Fillmore, On similarity and the diagonal of a matrix, Amer. Math. Monthly 76, 167-169 (1969).
- [2] P. Y. Wu, Sums of idempotent matrices. Linear Algebra Appl., 142, 43-54 (1990).
- [3] R. Slowik. Expressing Infinite Matrices as Sums of Idempotents. Ukranian Mathematical Journal. Vol 69, No 8. 1145-1152.(2017).
- [4] Z. Huang, H. Lin. On k -idempotent 0-1 matrices. Linear Algebra and its Applications. 597 (2020). 1987-197.
- [5] Ivan Gargate, Michael Gargate. Expressing finite-infinite matrices into products of commutators of finite order elements. arXiv:2004.09012v1. April 2020
- [6] N. H. Gustafson, On products of involutions, in: J.H. Ewing, F.W. Gehring (Eds.), Paul Halmos, Celebrating 50 Years of Mathematics, Springer-Verlag, New York, 1991, pp. 237-255.
- [7] R. Slowik. Expressing infinite matrices as product of involutions. Linear Algebra Appl. 438 (2013) 399-404.
- [8] M. Jeter, W. Pye, Nonnegative (s,t)-potent matrices. Linear Algebra Appl. 45. (1982), 109-121.
- [9] R. Huang, J. Liu, L. Zhu. A structural characterization of real k -potent matrices. Linear and Multilinear Algebra, 59(4). 433-439, (2011).
- [10] B. Zheng, Decomposition of matrices into commutators of involutions, Linear Algebra Appl. 347 (2002) 1-7.
- [11] Hou, X. Decomposition of infinite matrices into products of commutators of involutions. Linear Algebra and Its Applications. doi:10.1016/j.laa.2018.11.001
- [12] W. H. Gustafson, P. R. Halmos, H. R. Adjavi, Products of involutions. Collection of articles dedicated to Olga Taussky Todd., Linear Algebra Appl. 13, 157-163 (1976).

UTFPR, CAMPUS PATO BRANCO, RUA VIA DO CONHECIMENTO KM 01, 85503-390
PATO BRANCO, PR, BRAZIL

E-mail address: `ivangargate@utfpr.edu.br`

UTFPR, CAMPUS PATO BRANCO, RUA VIA DO CONHECIMENTO KM 01, 85503-390
PATO BRANCO, PR, BRAZIL

E-mail address: `michaelgargate@utfpr.edu.br`