Description of the newly observed Ξ_c^0 states as molecular states

HongQiang Zhu¹, NaNa Ma², and Yin Huang^{3a}

¹ College of Physics and Electronic Engineering, Chongqing Normal Universit

² School of Nuclear Science and Technology, Lanzhou University, 730000 Lanzhou, People's Republic of China

³ School of Physical Science and Technology, Southwest Jiaotong University, Chengdu 610031, China

Received: date / Revised version: date

Abstract. Very recently, three new structures $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ at the invariant mass spectrum of $\Lambda_c^+K^-$ observed by the LHCb Collaboration triggers a hot discussion about their inner structure. In this work, we study the strong decay mode of the newly observed Ξ_c assuming that the Ξ_c is a $\bar{D}\Lambda - \bar{D}\Sigma$ molecular state. With the possible quantum numbers $J^p = 1/2^{\pm}$ and $3/2^{\pm}$, the partial decay widths of the $\bar{D}\Lambda - \bar{D}\Sigma$ molecular state into the $\Lambda_c^+K^-$, $\Sigma_c^+K^-$, and $\Xi_c^+\pi^-$ final states through hadronic loop are calculated with the help of the effective Lagrangians. By comparison with the LHCb observation, the current results of total decay width support the $\Xi(2923)^0$ as $\bar{D}\Lambda - \bar{D}\Sigma$ molecule while the decay width of the $\Xi_c(2938)^0$ and $\Xi(2964)^0$ can not be well reproduced in the molecular state picture. In addition, the calculated partial decay widths with S wave $\bar{D}\Lambda - \bar{D}\Sigma$ molecular state picture indicate that allowed decay modes, $\Xi_c^+\pi^-$, may have the biggest branching ratios for the $\Xi_c(2923)$. The experimental measurements for this strong decay process could be a crucial test for the molecule interpretation of the $\Xi_c(2923)$.

PACS. 13.60.Le decay widths – 13.85.Lg mass spectrum – 25.30.-c molecular state

1 Introduction

During the past several decades, many narrow baryons with a heavy charm quark, a light up or down quark, and a strange quark have been reported by the LHCb,CDF Collaboration and so on [1]. Very recently, three other neutral resonances Ξ_c^{*0} named $\Xi_c(2923)^0$, $\Xi_c(2939)^0$, and $\Xi_c(2965)^0$ have been observed in the $K^-\Lambda_c^+$ mass spectra by the LHCb Collaboration [2]. The observed resonance masses and widths are

$$\begin{split} \mathbf{M} &= 2923.04 \pm 0.25(stat) \pm 0.20(syst) \pm 0.14(\Lambda_c^+) \ \mathrm{MeV} \\ \Gamma &= 7.1 \pm 0.8(stat) \pm 1.8(syst) \ \mathrm{MeV}, \end{split}$$

$$M = 2938.55 \pm 0.21(stat) \pm 0.17(syst) \pm 0.14(\Lambda_c^+) \text{ MeV}$$

$$\Gamma = 10.2 \pm 0.8(stat) \pm 1.1(syst) \text{ MeV},$$

$$M = 2964.88 \pm 0.26(stat) \pm 0.14 \pm (syst)0.14(\Lambda_c^+) \text{ MeV}$$

$$\Gamma = 14.1 \pm 0.9(stat) \pm 1.3(syst) \text{ MeV},$$

respectively. From the observed decay mode, the isospin of these three states are 1/2. Although the quantum numbers of these states remain undetermined, it is very helpful to understand the spectroscopy of the heavy baryons containing c and s quark.

Due to their observed decay mode, the new structures $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ contain at least three different valence quark components. In other word, these states may be candidates of conventional three-quark state. Indeed, the QCD sum rule suggests that the newly observed states $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ are most likely to be considered as the *P*-wave Ξ'_c baryons with the spin-parity $J^p = 1/2^-$ or $J^p = 3/2^-$ [3]. In Ref. [4] the $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ ware suggested to be $1P \ \Xi'_c$ state with spin-parity $J^P = 3/2^-$ or $J^P = 5/2^$ in the chiral quark model. In Ref. [5] the two-body strong decays of the $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ were calculated by employing the ${}^{3}P_0$ approach with the conclusion that the $\Xi_c(2923)^0$ can be regarded as the $2S \ \Xi'_c$ state. The lattice QCD calculation was also performed and try to determine their quantum numbers [6].

Although the authors in Refs. [3,4,5,6] try to assign these states into the conventional three-quark frames, it is obvious that the inner structure of $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ are not finally determined. And another interpretation is treating them as $D\Lambda - D\Sigma$ molecular states, because the smallest mass gaps between the newly observed Ξ_c baryons and the ground Ξ_c , about 450 MeV, is large enough to excite a light quark-antiquark pair to form a hadronic molecular. Indeed, it is shown in Refs. [7, 8,9] that the interaction between D meson and Λ or Σ

^a Email address: huangy2019@swjtu.edu.cn

baryon is strong enough to form a bound state with a mass about 2930 MeV.

The key point in this work is to explain whether the $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ can be considered as a molecular state. Here, we will consider the strong $\Xi_c^* \to \Lambda_c^+ K^-, \Sigma_c^+ K^-$, and $\Xi_c^+ \pi^-$ decays of the $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ with the possible quantum num-

 $\Sigma (2938)^{\circ}$, and $\Sigma (2904)^{\circ}$ with the possible quantum numbers $J^{p} = 1/2^{\pm}$ and $3/2^{\pm}$ using an effective Lagrangian approach. The approach is based on the hypothesis that the Ξ_{c}^{*} is a hadronic molecule state of DA- $D\Sigma$. The coupling of the Ξ_{c}^{*} to the constituents is described by the effective Lagrangian. The corresponding coupling constant $g_{\Xi_{c}^{*}AD}$ and $g_{\Xi_{c}^{*}\Sigma D}$ are determined by the compositeness condition Z = 0 [10,11,12,13,14], which implies that the renormalization constant of the hadron wave function is set equal to zero. By constructing a phenomenological Lagrangian including the couplings of the bound state to its constituents and the constituents with other particles we calculated one-loop diagrams describing different decays of the molecular states.

This work is organized as follows. The theoretical formalism is explained in Sec. 2. The predicted partial decay widths are presented in Sec. 3, followed by a short summary in the last section.

2 FORMALISM AND INGREDIENTS

In the molecular scenario, the details of the calculations for $\Xi_c^0 \to \Lambda_c^+ K^-$, $\Xi_c^0 \to \Sigma_c^+ K^-$, and $\Xi_c^0 \to \Xi_c^+ \pi^-$ are presented for Ξ_c state with two different total angular momentum J possibilities. The molecular structure of the Ξ_c^* baryon with quantum numbers $J^p = 1/2^{\pm}$ is described by the Lagrangian [14]

$$\mathcal{L}_{\Xi_c^*}(x) = g_{\Xi_c^* D\Lambda} \int d^4 y \Phi(y^2) D^0(x + \omega_\Lambda y) \Gamma \Lambda(x - \omega_{D^0} y)$$

$$\times \bar{\Xi}_c^*(x) + g_{\Xi_c^* D\Sigma} \int d^4 y \Phi(y^2) [\sqrt{\frac{2}{3}} D^0(x + \omega_{\Sigma^0} y)$$

$$\times \Gamma \Sigma^0(x - \omega_{D^0} y) - \sqrt{\frac{1}{3}} D^+(x + \omega_{\Sigma^-} y)$$

$$\times \Gamma \Sigma^-(x - \omega_{D^+} y)] \bar{\Xi}_c^*(x). \tag{1}$$

while for the choice $J^p = 3/2^{\pm}$ the Lagrangian contains a derivative

$$\mathcal{L}_{\Xi_{c}^{*}}(x) = g_{\Xi_{c}^{*}DA} \int d^{4}y \Phi(y^{2}) D^{0}(x + \omega_{A}y) \Gamma \partial_{\mu} \Lambda(x - \omega_{D^{0}}y)$$

$$\times \bar{\Xi}_{c}^{*\mu}(x) + g_{\Xi_{c}^{*}D\Sigma} \int d^{4}y \Phi(y^{2}) [\sqrt{\frac{2}{3}} D^{0}(x + \omega_{\Sigma^{0}}y)$$

$$\times \Gamma \partial_{\mu} \Sigma^{0}(x - \omega_{D^{0}}y) - \sqrt{\frac{1}{3}} D^{+}(x + \omega_{\Sigma^{-}}y)$$

$$\times \Gamma \partial_{\mu} \Sigma^{-}(x - \omega_{D^{+}}y)] \bar{\Xi}_{c}^{*\mu}(x). \qquad (2)$$

where Γ is the corresponding Dirac matrix related to the spin-parity of the Ξ_c^* . In particular, for $J^p = 1/2^+, 3/2^-$ we have $\Gamma = \gamma^5$ while for $J^p = 1/2^-, 3/2^+$ the Dirac

structure $\Gamma = 1$. In the Lagrangian, an effective correlation function $\Phi(y^2)$ is introduced to reflect the distribution of two constituents in the hadronic molecular Ξ_c^* state. The introduced correlation function also makes the Feynman diagrams finite in the ultraviolet region of Euclidean space, which indicates that the Fourier transformation of the correlation function should drop fast enough in the ultraviolet region. Here we choose the Fourier transformation of the correlation in the Gaussian form [10, 11, 12, 13, 14],

$$\Phi(p_E^2) \doteq \exp(-p_E^2/\alpha^2) \tag{3}$$

with α being the size parameter which characterize the distribution of components inside the molecule. Though the value of α could not be determined from first principles, it is usually chosen to be about 1 GeV in the literature [10, 11, 12, 13, 14]. In this work, we set $\Lambda = 1.0$ GeV.

With the help of the effective Lagrangian in Eq.(1) and Eq.(2), we can obtain the self energy of the Ξ_c^*

$$\begin{split} \Sigma_{\Xi_{c}^{*}}^{1/2}(k_{0}) &= \int \frac{d^{4}k_{1}}{(2\pi)^{4}} \{g_{\Xi_{c}^{*}AD}^{2} \Phi^{2}[(k_{1}-k_{0}\omega_{A})^{2}] \Gamma \frac{k_{1}+m_{A}}{k_{1}^{2}-m_{A}^{2}} \Gamma \\ &\times \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{0}}^{2}} + g_{\Xi_{c}^{*}\Sigma D}^{2} [\frac{2}{3} \Phi^{2}[(k_{1}-k_{0}\omega_{\Sigma^{0}})^{2}] \\ &\times \Gamma \frac{k_{1}+m_{\Sigma^{0}}}{k_{1}^{2}-m_{\Sigma^{0}}^{2}} \Gamma \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{0}}^{2}} + \frac{1}{3} \Phi^{2}[(k_{1}-k_{0}\omega_{\Sigma^{-}})^{2}] \\ &\times \Gamma \frac{k_{1}+m_{\Sigma^{-}}}{k_{1}^{2}-m_{\Sigma^{-}}^{2}} \Gamma \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{+}}^{2}}] \} \quad (4) \\ \Sigma_{\Xi_{c}^{*}}^{3/2}(k_{0}) &= \int \frac{d^{4}k_{1}}{(2\pi)^{4}} \{g_{\Xi_{c}^{*}AD}^{2} \Phi^{2}[(k_{1}-k_{0}\omega_{A})^{2}] \Gamma \frac{k_{1}+m_{A}}{k_{1}^{2}-m_{A}^{2}} \Gamma \\ &\times \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{0}}^{2}} + g_{\Xi_{c}^{*}\Sigma D}^{2} [\frac{2}{3} \Phi^{2}[(k_{1}-k_{0}\omega_{\Sigma^{0}})^{2}] \\ &\times \Gamma \frac{k_{1}+m_{\Sigma^{0}}}{k_{1}^{2}-m_{\Sigma^{0}}^{2}} \Gamma \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{0}}^{2}} + \frac{1}{3} \Phi^{2}[(k_{1}-k_{0}\omega_{\Sigma^{-}})^{2}] \\ &\times \Gamma \frac{k_{1}+m_{\Sigma^{0}}}{k_{1}^{2}-m_{\Sigma^{-}}^{2}} \Gamma \frac{1}{(k_{1}-k_{0})^{2}-m_{D^{+}}^{2}}] k_{1}^{\mu} k_{1}^{\nu} \quad (5) \end{split}$$

where $k_0^2 = m_{\Xi_c^*}^2$ with $k_0, m_{\Xi_c}^*$ denoting the four momenta and mass of the Ξ_c^* , respectively, k_1, m_D , and m_A, m_Σ are the four-momenta, the mass of the *D* meson, and the mass of the *A* baryon, the mass of the *\Sigma* baryon, respectively. The coupling constant $g_{\Xi_c^*A\bar{D}}$ and $g_{\Xi_c^*\Sigma\bar{D}}$ is determined by the compositeness condition [15,16]. It implies that the renormalization constant of the hadron wave function is set equal to zero with

$$Z_{\Xi_c^*} = x_{D\Lambda} + x_{D\Sigma} - \frac{d\Sigma_{\Xi_c^*}^{1/2,3/2-T}}{d\not_{k_0}}|_{\not_{k_0} = m_{\Xi_c^*}} = 0.$$
(6)

where the x_{AB} is the probability to find the Ξ_c^* in the hadronic state AB with the normalization $x_{DA} + x_{D\Sigma} =$ 1.0. And the $\Sigma_{\Xi_c^*}^{3/2-T}$ is the transverse part of the selfenergy operator $\Sigma_{\Xi_c^*}^{3/2}$, related to $\Sigma_{\Xi_c^*}^{3/2}$ via

$$\Sigma_{\Xi_b^*}^{3/2}(k_0) = (g_{\mu\nu} - \frac{k_0^{\mu}k_0^{\nu}}{k_0^2})\Sigma_{\Xi_c^*}^{3/2-T} + \cdots .$$
 (7)



Fig. 1. Self-energy of the Ξ_c^* states..

p

Fig. (2) shows the hadronic decay of the $\Lambda D - \Sigma D$ molecular state into the $\Lambda_c^+ K^-, \Xi_c^+ \pi^-$, and $\Sigma_c^+ K^-$ final states occuring by exchanging nucleon, D_s^* meson, and D^* meson. To compute the amplitudes of the diagrams shown in Fig.(2), we need the effective Lagrangian densities for the relevant interaction vertices. In Refs. [17,18], coupling of the vector meson to charm baryons are described from effective Lagrangians, which are obtained using the hidden gauge formalism and assuming SU(4) symmetry:

Fig. 2. Feynman diagrams for the $\Xi_c^* \to \Lambda_c^+ K^-, \Xi_c \pi$, and $\Sigma_c \bar{K}$ decay process.

$$\mathcal{L}_{VBB} = \frac{g_1}{4} \sum_{i,j,k,l=1}^{4} \bar{B}_{ijk} \gamma^{\mu} (V_{\mu,l}^K B^{ijl} + 2V_{\mu,l}^j B^{ilk}), \quad (8)$$

where the coupling constant g = 6.6 is from Ref. [17]. The symbol V_{μ} represents the vector fields of the 16-plet of the ρ , given by

$$V_{\mu} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\rho^{0} + \omega) & \rho^{+} & K^{*+} \bar{D}^{*0} \\ \rho^{-} & \frac{1}{\sqrt{2}}(-\rho^{0} + \omega) & K^{*0} & D^{*-} \\ K^{*-} & \bar{K}^{*0} & \phi & D^{*-}_{s} \\ D^{*0} & D^{*+} & D^{*+}_{s} J/\psi \end{pmatrix}_{\mu}, \quad (9)$$

$$\begin{split} B^{121} &= p, \qquad B^{122} = n, \qquad B^{132} = \frac{1}{\sqrt{2}} \varSigma^0 - \frac{1}{\sqrt{6}} \Lambda, \\ B^{213} &= \sqrt{\frac{2}{3}} \Lambda, \qquad B^{231} = \frac{1}{\sqrt{2}} \varSigma^0 + \frac{1}{\sqrt{6}} \Lambda, \qquad B^{232} = \varSigma^-, \\ B^{233} &= \varXi^-, B^{311} = \varSigma^+, \qquad B^{313} = \varXi^0, \qquad B^{141} = -\varSigma^{++}_c \\ B^{142} &= \frac{1}{\sqrt{2}} \varSigma^+_c + \frac{1}{\sqrt{6}} \Lambda_c, \qquad B^{143} = \frac{1}{\sqrt{2}} \varXi^\prime_c + -\frac{1}{\sqrt{6}} \varXi^+_c, \\ B^{241} &= \frac{1}{\sqrt{2}} \varSigma^+_c - \frac{1}{\sqrt{6}} \Lambda_c, \qquad B^{242} = \varSigma^0_c, \end{split}$$

and B is the tensor of baryons belonging to the 20-plet of

$$B^{243} = \frac{1}{\sqrt{2}} \Xi_c^{\prime 0} + \frac{1}{\sqrt{6}} \Xi_c^0, \qquad B^{341} = \frac{1}{\sqrt{2}} \Xi_c^{,+} + \frac{1}{\sqrt{6}} \Xi_c^{,+},$$

$$B^{124} = \sqrt{\frac{2}{3}} \Lambda_c, \qquad B^{234} = \sqrt{\frac{2}{3}} \Xi_c^0, \qquad B^{314} = \sqrt{\frac{2}{3}} \Xi_c^{,+},$$

$$B^{342} = \frac{1}{\sqrt{2}} \Xi_c^{\prime 0} - \frac{1}{\sqrt{6}} \Xi_c^0, \qquad B^{343} = \Omega_c^0,$$

$$B^{144} = \Xi_{cc}^{++}, \qquad B^{244} = -\Xi_{cc}^{+}, \qquad B^{344} = \Omega_{cc}, \qquad (10)$$

where the indices i, j, k of B^{ijk} denote the quark content of the baryon fields with the identification $1 \leftrightarrow u, 2 \leftrightarrow d, 3 \leftrightarrow s$, and $4 \leftrightarrow c$.

To evaluate the diagrams in Fig.(2), in addition to the Lagrangian in Eq.(1), Eq.(2), and Eq.(8), the following effective Lagrangians, responsible for vector mesons and pseudoscalar mesons interactions are needed as well [17]

$$\mathcal{L}_{PPV} = \frac{i}{4} g_h \langle [\partial^\mu P, P] V_\mu \rangle, \qquad (11)$$

where P is the SU(4) pseudoscalar meson matrices, and $\langle ... \rangle$ in the trace over the SU(4) matrices. The meson matrices are [17]

$$P = \sqrt{2} \begin{pmatrix} \frac{\pi^{0}}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} + \frac{\eta'}{\sqrt{3}} & \pi^{+} & K^{+} & D^{0} \\ \pi^{-} & -\frac{\pi^{0}}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} + \frac{\eta'}{\sqrt{3}} & K^{0} & -D^{-} \\ K^{-} & \bar{K}^{0} & -\sqrt{\frac{2}{3}}\eta + \frac{\eta'}{\sqrt{3}} & D^{-}_{s} \\ D^{0} & -D^{+} & D^{+}_{s} & \eta_{c} \end{pmatrix}$$
(12)

The coupling g_h is fixed from the strong decay width of $K^* \to K\pi$. With the help of Eq. (12), the two-body decay width $\Gamma(K^{*+} \to K^0\pi^+)$ is related to g_h as

$$\Gamma(K^{*+} \to K^0 \pi^+) = \frac{g_h^2}{6\pi m_{K^{*+}}^2} \mathcal{P}_{\pi K^*}^3 = \frac{2}{3} \Gamma_{K^{*+}}, \quad (13)$$

(D + 2E)a

~

where $\mathcal{P}_{\pi K^*}$ is the three-momentum of the π in the rest frame of the K^* . Using the experimental strong decay width($\Gamma_{K^{*+}} = 50.3 \pm 0.8 \text{ MeV}$) and the masses of the particles needed in the present work are listed in Table. (1)we obtain g = 4.64 [1].

Table 1. Masses of the particles needed in the present work (in units of MeV).

Λ	Λ_c^+	Σ_c^+	Σ_c^0	Ξ_c^+	D^0
1115.683	2286.46	2452.90	2453.75	2467.93	1864.83
p	n	K^0	K^{\pm}	π^{\pm}	D^{\pm}
938.272	939.565	497.611	493.68	139.57	1869.65
Ξ_c^0	$D_s^{*\pm}$	$D^{*\pm}$	D^{*0}	Σ^+	Σ^0
2470.91	2112.1	2010.26	2006.85	1189.37	1192.642

The vertexes for the meson-baryon interaction are needed and the form in the SU(3) sector is given by the chiral Lagrangian [19]

$$\mathcal{L}_{\phi\mathcal{B}\mathcal{B}} = \frac{F}{2} \langle \bar{\mathcal{B}}\gamma_{\mu}\gamma_{5}[u^{\mu},\mathcal{B}] \rangle + \frac{D}{2} \langle \bar{\mathcal{B}}\gamma_{\mu}\gamma_{5}\{u^{\mu},\mathcal{B}\} \rangle, \quad (14)$$

where F = 0.51, D = 0.75 [19] and at lowest order in the pseudoscalar field $u^{\mu} = -\sqrt{2}\partial^{\mu}\phi/f$, with f = 93 MeV. And \mathcal{B} and ϕ is now the SU(3) matrix of the baryon octet and meson, respectively.

$$\mathcal{B} = \begin{pmatrix} \frac{1}{\sqrt{2}} \Sigma^0 + \frac{1}{\sqrt{6}} \Lambda & \Sigma^+ & p \\ \Sigma^- & -\frac{1}{\sqrt{2}} \Sigma^0 + \frac{1}{\sqrt{6}} \Lambda & n \\ \Xi^- & \Xi^0 & -\frac{2}{\sqrt{6}} \Lambda \end{pmatrix}.$$
 (15)

$$\phi = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}}\eta \end{pmatrix}.$$
 (16)

Moreover, the effective Lagrangians for the DNA_c and $DN\Sigma_c$ couplings are [20,21]

$$\mathcal{L}_{DNA_c} = ig_{A_c p D} \bar{A}_c \gamma_5 p D^0, \qquad (17)$$

$$\mathcal{L}_{DN\Sigma_c} = -ig_{\Sigma_c pD}\bar{N}\gamma_5 \boldsymbol{\tau} \cdot \boldsymbol{\Sigma}_c D, \qquad (18)$$

with au being the usual Pauli matrices. The coupling constant $g_{A_cpD} = 10.7^{+5.3}_{-4.3}$ and $g_{\Sigma_cpD} = -2.69$ are borrowed from Refs. [20,21,22], where we take the central values $g_{\Lambda_c pD} = 10.7$ in our calculation.

With the above vertices, the amplitudes of the triangle diagrams of Fig. (2), evaluated in the center of mass frame of final states, are

$$\mathcal{M}_{a} = (i)^{3} \frac{g_{1}g_{h}g_{\Xi_{c}^{*}\Lambda D^{0}}}{2} \int \frac{d^{4}q}{(2\pi)^{4}} \Phi[(k_{1}\omega_{\Lambda} - k_{2}\omega_{D})^{2}]$$

$$\times \bar{u}(p_{2})\gamma_{\mu} \frac{(\not{k}_{2} + m_{\Lambda})}{k_{2}^{2} - m_{\Lambda}^{2}} \Gamma\{u(k_{0}), ik_{2\rho}u^{\rho}(k_{0})\}$$

$$\times \frac{1}{k_{1}^{2} - m_{D^{0}}^{2}} (p_{1}^{\nu} + k_{1}^{\nu}) \frac{(-g^{\mu\nu} + q^{\mu}q^{\nu}/m_{D_{s}^{*}}^{2})}{q^{2} - m_{D_{s}^{*}}^{2}},$$

$$\begin{split} \mathcal{M}_{b} &= -(i)^{3} \frac{(D+3F)g_{A,p}Dg_{\Xi_{c}^{*}AD^{0}}}{2\sqrt{3}f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ &\times \varPhi[(k_{1}\omega_{A}-k_{2}\omega_{D})^{2}]\bar{u}(p_{2})\frac{(\not{q}-m_{N})}{q^{2}-m_{N}^{2}} \not{p}_{1} \\ &\times \frac{\not{k}_{2}+m_{A}}{k_{2}^{2}-m_{A}^{2}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{0}}^{2}}, \\ \mathcal{M}_{c} &= -(i)^{3} \frac{g_{1}g_{h}g_{\Xi_{c}^{*}AD^{0}}}{k_{2}^{2}-m_{A}^{2}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\} \\ &\times \frac{1}{k_{1}^{2}-m_{D^{0}}^{2}} (p_{1}^{\prime\prime}+k_{1}^{\prime\prime})\frac{(-g^{\mu\nu}+q^{\mu}q^{\nu}/m_{D^{*+}}^{2})}{q^{2}-m_{D^{*+}}^{2}}, \\ \mathcal{M}_{d} &= -(i)^{3} \frac{(D+3F)g_{\Sigma_{c}^{*}Dg_{\Xi_{c}^{*}AD^{0}}}{2\sqrt{3}f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ &\times \frac{1}{k_{1}^{2}-m_{D^{0}}^{2}} (p_{1}^{\prime\prime}+k_{1}^{\prime\prime})\frac{(-g^{\mu\nu}+q^{\mu}q^{\nu}/m_{D^{*+}}^{2})}{q^{2}-m_{D^{*+}}^{2}}, \\ \mathcal{M}_{d} &= -(i)^{3} \frac{(D+3F)g_{\Sigma_{c}^{*}Dg_{\Xi_{c}^{*}AD^{0}}}{2\sqrt{3}f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ &\times \oint[(k_{1}\omega_{A}-k_{2}\omega_{D})^{2}]\bar{u}(p_{2})\frac{(\not{q}-m_{D})}{q^{2}-m_{N}^{2}}} \not{p}_{1} \\ &\times \frac{\not{k}_{2}+m_{A}}{k_{2}^{2}-m_{A}^{2}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{0}}^{2}}, \\ \mathcal{M}_{e1} &= (i)^{3} \frac{(F-D)g_{A,c}pDg_{\Xi_{c}^{*}\Sigma^{0}D^{0}}}{2\sqrt{3}f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ &\times \varPhi[(k_{1}\omega_{\Sigma^{-}}-k_{2}\omega_{D^{0}})^{2}]\bar{u}(p_{2})\frac{(\not{q}-m_{D})}{q^{2}-m_{D}^{2}}} \not{p}_{1} \\ &\times \frac{\not{k}_{2}+m_{\Sigma^{0}}}{k_{2}^{2}-m_{\Sigma^{0}}^{2}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{0}}^{2}}, \\ \mathcal{M}_{e2} &= -(i)^{3} \frac{(F-D)g_{A,c}pDg_{\Xi_{c}^{*}\Sigma^{-}D^{+}}}{\sqrt{6}f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ &\times \varPhi[(k_{1}\omega_{\Sigma^{-}}-k_{2}\omega_{D^{+}})^{2}]\bar{u}(p_{2})\frac{(\not{q}-m_{D})}{q^{2}-m_{D}^{2}}} \not{p}_{1} \\ &\times \frac{\not{k}_{2}+m_{\Sigma^{-}}}{k_{2}^{2}-m_{\Sigma^{-}}^{2}}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{+}}^{2}}, \\ \mathcal{M}_{f1} &= (i)^{3} \frac{g_{1}gg_{\Xi_{c}^{*}\Sigma^{-}D^{+}}}{q^{2}-m_{\Sigma^{-}}^{2}}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{+}}^{2}} \\ &\times [p_{1}^{*}+k_{1}^{*})\frac{(-g^{\mu\nu}+q^{\mu}q^{\nu}/m_{D^{++})}}{q^{2}-m_{D^{+}}^{2}}}, \\ \mathcal{M}_{f2} &= -(i)^{3} \frac{g_{1}gg_{\Xi_{c}^{*}\Sigma^{-}D^{+}}}{q^{2}-m_{\Sigma^{-}}^{2}}} \Gamma\{u(k_{0}),ik_{2\rho}u^{\rho}(k_{0})\}\frac{1}{k_{1}^{2}-m_{D^{+}}^{2}} \\ &\times [p_{1}^{*}+k_{1}^{*})\frac{(-g^{\mu\nu}+q^{\mu}q^{\nu}/m_{D$$

$$\mathcal{M}_{g2} = (i)^{3} \frac{\sqrt{6}(F-D)g_{\Sigma_{c}pD}g_{\Xi_{c}^{*}\Sigma^{-}D^{+}}}{3f} \int \frac{d^{4}q}{(2\pi)^{4}} \\ \times \varPhi[(k_{1}\omega_{\Sigma^{-}} - k_{2}\omega_{D^{+}})^{2}]\bar{u}(p_{2})\frac{(\not{q} - m_{N})}{q^{2} - m_{N}^{2}} \\ \times \not{p}_{1}\frac{\not{k}_{2} + m_{\Sigma^{-}}}{k_{2}^{2} - m_{\Sigma^{-}}^{2}} \Gamma\{u(k_{0}), ik_{2\rho}u^{\rho}(k_{0})\} \\ \times \frac{1}{k_{1}^{2} - m_{D^{+}}^{2}}.$$
(19)

Once the amplitudes are determined, the corresponding partial decay widths can be obtained, which read,

$$\Gamma(\Xi_c^* \to MB) = \frac{1}{2J+1} \frac{1}{8\pi} \frac{|\boldsymbol{p}_1|}{m_{\Xi_c^*}^2} \overline{|\mathcal{M}|^2}, \qquad (20)$$

where J is the total angular momentum of the Ξ_b^* state, the $|\mathbf{p}_1|$ is the three-momenta of the decay products in the center of mass frame, the overline indicates the sum over the polarization vectors of the final hadrons, and MBdenotes the decay channel of MB, i.e., $\Lambda_c \bar{K}$, $\Xi_c \pi$, $\Sigma_c \bar{K}$.

3 RESULTS AND DISCUSSIONS



Fig. 3. The coupling constants of the Ξ_c^* state with different J^P assignments as a function of the parameter x_{DA} . And the x_{DA} is the probability to find the Ξ_c^* in the hadronic state DA.

In this work, the main ideal is to explain whether the $\Xi_c(2923)^0$, $\Xi(2938)^0$, and $\Xi(2964)^0$ can be considered as a $DA - D\Sigma$ molecular state by computing allowed twobody strong decay modes. To compute the partial decay widths of the Ξ_c^* , we first need the information of the coupling constants related to the molecular state and its components. In Fig. 3 we show the results for the coupling constant $g_{\Xi_c^*AD}$ and $g_{\Xi_c^*\Sigma D}$ of the various Ξ_c^* states for different spin-parity assignments and for a variation of the size parameter x_{DA} in a range of 0.0 - 1.0. In the discussed x_{DA} range, the coupling constant $g_{\Xi_c^*\Sigma D}$ decreases, while the coupling constant $g_{\Xi_c^*AD}$ increases. The results in Fig. 3 also show that the coupling constants increase of the x_{DA} for the $J^p = 3/2^-$ case, where the Ξ_c^* is a D-wave $DA - D\Sigma$ molecular state. For the case of $J^p = 1/2^-$, the coupling constants increases (or decreases) but relatively slowly compared with that coupling constants in the case of $J^p = 3/2^-$.



Fig. 4. The total decay width with different spin-parity assignments for the various Ξ_c^* as a function of the parameter x_{DA} . The cyan bands denote the experimental total width. [2].

We show the dependence of the total decay width on the x_{DA} in Fig. 4. The total decay widths increase with x_{DA} for the $J^P = 1/2^+$ and $J^P = 3/2^{\pm}$ assignments. For the $J^P = 1/2^-$ assignments, we found that the line shape of the the total decay widths are huge different and the total decay widths first increases, then decreases but very slowly. A possible explanation for this may be that for an S-wave loosely bound state the effective coupling strength of the bound state to its components is more sensitive to the inner structure than the effective coupling strength of another possible molecular state, such as Pwave molecular state, relative to their inner component. This is why people often focus on the bound state from Swave interaction and assume the P- and D-wave bound state should be difficult to form from hadron-hadron interaction [23].

From the Fig. 4, one find that the predicted total decay widths for the $\Xi_c(2964)$ state and $\Xi_c(2938)$ state in the four spin-parity assignments are all smaller than the experimental total width. Such results disfavor the assignment of these two states as $DA - D\Sigma$ molecular state. For the $\Xi_c(2923)$ state, the predicted total decay width is much smaller than the experimental total width in the case of $J^P = 1/2^+$, which disfavors such a spin-parity assignment for the $\Xi_c(2923)$ in the $D\Lambda - D\Sigma$ molecular picture. For the case of $J^P = 3/2^+$, Since the estimated total decay widths is much smaller than the experimental total width, this case can be completely excluded as well. The $J^P = 3/2^-$ case is also disfavored due to the smallest width predicted. Hence, only the $\Xi_c(2923)$ can be considered as the molecular states composed of $D\Lambda$ and $D\Sigma$ components by comparison with the total decay width experimentally measured. The results in Fig. 4 also show that the total decay width of the $\Xi_c(2923)$ can not be reproduced when only consider $\Xi_c(2923)$ as pure $D\Lambda$ or pure $D\Sigma$ molecular state.



Fig. 5. Partial decay widths of $\Xi_c(2923)^0 \to \Lambda_c^+ K^-$ (red solid line) and $\Xi_c(2923)^0 \to \Xi_c \pi$ (black dash line) with $J^P = 1/2^-$ as a function of the parameter x_{DA} . The cyan bands denote the experimental total width. [2].

The partial decay widths of $\Xi_c(2923)^0 \to \Lambda_c^+ K^-$ and $\Xi_c(2923)^0 \to \Xi_c \pi$ with $J^P = 1/2^-$ assignment as a function of the parameter x_{DA} are presented in Fig. 5. It is found that the transition $\Xi_c(2923)^0 \to \Xi_c \pi$ is main decay channel, which almost saturates the total width of $\Xi_c(2923)$. However, the transition $\Xi_c(2923)^0 \to \Lambda_c^+ K^$ give minor contributions. The decay width $\Xi_c(2923)^0 \to \Xi_c \pi$ is very different from that in the constituent quark model [3,4,5] if we assign the S-wave $DA - D\Sigma$ bound state as $\Xi_c(2923)$. Future experimental measurements of such a process can be quite useful to test the different interpretations of the $\Xi_c(2923)$.

4 Summary

In this work, the S-wave $D\Lambda - D\Sigma$ molecular states were studied by calculating their allowed two body strong decay to investigate whether the three new narrow Ξ_c^* baryons, $\Xi^*(2923)$, $\Xi_c^*(2938)$, and $\Xi_c^*(2964)$ can be understood as $D\Lambda - D\Sigma$ molecules. With the coupling constants obtained by the composition condition, the decays through hadronic loops are calculated in a phenomenological effective Lagrangian approach. The total decay widths can be well reproduced with the assumption that the $\Xi^*(2923)$ as S-wave $DA - D\Sigma$ bound state with $J^P = 1/2^-$, which decay channels are $\Lambda_c^+ K^-$ and $\Xi_c \pi$. The other newly reported Ξ_c^* states cannot be accommodated in the current molecular picture. If the $\Xi^*(2923)$ is pure $DA - D\Sigma$ molecule, the transition strength of $\Xi_c(2923)^0 \to \Xi_c \pi$ is quite different from that in the constituent quark model [3, 4,5] and the decay width almost saturates the total width of $\Xi_c(2923)$. Future experimental measurements of such a process can be quite useful to test the different interpretations of the $\Xi_c(2923)$.

Acknowledgments

This work is partly supported by the Development and Exchange Platform for Theoretic Physics of Southwest Jiaotong University in 2020(Grants No.11947404). We acknowledge the supported by the National Science Foundation of Chongqing (Grant No. cstc2019jcyj-msxm0953), the Science and Technology Research Program of Chongqing Municipal Education Commission (Grant No. KJQN201800510). NaNa Ma also acknowledge the supported by the National Natural Science Foundation of China(No. 11947229) and Postdoctoral Science Foundation of China (No. 2019M663853)

References

- M. Tanabashi *et al.* [Particle Data Group], Review of Particle Physics, Phys. Rev. D 98, 030001 (2018).
- R. Aaij *et al.* [LHCb], Observation of new Z⁰_c baryons decaying to A⁺_cK⁻, [arXiv:2003.13649 [hep-ex]].
- H. Yang, H. Chen and Q. Mao, Identifying the \(\mathcal{E}_c^0\) baryons observed by LHCb as P-wave \(\mathcal{E}_c'\) baryons, [arXiv:2004.00531 [hep-ph]].
- 4. K. Wang, L. Xiao and X. Zhong, Understanding the newly observed Ξ_c^0 states through their decays, [arXiv:2004.03221 [hep-ph]].
- 5. Qi-Fang. Lü, Canonical interpretations of the newly observed $\Xi_c(2923)^0$, $\Xi_c(2939)^0$ and $\Xi_c(2965)^0$ resonances, [arXiv:2004.02374 [hep-ph]].
- H. Bahtiyar, K. U. Can, G. Erkol, P. Gubler, M. Oka and T. T. Takahashi, Charmed baryon spectrum from lattice QCD near the physical point, [arXiv:2004.08999 [hep-lat]].
- C. E. Jimenez-Tejero, A. Ramos and I. Vidana, Dynamically generated open charmed baryons beyond the zero range approximation, Phys. Rev. C 80, 055206 (2009).
- Q. X. Yu, R. Pavao, V. R. Debastiani and E. Oset, Description of the *\(\mathcal{E}_c\)* and *\(\mathcal{E}_b\)* states as molecular states, Eur. Phys. J. C **79**, 167 (2019).
- J. Nieves, R. Pavao and L. Tolos, Ξ_c and Ξ_b excited states within a SU(6)_{lsf}× HQSS model, Eur. Phys. J. C 80, 22 (2020).
- A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, Strong and radiative decays of the D^{*}_{s0}(2317) meson in the DK-molecule picture, Phys. Rev. D 76, 014005 (2007).
- 11. A. Faessler, T. Gutsche, V. E. Lyubovitskij and Y. L. Ma, D^*K molecular structure of the $D_{s1}(2460)$ meson, Phys. Rev. D **76**, 114008 (2007).

- 12. Y. Dong, A. Faessler, T. Gutsche and V. E. Lyubovitskij, Strong two-body decays of the $\Lambda_c(2940)^+$ in a hadronic molecule picture, Phys. Rev. D **81**, 014006 (2010).
- 13. Y. Dong, A. Faessler, T. Gutsche, S. Kumano and V. E. Lyubovitskij, Radiative decay of $\Lambda_c(2940)^+$ in a hadronic molecule picture, Phys. Rev. D 82, 034035 (2010).
- Y. Dong, A. Faessler, T. Gutsche and V. E. Lyubovitskij, Charmed baryon Sigmac(2800) as a ND hadronic molecule, Phys. Rev. D 81, 074011 (2010).
- 15. S. Weinberg, Elementary particle theory of composite particles, Phys. Rev. **130**, 776 (1963).
- A. Salam, Lagrangian theory of composite particles, Nuovo Cim. 25, 224 (1962).
- J. Hofmann and M. F. M. Lutz, Coupled-channel study of crypto-exotic baryons with charm, Nucl. Phys. A **763**, 90 (2005).
- 18. G. Montaña, A. Feijoo and à. Ramos, A meson-baryon molecular interpretation for some Ω_c excited states, Eur. Phys. J. A 54, 64 (2018).
- E. J. Garzon and E. Oset, Effects of pseudoscalar-baryon channels in the dynamically generated vector-baryon resonances, Eur. Phys. J. A 48, 5 (2012).
- 20. J. J. Xie, Y. B. Dong and X. Cao, Role of the Λ_c^+ (2940) in the $\pi^- p \to D^- D^0 p$ reaction close to threshold, Phys. Rev. D **92**, 034029 (2015).
- 21. Y. Huang, J. He, J. J. Xie and L. S. Geng, Production of the $\Lambda_c(2940)$ by kaon-induced reactions on a proton target, Phys. Rev. D **99**, 014045 (2019).
- A. Khodjamirian, C. Klein, T. Mannel and Y. M. Wang, How much charm can PANDA produce?, Eur. Phys. J. A 48, 31 (2012).
- 23. J. He, Understanding spin parity of $P_c(4450)$ and Y(4274) in a hadronic molecular state picture, Phys. Rev. D **95**, 074004 (2017).

