A REMARK ON THE LAPLACIAN FLOW AND THE MODIFIED LAPLACIAN CO-FLOW IN G_2 -GEOMETRY

LUCIO BEDULLI AND LUIGI VEZZONI

ABSTRACT. We observe that the DeTurck Laplacian flow of G_2 -structures introduced by Bryant and Xu as a gauge fixing of the Laplacian flow can be regarded as a flow of G_2 -structures (not necessarily closed) which fits in the general framework introduced by Hamilton in [4].

1. INTRODUCTION

In [1] Bryant introduced a geometric flow in G₂-geometry which evolves an initial closed G₂-structure φ_0 in the direction of its Laplacian.

Given a compact 7-dimensional manifold with a closed G₂-structure (M, φ_0) , a Laplacian flow is a solution to the evolution equation

(1)
$$\frac{\partial}{\partial t}\varphi_t = \Delta_{\varphi_t}\varphi_t, \quad d\varphi_t = 0, \quad \varphi_{|t=0} = \varphi_0.$$

The well-posedness of equation (1) is proved in [2] by applying the Nash-Moser theorem to the gauge fixing

(2)
$$\frac{\partial}{\partial t}\varphi_t = \Delta_{\varphi_t}\varphi_t + \mathcal{L}_{V(\varphi_t)}\varphi_t, \quad d\varphi_t = 0, \quad \varphi_{|t=0} = \varphi_0$$

where \mathcal{L} is the Lie derivative and $V: C^{\infty}(M, \Lambda^3_+) \to C^{\infty}(M, TM)$ is a first order differential operator which depends on the choice of a connection on M. Here Λ^3_+ denotes the open subbundle of Λ^3 of G₂-structures on M. A solution to (2) is usually called a *DeTurck Laplacian flow*.

A DeTurck Laplacian flow φ_t is also a solution to

(3)
$$\frac{\partial}{\partial t}\varphi_t = dd^*_{\varphi_t}\varphi_t + d\iota_{V(\varphi_t)}\varphi_t, \quad \varphi_{|t=0} = \varphi_0.$$

In the present note we observe that equation (3) fits in the general framework introduced by Hamilton in [4]. As a direct consequence we have the following theorem which in particular implies the well-posedness of (2)

Theorem 1.1. Let (M, φ_0) be a compact 7-dimensional manifold with a G₂-structure. Then equation (3) has a unique short-time solution.

In [5] Karigiannis, McKay and Tsui introduced the Laplacian co-flow as the solution to the evolution equation $(1 - 1)^{-1}$

(4)
$$\frac{\partial}{\partial t}(*_{\varphi_t}\varphi_t) = -\Delta_{\varphi_t} *_{\varphi_t}\varphi_t, \quad d *_{\varphi_t}\varphi_t = 0, \quad \varphi_{|t=0} = \varphi_0.$$

where in this case φ_0 is supposed to be co-closed with respect to the metric induced by itself. The well-posedness of this last equation is still an open problem and Grigorian introduced in [3] the following modification

(5)
$$\frac{\partial}{\partial t}(*_{\varphi_t}\varphi_t) = \Delta_{\varphi_t} *_{\varphi_t} \varphi_t + 2d((A - \operatorname{Tr}(T(\varphi_t))\varphi_t), \quad d *_{\varphi_t} \varphi_t = 0, \quad \varphi_{|t=0} = \varphi_0,$$

Date: May 8, 2020.

This work was supported by GNSAGA of INdAM.

where A is a constant and $T(\varphi_t)$ is the torsion of φ_t . In [3] it is proved the well-posedness of (5) following the same approach of Bryant in [1] by applying the Nash-Moser theorem to the gauge fixing

(6)
$$\frac{\partial}{\partial t}(*_{\varphi_t}\varphi_t) = \Delta_{\varphi_t}*_{\varphi_t}\varphi_t + 2d((A - \operatorname{Tr}(T(\varphi_t))\varphi_t) + \mathcal{L}_{V(\varphi_t)}\varphi_t, \quad d*_{\varphi_t}\varphi_t = 0, \quad \varphi_{|t=0} = \varphi_0,$$

Any solution to this last equation (7) satsfies

(7)
$$\frac{\partial}{\partial t}(*_{\varphi_t}\varphi_t) = dd_{\varphi_t}^* *_{\varphi_t}\varphi_t + 2d((A - \operatorname{Tr}(T(\varphi_t))\varphi_t) + d\iota_{V(\varphi_t)}\varphi_t, \quad \varphi_{|t=0} = \varphi_0$$

Analogously to theorem 1.1 we have

Theorem 1.2. Let (M, φ_0) be a compact 7-dimensional manifold with a G₂-structure. Then equation (7) has a unique short-time solution.

2. Proof of the results

Both theorems 1.1 and 1.2 can be proved by using the following set-up introduced by Hamilton in [4].

Let M be an oriented compact manifold, F a vector bundle over M, U an open subbundle of F and

$$E\colon C^{\infty}(M,U)\to C^{\infty}(M,F)$$

a second order differential operator. For $f \in C^{\infty}(M, U)$, we denote by $DE(f): C^{\infty}(M, F) \to C^{\infty}(M, F)$ the linearization of E at f and by $\sigma DE(f)$ the principal symbol of DE(f).

Definition 2.1. An integrability condition for *E* is a first order linear differential operator

$$L\colon C^{\infty}(M,F)\to C^{\infty}(M,G)$$

where G is another vector bundle over M, such that L(E(f)) = 0 for all $f \in C^{\infty}(M, U)$, and all the eigenvalues of $\sigma DE(f)$ restricted to ker σL have strictly positive real part.

Theorem 2.2 (Hamilton [4, Theorem 5.1]). Assume that E admits an integrability condition. Then for every $f_0 \in C^{\infty}(M, U)$ the geometric flow

(8)
$$\frac{\partial f}{\partial t} = E(f), \quad f(0) = f_0,$$

has a unique short-time solution.

Now we can focus on the set-up of theorem 1.1. Here we consider

$$F = \Lambda^3, \quad U = \Lambda^3_+, \quad G = \Lambda^4, \quad E(\varphi) = dd^*_{\varphi}\varphi + d\iota_{V(\varphi)}\varphi, \quad L = d \colon C^{\infty}(M, \Lambda^3) \to C^{\infty}(M, \Lambda^4).$$

From [2] it follows that for every $\varphi \in C^{\infty}(M, U)$ and every closed $\psi \in C^{\infty}(M, \Lambda^3)$ we have

$$DE(\varphi)(\psi) = -\Delta_{\varphi}\psi + \text{l.o.t.}$$

Hence all the assumptions of Hamilton's theorem 2.2 are satisfied and theorem 1.1 follows.

Notice that if the starting form φ_0 is closed, then the solution to (3) is closed for every t since

$$d\frac{\partial}{\partial t}\varphi = 0.$$

Therefore if φ_0 is closed, the unique solution φ_t to (3) solves also the DeTurck-Laplacian flow (2) and the short-time existence of the DeTurck-Laplacian flow (2) can be deduced from Theorem 1.1.

About the proof of theorem 1.2 we set

$$F = \Lambda^4, \quad U = \Lambda^4_+, \quad G = \Lambda^5, \quad E(*_{\varphi}\varphi) = dd^*_{\varphi}\varphi + d\iota_{V(\varphi)} + 2d((A - \operatorname{Tr}(T(\varphi))\varphi), L = d: C^{\infty}(M, \Lambda^4) \to C^{\infty}(M, \Lambda^5).$$

From [3] it follows

$$DE(*_{\varphi}\varphi)(\psi) = -\Delta_{\varphi}\psi + \text{l.o.t.}$$

for every closed $\psi \in C^{\infty}(M, \Lambda^4)$ and the proof of theorem 1.2 follows.

Acknowledgments. We would like to thank Jason Lotay for useful conversations.

References

- R. BRYANT, Some remarks on G₂-structures, Proceedings of Gokova Geometry/Topology Conference, Gokova 2006, 75–109.
- [2] R. BRYANT, F. XU, Laplacian Flow for Closed G_2 -Structures: Short Time Behavior. arXiv:1101.2004.
- [3] S. GRIGORIAN, Short-time behaviour of a modified Laplacian coflow of G_2 -structures. Adv. Math. 248 (2013), 378–415.
- [4] R. S. HAMILTON, Three-manifolds with positive Ricci curvature. J. Differential Geom. 17 (1982), no. 2, 255–306.
- [5] S. KARIGIANNIS, B. MCKAY, M.-P. TSUI, Soliton solutions for the Laplacian coflow of some G₂ structures with symmetry, *Differential Geom. Appl.* **30** (2012), 318–333.

DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA, UNIVERSITÀ DELL'AQUILA, VIA VETOIO, 67100 L'AQUILA, ITALY

E-mail address: lucio.bedulli@univaq.it

Dipartimento di Matematica G. Peano, Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italy

E-mail address: luigi.vezzoni@unito.it