Constraining Alfvénic Turbulence with Helicity Invariants

Swadesh M. Mahajan,^{1*} and Manasvi Lingam²[†]

¹Institute for Fusion Studies, The University of Texas at Austin, Austin, TX 78712 ²Department of Aerospace, Physics and Space Sciences, Florida Institute of Technology, Melbourne FL 32901, USA

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

In this paper, we study the constraints imposed by the invariants (generalized helicities and energy) of extended magnetohydrodynamics on some global characteristics of turbulence. We show that the global turbulent kinetic and magnetic energies will approach equipartition only under certain circumstances that depend on the ratio of the generalized helicities. In systems with minimal thermal energy, we demonstrate that the three invariants collectively determine the characteristic length scale associated with Alfvénic turbulence.

Key words: (Sun:) solar wind - turbulence - plasmas - magnetic fields

1 INTRODUCTION

The existence of (generalized) magnetic helicity constraints introduces a fundamental distinction between the Navier-Stokes fluid turbulence and the low-frequency Alfvénic turbulence realized in magnetized plasma systems. The resultant Alfvénic turbulence has been widely investigated within the context of the simplest model namely, ideal magnetohydrodynamics (MHD) - but also for models that are often collectively known as "beyond MHD" or extended MHD (Goedbloed & Poedts 2004; Freidberg 2014).

The scientific literature is replete with examples wherein helicity invariants have been exploited to find new relaxed states. The most famous among them are the so-called Woltjer-Taylor states of ideal MHD ($\nabla \times \mathbf{B} = \mu \mathbf{B}$) that are obtained by minimizing the magnetic energy $\langle |\mathbf{B}|^2 \rangle$ while holding the magnetic helicity $h_m = \langle \mathbf{A} \cdot \mathbf{B} \rangle$ fixed (Woltjer 1958; Taylor 1974; Berger 1999; Matthaeus et al. 2012); henceforth, we shall make use of the notation $\langle \dots \rangle = \int d^3x$ for the sake of simplicity.

A very crucial role played by the constancy of h_m in the evolution of MHD turbulence was identified in early MHD simulations as well as in analytical models: it permitted the inverse cascading of magnetic helicity in 3D models (Frisch et al. 1975; Pouquet et al. 1976),¹ whereas in standard fluid turbulence, the transfer of energy and helicity is typically from larger to smaller scales (Moffatt 1978; Krause & Raedler 1980; Biskamp 2003; Zhou et al. 2004; Brandenburg & Subramanian 2005; Galtier 2018). The latter feature was inherent in the famous conjecture of Andrey Nikolaevich Kolmogorov (Kolmogorov 1941) that led to the equally famous scaling law $E_k \sim k^{-5/3}$ for the kinetic energy spectrum E_k (Frisch 1995;

Biskamp 2003; Davidson 2013). Another seminal result in the realm of MHD turbulence is the Iroshnikov-Kraichnan theory (Iroshnikov 1963; Kraichnan 1965), which modelled turbulent fluctuations as weakly interacting Alfvénic wave packets and yielded the magnetic energy spectrum $E_k \sim k^{-3/2}$ (Biskamp 2003).

This paper, although motivated by Kolmogorov's legacy, will dwell on precisely those features of Alfvénic turbulence that are absent in Navier-Stokes systems. The goal, in the spirit of Kolmogorov (1941), is to obtain results of maximal simplicity and, hopefully, of considerable generality that are potentially valid for all Alfvénic turbulence irrespective of its origination and evolution. More precisely, we will delve into constraints on Alfvénic turbulence imposed by the helicity and energy invariants of extended MHD, and thereby extend prior analyses along similar lines (Ohsaki et al. 2001, 2002; Mahajan et al. 2002); see also Helander (2017).

As we shall show henceforth, this line of enquiry yields several results of broad scope and interest: (1) Total turbulent energy in each channel - namely, magnetic (E_m) , kinetic (E_{kin}) and thermal (E_{th}) - is determined by a single attribute of turbulence, namely, a characteristic length scale $(L_T = K_T^{-1})$, (2) Complete expressions in terms of a single unknown parameter for all these energies in terms of the invariants and L_T , thus enabling us to predict, for instance, the relative energy distribution.

2 INVARIANTS OF EXTENDED MHD

For the sake of simplicity, we will concentrate on a two component (electron-ion) quasineutral plasma. Under the assumption of isotropic pressure with an adiabatic equation of state ($p \propto n^{\gamma}$), each component obeys the following equation of motion (Steinhauer & Ishida 1997; Mahajan 2003; Mahajan & Lingam 2015):

$$\frac{\partial}{\partial t} \mathbf{P}_{\beta} = \mathbf{v}_{\beta} \times \mathbf{\Omega}_{\beta} - \nabla \psi_{\beta}, \tag{1}$$

^{*} E-mail: mahajan@mail.utexas.edu

[†] E-mail: mlingam@fit.edu

¹ However, at scales smaller than the electron skin depth, the inverse cascade of helicity is transformed into a direct cascade as per theory and simulations (Miloshevich et al. 2017, 2018).

2 Mahajan & Lingam

where $\mathbf{P}_{\beta} = \mathbf{A} + (m_{\beta}c/q_{\beta})\mathbf{v}_{\beta}$ is proportional to the canonical momentum, $\mathbf{\Omega}_{\beta} = \nabla \times \mathbf{P}_{\beta} = \mathbf{B} + (m_{\beta}c/q_{\beta})\nabla \times \mathbf{v}_{\beta}$ represents the generalized vorticity for the species β with mass and charge of m_{β} and q_{β} , and $\psi_{\beta} = c/q_{\beta}(h_{\beta} + 1/2m_{\beta}v_{\beta}^{2} + q_{\beta}\phi)$ encompasses all of the gradient forces; note that h_{β} is the specific enthalpy, and ϕ is the electrostatic potential.

Taking the curl of Eq. (1) yields the canonical vortical dynamics (Mahajan & Yoshida 1998; Mahajan 2003; Andrés et al. 2014) that is epitomized by

$$\frac{\partial \mathbf{\Omega}_{\beta}}{\partial t} = \nabla \times \left(\mathbf{v}_{\beta} \times \mathbf{\Omega}_{\beta} \right),\tag{2}$$

The low frequency behavior of this system of ideal fluid equations, which is closed via the Ampère's law,

$$\nabla \times \mathbf{B} = (4\pi/c)\mathbf{J}, \qquad \mathbf{J} = \sum q_{\beta} n_{\beta} \mathbf{v}_{\beta}, \tag{3}$$

is the object of this investigation. Straightforward manipulation of (1)-(3) yields the following three constants of motion: the total energy

$$E = \left\langle \frac{B^2}{2} + \frac{1}{2} \sum_{\beta} n_{\beta} m_{\beta} v_{\beta}^2 + \frac{p}{(\gamma - 1)} \right\rangle,\tag{4}$$

where p is the total pressure, and two generalized helicities (GH),

$$H_{\beta} = \frac{1}{2} \langle \mathbf{P}_{\beta} \cdot \mathbf{\Omega}_{\beta} \rangle, \tag{5}$$

associated with each species. For a perfectly conducting system of n dynamical species, there exist a total of (n + 1) bilinear invariants (Mahajan & Lingam 2015). Although it is self-evident, it must nevertheless be emphasized that in any magneto-fluid system, unless the fluid inertia is neglected, it is the generalized helicity H_{β} , and *not* the magnetic helicity $H_m = \langle \mathbf{A} \cdot \mathbf{B} \rangle$ that is conserved; for instance, the conservation of H_m in MHD and in Hall MHD holds true because electron inertia is ignored (Turner 1986).

To study the constrained Alfvén dynamics (including turbulence), it is convenient to work in the equivalent one fluid variables, viz., the center-of-mass velocity \mathbf{V} , and the current \mathbf{J} defined below:

$$\mathbf{V} = \frac{m\mathbf{v}_{\mathbf{e}} + M\mathbf{v}_{\mathbf{i}}}{m+M} = \mu_{e}\mathbf{v}_{\mathbf{e}} + \mu_{i}\mathbf{v}_{\mathbf{i}}, \quad \mathbf{J} = ne(\mathbf{v}_{\mathbf{i}} - \mathbf{v}_{\mathbf{e}})$$
(6)

where the two species are identified as electrons (mass *m* and charge -e) and protons (mass *M* and charge *e*). However, we shall not rewrite (1)-(2) explicitly in terms of **V** and **J** because, in what follows, we will focus only on the invariants (4)-(5). The electron and ion helicity invariants translate into the new variables as

$$2H_e = \left\langle \hat{\mathbf{A}} \cdot \hat{\mathbf{B}} + (m/e)^2 \mathbf{V} \cdot \nabla \times \mathbf{V} - 2(m/e) \mathbf{V} \cdot \hat{\mathbf{B}} \right\rangle$$
(7)

$$2H_i = \left\langle \hat{\mathbf{A}} \cdot \hat{\mathbf{B}} + (M/e)^2 \mathbf{V} \cdot \nabla \times \mathbf{V} + 2(M/e) \mathbf{V} \cdot \hat{\mathbf{B}} \right\rangle$$
(8)

where $\hat{\mathbf{A}} = \mathbf{A} + \lambda_e^2 \nabla \times \mathbf{B}$ is the vector potential modified by the contribution stemming from a finite electron skin depth ($\lambda_e^2 = c^2/\omega_{pe}^2$); in other words, the second term in $\hat{\mathbf{A}}$ is obtained after using the Ampère's law given by (3); it is also derivable by means of the Hamiltonian or Lagrangian formulations from the parent two-fluid model (Keramidas Charidakos et al. 2014; Abdelhamid et al. 2015; Lingam et al. 2015; D'Avignon et al. 2016).

Evidently, both helicities comprise of their purely magnetic $(\hat{A} \cdot \hat{B})$, purely kinematic $(V \cdot \nabla \times V)$, and the mixed (i.e., cross) $(V \cdot \hat{B})$ components - the chief difference is that the contribution of the kinematic and mixed parts can be far more dominant for the

protons (due to $M \gg m$). For further analysis, it is much more transparent to construct the invariant combinations:

$$H_{+} = 2\mu_{i}H_{i} + 2\mu_{e}H_{e} = \left\langle \hat{\mathbf{A}} \cdot \hat{\mathbf{B}} \right\rangle + \frac{m}{M}\lambda_{i}^{2} \left\langle \mathbf{V} \cdot \nabla \times \mathbf{V} \right\rangle$$
(9)

$$H_{-} = 2H_{i} - 2H_{e} = \lambda_{i}^{2} \left\langle \mathbf{V} \cdot \nabla \times \mathbf{V} \right\rangle + 2\lambda_{i} \left\langle \mathbf{V} \cdot \hat{\mathbf{B}} \right\rangle$$
(10)

where the magnetic field has been normalized to some ambient field strength B_0 and and the velocity field is measured in terms of the corresponding Alfvén speed V_A (where $V_A^2 = B_0^2/(4\pi nM)$), i.e., we have used Alfvénic units (Mahajan & Lingam 2015). Notice that, aside from the normalized fields, the only basic parameter is the ion skin depth λ_i (where $\lambda_i^2 = c^2/\omega_{pi}^2$) that defines the intrinsic length scale of the system. Of course, the existence of the term proportional to (m/M) serves as a reminder that the electron inertia is not (yet) neglected and the electron length scale ($\lambda_e = \sqrt{(m/M)\lambda_i}$) appears in H_+ and in the variables $\hat{\mathbf{A}}$ and $\hat{\mathbf{B}}$. With the above choice of normalization, the helicities acquire the dimensions of length. In what follows, we shall utilize the dimensionless helicities defined to be $h_{\pm} = H_{\pm}/\lambda_i$.

These invariants h_+ and h_- act in concordance to constrain the total magnetic and kinetic energies of the system. We suppose that the system is embedded in an ambient magnetic field such that $\hat{\mathbf{B}} = \hat{\mathbf{B}}_0 + \hat{\mathbf{b}}$, and a similar expression can be constructed for the vector potential $\hat{\mathbf{A}}$.² For the time being, we analyze the case where there exists no ambient flow, implying that \mathbf{v} fully represents the velocity field (i.e., we have $\mathbf{V}=\mathbf{v}$). Note that $\langle \hat{\mathbf{A}} \cdot \hat{\mathbf{B}} \rangle$ will acquire a contribution of the form $\langle \hat{\mathbf{A}}_0 \cdot \hat{\mathbf{B}}_0 \rangle = H_0$, while the linear terms will vanish on integration (Krause & Raedler 1980). Written fully in terms of the (normalized) fluctuating fields denoted by lowercase boldface letters, our normalized invariant equations become

$$h = h_{+} - h_{0} = \frac{\left\langle \hat{\mathbf{a}} \cdot \hat{\mathbf{b}} \right\rangle}{\lambda_{i}} + \frac{m}{M} \lambda_{i} \left\langle \mathbf{v} \cdot \nabla \times \mathbf{v} \right\rangle, \tag{11}$$

$$h_{-} = \lambda_{i} \left\langle \mathbf{v} \cdot \nabla \times \mathbf{v} \right\rangle + 2 \left\langle \mathbf{v} \cdot \hat{\mathbf{b}} \right\rangle$$
(12)

3 HELICITY CONSTRAINTS ON TURBULENCE

From this point onward, our analysis will be purely algebraic and qualitative, as it relies essentially on heuristic considerations. In this paper, we will neglect the electron scale length ($\lambda_e = 0$) for the sake of simplicity, although electron inertia can be readily reintroduced; in other words, we investigate the Hall MHD regime (Goedbloed & Poedts 2004). In this scenario, the second term on the RHS of (11) becomes vanishingly small, and we end up with $\hat{\mathbf{a}} = \mathbf{a}$ and $\hat{\mathbf{b}} = \mathbf{b}$. We wish to figure out the constraints imposed on Alfvénic turbulence by the invariance of h and h_- .

Now, we introduce a characteristic length scale L_T for the turbulent magnetic field **b**; the equivalent wave number is $K_T = 1/L_T$. More specifically, because **b** = $\nabla \times \mathbf{a}$ is valid, we will invoke a phenomenological scaling of the form **b** ~ $K_T \mathbf{a}$ or **a** ~ $K_T^{-1}\mathbf{b}$. In other words, one may interpret K_T as the measure of the gradient associated with **b**; a similar approach was introduced for the turbulent velocity in Pouquet et al. (1976, pg. 348). Note, however, that this mathematical expression is valid *sensu stricto* if **b** is

 $^{^2}$ We implicitly presume that the functions \hat{B} and \hat{A} are well-behaved and that the term $\left<\hat{A}_0\cdot\hat{B}_0\right>$ is finite.



Figure 1. The equipartition length scale (k_c) as a function of the helicity ratio h_-/h . There are two different solutions for k_c , namely, k_+ and k_- depending on whether s_+ or s_- is adopted. We have plotted k_+ and k_- for two choices of α , viz. $\alpha = 1$ and $\alpha = 0.1$.

specified to be a Beltrami field, with K_T serving as the corresponding Beltrami parameter; this ansatz is not unreasonable because a number of publications model the turbulent fields as Arn'old-Beltrami-Childress fields (Childress & Gilbert 1995; Brandenburg & Subramanian 2005). By utilizing the relationship $\mathbf{a} = K_T^{-1}\mathbf{b}$ introduced above, (11) reduces to

$$h \approx \frac{\langle \mathbf{b} \cdot \mathbf{b} \rangle}{k_T} \tag{13}$$

where $k_T = \lambda_i K_T$ is the inverse of turbulent scale length measured in units of the ion skin depth. Rewriting (13) yields an estimate for the magnetic energy

$$E_m = \frac{\langle \mathbf{b} \cdot \mathbf{b} \rangle}{2} \approx \frac{hk_T}{2} \tag{14}$$

Following the same procedure we obtain

$$h_{-} \approx 2k_T E_{\rm kin} + 4\alpha \sqrt{E_{\rm kin}} \sqrt{E_m} \tag{15}$$

where $E_{\rm kin} = \langle \mathbf{v} \cdot \mathbf{v} \rangle / 2$. The second term on the RHS represents an alignment condition of sorts, because we suppose that the dimensionless factor α captures the "projection" of one turbulent field on the other. This approach is inspired by the fact that, in a special class of exact solutions of nonlinear Alfvén waves, **b** is linearly proportional to **v**; see, for instance, Walén (1944); Mahajan & Krishan (2005); Mahajan & Miura (2009); Abdelhamid et al. (2016). A more general and rigorous strategy for obtaining this term relies upon invoking the well-known Cauchy-Bunyakovsky-Schwarz inequality (Steele 2004), which yields

$$|\langle \mathbf{b} \cdot \mathbf{v} \rangle|^2 \le \langle \mathbf{b} \cdot \mathbf{b} \rangle \langle \mathbf{v} \cdot \mathbf{v} \rangle \tag{16}$$

and subsequently replacing the inequality in this expression with an equality involving the phenomenological dimensionless factor α that implicitly obeys $0 \le \alpha \le 1$ as follows:

$$|\langle \mathbf{b} \cdot \mathbf{v} \rangle|^2 = \alpha^2 \langle \mathbf{b} \cdot \mathbf{b} \rangle \langle \mathbf{v} \cdot \mathbf{v} \rangle \tag{17}$$

Lastly, we make use of the definitions of E_{kin} and E_m introduced earlier, and take the square root of the above equation to obtain the second term on the RHS of (15).

By utilizing (14), (15) is readily solved for

$$E_{\rm kin} \approx \frac{s_{\pm}^2}{k_T}, \quad s_{\pm} = \frac{-\alpha\sqrt{h} \pm \sqrt{\alpha^2 h + h_-}}{\sqrt{2}} \tag{18}$$

where *s* depends on the constants of motion and the parameter α ; it will be regulated by the detailed nature of turbulence. The estimates for the global turbulent magnetic and kinetic energies, i.e., (14) and (18), are rather robust for all Alfvénic turbulence accessible within the two-fluid equations and constitutes one of the salient results in the paper. It can be readily verified that the dominant behavior, contained in the scaling,

$$E_m \propto k_T, \qquad E_{\rm kin} \propto \frac{1}{k_T}$$
 (19)

holds true (with some corrections on the order of $k_T \lambda_e$) even when the electron dynamics is retained.

Independent of details, Alfvénic turbulence is strongly constrained by the ideal invariants of the system. For instance, these systems must obey a definitive, verifiable proportionality emerging from (19):

$$\frac{E_m}{E_{\rm kin}} \propto k_T^2 \tag{20}$$

Therefore, the short-scale turbulence ought to be much richer in magnetic energy while the portion of kinetic energy increases (in relative terms) as one moves toward longer scales (see Shaikh & Shukla 2009, Fig. 3). To put it differently, from (20) we see that the two energies may be displaced from equipartition. This lack of equipartition is commonly observed in studies of Hall MHD turbulence, dynamos, and "reverse" dynamos (Mininni et al. 2003; Krishan & Mahajan 2004; Mahajan et al. 2005; Mininni et al. 2007; Lingam & Mahajan 2015; Lingam & Bhattacharjee 2016a,b). In particular, the above behavior is consistent with numerical simulations of magnetic and kinetic energy spectra, as seen from Miura & Araki (2014, Fig. 2) and Stawarz & Pouquet (2015, Fig. 4); note, however, that the plots in these publications investigate energy spectra and not the global energy budgets. Furthermore, a number of MHD turbulence simulations (Wang et al. 2011; Oughton et al. 2015) as well as observations of the (turbulent) solar wind (Matthaeus & Goldstein 1982; Grappin et al. 1991; Salem et al. 2009; Chen et al. 2013) have revealed an "excess" of magnetic energy at small scales as well as differences in the slopes of magnetic and kinetic spectra (Boldyrev et al. 2011); the theoretical calculations by Abdelhamid et al. (2016) suggest that this feature is a generic characteristic of extended MHD. As the invariants are the defining "labels" for a given system, once they are specified, we can determine explicit estimates for both E_m and E_{kin} .

Notice that, although the estimate for E_m is rather simply related to h (which is essentially the magnetic helicity), E_{kin} has two solutions s_{\pm} . It is straightforward to verify that it is s_{\pm} that must correspond to conventional MHD turbulence - in the limit $h_{-} \ll h$,

$$s_{+} \simeq \frac{h_{-}}{2\sqrt{2}\alpha\sqrt{h}} \tag{21}$$

The complete expression for s_+ is the relevant expression for conventional Hall MHD. The larger root s_- in terms of magnitude will consequently yield a higher kinetic energy. Thus, it must be emphasized that, for a given set of helicities, there are two distinct turbulent energy states: $[E_m, E_{kin}(s_+)]$ and $[E_m, E_{kin}(s_-)]$. The ratio E_m/E_{kin} is physically relevant since it represents the ratio of the magnetic and kinetic energies. It is possible for this ratio to attain values both greater and smaller than unity. The critical turbulent



Figure 2. The turbulent length scale is shown for different solutions and values of h_-/h and E/h. The red, black and blue curves correspond to E/h = 100, E/h = 1 and E/h = 0.01, respectively. The unbroken and dotted curves correspond to selecting the positive branch of (26) with $k_c = k_+$ and $k_c = k_-$ respectively, whereas the dot-dashed and dashed curves constitute the positive branch of (26) with $k_c = k_+$ and $k_c = k_-$ respectively. In all cases depicted herein, we have adopted $\alpha = 1$ for simplicity.

length scale (k_c) at which equipartition is obtained is found by solving for $E_m/E_{kin} = 1$, thus leading us to

$$k_{c} = \frac{\sqrt{2}|s_{\pm}|}{|\sqrt{h}|} = |\alpha| \left(\sqrt{1 + \frac{h_{-}}{\alpha^{2}h}} \pm 1 \right).$$
(22)

As expected, there are two different critical length scales at which equipartition of kinetic and magnetic energies is achieved. An interesting point that emerges from the above formula is that k_c depends only on the ratio $\Gamma = h_-/h$ and not the individual helicities; aside from this ratio, it also depends on α . For the case with $\Gamma \gg 1$, we determine that both roots converge to $k_c \approx \sqrt{h_-/h}$. Note, however, that this solution is physically problematic because it corresponds to $L_T \ll \lambda_i$ - in this regime, Hall MHD is not accurate because electron inertia effects (neglected herein) come into play.

On the other hand, when we consider $\Gamma \ll 1$, we find that two divergent values for k_c follow - we obtain $k_{c1} \approx 2$ for one branch and $k_{c2} \approx h_-/(2|\alpha|h)$ in the other. For the first branch, we arrive at $L_{c1} \approx \lambda_i/2$, whereas the second branch yields $L_{c2} \gg \lambda_i$. Thus, for $\Gamma \ll 1$, there is a manifest bifurcation of the equipartition length scales: one of them is comparable to the ion skin depth, while the other is much larger than λ_i , and presumably comparable to the characteristic system length scale (Yoshida et al. 2004). The combination of $\Gamma \ll 1$ and $L_{c2} \gg \lambda_i$ essentially means that the system is dominated by the magnetic helicity (as opposed to the cross helicity and fluid helicity) and that equipartition is being achieved at macroscopic scales *sensu lato*. Hence, this regime is consistent with an ideal MHD-like picture, wherein large-scale behavior and magnetic helicity are dominant. The different values of k_c as a function of h_-/h and α are depicted in Fig. 1.

Until now, our analysis has concentrated only on the constraints on the magnetic and kinetic energy imposed by the helicity invariants. Let us now examine these results in conjunction with the conservation of energy, which in the language of preceding considerations, becomes (after having subtracted ambient field energy)



Figure 3. The minimum value of E/h that is sufficient to yield real values of k_T is plotted as a function of the helicity ratio h_-/h for different choices of α .

$$E = \left\langle \frac{\mathbf{b} \cdot \mathbf{b} + \mathbf{v} \cdot \mathbf{v}}{2} + \frac{p}{\gamma - 1} \right\rangle = E_m + E_{\text{kin}} + E_{\text{th}}, \tag{23}$$

where *E* is a constant that denotes the difference between the total energy and ambient magnetic energy. The obvious inference is that having already estimated E_m and E_{kin} in (14) and (18), we find that (23) allows us to calculate the turbulent thermal energy in terms of the three invariants (*h*, *h*₋ and *E*) of the system.

If we specialize to the special case where the turbulent kinetic energy is negligible for an incompressible plasma, we see that (23) reduces to

$$E = E_m + E_{\rm kin},\tag{24}$$

and thereby imposes an additional constraint on the system. In fact, we end up constraining the characteristic scale length of turbulence as follows:

$$k_T = \lambda_i K_T = \frac{\lambda_i}{L_T} = \frac{E}{h} \pm \sqrt{\frac{E^2}{h^2} - \frac{s_{\pm}^2}{2h}}.$$
 (25)

It is far more transparent to rewrite (25) in terms of k_c because we end up with

$$k_T = \frac{E}{h} \left[1 \pm \sqrt{1 - \left(\frac{k_c h}{2E}\right)^2} \right]$$
(26)

Although this expression looks deceptively simple, it is quite complex. It has a dependence on α , E/h and h_-/h via k_c . Moreover, there are four solutions in total: 2 arising from the \pm in the right-hand-side of (26) and 2 more from the fact that k_c has two different branches as seen from (22). After fixing α , we have plotted k_T in Fig. 2. Note that not all of the 4 solutions are guaranteed to be real, as seen from inspecting this figure.

In order for k_T to be real-valued, the following inequality must hold true:

$$\left|\frac{k_c h}{2E}\right| \le 1,\tag{27}$$

which imposes a constraint on the parameter space of $\{E, h, h_-\}$. Hence, depending on the parameters adopted, it is possible for k_T to have either 4 real roots, 2 real roots and 1 complex-conjugate pair, or 0 real roots and 2 complex-conjugate pairs (see Fig. 2 for an example). These results, as embodied by the equations (25) and (26), are qualitatively similar, albeit derived from a more generic standpoint, to the generalized magneto-Bernoulli mechanism elucidated in Ohsaki et al. (2001); Mahajan et al. (2002); Shatashvili et al. (2019); it has been proposed that this mechanism may constitute a viable explanation for solar flares (Kagan & Mahajan 2010), as opposed to classic paradigms such as fast magnetic reconnection (Biskamp 2000; Shibata & Magara 2011; Comisso et al. 2016).

Note that (27) provides us with another means of envisioning k_c . It is not only the length scale at which the ratio of magnetic and kinetic energies equals unity, but also the length scale that fulfills the criterion $L_{c1} \ge \lambda_i |h/E|/2$; here, recall that L_{c1} is constructed from the larger root of k_c , which we had dubbed k_{c1} . By utilizing (27), we have plotted the minimum value of E/h that suffices to ensure that k_T is real-valued; this lower bound depends on both the helicity ratio h_-/h and α .

Upon inspecting (25), we find that k_T is determined almost wholly in terms of the three constants of motion. We notice that there are two different solutions for k_T for a given choice of s_{\pm} , thereby giving rise to multiple length scales that can differ considerably in magnitude. In the above setting, it would seem then that the turbulence in each incompressible Alfvénic system, which is defined by its three invariants, should give rise to characteristic length scales that are fully determined or severely constrained by the invariants.

4 DISCUSSION

It makes intuitive sense that integral invariants (helicities and energy) would consequently set constraints on the global (i.e., integral) magnetic, kinetic and thermal energies; in fact, they may even formally determine them. Thus, we have obtained explicit relationships between E_m , $E_{\rm kin}$ and $E_{\rm th}$ on the one hand and L_T on the other, but the essentially heuristic arguments developed in this paper cannot give any specifications for the parameter α , which measures the degree of alignment of the two turbulent fields.

Likewise, it is natural to contend that our analysis would not directly yield the *k*-spectrum of E_m , E_{kin} and E_{th} . This information can seemingly emerge only via detailed studies of the Alfvénic dynamics, which has been a most active field of investigation in the physics of turbulent plasmas. Thus, to reiterate, our work does not examine the consequences for energy spectra, as it focuses on the *global* energy budgets. The details of the energy spectrum for the various regimes of extended MHD have been explored by Abdelhamid et al. (2016) in the context of the solar wind. It was shown therein that the spectrum in the MHD regime obeys a Kolmogorov scaling as opposed to the Iroshnikov-Kraichnan scaling, in agreement with prior theoretical and empirical results (Goldstein et al. 1995; Bruno & Carbone 2016; Sahraoui et al. 2020).

By simply harnessing the fundamental plasma invariants (helicities and energy), we were able to formulate certain interesting and possibly generic results for Alfvénic turbulence. For a specific set of invariants that remain invariant during whatever dynamics the system undergoes (thus serving as a "label"), all three components of the total turbulent energy are potentially dictated by a single feature of turbulence that embodies the length scale L_T associated with the small-scale magnetic field; conversely, one may interpret this length scale as being fully determined if the trio of invariants are specified, as seen from (26).

Expressing the turbulent energies in terms of L_T also enabled us to deduce some basic constraints on their magnitudes. Although our analysis was expressly concerned with global quantities, we found that our results are compatible with spectral relationships that have been identified in the Hall regime via numerical simulations; while this fact does not validate our predictions, it bolsters their credibility. Apart from establishing some fundamental intrinsic features of Alfvénic turbulence as described hitherto, our results can motivate as well as provide a check on detailed simulations.

Lastly, our analysis is valid over a broad range of turbulent scale lengths, namely, $L_{eq}^{-1} \ll k_T = \lambda_i K_T \ll M/m$, where L_{eq} is some equilibrium scale length that typically encapsulates the system size. It can be readily extended to and beyond the electron skin depth (λ_e) , but it seems relatively unlikely that the characteristic scale for Alfvénic turbulence would enter this regime.

ACKNOWLEDGEMENTS

We thank our reviewer, Mitchell Berger, for the positive and insightful report. This work was partially supported by the US-DOE grant DE-FG02-04ER54742.

REFERENCES

- Abdelhamid H. M., Kawazura Y., Yoshida Z., 2015, JPhA, 48, 235502
- Abdelhamid H. M., Lingam M., Mahajan S. M., 2016, ApJ, 829, 87
- Andrés N., Gonzalez C., Martin L., Dmitruk P., Gómez D., 2014, PhPl, 21, 122305
- Berger M. A., 1999, PPCF, 41, B167
- Biskamp D., 2000, Magnetic Reconnection in Plasmas. Cambridge University Press
- Biskamp D., 2003, Magnetohydrodynamic Turbulence. Cambridge University Press
- Boldyrev S., Perez J. C., Borovsky J. E., Podesta J. J., 2011, ApJ, 741, L19 Brandenburg A., Subramanian K., 2005, Phys. Rep., 417, 1
- Bruno R., Carbone V., 2016, Turbulence in the Solar Wind. Lecture Notes in Physics Vol. 928, Springer, doi:10.1007/978-3-319-43440-7
- Chen C. H. K., Bale S. D., Salem C. S., Maruca B. A., 2013, ApJ, 770, 125
- Childress S., Gilbert A. D., 1995, Stretch, Twist, Fold: The Fast Dynamo. Springer-Verlag
- Comisso L., Lingam M., Huang Y. M., Bhattacharjee A., 2016, PhPl, 23, 100702
- D'Avignon E. C., Morrison P. J., Lingam M., 2016, PhPl, 23, 062101
- Davidson P. A., 2013, Turbulence in rotating, stratified and electrically conducting fluids. Cambridge University Press
- Freidberg J. P., 2014, Ideal MHD. Cambridge University Press
- Frisch U., 1995, Turbulence. The legacy of A.N. Kolmogorov. Cambridge University Press
- Frisch U., Pouquet A., Leorat J., Mazure A., 1975, JFM, 68, 769
- Galtier S., 2018, JPhA, 51, 293001
- Goedbloed J. P. H., Poedts S., 2004, Principles of Magnetohydrodynamics. Cambridge University Press
- Goldstein M. L., Roberts D. A., Matthaeus W. H., 1995, ARA&A, 33, 283
- Grappin R., Velli M., Mangeney A., 1991, AnGeo, 9, 416
- Helander P., 2017, JPIPh, 83, 715830401
- Iroshnikov P. S., 1963, Azh, 40, 742
- Kagan D., Mahajan S. M., 2010, MNRAS, 406, 1140
- Keramidas Charidakos I., Lingam M., Morrison P. J., White R. L., Wurm A., 2014, PhPl, 21, 092118
- Kolmogorov A., 1941, Akademiia Nauk SSSR Doklady, 30, 301
- Kraichnan R. H., 1965, PhFl, 8, 1385

6 Mahajan & Lingam

- Krause F., Raedler K. H., 1980, Mean-field magnetohydrodynamics and dynamo theory. Pergamon Press
- Krishan V., Mahajan S. M., 2004, JGRA, 109, A11105
- Lingam M., Bhattacharjee A., 2016a, MNRAS, 460, 478
- Lingam M., Bhattacharjee A., 2016b, ApJ, 829, 51
- Lingam M., Mahajan S. M., 2015, MNRAS, 449, L36
- Lingam M., Morrison P. J., Miloshevich G., 2015, PhPl, 22, 072111
- Lingam M., Miloshevich G., Morrison P. J., 2016, PhLA, 380, 2400
- Mahajan S. M., 2003, Phys. Rev. Lett., 90, 035001
- Mahajan S. M., Krishan V., 2005, MNRAS, 359, L27
- Mahajan S. M., Lingam M., 2015, PhPl, 22, 092123
- Mahajan S., Miura H., 2009, JPIPh, 75, 145
- Mahajan S. M., Yoshida Z., 1998, Phys. Rev. Lett., 81, 4863
- Mahajan S. M., Nikol'skaya K. I., Shatashvili N. L., Yoshida Z., 2002, ApJ, 576, L161
- Mahajan S. M., Shatashvili N. L., Mikeladze S. V., Sigua K. I., 2005, ApJ, 634, 419
- Matthaeus W. H., Goldstein M. L., 1982, J. Geophys. Res., 87, 6011
- Matthaeus W. H., Montgomery D. C., Wan M., Servidio S., 2012, JTurb, 13, 37
- Miloshevich G., Lingam M., Morrison P. J., 2017, NJPh, 19, 015007
- Miloshevich G., Morrison P. J., Tassi E., 2018, PhPl, 25, 072303
- Mininni P. D., Gómez D. O., Mahajan S. M., 2003, ApJ, 587, 472
- Mininni P. D., Alexakis A., Pouquet A., 2007, JPIPh, 73, 377
- Miura H., Araki K., 2014, PhPl, 21, 072313
- Moffatt H. K., 1978, Magnetic field generation in electrically conducting fluids. Cambridge University Press
- Ohsaki S., Shatashvili N. L., Yoshida Z., Mahajan S. M., 2001, ApJ, 559, L61
- Ohsaki S., Shatashvili N. L., Yoshida Z., Mahajan S. M., 2002, ApJ, 570, 395
- Oughton S., Matthaeus W. H., Wan M., Osman K. T., 2015, RSPTA, 373, 20140152
- Pouquet A., Frisch U., Leorat J., 1976, JFM, 77, 321
- Sahraoui F., Hadid L., Huang S., 2020, RvMPP, 4, 4
- Salem C., Mangeney A., Bale S. D., Veltri P., 2009, ApJ, 702, 537
- Shaikh D., Shukla P. K., 2009, Phys. Rev. Lett., 102, 045004
- Shatashvili N. L., Mahajan S. M., Berezhiani V. I., 2019, Ap&SS, 364, 148
- Shibata K., Magara T., 2011, LRSP, 8, 6
- Stawarz J. E., Pouquet A., 2015, PhRvE, 92, 063102
- Steele J. M., 2004, The Cauchy-Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities. Cambridge University Press
- Steinhauer L. C., Ishida A., 1997, Phys. Rev. Lett., 79, 3423
- Taylor J. B., 1974, Phys. Rev. Lett., 33, 1139
- Turner L., 1986, ITPS, 14, 849
- Walén C., 1944, ArMAF, 30A, 1
- Wang Y., Boldyrev S., Perez J. C., 2011, ApJ, 740, L36
- Woltjer L., 1958, PNAS, 44, 489
- Yoshida Z., Mahajan S. M., Ohsaki S., 2004, PhPl, 11, 3660
- Zhou Y., Matthaeus W. H., Dmitruk P., 2004, RvMP, 76, 1015

This paper has been typeset from a TEX/LATEX file prepared by the author.