An Alexander type invariant for doodles

B. Cisneros, M. Flores, J. Juyumaya and C. Roque-Márquez

Abstract

We construct an Alexander type invariant for oriented doodles from a deformation of the Tits representation of the twin group and from the Chebyshev polynomials of second kind. Similar to the Alexander polynomial, our invariant vanishes on unlinked doodles with more than one component. We also include values of our invariant on several doodles.

2020 Mathematics Subject Classification. Primary: 57K14, 57K12, 20F36. *Keywords*: doodle; twin group; Coxeter group; Alexander polynomial; Chebyshev polynomials.

1 Introduction

Doodles were firstly introduced by Fenn and Taylor [9] as a finite collection of embedded circles on the 2-sphere with no triple intersections. Later, Khovanov [19] pointed out that it was more natural to consider doodles as immersed circles on the 2-sphere than as simple curves. In this manner, some concepts of knot theory were successfully transferred to doodle theory. In particular, he obtained an associated group to a doodle, called the doodle group, in analogue to the fundamental group of the complement of a link. In contrast with knot theory, equivalence of doodles just considers a planar version of the first and the second Reidemeister moves (see Figure 1); the lack of the third Reidemeister move leads to significant differences in comparison with classical knot theory. For instance, every class of a doodle has a unique representative with minimal number of crossings [19]; the calculation of Vassiliev invariants is more complicated than classical knots, but Vassiliev invariants classify doodles, problem that remains open for knots [23]. Extended to a more general setting, recent work by Bartholomew-Fenn-Kamada-Kamada [4] consider doodles on closed oriented surfaces of any genus, which can be considered as virtual links analogue for doodles; in [5] they give a complete invariant for virtual doodles¹.

¹In Bartholomew's webpage [3] is shown a table of virtual doodles up to 10 crossings.

The algebraic counterparts of doodles are the so-called *twin groups*, terminology due to Khovanov. The role of these groups in doodle theory is similar to that of Artin's braid groups in knot theory. These groups have appeared under different names and contexts in the literature: cartographical Grothendieck groups [26], quantum symmetric groups [10], flat braid groups² [21], traid groups [13] and planar braid groups [11, 24]. Further, the twin group on n strands, denoted by T_n , is a right angled Coxeter group generated by n-1 involutions and its elements can be depicted as planar braids on n strands. The permutation induced by a planar braid defines a natural epimorphism from T_n to the symmetric group on n symbols, whose kernel is called the *pure twin group*. So far, the twin group and pure twin group have been studied in the last years, though there still exist many open questions about these groups. For details on recent developments, refer to [2, 8, 11, 13, 15, 14, 24].

The first evidence of the close relationship between doodles and twins is by the straightforward process of obtaining a doodle from a given twin by 'closing' the twin. In the reverse process, Khovanov [19] proved the Alexander theorem for doodles in which any doodle on the 2-sphere is the closure of a twin, and recently Gotin [12] has established the Markov theorem for doodles on the 2-sphere in which he defines the corresponding Markov moves for twins. These two theorems invite to the construction of doodle invariants in a similar way it has been done in classical knot theory. For instance, the construction of an Alexander and Jones polynomials for doodles.

The purpose of this paper is the construction of an Alexander type invariant for doodles. We follow Burau's work to compute the Alexander polynomial for links via certain representation of the braid group [7], applied in the context of twins and doodles.

More precisely, we define a representation $\psi_n : T_n \to GL_{n-1}(\mathbb{Z}(x))$ which is a deformation of the Tits representation of T_n . Then, using ψ_n we construct a function $f_n : T_n \to \mathbb{Z}(x)$ which defines an invariant of doodles up to a factor x^{2k} , i.e., if two twins α, β have equivalent closures $\hat{\alpha} \sim \hat{\beta}$ as doodles, then their images $f_n(\alpha), f_m(\beta)$ differ by a multiple of x^{2k} for some $k \in \mathbb{Z}$ (see Lemma 4.9). Thus, we obtain an invariant $\mathcal{Q}(D)$ for a doodle D as the smallest degree polynomial over all images under functions f_n of twins with closure D (Theorem 4.13). Furthermore, we verify that these functions satisfy the skein relation

$$f_n\left(\left(\begin{array}{c} \\ \\ \\ \end{array}\right) - f_n\left(\left(\begin{array}{c} \\ \\ \\ \end{array}\right) \right) = (x^2 - 1)\left(f_n\left(\left(\begin{array}{c} \\ \\ \\ \end{array}\right) \right) - f_n\left(\left(\begin{array}{c} \\ \\ \\ \end{array}\right) \right)\right).$$

It is worth noting that Chebyshev polynomials of second kind appeared unexpectedly in the construction of the functions f_n . Observe that it is not the first time where Chebyshev polynomials of second kind emerge in low dimensional topology. For instance, they play an important role defining Jones-Wenzl projectors [27].

²This term was later used in [17] to an entirely different object

The paper is organized as follows. In Section 2, we give some preliminaries on doodles and twins; we recall the Tits representation of a Coxeter group and list necessary properties of Chebyshev polynomials. In Section 3, we study a deformation of the Tits representation of T_n as a Coxeter group and some of its features. In Section 4, we investigate the behavior of the functions f_n under Markov moves, obtaining the definition of our polynomial invariant Q. Finally, we give some examples of computations in Section 5.

Acknowledgments

The authors would like to thank Jacob Mostovoy and Jesús González for useful conversations. The first author was supported by Cátedras-CONACYT (project no. 61). The second author was partially supported by grant FONDECYT 11170305. The third author was supported, in part, by grant FONDECYT 1180036. The fourth author was partially supported by a CONACYT Postdoctoral Fellowship. This work is a result of two Mexican-Chilean meetings celebrated at Oaxaca and at Valparaiso founded by FONDECYT 11170305 and FONDECYT 1180036.

2 Preliminaries

In the present section, we give the necessary background on doodles, twins, Tits representation and Chebyshev polynomials that will be used along the paper. This section contains nothing essentially new.

2.1 Doodles and twin group

Definition 2.1 (cf. Khovanov, [19]). A *doodle* is an immersion $D : \bigsqcup_n S^1 \to S^2$ of a disjoint union of *n* circles into the 2-sphere³ with no triple or higher multiple intersections; *n* is the number of components. An *oriented doodle* is a doodle in which each component is oriented. We denote the set of oriented doodles by \mathcal{D} .

Throughout this work we will only be concerned with oriented doodles, hence we will omit the word oriented in most of the cases.

Two doodles D and D' are *equivalent*, denoted by $D \sim D'$, if there exists a homotopy between D and D' in which no triple intersections are produced throughout the homotopy. Equivalently⁴, $D \sim D'$ if they can be transformed into each other through isotopies of S^2

³In [4], doodles in S^2 are called *planar doodles*.

⁴On *regular* doodles, i.e., all its multiple points are transversal double points [4].

and a finite sequence of local moves R1 and R2, see Figure 1. These moves will be called the first and the second Reidemeister moves.



Figure 1: Reidemeister moves.

The doodle in Figure 2a is called the trivial doodle. The first non-trivial doodle is the Borromean doodle, shown in Figure 2b, with six crossings and three components. Following [4], the doodle in Figure 2c is called the 4-poppy doodle and it has eight crossings; it is the first non-trivial 1-component doodle [4, Theorem 4.2].



The algebraic counterparts of doodles are twin groups. More precisely, these groups are to doodle theory as braid groups are to knot theory.

Definition 2.2. The *twin group* T_n is the group presented by generators t_1, \ldots, t_{n-1} and the relations:

$$t_i^2 = 1$$
 for all $1 \le i \le n - 1$, (2.1)

$$t_i t_j = t_j t_i$$
 for all $1 \le i, j \le n - 1$ with $|i - j| > 1$. (2.2)

In [18], Khovanov gives a geometrical interpretation of T_n . To be precise, the elements of T_n can be regarded as planar braids, called twins. Namely, a *twin* on n strands is a collection of n descending arcs in $\mathbb{R} \times [0, 1]$ which no three arcs have a point in common. In particular, the generator t_i is represented by the following "elementary" twin:

Relations (2.1) and (2.2) are represented, respectively, as follows:



For $n \geq 1$, let $\iota_n^R : T_n \hookrightarrow T_{n+1}$ be the natural inclusion given by $\iota_n^R(t_i) = t_i$, and let $\iota_n^L : T_n \hookrightarrow T_{n+1}$ be the inclusion given by $\iota_n^L(t_i) = t_{i+1}$, for $1 \leq i \leq n-1$ in both cases. Thus, geometrically ι_n^R (resp. ι_n^L) adds a vertical strand to the right (resp. left) of the twin, see Figure 4. We denote $\iota^R = {\iota_n^R}_{n\geq 1}$ and $\iota^L = {\iota_n^L}_{n\geq 1}$ to the systems of inclusions, and we will simply write $\iota^R(\beta)$ for $\iota_n^R(\beta)$ (resp. $\iota^L(\beta)$ for $\iota_n^L(\beta)$) if $\beta \in T_n$ for some $n \geq 1$. Let T_∞ denote the inductive limit associated to ι^R .

$$\begin{array}{c|c} & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & &$$

Figure 4: Inclusions ι_n^R and ι_n^L .

2.2 Alexander and Markov theorems for doodles

The *closure* $\hat{\beta}$ of the twin β is the doodle obtained by connecting its upper and lower ends respectively as shown in Figure 5. Knovanov proved the analogue to the classical Alexander theorem for doodles.

Theorem 2.3 (Khovanov, [19, Theorem 2.1]). Any doodle on the 2-sphere is the closure of a twin.

Theorem 2.3 says the map $\beta \mapsto \hat{\beta}$ from T_{∞} onto \mathcal{D} is surjective. However, this map is highly not injective, for instance the twins $t_1 \in T_2$ and $t_1t_2 \in T_3$ have the same closure. Recently in [12], Gotin proves the analogue to the Markov theorem for doodles filling the gap in this correspondence. In order to state this theorem, we set

$$t_{n,i} := t_n t_{n-1} \cdots t_{n-i+1} t_{n-i} t_{n-i+1} \cdots t_{n-1} t_n, \qquad (2.4)$$

$$t_{1,i} := t_1 t_2 \cdots t_i t_{i+1} t_i \cdots t_2 t_1 \tag{2.5}$$

for $0 \le i \le n-1$.



Figure 5: The Borromean doodle as the closure $(t_1t_2)^3$.

Theorem 2.4 (Gotin, [12, Theorem 4.1]). Let γ and γ' be two twins with D and D' their respective closures. Then, D and D' are equivalent doodles if and only if γ can be transformed into γ' by a finite sequence of the following moves:

- $M0 : \iota^R(\beta) \sim \iota^L(\beta)$ for any $\beta \in T_n$,
- $M1 : \alpha\beta \sim \beta\alpha$ for any $\alpha, \beta \in T_n$,
- $M2 : \beta \sim \iota^{R}(\beta)t_{n,i}$ for any $\beta \in T_{n}$ and $0 \leq i \leq n-1$,
- M3 : $\beta \sim \iota^L(\beta) t_{1,i}$ for any $\beta \in T_n$ and $0 \leq i \leq n-1$.

We call these moves, the Gotin-Markov moves.

<u> </u>	I I	
β	$\stackrel{M0}{\sim}$	eta

Figure 6: M0 move.

Remark 2.5. When i = 0 in M2 we have $t_{n,0} = t_n$, then the M2 move becomes the classical second Markov move.

The Gotin-Markov moves define an equivalence relation on T_{∞} , denoted by \sim_G . Thus, Theorem 2.3 together with Theorem 2.4 implies the bijection

$$\mathcal{D}/\sim \quad \longleftrightarrow \quad T_\infty/\sim_G.$$



Figure 7: M2 and M3 moves.

This bijection says that constructing an invariant for doodles it reduces to find a family $f = \{f_n : T_n \to S\}_{n \ge 1}$, where S is a well understood set and the functions f_n are invariant under the Gotin-Markov moves. In Section 4, we will construct a such family by taking $S = \mathbb{Z}(x)$ and certain f_n defined in terms of a deformation of the Tits representation of the twin group.

2.3 Tits representation of Coxeter groups

We describe briefly some properties of Coxeter groups and the particular case for twin groups. Let S be a finite set; a *Coxeter matrix* on S is a symmetric matrix $M = (m_{s,t})_{s,t\in S}$ such that $m_{s,t} \in \{2, 3, ...\} \cup \{\infty\}$ and $m_{s,s} = 1$. The Coxeter group associated to M is the group defined by the following presentation:

$$W(M) := \langle S \mid (st)^{m_{s,t}} \text{ for all } s, t \text{ with } m_{s,t} \neq \infty \rangle.$$

Tits constructed a faithful linear representation for any Coxeter group in the following way. Let $S = \{s_1, s_2, \ldots, s_n\}$ be a finite set, M be a Coxeter matrix on S and V be a real vector space with basis $\{e_1, \ldots, e_n\}$. Now define the symmetric bilinear form B on V by

$$B(e_i, e_j) = \begin{cases} -\cos(\frac{\pi}{m_{i,j}}) & \text{if } m_{s_i, s_j} \neq \infty, \\ -1 & \text{if } m_{s_i, s_j} = \infty. \end{cases}$$

The linear map $\sigma_i : V \to V$ given by $\sigma_i(v) = v - 2B(e_i, v)e_i$, defines an automorphism of V.

Theorem 2.6 (Tits, see [6, p. 96]). The map $\rho : W(M) \to GL(V)$ defined through $\rho(s_i) = \sigma_i$ is a faithful representation of W(M).

The representation ρ above is called the Tits (or geometrical) representation of W(M). Let us denote by I_m the identity matrix of size m. In particular for T_n , the Tits representation ρ is given by $t_i \mapsto \sigma_i$ where

$$\sigma_1 = \begin{pmatrix} -1 & 0 & \\ 2 & 1 & \\ \hline & 0 & I_{n-3} \end{pmatrix}, \qquad \sigma_{n-1} = \begin{pmatrix} I_{n-3} & 0 & \\ \hline & 1 & 2 \\ 0 & 0 & -1 \end{pmatrix}$$

and for 1 < i < n - 1,

	(I_{i-2})		0			
		1	2	0		
$\sigma_i =$	0	0	-1	0	0	
		0	2	1		
	$\sqrt{-0}$		0		I_{n-i-2}	

2.4 Chevyshev Polynomials

There are two types of Chebyshev polynomials: the first and the second kind. For the purpose of this paper, we only need to recall the definition of the Chebyshev polynomials of second kind. For a more detailed treatment on Chebyshev polynomials see [20].

Definition 2.7. The Chebyshev polynomial of second kind $U_n(z)$ is the polynomial of degree n in one variable z defined by the recurrence relation

$$U_n(z) = 2zU_{n-1}(z) - U_{n-2}(z), (2.6)$$

and the initial conditions

$$U_0(z) = 1$$
 and $U_1(z) = 2z.$ (2.7)

In the rest of the paper we simply refer to the polynomials U_i 's as the Chebyshev polynomials. In Section 4, we prove that certain polynomials associated to a family of twins can be written in terms of the Chebyshev polynomials (Corollary 4.4).

3 A Deformed Tits Representation for the Twin Group

The main purpose of this section is to define a deformation of the Tits representation and to give some of its properties.

For $n \ge 3$, we consider the following $(n-1) \times (n-1)$ matrices defined over the ring $\Lambda := \mathbb{Z}(x, y)$

$$V_{1} = \begin{pmatrix} -1 & 0 & \\ x & 1 & \\ \hline & 0 & I_{n-3} \end{pmatrix}, \qquad V_{n-1} = \begin{pmatrix} I_{n-3} & 0 & \\ \hline & 1 & y \\ 0 & 0 & -1 \end{pmatrix}$$

and for 1 < i < n - 1,

$$V_i = \begin{pmatrix} I_{i-2} & 0 & 0 \\ 1 & y & 0 & \\ 0 & 0 & -1 & 0 & 0 \\ 0 & x & 1 & \\ \hline 0 & 0 & I_{n-i-2} \end{pmatrix}.$$

For n = 2, we set $V_1 := -1$.

Lemma 3.1. *For any* $n \ge 3$ *and* $1 \le i \le n - 2$ *,*

$$V_i V_{i+1} V_i - V_{i+1} V_i V_{i+1} = (xy - 1)(V_i - V_{i+1}).$$

Proof. It suffices to verify the case i = 2 and n = 4, which follows from a direct computation.

Matrices V_i 's are an analogue to the matrices used in the reduced Burau representation of the braid group [7]. In fact, we have the following proposition.

Proposition 3.2. For $n \ge 2$, the function $\psi_n : T_n \longrightarrow GL_{n-1}(\Lambda)$ defined by $\psi_n(t_i) = V_i$ is a representation of the twin group.

Proof. The proof follows by checking the defining relations of the twin group are satisfied by the matrices V_i 's. For n = 3 the checking results from a direct computation. In the general case, it is enough to check for n = 4, which follows again by direct computation:

$$(\psi_4(t_2))^2 = \begin{pmatrix} 1 & y & 0 \\ 0 & -1 & 0 \\ 0 & x & 1 \end{pmatrix} \begin{pmatrix} 1 & y & 0 \\ 0 & -1 & 0 \\ 0 & x & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and

$$\begin{split} \psi_4(t_1)\psi_4(t_3) &= \begin{pmatrix} -1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & y \\ 0 & 0 & -1 \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 \\ x & 1 & y \\ 0 & 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & y \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} -1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \psi_4(t_3)\psi_4(t_1). \end{split}$$

Remark 3.3. Specializing x = y = 2, the matrices V_i 's become the matrices σ_i 's seen in Subsection 2.3. Thus, ψ_n can be regarded as a two parameters deformation of the Tits representation of the twin group T_n .

Observe that,

$$\psi_{n+1}(\iota^R(t_i)) = \begin{pmatrix} \psi_n(t_i) & 0\\ v^R(t_i) & 1 \end{pmatrix} = \begin{pmatrix} V_i & 0\\ v^R(t_i) & 1 \end{pmatrix} \qquad 1 \le i \le n-1, \qquad (3.1)$$

where $v^{R}(t_{i})$ is the row of length n-1 equal to 0 if i < n-1 and $(0, \ldots, 0, x)$ if i = n-1. This implies that for $n \ge 2$ and $\beta \in T_{n}$,

$$\psi_{n+1}(\iota^R(\beta)) = \begin{pmatrix} \psi_n(\beta) & 0\\ v^R(\beta) & 1 \end{pmatrix}, \qquad (3.2)$$

where $v^{R}(\beta)$ is a row of length n - 1 over Λ depending on β . In Corollary 3.7 we will see how to compute $v^{R}(\beta)$.

Remark 3.4. The analogous of Equation (3.2) for the inclusion ι^L is

$$\psi_{n+1}(\iota^{L}(\beta)) = \begin{pmatrix} 1 & v^{L}(\beta) \\ 0 & \psi_{n}(\beta) \end{pmatrix}, \qquad (3.3)$$

where $v^{L}(\beta)$ is a row of length n-1 over Λ depending on β .

In order to compute the values of ψ_n on the elements defined in 2.4 and 2.5, we need to introduce the following notation: given a matrix $M = (m_{r,s})$ of size $k \times l$, we denote $M^{\tau} = (m'_{r,s})$ the matrix whose entries are $m'_{r,s} := m_{k-r+1,l-s+1}$. For $n \ge 2$, let $A_n = (a_{i,j})$, $\tilde{Z}_n = (\tilde{z}_{i,j})$ be the $n \times n$ matrices given by

$$a_{i,j} := \begin{cases} -y^{j-1} & i = 1, \ 1 \le j \le n, \\ 0 & i < j-1, \\ x & i = j+1, \ 1 \le j \le n-1, \\ y^{j-i}(xy-1) & 1 < j \le i, \end{cases} \quad \tilde{z}_{i,j} := \begin{cases} xy^j - 2y^{j-1} & i = 1, \\ x^iy^j - x^{i-1}y^{j-1} & 1 < i < n, \\ -x^{n-1}y^{j-1} & i = n. \end{cases}$$

Finally, we set $Z_n := \tilde{Z}_n + I_n$. For instance, for n = 3 we have

$$A_{3} = \begin{pmatrix} -1 & -y & -y^{2} \\ x & xy - 1 & xy^{2} - y \\ 0 & x & xy - 1 \end{pmatrix}, \qquad Z_{3} = \begin{pmatrix} xy - 1 & xy^{2} - 2y & xy^{3} - 2y^{2} \\ x^{2}y - x & x^{2}y^{2} - xy + 1 & x^{2}y^{3} - xy^{2} \\ -x^{2} & -x^{2}y & -x^{2}y^{2} + 1 \end{pmatrix}.$$

Let $M_{k,l}(\Lambda)$ denote the ring of $k \times l$ matrices over Λ . In the lemma below we use the natural map $\theta : M_{k,l}(\Lambda) \to M_{k,l}(\Lambda)$ induced from the \mathbb{Z} -automorphism of Λ that interchanges x with y. **Lemma 3.5.** For any $n \ge 2$ and $1 \le i \le n - 1$,

- (a) $\psi_{n+1}(t_1\cdots t_n)=A_n$,
- (b) $\psi_{n+1}(t_{n,i}) = B_{n,i}$,
- (c) $\psi_{n+1}(t_{1,i}) = C_{n,i}$,

where $B_{n,i}$ and $C_{n,i}$ are defined as follows:

$$B_{n,i} = \begin{cases} \left(\begin{array}{c|c} I_{n-i-1} & Y_{i+1} \\ \hline 0 & Z_{i+1} \end{array} \right) & 1 \le i \le n-2, \\ Z_n & i = n-1, \end{cases} \qquad C_{n,i} = \begin{cases} \left(\begin{array}{c|c} \theta(Z_{i+1})^{\tau} & 0 \\ \hline \theta(Y_{i+1})^{\tau} & I_{n-i-1} \end{array} \right) & 1 \le i \le n-2, \\ \theta(Z_n)^{\tau} & i = n-1, \end{cases}$$

and Y_i is the $(n - i - 1) \times i$ matrix given by

$$Y_i = \begin{pmatrix} \vdots \\ y \quad y^2 \quad \cdots \quad y^i \end{pmatrix},$$

where vertical dots mean a block of 0's for $1 \le i < n - 2$.

Proof. (a) It follows by induction on n and Equation (3.2).

(b) For $1 \le i \le n-2$, we proceed by induction over *i*. For i = 1 the result is straightforward. We want to prove $\psi_{n+1}(t_{n,i+1}) = B_{n,i+1}$. Using Lemma 3.1, we obtain

$$\psi_{n+1}(t_{n,i+1}) = \psi_{n+1}(t_{n-i-1})\psi_{n+1}(t_{n,i})\psi_{n+1}(t_{n-i-1}) -(xy-1)(\psi_{n+1}(t_{n-i-1})-\psi_{n+1}(t_{n,i})).$$
(3.4)

Since $\psi_{n+1}(t_{n,i}) = B_{n,i}$ by induction hypothesis, we have

$$\psi_{n+1}(t_{n,i}) = \begin{pmatrix} I_{n-i-3} & & & \\ & 1 & 0 & 0 & \\ & 0 & 1 & y & r_1 \\ & 0 & 0 & xy - 1 & r_2 \\ \hline & & c_1 & A \end{pmatrix},$$

where r_1 and r_2 are row vectors of length *i*, c_1 is a column vector of length *i* and *A* is a $i \times i$ matrix. Hence

$$\psi_{n+1}(t_{n-i-1})\psi_{n+1}(t_{n,i})\psi_{n+1}(t_{n-i-1}) = \begin{pmatrix} I_{n-i-3} & & & \\ & 1 & xy^2 & y^2 & yr_1 \\ & 0 & -xy+1 & -y & -r_1 \\ & 0 & 2x^2y - 2x & 2xy - 1 & xr_1 + r_2 \\ \hline & & xc_1 & c_1 & A \end{pmatrix}$$

and

$$\psi_{n+1}(t_{n-i-1}) - \psi_{n+1}(t_{n,i}) = \begin{pmatrix} & & & & \\ 0 & y & 0 & & \\ 0 & -2 & -y & -r_1 \\ 0 & x & 2 - xy & -r_2 \\ \hline & & -c_1 & I_i - A \end{pmatrix}.$$

Replacing the expressions above in Equation (3.4), we obtain

$$\psi_{n+1}(t_{n,i+1}) = \begin{pmatrix} I_{n-i-3} & & & \\ & 1 & y & y^2 & yr_1 \\ & 0 & xy-1 & y(xy-2) & (xy-2)r_1 \\ & 0 & x^2y-x & y(x^2y-x)+1 & xr_1+xyr_2 \\ \hline & & xc_1 & xyc_1 & (1-xy)I_i - xyA \end{pmatrix}.$$

Finally, using that $xyr_2 = (x^2y - 2x)r_1$, we obtain that $\psi_{n+1}(t_{n,i+1}) = B_{n,i+1}$. For the case i = n - 1, the result follows by using Lemma 3.1 and the formula for $B_{n,n-2}$.

(c) This case follows by Lemma 3.1 and a similar calculation as in (b).

From now on, we take x = y. Thus, the representation ψ_n is defined over $\Lambda = \mathbb{Z}(x)$. For $n \ge 2$ define the vectors F^R and F^L of Λ^n as follows:

$$F^{R} := \left(U_{0}\left(\frac{1}{x}\right), U_{1}\left(\frac{1}{x}\right), \dots, U_{n-2}\left(\frac{1}{x}\right), U_{n-1}\left(\frac{1}{x}\right)\right) \in \Lambda^{n}$$

and

$$F^{L} := \left(U_{n-1}\left(\frac{1}{x}\right), U_{n-2}\left(\frac{1}{x}\right), \dots, U_{1}\left(\frac{1}{x}\right), U_{0}\left(\frac{1}{x}\right)\right) \in \Lambda^{n},$$

where $U_i(z)$ is the *i*-th Chebyshev polynomial.

Proposition 3.6. For any $\beta \in T_n$, the vectors F^R and F^L satisfy the following equations:

(a) $F^R \psi_{n+1}(\iota^R(\beta)) = F^R$, (b) $F^L \psi_{n+1}(\iota^L(\beta)) = F^L$.

Proof. (a) It suffices to prove it for matrices of the form (3.1). Expanding equations

$$F^{R}\begin{pmatrix} V_{i} & 0\\ v^{R}(t_{i}) & 1 \end{pmatrix} = F^{R} \quad \text{for } 0 \leq i \leq n-1 ,$$

it turns out to be equivalent to the system of equations:

$$-2U_0\left(\frac{1}{x}\right) + xU_1\left(\frac{1}{x}\right) = 0, \qquad (3.5)$$

$$xU_{i-1}\left(\frac{1}{x}\right) - 2U_i\left(\frac{1}{x}\right) + xU_{i+1}\left(\frac{1}{x}\right) = 0. \qquad (2 \le i \le n-2)$$
(3.6)

Now, Equation (3.5) is satisfied by the initial conditions (2.7) of the Chebyshev polynomials, and Equation (3.6) is exactly their recurrence relation (2.6). So, the proof is concluded.

(b) This case is treated similarly.

The next corollary allow us to compute vectors $v^R(\beta)$, $v^L(\beta)$ on matrices (3.2), (3.3), respectively. This will be useful to understand the compatibility of the representation ψ_n with the inclusions ι^R and ι^L , essential on the Gotin-Markov moves M0, M2, M3.

Corollary 3.7. Let $\beta \in T_n$ and let $(a_{1,k} a_{2,k} \cdots a_{n-1,k})$ be the k-th column of the matrix $\psi_n(\beta) - I_{n-1}$ for $1 \le k \le n-1$. Then, the entries of $v^R(\beta) = (b_1 \, b_2 \cdots b_{n-1})$ in matrix (3.2) and $v^{L}(\beta) = (c_1 c_2 \cdots c_{n-1})$ in matrix (3.3) satisfy:

(a)
$$-U_{n-1}\left(\frac{1}{x}\right)b_k = \sum_{i=1}^{n-1} U_{i-1}\left(\frac{1}{x}\right)a_{i,k},$$

(b) $-U_{n-1}\left(\frac{1}{x}\right)c_k = \sum_{i=1}^{n-1} U_{n-2-i}\left(\frac{1}{x}\right)a_{i,k}.$

Proof. We only prove (a), a similar argument works for (b). By Proposition 3.6,

$$F^{R}\left(\begin{array}{cc}\psi_{n}(\beta) & 0\\v^{R}(\beta) & 1\end{array}\right) = F^{R}.$$

Because $F^R I_n = F^R$, we obtain

$$F^{R}\left(\begin{array}{cc}\psi_{n}(\beta) - I_{n-1} & 0\\ v^{R}(\beta) & 0\end{array}\right) = 0.$$
(3.7)

Writing

$$\left(\begin{array}{c} \psi_n(\beta) - I_{n-1} \\ v^R(\beta) \end{array}\right) = \left(\begin{array}{cccc} a_{11} & a_{12} & \cdots & a_{1,n-1} \\ a_{21} & a_{22} & \cdots & a_{2,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n-1,1} & a_{n-1,2} & \cdots & a_{n-1,n-1} \\ b_1 & b_2 & \cdots & b_{n-1} \end{array}\right),$$

and expanding the product of Equation (3.7), we obtain the equations

$$a_{1,k} + U_1\left(\frac{1}{x}\right)a_{2,k} + U_2\left(\frac{1}{x}\right)a_{3,k} + \dots + U_{n-2}\left(\frac{1}{x}\right)a_{n-1,k} + U_{n-1}\left(\frac{1}{x}\right)b_k = 0,$$

where $1 \le k \le n - 1$. These equations imply the claim (a).

4 An Invariant of Doodles

This is the main section of the paper and consists in two subsections. In the first one we introduce a family of functions f_n 's and we study its behavior under the Gotin-Markov moves. Also, we give a a skein relation that is satisfied by f_n . In the next subsection we present our polynomial invariant for doodles Q.

4.1 The function f_n

Definition 4.1. For all $n \ge 1$, we define the polynomial $P_n(x) \in \Lambda$ by the formula

$$P_n(x) = \det(\psi_{n+1}(t_1 \dots t_n) - I_n).$$

We set $P_0(x) = 1$.

Proposition 4.2. For $n \ge 2$, the polynomials P_n satisfy the following recurrence relation:

$$P_n(x) = -2P_{n-1}(x) - x^2 P_{n-2}(x).$$

Proof. By Lemma 3.5, $P_n(x) = \det(A_n - I_n)$, where

$$A_n - I_n = \begin{pmatrix} -2 & -x & -x^2 & \cdots & -x^{n-2} & -x^{n-1} \\ x & x^2 - 2 & x(x^2 - 1) & \cdots & x^{n-3}(x^2 - 1) & x^{n-2}(x^2 - 1) \\ 0 & x & x^2 - 2 & \cdots & x^{n-4}(x^2 - 1) & x^{n-3}(x^2 - 1) \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & x & x^2 - 2 & x(x^2 - 1) \\ 0 & 0 & \cdots & 0 & x & x^2 - 2 \end{pmatrix}.$$

Then, expanding the determinant with respect the last row, we get

$$P_n(x) = (x^2 - 2)P_{n-1}(x) - x\det(M),$$
(4.1)

where M is the following $(n-1) \times (n-1)$ matrix

$$M = \begin{pmatrix} -2 & -x & -x^2 & \cdots & -x^{n-3} & -x^{n-1} \\ x & x^2 - 2 & x(x^2 - 1) & \cdots & x^{n-4}(x^2 - 1) & x^{n-2}(x^2 - 1) \\ 0 & x & x^2 - 2 & \cdots & x^{n-5}(x^2 - 1) & x^{n-3}(x^2 - 1) \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & x & x^2 - 2 & x^2(x^2 - 1) \\ 0 & 0 & \cdots & 0 & x & x(x^2 - 1) \end{pmatrix}.$$

We have that det(M) = x det(M'), where

$$M' = \begin{pmatrix} -2 & -x & -x^2 & \cdots & -x^{n-3} & -x^{n-2} \\ x & x^2 - 2 & x(x^2 - 1) & \cdots & x^{n-4}(x^2 - 1) & x^{n-3}(x^2 - 1) \\ 0 & x & x^2 - 2 & \cdots & x^{n-5}(x^2 - 1) & x^{n-4}(x^2 - 1) \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & x & x^2 - 2 & x(x^2 - 1) \\ 0 & 0 & \cdots & 0 & x & x^2 - 1 \end{pmatrix}.$$

Finally, it is not difficult to prove that $det(M') = P_{n-1}(x) + P_{n-2}(x)$. Then, the result follows by replacing it in Equation (4.1).

Remark 4.3. Observe that by Proposition (4.2), $P_n(x) \in \mathbb{Z}[x^2]$ for all n.

Corollary 4.4. For any $n \ge 0$,

$$P_n(x) = (-x)^n U_n\left(\frac{1}{x}\right).$$

Proof. Firstly, we have that $P_0(x) = 1 = (-x)^0 U_0\left(\frac{1}{x}\right)$ and $P_1(x) = -2 = -xU_1\left(\frac{1}{x}\right)$. Secondly, by the recurrence equation of the Chebyshev polynomials (2.6), we have

$$U_n\left(\frac{1}{x}\right) = \frac{2}{x}U_{n-1}\left(\frac{1}{x}\right) - U_{n-2}\left(\frac{1}{x}\right)$$

which implies $(-x)^n U_n\left(\frac{1}{x}\right) = -2(-x)^{n-1}U_{n-1}\left(\frac{1}{x}\right) - x^2(-x)^{n-2}U_{n-2}\left(\frac{1}{x}\right)$. That is, the polynomial $(-x)^n U_n\left(\frac{1}{x}\right)$ satisfies the same defining rules of $P_n(x)$, hence they are all equal.

Definition 4.5. Let $f_n : T_n \to \Lambda$ be the map defined by

$$f_n(\beta) := \begin{cases} 1 & \text{if } n = 1, \\ \\ \frac{\det(\psi_n(\beta) - I_{n-1})}{(-x)^{n-1}U_{n-1}(\frac{1}{x})} & \text{if } n \ge 2. \end{cases}$$

In particular $f_n(1) = 0$ and $f_n(t_1t_2...t_{n-1}) = 1$ for any $n \ge 2$.

The rest of the subsection is devoted to investigating the compatibility of functions f_n 's under the Gotin-Markov moves, it is divided into Lemmas 4.6, 4.8, 4.9 and Proposition 4.11. Some of the proofs might be troublesome by their length, nevertheless the calculations are straightforward and we show the required steps. In the beginning of Subsection 4.2, it is summarized the properties proved in the rest of this subsection.

Lemma 4.6. For any $n \ge 2$ and $\alpha, \beta \in T_n$,

- (a) $f_{n+1}(\iota^R(\beta)) = f_{n+1}(\iota^L(\beta)) = 0,$
- (b) $f_n(\alpha^{-1}\beta\alpha) = f_n(\beta)$.

Proof. The proof of (a) follows by Equation (3.2) and Equation (3.3). The proof of (b) follows from conjugacy properties of determinants.

Notation 4.7. In subsequent proofs, $C_{i,j}(k)$ (resp. $R_{i,j}(k)$) denotes the elementary operation which replaces the *i*-th column C_i (resp. row R_i) by the sum $C_i + kC_j$ (resp. $R_i + kR_j$). For $k \neq 0$, $C_i(k)$ (resp. $R_i(k)$) denotes the elementary operation which multiplies by k the *i*-th column (resp. row), and $C_{i,j}$ (resp. $R_{i,j}$) denotes the elementary operation that exchanges *i*-th and *j*-th columns (resp. rows).

Lemma 4.8. For any $n \ge 2$ and $\alpha, \beta \in T_n$,

- (a) $f_{n+1}(\iota^R(\beta)t_n) = f_n(\beta)$,
- (b) $f_{n+1}(\iota^{L}(\beta)t_{1}) = f_{n}(\beta).$

Proof. (a) The equation $f_{n+1}(\iota^R(\beta)t_n) = f_n(\beta)$ is equivalent to

$$U_n\left(\frac{1}{x}\right)\det(\psi_{n+1}(\iota^R(\beta)t_n) - I_n) = -xU_n\left(\frac{1}{x}\right)\det(\psi_n(\beta) - I_{n-1}).$$
(4.2)

Let

$$\begin{pmatrix} \psi_n(\beta) & 0\\ v^R(\beta) & 1 \end{pmatrix} = \begin{pmatrix} A & B & 0\\ C & D & 0\\ E & F & 1 \end{pmatrix}, \quad \text{(cf. [16, Lemma 3.12])}$$
(4.3)

where A is a square matrix of size (n-2), B is a column of height n-2, C and E are rows of length n-2, and $D, F \in \Lambda$. By Equation (3.2),

$$\psi_{n+1}(\iota^{R}(\beta)t_{n}) = \begin{pmatrix} \psi_{n}(\beta) & 0 \\ v^{R}(\beta) & 1 \end{pmatrix} \begin{pmatrix} I_{n-2} & 0 \\ 0 & 1 & x \\ 0 & 0 & -1 \end{pmatrix}$$
$$= \begin{pmatrix} A & B & 0 \\ C & D & 0 \\ E & F & 1 \end{pmatrix} \begin{pmatrix} I_{n-2} & 0 \\ 0 & 1 & x \\ 0 & 0 & -1 \end{pmatrix}$$
$$= \begin{pmatrix} A & B & xB \\ C & D & xD \\ E & F & xF - 1 \end{pmatrix}.$$
(4.4)

Subtracting the identity to (4.4), we have

$$\psi_{n+1}(\iota^R(\beta)t_n) - I_n = \begin{pmatrix} A - I_{n-2} & B & xB \\ C & D - 1 & xD \\ E & F & xF - 2 \end{pmatrix} \xrightarrow{C_{n,n-1}(-x)} \cdots$$
$$\cdots \rightarrow \begin{pmatrix} A - I_{n-2} & B & 0 \\ C & D - 1 & x \\ E & F & -2 \end{pmatrix} = M.$$
(4.5)

Hence, $det(\psi_{n+1}(\iota^R(\beta)t_n) - I_n) = det(M)$. Observe that

$$\begin{pmatrix} \psi_n(\beta) - I_{n-1} \\ v^R(\beta) \end{pmatrix} = \begin{pmatrix} A - I_{n-2} & B \\ C & D - 1 \\ E & F \end{pmatrix}.$$

Then,

$$U_{n-1}\left(\frac{1}{x}\right)\det(\psi_{n+1}(\iota^{R}(\beta)t_{n}) - I_{n}) = U_{n-1}\left(\frac{1}{x}\right)\det(M)$$

= $\det\begin{pmatrix}A - I_{n-2} & B & 0\\ C & D - 1 & x\\ U_{n-1}\left(\frac{1}{x}\right)E & U_{n-1}\left(\frac{1}{x}\right)F & -2U_{n-1}\left(\frac{1}{x}\right)\end{pmatrix}$
= $\det(N).$

For $1 \le i \le n-2$, if we add $U_{i-1}\left(\frac{1}{x}\right)$ times the *i*-th row of N to its *n*-th row, by Lemma 3.7 and the recurrence relation of Chebyshev polynomials (2.6), we have

$$\det(N) = \det \begin{pmatrix} A - I_{n-2} & B & 0 \\ C & D - 1 & x \\ 0 & 0 & -2U_{n-1}\left(\frac{1}{x}\right) + xU_{n-2}\left(\frac{1}{x}\right) \end{pmatrix}$$
$$= (-2U_{n-1}\left(\frac{1}{x}\right) + xU_{n-2}\left(\frac{1}{x}\right))\det(\psi_n(\beta) - I_{n-1})$$
$$= -xU_n\left(\frac{1}{x}\right)\det(\psi_n(\beta) - I_{n-1}).$$

Obtaining the Equation (4.2) as desired.

(b) This case is treated similarly by Remark 3.4.

Lemma 4.9. For any $n \ge 3$, $\beta \in T_n$ and $1 \le i \le n-1$,

(a) $f_{n+1}(\iota^R(\beta)t_{n,i}) = x^{2i}f_n(\beta),$

(b)
$$f_{n+1}(\iota^L(\beta)t_{1,i}) = x^{2i}f_n(\beta).$$

Proof. (a) We will prove it for $1 \le i \le n-2$. The proof of the case i = n-1 is treated similarly with the corresponding changes for the matrix $\psi_{n+1}(t_{n,n-1})$ (see Lemma 3.5). The equation $f_{n+1}(\iota^R(\beta)t_{n,i}) = x^{2i}f_n(\beta)$ is equivalent to

$$U_{n-1}\left(\frac{1}{x}\right)\det(\psi_{n+1}(\iota^{R}(\beta)t_{n,i}-I_{n})) = -x^{2i+1}U_{n}\left(\frac{1}{x}\right)\det(\psi_{n}(\beta-I_{n-1})).$$
(4.6)

As in Equation (4.3) of the previous lemma, we express $\psi_{n+1}(\iota^R(\beta))$ in blocks as

$$\begin{pmatrix} \psi_n(\beta) & 0\\ v^R(\beta) & 1 \end{pmatrix} = \begin{pmatrix} A & B & 0\\ C & D & 0\\ E & F & 1 \end{pmatrix},$$
(4.7)

where A is a square matrix of size (n - i - 2), C is a $(i + 1) \times (n - i - 2)$ matrix, D is a square matrix of size (i + 1), B and F are rows of length i + 1 and E is a row of length (n-i-2). Note that for the case i = n-2, we omit the blocks A, C and E. For every block, we write its entries by the same letter, for instance $B = (B_l)_{1 \le l \le i+1}$. Using Lemma 3.5, we know the description of $\psi_{n+1}(t_{n,i})$, obtaining the product

$$\psi_{n+1}(\iota^R(\beta)t_{n,i}) = \begin{pmatrix} A & B' \\ C & D' \\ E & F' \end{pmatrix},$$

where A, C, E are as in (4.7), B' and F' are rows of length i+2, and D' is a $(i+1) \times (i+2)$ matrix. By direct computation, we obtain that the entries of the blocks are the following:

$$B_{l}' = \begin{cases} B_{1} & l = 1, \\ xB_{1} + (x^{2} - 1)B_{2} + \sum_{j=1}^{i-1} x^{j}(x^{2} - 1)B_{j+2} & l = 2, \\ x^{l-1}B_{1} + x^{l-2}(x^{2} - 2)B_{2} + \sum_{j=1}^{i-1} [x^{j+l-2}(x^{2} - 1) + \delta_{j}^{l-2}]B_{j+2} & 3 \le l \le i+1 \\ x^{i+1}B_{1} + x^{i}(x^{2} - 2)B_{2} + \sum_{j=1}^{i-1} x^{j+i}(x^{2} - 1)B_{j+2} & l = i+2, \end{cases}$$

$$D_{k,l}' = \begin{cases} D_{k,1} & 1 \le k \le i, \ l = 1, \\ xD_{k,1} + (x^2 - 1)D_{k,2} + \sum_{j=1}^{i-1} x^j (x^2 - 1)D_{k,j+2} & 1 \le k \le i, \ l = 2, \\ x^{l-1}D_{k,1} + x^{l-2}(x^2 - 2)D_{k,2} + \sum_{j=1}^{i-1} [x^{j+l-2}(x^2 - 1) + \delta_j^{l-2}]D_{k,j+2} & 1 \le k \le i, \ 3 \le l \le i+1, \\ x^{i+1}D_{k,1} + x^i (x^2 - 2)D_{k,2} + \sum_{j=1}^{i-1} x^{j+i} (x^2 - 1)D_{k,j+2} & 1 \le k \le i, \ l = i+2, \end{cases}$$

$$F_l' = \begin{cases} F_1 & l = 1, \\ xF_1 + (x^2 - 1)F_2 + \sum_{j=1}^{i-1} x^j (x^2 - 1)F_{j+2} - x^i & l = 2, \\ x^{l-1}F_1 + x^{l-2}(x^2 - 2)F_2 + \sum_{j=1}^{i-1} [x^{j+l-2}(x^2 - 1) + \delta_j^{l-2}]F_{j+2} - x^{i+l-2} & 3 \le l \le i+1, \\ x^{i+1}F_1 + x^i (x^2 - 2)F_2 + \sum_{j=1}^{i-1} x^{j+i} (x^2 - 1)F_{j+2} - x^{2i} + 1 & l = i+2. \end{cases}$$

Subtracting the identity, we obtain the matrix $\psi_{n+1}(\iota^R(\beta)t_{n,i}) - I_n$. Then, applying a sequence of elementary operations to this matrix, we obtain

$$J = \begin{pmatrix} 0 \\ \psi_n(\beta) - I_{n-1} & x \\ x^2 - 2 \\ x(x^2 - 1) \\ \vdots \\ x^{i-1}(x^2 - 1) \\ \hline v^R(\beta) & -x^i \end{pmatrix}.$$

More precisely, we have

$$(\psi_{n+1}(\iota^R(\beta)t_{n,i}) - I_n) \xrightarrow{C_{n,n-i}(-x^i)} \bullet \xrightarrow{R_n(-x^{-i})} \bullet \xrightarrow{C_{n-i-2+l,n-i}(-x^{l-2})} \bullet \xrightarrow{C_{n-i-2+l,n}(x^{l-2})} \cdots$$

$$\cdots \to \bullet \xrightarrow{C_{n-i,n-i-1}(-x)} \bullet \xrightarrow{C_{n-i,n}(-(x^2-1))} \bullet \xrightarrow{C_{n-i,n-i+j}(-x^j(x^2-1))} \bullet \xrightarrow{C_{n,n-i}} J.$$

By Corollary 3.7, we know the description of $v^R(\beta)$ in terms of the rows of $\psi_n(\beta) - I_{n-1}$ and the Chebyshev polynomials. Thus we obtain the matrix

$$K = \begin{pmatrix} 0 & & \\ x_{n} & & \\ & x^{2} - 2 & & \\ & & x(x^{2} - 1) & \\ & & \vdots & \\ & & & \\ \hline & & & \\ 0 & & xU_{n-i-2}\left(\frac{1}{x}\right) + (x^{2} - 2)U_{n-i-1}\left(\frac{1}{x}\right) + \sum_{k=1}^{i-1} x^{i-k}(x^{2} - 1)U_{n-1-k}\left(\frac{1}{x}\right) - x^{i}U_{n-1}\left(\frac{1}{x}\right) \end{pmatrix},$$

by applying the following sequence of elementary operations to J,

$$J \xrightarrow{R_n\left(U_{n-1}\left(\frac{1}{x}\right)\right)} \bullet \xrightarrow{R_{n,j}\left(U_{j-1}\left(\frac{1}{x}\right)\right)} K.$$

Using iteratively the recurrence relation of the Chebyshev polynomials (2.6), we obtain

$$-x^{i+1}U_n\left(\frac{1}{x}\right) = xU_{n-i-2}\left(\frac{1}{x}\right) + (x^2 - 2)U_{n-i-1}\left(\frac{1}{x}\right) + \sum_{k=1}^{i-1} x^{i-k}(x^2 - 1)U_{n-1-k}\left(\frac{1}{x}\right) - x^iU_{n-1}\left(\frac{1}{x}\right).$$

Therefore,

$$K = \begin{pmatrix} 0 \\ x \\ \psi_n(\beta) - I_{n-1} & x^2 - 2 \\ x(x^2 - 1) \\ \vdots \\ x^{i-1}(x^2 - 1) \\ 0 & -x^{i+1}U_n\left(\frac{1}{x}\right) \end{pmatrix}.$$

Finally, following the sequence of elementary operations that transform $\psi_{n+1}(\iota^R(\beta)t_{n,i} - I_n)$ into K, we conclude that:

$$\det(\psi_{n+1}(\iota^R(\beta)t_{n,i}) - I_n) = \frac{x^i}{U_{n-1}\left(\frac{1}{x}\right)} \det(K)$$
$$= -x^{2i+1} \frac{U_n\left(\frac{1}{x}\right)}{U_{n-1}\left(\frac{1}{x}\right)} \det(\psi_n(\beta) - I_{n-1}),$$

and Equation (4.6) follows.

(b) This case is treated similarly to (a) by Remark 3.4.

Example 4.10. The twins $\alpha = (t_1t_2)^3 \in T_3$ and $\beta = (t_1t_2)^3t_3t_2t_3 \in T_4$ are Gotin-Markov equivalent. Let us compute their images under f_n . From the definition,

$$\psi_3(t_1) = \begin{pmatrix} -1 & 0 \\ x & 0 \end{pmatrix}$$
 and $\psi_3(t_2) = \begin{pmatrix} 1 & x \\ 0 & -1 \end{pmatrix}$.

It follows that

$$\psi_3(t_1t_2)^3 = \left(\begin{array}{cc} -x^4 + 3x^2 - 1 & -x^5 + 4x^3 - 3x \\ x^5 - 4x^3 + 3x & x^6 - 5x^4 + 6x^2 - 1 \end{array}\right).$$

Therefore

$$\det(\psi_3(\alpha) - I_2) = \det\left(\begin{array}{cc} -x^4 + 3x^2 - 2 & -x^5 + 4x^3 - 3x \\ x^5 - 4x^3 + 3x & x^6 - 5x^4 + 6x^2 - 2 \end{array}\right)$$
$$= -x^6 + 6x^4 - 9x^2 + 4$$
$$= (-x^2 + 4)(x^4 - 2x^2 + 1).$$

Since $U_2(z) = -1 + 4z^2$, then

$$f_3(\alpha) = \frac{\det(\psi_3(\alpha) - I_2)}{(-x)^2 U_2\left(\frac{1}{x}\right)} = \frac{(-x^2 + 4)(x^4 - 2x^2 + 1)}{-x^2 + 4} = x^4 - 2x^2 + 1.$$

On the other hand,

$$\psi_4(t_1) = \begin{pmatrix} -1 & 0 & 0 \\ x & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \psi_4(t_2) = \begin{pmatrix} 1 & x & 0 \\ 0 & -1 & 0 \\ 0 & x & 0 \end{pmatrix}, \quad \psi_4(t_3) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & x \\ 0 & 0 & -1 \end{pmatrix}.$$

It follows that

$$\psi_4((t_1t_2)^3t_3t_2t_3) = \begin{pmatrix} -x^4 + 3x^2 - 1 & -x^7 + 4x^5 - 4x^3 + 2x & -x^8 + 5x^6 - 8x^4 + 5x^2 \\ x^5 - 4x^3 + 3x & x^8 - 5x^6 + 7x^4 - 4x^2 + 1 & x^9 - 6x^7 + 12x^5 - 10x^3 + 2x \\ x^4 - x^2 & x^7 - 2x^5 + 2x^3 - 2x & x^8 - 3x^6 + 4x^4 - 3x^2 + 1 \end{pmatrix}.$$

Therefore

$$\det(\psi_4(\beta) - I_3) = 4x^8 - 16x^6 + 20x^4 - 8x^2 = 4x^2(x^2 - 2)(x^2 - 1)^2.$$

Since $U_3(z) = -4z + 8z^3$, then

$$f_4(\beta) = \frac{\det(\psi_4(\beta) - I_3)}{(-x)^3 U_3\left(\frac{1}{x}\right)} = \frac{4x^2(x^2 - 2)(x^2 - 1)^2}{4x^2 - 8} = x^2(x^2 - 1)^2 = x^6 - 2x^4 + x^2.$$

Hence, $f_4(\beta) = x^2 f_3(\alpha)$.

Note that with Lemmas 4.6, 4.8 and 4.9, we have actually proved that if $\alpha \in T_n$ and $\beta \in T_m$ are Gotin-Markov equivalent, then $f_n(\alpha)$ and $f_m(\beta)$ are equal up to some factor x^{2k} for some $k \in \mathbb{Z}$.

Proposition 4.11. The function f_n satisfies the skein relation:

$$f_n\left(\left(\begin{array}{c} \\ \\ \\ \\ \end{array}\right)\right) - f_n\left(\left(\begin{array}{c} \\ \\ \\ \\ \\ \end{array}\right)\right) = (x^2 - 1)\left(f_n\left(\left(\begin{array}{c} \\ \\ \\ \\ \\ \end{array}\right)\right) - f_n\left(\left(\begin{array}{c} \\ \\ \\ \\ \\ \\ \end{array}\right)\right)\right).$$
(4.8)

Proof. By (b) of Lemma 4.6, the skein relation (4.8) is equivalent to the equation

$$f_n(\beta t_i t_{i+1} t_i) - f_n(\beta t_{i+1} t_i t_{i+1}) = (x^2 - 1)(f_n(\beta t_i) - f_n(\beta t_{i+1})),$$

for some $\beta \in T_n$. Therefore, (4.8) is equivalent to prove the formula

$$\det(\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1}) - \det(\psi_n(\beta t_{i+1} t_i t_{i+1}) - I_{n-1}) = (x^2 - 1) [\det(\psi_n(\beta t_i) - I_{n-1}) - \det(\psi_n(\beta t_{i+1}) - I_{n-1})].$$
(4.9)

The corresponding matrix for $\psi_n(t_i t_{i+1} t_i)$ is

$$\psi_n(t_i t_{i+1} t_i) = \begin{pmatrix} I_{i-2} & 0 & 0 \\ 1 & x^3 & x^2 & 0 \\ 0 & 0 & -(x^2 - 1) & -x & 0 & 0 \\ 0 & x(x^2 - 2) & x^2 - 1 & 0 \\ 0 & x^2 & x & 1 \\ \hline 0 & 0 & I_{n-i-3} \end{pmatrix}.$$

Then, we may write $\psi_n(\beta)$ in blocks as follows:

$$\psi_n(\beta) = \begin{pmatrix} A & B & C \\ D & E & F \\ G & H & J \end{pmatrix} = \begin{pmatrix} A & B_1 & B_2 & B_3 & B_4 & C \\ \hline & & \vdots & & \\ D & E_{k,1} & E_{k,2} & E_{k,3} & E_{k,4} & F \\ \hline & & \vdots & & \\ \hline & & G & H_1 & H_2 & H_3 & H_4 & J \end{pmatrix}, \qquad (1 \le k \le 4)$$

where A is square matrix of size (i-2), B is a $(i-2) \times (4)$ matrix, C is a $(i-2) \times (n-i-3)$ matrix, D is a $4 \times (i-2)$ matrix, E is a square matrix of size 4, F is a $4 \times (n-i-3)$ matrix, G is a $(n-i-3) \times (i-2)$ matrix, H is a $(n-i-3) \times 4$ matrix and J is a square matrix of size (n-i-3). For $1 \le l \le 4$, we write B_l and H_l for the columns of B and H, respectively. The resulting product of $\psi_n(\beta)$ and $\psi_n(t_it_{i+1}t_i)$ is:

$$\begin{pmatrix} A & B_1 & x^3B_1 - (x^2 - 1)B_2 + x(x^2 - 2)B_3 + x^2B_4 & x^2B_1 - xB_2 + (x^2 - 1)B_3 + xB_4 & B_4 & C \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots \\ D & E_{k,1} & x^3E_{k,1} - (x^2 - 1)E_{k,2} + x(x^2 - 2)E_{k,3} + x^2E_{k,4} & x^2E_{k,1} - xE_{k,2} + (x^2 - 1)E_{k,3} + xE_{k,4} & E_{k,4} & F \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline G & H_1 & x^3H_1 - (x^2 - 1)H_2 + x(x^2 - 2)H_3 + x^2H_4 & x^2H_1 - xH_2 + (x^2 - 1)H_3 + xH_4 & H_4 & J \end{pmatrix}.$$

Subtracting the identity, we obtain $\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1}$. Then, we apply the following sequence of elementary operations

$$\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1} \xrightarrow{C_{i,i+1}(-x)} \bullet \xrightarrow{C_{i+1,i-1}(-x^2)} \bullet \xrightarrow{C_{i+1,i}(x)} \bullet \xrightarrow{C_{i+1,i+2}(-x)} \cdots \cdots \\ \cdots \to \bullet \xrightarrow{C_{i,i+1}(-x)} \bullet \xrightarrow{C_{i+1}(-1)} M,$$

where M coincides with $\psi_n(\beta) - I_{n-1}$ except for the *i*-th and (i + 1)-th columns. More precisely, if we write $\psi_n(\beta) - I_{n-1} = (a_1, a_2, \dots, a_{n-1})$ and $M = (m_1, m_2, \dots, m_{n-1})$ in vector columns $a_i, m_i \in \Lambda^{n-1}$ for $1 \le i \le n-1$, then

$$m_{j} = \begin{cases} a_{i} + xu & \text{for } j = i, \\ a_{i+1} + u & \text{for } j = i+1, \\ a_{j} & \text{otherwise,} \end{cases} \quad \text{where } u = \begin{pmatrix} 0 \\ -x^{2} \\ x \\ -x^{2} + 2 \\ -x \\ 0 \end{pmatrix}^{i \cdot \text{th}}.$$

By the sequence of elementary operations applied to $\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1}$, it follows that $\det(\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1}) = -\det(M)$.

On the other hand, for the term $\psi_n(t_{i+1}t_it_{i+1})$ given by

$$\psi_n(t_{i+1}t_it_{i+1}) = \begin{pmatrix} I_{i-2} & 0 & 0 \\ 1 & x & x^2 & 0 \\ 0 & 0 & x^2 - 1 & x(x^2 - 2) & 0 & 0 \\ 0 & -x & -(x^2 - 1) & 0 \\ 0 & x^2 & x^3 & 1 \\ \hline 0 & 0 & I_{n-i-3} \end{pmatrix},$$

the resulting product of $\psi_n(\beta)$ and $\psi_n(t_{i+1}t_it_{i+1})$ is

$$\begin{pmatrix} A & B_1 & xB_1 + (x^2 - 1)B_2 - xB_3 + x^2B_4 & x^2B_1 + x(x^2 - 2)B_2 - (x^2 - 1)B_3 + x^3B_4 & B_4 & C \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots \\ D & E_{k,1} & xE_{k,1} + (x^2 - 1)E_{k,2} - xE_{k,3} + x^2E_{k,4} & x^2E_{k,1} + x(x^2 - 2)E_{k,2} - (x^2 - 1)E_{k,3} + x^3E_{k,4} & E_{k,4} & F \\ \hline \vdots & \vdots & & \vdots & & \vdots \\ G & H_1 & xH_1 + (x^2 - 1)H_2 - xH_3 + x^2H_4 & x^2H_1 + x(x^2 - 2)H_2 - (x^2 - 1)H_3 + x^3H_4 & H_4 & J \end{pmatrix}.$$

In the same way as before, we subtract the identity and apply the following sequence of elementary operations

$$\psi_n(\beta t_{i+1}t_it_{i+1}) - I_{n-1} \xrightarrow{C_{i+1,i}(-x)} \bullet \xrightarrow{C_{i,i-1}(-x)} \bullet \xrightarrow{C_{i,i+1}(x)} \bullet \xrightarrow{C_{i,i+2}(-x^2)} \cdots \cdots \\ \cdots \to \bullet \xrightarrow{C_{i+1,i}(-x)} \bullet \xrightarrow{C_{i}(-1)} M',$$

where M' coincides with $\psi_n(\beta) - I_{n-1}$ except for the *i*-th and (i+1)-th columns. Namely, $M' = (m'_1, \ldots, m'_{n-1})$ with

$$m'_{j} = \begin{cases} a_{i} + v & \text{for } j = i, \\ a_{i+1} + xv & \text{for } j = i+1, \\ a_{j} & \text{otherwise}, \end{cases} \quad \text{where } v = \begin{pmatrix} 0 \\ -x \\ -x^{2} + 2 \\ x \\ -x^{2} \\ 0 \end{pmatrix}^{i\text{-th}}.$$

The determinant is computed as $\det(\psi_n(\beta t_{i+1}t_it_{i+1}) - I_{n-1}) = -\det(M')$. Consequently, on the left-hand side of Equation (4.9) we have

$$\det(\psi_n(\beta t_i t_{i+1} t_i) - I_{n-1}) - \det(\psi_n(\beta t_{i+1} t_i t_{i+1}) - I_{n-1}) = -\det(M) + \det(M').$$
(4.10)

Recalling that $\psi_n(\beta) - I_{n-1} = (a_1, \dots, a_{n-1})$, in what follows we write dots for entries a_j with $j \neq i, i+1$. Thus we get

$$-\det(M) + \det(M') = -\det(\dots, a_i + xu, a_{i+1} + u, \dots) + \det(\dots, a_i + v, a_{i+1} + xv, \dots)$$

= det(..., v - xu, a_{i+1}, \dots) + det(..., a_i, xv - u, \dots)
= (x^2 - 1)[det(\dots, w, a_{i+1}, \dots) - det(\dots, a_i, w', \dots)], (4.11)

where w, w' are column vectors given by

$$w = \begin{pmatrix} 0 \\ x \\ -2 \\ x \\ 0 \\ 0 \end{pmatrix}^{i-\text{th}} \quad \text{and} \quad w' = \begin{pmatrix} 0 \\ 0 \\ x \\ -2 \\ x \\ 0 \end{pmatrix}^{i-\text{th}}. \quad (4.12)$$

Now we work the right-hand side of Equation (4.9). The resulting product of $\psi_n(\beta)$ and $\psi_n(t_i)$ is:

$$\psi_n(\beta t_i) = \begin{pmatrix} A & B_1 & xB_1 - B_2 + xB_3 & B_3 & B_4 & C \\ \hline & \vdots & \vdots & \vdots & \vdots & \vdots \\ D & E_{k,1} & xE_{k,1} - E_{k,2} + xE_{k,3} & E_{k,3} & E_{k,4} & F \\ \hline & \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline & G & H_1 & xH_1 - H_2 + xH_3 & H_3 & H_4 & J \end{pmatrix}.$$

In a similar fashion, we subtract the identity and apply the following sequence of elementary operations

$$\psi_n(\beta t_i) - I_{n-1} \xrightarrow{C_{i,i-1}(-x)} \bullet \xrightarrow{C_{i,i+1}(-x)} \bullet \xrightarrow{C_i(-1)} P,$$

where P coincides with $\psi_n(\beta) - I_{n-1}$ except for the *i*-th column. Namely, $P = (p_1, \ldots, p_{n-1})$ with

$$p_j = \begin{cases} a_i - w & \text{for } j = i, \\ a_j & \text{otherwise,} \end{cases} \qquad (1 \le j \le n - 1)$$

where w is as in Equation (4.12). By the sequence of elementary operations, it follows that $\det(\psi_n(\beta t_i) - I_{n-1}) = -\det(P)$.

For the other term in the right-hand side of (4.9), the resulting product of $\psi_n(\beta)$ and $\psi_n(t_{i+1})$ is:

$$\psi_n(\beta t_{i+1}) = \begin{pmatrix} A & B_1 & B_2 & xB_2 - B_3 + xB_4 & B_4 & C \\ \hline & \vdots & \vdots & \vdots & \vdots \\ D & E_{k,1} & E_{k,2} & xE_{k,2} - E_{k,3} + xE_{k,4} & E_{k,4} & F \\ \hline & \vdots & \vdots & \vdots & \vdots \\ \hline G & H_1 & H_2 & xH_2 - H_3 + xH_4 & H_4 & J \end{pmatrix}.$$

Following the similar procedure of subtract the identity and applying a sequence of elementary operations, we have

$$\psi_n(\beta t_i) - I_{n-1} \xrightarrow{C_{i+1,i}(-x)} \bullet \xrightarrow{C_{i+1,i+2}(-x)} \bullet \xrightarrow{C_{i+1}(-1)} P'_{i+1,i+2}(-x)$$

where $P' = (p'_1, \ldots, p'_{n-1})$ with

$$p'_{j} = \begin{cases} a_{i+1} - w' & \text{for } j = i+1, \\ a_{j} & \text{otherwise,} \end{cases} \qquad (1 \le j \le n-1)$$

and w' is as in Equation (4.12). By the sequence of elementary operations applied to $\psi_n(\beta t_{i+1}) - I_{n-1}$, its determinant is computed as $\det(\psi_n(\beta t_{i+1}) - I_{n-1}) = -\det(P')$. Calculating the right-hand side of Equation (4.9), we obtain

$$\det(\psi_n(\beta t_i) - I_{n-1}) - \det(\psi_n(\beta t_{i+1}) - I_{n-1}) = -\det(P) + \det(P').$$
(4.13)

Simplifying,

$$-\det(P) + \det(P') = -\det(\dots, a_i - w, a_{i+1}, \dots) + \det(\dots, a_i, a_{i+1} - w', \dots)$$
$$= \det(\dots, w, a_{i+1}, \dots) - \det(\dots, a_i, w', \dots)$$
$$= \frac{-\det(M) + \det(M')}{(x^2 - 1)},$$

where the last equality is given by (4.11). Therefore, by Equations (4.10), (4.13) we obtain Equation (4.9), which is equivalent to the skein relation (4.8). \Box

4.2 The invariant Q

In subsection 4.1 we investigated how functions $f_n : T_n \to \mathbb{Z}(x)$ behave under Gotin-Markov moves. It is well summarized in the following pictures:

$$\begin{array}{l} \mathbf{i} \quad f_{n+1}\left(\begin{array}{c} \boxed{\beta} \\ \hline \beta \end{array} \right) = 0 = f_{n+1}\left(\left| \begin{array}{c} \boxed{\beta} \\ \hline \beta \end{array} \right), \\ \mathbf{ii} \quad f_n\left(\begin{array}{c} \hline \alpha \\ \hline \beta \end{array} \right) = f_n\left(\begin{array}{c} \hline \beta \\ \hline \alpha \end{array} \right), \\ \mathbf{iii} \quad f_{n+1}\left(\begin{array}{c} \boxed{\beta} \\ \hline \end{array} \right) = f_n\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) = f_{n+1}\left(\left| \begin{array}{c} \boxed{\beta} \\ \hline \end{array} \right), \\ \mathbf{iv} \quad f_{n+1}\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) = x^{2i}f_n\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) = f_{n+1}\left(\left| \begin{array}{c} \hline \beta \\ \hline \end{array} \right), \\ \mathbf{v} \quad f_n\left(\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) \right) = x^{2i}f_n\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) = (x^2 - 1)\left(f_n\left(\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) \right) - f_n\left(\left(\begin{array}{c} \hline \beta \\ \hline \end{array} \right) \right), \\ \end{array} \right)$$

for any $\alpha, \beta \in T_n$.

From properties above, it is clear that if we have two equivalent doodles $D \sim D'$, and α, β two twins with closures D and D' respectively. Then, $\alpha \sim_G \beta$ are Gotin-Markov equivalent and the corresponding polynomials $f_n(\alpha)$ and $f_m(\beta)$ are the same up to a factor x^{2k} for some integer k. Furthermore, by the skein relation (4.8), all the polynomials of a doodle D are polynomials in $\mathbb{Z}[x^2]$. In this way, it is natural to define our invariant as follows.

Definition 4.12. For a doodle $D \in \mathcal{D}$, let I_D be the ideal in $\mathbb{Z}[x^2]$ generated by all the polynomials $f_n(\alpha)$ with $\hat{\alpha} \sim D$. We define $\mathcal{Q} : \mathcal{D} \longrightarrow \mathbb{Z}[x^2]$ given by

$$\mathcal{Q}(D) := \gcd(I_D), \tag{4.14}$$

where gcd is the generator of the ideal I_D .

As a direct consequence of all the previous discussion, we have our main result.

Theorem 4.13. The function Q is a polynomial invariant of oriented doodles.

Remark 4.14. Despite the function f_n satisfies the skein relation (4.8), it is important to notice that the invariant Q does not. It is a consequence of reducing the degrees of the polynomials by taking the gcd.

Let us calculate a simple example of the invariant Q.

Example 4.15. According to [4, Theorem 4.2], the first non-trivial doodle has 6 crossings, and is the Borromean doodle (see Figure 2b). Let us compute its polynomial under Q. In Example 4.10, we compute two associated polynomials to the Borromean doodle

through two Gotin-Markov equivalent twins. Furthermore, for any twin β with closure the Borromean doodle, we have that $\deg(f_3((t_1t_2)^3)) \leq \deg(f_m(\beta))$. Thus,

$$\mathcal{Q}\Big(\bigodot\Big) = x^4 - 2x^2 + 1.$$

Remark 4.16. For a doodle D, if there exists a twin α_0 with closure the minimal representative D_0 of D, then Q(D) is obtained with the corresponding twin α_0 , i.e., $Q(D) = f_n(\alpha_0)$. In general, not every minimal representative of a doodle is realizable immediately by the closure of a twin, might be necessary to produce new double points in the diagram.

Analogously to the classical Alexander polynomial of links, the invariant Q vanishes on multi-component doodles with 'unlinked' components.

Proposition 4.17. Let $D = D_1 \bigsqcup D_2$ be a doodle with two disjoint components, then Q(D) = 0.

Proof. If one of the components is the trivial doodle, the result follows easily from the skein relation (4.8), or by definition of f_n and Lemma 4.6. Let $\alpha \in T_n$ and $\beta \in T_m$ be two twins such that $\alpha^{-1} = D_1$ and $\beta = D_2$, then adding vertical strands on the left or right, we have $\iota^R(\alpha), \iota^L(\beta) \in T_{n+m}$ such that $\iota^R(\alpha)\iota^L(\beta) = \iota^L(\beta)\iota^R(\alpha) = D$. By Equation (3.2) and Remark 3.4 it follows

$$\psi_{n+m}(\iota^R(\alpha)\iota^L(\beta)) = \begin{pmatrix} \psi_n(\alpha) & 0 & 0\\ v^R(\alpha) & 1 & v^L(\beta)\\ 0 & 0 & \psi_m(\beta) \end{pmatrix}.$$

Thus, the *n*-th column of $\psi_{n+m}(\iota^R(\alpha)\iota^L(\beta)) - I_{n+m-1}$ is zero and $f_{n+m}(\iota^R(\alpha)\iota^L(\beta))$ vanishes. Since every polynomial associated to the doodle is the same, up to a factor x^{2k} for some integer k, it follows $\mathcal{Q}(D) = 0$.

5 Some Computations

In this section we compute the invariant Q for many doodles found in the literature [4, 9, 19, 22, 23, 25]. The results are summarized in a table in Subsection 5.2.

5.1 Families of doodles

In [4], Bartholomew-Fenn-Kamada-Kamada give a geometric construction of infinite families of doodles from polygons. We only recall one of those examples in detail and the rest is described vaguely or in terms of twins. We refer to [4] for the original constructions.

Example 5.1. Define the doodle B_n as follows. Start with two concentric *n*-gons with vertices X_1, \ldots, X_n and Y_1, \ldots, Y_n , respectively, and add the missing edges to construct the triangles $X_iY_iX_{i+1}$ for $1 \le i \le n$ cyclically modulo *n* (see Figure 8). All vertices of the resulting diagram have valency four, and it is natural how to smooth the edges to obtain the doodle B_n . By the symmetry of the diagram, it is easy to see that $\widehat{(t_1t_2)^n}$ is B_n for any *n*. In particular, when *n* is divisible by 3, the twin $(t_1t_2)^n$ induces the trivial permutation, i.e., is an element of the pure twin group in 3 strands. Thus, the number of components of B_n is three if *n* is divisible by 3, called the *n*-generalized Borromean doodle; otherwise, B_n has one component, called the *n*-poppy doodle.

Computations prompt the invariant Q never vanishes on this family and its restriction to the family is very strong such that distinguishes, i.e., if $Q(B_n) = Q(B_m)$, then n = m.



Figure 8: Generalized Borromean and *n*-poppy doodles.

Example 5.2. The doodle C_n^1 is constructed from the previous example D_n adding a circle as in Figure 9a. In terms of twins, C_n^1 is the closure of the twin $(t_1t_2t_3t_2)^n$. The doodle C_n^1 has four components if n is divisible by 3, otherwise C_n^1 has two components.

The doodle C_n^1 can be generalized to the doodle D_n with r circles added. Written in twins, define the doodle C_n^r as the closure of the twin $(t_1(t_2 \cdots t_{r+1})t_{r+2}(t_{r+1} \cdots t_2))^n$. In this case, the doodle C_n^r has r+3 components if n is divisible by 3, otherwise C_n^r has r+1 components.

Testing the invariant under different families of doodles, unfortunately we found Q vanishes for an infinite family of doodles.

Proposition 5.3. For integers $r \ge 1$ and $n \ge 3$, $\mathcal{Q}(C_n^r) = 0$.



Figure 9: Doodles C_n^r .

Proof. The doodle C_n^r is the closure of the twin $\alpha_r^n = (t_1(t_2 \cdots t_{r+1})t_{r+2}(t_{r+1} \cdots t_2))^n$. Applying induction over r and using iteratively Lemma 3.1, the matrix $\psi_{r+3}(\alpha_r)^n - I_{r+2}$ has r+1 columns equal up to a power of x. More precisely,

$$\psi_{r+3}(\alpha_r)^n - I_{r+2} = (c_1, x^r c_{r+2}, x^{r-1} c_{r+2}, \dots, x c_{r+2}, c_{r+2})$$

written in columns. Hence $det(\psi_{r+3}(\alpha_r)^n - I_{r+2}) = 0$ and the result follows.

Example 5.4. Another infinite family of doodles related to C_n^r (by the geometric construction given in [4]) is the family D_n^1 . It consists of a circle as the skeleton and an array of n trivial doodles O_1, \ldots, O_n overlapping as in Figures 10a,10b. Doodles D_n^r are a natural generalization of D_n^1 by adding r concentric circles in the skeleton. We do not know a general formula of D_n^r written in twins, but for small values of r and n, we have the invariant Q vanishes.



Figure 10: Doodles D_n^r .

5.2 Table

We follow a format very similar to the table of knots given in [1]. The number in the first line following the picture of the doodle, stands for the number of crossings and components of the minimal representative. The second line is the twin representation of the doodle, where the number k denotes the generator t_k . The last line codified the polynomial invariant $Q(D) \in \mathbb{Z}[x^2]$, the first number in curly brackets denotes the half of the

maximum degree of the polynomial and the next sequence in parenthesis denotes the coefficients in even powers, from higher to lower degree. For instance, $\{7\}(1, 2, -1, -2, 1)$ denotes the polynomial $x^{14} + 2x^{12} - x^{10} - 2x^8 + x^6$.

One component doodles







$$: \{11\}(1, -6, 11, -4, -7, 8, -1, -2, 1)$$



$$: \{10\}(1, -24, 218, -960, 2251, -2880, 1962, -648, 81)$$



 $\begin{array}{c} \cdot 21 \\ \cdot \ 7654323(54)^2 654321 \\ 765432(43)^2 5654321 \\ 543 \end{array}$

 $: \{12\}(4, -4, 1)$



 $: \{12\}(1, -22, 211, -1158, 4013, \\ -9142, 13820, -13672, 8518, -3108, \\ 581, -42, 1)$



 $: \{6\}(64, -320, 592, -496, 184, -24, 1)$



 $: \{10\}(1, -18, 137, -574, 1444, -2232, 2083, -1106, 295, -30, 1)$



 $: \{13\}(1, -24, 254, -1560, 6157, \\ -16336, 29618, -36568, 30086, \\ -15792, 4900, -784, 49)$



 $: \{15\}(1, -28, 352, -2624, 12904, \\ -44064, 107104, -186880, 233108, \\ -204528, 122464, -47616, 11088, \\ -1344, 64)$

Multi-component doodles





References

- [1] C. C. Adams. *The knot book*. American Mathematical Society, Providence, RI, 2004. An elementary introduction to the mathematical theory of knots, Revised reprint of the 1994 original.
- [2] V. Bardakov, M. Singh, and A. Vesnin. Structural aspects of twin and pure twin groups. *Geom. Dedicata*, 203:135–154, 2019.

- [3] A. Bartholomew. Virtual Doodles. http://www.layer8.co.uk/maths/ doodles/index.htm, 2018. Updated: 2018-10-06.
- [4] A. Bartholomew, R. Fenn, N. Kamada, and S. Kamada. Doodles on surfaces. *J. Knot Theory Ramifications*, 27(12):1850071, 26, 2018.
- [5] A. Bartholomew, R. Fenn, N. Kamada, and S. Kamada. On Gauss codes of virtual doodles. *J. Knot Theory Ramifications*, 27(11):1843013, 26, 2018.
- [6] N. Bourbaki. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [7] W. Burau. Über Zopfgruppen und gleichsinnig verdrillte Verkettungen. *Abh. Math. Sem. Univ. Hamburg*, 11(1):179–186, 1935.
- [8] S. Dey and K. Gongopadhyay. Commutator subgroups of twin groups and Grothendieck's cartographical groups. *J. Algebra*, 530:215–234, 2019.
- [9] R. Fenn and P. Taylor. Introducing doodles. In *Topology of low-dimensional manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977)*, volume 722 of *Lecture Notes in Math.*, pages 37–43. Springer, Berlin, 1979.
- [10] M. Gerstenhaber, A. Giaquinto, and S. D. Schack. Quantum symmetry. In *Quantum groups (Leningrad, 1990)*, volume 1510 of *Lecture Notes in Math.*, pages 9–46. Springer, Berlin, 1992.
- [11] J. González, J. L. León-Medina, and C. Roque. Linear motion planning with controlled collisions and pure planar braids. *arXiv e-prints*, page arXiv:1902.06190, Feb. 2019.
- [12] K. Gotin. Markov theorem for doodles on two-sphere. *arXiv e-prints*, page arXiv:1807.05337, Jul 2018.
- [13] N. L. Harshman and A. C. Knapp. Anyons from three-body hard-core interactions in one dimension. Ann. Physics, 412:168003, 18, 2020.
- [14] T. Kanta Naik, N. Nanda, and M. Singh. Conjugacy classes and automorphisms of twin groups. *Forum Math.*, 2020. Accepted.
- [15] T. Kanta Naik, N. Nanda, and M. Singh. Some remarks on twin groups. J. Knot Theory Ramifications, 2020. Accepted.

- [16] C. Kassel and V. Turaev. *Braid groups*, volume 247 of *Graduate Texts in Mathematics*. Springer, New York, 2008. With the graphical assistance of Olivier Dodane.
- [17] L. H. Kauffman and S. Lambropoulou. Virtual braids. Fund. Math., 184:159–186, 2004.
- [18] M. Khovanov. Real $K(\pi, 1)$ arrangements from finite root systems. *Math. Res. Lett.*, 3(2):261–274, 1996.
- [19] M. Khovanov. Doodle groups. Trans. Amer. Math. Soc., 349(6):2297–2315, 1997.
- [20] J. C. Mason and D. C. Handscomb. *Chebyshev polynomials*. Chapman & Hall/CRC, Boca Raton, FL, 2003.
- [21] A. B. Merkov. Vassiliev invariants classify flat braids. In *Differential and symplectic topology of knots and curves*, volume 190 of *Amer. Math. Soc. Transl. Ser.* 2, pages 83–102. Amer. Math. Soc., Providence, RI, 1999.
- [22] A. B. Merkov. Vassiliev invariants of doodles, ornaments, etc. *Publ. Inst. Math.* (*Beograd*) (*N.S.*), 66(80):101–126, 1999. Geometric combinatorics (Kotor, 1998).
- [23] A. B. Merkov. Vassiliev invariants classify plane curves and sets of curves without triple intersections. *Mat. Sb.*, 194(9):31–62, 2003.
- [24] J. Mostovoy and C. Roque-Márquez. Planar pure braids on six strands. J. Knot Theory Ramifications, 29(1):1950097, 11, 2020.
- [25] V. A. Vassiliev. On finite order invariants of triple point free plane curves. In *Differential topology, infinite-dimensional Lie algebras, and applications*, volume 194 of *Amer. Math. Soc. Transl. Ser. 2*, pages 275–300. Amer. Math. Soc., Providence, RI, 1999.
- [26] V. A. Voevodsky. Flags and grothendieck cartographical group in higher dimensions. *CSTARCI Math. Preprints*, 1990.
- [27] Z. Wang. Topological quantum computation, volume 112 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2010.

INSTITUTO DE MATEMÁTICAS UNIVERSIDAD DE VALPARAÍSO, GRAN BRETAÑA 1111, VALPARAÍSO 2340000, CHILE. marcelo.flores@uv.cl juyumaya@uvach.cl

INSTITUTO DE MATEMÁTICAS DE LA UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO LEÓN NO.2, ALTOS, OAXACA DE JUÁREZ 68000, MÉXICO bruno@im.unam.mx

croque@im.unam.mx