

# A REMARK ON APPROXIMATING PERMANENTS OF POSITIVE DEFINITE MATRICES

ALEXANDER BARVINOK

May 12, 2020

ABSTRACT. Let  $A$  be an  $n \times n$  positive definite Hermitian matrix with all eigenvalues between 1 and 2. We represent the permanent of  $A$  as the integral of some explicit log-concave function on  $\mathbb{R}^{2n}$ . Consequently, there is a fully polynomial randomized approximation scheme (FPRAS) for  $\text{per } A$ .

## 1. INTRODUCTION AND MAIN RESULTS

Let  $A = (a_{ij})$  be an  $n \times n$  complex matrix. The *permanent* of  $A$  is defined as

$$\text{per } A = \sum_{\sigma \in S_n} \prod_{k=1}^n a_{k\sigma(k)},$$

where  $S_n$  is the symmetric group of all  $n!$  permutations of the set  $\{1, \dots, n\}$ . Recently, there was some interest in efficient computing (approximating)  $\text{per } A$ , when  $A$  is a positive definite Hermitian matrix (as is known, in that case  $\text{per } A$  is real and non-negative), see [A+17] and reference therein. In particular, Anari et al. construct in [A+17] a deterministic algorithm approximating the permanent of a positive semidefinite  $n \times n$  Hermitian matrix  $A$  within a multiplicative factor of  $c^n$  for  $c = e^{1+\gamma} \approx 4.84$ , where  $\gamma \approx 0.577$  is the Euler constant.

In this note, we show that there is a fully polynomially randomized approximation scheme (FPRAS) for permanents of positive definite matrices with the eigenvalues between 1 and 2. Namely, we represent  $\text{per } A$  for such a matrix  $A$  as an integral of an explicitly constructed log-concave function  $f_A : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ , so that a Markov Chain Monte Carlo algorithm can be applied to efficiently approximate

$$\int_{\mathbb{R}^{2n}} f_A(x) dx = \text{per } A,$$

---

1991 *Mathematics Subject Classification.* 15A15, 15A57, 68W20, 60J22, 26B25.

*Key words and phrases.* permanent, positive definite matrices, log-concave measures.

This research was partially supported by NSF Grant DMS 1855428.

see [LV07].

We consider the space  $\mathbb{C}^n$  with the standard norm

$$\|z\|^2 = |z_1|^2 + \dots + |z_n|^2, \quad \text{where } z = (z_1, \dots, z_n).$$

We identify  $\mathbb{C}^n = \mathbb{R}^{2n}$  by identifying  $z = x + iy$  with  $(x, y)$ . For a complex matrix  $L = (l_{jk})$ , we denote by  $L^* = (l_{jk}^*)$  its conjugate, so that

$$l_{jk}^* = \overline{l_{kj}} \quad \text{for all } j, k.$$

We prove the following main result.

**(1.1) Theorem.** *Let  $A$  be an  $n \times n$  positive definite matrix with all eigenvalues between 1 and 2. Let us write  $A = I + B$ , where  $I$  is the  $n \times n$  identity matrix and  $B$  is an  $n \times n$  positive semidefinite Hermitian matrix with eigenvalues between 0 and 1. Further, we write  $B = LL^*$ , where  $L = (l_{jk})$  is an  $n \times n$  complex matrix. We define linear functions  $\ell_1, \dots, \ell_n : \mathbb{C}^n \rightarrow \mathbb{C}$  by*

$$\ell_j(z) = \sum_{k=1}^n l_{jk} z_k \quad \text{for } z = (z_1, \dots, z_n).$$

Let us define  $f_A : \mathbb{C}^n \rightarrow \mathbb{R}_+$  by

$$f_A(z) = \frac{1}{\pi^n} e^{-\|z\|^2} \prod_{j=1}^n (1 + |\ell_j(z)|^2).$$

(1) *Identifying  $\mathbb{C}^n = \mathbb{R}^{2n}$ , we have*

$$\text{per } A = \int_{\mathbb{R}^{2n}} f_A(x, y) \, dx dy.$$

(2) *The function  $f_A : \mathbb{R}^{2n} \rightarrow \mathbb{R}_+$  is log-concave, that is,*

$$f_A(\alpha x_1 + (1 - \alpha)x_2) \geq f_A^\alpha(x_1) f_A^{1-\alpha}(x_2)$$

for any  $x_1, x_2 \in \mathbb{R}^{2n}$  and any  $0 \leq \alpha \leq 1$ .

## 2. PROOFS

We start with a known integral representation of the permanent of a positive semidefinite matrix.

**(2.1) The integral formula.** Let  $\mu$  be the Gaussian probability measure in  $\mathbb{C}^n$  with density

$$\frac{1}{\pi^n} e^{-\|z\|^2} \quad \text{where} \quad \|z\|^2 = |z_1|^2 + \dots + |z_n|^2 \quad \text{for} \quad z = (z_1, \dots, z_n).$$

Let  $\ell_1, \dots, \ell_n : \mathbb{C}^n \rightarrow \mathbb{C}$  be linear functions and let  $B = (b_{jk})$  be the  $n \times n$  matrix,

$$b_{jk} = \mathbf{E} \ell_j \overline{\ell_k} = \int_{\mathbb{C}^n} \ell_j(z) \overline{\ell_k(z)} d\mu(z) \quad \text{for} \quad j, k = 1, \dots, n.$$

Hence  $B$  is a Hermitian positive semidefinite matrix and the Wick formula (see, for example, Section 3.1.4 of [Ba16]) implies that

$$(2.1.1) \quad \text{per } B = \mathbf{E} (|\ell_1|^2 \dots |\ell_n|^2) = \int_{\mathbb{C}^n} |\ell_1(z)|^2 \dots |\ell_n(z)|^2 d\mu(z).$$

Next, we need a simple lemma.

**(2.2) Lemma.** *Let  $q : \mathbb{R}^n \rightarrow \mathbb{R}_+$  be a positive semidefinite quadratic form. Then the function*

$$h(x) = \ln(1 + q(x)) - q(x)$$

*is concave.*

*Proof.* It suffices to check that the restriction of  $h$  onto any affine line  $x(\tau) = \tau a + b$  with  $a, b \in \mathbb{R}^n$  is concave. Thus we need to check that the univariate function

$$G(\tau) = \ln(1 + (\alpha\tau + \beta)^2 + \gamma^2) - (\alpha\tau + \beta)^2 - \gamma^2 \quad \text{for} \quad \tau \in \mathbb{R},$$

where  $\alpha \neq 0$ , is concave, for which it suffices to check that  $G''(\tau) \leq 0$  for all  $\tau$ . Via the affine substitution  $\tau := (\tau - \beta)/\alpha$ , it suffices to check that  $g''(\tau) \leq 0$ , where

$$g(\tau) = \ln(1 + \tau^2 + \gamma^2) - (\tau^2 + \gamma^2).$$

We have

$$g'(\tau) = \frac{2\tau}{1 + \tau^2 + \gamma^2} - 2\tau$$

and

$$\begin{aligned} g''(\tau) &= \frac{2(1 + \tau^2 + \gamma^2) - 4\tau^2}{(1 + \tau^2 + \gamma^2)^2} - 2 \\ &= \frac{2(1 + \tau^2 + \gamma^2) - 4\tau^2 - 2(1 + \tau^2 + \gamma^2)^2}{(1 + \tau^2 + \gamma^2)^2} \\ &= \frac{2 + 2\tau^2 + 2\gamma^2 - 4\tau^2 - 2 - 2\tau^4 - 2\gamma^4 - 4\tau^2 - 4\gamma^2 - 4\tau^2\gamma^2}{(1 + \tau^2 + \gamma^2)^2} \\ &= -\frac{6\tau^2 + 2\gamma^2 + 2\tau^4 + 2\gamma^4 + 4\tau^2\gamma^2}{(1 + \tau^2 + \gamma^2)^2} \leq 0 \end{aligned}$$

and the proof follows. □

**(2.3) Proof of Theorem 1.1.** We have

$$\text{per } A = \text{per}(I + B) = \sum_{J \subset \{1, \dots, n\}} \text{per } B_J,$$

where  $B_J$  is the principal  $|J| \times |J|$  submatrix of  $B$  with row and column indices in  $J$  and where we agree that  $\text{per } B_\emptyset = 1$ . Let us consider the Gaussian probability measure in  $\mathbb{C}^n$  with density  $\pi^{-n} e^{-\|z\|^2}$ . By (2.1.1), we have

$$\text{per } B_J = \mathbf{E} \prod_{j \in J} |\ell_j(z)|^2$$

and hence

$$\text{per } A = \mathbf{E} \prod_{j=1}^n (1 + |\ell_j(z)|^2) = \int_{\mathbb{R}^{2n}} f_A(x, y) \, dx dy,$$

and the proof of Part (1) follows.

We write

$$e^{-\|z\|^2} \prod_{j=1}^n (1 + |\ell_j(z)|^2) = e^{-q(z)} \prod_{j=1}^n (1 + |\ell_j(z)|^2) e^{-|\ell_j(z)|^2},$$

$$\text{where } q(z) = \|z\|^2 - \sum_{j=1}^n |\ell_j(z)|^2.$$

By Lemma 2.2 each function  $(1 + |\ell_j(z)|^2) e^{-|\ell_j(z)|^2}$  is log-concave on  $\mathbb{R}^{2n} = \mathbb{C}^n$  and hence to complete the proof of Part (2) it suffices to show that  $q$  is a positive semidefinite Hermitian form. To this end, we consider the Hermitian form

$$\begin{aligned} p(z) &= \sum_{j=1}^n |\ell_j(z)|^2 = \sum_{j=1}^n \left| \sum_{k=1}^n l_{jk} z_k \right|^2 = \sum_{j=1}^n \sum_{1 \leq k_1, k_2 \leq n} l_{jk_1} \overline{l_{jk_2}} z_{k_1} \overline{z_{k_2}} \\ &= \sum_{1 \leq k_1, k_2 \leq n} c_{k_1 k_2} z_{k_1} \overline{z_{k_2}}, \end{aligned}$$

where

$$c_{k_1 k_2} = \sum_{j=1}^n l_{jk_1} \overline{l_{jk_2}} \quad \text{for } 1 \leq k_1, k_2 \leq n.$$

Hence for the matrix  $C = (c_{k_1 k_2})$  of  $p$ , we have  $C = \overline{L^* L}$ . We note that  $B = LL^*$  and that the eigenvalues of  $B$  lie between 0 and 1. Therefore, the eigenvalues of  $L^* L$  lie between 0 and 1 (in the generic case, when  $L$  is invertible, the matrices  $LL^*$  and  $L^* L$  are similar). Consequently, the eigenvalues of  $C$  lie between 0 and 1 and hence the Hermitian form  $q(z)$  with matrix  $I - C$  is positive semidefinite, which completes the proof of Part (2).  $\square$

## REFERENCES

- [A+17] N. Anari, L. Gurvits, S.O. Gharan, and A. Saberi, *Simply exponential approximation of the permanent of positive semidefinite matrices*, 58th Annual IEEE Symposium on Foundations of Computer Science – FOCS 2017, IEEE Computer Soc., Los Alamitos, CA, 2017, pp. 914–925.
- [Ba16] A. Barvinok, *Combinatorics and Complexity of Partition Functions*, Algorithms and Combinatorics, 30, Springer, Cham, 2016.
- [LV07] L. Lovász and S. Vempala, *The geometry of logconcave functions and sampling algorithms*, Random Structures & Algorithms **30** (2007), no. 3, 307–358.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043,  
USA

*E-mail address:* `barvinok@umich.edu`