

Convergence to consensus for a Hegselmann-Krause-type model with distributed time delay

Alessandro Paolucci*

Abstract

In this paper we study a Hegselmann-Krause opinion formation model with distributed time delay and positive influence functions. Through a Lyapunov functional approach, we provide a consensus result under a smallness assumption on the initial delay. Furthermore, we analyze a transport equation, obtained as mean-field limit of the particle one. We prove global existence and uniqueness of the measure-valued solution for the delayed transport equation and its convergence to consensus under a smallness assumption on the delay, using a priori estimates which are uniform with respect to the number of agents.

Keywords and Phrases: Hegselmann-Krause model, opinion formation, delay, consensus

1 Introduction

In recent years, many researchers have focused their attention to multi-agent systems. One aspect of these models is the natural self-aggregation, which has been studied in different fields such as biology [1], robotics [12], sociology, economics [19], computer science, control theory [21, 22, 28], social sciences [26, 27] and many other areas. In these last decades a large number of mathematical models has been proposed to study the consensus behavior. First order models, such as the Hegselmann-Krause model [16], have been proposed to study opinion formation. We mention also [17], in which bounded confidence yields the so-called clustering phenomenon. Second order models, in particular Cucker-Smale model [11], have been studied by many authors [13, 14, 24], in order to describe, for example, flocking of birds, swarming of bacteria, or schooling of fishes.

In addition, it is reasonable to introduce a delay in the model as a reaction time or simply as a time to receive the information from outside, in order to let the dynamics more realistic. For first order models, we refer to [5, 8, 10], while for delayed Cucker-Smale-type models we mention [6, 7, 15, 25]. In particular, in very recent papers (see [9, 18, 23]), the authors analyzed modified Cucker-Smale models with distributed time delay, thanks to which agents are influenced by the other ones on a time interval $[t - \tau(t), t]$.

Furthermore, delayed and non-delayed kinetic and transport equations associated to the particle multi-agent systems have been studied in [2, 3, 4, 6, 8, 9].

In this paper, we are interested in the evolution of opinions among N agents, with $N \in \mathbb{N}$. Let $x_i \in \mathbb{R}^d$ be the opinion of the i -th agent, for any $i = 1, \dots, N$. Then, the dynamics is given

*Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università di L'Aquila, Via Vetoio, Loc. Coppito, 67010 L'Aquila Italy (alessandro.paolucci2@graduate.univaq.it).

by the following Hegselmann-Krause-type model:

$$\begin{aligned} \frac{dx_i(t)}{dt} &= \frac{1}{Nh(t)} \sum_{j \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ij}(t; s) (x_j(s) - x_i(t)) ds, \quad t > 0 \\ x_i(s) &= x_{i,0}(s), \quad s \in [-\tau(0), 0], \end{aligned} \quad (1.1)$$

where $\tau : [0, +\infty) \rightarrow (0, +\infty)$ is the time delay. It is a function in $W^{1,\infty}([0, T])$, for any $T > 0$ and we assume that $\tau(t) \geq \tau_*$ for some $\tau_* > 0$, and

$$\tau'(t) \leq 0, \quad \forall t \geq 0. \quad (1.2)$$

This implies that $\tau(t) \leq \tau(0)$, for any $t \geq 0$. We stress the fact that constant delays $\tau(t) \equiv \bar{\tau} > 0$ are allowed.

Motivated by [11, 17, 20], we take the communication rates $a_{ij}(t; s)$ either of the form

$$a_{ij}(t; s) = \psi(|x_j(s) - x_i(t)|), \quad (1.3)$$

for any $i, j \in \{1, \dots, N\}$, where $\psi : [0, +\infty) \rightarrow (0, +\infty)$ is a non-increasing function, or

$$a_{ij}(t; s) = \frac{N\psi(|x_j(s) - x_i(t)|)}{\sum_{k=1}^N \psi(|x_k(s) - x_i(t)|)}, \quad \forall t \geq 0. \quad (1.4)$$

Without loss of generality, we can assume that $\psi(0) = 1$. We notice that in both cases we have that

$$\frac{1}{N} \sum_{j=1}^N a_{ij}(t; s) \leq 1, \quad \forall t \geq 0. \quad (1.5)$$

Moreover, $\alpha : [0, \tau(0)] \rightarrow [0, +\infty)$ is a weight function which satisfies

$$\underline{A} := \int_0^{\tau_*} \alpha(s) ds > 0.$$

Furthermore, we define for any $t \geq 0$

$$h(t) := \int_0^{\tau(t)} \alpha(s) ds. \quad (1.6)$$

Remark 1.1. We notice that if $\alpha(s) = \delta_{\tau(t)}(s)$, then system (1.1) can be rewritten as

$$\begin{aligned} \frac{dx_i(t)}{dt} &= \frac{1}{N} \sum_{j \neq i} a_{ij}(t; t - \tau(t)) (x_j(t - \tau(t)) - x_i(t)), \\ x_i(s) &= x_{i,0}(s), \quad s \in [-\tau(0), 0], \end{aligned}$$

which is already analyzed in [8].

We define, now, the following quantity:

$$d_X(t) := \max_{1 \leq i, j \leq N} |x_i(t) - x_j(t)|.$$

Definition 1.2. We say that a solution $\{x_i(t)\}_{i=1, \dots, N}$ to (1.1) converges to consensus if

$$\lim_{t \rightarrow +\infty} d_X(t) = 0.$$

We will prove the following consensus result.

Theorem 1.3. *Let $\{x_i(t)\}_{i=1}^N$ be the solution to (1.1). Suppose that*

$$\left(e^{\tau(0)} - 1\right) h(0) \leq \frac{A\psi(2R)^3}{2 + \psi(2R)^2}. \quad (1.7)$$

Then, there exist two positive constants C, K such that

$$d_X(t) \leq Ce^{-Kt}, \quad \forall t \geq 0. \quad (1.8)$$

Remark 1.4. *Here, we stress the fact that the quantity*

$$\left(e^{\tau(0)} - 1\right) \int_0^{\tau(0)} \alpha(s) ds$$

is increasing with respect to $\tau(0)$. Then, (1.7) represents a smallness assumption on $\tau(0)$. Moreover, the right-hand side of (1.7) is increasing with respect to $\psi(2R)$. Therefore, we observe that if R is small enough and/or the decay of ψ is not too fast, then the quantity

$$\frac{\psi(2R)^3}{2 + \psi(2R)^2}$$

becomes large and consensus occurs for more values of $\tau(0)$.

The transport equation associated to (1.1) can be obtained as mean-field limit of the particle system (1.1) when $N \rightarrow +\infty$. Let $\mathcal{M}(\mathbb{R}^d)$ be the set of probability measures on the space \mathbb{R}^d . Then, the transport equation associated to (1.1) reads as

$$\begin{aligned} \partial_t \mu_t + \operatorname{div} \left(\frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s] ds \mu_t \right) &= 0, \quad x \in \mathbb{R}^d, \quad t \geq 0 \\ \mu_s &= g_s \quad s \in [-\tau(0), 0], \end{aligned} \quad (1.9)$$

where F is given by either

$$F[\mu_s](x) = \int_{\mathbb{R}^d} \psi(|x-y|)(y-x) d\mu_s(y), \quad (1.10)$$

or

$$F[\mu_s](x) = \frac{\int_{\mathbb{R}^d} \psi(|x-y|)(y-x) d\mu_s(y)}{\int_{\mathbb{R}^d} \psi(|x-y|) d\mu_s(y)}, \quad (1.11)$$

according to the choice of (1.3) and (1.4). Furthermore, we take $g_s \in \mathcal{C}([-\tau(0), 0]; \mathcal{M}(\mathbb{R}^d))$.

Definition 1.5. *Let $T > 0$. We say that $\mu_t \in \mathcal{C}([0, T]; \mathcal{M}(\mathbb{R}^d))$ is a weak solution to (1.9) on the time interval $[0, T)$ if for all $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d \times [0, T))$ we have the following result:*

$$\int_0^T \int_{\mathbb{R}^d} \left(\partial_t \varphi + \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](x) ds \cdot \nabla_x \varphi \right) d\mu_t(x) dt + \int_{\mathbb{R}^d} \varphi(x, 0) dg_0(x) = 0, \quad (1.12)$$

where $F[\mu_s]$ is defined as in (1.10) or (1.11).

We will prove the following theorem.

Theorem 1.6. *Let $\mu_t \in \mathcal{C}([0, T]; \mathcal{P}_1(\mathbb{R}^d))$ be a weak solution to (1.9), with compactly supported initial measure $g_s \in \mathcal{C}([-\tau(0), 0]; \mathcal{P}_1(\mathbb{R}^d))$ and let F as in (1.10) or (1.11). Suppose that*

$$\left(e^{\tau(0)} - 1\right) h(0) \leq \frac{A\psi(2R)^3}{2 + \psi(2R)^2}. \quad (1.13)$$

Then, there exists a constant $C > 0$ independent of t such that

$$d_X(\mu_t) \leq \left(\max_{s \in [-\tau(0), 0]} d_X(g_s) \right) e^{-Ct}, \quad (1.14)$$

for all $t \geq 0$, where

$$d_X(\mu_t) := \text{diam supp } \mu_t.$$

The paper is organized as follows. In Section 2 we study the consensus behavior of solution to (1.1), after assuming an upper-bound on the initial delay $\tau(0)$, namely we will prove Theorem 1.3. In Section 3 we focus our attention on system (1.9) and we study the existence and uniqueness of the solution and its convergence to consensus.

2 Consensus results

We notice that d_X may be not differentiable at some $t \geq 0$. Then, we will use a suitable generalized derivative. We define the upper Dini derivative of a continuous function F as follows:

$$D^+ F(t) := \limsup_{h \rightarrow 0^+} \frac{F(t+h) - F(t)}{h}.$$

Before studying the convergence to consensus of the solution to (1.1), we state the following lemma.

Lemma 2.1. *Let $\{x_i(t)\}_{i=1}^N$ be a solution to (1.1). Suppose that the initial functions $x_{i,0}(s)$ are continuous on the time interval $[-\tau(0), 0]$ for all $i = 1, \dots, N$. Set*

$$R := \max_{s \in [-\tau(0), 0]} \max_{1 \leq i \leq N} |x_i(s)|.$$

Then,

$$\max_{1 \leq i \leq N} |x_i(t)| \leq R \quad (2.15)$$

for all $t \geq 0$.

Proof. Let $\epsilon > 0$ and define $R_\epsilon := R + \epsilon$. Set

$$S^\epsilon = \left\{ t > 0 : \max_{1 \leq i \leq N} |x_i(s)| < R_\epsilon, \quad \forall s \in [0, t] \right\}.$$

By continuity, $S^\epsilon \neq \emptyset$. Denote $T^\epsilon := \sup S^\epsilon$ and assume by contradiction that $T^\epsilon < +\infty$. Then,

$$\lim_{t \rightarrow T^{\epsilon-}} \max_{1 \leq i \leq N} |x_i(t)| = R^\epsilon. \quad (2.16)$$

On the other hand, we have that for any $t \leq T^\epsilon$,

$$\begin{aligned}
\frac{1}{2}D^+|x_i(t)|^2 &\leq \left\langle x_i(t), \frac{dx_i(t)}{dt} \right\rangle \\
&= \left\langle x_i(t), \frac{1}{Nh(t)} \sum_{j \neq i} \int_{t-\tau(t)}^t \alpha(t-s)a_{ij}(t;s)(x_j(s) - x_i(t))ds \right\rangle \\
&= \frac{1}{Nh(t)} \sum_{j \neq i} \int_{t-\tau(t)}^t \alpha(t-s)a_{ij}(t;s) \langle x_i(t), x_j(s) - x_i(t) \rangle ds \\
&= \frac{1}{Nh(t)} \sum_{j \neq i} \int_{t-\tau(t)}^t \alpha(t-s)a_{ij}(t;s) (\langle x_i(t), x_j(s) \rangle - |x_i(t)|^2) ds \\
&\leq \frac{1}{Nh(t)} \sum_{j \neq i} \int_{t-\tau(t)}^t \alpha(t-s)a_{ij}(t;s) |x_i(t)| (|x_j(s)| - |x_i(t)|) ds.
\end{aligned}$$

Using (1.5) and the fact that $t \leq T^\epsilon$ yield

$$\frac{1}{2}D^+|x_i(t)|^2 \leq \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s)ds |x_i(t)|(R^\epsilon - |x_i(t)|) = |x_i(t)|(R^\epsilon - |x_i(t)|).$$

Hence, we have that

$$D^+|x_i(t)| \leq R^\epsilon - |x_i(t)|.$$

By Gronwall inequality, we obtain

$$|x_i(t)| \leq e^{-t} (|x_i(0)| - R^\epsilon) + R^\epsilon < R^\epsilon.$$

Therefore,

$$\lim_{t \rightarrow T^\epsilon} \max_{1 \leq i \leq N} |x_i(t)| < R^\epsilon,$$

which is in contradiction with (2.16). Moreover, since ϵ is arbitrary, we obtain (2.15). \blacksquare

Remark 2.2. Thanks to the previous lemma, we can find a control on $a_{ij}(t;s)$ from below. Indeed, for any $i, j \in \{1, \dots, N\}$, for any $t \geq 0$ and $s \in [t - \tau(t), t]$, we have that

$$|x_j(s) - x_i(t)| \leq |x_j(s)| + |x_i(t)| \leq 2R.$$

Hence, from (1.3) and (1.4), we can deduce that

$$a_{ij}(t;s) \geq \psi(2R), \quad \forall t \geq 0. \quad (2.17)$$

Lemma 2.3. Let $\{x_i(t)\}_{i=1}^N$ be the solution to (1.1). Moreover, define

$$\gamma(t) := \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) \int_s^t \max_{1 \leq k \leq N} \left| \frac{dx_k(z)}{dz} \right| dz ds, \quad \forall t \geq 0. \quad (2.18)$$

Then,

$$D^+d_X(t) \leq \frac{2}{\psi(2R)}\gamma(t) - \psi(2R)d_X(t), \quad \forall t \geq 0. \quad (2.19)$$

Proof. Due to continuity of $x_i(t)$, for any $i \in \{1, \dots, N\}$, there exists a sequence of times $\{t_k\}_{k \in \mathbb{N}}$ such that

$$\bigcup_{k \in \mathbb{N}} [t_k, t_{k+1}) = [0, +\infty),$$

and for each $k \in \mathbb{N}$ and for any $t \in (t_k, t_{k+1})$ there exist $i, j \in \{1, \dots, N\}$ such that

$$d_X(t) = |x_i(t) - x_j(t)|.$$

Hence, we have that

$$\begin{aligned} \frac{1}{2} D^+ d_X^2(t) &\leq \left\langle x_i(t) - x_j(t), \frac{dx_i(t)}{dt} - \frac{dx_j(t)}{dt} \right\rangle \\ &= \frac{1}{Nh(t)} \left\langle x_i(t) - x_j(t), \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ik}(t; s) (x_k(s) - x_i(t)) ds \right\rangle \\ &\quad - \frac{1}{Nh(t)} \left\langle x_i(t) - x_j(t), \sum_{k \neq j} \int_{t-\tau(t)}^t \alpha(t-s) a_{jk}(t; s) (x_k(s) - x_j(t)) ds \right\rangle \\ &=: I_1 + I_2. \end{aligned} \tag{2.20}$$

Now, I_1 and I_2 can be rewritten in the following way:

$$\begin{aligned} I_1 &= \frac{1}{Nh(t)} \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ik}(t; s) \langle x_i(t) - x_j(t), x_k(s) - x_k(t) \rangle ds \\ &\quad + \frac{1}{Nh(t)} \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ik}(t; s) \langle x_i(t) - x_j(t), x_k(t) - x_i(t) \rangle ds \end{aligned} \tag{2.21}$$

and

$$\begin{aligned} I_2 &= -\frac{1}{Nh(t)} \sum_{k \neq j} \int_{t-\tau(t)}^t \alpha(t-s) a_{jk}(t; s) \langle x_i(t) - x_j(t), x_k(s) - x_k(t) \rangle ds \\ &\quad - \frac{1}{Nh(t)} \sum_{k \neq j} \int_{t-\tau(t)}^t \alpha(t-s) a_{jk}(t; s) \langle x_i(t) - x_j(t), x_k(t) - x_j(t) \rangle ds. \end{aligned}$$

We observe (as in [8]) that for any $t \geq 0$,

$$\langle x_i(t) - x_j(t), x_k(t) - x_i(t) \rangle \leq 0, \quad \forall k \in \{1, \dots, N\}.$$

Moreover, we notice that for any $i, j \in \{1, \dots, N\}$

$$a_{ij}(t; s) \leq \frac{1}{\psi(2R)} \tag{2.22}$$

in both cases (1.3) and (1.4). Indeed, if a_{ij} are as in (1.4), for any $i, j = 1, \dots, N$, then we obtain (2.22), using (2.17) and the fact that ψ is a non-increasing function with $\psi(0) = 1$. Moreover, if we take a_{ij} as in (1.3), then (2.22) immediately follows, using the fact that $a_{ij}(t; s) \leq 1$, for any $i, j = 1, \dots, N$, and $\psi(2R) \leq 1$. Therefore, using (2.17) and (2.22) in (2.21) yield

$$\begin{aligned} I_1 &\leq \frac{1}{Nh(t) \psi(2R)} \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) |x_k(s) - x_k(t)| ds \\ &\quad + \frac{\psi(2R)}{N} \sum_{k=1}^N \langle x_i(t) - x_j(t), x_k(t) - x_i(t) \rangle. \end{aligned} \tag{2.23}$$

As before, we observe that for any $t \geq 0$

$$-\langle x_i(t) - x_j(t), x_k(t) - x_j(t) \rangle \leq 0, \quad \forall k \in \{1, \dots, N\}.$$

Hence, using again (2.17) and (2.22), we can obtain a similar estimate for I_2 , namely

$$\begin{aligned} I_2 \leq & \frac{1}{Nh(t)} \frac{d_X(t)}{\psi(2R)} \sum_{k \neq j} \int_{t-\tau(t)}^t \alpha(t-s) |x_k(s) - x_k(t)| ds \\ & + \frac{\psi(2R)}{N} \sum_{k=1}^N \langle x_i(t) - x_j(t), x_j(t) - x_k(t) \rangle. \end{aligned} \quad (2.24)$$

Using (2.23) and (2.24) in (2.20), we have that

$$\frac{1}{2} D^+ d_X(t)^2 \leq \frac{2}{Nh(t)} \frac{d_X(t)}{\psi(2R)} \sum_{k=1}^N \int_{t-\tau(t)}^t \alpha(t-s) |x_k(s) - x_k(t)| ds - \psi(2R) d_X(t)^2. \quad (2.25)$$

Moreover, we notice that, for $s < t$,

$$\sum_{k=1}^N |x_k(s) - x_k(t)| \leq \sum_{k=1}^N \int_s^t \left| \frac{dx_k(z)}{dz} \right| dz \leq N \int_s^t \max_{1 \leq k \leq N} \left| \frac{dx_k(z)}{dz} \right| dz.$$

Substituting this estimate in (2.25), we obtain

$$\frac{1}{2} D^+ d_X(t)^2 \leq \frac{2d_X(t)}{\psi(2R)} \gamma(t) - \psi(2R) d_X(t)^2,$$

which yields (2.19). ■

Lemma 2.4. *Let $\{x_i(t)\}_{i=1}^N$ be the solution to (1.1). Then, for any $t \geq 0$*

$$\max_{1 \leq i \leq N} \left| \frac{dx_i(t)}{dt} \right| \leq \frac{1}{\psi(2R)} \gamma(t) + \frac{1}{\psi(2R)} d_X(t). \quad (2.26)$$

Proof. We have that for any $i \in \{1, \dots, N\}$,

$$\begin{aligned} \left| \frac{dx_i(t)}{dt} \right| \leq & \frac{1}{Nh(t)} \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ik}(t; s) |x_k(s) - x_k(t)| ds \\ & + \frac{1}{Nh(t)} \sum_{k \neq i} \int_{t-\tau(t)}^t \alpha(t-s) a_{ik}(t; s) |x_k(t) - x_i(t)| ds. \end{aligned}$$

Using (2.22) yields

$$\left| \frac{dx_i(t)}{dt} \right| \leq \frac{1}{\psi(2R)} \gamma(t) + \frac{1}{\psi(2R)} d_X(t).$$

Taking the maximum, we obtain (2.26). ■

Now, we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Define the following Lyapunov functional:

$$\mathcal{L}(t) := d_X(t) + \beta \int_0^{\tau(t)} \alpha(s) \int_{t-s}^t e^{-(t-\sigma)} \int_{\sigma}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp d\sigma ds,$$

with $\beta > 0$. Then,

$$\begin{aligned} D^+ \mathcal{L}(t) &= D^+ d_X(t) + \beta \tau'(t) \alpha(\tau(t)) \int_{t-\tau(t)}^t e^{-(t-\sigma)} \int_{\sigma}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp d\sigma \\ &\quad - \beta \int_0^{\tau(t)} \alpha(s) e^{-s} \int_{t-s}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp ds \\ &\quad - \beta \int_0^{\tau(t)} \alpha(s) \int_{t-s}^t e^{-(t-\sigma)} \int_{\sigma}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp d\sigma ds \\ &\quad + \beta \max_{1 \leq k \leq N} \left| \frac{dx_k(t)}{dt} \right| \int_0^{\tau(t)} \alpha(s) \int_{t-s}^t e^{-(t-\sigma)} d\sigma ds. \end{aligned}$$

Using $\underline{A} \leq h(t) \leq h(0)$ and $\tau'(t) \leq 0$, we deduce

$$\begin{aligned} D^+ \mathcal{L}(t) &\leq D^+ d_X(t) - \beta e^{-\tau(0)} \underline{A} \gamma(t) - \beta \int_0^{\tau(t)} \alpha(s) \int_{t-s}^t e^{-(t-\sigma)} \int_{\sigma}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp d\sigma ds \\ &\quad + \beta h(0) (1 - e^{-\tau(0)}) \max_{1 \leq k \leq N} \left| \frac{dx_k(t)}{dt} \right|. \end{aligned}$$

Now, since (2.19) and (2.26) hold, we have that

$$\begin{aligned} D^+ \mathcal{L}(t) &\leq \left(\frac{2}{\psi(2R)} - \beta e^{-\tau(0)} \underline{A} + \beta h(0) (1 - e^{-\tau(0)}) \frac{1}{\psi(2R)} \right) \gamma(t) \\ &\quad + \left(-\psi(2R) + \beta h(0) \frac{1 - e^{-\tau(0)}}{\psi(2R)} \right) d_X(t) - \beta \int_0^{\tau(t)} \alpha(s) \int_{t-s}^t e^{-(t-\sigma)} \int_{\sigma}^t \max_{1 \leq k \leq N} \left| \frac{dx_k(\rho)}{d\rho} \right| dp d\sigma ds. \end{aligned}$$

We want to show that for $\tau(0)$ sufficiently small we obtain the existence of $K > 0$ such that

$$D^+ \mathcal{L}(t) \leq -K \mathcal{L}(t), \quad \forall t \geq 0. \quad (2.27)$$

This is true if the following two conditions hold:

$$\frac{2}{\psi(2R)} - \beta e^{-\tau(0)} \underline{A} + \beta h(0) (1 - e^{-\tau(0)}) \frac{1}{\psi(2R)} \leq 0, \quad (2.28)$$

$$-\psi(2R) + \beta h(0) \frac{1 - e^{-\tau(0)}}{\psi(2R)} < 0. \quad (2.29)$$

The inequality (2.29) is satisfied for

$$\beta < \frac{\psi(2R)^2}{h(0)(1 - e^{-\tau(0)})}. \quad (2.30)$$

Now, in order to have (2.28), we need

$$h(0) (e^{\tau(0)} - 1) < \underline{A} \psi(2R).$$

Hence, (2.28) is satisfied if

$$\beta \geq \frac{2}{e^{-\tau(0)} \underline{A}\psi(2R) - h(0)(1 - e^{-\tau(0)})}. \quad (2.31)$$

Then, in order to have the existence of the parameter $\beta > 0$ such that (2.30) and (2.31) hold, we need

$$\frac{2}{e^{-\tau(0)} \underline{A}\psi(2R) - h(0)(1 - e^{-\tau(0)})} < \frac{\psi(2R)^2}{h(0)(1 - e^{-\tau(0)})},$$

which is true for any $\tau(0)$ satisfying (1.7). Choosing

$$K = \min \left\{ \beta, \psi(2R) - \beta h(0) \frac{1 - e^{-\tau(0)}}{\psi(2R)} \right\},$$

we obtain (2.27). We notice that since β satisfies (2.30), then $K > 0$. This implies immediately (1.8). Hence, the theorem is proved. ■

3 Consensus of solution to (1.9)

In this section we want to analyse the transport equation (1.9) associated to (1.1), obtained as mean-field limit of the particle system when $N \rightarrow +\infty$. To do so, we consider ψ Lipschitz continuous and we denote by L its Lipschitz constant.

Before proving the existence and uniqueness of solutions to (1.9), we first recall some tools on probability spaces and measures.

Definition 3.1. Let $\mu, \nu \in \mathcal{M}(\mathbb{R}^d)$ be two probability measures on \mathbb{R}^d . We define the 1-Wasserstein distance between μ and ν as

$$d_1(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y| d\pi(x, y),$$

where $\Pi(\mu, \nu)$ is the space of all couplings for μ and ν , namely all those probability measures on \mathbb{R}^{2d} having as marginals μ and ν :

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} \varphi(x) d\pi(x, y) = \int_{\mathbb{R}^d} \varphi(x) d\mu(x), \quad \int_{\mathbb{R}^d \times \mathbb{R}^d} \varphi(y) d\pi(x, y) = \int_{\mathbb{R}^d} \varphi(y) d\nu(y),$$

for all $\varphi \in \mathcal{C}_b(\mathbb{R}^d)$.

It's well-known that $(\mathcal{P}_1(\mathbb{R}^d), d_1)$ (where \mathcal{P}_1 is the space of all probability measures with finite first-order moment) is a complete metric space. Moreover, in order to prove the existence of solution to (1.9), we need the following definition.

Definition 3.2. Let μ be a Borel measure on \mathbb{R}^d and let $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a measurable map. We define the push-forward of μ via T as the measure

$$T\#\mu(A) := \mu(T^{-1}(A)),$$

for all Borel sets $A \subset \mathbb{R}^d$.

Then, we have the following theorem.

Theorem 3.3. *Consider the system (1.9) with $g_s \in \mathcal{C}([-\tau(0), 0]; \mathcal{P}_1(\mathbb{R}^d))$. Suppose that there exists a constant $R > 0$ such that*

$$\text{supp } g_t \in B^d(0, R),$$

for all $t \in [-\tau(0), 0]$, where $B^d(0, R)$ denotes the ball of radius R in \mathbb{R}^d centered at the origin. Then, for any $T > 0$ there exists a unique weak solution $\mu_t \in \mathcal{C}([0, T]; \mathcal{P}_1(\mathbb{R}^d))$ of (1.9) in the sense of (1.12). Moreover, μ_t is uniformly compactly supported and

$$\mu_t = X(t; \cdot) \# \mu_0, \quad (3.32)$$

where $X(t; \cdot)$ is the solution of the characteristic system associated to (1.9) for any $t \in [0, T]$.

Proof. First of all we claim that for any $t \in [0, T]$, there exist two positive constants $C, K > 0$ such that

$$\left| \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](x) ds - \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](\tilde{x}) ds \right| \leq C|x - \tilde{x}|,$$

for any $x, \tilde{x} \in B^d(0, R)$, and

$$\left| \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](x) ds \right| \leq K,$$

for all $x \in B^d(0, R)$, with F as in (1.10) or in (1.11). The proof of this claim is very similar to [8, Lemma 3.4]. Then, from [2, Theorem 3.10], we deduce that there exists a unique weak solution to (1.9) in the sense of (1.12) and it exists as long as μ_t is compactly supported. Hence, we need to estimate the growth of support. To do so, we set

$$R_X[\mu_t] := \max_{x \in \text{supp } \mu_t} |x|,$$

for $t \in [0, T]$ and we define

$$R_X(t) := \max_{-\tau(0) \leq s \leq t} R_X[\mu_s].$$

Now, we proceed by steps. We consider $t \in [0, \tau_*]$ and we construct the system of characteristics $X(t; x) : [0, \tau_*] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ associated to (1.9):

$$\begin{aligned} \frac{dX(t; x)}{dt} &= \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](X(s; x)) ds, \\ X(0; x) &= x, \quad x \in \mathbb{R}^d. \end{aligned} \quad (3.33)$$

We notice that the system (3.33) is well-defined, since the velocity field

$$\frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s] ds$$

is locally Lipschitz and locally bounded. Then, arguing as in Lemma 2.1, we have that

$$\frac{d|X(t; x)|}{dt} \leq R_X(t) - |X(t; x)|,$$

which yields

$$R_X(t) < R_X(0),$$

for any $t \in [0, \tau_*]$. Thus, we obtain a unique solution μ_t to (1.9) on the time interval $[0, \tau_*]$. We can iterate this process on all the intervals of the type $[k\tau_*, (k+1)\tau_*]$, with $k = 1, 2, \dots$, until we reach the final time T . Moreover, following [2], it's possible to find a measure μ_t which satisfies (3.32) and this is equivalent to the definition of weak solution (1.12). ■

3.1 Consensus behavior

In this subsection we will prove the consensus behavior of the solution to (1.9), with F as in (1.10) or (1.11). To do so, we firstly need the following stability result.

Lemma 3.4. *Let $\mu_t^1, \mu_t^2 \in \mathcal{C}([0, T]; \mathcal{P}_1(\mathbb{R}^d))$ be two weak solutions to (1.9), with compactly supported initial data $g_s^1, g_s^2 \in \mathcal{C}([-\tau(0), 0]; \mathcal{P}_1(\mathbb{R}^d))$ respectively. Then, there exists a constant $C > 0$ depending only on T such that*

$$d_1(\mu_t^1, \mu_t^2) \leq C \max_{s \in [-\tau(0), 0]} d_1(g_s^1, g_s^2), \quad (3.34)$$

for any $t \in [0, T]$.

Proof. For $i = 1, 2$ let $X^i(t; x) : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be the characteristics associated to (1.9), which obey to

$$\begin{aligned} \frac{dX^i(t; x)}{dt} &= \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) F[\mu_s](X^i(s; x)) ds, \\ X^i(0; x) &= x, \end{aligned}$$

for any $x \in \mathbb{R}^d$. We remember that the characteristics X^i are well-defined in $[0, T]$ since, by Theorem 3.3, μ_t^i have uniformly compact support on such interval. Then, we have that

$$\mu_t^i = X^i(t; \cdot) \# \mu_s^i, \quad \forall t, s \in [0, T].$$

Moreover, as before, we define

$$R_{i; X}^T := \max_{s \in [-\tau(0), T]} R_X[\mu_s^i].$$

Then, we choose an optimal transport map between μ_0^1 and μ_0^2 with respect to d_1 (call it $S_0(x)$) such that $\mu_0^2 = S_0 \# \mu_0^1$ and

$$d_1(\mu_0^1, \mu_0^2) = \int_{\mathbb{R}^d} |x - S_0(x)| d\mu_0^1(x).$$

Moreover, we define the map T^t for any $t \in [0, T]$ as

$$T^t := X^2(t; \cdot) \circ S_0 \circ X^1(t; \cdot)^{-1}. \quad (3.35)$$

Therefore, we can write

$$T^t \# \mu_t^1 = \mu_t^2, \quad \forall t \in [0, T]$$

and

$$d_1(\mu_t^1, \mu_t^2) \leq \int_{\mathbb{R}^d} |x - T^t(x)| d\mu_t^1(x) := u(t).$$

Using (3.35) yields

$$u(t) = \int_{\mathbb{R}^d} |X^1(t; x) - X^2(t; S_0(x))| d\mu_0^1(x).$$

Moreover, we extend the definition of T^t on the interval $[-\tau(0), 0]$ and we define $u(t)$ for $t \in [-\tau(0), 0]$ as

$$u(t) := d_1(g_t^1, g_t^2) = \int_{\mathbb{R}^d} |x - T^t(x)| dg_t^1(x).$$

Now, differentiating $u(t)$ and using (3.35), we obtain

$$\frac{du(t)}{dt} \leq \frac{1}{h(t)} \int_{\mathbb{R}^d} \int_{t-\tau(t)}^t \alpha(t-s) |F[\mu_s^1](x) - F[\mu_s^2](T^t(x))| ds d\mu_t^1(x) =: J.$$

We consider, now, the case of F as in (1.10). Then,

$$\begin{aligned} & |F[\mu_s^1](x) - F[\mu_s^2](T^t(x))| \\ & \leq \int_{\mathbb{R}^d} |\psi(|x-y|)(y-x) - \psi(|T^t(x) - T^s(y)|)(T^s(y) - T^t(x))| d\mu_s^1(y) \\ & \leq \int_{\mathbb{R}^d} |\psi(|x-y|) - \psi(|T^t(x) - T^s(y)|)| \cdot |y-x| d\mu_s^1(y) \\ & \quad + \int_{\mathbb{R}^d} \psi(|T^t(x) - T^s(y)|) \cdot |y-x - (T^s(y) - T^t(x))| d\mu_s^1(y) \\ & = (1) + (2). \end{aligned}$$

Now,

$$\begin{aligned} (1) & \leq L \int_{\mathbb{R}^d} |x-y - T^t(x) + T^s(y)| \cdot |y-x| d\mu_s^1(y) \\ & \leq L(|x| + R_{1;X}^T) \left[|x - T^t(x)| + \int_{\mathbb{R}^d} |y - T^s(y)| d\mu_s^1(y) \right], \end{aligned}$$

and

$$(2) \leq |x - T^t(x)| + \int_{\mathbb{R}^d} |y - T^s(y)| d\mu_s^1(y).$$

Therefore, there exists a constant $C > 0$ depending only on T such that

$$J \leq C \left(u(t) + \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s) u(s) ds \right).$$

Now, if we take F as in (1.11), we have that

$$\begin{aligned} & |F[\mu_s^1](x) - F[\mu_s^2](T^t(x))| \\ & = \left| \frac{\int_{\mathbb{R}^d} \psi(|x-y|)(y-x) d\mu_s^1(y)}{\int_{\mathbb{R}^d} \psi(|x-y|) d\mu_s^1(y)} - \frac{\int_{\mathbb{R}^d} \psi(|T^t(x)-y|)(y-T^t(x)) d\mu_s^2(y)}{\int_{\mathbb{R}^d} \psi(|T^t(x)-y|) d\mu_s^2(y)} \right| \\ & \leq \frac{1}{\psi(R_{1;X}^T)} \left| \int_{\mathbb{R}^d} \psi(|x-y|)(y-x) d\mu_s^1(y) - \int_{\mathbb{R}^d} \psi(|T^t(x)-y|)(y-T^t(x)) d\mu_s^2(y) \right| \\ & \quad + \frac{1}{\psi(R_{1;X}^T)\psi(R_{2;X}^T)} \left| \int_{\mathbb{R}^d} \psi(|T^t(x)-y|)(y-T^t(x)) d\mu_s^2(y) \right| \\ & \quad \times \left| \int_{\mathbb{R}^d} \psi(|x-y|) d\mu_s^1(y) - \int_{\mathbb{R}^d} \psi(|T^t(x)-y|) d\mu_s^2(y) \right|. \end{aligned}$$

As before we have that

$$\left| \int_{\mathbb{R}^d} \psi(|x-y|)(y-x)d\mu_s^1(y) - \int_{\mathbb{R}^d} \psi(|T^t(x)-y|)(y-T^t(x))d\mu_s^2(y) \right| \leq [(|x| + R_{1;X}^T)L + 1] (|x - T^t(x)| + u(s)).$$

Furthermore,

$$\left| \int_{\mathbb{R}^d} \psi(|T^t(x)-y|)(y-T^t(x))d\mu_s^2(y) \right| \leq R_{2;X}^T + |T^t(x)|,$$

and

$$\left| \int_{\mathbb{R}^d} \psi(|x-y|)d\mu_s^1(y) - \int_{\mathbb{R}^d} \psi(|T^t(x)-y|)d\mu_s^2(y) \right| \leq L(|x - T^t(x)| + u(s)).$$

Hence, we obtain again the existence of a constant $C > 0$ depending only on L and T such that

$$\frac{du(t)}{dt} \leq C \left(u(t) + \frac{1}{h(t)} \int_{t-\tau(t)}^t \alpha(t-s)u(s)ds \right).$$

Denote

$$\bar{u} = \max_{s \in [-\tau(0), 0]} u(s) = \max_{s \in [-\tau(0), 0]} d_1(g_s^1, g_s^2),$$

and define $w(t) := e^{-Ct}u(t)$. Then, we have that

$$\frac{dw(t)}{dt} \leq \frac{C}{h(t)} \int_{-\tau(0)}^t \alpha(t-s)w(s)ds. \quad (3.36)$$

Thus, we can rewrite (3.36) as

$$\frac{dw(t)}{dt} \leq K\tau(0)\bar{u} + K \int_0^t w(s)ds,$$

for some $K > 0$. This gives us the following estimate:

$$w(t) \leq \tilde{K}\bar{u}, \quad \forall t \in [0, T],$$

for some $\tilde{K} > 0$. Then, by definition of w we have

$$d_1(\mu_t^1, \mu_t^2) \leq u(t) \leq \tilde{K}e^{CT}\bar{u}, \quad \forall t \in [0, T],$$

which gives us the thesis of this lemma. ■

We are finally ready to prove Theorem 1.6.

Proof of Theorem 1.6. Fixed $g_s \in \mathcal{C}([-\tau(0), 0]; \mathcal{P}_1(\mathbb{R}^d))$, we construct the family of N -particle approximations of g_s , which is a family $\{g_s^N\}_{N \in \mathbb{N}}$ such that

$$g_s^N = \sum_{i=1}^N \delta(x - x_i^0(s)),$$

where $x_i^0 \in \mathcal{C}([-\tau(0), 0]; \mathbb{R}^d)$ satisfy

$$\max_{s \in [-\tau(0), 0]} d_1(g_s^N, g_s) \rightarrow 0, \quad \text{as } N \rightarrow +\infty.$$

Moreover, let $\{x_i^N\}$ be the solution to (1.1), with initial data $x_i(s) = x_i^0(s)$ for any $s \in [-\tau(0), 0]$ and we denote

$$\mu_t^N := \sum_{i=1}^N \delta(x - x_i^N(t)),$$

for any $t \in [0, T]$, which is a weak solution to (1.9). Now, since (1.13) holds, then we know that there exists a constant $C > 0$ such that

$$d_X(t) \leq d_X(0)e^{-Ct} \leq \left(\max_{s \in [-\tau(0), 0]} d_X(s) \right) e^{-Ct},$$

for any $t \geq 0$. Fixing $T \geq 0$, by Lemma 3.4 we have that there exists a constant $K > 0$ independent of N such that

$$d_1(\mu_t, \mu_t^N) \leq K \max_{s \in [-\tau(0), 0]} d_1(g_s, g_s^N),$$

for any $t \in [0, T]$, where μ_t is the weak solution to (1.9) with initial measure g_s . Sending $N \rightarrow +\infty$ we have that $d_X(t) \rightarrow d_X(\mu_t)$ and for any $s \in [-\tau(0), 0]$, $d_X(g_s) = d_X(s)$. This gives (1.14) for any $t \in [0, T]$. Since T can be chosen arbitrarily, then the theorem is proved. ■

4 Acknowledgments

The research of the author is partially supported by the GNAMPA 2019 project *Modelli alle derivate parziali per sistemi multi-agente* (INdAM).

References

- [1] S. Camazine, J. L. Deneubourg, N.R. Franks, J. Sneyd, G. Theraulaz and E. Bonabeau, *Self-Organization in Biological Systems*, Princeton University Press, Princeton, NJ, 2001.
- [2] J. Cañizo, J. Carrillo and J. Rosado, A well-posedness theory in measures for some kinetic models of collective motion, *Math. Mod. Meth. Appl. Sci.*, 21(3), (2011), 515–539.
- [3] C. Canuto, F. Fagnani and P. Tilli, A Eulerian approach to the analysis of rendez-vous algorithms, *IFAC Proceedings Volumes*, Vol. 41, (2008), 9039–9044.
- [4] C. Canuto, F. Fagnani and P. Tilli, An Eulerian approach to the analysis of Krause’s consensus models, *SIAM J. Control Optim.*, 50, (2012), 243–265.
- [5] F. Ceragioli and P. Frasca. Continuous and discontinuous opinion dynamics with bounded confidence, *Nonlinear Anal. Real World Appl.*, 13, (2012), 1239–1251.
- [6] Y.-P. Choi and J. Haskovec, Cucker-Smale model with normalized communication weights and time delay, *Kinet. Relat. Models*, 10, (2017), 1011–1033.
- [7] Y.-P. Choi and Z. Li, Emergent behavior of Cucker-Smale flocking particles with heterogeneous time delays, *Appl. Math. Lett.*, 86, (2018), 49–56.
- [8] Y.-P. Choi, A. Paolucci and C. Pignotti, Consensus of the Hegselmann-Krause opinion formation model with time delay, *Preprint 2019 arXiv:1909.02795*.

- [9] Y.-P. Choi and C. Pignotti, Emergent behavior of Cucker-Smale model with normalized weights and distributed time delays, *Netw. Heterog. Media*, Vol. 14 (2019), pp. 789-804.
- [10] Y.-P. Choi and C. Pignotti, Exponential synchronization of Kuramoto oscillators with time delayed coupling, *Preprint 2019, arXiv:1910.00980*.
- [11] F. Cucker and S. Smale, Emergent behaviour in flocks, *IEEE Transactions on Automatic Control*, 52, (2007), 852–862.
- [12] J. P. Desai, J. P. Ostrowski, V. Kumar, Modeling and control of formations of nonholonomic mobile robots, *IEEE Trans. Robot. Automat.*, 17, (2001), 905-908.
- [13] S.Y. Ha and J. G. Liu, A simple proof of the Cucker-Smale flocking dynamics and mean-field limit, *Commun. Math. Sci.*, 7, (2009) 297–325.
- [14] S.Y. Ha and E. Tadmor, From particle to kinetic and hydrodynamic descriptions of flocking, *Kinet. Relat. Models*, 1, (2008) 415–435.
- [15] J. Haskovec and I. Markou, Delayed Cucker-Smale model with and without noise revisited, *Preprint 2018, arXiv:1810.01084*.
- [16] R. Hegselmann and U. Krause, Opinion dynamics and bounded confidence models, analysis, and simulation, *J. Artif. Soc. Soc. Simul.*, 5, (2002), 1–24.
- [17] P. E. Jabin and S. Motsch, Clustering and asymptotic behavior in opinion formation, *J. Differential Equations*, 257, (2014), 4165–4187.
- [18] Z. Liu, X. Li, Y. Liu, X. Wang, Asymptotic Flocking Behavior of the General Finite-Dimensional Cucker–Smale Model with Distributed Time Delays, *Bull. Malays. Math. Sci. Soc.* (2020). <https://doi.org/10.1007/s40840-020-00917-8>
- [19] G. A. Marsan, N. Bellomo and M. Egidi, Towards a mathematical theory of complex socio-economical systems by functional subsystems representation, *Kinet. Relat. Models*, 1, (2008), 249–278.
- [20] S. Motsch, E. Tadmor, A New Model for Self-organized Dynamics and Its Flocking Behavior, *J Stat Phys*, 144, (2011), 923–947.
- [21] B. Piccoli, N. Pouradier Duteil and E. Trélat, Sparse control of Hegselmann-Krause models: Black hole and declustering, *Preprint 2018, arXiv:1802.00615*.
- [22] B. Piccoli, F. Rossi and E. Trélat, Control to flocking of the kinetic Cucker-Smale model, *SIAM J. Mathematical Analysis*, 47(6), (2015), 4685–4719.
- [23] C. Pignotti and I. Reche Vallejo, Asymptotic analysis of a Cucker-Smale system with leadership and distributed delay, in *Trends in Control Theory and Partial Differential Equations* (eds. F. Alabau-Boussouira, F. Ancona, A. Porretta and C. Sinestrari), Springer Indam Ser., 32, (2019), 233-253.
- [24] C. Pignotti and I. Reche Vallejo, Flocking estimates for the Cucker-Smale model with time lag and hierarchical leadership, *J. Math. Anal. Appl.*, 464, (2018), 1313-1332.

- [25] C. Pignotti and E. Trélat, Convergence to consensus of the general finite-dimensional Cucker-Smale model with time-varying delays, *Commun. Math. Sci.*, 16, (2018), 2053–2076.
- [26] S. Y. Pilyugin and M. C. Campi, Opinion formation in voting processes under bounded confidence, *Netw. Heterog. Media*, 14, (2019), 617–632.
- [27] G. Toscani, Kinetic models of opinion formation, *Commun. Math. Sci.*, 4(3), 481–496, 09 2006.
- [28] S. Wongkaew, M. Caponigro and A. Borzì, On the control through leadership of the Hegselmann-Krause opinion formation model, *Math. Models Methods Appl. Sci.*, 25, (2015), 565–585.