ON THE SECOND EIGENVALUE OF RANDOM BIPARTITE BIREGULAR GRAPHS

YIZHE ZHU

ABSTRACT. We consider the spectral gap of a uniformly chosen random (d_1,d_2) -biregular bipartite graph G with $|V_1|=n, |V_2|=m$, where d_1,d_2 could possibly grow with n and m. Let A be the adjacency matrix of G. Under the assumption that $d_1 \geq d_2$ and $d_2 = O(n^{2/3})$, we show that $\lambda_2(A) = O(\sqrt{d_1})$ with high probability. As a corollary, combining the results from [53], we showed that the second singular value of a uniform random d-regular digraph is $O(\sqrt{d})$ for $1 \leq d \leq n/2$ with high probability. Assuming d_2 is fixed and $d_1 = O(n^2)$, we further prove that for a random (d_1, d_2) -biregular bipartite graph, $|\lambda_i^2(A) - d_1| = O(\sqrt{d_1})$ for all $2 \leq i \leq n+m-1$ with high probability. The proofs of the two results are based on the size biased coupling method introduced in [14] for random d-regular graphs and several new switching operations we defined for random bipartite biregular graphs.

1. Introduction

An expander graph is a sparse graph that has strong connectivity properties and exhibits rapid mixing. Expander graphs play an important role in computer science, including sampling, complexity theory, and the design of error-correcting codes (see [30, 2]). When a graph is d-regular, i.e., each vertex has degree d, quantification of expansion is possible based on the eigenvalues of the adjacency matrix. Let A be the adjacency matrix of a d-regular graph. The first eigenvalue $\lambda_1(A)$ is always d. The second eigenvalue in absolute value $\lambda(A) = \max\{\lambda_2(A), -\lambda_n(A)\}$ is of particular interest, since the difference between d and λ , also known as the spectral gap, provides an estimate on the expansion property of the graph. The study of the spectral gap in d-regular graphs with fixed d had the first breakthrough in the Alon-Boppana bound. It was proved in [1, 49] that for d-regular graph $\lambda(A) \geq 2\sqrt{d-1} - o(1)$. Regular graphs with $\lambda(A) \leq 2\sqrt{d-1}$ are called Ramanujan. In [23], Friedman proved Alon's conjecture in [1] that for the uniform model of random d-regular graph with fixed $d \geq 3$, $\lambda(A) \leq 2\sqrt{d-1} + \varepsilon$ asymptotically almost surely for any $\varepsilon > 0$. The result implies almost all random regular graphs are nearly Ramanujan. Bordenave [8] gave a simpler proof that $\lambda(A) \leq 2\sqrt{d-1} + \varepsilon_n$ for a sequence $\varepsilon_n \to 0$ asymptotically almost surely. Very recently, in [32], this estimate was improved to $\lambda(A) \leq 2\sqrt{d-1} + O(n^{-c})$.

A generalization of Alon's question is to consider the spectral gap of random d-regular graphs when d grows with n. In [10] the authors showed that for $d = o(n^{1/2})$, a uniformly distributed random d-regular graph satisfies $\lambda(A) = O(\sqrt{d})$ with high probability. The authors worked with random regular multigraphs drawn from the configuration model and translated the result for the uniform model by the contiguity argument, which hit a barrier at $d = o(n^{1/2})$. The range of d for the bound $\lambda(A) = O(\sqrt{d})$ was extended to $d = O(n^{2/3})$ in [14] by proving concentration results directly for the uniform model. In [53] it was proved that the $O(\sqrt{d})$ bound holds for $n^{\varepsilon} \leq d \leq n/2$ with $\varepsilon \in (0,1)$. Vu in [55, 56] conjectured that $\lambda(A) = (2+o(1))\sqrt{d(1-d/n)}$ with high probability when $d \leq n/2$ and d tends to infinity with n. The combination of the results in [14] and [53] confirms Vu's conjecture up to a multiplicative constant. Recently, the authors in [5] showed that for $n^{\varepsilon} \leq d \leq n^{2/3-\varepsilon}$ with any $\varepsilon > 0$, $\lambda(A) = (2+o(1))\sqrt{d-1}$ with high probability. Later, it was proved in [50] that Vu's conjecture holds for $\log^{10}(n) \ll d \leq cn$, where c is a small constant. In [28], the same bound was proved for $n^{2/3} \ll d < n/2$, which settled this conjecture in the regime $\log^{10} n \ll d < n/2$.

1.1. Random bipartite biregular graphs. In many applications, one would like to construct bipartite expander graphs with two unbalanced disjoint vertex sets, among which bipartite biregular graphs are of particular interest. An (n, m, d_1, d_2) -bipartite biregular graph is a bipartite graph $G = (V_1, V_2, E)$ where

Date: July 18, 2022.

 $^{2000\ \}textit{Mathematics Subject Classification}.\ \textit{Primary 60C05},\ 60B20;\ \textit{Secondary 05C50}.$

Key words and phrases. random bipartite biregular graph, spectral gap, switching, size biased coupling.

 $|V_1|=n, |V_2|=m$ and every vertex in V_1 has degree d_1 and every vertex in V_2 has degree d_2 . Note that we must have $nd_1=md_2=|E|$. When the number of vertices is clear, we call it a (d_1,d_2) -biregular bipartite graph for simplicity. Let $X \in \{0,1\}^{n \times m}$ be a matrix indexed by $V_1 \times V_2$ such that $X_{ij}=1$ if and only if $(i,j) \in E$. The adjacency matrix of a (d_1,d_2) -biregular bipartite graph with $V_1=[n],V_2=[m]$ can be written as

$$A = \begin{bmatrix} 0 & X \\ X^{\top} & 0 \end{bmatrix}.$$

All eigenvalues of A come in pairs as $\{-\lambda,\lambda\}$, where $|\lambda|$ is a singular value of X along with at least |n-m| zero eigenvalues. It's easy to see $\lambda_1(A) = -\lambda_{n+m}(A) = \sqrt{d_1d_2}$. The difference between $\sqrt{d_1d_2}$ and $\lambda_2(A)$ is called the spectral gap for the bipartite biregular graph. The spectral gap of bipartite biregular graphs has found applications in error correcting codes, matrix completion and community detection, see for example [52, 51, 26, 9, 11]. Previous works of [22, 39] showed an analog of Alon-Boppana bound for bipartite biregular graph: for any sequence of (d_1, d_2) -biregular bipartite graphs with fixed d_1 and d_2 , as the number of vertices tends to infinity, for any $\varepsilon > 0$, $\lim \inf_{n \to \infty} \lambda_2 \ge \sqrt{d_1 - 1} + \sqrt{d_2 - 1} - \varepsilon$. In [22], a (d_1, d_2) -biregular bipartite graph is defined to be Ramanujan if $\lambda_2 \le \sqrt{d_1 - 1} + \sqrt{d_2 - 1}$. It was shown in [45] that there exist infinite families of (d_1, d_2) -biregular bipartite Ramanujan graphs for every $d_1, d_2 \ge 3$. Very recently, for fixed d_1 and d_2 , [9] showed that almost all (d_1, d_2) -biregular bipartite graphs are almost Ramanujan in the sense that $\lambda_2 \le \sqrt{d_1 - 1} + \sqrt{d_2 - 1} + \varepsilon_n$ for a sequence $\varepsilon_n \to 0$ asymptotically almost surely.

In this paper, we consider the spectral gap of a uniformly chosen random (d_1, d_2) -biregular bipartite graph with $V_1 = [n], V_2 = [m]$, where d_1, d_2 can possibly grow with n and m. Without loss of generality, we assume $d_1 \geq d_2$. Let $X \in \mathbb{R}^{n \times m}$ be the biadjacency matrix of a bipartite biregular graph. Namely, $X_{uv} = 1$ if $(u, v) \in E, u \in V_1$ and $v \in V_2$. Let $\mathbf{1}_m = (1, \dots, 1)^{\top} \in \mathbb{R}^m$. Since $X^{\top}X\mathbf{1}_m = d_1d_2\mathbf{1}_m$, the largest singular value satisfies $\sigma_1(X) = \sqrt{d_1d_2}$. Moreover, the second eigenvalue of A in (1.1) is equal to $\sigma_2(X)$.

Our first result is the second eigenvalue bound, which is an extension of [9] to the case where d_1, d_2 can possibly grow with n, m.

Theorem 1.1. Let A be the adjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph with $d_1 \ge d_2$. For any K > 0, if $d_2 \le \frac{1}{2}n^{2/3}$, then there exists a constant $\alpha > 0$ depending only on K such that

(1.2)
$$\mathbb{P}\left(\lambda_2(A) \le \alpha \sqrt{d_1}\right) \ge 1 - m^{-K} - e^{-m}.$$

A d-regular digraph on n vertices is a digraph with each vertex having d in-neighbors and d out-neighbors. We can interpret the biadjacency matrix X of a random (n, n, d, d)-bipartite biregular graph as the adjacency matrix of a random d-regular digraph. Therefore the following corollary holds.

Corollary 1.2. Let A be the adjacency matrix of a uniform random d-regular digraph on n vertices. For any K > 0, if $d \le \frac{1}{2}n^{2/3}$, then there exists a constant $\alpha > 0$ depending only on K such that

(1.3)
$$\mathbb{P}\left(\sigma_2(A) \le \alpha \sqrt{d}\right) \ge 1 - n^{-K} - e^{-n}.$$

For fixed d, it was proved in [9] that $\sigma_2(A) \leq 2\sqrt{d-1} + o(1)$ with high probability. In [53] the authors showed $\sigma_2(A) = O(\sqrt{d})$ with high probability when $n^{\varepsilon} \leq d \leq \frac{n}{2}$. Combining Corollary 1.2 and their result, we confirm a conjecture in [15] that a uniform d-regular digraph has $\sigma_2(A) = O(\sqrt{d})$ for $1 \leq d \leq n/2$ with high probability. Although it was not stated in [15], analogous to Vu's conjecture [55, 56], a more precise version of the conjecture can be formulated as $\sigma_2(A) \leq (2 + o(1))\sqrt{d-1}$ for $1 \ll d \leq n/2$.

Order the eigenvalue of A in modulus as $|\lambda_1(A)| \ge |\lambda_2(A)| \ge \cdots \ge |\lambda_n(A)|$. It is also shown in [17] that for random d-regular digraph, $|\lambda_2(A)| \le \sqrt{d} + \varepsilon$ with high probability for any $\varepsilon > 0$. Note that for a d-regular digraph, $\lambda_1(A) = \sigma_1(A) = d$. By Weyl's inequality between eigenvalues and singular values [31, Theorem 3.3.2]), we have $|\lambda_1(A)\lambda_2(A)| \le \sigma_1(A)\sigma_2(A)$, and $|\lambda_2(A)| \le \sigma_2(A)$ for any d-regular digraph. Therefore, Corollary 1.2 and [53, Theorem B] together also imply $|\lambda_2(A)| = O(\sqrt{d})$ with high probability. Very recently, for the permutation model of random regular digraphs, it was proved in [18] that $|\lambda_2(A)| \le (1+\varepsilon)\sqrt{d}$ when $1 \le d \le n^{o(1)}$.

Now we assume d_2 is a bounded constant and d_1 could grow with n, we refine our estimate as follows.

Theorem 1.3. Let A be the adjacency matrix of a random (n, m, d_1, d_2) -bipartite biregular graph with $d_1 \ge d_2$. Let d_2 be a fixed constant independent of n, and $n \ge 4d_2$. For any constants $K, C_1 > 0$, if $d_1 \le C_1 n^2$, there exists $\alpha > 0$ depending only on C_1, d_2, K such that

(1.4)
$$\mathbb{P}\left(\max_{2 \le i \le m+n-1} |\lambda_i^2(A) - d_1| \le \alpha \sqrt{d_1}\right) \ge 1 - n^{-K} - e^{-n}.$$

We see from (1.4) that the absolute values of all nontrivial eigenvalues of A are concentrated around $\sqrt{d_1}$, which is an improved estimate compared to Theorem 1.1. Theorem 1.3 is also used in [20] to study the global eigenvalue fluctuation of random bipartite biregular graphs. We believe the conditions on d_1, d_2 are only technical assumptions due to the limitation of the method we used. It's possible that by other methods one can extend the range of d_1, d_2 such that (1.4) still holds.

2. Size biased coupling

A key ingredient in the proof of our main results is to show the concentration of linear functions for a random matrix. In this section, we collect some useful results on size biased coupling. For more background on size biased coupling, see the survey [3] and Section 3 in [14].

Let X be a nonnegative random variable with $\mu = \mathbb{E}X > 0$. We say X^s has the X-size biased distribution if $\mathbb{E}[Xf(X)] = \mu \mathbb{E}[f(X^s)]$ for all functions f such that the left hand side above exists. We say a pair of random variable (X, X^s) defined on a common probability space is a size biased coupling for X when X^s has the X-size biased distribution. The following lemma for the sum of indicator random variables will be convenient in our setting.

Lemma 2.1 (Lemma 3.1 in [14]). Let $X_i = a_i F_i$ where F_i is a non-constant random variable taking values in $\{0,1\}$ and $a_i \geq 0$. Let $X = \sum_{i=1}^n X_i$. Let $(X_1^{(i)}, \ldots, X_n^{(i)})$ be random variables such that $X_i^{(i)} = a_i$ and $(X_j^{(i)})_{j \neq i}$ are distributed as $(X_j^{(i)})_{j \neq i}$ conditioned on $F_i = 1$. Independent of everything else, choose a random index I such that $\mathbb{P}(I = i) = \mathbb{E}X_i/\mathbb{E}X$ for $1 \leq i \leq n$. Then $X^s = \sum_{i=1}^n X_i^{(I)}$ has the size biased distribution of X.

The following result in [14] provides concentration inequalities from the construction of size biased couplings. For $x \in \mathbb{R}$, define $(x)_+ := \max\{x, 0\}$.

Lemma 2.2 (Theorem 3.4. in [14]). Let (X, X^s) be a size biased coupling with $\mathbb{E}X = \mu$, \mathcal{B} be an event on which $X^s - X \leq c$. Let $D = (X^s - X)_+$ and suppose $\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X] \leq \tau^2/\mu$ almost surely. Define $h(x) = (1+x)\log(1+x) - x$, $x \geq -1$. Then the following holds:

- (1) If $\mathbb{P}[\mathcal{B} \mid X^s] \ge p$ almost surely, then for $x \ge 0$, $\mathbb{P}\left(X \frac{\mu}{p} \ge x\right) \le \exp\left(-\frac{\tau^2}{pc^2}h\left(\frac{pcx}{\tau^2}\right)\right)$.
- (2) If $\mathbb{P}[\mathcal{B} \mid X] \ge p$ almost surely, then for $x \ge 0$, $\mathbb{P}(X p\mu \le -x) \le \exp\left(-\frac{\tau^2}{c^2}h\left(\frac{cx}{\tau^2}\right)\right)$.

3. SWITCHING FOR BIPARTITE BIREGULAR GRAPHS

The method of switchings, developed by McKay and Wormald [46, 57], has been used to approximately enumerate regular graphs, counting subgraphs in random regular graphs, see for example [47, 48, 35, 36]. In recent years, combined with other random matrix techniques, switching has become a useful tool to study the spectra of random regular graphs [33, 14, 7, 4, 6, 5]. It was also applied to spectral analysis of other random graph models including random regular digraphs [15, 16, 13, 40, 41, 42, 43] and random bipartite biregular graphs [59, 58]. The switching operation defines a natural Markov chain called the "switch chain", which is often used to sample random graphs. In [54], the authors derived Poincaré inequalities for the switch chain on d-regular bipartite graphs when $3 \le d \le cn$. It is known that such functional inequalities imply corresponding concentration inequalities [37], which provides a possible approach to study the spectral gap of uniform random regular bipartite graphs.

In this section, we introduce the switching operations on bipartite biregular graphs, which are different from [59] and involve more vertices. Our definition of switchings is an analog of the "double switchings" defined for regular graphs in Section 4 of [14], and is suitable for bipartite biregular graphs. The switchings will be used to construct a coupling between X and $X^{(u_1v_1)}$, where X is the biadjacency matrix of a random bipartite biregular graph and $X^{(u_1v_1)}$ is the distribution of X conditioned on $X_{u_1v_1} = 1$.

Definition 3.1 (valid switchings). Assume $X_{u_1v_2} = X_{u_2v_1} = X_{u_3v_3} = 1$ and $X_{u_1v_1} = X_{u_2v_3} = X_{u_3v_2} = 0$. We define $(u_1, u_2, u_3, v_1, v_2, v_3)$ to be a valid forward switching for X as follows. After switching in the graph G, the edges u_1v_1, u_2v_3, u_3v_2 are added and the edges u_1v_2, u_2v_1, u_3v_3 are removed. In a similar way, suppose $X_{u_1v_2} = X_{u_2v_1} = X_{u_3v_3} = 0$ and $X_{u_1v_1} = X_{u_2v_3} = X_{u_3v_2} = 1$, define $(u_1, u_2, u_3, v_1, v_2, v_3)$ to be a valid backward switching for X if after switching, the edges u_1v_2, u_2v_1, u_3v_3 are added and the edges u_1v_1, u_2v_3, u_3v_2 are removed. In both of the forward and backward switchings, we assume $u_1, u_2, u_3, v_1, v_2, v_3$ are distinct vertices.

See Figure 1 for an example for a valid forward switching. By reversing the arrow in Figure 1, we obtain a valid backward switching from the right to the left. The following lemma estimates the number of valid forward and backward switchings.




FIGURE 1. a valid forward switching

Lemma 3.2. Let $s_{u_1v_1}(G)$ and $t_{u_1u_1}(G)$ be the number of valid forward and backward switchings of the form $(u_1,\cdot,\cdot,v_1,\cdot,\cdot)$, respectively. Then the following inequalities hold:

- (1) If $X_{u_1v_1} = 0$, $d_1^2d_2(n 2d_2) \le s_{u_1v_1}(G) \le d_1^2d_2(n d_2)$. (2) If $X_{u_1v_1} = 1$, $d_1^2(n d_2)(n 2d_2) \le t_{u_1v_1}(G) \le d_1^2(n d_2)^2$.

Proof. Let $\mathcal{N}(v)$ be the set of neighborhood of a vertex v in the graph G. Define $\overline{\mathcal{N}}(v) = V_2 \setminus \mathcal{N}(v)$ if $v \in V_1$, and $\mathcal{N}(v) = V_1 \setminus \mathcal{N}(v)$ if $v \in V_2$. Fix u_1, v_1 and assume $X_{u_1v_1} = 0$. By choosing $u_2 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_1), u_3 \in \mathcal{N}(v_1)$ $\overline{\mathcal{N}}(v_2), v_3 \in \mathcal{N}(u_3)$, we have $d_1^2 d_2(n-d_2)$ many tuples, which gives the upper bound on $s_{u_1v_1}(G)$. Among those $d_1^2d_2(n-d_2)$ many tuples, a tuple is a valid forward switching if and only if $v_3 \in \overline{\mathcal{N}}(u_2)$. By choosing $u_2 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_1), v_3 \in \mathcal{N}(u_2), u_3 \in \mathcal{N}(v_3),$ we have at most $d_1^2 d_2^2$ many tuples that are not valid forward switchings. Therefore $s_{u_1v_1}(G) \geq d_1^2d_2(n-d_2) - d_1^2d_2^2 = d_1^2d_2(n-2d_2)$. This completes the proof for the first claim. Now assume $X_{u_1v_1} = 1$. By choosing $u_2 \in \overline{\mathcal{N}}(v_1), v_3 \in \mathcal{N}(u_2), u_3 \in \overline{\mathcal{N}}(v_3), v_2 \in \mathcal{N}(u_3),$ we have $d_1^2(n-d_2)^2$ many tuples, giving the upper bound of $t_{u_1v_1}(G)$. Among those tuples, a tuple is not a valid backward switching if and only if $v_2 \in \mathcal{N}(u_1)$. We can bound the number of invalid tuples by choosing $v_2 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1), v_3 \in \mathcal{N}(u_2), u_3 \in \mathcal{N}(v_2), \text{ which has at most } d_1^2 d_2(n-d_2) \text{ many.}$ Therefore the second claim holds.

Let \mathcal{G} be the collection of the biadjacency matrices of all (n, m, d_1, d_2) -bipartite biregular graphs. For fixed $u_1 \in [n], v_1 \in [m]$, let $\mathcal{G}_{u_1v_1}$ be the subset of \mathcal{G} such that $X_{u_1v_1} = 1$. We construct an edge-weighted bipartite graph \mathfrak{G}_0 on two vertex class \mathcal{G} and $\mathcal{G}_{u_1v_1}$ as follows:

- If $X \in \mathcal{G}$ with $X_{u_1v_1} = 0$, form an edge of weight 1 between X and every element of $\mathcal{G}_{u_1v_1}$ that is a result of a valid forward switching from X.
- If $X \in \mathcal{G}$ with $X_{u_1v_1} = 1$, form an edge of weight $d_1^2d_2(n d_2)$ between X and its identical copy in $\mathcal{G}_{u_1v_1}$.

Define an edge-weighted bipartite graph with two disjoint vertex sets V_1, V_2 to be a (w_1, w_2) -biregular graph if the degree (sum of weights from adjacent edges) of each vertex in V_1 is w_1 and the degree of each vertex in V_2 is w_2 . The following lemma shows we can embed (allowing edge weight to increase) \mathfrak{G}_0 into a weighted bipartite biregular graph.

Lemma 3.3. In \mathfrak{G}_0 , the following holds:

- (1) Every vertex in \mathcal{G} has degree between $d_1^2d_2(n-2d_2)$ and $d_1^2d_2(n-d_2)$.
- (2) Every vertex in $\mathcal{G}_{u_1v_1}$ has degree between $d_1^2(n-d_2)^2$ and $d_1^2n(n-d_2)$.
- (3) \mathfrak{G}_0 can be embedded into a weighted bipartite biregular graph \mathfrak{G} on the same vertex sets, with vertices in \mathcal{G} having degree $d_1^2d_2(n-d_2)$ and vertices in $\mathcal{G}_{u_1v_1}$ having degree $d_1^2n(n-d_2)$.

Proof. Claim (1) follows from our construction of \mathfrak{G}_0 and the first claim in Lemma 3.2. Every X in \mathcal{G}_{u_1,v_1} with the corresponding graph G is connected to $t_{u_1v_1}(G)$ many vertices in \mathcal{G} , with each edge of weight 1. It is also connected to its identical copy in \mathcal{G} with edge weight $d_1^2d_2(n-d_2)$. Then Claim (2) follows from the second claim in Lemma 3.2. To construct \mathfrak{G} , we start with \mathfrak{G}_0 and add edges as follows. Go through the vertices of \mathcal{G} and for each vertex with degree less than $d_1^2d_2(n-d_2)$, arbitrarily make edges or increase edge weights from the vertex to vertices in $\mathcal{G}_{u_1v_1}$ with degree less than $d_1^2n(n-d_2)$. Continue this procedure until either all vertices in \mathcal{G} have degree $d_1^2d_2(n-d_2)$ or all vertices in \mathcal{G}_{u_1,v_1} have degree $d_1^2n(n-d_2)$. We claim now \mathfrak{G} is bipartite biregular. From the distribution of uniform random bipartite biregular graphs, the probability of a bipartite biregular graph containing any edge u_1v_1 is $\frac{d_2}{n}$, we have $\frac{|\mathcal{G}_{u_1v_1}|}{|\mathcal{G}|} = \frac{d_2}{n}$. If all degrees in \mathcal{G} are $d_1^2d_2(n-d_2)$, and all degrees in $\mathcal{G}_{u_1,u_2,v_1}$ are at most $d_1^2n(n-d_2)$, then $|\mathcal{G}|d_1^2d_2(n-d_2) \leq |\mathcal{G}_{u_1v_1}|d_1^2n(n-d_2) = |\mathcal{G}|d_1^2d_2(n-d_2)$. So all vertices in $\mathcal{G}_{u_1v_1}$ must have degree exactly $d_1^2n(n-d_2)$. In the same way, if all the degrees in $\mathcal{G}_{u_1v_1}$ are $d_1^2n(n-d_2)$, then the degrees in \mathcal{G} are exactly $d_1^2d_2(n-d_2)$. Therefore we can embed \mathfrak{G}_0 into a weighted biregular bipartite graph \mathfrak{G} .

In the graph \mathfrak{G} , we uniformly choose a random biadjacency matrix X in \mathcal{G} and consider $X^{(u_1v_1)}$ to be the element in $\mathcal{G}_{u_1v_1}$ given by walking from X along an edge with probability proportionate to its weight. Since \mathfrak{G} is bipartite biregular, $X^{(u_1v_1)}$ is uniformly distributed in the vertex set $\mathcal{G}_{u_1v_1}$. Lemma 3.3 yields a coupling of X and $X^{(u_1v_1)}$ that satisfies

(3.1)
$$\mathbb{P}\left(X, X^{(u_1v_1)} \text{ are identical or differ by a switching } \mid X^{(u_1v_1)}\right) \ge 1 - \frac{d_1}{m},$$

(3.2)
$$\mathbb{P}\left(X, X^{(u_1v_1)} \text{ are identical or differ by a switching } | X\right) \geq 1 - \frac{d_1}{m - d_1}.$$

To see (3.1) holds, note that \mathfrak{G}_0 is embedded into \mathfrak{G} , from the definition of \mathfrak{G}_0 , if the walk from X to $X^{(u_1v_1)}$ is chosen from an edge in \mathfrak{G}_0 , then $X, X^{(u_1v_1)}$ are identical or differ by a switching. Therefore the left hand side of (3.1) is lower bounded by the probability that an edge adjacent to $X^{(u_1v_1)}$ is chosen from \mathcal{G}_0 . From (2) and (3) in Lemma 3.3, such probability is at least $\frac{d_1^2(n-d_2)^2}{d_1^2n(n-d_2)} = 1 - \frac{d_2}{n} = 1 - \frac{d_1}{m}$. Similarly, (3.2) holds.

4. Concentration for linear functions of X

Let Q be a $n \times m$ matrix and X be the biadjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph. Define a linear function for entries of X as $f_Q(X) := \sum_{u \in [n], v \in [m]} Q_{uv} X_{uv}$. In this section, we will use the coupling we constructed in Section 3 together with Lemma 2.1 to construct a size biased coupling of the linear function $f_Q(X)$.

From the distribution of X, we have $\mathbb{E}X_{uv} = \frac{d_1}{m}$ for any $u \in [n], v \in [m]$. We define the following two parameters:

(4.1)
$$\mu := \mathbb{E} f_Q(X) = \frac{d_1}{m} \sum_{u \in [n], v \in [m]} Q_{uv}, \quad \tilde{\sigma}^2 := \mathbb{E} f_{Q \circ Q}(X) = \frac{d_1}{m} \sum_{u \in [n], v \in [m]} Q_{uv}^2.$$

Theorem 4.1. Let X be the biadjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph. Let Q be a $n \times m$ matrix with all entries in [0, a]. Let $p = 1 - \frac{d_1}{m}$ and $p' = 1 - \frac{d_1}{m-d_1}$. Then for all $t \ge 0$,

(4.2)
$$\mathbb{P}\left(f_Q(X) - \frac{\mu}{p} \ge t\right) \le \exp\left(-\frac{\tilde{\sigma}^2}{3pa^2}h\left(\frac{pat}{\tilde{\sigma}^2}\right)\right),$$

(4.3)
$$\mathbb{P}\left(f_Q(X) - p'\mu \le -t\right) \le \exp\left(-\frac{\tilde{\sigma}^2}{3a^2}h\left(\frac{at}{\tilde{\sigma}^2}\right)\right).$$

Proof. We construct a size biased coupling based on the analysis of switchings in Section 3. Choose a vertex $X \in \mathcal{G}$ uniformly at random and walk through an edge adjacent to the vertex with probability proportional to its weight. We then obtain a uniform random element $X^{(u_1v_1)}$ in $\mathcal{G}_{u_1v_1}$. The matrix $X^{(u_1v_1)}$ is distributed as X conditioned on the event $X_{u_1v_1} = 1$. Independently of X, we choose a random variable (U_1, V_1) such that for all $u \in [n], v \in [m]$,

(4.4)
$$\mathbb{P}(U_1 = u, V_1 = v) = \frac{Q_{uv}}{\sum_{u \in [n], v \in [m]} Q_{uv}}.$$

Define $X' = X^{(U_1,V_1)}$. Since X_{uv} is an indicator random variable for all $u \in [n], v \in [m]$, by Lemma 2.1, the pair $(f_Q(X), f_Q(X'))$ is a size biased coupling.

Let $S(u_1, v_1)$ be the set of all tuples (u_2, u_3, v_2, v_3) such that $(u_1, u_2, u_3, v_1, v_2, v_3)$ is a valid forward switching for X. Let $X(u_1, u_2, u_3, v_1, v_2, v_3)$ be the matrix obtained from X by a valid forward switching $(u_1, u_2, u_3, v_1, v_2, v_3)$. Then for any $(u_2, u_3, v_2, v_3) \in S(U_1, V_1)$, assuming $X_{U_1, V_1} = 0$ and conditioned on X, U_1 and V_1 , the element $X' \in \mathcal{G}_{U_1, V_1}$ is equally likely to be $X(U_1, u_2, u_3, V_1, v_2, v_3)$ for any tuple $(u_2, u_3, v_2, v_3) \in S(X, U_1, V_1)$. Recall from Lemma 3.3, each vertex in \mathcal{G} has degree $d_1^2 d_2(n - d_2)$ in the graph \mathfrak{G} . By our construction of the coupling (X, X'), we have for any tuple $(u_2, u_3, v_2, v_3) \in S(U_1, V_1)$,

$$(4.5) \mathbb{P}(X' = X(U_1, u_2, u_3, V_1, v_2, v_3) \mid U_1, V_1, X, X_{U_1, V_1} = 0) = \frac{1}{d_1^2 d_2 (n - d_2)}.$$

For any valid forward switching $(u_1, u_2, u_3, v_1, v_2, v_3)$,

$$f_Q(X(u_1, u_2, u_3, v_1, v_2, v_3)) - f_Q(X) = Q_{u_1v_1} + Q_{u_2v_3} + Q_{u_3v_2} - Q_{u_1v_2} - Q_{u_2v_1} - Q_{u_3v_3} \le 3a.$$

Let \mathcal{B} be the event that the edge chosen in the random walk on \mathfrak{G} from X to X' belongs to the subgraph \mathfrak{G}_0 . By (3.1) and (3.2), $\mathbb{P}(\mathcal{B} \mid X') \geq p$ and $\mathbb{P}(\mathcal{B} \mid X) \geq p'$, where p, p' are the parameters in the statement of Theorem 4.1. Therefore $f_Q(X') - f_Q(X) \leq 3a$ on the event \mathcal{B} . Let $\overline{\mathcal{S}}(u_1, v_1)$ be the set of (u_2, u_3, v_2, v_3) such that $u_2 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_1), u_3 \in \overline{\mathcal{N}}(v_2), v_3 \in \mathcal{N}(u_3)$. Then $\overline{\mathcal{S}}(u_1, v_1)$ has size $d_1^2 d_2(n - d_2)$ and $\mathcal{S}(U_1, V_1) \subset \overline{\mathcal{S}}(u_1, v_1)$. Let $D = (f_Q(X') - f_Q(X))_+$. Then from (4.5),

$$\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X, U_{1}, V_{1}]$$

$$= \frac{1}{d_{1}^{2}d_{2}(n - d_{2})} \sum_{(u_{2}, u_{3}, v_{2}, v_{3}) \in \mathcal{S}(U_{1}, V_{1})} (f_{Q}(X(U_{1}, u_{2}, u_{3}, V_{1}, v_{2}, v_{3}) - f_{Q}(X))_{+}$$

$$\leq \frac{1}{d_{1}^{2}d_{2}(n - d_{2})} \sum_{(u_{2}, u_{3}, v_{2}, v_{3}) \in \overline{\mathcal{S}}(U_{1}, V_{1})} (Q_{U_{1}V_{1}} + Q_{u_{2}v_{3}} + Q_{u_{3}v_{2}}).$$

Taking the expectation over U_1, V_1 , from (4.4), we have

$$\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X] \\
\leq \sum_{u_{1} \in [n], v_{1} \in [m]} \frac{Q_{u_{1}u_{2}}}{\sum_{u, v} Q_{uv}} \left(\frac{1}{d_{1}^{2}d_{2}(n - d_{2})} \sum_{(u_{2}, u_{3}, v_{2}, v_{3}) \in \overline{S}(u_{1}, v_{1})} (Q_{u_{1}v_{1}} + Q_{u_{2}v_{3}} + Q_{u_{3}v_{2}}) \right) \\
= \frac{1}{d_{1}^{2}(n - d_{2})n\mu} \sum_{\substack{u_{1} \in [n], v_{1} \in [m] \\ (u_{2}, u_{3}, v_{2}, v_{3}) \in \overline{S}(u_{1}, v_{1})}} (Q_{u_{1}v_{1}}^{2} + Q_{u_{1}v_{1}}Q_{u_{2}v_{3}} + Q_{u_{1}v_{1}}Q_{u_{3}v_{2}}) \\$$

The first term in the sum (4.6) satisfies

(4.7)
$$\sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)}} Q_{u_1 v_1}^2 = \sum_{u_1 \in [n], v_1 \in [m]} d_1^2 d_2 (n - d_2) Q_{u_1 u_2}^2 = d_1^2 n (n - d_2) \tilde{\sigma}^2.$$

For the second term in (4.6), by Cauchy's inequality

$$(4.8) \sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)}} Q_{u_1 v_1} Q_{u_2 v_3} \le \left(\sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)}} Q_{u_1 v_1}^2 \right)^{1/2} \left(\sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)}} Q_{u_2 v_3}^2 \right)^{1/2}.$$

For a given (u_2, v_3) , there are $d_1^2 d_2(n - d_2)$ many (u_1, u_3, v_1, v_2) such that $(u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)$. Hence

For a given
$$(u_2, v_3)$$
, there are $d_1^2 d_2(n - d_2)$ many (u_1, u_3, v_1, v_2) such that $(u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}$ (4.9)
$$\sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{\mathcal{S}}(u_1, v_1)}} Q_{u_2 v_3}^2 = d_1^2 d_2(n - d_2) \sum_{u_2 \in [n], v_3 \in [m]} Q_{u_2 v_3}^2 = d_1^2 n(n - d_2) \tilde{\sigma}^2.$$

Therefore with (4.7), the left hand side of (4.8) is bounded by $d_1^2 n(n-d_2)\tilde{\sigma}^2$. By the same argument, the third term in (4.6)

$$\sum_{\substack{u_1 \in [n], v_1 \in [m] \\ (u_2, u_3, v_2, v_3) \in \overline{S}(u_1, v_1)}} Q_{u_1 v_1} Q_{u_3 v_2} \le d_1^2 n(n - d_2) \tilde{\sigma}^2.$$

Altogether we have (4.6) satisfies $\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X] \leq \frac{3}{d_1^2(n-d_2)n\mu} \cdot d_1^2n(n-d_2)\tilde{\sigma}^2 = \frac{3\tilde{\sigma}^2}{\mu}$. (4.2) and (4.3) then follow from Theorem 2.2 by taking $\tau^2 = 3\tilde{\sigma}^2, c = 3a$.

Corollary 4.2. Let X be the biadjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph. Let Q be a real $n \times m$ matrix with all entries in [0, a]. Let $c_0 = \frac{1}{3}(1 - \frac{d_1}{m}), \gamma_0 = \frac{d_1}{m - d_1}$. Then for all $t \geq 0$, we have

(4.10)
$$\mathbb{P}(f_Q(X) - \mu \ge \gamma_0 \mu + t) \le \exp\left(-c_0 \frac{\tilde{\sigma}^2}{a^2} h\left(\frac{at}{\tilde{\sigma}^2}\right)\right).$$

(4.11)
$$\mathbb{P}(|f_Q(X) - \mu| \ge \gamma_0 \mu + t) \le 2 \exp\left(-\frac{c_0 t^2}{2(\tilde{\sigma}^2 + at/3)}\right).$$

Proof. Recall $p=1-\frac{d_1}{m}$, $p'=1-\frac{d_1}{m-d_1}$ from Theorem 4.1. We have $c_0=\frac{p}{3}$ and $\gamma_0=\frac{1}{p}-1$. It is shown in the proof of Proposition 2.3 (c) in [14] that for any $p \in [0,1]$ and $x \ge 0$, $p^{-1}h(px) \ge ph(x)$. Then from (4.2), for all t > 0,

$$(4.12) \mathbb{P}(f_Q(X) - \mu \ge \gamma_0 \mu + t) \le \exp\left(-\frac{\tilde{\sigma}^2}{3pa^2} h\left(\frac{pat}{\tilde{\sigma}^2}\right)\right) \le \exp\left(-c_0 \frac{\tilde{\sigma}^2}{a^2} h\left(\frac{at}{\tilde{\sigma}^2}\right)\right).$$

Therefore (4.10) holds. Note that $\gamma_0 - 1 + p' = \frac{1}{1 - d_1/m} - 1 - \frac{d_1}{m - d_1} = 0$. Then from (4.3),

$$(4.13) \mathbb{P}(f_Q(X) - \mu \le -\gamma_0 \mu - t) \le \exp\left(-\frac{\tilde{\sigma}^2}{3a^2} h\left(\frac{at}{\tilde{\sigma}^2}\right)\right) \le \exp\left(-c_0 \frac{\tilde{\sigma}^2}{a^2} h\left(\frac{at}{\tilde{\sigma}^2}\right)\right).$$

From (4.12) and (4.13), with the inequality $h(x) \ge \frac{x^2}{2(1+x/3)}$ for $x \ge 0$, we obtain (4.11).

5. The Kahn-Szemerédi argument

The Kahn-Szemerédi argument was first introduced in [24] to prove the second eigenvalue of a random d-regular graph is $O(\sqrt{d})$ with high probability. Later on, it has been applied to a wide range of random graph models to provide the upper bound on top eigenvalues (see for example [25, 10, 21, 12, 34, 44, 19, 38, 14, 53, 29, 60]). We will use the concentration inequalities from Section 4 and the Kahn-Szemerédi argument [24] to prove the upper bound on $\sigma_2(X)$, which directly implies the bound for $\lambda(A)$ in Theorem 1.1.

For fixed
$$x \in S^{n-1}, y \in S_0^{m-1}$$
, we have
$$\langle x, Xy \rangle = \sum_{u \in [n], v \in [m]} X_{uv} x_u y_v.$$

Define the set of light and heavy couples as

$$\mathcal{L}(x,y) = \{(u,v) : |x_u y_v| \le \sqrt{d_1}/m\}, \quad \mathcal{H}(x,y) = \{(u,v) : |x_u y_v| > \sqrt{d_1}/m\}.$$

By taking $Q = xy^{\top}$ we can decompose the linear form $f_Q(X)$ as

$$f_{xy^{\top}}(A) = \langle x, Xy \rangle = f_{\mathcal{L}(x,y)}(X) + f_{\mathcal{H}(x,y)}(X),$$

where

(5.2)
$$f_{\mathcal{L}(x,y)}(X) = \sum_{(u,v)\in\mathcal{L}(x,y)} x_u y_v X_{uv}, \quad f_{\mathcal{H}(x,y)}(X) = \sum_{(u,v)\in\mathcal{H}(x,y)} x_u y_v X_{uv}.$$

To apply the concentration inequality, we first estimate the mean of the light part.

Lemma 5.1. For any fixed $x \in S^{n-1}$, $y \in S_0^{m-1}$, we have $|\mathbb{E}f_{\mathcal{L}(x,y)}(X)| \leq \sqrt{d_1}$.

Proof. Since $\mathbb{E}X_{uv} = \frac{d_1}{m}$ for any $u \in [n], v \in [m]$,

$$|\mathbb{E}f_{\mathcal{L}(x,y)}(X)| = \frac{d_1}{m} \left| \sum_{(u,v)\in\mathcal{L}(x,y)} x_u y_v \right| \le \frac{d_1}{m} \sum_{u\in[n]} \left| \sum_{v:(u,v)\in\mathcal{L}(x,y)} x_u y_v \right|.$$

For any $y \in S_0^{m-1}$, we have $\sum_{v \in [m]} y_v = 0$ and it implies for fixed $u \in [n]$,

$$\sum_{v:(u,v)\in\mathcal{L}(x,y)} x_u y_v = -\sum_{v:(u,v)\in\mathcal{H}(x,y)} x_u y_v.$$

Then $|\mathbb{E}f_{\mathcal{L}(x,y)}(X)| \leq \frac{d_1}{m} \sum_{(u,v) \in \mathcal{H}(x,y)} \frac{|x_u y_v|^2}{\sqrt{d_1}/m} \leq \sqrt{d_1}$.

We further split $\mathcal{L}(x,y)$ into two parts as $\mathcal{L}(x,y) = \mathcal{L}_{+}(x,y) \cup \mathcal{L}_{-}(x,y)$ where

$$\mathcal{L}_{+}(x,y) = \{ u \in [n], v \in [m] : 0 \le x_{u}y_{v} \le \sqrt{d_{1}}/m \}, \quad \mathcal{L}_{-}(x,y) = \mathcal{L} \setminus \mathcal{L}_{+}(x,y),$$

$$f_{\mathcal{L}_{+}(x,y)}(X) = \sum_{(u,v) \in \mathcal{L}_{+}(x,y)} x_{u}y_{v}X_{uv}, \quad f_{\mathcal{L}_{-}(x,y)}(X) = \sum_{(u,v) \in \mathcal{L}_{-}(x,y)} x_{u}y_{v}X_{uv}.$$

Lemma 5.2. Let $c_0 = \frac{1}{3}(1 - \frac{d_1}{m}), \gamma_0 = \frac{d_1}{m - d_1}$. For any fixed $(x, y) \in S^{n-1} \times S_0^{m-1}$, and $\beta \ge 2\gamma_0 \sqrt{d_2}$,

$$(5.3) \mathbb{P}\left(|f_{\mathcal{L}(x,y)}(X)| \ge (\beta+1)\sqrt{d_1}\right) \le 4\exp\left(-\frac{3c_0\beta^2m}{24+4\beta}\right).$$

Proof. We have by Cauchy's inequality,

(5.4)
$$\mu := \mathbb{E} f_{\mathcal{L}_{+}(x,y)}(X) \le \frac{d_{1}}{m} \sum_{u \in [n], v \in [m]} |x_{u}y_{v}| \le d_{1} \sqrt{n/m}.$$

And $\tilde{\sigma}^2 := \sum_{(u,v) \in \mathcal{L}_+(x,y)} |x_u y_v|^2 \mathbb{E} X_{uv} \le \frac{d_1}{m} \sum_{u \in [n], v \in [m]} |x_u y_v|^2 = \frac{d_1}{m}$. For any $\beta \ge 2\gamma_0 \sqrt{d_2}$, from (5.4) we have $(\beta/2)\sqrt{d_1} \ge \gamma_0 d_1 \sqrt{n/m} \ge \gamma_0 \mu$. Then by (4.11),

$$\mathbb{P}\left(|f_{\mathcal{L}_{+}(x,y)}(X) - \mathbb{E}f_{\mathcal{L}_{+}(x,y)}(X)| \geq (\beta/2)\sqrt{d_{1}}\right) \\
= \mathbb{P}\left(|f_{\mathcal{L}_{+}(x,y)}(X) - \mathbb{E}f_{\mathcal{L}_{+}(x,y)}(X)| \geq \gamma_{0}\mu + (\beta/2)\sqrt{d_{1}} - \gamma_{0}\mu\right) \\
\leq 2\exp\left(-\frac{c_{0}\left(\frac{\beta}{2}\sqrt{d_{1}} - \gamma_{0}\mu\right)^{2}}{\frac{2d_{1}}{m} + \frac{2}{3}\frac{\sqrt{d_{1}}}{\sqrt{m}}\left(\frac{\beta}{2}\sqrt{d_{1}} - \gamma_{0}\mu\right)}\right) \leq 2\exp\left(-\frac{3c_{0}\beta^{2}m}{24 + 4\beta}\right),$$

where in the last inequality we use the fact that $t \mapsto \frac{t^2}{a+bt}$ with a,b>0 is increasing for $t\geq 0$. The same inequality holds for $\mathcal{L}_{-}(x,y)$. Therefore (5.3) holds.

We now consider the heavy part. We will follow the notations used in Section 6 of [14] and begin with the definition of the discrepancy property for X.

Definition 5.3 (discrepancy property for X). Let X be the biadjacency matrix of a (n, m, d_1, d_2) -bipartite biregular graph. For any $S \subset [n]$, $T \subset [m]$, define $e(S,T) := \sum_{u \in S, v \in T} X_{uv}$. We say A satisfies the discrepancy property with $\delta \in (0,1), \kappa_1 < 1$ and $\kappa_2 \geq 0$, denoted by $\mathrm{DP}(\delta, \kappa_1, \kappa_2)$, if for all non-empty subsets $S \subset [n], T \subset [m]$, at least one of the following holds:

$$\begin{array}{l} (1) \ e(S,T) \leq \kappa_1 \delta |S| |T|, \\ (2) \ e(S,T) \log \left(\frac{e(S,T)}{\delta |S| |T|} \right) \leq \kappa_2 (|S| \vee |T|) \log \left(\frac{em}{|S| \vee |T|} \right). \end{array}$$

Since X is not Hermitian, following the same proof from Lemma 6.4 in [14] with some modification for our model, we obtain the following lemma.

Lemma 5.4. Let X be the biadjacency matrix of a random (n, m, d_1, d_2) -bipartite biregular graph. For any K>0, with probability at least $1-m^{-K}$, the discrepancy property $DP(\delta,\kappa_1,\kappa_2)$ holds for X with $\delta=\frac{d_1}{m}$, $\kappa_1 = e^2 (1 + \gamma_0)^2, \text{ and } \kappa_2 = \frac{2}{c_0} (1 + \gamma_0)(K + 4).$

Proof. Let $Q = \mathbf{1}_S \mathbf{1}_T^{\mathsf{T}}$, where $\mathbf{1}_S \in \{0,1\}^n$ and $\mathbf{1}_T \in \{0,1\}^m$ are the indicator vectors of the set S and T, respectively. We have

$$f_Q(X) = \sum_{u \in [n], v \in [m]} (\mathbf{1}_S)_u (\mathbf{1}_T)_v X_{uv} = e(S, T).$$

Denote $\mu(S,T) := \mathbb{E}e(S,T) = \frac{d_1}{m}|S||T| = \delta|S||T|$. Recall $\gamma_0 = \frac{d_1}{m-d_1}$. For fixed K > 0, let $\gamma_1 = e^2(1+\gamma_0)^2 - 1$ and $\gamma = \gamma(S,T,m) = \max(\gamma^*,\gamma_1)$, where γ^* is the unique x such that

(5.6)
$$c_0 h(x - \gamma_0) \mu(S, T) = (K + 4)(|S| \vee |T|) \log \left(\frac{em}{|S| \vee |T|}\right).$$

Note that we have $\gamma \geq \gamma_1 \geq \gamma_0$. Taking a=1 and $\tilde{\sigma}^2 = \frac{d_1}{m} \sum_{u \in [n], v \in [m]} Q_{uv}^2 = \frac{d_1}{m} |S| |T| = \mu(S,T)$ in (4.10),

$$\mathbb{P}(e(S,T) \ge (1+\gamma)\mu(S,T)) = \mathbb{P}(f_Q(X) \ge (1+\gamma)\mathbb{E}f_Q(X))$$

$$\le \exp\left(-c_0\tilde{\sigma}^2 h\left(\frac{(\gamma-\gamma_0)\mu(S,T)}{\tilde{\sigma}^2}\right)\right) = \exp\left(-c_0\mu(S,T)h\left(\gamma-\gamma_0\right)\right).$$

Then

$$\mathbb{P}(\exists S \subset [n], T \subset [m], |S| = s, |T| = t, e(S, T) \ge (1 + \gamma)\mu(S, T))$$

$$\leq \sum_{S \subset [n], |S| = s} \sum_{T \subset [m], |T| = t} \exp(-c_0 h(\gamma - \gamma_0)\mu(S, T))$$

$$\leq \sum_{S \subset [n], |S| = s} \sum_{T \subset [m], |T| = t} \exp(-c_0 h(\gamma^* - \gamma_0)\mu(S, T))$$

$$= \binom{n}{s} \binom{m}{t} \exp\left(-(K + 4)(s \lor t) \log\left(\frac{em}{s \lor t}\right)\right) \le \left(\frac{ne}{s}\right)^s \left(\frac{me}{t}\right)^t \exp\left(-(K + 4)(s \lor t) \log\left(\frac{em}{s \lor t}\right)\right)$$

$$\leq \exp\left(-(K + 2)(s \lor t) \log\left(\frac{em}{s \lor t}\right)\right) \le \exp(-(K + 2) \log(em)),$$

where in the third line we use the fact that h(x) is increasing for $x \ge 0$, and in the last line we use the fact that $x \mapsto x \log(e/x)$ is increasing on [0, 1]. Taking a union bound over all $s \in [n], t \in [m]$, we have

$$(5.7) \mathbb{P}(\exists S \in [n], T \in [m], e(S, T) \ge (1 + \gamma)\mu(S, T)) \le nm \exp(-(K + 2)\log(em)) \le m^{-K}.$$

If the subsets S, T satisfy $\gamma(S, T, m) = \gamma_1$, then from (5.7) with probability at least $1 - m^{-K}$,

$$e(S,T) \le (1+\gamma_1)\mu(S,T) = e^2(1+\gamma_0)^2\delta|S||T|.$$

Hence Case 1 holds with $\kappa_1 = e^2(1+\gamma_0)^2$. Now assume the subsets S,T satisfy $\gamma(S,T,m) = \gamma^*$. From (5.7), with probability at least $1-m^{-K}$, $e(S,T) \leq (1+\gamma^*)\mu(S,T)$. Therefore from (5.6),

(5.8)
$$\frac{c_0}{1+\gamma^*}h(\gamma^*-\gamma_0)e(S,T) \le (K+4)(|S| \lor |T|)\log\left(\frac{em}{|S| \lor |T|}\right).$$

It is shown in the proof of Lemma 6.4 in [14] that

$$\frac{h(\gamma^* - \gamma_0)}{1 + \gamma_*} \geq \frac{1}{2(1 + \gamma_0)} \log \frac{e(S, T)}{\mu(S, T)} = \frac{1}{2(1 + \gamma_0)} \log \frac{e(S, T)}{\delta |S| |T|}.$$

Together with (5.8), it implies that when $\gamma^* \geq \gamma_1$,

$$e(S,T)\log\frac{e(S,T)}{\delta|S||T|} \le \frac{2}{c_0}(1+\gamma_0)(K+4)(|S|\vee|T|)\log\frac{em}{|S|\vee|T|}$$

Then Case 2 follows with $\kappa_2 = \frac{2}{c_0}(1+\gamma_0)(K+4)$. This completes the proof.

Assuming the discrepancy property holds for X, the following lemma implies that the contribution from heavy tuples is $O(\sqrt{d_1})$. Since the proof is very similar to the proof of Lemma 6.6 in [14], we omit the details.

Lemma 5.5. Let X be the biadjacency matrix of a (n, m, d_1, d_2) -bipartite biregular graph. Suppose X has the discrepancy property $DP(\delta, \kappa_1, \kappa_2)$ with $\delta, \kappa_1, \kappa_2$ given in Lemma 5.4. Then for any $(x, y) \in S^{n-1} \times S_0^{m-1}$, we have $f_{\mathcal{H}(x,y)}(X) \leq \alpha_0 \sqrt{d_1}$ with $\alpha_0 = 48 + 32\kappa_1 + 64\kappa_2 \left(1 + \frac{1}{\kappa_1 \log \kappa_1}\right)$.

Now we finish the proof of Theorem 1.1 with the ε -net argument. The following lemma is standard and we omit the proof.

Lemma 5.6. For $\varepsilon \in (0,1/2)$, let $\mathcal{N}_{\varepsilon}$ be an ε -net of S^{n-1} and $\mathcal{N}_{\varepsilon}^{0}$ be an ε -net of S^{m-1}_{0} . Let X be the biadjacency matrix of a (n, m, d_1, d_2) -bipartite biregular graph. Then

$$\sigma_2(X) = \sup_{x \in S^{n-1}, y \in S_0^{m-1}} \langle x, Xy \rangle \le \frac{1}{1 - 2\varepsilon} \sup_{x \in \mathcal{N}_{\varepsilon}, y \in \mathcal{N}_{\varepsilon}^0} |\langle x, Xy \rangle|.$$

Proof of Theorem 1.1. Fix K > 0. By Lemma 5.4, with probability at least $1 - m^{-K}$, X has $DP(\delta, \kappa_1, \kappa_2)$ property. Let \mathcal{D} be the event that this property holds. Take $\varepsilon = 1/4$ in Lemma 5.6. By taking the union bound over $\mathcal{N}_{\varepsilon} \times \mathcal{N}_{\varepsilon}^{0}$, we have for any $\alpha > 0$,

$$(5.9) \mathbb{P}\left(\mathcal{D}\cap\left\{\sigma_2(X)\geq\alpha\sqrt{d_1}\right\}\right)\leq \sum_{x\in\mathcal{N}_\varepsilon,y\in\mathcal{N}_\varepsilon^0}\mathbb{P}\left(\mathcal{D}\cap\left\{|\langle x,Xy\rangle|\geq\alpha/2\sqrt{d_1}\right\}\right).$$

For any fixed $(x,y) \in S^{n-1} \times S_0^{m-1}$ and any $\alpha > 2\alpha_0$ with α_0 given in Lemma 5.5, from our analysis of the heavy couples, we have

$$\mathbb{P}\left(\mathcal{D}\cap\left\{|\langle x,Xy\rangle|\geq\alpha/2\sqrt{d_1}\right\}\right)\leq\mathbb{P}\left(\mathcal{D}\cap\left\{|f_{\mathcal{L}(x,y)}(X)|\geq\alpha/2\sqrt{d_1}-|f_{\mathcal{H}(x,y)}(X)|\right\}\right)$$
$$\leq\mathbb{P}\left(|f_{\mathcal{L}(x,y)}(X)|\geq(\alpha/2-\alpha_0)\sqrt{d_1}\right).$$

Take $\beta = \alpha/2 - \alpha_0 - 1$. When $\beta > 2\gamma_0\sqrt{d_2}$, from (5.3), we have

$$\mathbb{P}\left(|f_{\mathcal{L}(x,y)}(X)| \ge (\alpha/2 - \alpha_0)\sqrt{d_1}\right) \le 4\exp\left(-\frac{3c_0\beta^2 m}{24 + 4\beta}\right).$$

When $\varepsilon = 1/4$, there exist ε -nets $\mathcal{N}_{\varepsilon}$ and $\mathcal{N}_{\varepsilon}^0$ such that $|\mathcal{N}_{\varepsilon}| \leq 9^n$ and $|\mathcal{N}_{\varepsilon}^0| \leq 9^m$. Then

$$(5.10) \qquad \mathbb{P}\left(\mathcal{D}\cap\left\{\sigma_2(A)\geq\alpha\sqrt{d_1}\right\}\right)\leq 4\cdot 9^{n+m}\exp\left(-\frac{3c_0\beta^2m}{24+4\beta}\right)\leq 9^{2m+1}\exp\left(-\frac{3c_0\beta^2m}{24+4\beta}\right).$$

Then $c_0 \geq \frac{1}{6}, 2\gamma_0\sqrt{d_2} = \frac{2d_2\sqrt{d_2}}{n-d_2} \leq \frac{4d_2\sqrt{d_2}}{n} \leq \sqrt{2}$. Recall $\kappa_1 = e^2(1+\gamma_0)^2, \kappa_2 = \frac{2}{c_0}(1+\gamma_0)(K+4)$. We have $e^2 \leq \kappa_1 \leq 4e^2, \kappa_2 \leq 24(K+4)$. Then $\alpha_0 = 32\kappa_1 + 48 + 64\kappa_2\left(1+\frac{1}{\kappa_1\log\kappa_1}\right)$ is a constant depending on K. Note that $\alpha = 2\beta + 2\alpha_0 + 2$. Take $\beta > \sqrt{2} \geq 2\gamma_0\sqrt{d_2}$ to be a sufficiently large constant independent of m such that $9^{2m+1}\exp\left(-\frac{3c_0\beta^2m}{24+4\beta}\right) \leq e^{-m}$. Then α is a constant depending only on K. From (5.10),

$$\mathbb{P}\left(\sigma_2(X) \geq \alpha \sqrt{d_1}\right) \leq \mathbb{P}(\mathcal{D}^c) + \mathbb{P}\left(\mathcal{D} \cap \left\{\sigma_2(X) \geq \alpha \sqrt{d_1}\right\}\right) \leq m^{-K} + e^{-m}.$$

This completes the proof of Theorem 1.1.

6. Higher order switching

The size-biased coupling can be used to study any random variable that permits a size-biased coupling with good properties, see the general theorem stated in Lemma 2.2, and more applications of size-biased coupling in [14, 3, 27]. From Lemma 2.1, it's easy to construct such sized-biased coupling for linear forms of A, since it is a weighted sum of indicator random variables, and the switching operators were used to create a coupling that has good control of \mathcal{B} and D in the statement of Lemma 2.2. The limitation of this argument is that one cannot get a sharp constant in front of $\sqrt{d_1-1}+\sqrt{d_2-1}$ with the use of an ε -net. This size-biased coupling technique is used to prove concentration of x^TAy for a random matrix A and two fixed vectors x, y. After passing through the ε -net, the constants in front of $\sqrt{d_1-1}+\sqrt{d_2+1}$ will depend on the size of the net. It's possible that one could come up with a better size-biased coupling without taking a union bound over the ε -net to get the sharp constant.

In this section, we modify our switching operations and apply the previous analysis to $M := XX^{\top} - d_1I$. By applying an ε -net argument to M, we won't be able to capture the sharp dependence on d_2 , but it allows us to find a sharp dependence on d_1 .

Now we move on to the proof of Theorem 1.3. For a given $n \times n$ symmetric matrix Q, the linear function $f_Q(M)$ can be written as

$$f_Q(M) = \sum_{u_1, u_2 \in [n]} Q_{u_1 u_2} M_{u_1 u_2} = \sum_{u_1 \neq u_2, v \in [m]} Q_{u_1 u_2} X_{u_1 v} X_{u_2 v}.$$

We see that $f_Q(M)$ is a linear combination of n(n-1)m many indicator random variables $X_{u_1v}X_{u_2v}$. Then to construct a size biased coupling for $f_Q(M)$, according to Lemma 2.1, we need to construct a coupling $(X, X^{(u_1u_2v_1)})$ where $X^{(u_1u_2v_1)}$ is distributed as X conditioned on the event $X_{u_1v_1}X_{u_2v_1} = 1$.

To create a size biased coupling with the desired property, we introduce three types of switchings involving more vertices.

Definition 6.1 (valid Type 1 and Type 2 switchings). Assume $X_{u_1v_1} = X_{u_2v_2} = X_{u_3v_1} = X_{u_4v_3} = 1$, $X_{u_2v_1} = X_{u_3v_3} = X_{u_4v_2} = 0$. We define $(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ to be a valid Type 1 forward switching if after the switching, the edges u_2v_1, u_3v_3, u_4v_2 are added and the edges u_2v_2, u_3v_1, u_4v_3 are removed.

For valid Type 1 switchings, the appearance of the edge u_1v_1 is not changed. Similarly, if $X_{u_1v_2} = X_{u_2v_2} = X_{u_3v_1} = X_{u_4v_3} = 1, X_{u_1v_1} = X_{u_3v_3} = X_{u_4v_2} = 0$. We define $(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ to be a valid Type 2 forward switching. For valid Type 2 switchings, the appearance of the edge u_2v_1 is not changed. Similar to Definition 3.1, the backward switchings are defined accordingly.




Figure 2. valid Type 1 and Type 2 forward switchings for fixed u_1, u_2, v_1

See Figure 2 for an example of Definition 6.1. In addition, we define the following switching as a combination of Type 1 and Type 2 switchings.

Definition 6.2 (valid Type 3 switchings). Assume $X_{u_1v_4} = X_{u_2v_2} = X_{u_3v_1} = X_{u_4v_3} = X_{u_5v_1} = X_{u_6v_5} = 1$ and $X_{u_1v_1} = X_{u_2v_1} = X_{u_3v_3} = X_{u_4v_2} = X_{u_5v_5} = X_{u_6v_4} = 0$. we define $(u_1, u_2, u_3, u_4, u_5, u_6, v_1, v_2, v_3, v_4, v_5)$ to be a valid Type 3 forward switching if after the switching, the edges $u_1v_1, u_2v_1, u_3v_3, u_4v_2, u_5v_5, u_6v_4$ are added and the edges u_2v_1, u_3v_1, u_4v_3 are removed.




FIGURE 3. a valid Type 3 forward switching for fixed u_1, u_2, v_1

See Figure 3 for an example of Definition 6.2. We first estimate the number of all types of switchings for a given bipartite biregular graph. The argument is similar to the proof of Lemma 3.2.

Lemma 6.3. Consider an (n, m, d_1, d_2) -bipartite biregular graph G. Let $s_{u_1u_2v_1}^{(i)}(G), 1 \le i \le 3$ be the number of valid Type i forward switchings of the form

$$(u_1, u_2, \cdot, \cdot, v_1, \cdot, \cdot), \quad (u_1, u_2, \cdot, \cdot, v_1, \cdot, \cdot), \quad (u_1, u_2, \cdot, \cdot, \cdot, \cdot, \cdot, v_1, \cdot, \cdot, \cdot, \cdot),$$

respectively. Let $t_{u_1u_2v_1}^{(i)}(G)$ be the number of the corresponding valid Type i backward switchings for i = 1, 2, 3. Then the following inequalities hold:

(1) If
$$X_{u_1v_1} = 1, X_{u_2v_1} = 0$$
,

(6.1)
$$d_1^2(d_2-1)(n-2d_2) \le s_{u_1u_2v_1}^{(1)} \le d_1^2(d_2-1)(n-d_2).$$

If
$$X_{u_1v_1} = 0, X_{u_2v_1} = 1$$
,

(6.2)
$$d_1^2(d_2-1)(n-2d_2) \le s_{u_1u_2v_1}^{(2)} \le d_1^2(d_2-1)(n-d_2).$$

(2) If
$$X_{u_1v_1} = 0 = X_{u_2v_1} = 0$$
,

$$(6.3) d_1^4 d_2(d_2 - 1)(n - d_2)(n - 3d_2) \le s_{u_1 u_2 v_1}^{(3)} \le d_1^4 d_2(d_2 - 1)(n - d_2)^2.$$

(3) If
$$X_{u_1v_1} = X_{u_2v_1} = 1$$
,

$$(6.4) d_1^2(n-d_2)(n-2d_2) \le t_{u_1u_2v_1}^{(1)} \le d_1^2(n-d_2)^2,$$

(6.5)
$$d_1^2(n-d_2)(n-2d_2) \le t_{u_1u_2v_1}^{(2)} \le d_1^2(n-d_2)^2,$$

and

$$(6.6) d_1^4(n-d_2)^2(n-d_2-1)(n-3d_2) \le t_{u_1u_2v_1}^{(3)} \le d_1^4(n-d_2)^3(n-d_2-1).$$

Proof. For fixed u_1, u_2, v_1 , assume $X_{u_1v_1} = 1, X_{u_2v_1} = 0$. We first consider $s_{u_1u_2v_1}^{(1)}$. We bound the number of all possible Type 1 forward switching by choosing $u_3 \in \mathcal{N}(v_1)$ with $u_3 \neq u_1, v_2 \in \mathcal{N}(u_2), u_4 \in \overline{\mathcal{N}}(v_2)$, and $v_3 \in \mathcal{N}(u_4)$. Here we require $u_3 \neq u_1$ because after the switching, u_3, v_1 are not connected but u_1, v_1 are connected (see Figure 2). There are $d_1^2(d_2 - 1)(n - d_2)$ many choices in total. This gives the upper bound on $s_{u_1u_2v_1}^{(1)}$. For the lower bound, we consider the number of tuples that do not allow a valid Type 1 forward switching among the $d_1^2(d_2 - 1)(n - d_2)$ many choices, and we denote it by K. Then $s_{u_1u_2v_1}^{(1)} = d_1^2(d_2 - 1)(n - d_2) - K$. Any tuple chosen as above is not a valid switching if and only if $v_3 \in \mathcal{N}(u_3)$. We upper bound K by choosing $v_2 \in \mathcal{N}(u_2), u_3 \in \mathcal{N}(v_1)$ with $u_3 \neq u_1, v_3 \in \mathcal{N}(u_3)$ and $u_4 \in \mathcal{N}(v_3)$, making

at most $d_1^2d_2(d_2-1)$ many choices in total. Hence $K \leq d_1^2d_2(d_2-1)$ and $s_{u_1u_2v_1}^{(1)} \geq d_1^2(d_2-1)(n-2d_2)$. The estimate of Type 2 forward switchings follows from the symmetry of u_1 and u_2 . Therefore (6.1) and (6.2) hold

Now assume $X_{u_1v_1}=X_{u_2v_1}=1$. For fixed (u_1,u_2,v_1) , by choosing $u_3\in \overline{\mathcal{N}}(v_1), v_3\in \mathcal{N}(u_3), u_4\in \overline{\mathcal{N}}(v_3)$ and $v_2\in \mathcal{N}(u_4)$, there are at most $d_1^2(n-d_2)^2$ many choices. Among those tuples, the switching will fail if and only if $v_2\in \mathcal{N}(u_2)$. Let $t_{u_1u_2v_1}^{(1)}=d_1^2(n-d_2)^2-L$, where L is the number of invalid Type 1 backward switchings among the $d_1^2(n-d_2)^2$ many choices above. We can then bound L by choosing $u_3\in \overline{\mathcal{N}}(v_1), v_2\in \mathcal{N}(u_2), v_3\in \mathcal{N}(u_3)$ and $u_4\in \mathcal{N}(v_2)$, which has at most $d_1^2d_2(n-d_2)$ many choices. Hence $d_1^2(n-d_2)\leq t_{u_1u_2v_1}^{(1)}\leq d_1^2(n-d_2)^2$. The estimate of $t_{u_1u_2v_1}^{(2)}$ follows in the same way. Then (6.4) and (6.5) hold.

It remains to show (6.3) and (6.6). Assume $X_{u_1v_1} = X_{u_2v_1} = 0$. By choosing $u_3 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_2), u_4 \in \overline{\mathcal{N}}(v_2), v_3 \in \mathcal{N}(u_4), u_5 \in \mathcal{N}(v_1)$ with $u_5 \neq u_3, v_4 \in \mathcal{N}(u_1), u_6 \in \overline{\mathcal{N}}(v_4)$ and $v_5 \in \mathcal{N}(u_6)$, we have a total number of $d_1^2d_2(d_2-1)(n-d_2)^2$ tuples. Therefore $s_{u_1u_2v_1}^{(3)} \leq d_1^4d_2(d_2-1)(n-d_2)^2$. Among those tuples, a tuple is not valid Type 3 forward switching if and only if one of the two cases happens: (i) $v_3 \in \mathcal{N}(u_3)$ or (ii) $v_5 \in \mathcal{N}(u_5)$. Let $s_{u_1u_2v_1}^{(3)} = d_1^4d_2(d_2-1)(n-d_2)^2 - K$. For the first case, by choosing $u_3 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_2), v_3 \in \mathcal{N}(u_3), u_4 \in \mathcal{N}(v_3), u_5 \in \mathcal{N}(v_1)$ with $u_5 \neq u_3, v_4 \in \mathcal{N}(v_1), u_6 \in \overline{\mathcal{N}}(v_4)$ and $v_5 \in \mathcal{N}(u_5)$ there are at most $d_1^4d_2^2(d_2-1)(n-d_2)$ many tuples that are not valid. For the second case, by choosing $u_3 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_2), u_4 \in \overline{\mathcal{N}}(v_2), v_3 \in \mathcal{N}(u_4), u_5 \in \mathcal{N}(v_1)$ with $u_5 \neq u_3, v_4 \in \mathcal{N}(u_1), v_5 \in \mathcal{N}(u_5)$ and $u_6 \in \mathcal{N}(v_5)$ there are at most $d_1^4d_2^2(d_2-1)(n-d_2)$ many tuples that are not valid. From those two cases, we have $K \leq 2d_1^4d_2^2(d_2-1)(n-d_2)$ and $s_{u_1u_2v_1}^{(3)} \geq d_1^4d_2(d_2-1)(n-d_2)(n-3d_2)$. This implies (6.3).

Now assume $X_{u_1v_1}=X_{u_2v_1}=1$. For the upper bound on $t_{u_1u_2v_1}^{(3)}$, we choose $u_3\in \overline{\mathcal{N}}(v_1), v_3\in \mathcal{N}(u_3), u_4\in \overline{\mathcal{N}}(v_3), v_2\in \mathcal{N}(u_4), u_5\in \overline{\mathcal{N}}(v_1)$ with $u_5\neq u_3, v_5\in \mathcal{N}(u_5), u_6\in \overline{\mathcal{N}}(v_5)$ and $v_4\in \mathcal{N}(u_6)$. This gives a total number of $d_1^4(n-d_2)^3(n-d_2-1)$ choices. Among those choices, a tuple is not a valid Type 3 backward switching if and only if (i) $v_2\in \mathcal{N}(u_2)$ or (ii) $u_4\in \mathcal{N}(u_1)$. Let $t_{u_1u_2v_1}^{(3)}=d_1^4(n-d_2)^3(n-d_2-1)-L$. For the first case, we have at most $d_1^4d_2(n-d_2)^2(n-d_2-1)$ choices by choosing $u_3\in \overline{\mathcal{N}}(v_1), v_2\in \mathcal{N}(u_2), v_3\in \mathcal{N}(u_3), u_4\in \mathcal{N}(v_2), u_5\in \overline{\mathcal{N}}(v_1), u_5\neq u_3, v_5\in \mathcal{N}(u_5), u_6\in \overline{\mathcal{N}}(v_5)$ and $v_4\in \mathcal{N}(u_6)$. By a similar argument, there are at most $d_1^4d_2(n-d_2)^2(n-d_2-1)$ choices for the second case. Hence $L\leq 2d_1^4d_2(n-d_2)^2(n-d_2-1)$ and $t_{u_1u_2v_1}^{(3)}\geq d_1^4(n-d_2)^2(n-d_2-1)(n-3d_2)$. Therefore (6.6) holds.

Let \mathcal{G} be the collection of the biadjacency matrices of all (n, m, d_1, d_2) -bipartite biregular graphs and $\mathcal{G}_{u_1u_2v_1}$ be the subset of \mathcal{G} such that $X_{u_1v_1} = X_{u_2v_1} = 1$. Fix $u_1, u_2 \in [n]$ with $u_1 \neq u_2$ and $v_1 \in [m]$. We construct an edge-weighted bipartite graph \mathfrak{G}_0 on two vertex class \mathcal{G} and $\mathcal{G}_{u_1u_2v_1}$ as follows:

- If $X \in \mathcal{G}$ with $X_{u_1v_1} = 0, X_{u_2v_1} = 1$, then form an edge of weight $d_1^2d_2(n d_2)$ between X and every element of $\mathcal{G}_{u_1u_2v_1}$ that is a result of a valid Type 1 forward switching from X.
- If $X \in \mathcal{G}$ with $X_{u_1v_1} = 1, X_{u_2v_1} = 0$, then form an edge of weight $d_1^2d_2(n d_2)$ between X and every element of $\mathcal{G}_{u_1u_2v_1}$ that is a result of a valid Type 2 forward switching from X.
- If $X \in \mathcal{G}$ with $X_{u_1v_1} = X_{u_2v_1} = 0$, then form an edge of weight 1 between X and every element of $\mathcal{G}_{u_1u_2v_1}$ that is a result of a valid Type 3 forward switching from X.
- If $X \in \mathcal{G}$ with $X_{u_1v_1} = X_{u_2v_1} = 1$, then form an edge of weight $d_1^4d_2(d_2 1)(n d_2)^2$ between X and its identical copy in $G_{u_1u_2v_1}$.

By our construction of \mathfrak{G}_0 , the following lemma holds.

Lemma 6.4. In \mathfrak{G}_0 , the following holds:

- (1) Every element in \mathcal{G} has degree between $d_1^4d_2(d_2-1)(n-d_2)(n-3d_2)$ and $d_1^4d_2(d_2-1)(n-d_2)^2$.
- (2) Every element in $\mathcal{G}_{u_1u_2v_1}$ has degree between $d_1^4(n-d_2)^2(n-2d_2)(n-1)$ and $d_1^4(n-d_2)^2n(n-1)$.
- (3) \mathfrak{G}_0 can be embedded into a weighted bipartite biregular graph \mathfrak{G} on the same vertex sets, with vertices in \mathcal{G} having degree $d_1^4d_2(d_2-1)(n-d_2)^2$ and vertices in $\mathcal{G}_{u_1u_2v_1}$ having degree $d_1^4(n-d_2)^2n(n-1)$.

Proof. From Lemma 6.3, with the weight we assign to each edge, we have the following:

• If $X_{u_1,v_1} = 0, X_{u_2,v_2} = 1$ or $X_{u_1,v_1} = 1, X_{u_2,v_2} = 0$, then the degree of X in \mathfrak{G}_0 is between $d_1^4 d_2 (d_2 - 1)(n - d_2)(n - 2d_2)$ and $d_1^4 d_2 (d_2 - 1)(n - d_2)^2$.

- If $X_{u_1,v_1} = X_{u_2,v_2} = 0$, from (6.3), the degree of X in \mathfrak{G}_0 is between $d_1^4 d_2 (d_2 1)(n d_2)(n 3d_2)$ and $d_1^4 d_2 (d_2 1)(n d_2)^2$.
- If $X_{u_1,v_1} = X_{u_2,v_2} = 1$, then the degree of X in \mathfrak{G}_0 is exactly $d_1^4 d_2 (d_2 1)(n d_2)^2$.

Therefore the first claim of Lemma 6.4 holds. Now we turn to elements in $G_{u_1u_2v_1}$. For any $X \in \mathcal{G}_{u_1u_2v_1}$, X can be adjacent to elements X' such that

- X' is a result of a valid Type 1 or Type 2 backward switching from X, with edge weight $d_1^2d_2(n-d_2)$.
- X' is a result of a valid Type 3 backward switching from X, with edge weight 1.
- X' is an identical copy of X in \mathcal{G} , with edge weight $d_1^4 d_2 (d_2 1)(n d_2)^2$.

Then for each $X \in \mathcal{G}_{u_1u_2v_1}$, combining all the weighted edges from valid Type 1,2,3 backward switchings and the weighted edge from its identical copy, from (6.4), (6.5) and (6.6), the degree is at most

$$2d_1^2d_2(n-d_2) \cdot d_1^2(n-d_2)^2 + d_1^4(n-d_2)^3(n-d_2-1) + d_1^4d_2(d_2-1)(n-d_2)^2$$

= $d_1^4(n-d_2)^2n(n-1)$.

And similarly, its degree is at least $d_1^4(n-d_2)^2(n-2d_2)(n-1)$.

Then the second claim in Lemma 6.4 holds. It remains to prove the third claim. Since X is uniformly distributed, we have

$$\mathbb{P}(X_{u_1v_1} = X_{u_2v_1} = 1) = \frac{nd_1(d_2 - 1)}{n(n-1)m} = \frac{d_2(d_2 - 1)}{n(n-1)}$$

and

(6.7)
$$\frac{|\mathcal{G}_{u_1 u_2 v_1}|}{|\mathcal{G}|} = \frac{d_2(d_2 - 1)}{n(n - 1)}.$$

To construct \mathfrak{G} , we start with \mathfrak{G}_0 and add edges as follows. Go through the vertices in \mathcal{G} and for each vertex with degree less than $d_1^4d_2(d_2-1)(n-d_2)^2$, arbitrarily make edges from the vertex to vertices in $\mathcal{G}_{u_1,u_2,v_1}$ with degree less than $d_1^4(n-d_2)^2n(n-1)$. Continue this procedure until either all vertices in \mathcal{G} have degree $d_1^4d_2(d_2-1)(n-d_2)^2$ or all vertices in $\mathcal{G}_{u_1,u_2,v_1}$ have degree $d_1^4(n-d_2)^2n(n-1)$. Similar to the proof of Lemma 3.3, \mathfrak{G} is bipartite biregular. Therefore we have embedded \mathfrak{G}_0 into \mathfrak{G} .

Now we are able to construct a coupling with the desired property. In the graph \mathfrak{G} , Choose a uniform element X in \mathcal{G} and consider $X^{(u_1u_2v_1)}$ to be the element in $\mathcal{G}_{u_1u_2v_1}$ given by walking from X along an edge with probability proportional to its weight. Then $X^{(u_1u_2v_1)}$ is uniformly distributed in the vertex set $\mathcal{G}_{u_1u_2v_1}$. Lemma 6.4 yields a coupling of $(X, X^{(u_1u_2v_1)})$ that satisfies

(6.8)
$$\mathbb{P}(X, X^{(u_1 u_2 v_1)} \text{ are identical or differ by a switching } | X^{(u_1 u_2 v_1)}) \ge 1 - \frac{2d_2}{n},$$

(6.9)
$$\mathbb{P}(X, X^{(u_1 u_2 v_1)} \text{ are identical or differ by a switching } | X) \ge 1 - \frac{2d_2}{n - d_2}.$$

7. Concentration for linear functions of $XX^{\top} - d_1I$

For a given $n \times n$ symmetric matrix Q, we define the following linear function for X:

$$(7.1) g_Q(X) := \sum_{u_1, u_2 \in [n]} Q_{u_1 u_2} (XX^{\top} - d_1 I)_{u_1 u_2} = \sum_{u_1, u_2 \in [n], u_1 \neq u_2} \sum_{v_1 \in [m]} Q_{u_1 u_2} (X_{u_1 v_1} X_{u_2 v_1}).$$

In this section, we analyze $g_Q(X)$ by the size biased coupling and the switching operations we introduced in Section 6. To quantify the change of $g_Q(X)$ after a valid forward switching, we first introduce the notion of codegrees.

Definition 7.1 (codgeree). Let G be a bipartite graph with vertex sets $V_1 = [n], V_2 = [m]$ and biadjacency matrix X. Define the *codegree* of two vertices $i, j \in [n]$ in G as

$$\operatorname{codeg}(X, i, j) = |\{v \in [m] : (i, v) \in E(G) \text{ and } (j, v) \in E(G)\}|.$$

Equivalently, we have $\operatorname{codeg}(X, i, j) = \sum_{v \in [m]} X_{iv} X_{jv}$.

From Definition 7.1, Equation (7.1) can also be written as

(7.2)
$$g_Q(X) = \sum_{u_1, u_2 \in [n], u_1 \neq u_2} Q_{u_1 u_2} \cdot \operatorname{codeg}(X, u_1, u_2).$$

For any $u_1, u_2 \in [n], u_1 \neq u_2, v_1 \in [m]$, since X is uniformly distributed, we have $\mathbb{E}X_{u_1v_1}X_{u_2v_1} = \frac{d_1(d_2-1)}{n-1}$. Denote

(7.3)
$$\mu := \mathbb{E}g_Q(X) = \frac{d_1(d_2 - 1)}{n - 1} \sum_{u_1 \neq u_2} Q_{u_1 u_2}, \quad \tilde{\sigma}^2 := \frac{d_1(d_2 - 1)}{n - 1} \sum_{u_1 \neq u_2} Q_{u_1 u_2}^2.$$

Theorem 7.2. Let X be the biadjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph and $M = XX^{\top} - d_1I$. Let Q be an $n \times n$ symmetric matrix with all entries in [0, a] and $Q_{ii} = 0$ for $i \in [n]$. Let $\tilde{\sigma}^2$, μ be the parameters defined in (7.3). Denote $p = 1 - \frac{2d_2}{n}$, $p' = 1 - \frac{2d_2}{n-d_2}$. Then for all $t \geq 0$,

(7.4)
$$\mathbb{P}\left(g_Q(X) - \frac{\mu}{p} \ge t\right) \le \exp\left(-\frac{\tilde{\sigma}^2}{6d_2pa^2}h\left(\frac{pat}{2\tilde{\sigma}^2}\right)\right),$$

and

(7.5)
$$\mathbb{P}\left(g_Q(X) - p'\mu \le -t\right) \le \exp\left(-\frac{\tilde{\sigma}^2}{6d_2a^2}h\left(\frac{at}{2\tilde{\sigma}^2}\right)\right).$$

Proof. We now construct a size biased coupling for $g_Q(X)$ based the analysis of switchings in Section 6. Choose $X \in \mathcal{G}$ uniformly and walk through an edge with probability proportional to its weight. We then obtain a uniform random element $X^{(u_1u_2v_1)}$ in $\mathcal{G}_{u_1u_2v_1}$. The matrix $X^{(u_1u_2v_1)}$ is distributed as X conditioned on $X_{u_1v_1} = X_{u_2v_1} = 1$. Independently of X, we choose $(U_1, U_2) = (u_1, u_2)$ with probability

(7.6)
$$\mathbb{P}(U_1 = u_1, U_2 = u_2) = \frac{Q_{u_1 u_2}}{\sum_{u \neq v} Q_{uv}}$$

for all $u_1 \neq u_2$, and independently of everything choose $V_1 \in [m]$ uniformly. Set $X' = X^{(U_1,U_2,V_1)}$. By Lemma 2.1, and (7.1), the pair $(g_Q(X), g_Q(X'))$ is a size biased coupling. Let $\mathcal{S}_i(u_1, u_2, v_1)$ be the set of all tuples $(u_2, u_3, u_4, v_2, v_3)$ such that $(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ is a valid Type i forward switching for i = 1, 2 and let $\mathcal{S}_3(u_1, u_2, v_1)$ be the set of all tuples $(u_2, \dots, u_6, v_2, \dots, v_5)$ such that $(u_1, \dots, u_6, v_1, \dots, v_5)$ is a valid Type 3 forward switching. Let $X^{(i)}(u_1, \dots, u_4, v_1, v_2, v_3)$ be the matrix obtained from X by a valid forward Type i switching $(u_1, \dots, u_4, v_1, v_2, v_3)$ for i = 1, 2. Similarly, let $X^{(3)}(u_1, \dots, u_6, v_1, \dots, v_5)$ be the matrix obtained from X by a valid forward Type 3 switching $(u_1, \dots, u_6, v_1, \dots, v_5)$. From Lemma 6.4, the coupling of (X, X') can be described as follows. Assuming the product $X_{U_1V_1}X_{U_2V_1} = 0$ and conditioned on X, U_1, U_2, V_1 , there is exactly one non-empty set among $\{\mathcal{S}_i(U_1, U_2, V_1)\}_{1 \leq i \leq 3}$. The matrix X' takes the value $X^{(i)}(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ with probability

$$\frac{d_1^2d_2(n-d_2)}{d_1^4d_2(d_2-1)(n-d_2)^2} = \frac{1}{d_1^2(d_2-1)(n-d_2)}$$

for each $(u_1, \ldots, u_4, v_1, v_2, v_3) \in S_i(U_1, U_2, V_1)$ and i = 1, 2. And it takes the value $X^{(3)}(u_1, \ldots, u_6, v_1, \ldots, v_5)$ with probability $\frac{1}{d_1^3 d_2(d_2-1)(n-d_2)^2}$ for each $(u_1, \ldots, u_6, v_1, \ldots, v_5) \in S_3(u_1, u_2, v_1)$. For any valid Type 1 forward switching $(u_1, \ldots, u_4, v_1, v_2, v_3)$, denote $\tilde{X} := X^{(1)}(u_1, \ldots, u_4, v_1, v_2, v_3)$. From (7.2), we have

$$g_Q(X^{(1)}(u_1, \dots, u_4, v_1, v_2, v_3)) - g_Q(X) = \sum_{i \neq j, i, j \in [n]} Q_{ij} \left(\operatorname{codeg}(\tilde{X}, i, j) - \operatorname{codeg}(X, i, j) \right)$$

(7.7)
$$\leq \sum_{i \neq j, i, j \in [n]} Q_{ij} \left(\operatorname{codeg}(\tilde{X}, i, j) - \operatorname{codeg}(X, i, j) \right)_{+}.$$

Then to have an upper bound on $g_Q(\tilde{X}) - g_Q(X)$, it suffices to count the number of pairs (i, j) with increased codegrees after applying the switching. From Figure 2, the only new edges created in the valid Type 1 forward switching $(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ are u_2v_1, u_3v_3 and u_4v_2 . When the edge u_2v_1 is added, for any $u \in \mathcal{N}(v_1)$, the codegree of u and u_2 is increased by 1. Similarly, when the edge u_3v_3 is added, for any $u \in \mathcal{N}(v_2)$, the

codegree of u and u_4 is increased by 1. When the edge u_4v_2 is added, for any $u \in \mathcal{N}(v_3)$, the codegree of u and u_3 is increased by 1. Therefore (7.7) is bounded by

(7.8)
$$2\sum_{u\in\mathcal{N}(v_1)}Q_{uu_2} + 2\sum_{u\in\mathcal{N}(v_2)}Q_{uu_4} + 2\sum_{u\in\mathcal{N}(v_3)}Q_{uu_3} \le 6d_2a,$$

where the factor 2 comes from the symmetry of indices i, j in (7.7). Using the same argument, for any valid Type 2 forward switching $(u_1, \ldots, u_4, v_1, v_2, v_3)$,

$$(7.9) \quad g_Q(X^{(2)}(u_1, \dots, u_4, v_1, v_2, v_3)) - g_Q(X) \le 2 \sum_{u \in \mathcal{N}(v_1)} Q_{uu_1} + 2 \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4} + 2 \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3} \le 6d_2 a.$$

As can be seen from Figure 3, there are 6 new edges created in a valid Type 3 forward switching. We also have for any valid Type 3 forward switching $(u_1, \ldots, u_6, v_1, \ldots, v_5)$,

$$g_{Q}(X^{(3)}(u_{1},...,u_{6},v_{1},...,v_{5})) - g_{Q}(X)$$

$$\leq 2 \left(\sum_{u \in \mathcal{N}(v_{1})} Q_{uu_{1}} + \sum_{u \in \mathcal{N}(v_{1})} Q_{uu_{2}} + \sum_{u \in \mathcal{N}(v_{2})} Q_{uu_{4}} + \sum_{u \in \mathcal{N}(v_{3})} Q_{uu_{3}} + \sum_{u \in \mathcal{N}(v_{4})} Q_{uu_{6}} + \sum_{u \in \mathcal{N}(v_{5})} Q_{uu_{5}} \right)$$

$$(7.10) \leq 12d_{2}a.$$

If $X_{u_1v_1}=1, X_{u_2v_1}=0$, let $\overline{\mathcal{S}_1}(u_1, u_2, v_1)$ be the set of tuples (u_3, u_4, v_2, v_3) such that $v_2 \in \mathcal{N}(u_2), u_3 \in \mathcal{N}(v_1)$ with $u_3 \neq u_1, u_4 \in \overline{\mathcal{N}}(v_2)$, and $v_3 \in \mathcal{N}(u_4)$. From the proof of Lemma 6.3, there are $d_1^2(d_2-1)(n-d_2)$ many choices in total, hence $|\overline{\mathcal{S}_1}(u_1, u_2, v_1)| = d_1^2(d_2-1)(n-d_2)$ and we have $\mathcal{S}_1(u_1, u_2, v_1) \subset \overline{\mathcal{S}_1}(u_1, u_2, v_1)$. If the condition $X_{u_1v_1}=1, X_{u_2v_1}=0$ does not hold, set $\overline{\mathcal{S}_1}(u_1, u_2, v_1)=\emptyset$.

If the condition $X_{u_1v_1}=1, X_{u_2v_1}=0$ does not hold, set $\overline{\mathcal{S}_1}(u_1,u_2,v_1)=\emptyset$. If $X_{u_1v_1}=0, X_{u_2v_1}=1$, define $\overline{\mathcal{S}_2}(u_1,u_2,v_1)$ to be the set of tuples (u_3,u_4,v_2,v_3) such that $v_2\in\mathcal{N}(u_1)$, $u_3\in\mathcal{N}(v_1)$ with $u_3\neq u_1, u_4\in\overline{\mathcal{N}}(v_2)$, and $v_3\in\mathcal{N}(u_4)$. We have $|\overline{\mathcal{S}_2}(u_1,u_2,v_1)|=d_1^2(d_2-1)(n-d_2)$ and $\mathcal{S}_2(u_1,u_2,v_1)\subset\overline{\mathcal{S}_2}(u_1,u_2,v_1)$. If the condition does not hold, set $\overline{\mathcal{S}_2}(u_1,u_2,v_1)=\emptyset$.

If $X_{u_1v_1} = X_{u_2v_1} = 0$, let $\overline{S_3}(u_1, u_2, v_1)$ be the set of tuples $(u_3, u_4, u_5, u_6, v_2, v_3, v_4)$ such that $v_2 \in \mathcal{N}(u_2)$, $u_3 \in \mathcal{N}(v_1)$ with $u_3 \neq u_1$, $u_4 \in \overline{\mathcal{N}}(v_2)$, $v_3 \in \mathcal{N}(u_4)$, $u_6 \in \overline{\mathcal{N}}(v_4)$, $v_5 \in \mathcal{N}(u_6)$. It satisfies $|\overline{S_3}(u_1, u_2, v_1)| = d_1^4 d_2 (d_2 - 1)(n - d_2)^2$ and $S_3(u_1, u_2, v_1) \subset \overline{S_3}(u_1, u_2, v_1)$. If the condition does not hold, set $\overline{S_3}(u_1, u_2, v_1) = \emptyset$. Denote $D := (g_Q(X') - g_Q(X))_+$. We can write $\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X, U_1, U_2, V_1]$ as

$$\frac{1}{d_1^2(d_2-1)(n-d_2)} \sum_{(u_3,u_4,v_2,v_3)\in\mathcal{S}_1(u_1,u_2,v_1)} (g_Q(X^{(1)}(U_1,U_2,\ldots,V_1,v_2,v_3)) - g_Q(X))_+ \\
+ \frac{1}{d_1^2(d_2-1)(n-d_2)} \sum_{(u_3,u_4,v_2,v_3)\in\mathcal{S}_2(u_1,u_2,v_1)} (g_Q(X^{(2)}(U_1,U_2,\ldots,V_1,v_2,v_3)) - g_Q(X))_+ \\
(7.11) + \frac{1}{d_1^4d_2(d_2-1)(n-d_2)^2} \sum_{(u_3,\ldots,u_6,v_2,v_3,v_4)\in\mathcal{S}_3(u_1,u_2,v_1)} (g_Q(X^{(3)}(U_1,U_2,\ldots,V_1,\ldots,V_5)) - g_Q(X))_+.$$

Inequalities (7.8), (7.9) and (7.10) imply that (7.11) is bounded by

$$\begin{split} &\frac{2}{d_1^2(d_2-1)(n-d_2)} \sum_{(u_3,u_4,v_2,v_3) \in \overline{\mathcal{S}_1}(U_1,U_2,V_1)} \left(\sum_{u \in \mathcal{N}(V_1)} Q_{uU_2} + \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3} \right) \\ &+ \frac{2}{d_1^2(d_2-1)(n-d_2)} \sum_{(u_3,u_4,v_2,v_3) \in \overline{\mathcal{S}_2}(U_1,U_2,V_1)} \left(\sum_{u \in \mathcal{N}(V_1)} Q_{uU_1} + \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3} \right) \\ &+ \frac{2}{d_1^4 d_2(d_2-1)(n-d_2)^2} \sum_{(u_3,\dots,u_6,v_2,v_3,v_4) \in \overline{\mathcal{S}_3}(U_1,U_2,V_1)} \\ &\left(\sum_{u \in \mathcal{N}(V_1)} Q_{uU_1} + \sum_{u \in \mathcal{N}(V_1)} Q_{uU_2} + \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3} + \sum_{u \in \mathcal{N}(v_4)} Q_{uu_6} + \sum_{u \in \mathcal{N}(v_5)} Q_{uu_5} \right). \end{split}$$

Recall the distribution of U_1, U_2, V_1 in (7.6), and $\mu = \frac{d_1(d_2-1)}{n-1} \sum_{u_1 \neq u_2} Q_{u_1u_2}$. Taking expectation over $U_1, U_2, V_1, \mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X]$ is then upper bounded by the sum of the following three terms:

$$\frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} \sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1) \\ u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{\substack{u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}_1}(u_1, u_2, v_1)} Q_{u_1 u_2} \left(\sum_{u \in \mathcal{N}(v_1)} Q_{u u_2} + \sum_{u \in \mathcal{N}(v_2)} Q_{u u_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{u u_3} \right),$$

(7.13)

$$\frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} \sum_{\substack{u_1 \in [n] \\ v_1 \in \overline{\mathcal{N}}(u_1) \\ u_2 \in \mathcal{N}(v_1)}} \sum_{\substack{u_1 \in [n] \\ (u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}_2}(u_1, u_2, v_1)}} Q_{u_1 u_2} \left(\sum_{u \in \mathcal{N}(v_1)} Q_{u u_1} + \sum_{u \in \mathcal{N}(v_2)} Q_{u u_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{u u_3} \right),$$

and

(7.14)

$$\frac{2}{\mu d_1^4 n(n-1)(n-d_2)^2} \sum_{u_1 \in [n], v_1 \in \overline{\mathcal{N}}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)} \sum_{(u_3, \dots, v_4) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)} Q_{uu_2} \left(\sum_{u \in \mathcal{N}(v_1)} Q_{uu_1} + \sum_{u \in \mathcal{N}(v_1)} Q_{uu_2} + \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3} + \sum_{u \in \mathcal{N}(v_4)} Q_{uu_6} + \sum_{u \in \mathcal{N}(v_5)} Q_{uu_5} \right) \\
:= S_{31} + S_{32} + S_{33} + S_{34} + S_{35} + S_{36}.$$

In the following proof, we estimate the three terms (7.12), (7.13) and (7.14) separately. Write the sum in (7.12) as

(7.15)

$$\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{\substack{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1) \\ := S_{11} + S_{12} + S_{13}.}} \left(\sum_{u \in \mathcal{N}(v_1)} Q_{u_1 u_2} Q_{u u_2} + \sum_{u \in \mathcal{N}(v_2)} Q_{u_1 u_2} Q_{u u_4} + \sum_{u \in \mathcal{N}(v_3)} Q_{u_1 u_2} Q_{u u_3} \right)$$

By Cauchy's inequality, S_{11} is bounded by

(7.16)

$$\left(\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1) \\ u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{u_1 u_2}^2\right)^{1/2} \left(\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1) \\ u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{uu_2}^2\right)^{1/2}.$$

Since $|\overline{S_1}(u_1, u_2, v_1)| = d_1^2(d_2 - 1)(n - d_2)$ and $\tilde{\sigma}^2 = \frac{d_1(d_2 - 1)}{n - 1} \sum_{u_1 \neq u_2} Q_{u_1 u_2}^2$, the first factor in the product of (7.16) satisfies

(7.17)

$$\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{\substack{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1) \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{\substack{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1) \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} Q_{u_1 u_2}^2 = d_1^2 d_2 (d_2 - 1)(n - d_2) \sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} Q_{u_1 u_2}^2.$$

Since

$$\sum_{u_1 \in [n], v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)} Q_{u_1 u_2}^2 \leq \sum_{u_1, u_2 \in [n], v_1 \in \mathcal{N}(u_1)} Q_{u_1 u_2}^2 = d_1 \sum_{u_1 \neq u_2} Q_{u_1 u_2}^2,$$

(7.17) is then bounded by

(7.18)
$$d_1^3 d_2(d_2 - 1)(n - d_2) \sum_{u_1 \neq u_2} Q_{u_1 u_2}^2 = d_1^2 d_2(n - 1)(n - d_2) \tilde{\sigma}^2.$$

We also have

$$\sum_{u_1 \in [n], v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{uu_2}^2 \le \sum_{u_2 \in [n]} \sum_{u_1 \in [n], v_1 \in \mathcal{N}(u_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{uu_2}^2 \le d_1 d_2 \sum_{u \ne u_2} Q_{uu_2}^2 = \frac{d_2 (n-1)\tilde{\sigma}^2}{d_2 - 1},$$

where the second inequality comes from the fact that each $u \in [n]$ is counted d_1d_2 times in the sum. Then the second factor in (7.16) is bounded by

(7.19)
$$d_1^2(d_2-1)(n-d_2) \cdot \frac{d_2(n-1)\tilde{\sigma}^2}{d_2-1} = d_1^2 d_2(n-1)(n-d_2)\tilde{\sigma}^2.$$

From (7.16), (7.18) and (7.19), we have

$$(7.20) \frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} S_{11} \le \frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} \cdot d_1^2 d_2(n-1)(n-d_2) \tilde{\sigma}^2 = \frac{2\tilde{\sigma}^2}{\mu} \cdot \frac{d_2^2}{n}.$$

Similarly, by Cauchy's inequality,

$$S_{12} \leq [d_1^2 d_2 (n-1)(n-d_2)\tilde{\sigma}^2]^{1/2} \left(\sum_{u_1 \in [n], v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)} \sum_{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4}^2 \right)^{1/2}.$$

For a given (v_2, u_4) , by taking $v_3 \in \mathcal{N}(u_4), u_2 \in \mathcal{N}(v_2), v_1 \in \overline{\mathcal{N}}(u_2), u_1 \in \mathcal{N}(v_1), u_3 \in \mathcal{N}(v_1), u_3 \neq u_1$, there are at most $(m - d_1)d_1d_2^2(d_2 - 1)$ many tuples $(u_1, u_2, u_3, u_4, v_1, v_3)$ such that $(u_1, u_2, u_3, u_4, v_1, v_2, v_3) \in \overline{\mathcal{S}_1}(u_1, u_2, v_1)$. Hence

$$\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_2)} Q_{uu_4}^2 \leq (m - d_1) d_1 d_2^2(d_2 - 1) \sum_{v_2 \in [m], u_4 \in [n], u \in \mathcal{N}(v_2)} Q_{uu_4}^2$$

$$\leq (m-d_1)d_1^2d_2^2(d_2-1)\sum_{u\neq u_4}Q_{uu_4}^2 = (m-d_1)(n-1)d_1d_2^2\tilde{\sigma}^2 = d_1^2d_2(n-1)(n-d_2)\tilde{\sigma}^2.$$

Then

(7.21)
$$\frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} S_{12} \le \frac{2\tilde{\sigma}^2}{\mu} \cdot \frac{d_2^2}{n}$$

For the third term in (7.15), we have

$$S_{13} \leq [d_1^2(d_2 - 1)(n - 1)(n - d_2)\tilde{\sigma}^2]^{1/2} \left(\sum_{\substack{u_1 \in [n] \\ v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)}} \sum_{(u_3, u_4, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3}^2 \right)^{1/2}.$$

For fixed v_3, u_3 , by choosing $v_1 \in \mathcal{N}(u_3), u_1 \in \mathcal{N}(v_1), u_1 \neq u_3, u_2 \in \overline{\mathcal{N}}(v_1), v_2 \in \mathcal{N}(u_2), u_4 \in \mathcal{N}(u_3)$, there are at most $d_1^2 d_2(d_2 - 1)(n - d_2)$ many tuples $(u_1, u_2, u_3, u_4, v_1, v_2, v_3)$ such that $(u_1, u_2, u_3, u_4, v_1, v_2, v_3) \in \overline{\mathcal{S}}_1(u_1, u_2, v_1)$. Therefore by the same argument, we have

$$\sum_{u_1 \in [n], v_1 \in \mathcal{N}(u_1), u_2 \in \overline{\mathcal{N}}(v_1)} \sum_{(u_3, u_4, v_2, v_3) \in \overline{S}_1(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_3)} Q_{uu_3}^2 \le d_1^2 d_2 (n - d_2) (n - 1) \tilde{\sigma}^2,$$

and

(7.22)
$$\frac{2d_2}{\mu d_1^2 n(n-1)(n-d_2)} S_{13} \le \frac{2\tilde{\sigma}^2}{\mu} \cdot \frac{d_2^2}{n}.$$

From (7.20), (7.21) and (7.22), the term (7.12) is bounded by $\frac{6\tilde{\sigma}_2^2d_2^2}{\mu n}$. The bound on (7.12) also holds for (7.13) by the symmetric role of the two vertices u_1, u_2 .

Now it remains to estimate (7.14). Recall $|\overline{S}_3(u_1, u_2, v_1)| = d_1^4 d_2 (d_2 - 1)(n - d_2)^2$. By Cauchy's inequality, (7.23)

$$S_{31} \leq \left(\sum_{\substack{u_1 \neq u_2 \in [n] \\ v_1 \in [m]}} \sum_{(u_3, \dots, v_4) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{u_1 u_2}^2\right)^{1/2} \left(\sum_{\substack{u_1 \neq u_2 \in [n] \\ v_1 \in [m]}} \sum_{(u_3, \dots, v_4) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{u u_1}^2\right)^{1/2}$$

$$= \left[d_1^4 d_2 (n - d_2)^2 n (n - 1) \tilde{\sigma}^2\right]^{1/2} \left(\sum_{\substack{u_1 \neq u_2 \in [n], v_1 \in [m] \\ u_1 \neq u_2 \in [n], v_1 \in [m]}} \sum_{(u_3, \dots, v_4) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{u u_1}^2\right)^{1/2}.$$

For fixed v_1, u_1 , by taking $u_2 \in \overline{\mathcal{N}}(v_1), u_3 \in \mathcal{N}(v_1), v_2 \in \mathcal{N}(u_2), u_4 \in \overline{\mathcal{N}}(v_2), v_3 \in \mathcal{N}(u_4), u_5 \in \mathcal{N}(v_1), u_5 \neq u_3, v_4 \in \mathcal{N}(u_1), u_6 \in \overline{\mathcal{N}}(v_4)$ and $v_5 \in \mathcal{N}(u_6)$, there are at least $d_1^4 d_2 (d_2 - 1)(n - d_2)^3$ many choices of $(u_2, u_3, u_4, u_5, u_6, v_2, v_3, v_4, v_5)$ such that $(u_3, u_4, u_5, u_6, v_2, v_3, v_4, v_5) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)$. Then

$$\sum_{u_1 \neq u_2 \in [n], v_1 \in [m]} \sum_{(u_3, \dots, v_4) \in \overline{\mathcal{S}_3}(u_1, u_2, v_1)} \sum_{u \in \mathcal{N}(v_1)} Q_{uu_1}^2 \\
\leq d_1^4 d_2 (d_2 - 1)(n - d_2)^3 \sum_{u_1 \in [n], v_1 \in [m]} \sum_{u \in \mathcal{N}(v_1)} Q_{uu_1}^2 \\
= d_1^5 d_2 (d_2 - 1)(n - d_2)^3 \sum_{u \neq u_1} Q_{uu_1}^2 = d_1^4 d_2 (n - d_2)^3 (n - 1)\tilde{\sigma}^2.$$
(7.24)

Hence

(7.25)
$$\frac{2}{\mu d_1^4 n (n-1)(n-d_2)^2} S_{31} \le \frac{2\tilde{\sigma}^2}{\mu} \frac{d_2 \sqrt{n-d_2}}{\sqrt{n}} \le \frac{2\tilde{\sigma}^2 d_2}{\mu}.$$

Similarly, by the symmetry of u_1 and u_2 , we obtain $\frac{2}{\mu d_1^4 n(n-1)(n-d_2)^2} S_{32} \leq \frac{2\tilde{\sigma}^2 d_2}{\mu}$. Using the similar argument for the proof of (7.24), the upper bound in (7.25) holds for all other 4 terms in (7.14) as well. Therefore (7.14) is bounded by $\frac{12\tilde{\sigma}^2 d_2}{\mu}$. Combining the estimates for (7.12), (7.13) and (7.14), since $d_2 \leq n$, we then obtain

(7.26)
$$\mathbb{E}[D\mathbf{1}_{\mathcal{B}} \mid X] \le \frac{12\tilde{\sigma}^2 d_2^2}{\mu n} + \frac{12\tilde{\sigma}^2 d_2}{\mu} \le \frac{24\tilde{\sigma}^2 d_2}{\mu}.$$

By taking $\tau^2 = 24\tilde{\sigma}^2 d_2$, $c = 12d_2a$ in Theorem 2.2, the result follows.

Similar to Corollary 4.2, we obtain the following concentration inequalities from Theorem 7.2.

Corollary 7.3. Let
$$\gamma_0 = \frac{1}{p} - 1 = \frac{2d_2}{n - 2d_2}, c_0 = \frac{p}{6} = \frac{1}{6}(1 - \frac{2d_2}{n})$$
. We have

(7.27)
$$\mathbb{P}(g_Q(X) \ge (1 + \gamma_0)\mu + t) \le \exp\left(-c_0 \frac{\tilde{\sigma}^2}{d_2 a^2} h\left(\frac{at}{2\tilde{\sigma}^2}\right)\right),$$

(7.28)
$$\mathbb{P}(|g_Q(X) - \mu| \ge \gamma_0 \mu + t) \le 2 \exp\left(-\frac{c_0 t^2}{8d_2(\tilde{\sigma}^2 + \frac{1}{6}at)}\right).$$

8. Proof of Theorem 1.3

The remaining analysis is similar to the analysis in Section 5 for the proof of Theorem 1.1. We will only address the main differences. Let $M = XX^T - d_1I$. For convenience we now write (7.1) as

$$g_Q(X) = f_Q(M) = \sum_{u_1, u_2 \in [n]} Q_{u_1 u_2} M_{u_1 u_2}.$$

Recall $\lambda(M) = \sup_{x \in S_0^{n-1}} |\langle x, Mx \rangle|$. For fixed $x \in S_0^{n-1}$, we split the sum into light and heavy parts. Define light and heavy couples by

$$\mathcal{L}(x) = \{u, v \in [n] : |x_u x_v| \le \sqrt{d_1(d_2 - 1)}/n\}, \quad \mathcal{H}(x) = \{u, v \in [n] : |x_u x_v| > \sqrt{d_1(d_2 - 1)}/n\}.$$

We can decompose the linear form $f_O(M)$ as

$$f_{xx^{\top}}(M) = \langle x, Mx \rangle = f_{\mathcal{L}(x)}(M) + f_{\mathcal{H}(x)}(M),$$

where

(8.1)
$$f_{\mathcal{L}(x)}(M) = \sum_{(u,v)\in\mathcal{L}(x)} x_u x_v M_{uv}, \quad f_{\mathcal{H}(x)}(M) = \sum_{(u,v)\in\mathcal{H}(x)} x_u x_v M_{uv}.$$

Lemma 8.1. For any fixed $x \in S_0^{n-1}$, $\beta \ge 4\gamma_0\sqrt{d_1(d_2-1)}$, and $n \ge 2$,

(8.2)
$$\mathbb{P}\left(|f_{\mathcal{L}(x)}(M)| \ge (\beta+3)\sqrt{d_1(d_2-1)}\right) \le 4\exp\left(-\frac{3c_0\beta^2n}{8d_2(24+\beta)}\right).$$

Proof. Recall for $u \neq v$, $\mathbb{E} M_{uv} = \frac{d_1(d_2-1)}{n-1}$. For any fixed $x \in S_0^{n-1}$,

$$\begin{split} |\mathbb{E}f_{\mathcal{L}(x)}(M)| &\leq |\mathbb{E}\langle x, Mx\rangle| + |\mathbb{E}f_{\mathcal{H}(x)}(M)| \\ &\leq \left| x^{\top} \left(\mathbb{E}M - \frac{d_{1}(d_{2}-1)}{n} \mathbf{1}_{n} \mathbf{1}_{n}^{\top} \right) x \right| + \frac{d_{1}(d_{2}-1)}{n-1} \sum_{(u,v) \in \mathcal{H}(x)} |x_{u} x_{v}| \\ &\leq \left\| \mathbb{E}M - \frac{d_{1}(d_{2}-1)}{n} \mathbf{1}_{n} \mathbf{1}_{n}^{\top} \right\|_{F} + \frac{d_{1}(d_{2}-1)}{n-1} \sum_{(u,v) \in \mathcal{H}(x)} \frac{|x_{u} x_{v}|^{2}}{\sqrt{d_{1}(d_{2}-1)}/n} \\ &\leq \frac{d_{1}(d_{2}-1)}{\sqrt{n-1}} + \sqrt{d_{1}(d_{2}-1)} \frac{n}{n-1} \leq 3\sqrt{d_{1}(d_{2}-1)}. \end{split}$$

We split the set $\mathcal{L}(x)$ into two parts as $\mathcal{L}(x) = \mathcal{L}_{+}(x) \cup \mathcal{L}_{-}(x)$ where

$$\mathcal{L}_{+}(x) = \{u, v \in [n] : 0 \le x_{u} x_{v} \le \sqrt{d_{1}(d_{2} - 1)}/n\}, \quad \mathcal{L}_{-}(x) = \mathcal{L} \setminus \mathcal{L}_{+}(x).$$

Then

$$(8.3) |f_{\mathcal{L}(x)}(M) - \mathbb{E}f_{\mathcal{L}(x)}(M)| \le |f_{\mathcal{L}_{+}(x)}(M) - \mathbb{E}f_{\mathcal{L}_{+}(x)}(M)| + |f_{\mathcal{L}_{-}(x)}(M) - \mathbb{E}f_{\mathcal{L}_{-}(x)}(M)|.$$

Consider the first term in the right hand side of (8.3). By Cauchy's inequality,

$$\mu = \mathbb{E} f_{\mathcal{L}_+(x)}(M) \le \frac{d_1(d_2 - 1)}{n - 1} \sum_{uv} |x_u x_v| \le d_1(d_2 - 1) \frac{n}{n - 1} \le 2d_1(d_2 - 1).$$

Also we have $\tilde{\sigma}^2 = \sum_{uv \in \mathcal{L}_+(x)} |x_u x_v|^2 \mathbb{E} M_{uv} \le \frac{d_1(d_2 - 1)}{n - 1}$. Then by (7.28) with $a = \frac{\sqrt{d_1(d_2 - 1)}}{n}$, for any $\beta > 4\gamma_0 \sqrt{d_1(d_2 - 1)}$, we have

$$\mathbb{P}\left(|f_{\mathcal{L}_{+}(x)}(M)| - \mathbb{E}f_{\mathcal{L}_{+}(x)}(M)| \ge (\beta/2)\sqrt{d_{1}(d_{2}-1)}\right) \le 2\exp\left(-\frac{3c_{0}\beta^{2}n}{8d_{2}(24+\beta)}\right).$$

The same bound holds for the second term in (8.3). Then with probability at least $1 - 4 \exp\left(-\frac{3c_0\beta^2n}{8d_2(24+\beta)}\right)$,

$$|f_{\mathcal{L}(x)}(M)| \le |\mathbb{E}f_{\mathcal{L}(x)}(M)| + |f_{\mathcal{L}(x)}(M) - \mathbb{E}f_{\mathcal{L}(x)}(M)| \le (3+\beta)\sqrt{d_1(d_2-1)}$$
.

This completes the proof.

Similar to Lemma 5.4, the following lemma shows that the discrepancy property for $M = XX^{\top} - d_1I$ holds for a random bipartite biregular graph with high probability. The proof is similar, and we skip the details.

Lemma 8.2. Let $M = XX^{\top} - d_1I$ where X is the biadjacency matrix of a uniform random (n, m, d_1, d_2) -bipartite biregular graph. For any $K \geq 0$, With probability at least $1 - n^{-K}$, $\mathrm{DP}(\delta, \kappa_1, \kappa_2)$ holds for M with $\delta = \frac{d_1(d_2-1)}{n-1}$, $\kappa_1 = e^2(1+\gamma_0)^2$, and $\kappa_2 = \frac{8d_2}{c_0}(1+\gamma_0)(K+4)$.

Lemma 8.3. Let $M = XX^{\top} - d_1I$ where X is the biadjacency matrix of a (n, m, d_1, d_2) -bipartite biregular graph. Suppose M has $DP(\delta, \kappa_1, \kappa_2)$ with $\delta, \kappa_1, \kappa_2$ given in Lemma 8.2. Then there exists a constant α_0 depending on κ_1, κ_2 such that $f_{\mathcal{H}(x)}(M) \leq \alpha_0 \sqrt{d_1(d_2 - 1)}$, where $\alpha_0 = 16 + 64(\kappa_1 + 1) + 64\kappa_2 \left(1 + \frac{2}{\kappa_1 \log \kappa_1}\right)$.

Proof. Note that $\delta = \frac{d_1(d_2-1)}{n-1} \leq \frac{2d_1(d_2-1)}{n}$ for $n \geq 2$. The proof follows verbatim from [14, Lemma 6.6].

Now we are ready to prove Theorem 1.3 using the ε -net argument.

Lemma 8.4. For $\varepsilon \in (0, 1/2)$, let $\mathcal{N}_{\varepsilon}^0$ be an ε -net of S_0^{n-1} . Let X be the biadjacency matrix of a (n, m, d_1, d_2) -bipartite biregular graph and $M = XX^{\top} - d_1I$. Then

(8.4)
$$\lambda(M) \le \frac{1}{1 - 2\varepsilon} \sup_{x \in \mathcal{N}^0} |\langle x, Mx \rangle|.$$

Proof of Theorem 1.3. Fix K > 0. By Lemma 8.2, with probability at least $1 - n^{-K}$, M has $DP(\delta, \kappa_1, \kappa_2)$ property where the parameters $\delta, \kappa_1, \kappa_2$ are given in Lemma 8.2. Let \mathcal{D} be the event that this property holds. Then it suffices to show

$$\mathbb{P}\left(\mathcal{D}\cap\left\{\lambda(M)\geq\alpha\sqrt{d_1(d_2-1)}\right\}\right)\leq e^{-n}.$$

Take $\varepsilon = 1/4$ in (8.4). Then $\lambda(M) \leq 2 \sup_{x \in \mathcal{N}_0^0} |\langle x, Mx \rangle|$. We obtain

$$(8.5) \mathbb{P}\left(\mathcal{D}\cap\left\{\lambda(M)\geq\alpha\sqrt{d_1(d_2-1)}\right\}\right)\leq \sum_{x\in\mathcal{N}_2^0}\mathbb{P}\left(\mathcal{D}\cap\left\{|\langle x,Mx\rangle|\geq(\alpha/2)\sqrt{d_1(d_1-1)}\right\}\right).$$

For any fixed $x \in \mathcal{N}_{\varepsilon}$,

$$\mathbb{P}\left(\mathcal{D} \cap \left\{ |\langle x, Mx \rangle| \ge (\alpha/2)\sqrt{d_1(d_2 - 1)} \right\} \right)$$

$$\le \mathbb{P}\left(\mathcal{D} \cap \left\{ |f_{\mathcal{L}(x)}(M)| \ge (\alpha/2)\sqrt{d_1(d_2 - 1)} - |f_{\mathcal{H}(x)}(M)| \right\} \right)$$

$$\le \mathbb{P}\left(|f_{\mathcal{L}(x)}(M)| \ge (\alpha/2 - \alpha_0)\sqrt{d_1(d_2 - 1)} \right).$$

Take $\beta = \frac{1}{2}\alpha - \alpha_0 - 3$. When $\beta > 4\gamma_0\sqrt{d_2(d_1 - 1)}$, from (8.2), we have

$$(8.6) \mathbb{P}\left(\mathcal{D}\cap\left\{|\langle x,Mx\rangle|\geq (\alpha/2)\sqrt{d_1(d_2-1)}\right\}\right)\leq 2\exp\left(-\frac{3c_0\beta^2n}{8d_2(24+\beta)}\right).$$

Recall $\gamma_0 = \frac{2d_2}{n-2d_2}$, $c_0 = \frac{1}{6}(1-\frac{2d_2}{n})$ in Corollary 7.3. In the assumption of Theorem 1.3 we also have $d_2 \le n/4$ and $d_1 \le C_1 n^2$. Then it follows that $c_0 \ge \frac{1}{12}$, $\gamma_0 \le 1$ and

$$(8.7) 4\gamma_0 \sqrt{d_1(d_2-1)} = \frac{8d_2 \sqrt{d_1(d_2-1)}}{n-2d_2} \le \frac{8d_2 \sqrt{C_1(d_2-1)}n}{n-2d_2} \le 16d_2 \sqrt{(d_2-1)C_1}.$$

Also recall $\kappa_1=e^2(1+\gamma_0)^2$, $\kappa_2=\frac{8d_2}{c_0}(1+\gamma_0)(K+4)$ from Lemma 8.2. We have

$$e^2 \le \kappa_1 \le 4e^2$$
, $\kappa_2 \le 192d_2(K+4)$.

Then α_0 given in Lemma 8.3 is a bounded constant depending on d_2 and K. Since $|\mathcal{N}_{1/4}^0| \leq 9^n$, from (8.5) we have

(8.8)
$$\mathbb{P}\left(\mathcal{D}\cap\left\{\lambda(M)\geq\alpha\sqrt{d_1(d_2-1)}\right\}\right)\leq 2\cdot 9^n\exp\left(-\frac{3c_0\beta^2n}{8d_2(24+\beta)}\right).$$

Taking $\beta > 16d_2\sqrt{(d_2-1)C_1}$, then from (8.7), the condition $\beta > 4\gamma_0\sqrt{d_2(d_2-1)}$ holds for (8.6). We obtain from (8.8) that for sufficiently large β depending on d_2, C_1 ,

$$\mathbb{P}\left(\mathcal{D}\cap\left\{\lambda(M)\geq\alpha\sqrt{d_1(d_2-1)}\right\}\right)\leq 2\cdot 9^n\exp\left(-\frac{\frac{1}{4}\beta^2n}{8d_2(24+\beta)}\right)\leq e^{-n},$$

where $\alpha = 2(\beta + \alpha_0 + 3)$. We have $\mathbb{P}\left(\lambda(M) \ge \alpha \sqrt{d_1(d_2 - 1)}\right) \le n^{-K} + e^{-n}$. This completes the proof by taking a new constant $\alpha' = \alpha \sqrt{d_2 - 1}$.

Remark 8.5. In (8.8), we see that if d_2 is not a bounded constant, the probability bound cannot be o(1), since the union bound is taken over exponentially many points on a ε -net. This is a limitation of the method we use and we must assume d_2 is fixed in our proof.

Acknowledgements

The author thanks Ioana Dumitriu and Tobias Johnson for helpful discussion. This work is partially supported by NSF DMS-1949617. The author acknowledges support from NSF DMS-1928930 during his participation in the program "Universality and Integrability in Random Matrix Theory and Interacting Particle Systems" hosted by the Mathematical Sciences Research Institute in Berkeley, California during the Fall semester of 2021.

References

- [1] Noga Alon. Eigenvalues and expanders. Combinatorica, 6(2):83-96, 1986.
- [2] Noga Alon and Vitali D Milman. λ_1 , isoperimetric inequalities for graphs, and superconcentrators. *Journal of Combinatorial Theory*, Series B, 38(1):73–88, 1985.
- [3] Richard Arratia, Larry Goldstein, and Fred Kochman. Size bias for one and all. Probability Surveys, 16:1-61, 2019.
- [4] Roland Bauerschmidt, Jiaoyang Huang, Antti Knowles, and Horng-Tzer Yau. Bulk eigenvalue statistics for random regular graphs. The Annals of Probability, 45(6A):3626–3663, 2017.
- [5] Roland Bauerschmidt, Jiaoyang Huang, Antti Knowles, and Horng-Tzer Yau. Edge rigidity and universality of random regular graphs of intermediate degree. *Geometric and Functional Analysis*, pages 1–77, 2020.
- [6] Roland Bauerschmidt, Jiaoyang Huang, and Horng-Tzer Yau. Local Kesten-Mckay law for random regular graphs. Communications in Mathematical Physics, 369(2):523-636, 2019.
- [7] Roland Bauerschmidt, Antti Knowles, and Horng-Tzer Yau. Local semicircle law for random regular graphs. Communications on Pure and Applied Mathematics, 70(10):1898–1960, 2017.
- [8] Charles Bordenave. A new proof of Friedman's second eigenvalue theorem and its extension to random lifts. In *Annales scientifiques de l'Ecole normale supérieure*, 2019.
- [9] Gerandy Brito, Ioana Dumitriu, and Kameron Decker Harris. Spectral gap in random bipartite biregular graphs and applications. *Combinatorics, Probability and Computing*, 31(2):229–267, 2022.
- [10] Andrei Z Broder, Alan M Frieze, Stephen Suen, and Eli Upfal. Optimal construction of edge-disjoint paths in random graphs. SIAM Journal on Computing, 28(2):541–573, 1998.
- [11] Shantanu Prasad Burnwal and Mathukumalli Vidyasagar. Deterministic completion of rectangular matrices using asymmetric ramanujan graphs: Exact and stable recovery. *IEEE Transactions on Signal Processing*, 68:3834–3848, 2020.
- [12] Amin Coja-Oghlan and André Lanka. The spectral gap of random graphs with given expected degrees. the electronic journal of combinatorics, 16(1):R138, 2009.
- [13] Nicholas Cook. The circular law for random regular digraphs. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 55(4):2111–2167, 2019.
- [14] Nicholas Cook, Larry Goldstein, and Tobias Johnson. Size biased couplings and the spectral gap for random regular graphs. The Annals of Probability, 46(1):72–125, 2018.
- [15] Nicholas A Cook. Discrepancy properties for random regular digraphs. Random Structures & Algorithms, 50(1):23–58, 2017.
- [16] Nicholas A Cook. On the singularity of adjacency matrices for random regular digraphs. Probability Theory and Related Fields, 167(1-2):143-200, 2017.
- [17] Simon Coste. The spectral gap of sparse random digraphs. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 57(2):644–684, 2021.
- [18] Simon Coste, Gaultier Lambert, and Yizhe Zhu. The characteristic polynomial of sums of random permutations and regular digraphs. arXiv preprint arXiv:2204.00524, 2022.
- [19] Ioana Dumitriu, Tobias Johnson, Soumik Pal, and Elliot Paquette. Functional limit theorems for random regular graphs. Probability Theory and Related Fields, 156(3-4):921–975, 2013.
- [20] Ioana Dumitriu and Yizhe Zhu. Global eigenvalue fluctuations of random biregular bipartite graphs. arXiv preprint arXiv:2008.11760, 2020.
- [21] Uriel Feige and Eran Ofek. Spectral techniques applied to sparse random graphs. Random Structures & Algorithms, 27(2):251–275, 2005.
- [22] Keqin Feng and Wen-Ching Winnie Li. Spectra of hypergraphs and applications. Journal of number theory, 60(1):1–22, 1996.
- [23] Joel Friedman. A Proof of Alon's Second Eigenvalue Conjecture and Related Problems. Memoirs of the American Mathematical Society. American Mathematical Society, 2008.
- [24] Joel Friedman, Jeff Kahn, and Endre Szemerédi. On the second eigenvalue of random regular graphs. In Proceedings of the twenty-first annual ACM symposium on Theory of computing, pages 587–598, 1989.
- [25] Joel Friedman and Avi Wigderson. On the second eigenvalue of hypergraphs. Combinatorica, 15(1):43-65, 1995.
- [26] David Gamarnik, Quan Li, and Hongyi Zhang. Matrix completion from O(n) samples in linear time. In Conference on Learning Theory, pages 940–947, 2017.
- [27] Subhankar Ghosh and Larry Goldstein. Concentration of measures via size-biased couplings. Probability theory and related fields, 149(1):271–278, 2011.
- [28] Yukun He. Spectral gap and edge universality of dense random regular graphs. arXiv preprint arXiv:2203.07317, 2022.

- [29] Christopher Hoffman, Matthew Kahle, and Elliot Paquette. Spectral gaps of random graphs and applications. International Mathematics Research Notices, 05 2019.
- [30] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–561, 2006.
- [31] Roger A. Horn and Charles R. Johnson. Topics in Matrix Analysis. Cambridge University Press, 1991.
- [32] Jiaoyang Huang and Horng-Tzer Yau. Spectrum of random d-regular graphs up to the edge. arXiv preprint arXiv:2102.00963, 2021.
- [33] Tobias Johnson. Exchangeable pairs, switchings, and random regular graphs. The Electronic Journal of Combinatorics, 22(1):P1-33, 2015.
- [34] Raghunandan H Keshavan, Andrea Montanari, and Sewoong Oh. Matrix completion from a few entries. IEEE transactions on information theory, 56(6):2980–2998, 2010.
- [35] Jeong Han Kim, Benny Sudakov, and Van Vu. Small subgraphs of random regular graphs. Discrete Mathematics, 307(15):1961–1967, 2007.
- [36] Michael Krivelevich, Benny Sudakov, Van H Vu, and Nicholas C Wormald. Random regular graphs of high degree. Random Structures & Algorithms, 18(4):346–363, 2001.
- [37] Michel Ledoux. The concentration of measure phenomenon. Number 89. American Mathematical Soc., 2001.
- [38] Jing Lei and Alessandro Rinaldo. Consistency of spectral clustering in stochastic block models. *The Annals of Statistics*, 43(1):215–237, 2015.
- [39] Wen-Ching Winnie Li and Patrick Solé. Spectra of regular graphs and hypergraphs and orthogonal polynomials. *European Journal of Combinatorics*, 17(5):461–477, 1996.
- [40] Alexander Litvak, Anna Lytova, Konstantin Tikhomirov, Nicole Tomczak-Jaegermann, and Pierre Youssef. Structure of eigenvectors of random regular digraphs. Transactions of the American Mathematical Society, 371(11):8097–8172, 2019.
- [41] Alexander E Litvak, Anna Lytova, Konstantin Tikhomirov, Nicole Tomczak-Jaegermann, and Pierre Youssef. Adjacency matrices of random digraphs: singularity and anti-concentration. *Journal of Mathematical Analysis and Applications*, 445(2):1447–1491, 2017.
- [42] Alexander E Litvak, Anna Lytova, Konstantin Tikhomirov, Nicole Tomczak-Jaegermann, and Pierre Youssef. The smallest singular value of a shifted d-regular random square matrix. Probability Theory and Related Fields, 173(3-4):1301–1347, 2019.
- [43] Alexander E Litvak, Anna Lytova, Konstantin Tikhomirov, Nicole Tomczak-Jaegermann, and Pierre Youssef. Circular law for sparse random regular digraphs. Journal of the European Mathematical Society, 23(2):467–501, 2020.
- [44] Eyal Lubetzky, Benny Sudakov, and Van Vu. Spectra of lifted Ramanujan graphs. Advances in Mathematics, 227(4):1612–1645, 2011.
- [45] Adam W Marcus, Daniel A Spielman, and Nikhil Srivastava. Interlacing families I: Bipartite Ramanujan graphs of all degrees. Annals of Mathematics, pages 307–325, 2015.
- [46] Brendan D McKay. The expected eigenvalue distribution of a large regular graph. Linear Algebra and its Applications, 40:203–216, 1981.
- [47] Brendan D McKay. Subgraphs of random graphs with specified degrees. In Congressus Numerantium, volume 33, pages 213–223, 1981.
- [48] Brendan D McKay, Nicholas C Wormald, and Beata Wysocka. Short cycles in random regular graphs. the electronic journal of combinatorics, pages R66–R66, 2004.
- [49] Alon Nilli. On the second eigenvalue of a graph. Discrete Mathematics, 91(2):207-210, 1991.
- [50] Amir Sarid. The spectral gap of random regular graphs. arXiv preprint arXiv:2201.02015, 2022.
- [51] Michael Sipser and Daniel A Spielman. Expander codes. IEEE transactions on Information Theory, 42(6):1710–1722, 1996.
- [52] R Tanner. A recursive approach to low complexity codes. IEEE Transactions on information theory, 27(5):533-547, 1981.
- [53] Konstantin Tikhomirov and Pierre Youssef. The spectral gap of dense random regular graphs. The Annals of Probability, 47(1):362–419, 2019.
- [54] Konstantin Tikhomirov and Pierre Youssef. Sharp Poincaré and log-Sobolev inequalities for the switch chain on regular bipartite graphs. arXiv preprint arXiv:2007.02729, 2020.
- [55] Van Vu. Random discrete matrices. In Horizons of combinatorics, pages 257–280. Springer, 2008.
- [56] Van H Vu. Combinatorial problems in random matrix theory. In Proceedings ICM, volume 4, pages 489–508, 2014.
- [57] Nicholas C Wormald. Models of random regular graphs. London Mathematical Society Lecture Note Series, pages 239–298, 1999.
- [58] Kevin Yang. Bulk eigenvalue correlation statistics of random biregular bipartite graphs. arXiv preprint arXiv:1705.00083, 2017.
- [59] Kevin Yang. Local Marchenko-Pastur law for random bipartite graphs. arXiv preprint arXiv:1704.08672, 2017.
- [60] Zhixin Zhou and Yizhe Zhu. Sparse random tensors: Concentration, regularization and applications. Electronic Journal of Statistics, 15(1):2483–2516, 2021.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA IRVINE, IRVINE, CA 92697 Email address: yizhe.zhu@uci.edu