SINGULAR YAMABE METRICS BY EQUIVARIANT REDUCTION

ALI HYDER, ANGELA PISTOIA, AND YANNICK SIRE

ABSTRACT. We construct singular solutions to the Yamabe equation using a reduction of the problem in an equivariant setting. This provides a non-trivial geometric example for which the analysis is simpler than in Mazzeo-Pacard program. Our construction provides also a non-trivial example of a weak solution to the Yamabe problem involving an equation with (smooth) coefficients.

Contents

1. Introduction	1
2. Preliminaries	3
2.1. Function spaces	3
2.2. The singular solution	4
2.3. The linearized operator around the singular solution	4
3. A scheme of the proof	5
4. The linear operator L_{ε}	6
4.1. Injectivity of L_{ε} on $C_{\mu,\mathcal{D}}^{2,\alpha}(\Omega \setminus \Sigma)$	6
4.2. Uniform surjectivity of L_{ε} on $C_{\mu,\mathcal{D}}^{2,\alpha}(\Omega \setminus \Sigma)$	9
5. The non-linear term Q	12
6. Appendix	14
References	15

1. Introduction

We consider the semilinear elliptic equation

$$-\Delta_{\mathfrak{g}}u + hu = u^p, \ u > 0, \text{ on } (\mathcal{M}, \mathfrak{g})$$
(1)

where $(\mathcal{M}, \mathfrak{g})$ is a n-dimensional compact Riemannian manifold without boundary, h is a C^1 -real function on \mathcal{M} s.t. $-\Delta_{\mathfrak{g}} + h$ is coercive and p > 1.

We are interested in finding solutions which are singular at k-dimensional manifolds for some integer $k \geq 1$.

In the critical case, i.e. $p = 2_n^* - 1 := \frac{n+2}{n-2}$ when the equation (1) coincides with the Yamabe equation (for $h = R_{\mathfrak{g}}$ the scalar curvature of \mathcal{M}), solutions singular at isolated points and at k-dimensional manifolds are known provided k < (n-2)/2 (see [MP99, MS91, MP96, Sch88]).

In the present work, we provide a non trivial example of a geometric singular solution, in a much simpler setting than the original construction in [MP96]. Our idea is to rely on an equivariant reduction of the problem like the ones described for instance in [CP16].

Date: May 20, 2020.

²⁰¹⁰ Mathematics Subject Classification. Primary: 35J60, Secondary: 35C20, 58J60.

Key words and phrases. Singular solution, Yamabe problem, warped product manifold, equivariant solution.

A. Hyder was supported by SNSF Grant No. P400P2-183866. A. Pistoja was partially supported by project.

A. Hyder was supported by SNSF Grant No. P400P2-183866. A. Pistoia was partially supported by project Vain-Hopes within the program VALERE: VAnviteLli pEr la RicErca.

For any integer $0 \le k \le n-3$ let $2_{n,k}^* = \frac{2(n-k)}{n-k-2}$ be the (k+1)-st critical exponent. We remark that $2_{n,k}^* = 2_{n-k,0}^*$ is nothing but the critical exponent for the Sobolev embedding $H_{\mathfrak{g}}^1(\mathcal{M}) \hookrightarrow L_{\mathfrak{g}}^q(\mathcal{M})$, when $(\mathcal{M},\mathfrak{g})$ is a (n-k)-dimensional Riemannian manifold. In particular, $2_{n,0}^* = \frac{2n}{n-2}$ is the usual Sobolev critical exponent.

 $2_{n,0}^* = \frac{2n}{n-2}$ is the usual Sobolev critical exponent. In order to reduce the problem, we will consider the background manifold \mathcal{M} to be given by a warped product. Let (M,g) and (K,κ) be two riemannian manifolds of dimensions N and k, respectively. Let $\omega \in C^2(M)$, $\omega > 0$ be a given function. The warped product $\mathcal{M} = M \times_{\omega} K$ is the product (differentiable) n-dimensional (n = N + k) manifold $M \times K$ endowed with the riemannian metric $\mathfrak{g} = g + \omega^2 \kappa$. The function ω is called the warping function. For example, every surface of revolution (not crossing the axis of revolution) is isometric to a warped product, with M the generating curve, $K = S^1$ and $\omega(x)$ the distance from $x \in M$ to the axis of revolution.

It is not difficult to check that if $u \in C^2(M \times_{\omega} K)$ then

$$\Delta_{\mathfrak{g}}u = \Delta_{\mathfrak{g}}u + \frac{m}{\omega}g(\nabla_{\mathfrak{g}}u\nabla_{\mathfrak{g}}u) + \frac{1}{\omega^2}\Delta_{\kappa}u. \tag{2}$$

Assume h is invariant with respect to K, i.e. h(x,y) = h(x) for any $(x,y) \in M \times K$. If we look for solutions to (1) which are invariant with respect to K, i.e. u(x,y) = v(x) then by (2) we immediately deduce that u solves (1) if and only if v solves

$$-\Delta_g v - \frac{m}{\omega} g \left(\nabla_g v, \nabla_g v \right) + h v = v^p \quad \text{in } (M, g).$$
 (3)

or equivalently

$$-\operatorname{div}_q(\omega^N \nabla_q v) + \omega^N h v = \omega^N v^p, \ v > 0 \quad \text{in } (M, g).$$

It is clear that if v is a solution to problem (3) which is singular at a point $\xi_0 \in M$ then u(x,y)=v(x) is a solution to problem (1) which is singular only on the fiber $\{\xi_0\}\times K$, which is a k-dimensional submanifold of $M\times_{\omega}K$. It is important to notice that the fiber $\{\xi_0\}\times K$ is totally geodesic in $M\times_{\omega}K$ (and in particular a minimal submanifold of $M\times_{\omega}K$) if ξ_0 is a critical point of the warping function ω .

Therefore, we are lead to consider the more general anisotropic problem

$$-\operatorname{div}_{a}(a\nabla_{a}u) + ahu = au^{p}, \ u > 0 \quad \text{in } (M,g)$$

$$\tag{4}$$

where (M,g) is a N-dimensional compact Riemannian manifold, p>1, $h\in C^1(M)$ and $a\in C^2(M)$ with $\min_M a>0$. We will assume that the anisotropic operator $-\mathrm{div}_g\ (a\nabla_g u\)+ahu$ is coercive in $H^1(M)$. Our main result reads as follows.

Theorem 1.1. If $\frac{N}{N-2} , then the problem (4) has a solution which is singular at a point <math>\xi_0 \in M$.

As a consequence of the previous theorem and the above discussion, we deduce

Theorem 1.2. Assume that $(\mathcal{M}, \mathfrak{g})$ is a warped product $M \times_{\omega} K$. If $0 < k < \frac{n-2}{2}$ then there exists a solution invariant with respect to K of

$$-\Delta_{\mathfrak{g}}u + R_{\mathfrak{g}}u = u^{\frac{n+2}{n-2}}, \ u > 0, \ in (\mathcal{M}, \mathfrak{g})$$

which is singular on $\{\xi_0\} \times K$, where ξ_0 is any point on M. Furthermore, if ξ_0 is a critical point of ω then the submanifold $\{\xi_0\} \times K$ is minimal in M.

The proof of Theorem (1.1) follows the same strategy developed in [MP96]. In particular, we will replace the N-dimensional manifold M by a bounded smooth domain Ω in \mathbb{R}^N and we will focus on the Dirichlet boundary problem

$$\begin{cases}
-div(a\nabla u) + ahu = au^p & \text{in } \Omega \\
u = 0 & \text{on } \partial\Omega \\
u > 0 & \text{in } \Omega.
\end{cases}$$
(5)

Here $h \in C^1(M)$, $a \in C^2(\bar{\Omega})$ with $\min_{\Omega} a > 0$ and the anisotropic operator $-div(a\nabla u) + ahu$ is coercive in $H_0^1(\Omega)$. We will show the following result

Theorem 1.3. If $\frac{N}{N-2} , then the problem (5) has a solution which is singular at$ a point $\xi_0 \in \Omega$.

The modification in the arguments to solve the problem on the manifold instead of in the domain are minor and are described in the last section of [MP96].

The paper is organized as follows. The proof of Theorem 1.3 is carried out in Section 3 and relies on the linear theory studied in Section 4 together with a contraction mapping argument developed in Section 5. All the necessary technical tools are contained in Section 2 and in the Appendix 6.

2. Preliminaries

2.1. Function spaces. For $\sigma > 0$ we let N_{σ} to be the ball $B_{\sigma}(\xi_0)$. For $\alpha \in (0,1)$, $s \in (0,\sigma)$, $k \in \mathbb{N} \cup \{0\}$ and $\nu \in \mathbb{R}$ we define the seminorms

$$|w|_{k,\alpha,s} := \sum_{j=0}^{k} s^{j} \sup_{N_{s} \setminus N_{\frac{s}{2}}} |\nabla^{j} w| + s^{k+\alpha} \sup_{x,x' \in N_{s} \setminus N_{\frac{s}{2}}} \frac{|\nabla^{k} w(x) - \nabla^{k} w(x')|}{|x - x'|^{\alpha}}, \tag{6}$$

and the weighted Hölder norm ($\sigma > 0$ is fixed)

$$||w||_{C^{k,\alpha}_{\nu}} := |w|_{C^{k,\alpha}(\bar{\Omega}\setminus N_{\frac{\sigma}{2}})} + \sup_{0 \le s \le \sigma} s^{-\nu} |w|_{k,\alpha,s}.$$

The weighted Hölder space $C^{k,\alpha}_{\nu}(\Omega \setminus \Sigma)$ is defined by (here $\Sigma = \{\xi_0\}$)

$$C_{\nu}^{k,\alpha}(\Omega \setminus \Sigma) := \left\{ w \in C_{loc}^{k,\alpha}(\bar{\Omega} \setminus \Sigma) : \|w\|_{C_{\nu}^{k,\alpha}} < \infty \right\}.$$

The subspace of $C^{k,\alpha}_{\nu}(\Omega \setminus \Sigma)$ with Dirichlet boundary conditions will be denoted by

$$C^{k,\alpha}_{\nu,\mathcal{D}}(\Omega \setminus \Sigma) := \{ w \in C^{k,\alpha}_{\nu}(\Omega \setminus \Sigma) : w = 0 \text{ on } \partial\Omega \}.$$

The space $C_{\nu,\nu'}^{k,\alpha}(\mathbb{R}^N\setminus\{0\})$ is defined by

$$||w||_{C^{k,\alpha}_{\nu,\nu'}(\mathbb{R}^N\setminus\{0\})}:=||w||_{C^{k,\alpha}_{\nu}(B_2\setminus\{0\})}+\sup_{r>1}(r^{-\nu'}||w(r\cdot)||_{C^{k,\alpha}(\bar{B}_2\setminus B_1)}).$$

We now list some useful properties of the space $C^{k,\alpha}_{\nu}(\Omega \setminus \Sigma)$, see e.g. [MP96] and the book [PR00].

Lemma 2.1. The following properties hold.

- $\begin{array}{ll} \text{i)} & \textit{If } w \in C^{k+1,\alpha}_{\gamma}(\Omega \setminus \Sigma) \; \textit{then } \nabla w \in C^{k,\alpha}_{\gamma-1}(\Omega \setminus \Sigma). \\ \\ \text{ii)} & \textit{If } w \in C^{k+1,0}_{\gamma}(\Omega \setminus \Sigma) \; \textit{then } w \in C^{k,\alpha}_{\gamma}(\Omega \setminus \Sigma) \; \textit{for every } \alpha \in [0,1). \\ \\ \text{iii)} & \textit{For every } w_i \in C^{k,\alpha}_{\gamma_i}(\Omega \setminus \Sigma), \; i = 1,2, \; \textit{we have} \end{array}$

$$||w_1w_2||_{k,\gamma_1+\gamma_2,\alpha} \le C||w_1||_{k,\gamma_1,\alpha}||w_2||_{k,\gamma_2,\alpha},$$

for some C > 0 independent of w_1, w_2 .

iv) There exists C > 0 such that for every $w \in C^{k,\alpha}_{\gamma}(\Omega \setminus \Sigma)$ with w > 0 in $\bar{\Omega} \setminus \Sigma$ we have

$$||w^p||_{k,\gamma,\alpha} \le C||w||_{k,\gamma,\alpha}^p.$$

2.2. **The singular solution.** The building block for our theory is the existence of a singular solution with different behaviour at the origin and at infinity. The following theorem provides such a solution.

Theorem 2.2 ([MP96]). Suppose that $\frac{N}{N-2} . Then for every <math>\beta > 0$ there exists a unique radial solution u to

$$\begin{cases}
-\Delta u = u^p & \text{in } \mathbb{R}^N \setminus \{0\} \\
u > 0 & \text{in } \mathbb{R}^N \setminus \{0\} \\
\lim_{|x| \to 0} u(x) = \infty,
\end{cases}$$
(7)

such that

$$\lim_{r \to \infty} r^{N-2} u(r) = \beta, \quad \lim_{r \to 0^+} r^{\frac{2}{p-1}} u(r) = c_p := [k(p, N)]^{\frac{1}{p-1}},$$

where

$$k(p,N) = \frac{2}{p-1} \left(N - \frac{2p}{p-1} \right).$$

Let u be a singular radial solution to (7). Then $u_{\varepsilon}(x) := \varepsilon^{-\frac{2}{p-1}} u(\frac{x}{\varepsilon})$ is also a solution to (7). Note that

$$u_{\varepsilon}(x) \le C(\delta, u) \varepsilon^{N - 2 - \frac{2}{p - 1}} \quad \text{for } |x| \ge \delta,$$

which shows that $u_{\varepsilon} \to 0$ locally uniformly in $\mathbb{R}^N \setminus \{0\}$. Due to this scaling and the asymptotic behavior of u at infinity, for a given $\alpha > 0$, we can find a solution u_1 such that

$$r^2 u_1^{p-1}(r) \le \alpha$$
 on $(1, \infty)$.

2.3. The linearized operator around the singular solution. We consider the linearized operator

$$L_1 = \Delta + pu_1^{p-1}$$

where in polar coordinates we denote

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{N-1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \Delta_{\theta}.$$

Following [MP96], we recall that γ_j is an indicial root of L_1 at 0 if $L_1(|x|^{\gamma_j}\varphi_j) = o(|x|^{\gamma-2})$, where φ_j is the j-th eigenfunction of $-\Delta_\theta$ on S^{N-1} , that is $-\Delta_\theta \varphi_j = \lambda_j \varphi_j$,

$$\lambda_0 = 0$$
, $\lambda_j = N - 1$, for $j = 1, \dots, N$,

and so on. Setting

$$A_p := p \lim_{r \to 0} r^2 u_1^{p-1}(r) = pk(p, N).$$
(8)

we have that

$$\gamma_j^{\pm} = \frac{1}{2} \left[2 - N \pm \sqrt{(N-2)^2 + 4(\lambda_j - A_p)} \right].$$

For $\frac{N}{N-2} we have that (<math>\Re$ denotes the real part)

$$2 - N < -\frac{2}{p-1} < \Re(\gamma_0^-) \le \frac{2-N}{2} \le \Re(\gamma_0^+) < 0$$

and

$$\gamma_j^-<-\frac{2}{p-1}\quad\text{for }j\geq 1.$$

Since $\lim_{r\to\infty} r^2 u_1^{p-1}(r) = 0$, the indicial roots of L_1 at infinity are the same as for the Δ itself. These values are given by

$$\tilde{\gamma}_{j}^{\pm} = \frac{1}{2} \left[2 - N \pm \sqrt{(N-2)^2 + 4\lambda_{j}} \right].$$

We shall choose μ, ν in the region

$$\frac{-2}{p-1} < \nu < \min\left\{\frac{-2}{p-1} + 1, \Re(\gamma_0^-)\right\} \le \frac{2-N}{2} \le \Re(\gamma_0^+) < \mu < 0, \tag{9}$$

so that $\mu + \nu = 2 - N$.

We have the following propositions whose proofs can be found in [MP96].

Proposition 2.3. Let $w \in C^{2,\alpha}_{u,0}(\mathbb{R}^N \setminus \{0\})$ be a solution to $L_1w = 0$. Then $w \equiv 0$.

Proposition 2.4. Let $w \in C^{2,\alpha}_{\gamma,\gamma}(\mathbb{R}^N \setminus \{0\})$ be a solution to

$$\Delta w + \frac{A_p}{r^2} w = 0 \quad in \ \mathbb{R}^N \setminus \{0\},\,$$

where A_p is given by (8). If γ is not an indicial root of the operator $\Delta + \frac{A_p}{r^2}$ then $w \equiv 0$.

3. A SCHEME OF THE PROOF

Let $\Sigma = \{\xi_0\} \subset \Omega$. To construct a solution to (5) which is singular precisely at the point ξ_0 , we start by constructing an approximate solution to (5) which is singular exactly on Σ . For $\sigma > 0$ small (to be chosen later) let us first fix a non-negative cut-off function $\chi \in C_c^{\infty}(B_{\sigma}(\xi_0))$ such that $\chi = 1$ in $B_{\frac{\sigma}{2}}(\xi_0)$. An approximate solution \bar{u}_{ε} is defined by

$$\bar{u}_{\varepsilon}(x) = \chi(x)u_{\varepsilon}(x - \xi_0) = \varepsilon^{-\frac{2}{p-1}}\chi(x)u_1(\frac{x - \xi_0}{\varepsilon}).$$

We shall look for positive solutions of the form $u = \bar{u}_{\varepsilon} + v$. Then, v has to satisfy

$$L_{\varepsilon}v + f_{\varepsilon} + Q(v) = 0, \tag{10}$$

where the linear operator L_{ε} is

$$L_{\varepsilon}v := div(a\nabla v) + a[p\bar{u}_{\varepsilon}^{p-1} - h]v, \tag{11}$$

the error term is

$$f_{\varepsilon} := div(a\nabla \bar{u}_{\varepsilon}) - ah\bar{u}_{\varepsilon} + a\bar{u}_{\varepsilon}^{p}, \tag{12}$$

and the non-linear term Q is

$$Q(v) = a[|\bar{u}_{\varepsilon} + v|^p - \bar{u}_{\varepsilon}^p - p\bar{u}_{\varepsilon}^{p-1}v]. \tag{13}$$

To prove existence of solution to (10) we will use a fixed point argument on the space $C^{2,\alpha}_{\nu,\mathcal{D}}(\Omega\setminus\Sigma)$ for a suitable ν . We note that if $v\in C^{2,\alpha}_{\nu,\mathcal{D}}(\Omega\setminus\Sigma)$ solves (10) then by maximum principle we have that $\bar{u}_{\varepsilon}+v>0$ in Ω . This is a simple consequence of the fact that we will choose $\nu>-\frac{2}{p-1}$, and therefore, $\bar{u}_{\varepsilon}+v>0$ in a small neighborhood of Σ , thanks to the asymptotic behavior of \bar{u}_{ε} , around the origin and the coercivity assumption on h.

First of all, we estimate the size of the error term.

Lemma 3.1. The error f_{ε} satisfies

$$||f_{\varepsilon}||_{C^{0,\alpha}_{\gamma-2}} \le C_{\gamma} \max \left\{ \varepsilon^{1-\gamma-\frac{2}{p-1}}, \varepsilon^{N-\frac{2p}{p-1}} \right\} \quad \text{for } \varepsilon > 0 \text{ small},$$

for every $\gamma < 1 - \frac{2}{p-1}$.

Proof. We only estimate the first term in (6), the estimate for the second term should having the same order as of the first one. It follows that

$$u_{\varepsilon}(x-\xi_0) \approx \begin{cases} |x-\xi_0|^{-\frac{2}{p-1}} & \text{for } |x-\xi_0| \leq \varepsilon \\ \frac{\varepsilon^{N-\frac{2p}{p-1}}}{|x-\xi_0|^{N-2}} & \text{for } |x-\xi_0| \geq \varepsilon, \end{cases}$$

and

$$|\nabla u_{\varepsilon}(x-\xi_0)| \le C \begin{cases} |x-\xi_0|^{-\frac{2}{p-1}-1} & \text{for } |x-\xi_0| \le \varepsilon \\ \frac{\varepsilon^{N-\frac{2p}{p-1}}}{|x-\xi_0|^{N-1}} & \text{for } |x-\xi_0| \ge \varepsilon. \end{cases}$$

We write

$$f_{\varepsilon} = a[\Delta \bar{u}_{\varepsilon} + \bar{u}_{\varepsilon}^{p}] + \nabla \bar{u}_{\varepsilon} \cdot \nabla a - ah\bar{u}_{\varepsilon}.$$

Since $\chi \equiv 1$ in a small neighborhood of ξ_0 , we have

$$a|\Delta \bar{u}_{\varepsilon} + \bar{u}_{\varepsilon}^p| \le C\varepsilon^{N-\frac{2p}{p-1}}$$
 and $u_{\varepsilon}(x-\xi_0)|\nabla \chi \cdot \nabla a| \le \varepsilon^{N-\frac{2p}{p-1}}$.

Moreover,

$$|x - \xi_0|^{2-\gamma} |\nabla u_{\varepsilon}(x - \xi_0)| |\nabla a(x)| \le C \varepsilon^{1-\gamma - \frac{2}{p-1}},$$
$$|x - \xi_0|^{2-\gamma} \bar{u}_{\varepsilon}(x - \xi_0) < C \varepsilon^{2-\gamma - \frac{2}{p-1}}.$$

The lemma follows.

Next, we use the linear theory of L_{ε} developed in the Section 4 and, applying the inverse of L_{ε} , that is G_{ε} , we rewrite the above equation (10) as

$$v + G_{\varepsilon} f_{\varepsilon} + G_{\varepsilon} Q(v) = 0.$$

The crucial fact we shall use is that the norm of G_{ε} is uniformly bounded if ε is sufficiently small.

By Lemma 3.1, the error f_{ε} satisfies the estimate

$$||f_{\varepsilon}||_{0,\alpha,\nu-2} \le C\varepsilon^q, \quad q := \min\left\{N - \frac{2p}{p-1}, 1 - \nu - \frac{2}{p-1}\right\}.$$

Then, there exists $C_0 > 0$ such that $||G_{\varepsilon}f_{\varepsilon}||_{2,\alpha,\nu} \leq C_0 \varepsilon^q$. This suggests to work on the ball

$$\mathcal{B}_{\varepsilon,M} = \left\{ v \in C_{\nu}^{2,\alpha} : ||v||_{2,\alpha,\nu} \le M\varepsilon^q \right\},$$

for some $M > 2C_0$ large. In Section 5 we shall show that the map $v \mapsto G_{\varepsilon}[f_{\varepsilon} + Q(v)]$ is a contraction on the ball $\mathcal{B}_{\varepsilon,M}$ when M is large and ε is small enough. That will concludes our proof.

4. The linear operator L_{ε}

4.1. **Injectivity of** L_{ε} **on** $C^{2,\alpha}_{\mu,\mathcal{D}}(\Omega \setminus \Sigma)$. In this section we study injectivity of the linearized operator

$$L_{\varepsilon}w := div(a\nabla w) + a[p\bar{u}_{\varepsilon}^{p-1} - h]w.$$

We shall use the following notations:

$$\Omega_{\varepsilon} := \Omega \setminus B_{\varepsilon}(\xi_0), \quad f^+ := \max\{f, 0\}, \quad f^- := \min\{f, 0\}.$$

Lemma 4.1. After a suitable normalization of u_1 , the operator L_{ε} satisfies maximum principle in Ω_{ε} for $\varepsilon > 0$ small. More precisely, if $w \in H^1(\Omega_{\varepsilon})$ satisfies

$$\begin{cases} L_{\varepsilon}w \geq 0 & \text{in } \Omega_{\varepsilon} \\ w \leq 0 & \text{on } \partial \Omega_{\varepsilon}, \end{cases}$$

then $w \leq 0$ in Ω_{ε} .

Proof. The crucial fact we shall use is that the operator

$$v \mapsto -div(a\nabla v) + ahv,$$

is coercive, that is, for some $c_0 > 0$ we have

$$\int_{\Omega} a[|\nabla v|^2 + hv^2] dx \ge c_0 \int_{\Omega} |\nabla v|^2 dx \quad \text{for every } v \in H_0^1(\Omega).$$

Since $w \leq 0$ on the boundary $\partial \Omega_{\varepsilon}$, by extending w^+ by 0 on $B_{\varepsilon}(\xi_0)$ we see that $w^+ \in H_0^1(\Omega)$. Multiplying the inequality $L_{\varepsilon}w \geq 0$ by w^+ , and then integrating by parts we obtain

$$\int_{\Omega} a[|\nabla w^{+}|^{2} + h(w^{+})^{2} - p\bar{u}_{\varepsilon}^{p-1}(w^{+})^{2}]dx = 0.$$

We also have

$$\int_{\Omega} \frac{(w^+)^2}{|x - \xi_0|^2} dx \le \frac{4}{(N - 2)^2} \int_{\Omega} |\nabla w^+|^2 dx.$$

If we normalize u_1 in such a way that

$$pa\bar{u}_{\varepsilon}^{p-1} \le c_0 \frac{(N-2)^2}{8} \frac{1}{|x-\xi_0|^2}$$
 on Ω_{ε} ,

then we have

$$\int_{\Omega} |\nabla w^+|^2 dx = 0.$$

We conclude the lemma.

Remark 1. The above proof shows that L_{ε} satisfies maximum principle in $B_{\sigma}(\xi_0) \setminus B_{\varepsilon}(\xi_0)$ for $\varepsilon > 0$ small.

Lemma 4.2. Fix $\varepsilon_0 > 0$ such that L_{ε} satisfies maximum principle on Ω_{ε} . Let $2 - N < \gamma < 0$ be fixed. Let w_{ε} be a solution to $L_{\varepsilon}w_{\varepsilon} = f_{\varepsilon}$ on Ω_{ε} for some $f_{\varepsilon} \in C^{0,\alpha}_{\gamma-2}(\Omega_{\varepsilon})$, and $0 < \varepsilon \leq \varepsilon_0$. Assume that $w_{\varepsilon} = 0$ on $\partial\Omega$. Then there exists C > 0 such that

$$||w_{\varepsilon}||_{2,\alpha,\gamma} \le C\left(||f_{\varepsilon}||_{0,\alpha,\gamma-2} + \varepsilon^{-\gamma}||w_{\varepsilon}||_{C^{0}(\partial B_{\varepsilon}(\xi_{0}))}\right). \tag{14}$$

Proof. For $\phi(x) := |x - \xi_0|^{\gamma}$ we have

$$\Delta \phi(x) = c_{N,\gamma} |x - \xi_0|^{\gamma - 2}, \quad c_{N,\gamma} := \gamma(N + \gamma - 2) < 0.$$

Since

$$\nabla a \cdot \nabla \phi - ah\phi = O(|x - \xi_0|^{\gamma - 1}),$$

for $\sigma > 0$ small we have that

$$a\Delta\phi + \nabla a \cdot \nabla\phi - ah\phi \le \frac{c_{N,\gamma}}{2}a|x - \xi_0|^{\gamma - 2}$$
 on $B_{\sigma}(\xi_0)$.

This shows that for a suitable choice of u_1 , we have for some $\delta > 0$

$$L_{\varepsilon}\phi(x) \le -\delta|x-\xi_0|^{\gamma-2}$$
 on $\Omega := B_{\sigma}(\xi_0) \setminus B_{\varepsilon}(\xi_0)$.

Therefore, we can choose $c_{1,\varepsilon} \approx ||f_{\varepsilon}||_{0,\alpha,\gamma-2}$ so that

$$L_{\varepsilon}(w_{\varepsilon} + c_{1,\varepsilon}\phi) < 0$$
 on Ω .

We can also choose

$$c_{2,\varepsilon} \approx \varepsilon^{-\gamma} \|w_{\varepsilon}\|_{C^{0}(\partial B_{\varepsilon}(\xi_{0}))} + \|w_{\varepsilon}\|_{C^{0}(\partial B_{\sigma}(\xi_{0}))} =: c_{3,\varepsilon} + c_{4,\varepsilon},$$

so that

$$w_{\varepsilon} + (c_{1,\varepsilon} + c_{2,\varepsilon})\phi \geq 0$$
 on $\partial \Omega$.

Then by Maximum principle we have that (to get the other inequality use $-\phi$)

$$|w_{\varepsilon}| \leq (c_{1,\varepsilon} + c_{2,\varepsilon})\phi$$
 in Ω .

Since, $L_{\varepsilon}w_{\varepsilon}=f_{\varepsilon}$ in $\Omega_{\varepsilon}\setminus\Omega$, and $w_{\varepsilon}=0$ on $\partial\Omega$, we get that

$$|w_{\varepsilon}(x)| \lesssim (c_{1,\varepsilon} + c_{2,\varepsilon})$$
 for $x \in \Omega_{\varepsilon} \setminus \Omega$.

We claim that

$$c_{4,\varepsilon} \lesssim c_{3,\varepsilon} + ||f_{\varepsilon}||_{0,\alpha,\gamma-2}.$$

We assume by contradiction that the above claim is false. Then there exists a family of solutions $w_{\ell} = w_{\varepsilon_{\ell}}$ to $L_{\varepsilon_{\ell}} w_{\ell} = f_{\ell}$ with $0 < \varepsilon_{\ell} < \varepsilon_{0}$, $f_{\ell} \in C_{\gamma-2}^{0,\alpha}(\Omega_{\varepsilon_{\ell}})$, $w_{\ell} = 0$ on $\partial\Omega$ such that

$$c_{4,\varepsilon_{\ell}} = 1 \quad \text{and} \quad c_{3,\varepsilon_{\ell}} + ||f_{\ell}||_{0,\alpha,\gamma-2} \to 0.$$
 (15)

Then, up to a subsequence, $\Omega_{\varepsilon_{\ell}} \to \tilde{\Omega}_{\tilde{\varepsilon}}$, where

$$\tilde{\Omega}_{\tilde{\varepsilon}} = \Omega \setminus \{\xi_0\} \quad \text{if } \tilde{\varepsilon} = 0, \quad \text{and } \tilde{\Omega}_{\tilde{\varepsilon}} = \Omega_{\tilde{\varepsilon}} = \Omega \setminus B_{\tilde{\varepsilon}}(\xi_0) \quad \text{if } \tilde{\varepsilon} > 0.$$

From the estimates on w_{ℓ} we see that $w_{\ell} \to w$ in $\tilde{\Omega}_{\tilde{\varepsilon}}$. Moreover, w satisfies

$$\begin{cases} L_{\tilde{\varepsilon}}w = div(a\nabla w) - ahw + p\bar{u}_{\tilde{\varepsilon}}^{p-1}w = 0 & \text{in } \tilde{\Omega}_{\tilde{\varepsilon}} \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

Here, for $\tilde{\varepsilon} = 0$ the function $\bar{u}_{\tilde{\varepsilon}}$ is considered to be identically zero.

If $\tilde{\varepsilon} > 0$ then by Lemma 4.1 we get that $w \equiv 0$. Next we consider the case $\tilde{\varepsilon} = 0$. We have that $w(x) = O(|x - \xi_0|^{\gamma})$, and hence the singularity at ξ_0 is removable (note that $\gamma > 2 - N$), that is, $L_{\tilde{\varepsilon}}w = 0$ weakly in Ω . Thus, we can use coercivity hypothesis on h to conclude that $w \equiv 0$. This contradicts the first condition in (15).

In this way we have that there exists C > 0 independent of ε , but depending only on the right hand side of (14) such that

$$|w_{\varepsilon}| \leq C\phi$$
 in Ω_{ε} .

The desired estimate follows from Lemma 6.4 and a scaling argument (see e.g. [PR00, Chapter 2.2.1]).

Lemma 4.3. There exists $\varepsilon_0 > 0$ sufficiently small such that if $\varepsilon < \varepsilon_0$ then

$$L_{\varepsilon}: C^{2,\alpha}_{\mu,\mathcal{D}}(\Omega \setminus \Sigma) \to C^{0,\alpha}_{\mu-2}(\Omega \setminus \Sigma)$$

is injective.

Proof. We assume by contradiction that $L_{\varepsilon^{\ell}}$ is not injective for some $\varepsilon^{\ell} \to 0$. Let $w_{\ell} \in C^{2,\alpha}_{u,\mathcal{D}}(\Omega \setminus \Sigma)$ be a non-trivial solution to $L_{\varepsilon^{\ell}}w_{\ell} = 0$. We normalize w_{ℓ} so that

$$\max_{\partial \Omega_{\varepsilon^{\ell}}} \rho(x)^{-\mu} |w_{\ell}(x)| = (\varepsilon^{\ell})^{-\mu} \max_{\partial \Omega_{\varepsilon^{\ell}}} |w_{\ell}(x)| = 1, \tag{16}$$

where $\rho(x) = |x - \xi_0|$ in a small neighborhood of ξ_0 , and outside it is a smooth positive function. Then by Lemma 4.2 we get that

$$\sup_{\Omega_{\varepsilon^{\ell}}} \left(\rho(x)^{-\mu} |w_{\ell}(x)| + \rho(x)^{-\mu+1} |\nabla w_{\ell}(x)| \right) \le C. \tag{17}$$

We set

$$\tilde{w}_{\ell}(x) = (\varepsilon^{\ell})^{-\mu} w_{\ell}(\varepsilon^{\ell} x + \xi_0), \quad |x| < R_{\ell} := \frac{\sigma}{2\varepsilon^{\ell}}.$$

Then

$$\Delta \tilde{w}_{\ell}(x) + p u_1^{p-1} \tilde{w}_{\ell}(x) = f_{\ell}(x),$$

where

$$f_{\ell}(x) := (\varepsilon^{\ell})^{2-\mu} \left(h w_{\ell} - a^{-1} \nabla a \cdot \nabla w_{\ell} \right), \quad y := \varepsilon^{\ell} x + \xi_{0}$$
$$= (\varepsilon^{\ell})^{2} h(y) \tilde{w}_{\ell}(x) - \varepsilon^{\ell} a(y)^{-1} \nabla a(y) \cdot \nabla \tilde{w}_{\ell}(x).$$

It follows from (17) that $\tilde{w}_{\ell} \to \tilde{w}_{\infty}$ and $f_{\ell} \to 0$ in $C^2_{loc}(\mathbb{R}^N \setminus B_1)$ and $C^1_{loc}(\mathbb{R}^N \setminus B_1)$ respectively. Next we show that \tilde{w}_{ℓ} is bounded in $C^2_{loc}(B_2 \setminus \{0\})$. To this end it suffices to prove that

$$S_{\ell} = \sup_{B_2} \left(|x|^{-\mu} |\tilde{w}_{\ell}(x)| + |x|^{-\mu+1} |\nabla \tilde{w}_{\ell}(x)| \right) \le C.$$

We assume by contradiction that the above supremum is not uniformly bounded. Let $0 \neq x_{\ell} \in B_2$ be such that

$$S_{\ell} \approx |x_{\ell}|^{-\mu} |\tilde{w}_{\ell}(x_{\ell})| + |x_{\ell}|^{1-\mu} |\nabla \tilde{w}_{\ell}(x_{\ell})|.$$

We claim that $|x_{\ell}| \to 0$. On the contrary, if $x_{\ell} \to x_{\infty} \neq 0$, then setting $\bar{w}_{\ell} = \frac{w_{\ell}}{S_{\ell}}$ we see that $\bar{w}_{\ell} \to \bar{w}_{\infty}$, where

$$L_1\bar{w}_{\infty} = 0$$
 in $B_2 \setminus \{0\}$, $\bar{w}_{\infty} \equiv 0$ in $B_2 \setminus B_1$.

Therefore, $\bar{w}_{\infty} \equiv 0$ in B_2 , which contradicts to

$$|x_{\infty}|^{-\mu}|\bar{w}_{\infty}(x_{\infty})| + |x_{\infty}|^{1-\mu}|\nabla \bar{w}_{\infty}(x_{\infty})| \approx 1.$$

Thus we get that $x_{\ell} \to 0$.

Now we set

$$v_{\ell}(x) = \frac{r_{\ell}^{-\mu} \tilde{w}_{\ell}(r_{\ell}x)}{S_{\ell}}, \quad r_{\ell} := |x_{\ell}|.$$

Then, for every $\delta > 0$ and ℓ large we have

$$L_1 v_{\ell} = o_{\ell}(1), \quad |x|^{\mu} |v_{\ell}| + |x|^{1+\mu} |\nabla v_{\ell}| \le C \quad \text{for } \delta \le |x| \le \frac{1}{\delta}.$$

Therefore, up to a subsequence, $v_{\ell} \to v_{\infty}$ where v_{∞} satisfies

$$\Delta v_{\infty} + \frac{p\kappa(p, N)}{|x|^2} v_{\infty} = 0, \quad |v_{\infty}(x)| \le C|x|^{\mu} \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

Hence, by Proposition 2.4 we have $v_{\infty} \equiv 0$, a contradiction to $\max_{\partial B_1}(|v_{\infty}| + |\nabla v_{\infty}|) \approx 1$. This proves that $S_{\ell} \leq C$, and consequently we obtain that $\tilde{w}_{\ell} \to \tilde{w}_{\infty}$ in $C^1_{loc}(\mathbb{R}^N \setminus \{0\})$. Then the limit function \tilde{w}_{∞} would satisfy

$$\Delta \tilde{w}_{\infty} + p u_1^{p-1} \tilde{w}_{\infty} = 0 \text{ in } \mathbb{R}^N \setminus \{0\}, \quad \tilde{w}_{\infty} \in C^{2,\alpha}_{\mu,\mu}(\mathbb{R}^N \setminus \{0\}).$$

Then by Proposition 2.3 we have $\tilde{w}_{\infty} \equiv 0$, a contradiction to (16).

4.2. Uniform surjectivity of L_{ε} on $C_{\mu,\mathcal{D}}^{2,\alpha}(\Omega \setminus \Sigma)$. Instead of using general theory of edge operators as developed in [Maz91], we shall use the notes of Pacard [Paca, Pacb] and Pacard-Rivière [PR00] for edge operators with point singularity. Denoting $\rho(x) := |x - \xi_0|$, the weighted space $L_{\delta}^2(\Omega \setminus \Sigma)$ is defined by (we may also simply write L_{δ}^2 or $L_{\delta}^2(\Omega)$)

$$L^2_\delta(\Omega \setminus \Sigma) := \left\{ w \in L^2_{loc}(\Omega \setminus \Sigma) : \int_\Omega \rho^{-2-2\delta} |w|^2 dx < \infty \right\}.$$

Let $L^2_{-\delta}(\Omega \setminus \Sigma)$ be the dual of $L^2_{\delta}(\Omega \setminus \Sigma)$ with respect to the pairing

$$L^2_{\delta}(\Omega \setminus \Sigma) \times L^2_{-\delta}(\Omega \setminus \Sigma) \ni (w_1, w_2) \longrightarrow \int_{\Omega} w_1 w_2 \rho^{-2} dx.$$

We note that the following embedding is continuous

$$C^{k,\alpha}_{\gamma}(\Omega \setminus \Sigma) \hookrightarrow L^2_{\delta}(\Omega \setminus \Sigma) \quad \text{for } \delta < \gamma + \frac{N-2}{2}.$$

Lemma 4.4. Let $w \in L^2_{\delta}$ be a solution to

$$L_{\varepsilon}w = 0$$
 in $\Omega \setminus \{\xi_0\}$.

Then $w \in C^{2,\alpha}_{\delta-\frac{N-2}{2}}(\tilde{\Omega})$ for every $\tilde{\Omega} \subseteq \Omega$.

Proof. For $x_0 \in \Omega$ with $d(x_0, \partial\Omega) \geq |x_0 - \xi_0|$ we set

$$v(x) = w(x_0 + Rx), \quad R = \frac{1}{2}|x - \xi_0|, \quad |x| \le 1.$$

Then using the elliptic regularity for v, namely,

$$||v||_{C^0(B_{\frac{1}{2}})} \le C||v||_{L^2(B_1)},$$

one obtains that $|w(x)| \leq C|x|^{\delta - \frac{n-2}{2}}$ for x in a small neighborhood of ξ_0 . In fact, by elliptic regularity, this estimate also holds on compact sets in Ω . The lemma follows by a scaling argument and Schauder regularity.

The natural domain $D(L_{\varepsilon})$ of the operator L_{ε} is the set of functions $w \in L^2_{\delta}$ such that the distributional derivative $L_{\varepsilon}w$ is in $L^2_{\delta-2}$. More precisely, $w \in D(L_{\varepsilon})$ if there exists $f \in L^2_{\delta-2}$ such that w satisfies $L_{\varepsilon}w = f$ in the sense of distributions in $\Omega \setminus \Sigma$. However, in order to identify the adjoint of L_{ε} in a natural way, one has to consider a smaller space including the boundary condition w = 0 on $\partial\Omega$, which well-defined as a trace according to the next estimate.

Together with Lemma 6.3 and a rescaling argument (see e.g., [Paca, Proposition 1.2.1]) one can show that the following elliptic estimate holds: for $r_0 > 0$ with $B_{2r_0}(\xi_0) \subset \Omega$

$$\sum_{\ell=1}^{2} \|\nabla^{\ell} w\|_{L^{2}_{\delta-\ell}(B_{r_{0}}(\xi_{0}))} \le C(\|f\|_{L^{2}_{\delta-2}(B_{2r_{0}}(\xi_{0}))} + \|w\|_{L^{2}_{\delta}(B_{2r_{0}}(\xi_{0}))}). \tag{18}$$

In our next lemma we bound the weighted norm $||w||_{L^2_{\delta}}$ by L^2 norm of w and the weighted norm $||f||_{L^2_{\delta-2}}$ for some values of δ .

Lemma 4.5. Assume that $\delta - \frac{N-2}{2} \notin \{\Re \gamma_j^{\pm} : j = 0, 1, \dots\}$. Then there exists a compact set $K \subset \bar{\Omega} \setminus \{\xi_0\}$ and r > 0 such that

$$||w||_{L^{2}_{\delta}(B_{r}(\xi_{0}))} \le C(\varepsilon)(||f||_{L^{2}_{\delta-2}(\Omega)} + ||w||_{L^{2}(K)}). \tag{19}$$

Proof. Let R > 0 be such that $B_{4R}(\xi_0) \subset \Omega$. Applying Lemma 6.2 on the ball $B_R(\xi_0)$ we get that

$$||u||_{L^{2}_{\delta}(B_{R}(\xi_{0}))} \leq C_{1} \left[||h||_{L^{2}_{\delta-2}(B_{R}(\xi_{0}))}|| + ||\nabla a \cdot \nabla w||_{L^{2}_{\delta-2}(B_{R}(\xi_{0}))} + ||u||_{L^{2}(K)} \right],$$

for some $C_1 > 0$ and for some compact set $K \subset \bar{B}_R(\xi_0) \setminus \{\xi_0\}$. Since $\|\nabla a\| \in L^{\infty}(\Omega)$, it follows that

$$\lim_{r \to 0} \frac{\|\nabla a \cdot \nabla w\|_{L^2_{\delta-2}(B_r(\xi_0))}}{\|\nabla w\|_{L^2_{\delta-1}(B_r(\xi_0))}} = 0.$$

Therefore, for r > 0 small enough, the weighted norm $\|\nabla a \cdot \nabla w\|_{L^2_{\delta-2}(B_r(\xi_0))}$ can be absorbed one the left hand side, thanks to (18). On the region $B_R(\xi_0) \setminus B_r(\xi_0)$, the weighted norm $\|\nabla a \cdot \nabla w\|_{L^2_{\delta-2}}$ is equivalent to $\|\nabla a \cdot \nabla w\|_{L^2}$, and this can be controlled by

$$||w||_{L^2(B_{2R}(\xi_0)\setminus B_{\frac{r}{2}}(\xi_0)} + ||f||_{L^2(B_{2R}(\xi_0)\setminus B_{\frac{r}{2}}(\xi_0)}.$$

We conclude the lemma.

As a consequence of (18)-(19) one can prove the following lemma (see e.g., Chapter 9, [Pacb]).

Lemma 4.6. The operator $L_{\varepsilon}: L_{\delta}^2 \to L_{\delta-2}^2$ is Fredholm, provided $\delta - \frac{N-2}{2} \notin \{\Re \gamma_j^{\pm}: j = 0, 1, \dots\}$.

We shall fix $\delta > 0$ slightly bigger than $\mu + \frac{N-2}{2}$, where μ is fixed according to (9). Thanks to the previous comment on the domain of L_{ε} , the adjoint of the operator

$$L_{\varepsilon}: L_{-\delta}^2 \to L_{-\delta-2}^2 \tag{20}$$

is given by

$$L_{\delta+2}^2 \to L_{\delta}^2, \quad w \mapsto \rho^2 L_{\varepsilon}(w\rho^{-2}).$$
 (21)

Then the adjoint operator (21) is injective, and L_{ε} in (20) is surjective. Using the isomorphism

$$\rho^{2\delta}: L^2_{\tilde{\delta}} \to L^2_{2\delta + \tilde{\delta}}, \quad w \mapsto \rho^{2\delta} w,$$

we identify the adjoint operator as

$$L_{\varepsilon}^*: L_{-\delta+2}^2 \to L_{-\delta}^2, \quad w \mapsto \rho^{2-2\delta} L_{\varepsilon}(w\rho^{2\delta-2}).$$

Now we consider the composition

$$\mathcal{L} = L_{\varepsilon} \circ L_{\varepsilon}^* : L_{-\delta+2}^2 \to L_{-\delta-2}^2, \quad w \mapsto L_{\varepsilon}[\rho^{2-2\delta}L_{\varepsilon}(w\rho^{2\delta-2})].$$

Then \mathcal{L} is an isomorphism, and hence there exists a two sided inverse

$$\mathcal{G}_{\varepsilon}: L^2_{-\delta-2} \to L^2_{-\delta+2}.$$

Consequently, the right inverse of L_{ε} is given by $G_{\varepsilon} := L_{\varepsilon}^* \mathcal{G}_{\varepsilon}$. It follows that

$$G_{\varepsilon}: C^{0,\alpha}_{\nu-2}(\Omega \setminus \Sigma) \to C^{2,\alpha}_{\nu,\mathcal{D}}(\Omega \setminus \Sigma)$$

is bounded.

Lemma 4.7. Let $\varepsilon_0 > 0$ be as in Lemma 4.3. Then for $0 < \varepsilon < \varepsilon_0$ the system $L_{\varepsilon}w_1 = 0$, $w_1 = L_{\varepsilon}^* w_2$ with $w_1 \in C_{\nu,\mathcal{D}}^{2,\alpha}(\Omega \setminus \Sigma)$ and $w_2 \in C_{\nu+2,\mathcal{D}}^{4,\alpha}(\Omega \setminus \Sigma)$ has only trivial solution.

Proof. We set $w = \rho^{2\delta-2}w_2$. Then $L_{\varepsilon}[\rho^{2-2\delta}L_{\varepsilon}w] = 0$. Multiplying the equation by w and then integrating by parts we get

$$0 = \int_{\Omega} \rho^{2-2\delta} |L_{\varepsilon}w|^2 dx.$$

Since $\nu + 2\delta > \mu$, we have $w \in C^{2,\alpha}_{\nu+2\delta}(\Omega \setminus \Sigma) \subset C^{2,\alpha}_{\mu}(\Omega \setminus \Sigma)$. Then by Lemma 4.3 we get that w = 0, equivalently $w_1 = w_2 = 0$.

Lemma 4.8. There exists $\varepsilon_0 > 0$ small such that if $0 < \varepsilon < \varepsilon_0$, then the sequence of solutions $(w_{1,\varepsilon^{\ell}}) \subset C^{2,\alpha}_{\nu,\mathcal{D}}(\Omega \setminus \Sigma) \cap L^*_{\varepsilon^{\ell}}[C^{4,\alpha}_{\nu+2,\mathcal{D}}(\Omega \setminus \Sigma)]$ to $L_{\varepsilon^{\ell}}w_{1,\varepsilon^{\ell}} = f_{\varepsilon^{\ell}}$ is uniformly bounded in $C^{2,\alpha}_{\nu}(\Omega \setminus \Sigma)$, provided $(f_{\varepsilon^{\ell}})$ is uniformly bounded in $C^{0,\alpha}_{\nu-2}(\Omega \setminus \Sigma)$.

Proof. Assume by contradiction that the lemma is false. Then there exists a sequence $\varepsilon^{\ell} \to 0$ and $w_{1,\varepsilon^{\ell}} \in C^{2,\alpha}_{\nu,\mathcal{D}}(\Omega \setminus \Sigma) \cap L^*_{\varepsilon^{\ell}}[C^{4,\alpha}_{\nu+4,\mathcal{D}}(\Omega \setminus \Sigma)]$ with $L_{\varepsilon^{\ell}}w_{1,\varepsilon^{\ell}} = f_{\varepsilon^{\ell}}$ such that $\|f_{\varepsilon^{\ell}}\|_{C^{0,\alpha}_{\nu-2,\mathcal{D}}(\Omega \setminus \Sigma)} \le C$, and $(w_{1,\varepsilon^{\ell}})$ is not bounded in $C^{2,\alpha}_{\nu}(\Omega \setminus \Sigma)$. By Lemma 4.2

$$\|w_{1,\varepsilon^\ell}\|_{C^{2,\alpha}_\nu(\Omega_{\varepsilon^\ell})} \leq C + C \max_{\partial B_{\varepsilon^\ell}(\xi_0)} (\varepsilon^\ell)^{-\nu} \left(|w_{1,\varepsilon^\ell}| + \varepsilon^\ell |\nabla w_{1,\varepsilon^\ell}|\right) =: C + C S_{\varepsilon^\ell}.$$

We distinguish the following two cases.

Case 1 $S_{\varepsilon^{\ell}} \leq C$.

In this case we proceed as in the proof of Lemma 4.3. Let $x_{\ell} \in B_{\varepsilon^{\ell}}(\xi_0)$ be such that

$$\sup_{B_{\varepsilon^{\ell}}(\xi_{0})} \left(\rho^{-\nu} |w_{1,\varepsilon^{\ell}}| + \rho^{-\nu+1} |\nabla w_{1,\varepsilon^{\ell}}| \right) \approx |x_{\ell} - \xi_{0}|^{-\nu} \left(|w_{1,\varepsilon^{\ell}}(x_{\ell})| + |x_{\ell} - \xi_{0}| |\nabla w_{1,\varepsilon^{\ell}}(x_{\ell})| \right) =: S_{\ell} \to \infty.$$

Then necessarily $r_{\ell} := |x_{\ell} - \xi_0| = o(\varepsilon^{\ell})$. Setting

$$\tilde{w}_{1,\varepsilon^{\ell}}(x) := \frac{r_{\ell}^{-\nu} w_{1,\varepsilon^{\ell}}(r_{\ell}x + \xi_0)}{S_{\ell}}$$

one would get that $\tilde{w}_{1,\varepsilon^\ell} \to \tilde{w}_1 \not\equiv 0$ where

$$\tilde{L}_1 \tilde{w}_1 = 0$$
 in $\mathbb{R}^n \setminus \{0\}$, $r^{-\nu} |\tilde{w}_1| \le C$, $\tilde{L}_1 := \Delta + \frac{A_p}{r^2}$

where A_p is as in (8). Since ν does not coincide with indicial roots of \tilde{L}_1 , from Proposition 2.4 we get that $\tilde{w}_1 \equiv 0$, a contradiction.

Case 2 $S_{\varepsilon^{\ell}} \to \infty$.

In this case we set

$$\tilde{w}_{1,\varepsilon^{\ell}}(x) = (\varepsilon^{\ell})^{-\nu} \frac{w_{1,\varepsilon^{\ell}}(\varepsilon^{\ell}x + \xi_0)}{S_{\varepsilon^{\ell}}}.$$

Then $\max_{\partial B_1}(|\tilde{w}_{1,\varepsilon^{\ell}}|+|\nabla \tilde{w}_{1,\varepsilon^{\ell}}|)\approx 1$. Moreover, proceeding as before (see Lemma 4.3) we would get that $w_{1,\varepsilon^{\ell}}\to \tilde{w}_1\not\equiv 0$ where

$$L_1\tilde{w}_1 = 0$$
 in $\mathbb{R}^n \setminus \{0\}$, $r^{-\nu}|\tilde{w}_1| \le C$.

Since \tilde{w}_1 decays at infinity, its decay rate is determined by the indicial roots of L_1 (which are exactly the same as Δ) at infinity. In fact, \tilde{w}_1 would be bounded by r^{2-N} at infinity, see e.g., [MP96].

Since $w_{1,\varepsilon^{\ell}} \in L_{\varepsilon^{\ell}}^*[C_{\nu+2,\mathcal{D}}^{4,\alpha}(\Omega \setminus \Sigma)]$, we have $w_{1,\varepsilon^{\ell}} = \rho^{2-2\delta}L_{\varepsilon^{\ell}}w_{2,\varepsilon^{\ell}}$ for some $w_{2,\varepsilon^{\ell}} \in C_{\nu+2\delta,\mathcal{D}}^{4,\alpha}(\Omega \setminus \Sigma)$. Now we set

$$\tilde{w}_{2,\varepsilon^\ell}(x) := \frac{(\varepsilon^\ell)^{-\nu - 2\delta} w_{2,\varepsilon^\ell}(\varepsilon^\ell x + \xi_0)}{S_{\varepsilon^\ell}}.$$

Using that $2\delta + \nu > \mu$, and following the proof of Lemma 4.3, one can show that the family $\tilde{w}_{2,\varepsilon^{\ell}}$ converges to a limit function \tilde{w}_2 , where

$$L_1 \tilde{w}_2 = |x|^{2\delta - 2} \tilde{w}_1 \text{ in } \mathbb{R}^n \setminus \{0\}, |x|^{-\nu - 2\delta} |\tilde{w}_2| \le C.$$

Thus, $L_1[r^{2-2\delta}L_1\tilde{w}_2]=0$. We multiply this equation by \tilde{w}_2 and integrate it on \mathbb{R}^N . Then an integration by parts leads $L_1\tilde{w}_2=0$ (this is justified because of the decay of \tilde{w}_1 at infinity, provided we choose $\delta>0$ sufficiently close to $\mu+\frac{N-2}{2}$). Again, as $2\delta+\nu>\mu$, by Proposition 2.3 we have $\tilde{w}_2=\tilde{w}_1=0$, a contradiction.

5. The non-linear term Q

Lemma 5.1. Let $M_1 > 1$ be fixed. Then for $\varepsilon_0 << 1$ we have

$$||Q(v_1) - Q(v_2)||_{0,\alpha,\nu-2} \le \frac{1}{M_1} ||v_1 - v_2||_{2,\alpha,\nu}$$

for every $v_1, v_2 \in \mathcal{B}_{\varepsilon,M} := \left\{ v \in C_{\nu}^{2,\alpha} : ||v||_{2,\alpha,\nu} \le M\varepsilon^q \right\}$.

Proof. In Lemma 3.1, the error term f_{ε} is bounded by the maximum of two terms. If the maximum is the second term $\varepsilon^{N-2p/(p-1)}$, we argue as in [MP96]. Let us consider the case when the maximum is the first term. Let

$$q_1 := \left(N - \frac{2p}{p-1}\right) - \left(1 - \nu - \frac{2}{p-1}\right) = N + \nu - 3 > 0.$$

We start by showing that there exists $\tau > 0$ small (independent of $\varepsilon << 1$) such that

$$|v(x)| \le \frac{1}{10} \bar{u}_{\varepsilon}(x)$$
 for every $x \in B_{\tau_{\varepsilon}}(\xi_0), v \in \mathcal{B}_{\varepsilon,M},$ (22)

where

$$\tau_{\varepsilon} := \tau \varepsilon^{\frac{q_1}{\nu - 2 + N}} \to 0.$$

To prove this we recall that there exists c_1 , $c_2 > 1$ such that

$$\frac{1}{c_1} \le |x|^{\frac{2}{p-1}} u_{\varepsilon}(x) \le c_1 \quad \text{for } |x| \le \varepsilon,$$

$$\frac{1}{c_2} \le \varepsilon^{-N + \frac{2p}{p-1}} |x|^{N-2} u_{\varepsilon}(x) \le c_2 \quad \text{for } \varepsilon \le |x| \le \tau.$$

On the other hand,

$$\varepsilon^{-N + \frac{2p}{p-1}} \rho(x)^{-\nu} |v(x)| \le M.$$

As $\nu > \frac{-2}{p-1}$, we have (22) for some $\tau > 0$ small.

We have

$$Q(v_1) - Q(v_2) = a \int_0^1 \frac{d}{dt} |\bar{u}_{\varepsilon} + v_1 + t(v_1 - v_2)|^p dt - p\bar{u}_{\varepsilon}^{p-1}(v_1 - v_2)$$

$$= ap(v_1 - v_2) \int_0^1 (|\bar{u}_{\varepsilon} + v_1 + t(v_1 - v_2)|^{p-1} - \bar{u}_{\varepsilon}^{p-1}) dt$$

$$=: ap(v_1 - v_2) \int_0^1 Q(v_1, v_2) dt.$$

Next, using that

$$(1+r)^{p-1} = 1 + O(|r|)$$
 for $|r| \le \frac{1}{2}$,

we estimate for $x \in B_{\varepsilon}(\xi_0)$

$$|Q(v_1, v_2)|(x) \le C \bar{u}_{\varepsilon}(x)^{p-1} \frac{|v_1|(x) + |v_2|(x)}{\bar{u}_{\varepsilon}(x)}$$

$$\le C M \varepsilon^{\frac{2}{p-1} + \nu + q} \rho^{-2}(x)$$

$$= C M \varepsilon \rho(x)^{-2},$$

and for $x \in B_{\tau_{\varepsilon}}(\xi_0) \setminus B_{\varepsilon}(\xi_0)$

$$|Q(v_1, v_2)|(x) \le CM\rho(x)^{-2} \max\{\varepsilon, \varepsilon^{(N - \frac{2p}{p-1})(p-2) + q} \tau_{\varepsilon}^{(2-N)(p-2) + \nu + 2}\}$$

= $CM\rho(x)^{-2} o_{\varepsilon}(1)$.

Here we have used that the second term in the maximum is of the order ε^r for some r > 0. Indeed, from the definition of τ_{ε} , q and q_1 , the exponent of ε is

$$\left(N - \frac{2p}{p-1} \right) (p-2) + q + \left[(2-N)(p-2) + \nu + 2 \right] \left(1 - \frac{1}{N+\nu-2} \right)$$

$$= 1 - \frac{(2-N)(p-2) + \nu + 2}{N+\nu-2}$$

$$= \frac{(N-2)(p-1) - 2}{N+\nu-2}$$

$$> 0.$$

where the last inequality follows from $p > \frac{N}{N-2}$ and $N + \nu - 2 > 0$. Finally, as $\nu > -\frac{2}{p-1}$, we easily obtain for $x \in \Omega \setminus B_{\tau_{\varepsilon}}(\xi_0)$

$$|Q(v_1, v_2)|(x) \le C(\bar{u}_{\varepsilon}^{p-1} + |v_1|^{p-1} + |v_2|^{p-1})(x)$$

$$\le C\rho(x)^{-2} \left(\bar{u}_{\varepsilon}^{p-1}(x)\rho(x)^2 + M\varepsilon^{q(p-1)}\right)$$

$$= o_{\varepsilon}(1)\rho(x)^{-2}.$$

Combining these estimates we get for $\varepsilon << 1$

$$||Q(v_1) - Q(v_2)||_{0,0,\nu-2} = o_{\varepsilon}(1)||v_1 - v_2||_{0,0,\nu} = o_{\varepsilon}(1)||v_1 - v_2||_{2,\alpha,\nu}.$$

Next we estimate the weighted Hölder norm of $Q(v_1) - Q(v_2)$ with Hölder exponent $\alpha \le p-1$. For $0 < s < \sigma$ we write

$$s^{2-\nu+\alpha} \sup_{x,x' \in N_s \setminus N_{\frac{s}{2}}} \frac{|[Q(v_1) - Q(v_2)](x) - [Q(v_1) - Q(v_2)](x')|}{|x - x'|^{\alpha}}$$

$$\leq 4||Q(v_1) - Q(v_2)||_{0,0,\nu-2}$$

$$+ s^{2-\nu+\alpha} \sup_{x,x' \in N_s \setminus N_{\frac{s}{2}}, |x-x'| \leq \frac{s}{4}} \frac{|[Q(v_1) - Q(v_2)](x) - [Q(v_1) - Q(v_2)](x')|}{|x - x'|^{\alpha}}.$$

Notice that for $x, x' \in N_s \setminus N_{\frac{s}{2}}$ with $|x - x'| \leq \frac{s}{4}$, the line segment [x, y] joining x and y lies in $N_{2s} \setminus N_{\frac{s}{4}}$. The desired estimate follows on the ball $B_{\tau_{\varepsilon}}(\xi_0)$ by estimating $Q(v_1, v_2)(x) - Q(v_1, v_2)(x')$ using the following gradient bound (we are using that $|\bar{u}_{\varepsilon} + v|^{p-1}$ is C^1 in this region)

$$\nabla Q(v_1, v_2) = (p-1) \left[(\bar{u}_{\varepsilon} + v_1 + t(v_1 - v_2)^{p-2} - \bar{u}_{\varepsilon}^{p-2}) \right] \nabla \bar{u}_{\varepsilon}$$

$$+ (p-1) (\bar{u}_{\varepsilon} + v_1 + t(v_1 - v_2)^{p-2} \nabla [v_1 + t(v_1 - v_2)]$$

$$= O(1) \bar{u}_{\varepsilon}^{p-3} (|v_1| + |v_2|) |\nabla u_{\varepsilon}| + O(1) \bar{u}_{\varepsilon}^{p-2} (|\nabla v_1| + |\nabla v_2|).$$

In fact, gradient bounds can also be used for the region $B_{\tau_{\varepsilon}}(\xi_0)$ if $p \geq 2$. For 1 , one can use the following inequalty

$$||\phi|^{p-1}(x) - |\phi|^{p-1}(x')| \le |\phi(x) - \phi(x')|^{p-1} \le ||\nabla \phi||_{C^0([x,x'])}^{p-1}|x - x'|^{p-1},$$

with $\phi = \bar{u}_{\varepsilon}$ and $\phi = \bar{u}_{\varepsilon} + v_1 + t(v_1 - v_2)$.

We conclude the lemma.

6. Appendix

The following lemma can be proven in the spirit of [Paca, Proposition 1.5.1]

Lemma 6.1. For $d \in \mathbb{R}$ set

$$\delta_j := \Re\left(\left(\frac{N-2}{2}\right)^2 + \lambda_j - d\right)^{\frac{1}{2}}, \quad j \in \mathbb{N}.$$
 (23)

Then for $\delta \in \mathbb{R} \setminus \{\pm \delta_j : j = 0, 1, ...\}$ there exists $C = C(N, \delta)$ such that if u is a solution to

$$\Delta u + \frac{d}{|x|^2} u = f \quad in \ B_1 \setminus \{0\},$$

then

$$||u||_{L^2_{\delta}(B_1)} \le C(||f||_{L^2_{\delta-2}(B_1)} + ||u||_{L^2(B_1 \setminus B_{\frac{1}{2}})}).$$

Lemma 6.2. Let ζ be a continuous function in \bar{B}_1 . Let δ_j be given by (23) with $d = \zeta(0)$. Then for $\delta \in \mathbb{R} \setminus \{\pm \delta_j : j = 0, 1, ...\}$ there exists a compact set $K \subset \bar{B}_1 \setminus \{0\}$ and a constant C > 0 such that for every $u \in L^2_{\delta}(B_1)$ solving

$$\Delta u + \frac{\zeta}{|x|^2} u = f$$
 in $B_1 \setminus \{0\}$, $f \in L^2_{\delta}(B_1)$,

we have

$$||u||_{L^2_{\delta}(B_1)} \le C(||f||_{L^2_{\delta-2}(B_1)} + ||u||_{L^2(K)}).$$

Proof. We rewrite the equation as

$$\Delta u + \frac{\zeta(0)}{|x|^2} u = f + \tilde{f}, \quad \tilde{f} := \frac{\zeta(0) - \zeta}{|x|^2} u.$$

Then by Lemma 6.1 we get

$$||u||_{L^2_{\delta}(B_1)} \le C_1(||f||_{L^2_{\delta-2}(B_1)} + ||\tilde{f}||_{L^2_{\delta-2}(B_1)} + ||u||_{L^2(B_1 \setminus B_{\frac{1}{\lambda}})}).$$

Let r > 0 be sufficiently small so that $|\zeta - \zeta(0)| \leq \frac{1}{2C_1}$ on B_r . Then

$$\|\tilde{f}\|_{L^{2}_{\delta-2}(B_{1})} \leq \frac{1}{2C_{1}} \|u\|_{L^{2}_{\delta}(B_{r})} + C(r, \|\zeta\|_{L^{\infty}(B_{1})}) \|u\|_{L^{2}(B_{1} \setminus B_{r})}.$$

The proof follows by absorbing the term $||u||_{L^2_{\delta}(B_r)}$ on the left hand side, and taking $K = \bar{B}_1 \setminus B_r$.

Lemma 6.3 (L^2 estimate). Let Ω be a bounded open set in \mathbb{R}^n . Let $b_i \in L^{\infty}(\Omega)$ with

$$||b_i||_{L^{\infty}(\Omega)} \leq \Lambda, \quad i = 0, 1, \dots, n.$$

Let $u \in L^2(\Omega)$ be a weak solution solution to

$$\Delta u + \sum_{i=1}^{n} b_i \frac{\partial u}{\partial x_i} + b_0 u = f \quad in \ \Omega,$$

for some $f \in L^2(\Omega)$. Then for every $\tilde{\Omega} \subseteq \Omega$ there exists $C = C(\tilde{\Omega}, \Lambda)$ such that

$$||u||_{W^{2,2}(\tilde{\Omega})} \le C(||f||_{L^2(\Omega)} + ||u||_{L^2(\Omega)}).$$

Lemma 6.4 (Schauder estimate). Let Ω be a bounded open set in \mathbb{R}^n . Let $b_i \in C^{0,\alpha}(\Omega)$ with

$$||b_i||_{C^{0,\alpha}(\Omega)} \leq \Lambda, \quad i = 0, 1, \dots, n.$$

Let u be a classical solution to

$$\Delta u + \sum_{i=1}^{n} b_i \frac{\partial u}{\partial x_i} + b_0 u = f \quad in \ \Omega,$$

for some $f \in C^{0,\alpha}(\Omega)$. Then for every $\tilde{\Omega} \subseteq \Omega$ there exists $C = C(\tilde{\Omega}, \Lambda)$ such that

$$||u||_{C^{2,\alpha}(\tilde{\Omega})} \le C(||f||_{C^{0,\alpha}(\Omega)} + ||u||_{C^0(\Omega)}).$$

Additionally, if Ω is regular, $\partial\Omega$ has two components Γ_1 and Γ_2 , and if u=0 on Γ_1 then for $\tilde{\Omega} \in (\Omega \cup \Gamma_1)$ there exists $C = C(\tilde{\Omega}, \Lambda)$ such that

$$||u||_{C^{2,\alpha}(\tilde{\Omega})} \le C(||f||_{C^{0,\alpha}(\Omega)} + ||u||_{C^{0}(\Omega)}).$$

References

[CP16] Mónica Clapp and Angela Pistoia. Symmetries, Hopf fibrations and supercritical elliptic problems. In *Mathematical Congress of the Americas*, volume 656 of *Contemp. Math.*, pages 1–12. Amer. Math. Soc., Providence, RI, 2016.

[Maz91] Rafe Mazzeo. Elliptic theory of differential edge operators. I. Comm. Partial Differential Equations, 16(10):1615–1664, 1991.

[MP96] Rafe Mazzeo and Frank Pacard. A construction of singular solutions for a semilinear elliptic equation using asymptotic analysis. *J. Differential Geom.*, 44(2):331–370, 1996.

[MP99] Rafe Mazzeo and Frank Pacard. Constant scalar curvature metrics with isolated singularities. Duke Math. J., 99(3):353–418, 1999.

[MS91] Rafe Mazzeo and Nathan Smale. Conformally flat metrics of constant positive scalar curvature on subdomains of the sphere. J. Differential Geom., 34(3):581–621, 1991.

[Paca] F. Pacard. Analysis in weighted spaces. https://cel.archives-ouvertes.fr/cel-00392164/file/Pacard.pdf.

[Pacb] F. Pacard. Connected sum constructions in geometry and in nonlinear analysis. http://www.cmls.polytechnique.fr/perso/pacard/Publications/Lecture-Part-I.pdf.

[PR00] Frank Pacard and Tristan Rivière. Linear and nonlinear aspects of vortices, volume 39 of Progress in Nonlinear Differential Equations and their Applications. Birkhäuser Boston, Inc., Boston, MA, 2000. The Ginzburg-Landau model.

[Sch88] Richard M. Schoen. The existence of weak solutions with prescribed singular behavior for a conformally invariant scalar equation. Comm. Pure Appl. Math., 41(3):317–392, 1988.

(Ali Hyder) Department of Mathematics, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218

E-mail address: ahyder4@jhu.edu

(Angela Pistoia) Dipartimento SBAI, "Sapienza" Università di Roma, via Antonio Scarpa 16, 00161 Roma, Italy

E-mail address: angela.pistoia@uniroma1.it

(Yannick Sire) Department of Mathematics, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218

E-mail address: sire@jhu.edu